

A human monoclonal antibody blocking SARS-CoV-2 infection

Running Head: A cross-neutralizing human antibody targeting SARS-CoV and SARS-CoV-2

Chunyan Wang^{a,**}, Wentao Li^{a,**}, Dubravka Drabek^{b,c,**}, Nisreen M.A. Okba^d, Rien van Haperen^{b,c}, Albert D.M.E. Osterhaus^e, Frank J.M. van Kuppeveld^a, Bart L. Haagmans^d, Frank Grosveld^{b,c,\$} and Berend-Jan Bosch^{a,*, \$}

Virology Division, Department of Biomolecular Health Sciences, Faculty of Veterinary Medicine, Utrecht University, Utrecht, the Netherlands^a; Department of Cell Biology, Erasmus MC, Rotterdam, the Netherlands^b; Harbour Antibodies B.V., Rotterdam, Netherlands^c; Department of Molecular and Cell Biology, Department of Viroscience, Erasmus Medical Center, Rotterdam, the Netherlands^d. University of Veterinary Medicine, Hannover, Germany^e

* Address correspondence to Berend-Jan Bosch, b.j.bosch@uu.nl.

** These authors are joint first authors. \$ These authors are joint senior authors.

Abstract

The emergence of the novel human coronavirus SARS-CoV-2 in Wuhan, China has caused a worldwide epidemic of respiratory disease (COVID-19). Vaccines and targeted therapeutics for treatment of this disease are currently lacking. Here we report a human monoclonal antibody that neutralizes SARS-CoV-2 (and SARS-CoV). This cross-neutralizing antibody targets a communal epitope on these viruses and offers potential for prevention and treatment of COVID-19.

Main text

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the etiological agent of the coronavirus induced disease 19 (COVID-19) that emerged in China late 2019 and causing a worldwide epidemic¹. As of March 10th 2020, over 115,113 cases have been reported in 109 countries, of which 4,063 (3.5%) succumbed to the infection². SARS-CoV-2 belongs to the *Sarbecovirus* subgenus (genus *Betacoronavirus*, family *Coronaviridae*)³ together with SARS-CoV that emerged in 2002 causing approximately 8000 infections with a lethality of 10%. Both viruses crossed species barriers from an animal reservoir and can cause a life-threatening respiratory illness in humans. Presently no approved targeted therapeutics are available for COVID-19. Monoclonal antibodies targeting vulnerable sites on viral surface proteins are increasingly recognised as a promising class of drugs against infectious diseases and have shown therapeutic efficacy for a number of viruses^{4, 5}.

Coronavirus neutralizing antibodies primarily target the trimeric spike (S) glycoproteins on the viral surface that mediate entry into host cells. The S protein has two functional subunits that mediate cell attachment (the S1 subunit, existing of four core domains S1_A through S1_D) and fusion of the viral and cellular membrane (the S2 subunit). Potent neutralizing antibodies often target the receptor interaction site in S1, disabling receptor interactions⁶⁻¹¹. The spike proteins of SARS-CoV-2 (SARS2-S; 1,273 residues, strain Wuhan-Hu-1) and SARS-CoV (SARS-S, 1,255 residues, strain Urbani) are 77.5% identical by primary amino acid sequence, are structurally very similar^{12, 13} and commonly bind the human angiotensin converting enzyme 2 (ACE2) protein as a host receptor^{1, 14} through their S1_B domain. Receptor interaction is known to trigger

irreversible conformational changes in coronavirus spike proteins enabling membrane fusion¹⁵.

In order to identify SARS-CoV-2 neutralizing antibodies, ELISA-(cross)reactivity was assessed of antibody-containing supernatants of a collection of 51 SARS-S hybridoma's derived from immunized transgenic H2L2 mice that encode chimeric immunoglobulins with human variable heavy and light chains and constant regions of rat origin (Suppl.Fig.1). Four of 51 SARS-S hybridoma supernatants displayed ELISA-cross-reactivity with the SARS2-S1 subunit (S residues 1-681; Suppl.Fig.1), of which one (47D11) exhibited cross-neutralizing activity of SARS-S and SARS2-S pseudotyped VSV infection. The chimeric 47D11 H2L2 antibody was reformatted and recombinantly expressed as a fully human IgG1 isotype antibody for further characterization.

The human 47D11 antibody binds to cells expressing the full-length spike proteins of SARS-CoV and SARS-CoV-2 (Fig.1a). The 47D11 antibody was found to potently inhibit infection of VeroE6 cells with SARS-S and SARS2-S pseudotyped VSV with IC₅₀ values of 0.06 and 0.08 µg/ml (Fig.1b), respectively. Authentic infection of VeroE6 cells with SARS-CoV and SARS-CoV-2 was neutralized with IC₅₀ values of 0.19 and 0.57 µg/ml (Fig.1c). Using ELISA 47D11 was shown to target the S1_B receptor binding domain (RBD) of SARS-S and SARS2-S. 47D11 bound the S1_B of both viruses with similar affinities as shown by the ELISA-based half maximal effective concentration (EC₅₀) values (0.02 and 0.03 µg/ml, respectively; Fig.2a). ELISA-based binding affinity of 47D11 for the spike ectodomain (S_{ecto}) of SARS-CoV was higher relative to that of SARS-CoV-2 (EC₅₀ values: 0.018 and 0.15 µg/ml, respectively), despite equimolar antigen coating (Suppl.Fig.2). Congruent with the ELISA-reactivities, measurement of binding kinetics of 47D11 by biolayer interferometry showed that 47D11 binds SARS-S_{ecto} with higher affinity (equilibrium dissociation constant [K_D]: 0.745 nM) relative to SARS2-S_{ecto} (K_D 10.8 nM) whereas affinity for SARS-S1_B and SARS2-S1_B was in a similar range (16.1 and 9.6 nM, respectively, Suppl.Fig.3). This difference may originate from differences in epitope accessibility in SARS-S versus SARS2-S, as domain B can adopt a closed and open conformation in the prefusion spike homotrimer^{12, 13}. Remarkably, binding of 47D11 to SARS-S1_B and SARS2-S1_B did not compete with S1_B binding to the ACE2 receptor expressed at the cell surface as shown by flow cytometry (Fig.2b; Suppl.Fig.4) nor with S_{ecto} and S1_B binding to soluble ACE2

in solid-phase based assay (Suppl.Fig.5), whereas two SARS-S1 specific antibodies 35F4 and 43C6 that neutralize SARS-S (but not SARS2-S) pseudotyped VSV infection (Suppl.Fig.6) do block binding of SARS-S_{ecto} and SARS-S1_B to ACE2. Using a trypsin-triggered cell-cell fusion assay, 47D11 was shown to impair SARS-S and SARS2-S mediated syncytia formation (Suppl.Fig.7). Our data show that 47D11 neutralizes SARS-CoV and SARS-CoV-2 through a yet unknown mechanism that is different from receptor binding interference. Alternative mechanisms of coronavirus neutralization by RBD-targeting antibodies have been reported including spike inactivation through antibody-induced destabilization of its prefusion structure¹⁵, which may also apply for 47D11.

The SARS2-S1_B receptor binding domain (residues 338-506) consists of a core domain and a receptor binding subdomain (residues 438-498) looping out from the antiparallel betasheet core domain structure that directly engages the receptor. Compared to the S1_B core domain, the protein sequence identity of the S1_B receptor interacting subdomain of SARS-S and SARS2-S is substantially lower (46.7% versus 86.3%; Suppl.Fig.8 and Fig.2c). Potent neutralizing antibodies often target this receptor binding subdomain. However, due to common variations in this subdomain, these antibodies are often virus-specific and bind and neutralize related viruses poorly^{16, 17}. The cross-reactive nature of 47D11 indicates that the antibody is more likely to target the conserved core structure of the S1_B receptor binding domain. S1_B binding by 47D11 further away from the receptor binding interface explains its inability to compromise spike-receptor interaction.

In conclusion, this is the first report on a (human) monoclonal antibody that neutralizes SARS-CoV-2. 47D11 binds a conserved epitope on the spike receptor binding domain explaining its ability to cross-neutralize SARS-CoV and SARS-CoV-2, using a mechanism that is independent of receptor binding inhibition. This antibody will be useful for development of antigen detection tests and serological assays targeting SARS-CoV-2. Neutralizing antibodies can alter the course of infection in the infected host supporting virus clearance or protect an uninfected host that is exposed to the virus⁴. Hence, this antibody offers the potential to prevent and/or treat COVID-19, and possibly also other future emerging diseases in humans caused by viruses from the *Sarbecovirus* subgenus.

References

1. Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*, 1-4 (2020).
2. World Health Organization. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200310-sitrep-50-covid-19.pdf?sfvrsn=55e904fb_2.
3. Coronaviridae Study Group of the International Committee on Taxonomy of Viruses. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat. Microbiol.* (2020).
4. Prabakaran, P. *et al.* Potent human monoclonal antibodies against SARS CoV, Nipah and Hendra viruses. *Expert opinion on biological therapy* **9**, 355-368 (2009).
5. Saphire, E. O., Schendel, S. L., Gunn, B. M., Milligan, J. C. & Alter, G. Antibody-mediated protection against Ebola virus. *Nat. Immunol.* **19**, 1169-1178 (2018).
6. Reguera, J. *et al.* Structural bases of coronavirus attachment to host aminopeptidase N and its inhibition by neutralizing antibodies. *PLoS Pathog.* **8**, e1002859 (2012).
7. Yu, X. *et al.* Structural basis for the neutralization of MERS-CoV by a human monoclonal antibody MERS-27. *Sci. Rep.* **5**, 13133 (2015).
8. Prabakaran, P. *et al.* Structure of severe acute respiratory syndrome coronavirus receptor-binding domain complexed with neutralizing antibody. *J. Biol. Chem.* **281**, 15829-15836 (2006).
9. Hwang, W. C. *et al.* Structural basis of neutralization by a human anti-severe acute respiratory syndrome spike protein antibody, 80R. *J. Biol. Chem.* **281**, 34610-34616 (2006).
10. Rockx, B. *et al.* Structural basis for potent cross-neutralizing human monoclonal antibody protection against lethal human and zoonotic severe acute respiratory syndrome coronavirus challenge. *J. Virol.* **82**, 3220-3235 (2008).
11. Widjaja, I. *et al.* Towards a solution to MERS: protective human monoclonal antibodies targeting different domains and functions of the MERS-coronavirus spike glycoprotein. *Emerg. Microbes Infect.* **8**, 516-530 (2019).
12. Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* (2020).
13. Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* (2020).
14. Li, W. *et al.* Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature* **426**, 450-454 (2003).

- 148 15. Walls, A. C. *et al.* Unexpected receptor functional mimicry elucidates activation of
149 coronavirus fusion. *Cell* **176**, 1026-1039. e15 (2019).
- 150 16. Tian, X. *et al.* Potent binding of 2019 novel coronavirus spike protein by a SARS
151 coronavirus-specific human monoclonal antibody. *Emerging Microbes & Infections* **9**,
152 382-385 (2020).
- 153 17. Menachery, V. D. *et al.* A SARS-like cluster of circulating bat coronaviruses shows
154 potential for human emergence. *Nat. Med.* **21**, 1508 (2015).
- 155 18. Li, F., Li, W., Farzan, M. & Harrison, S. C. Structure of SARS coronavirus spike
156 receptor-binding domain complexed with receptor. *Science* **309**, 1864-1868 (2005).
- 157 19. Raj, V. S. *et al.* Dipeptidyl peptidase 4 is a functional receptor for the emerging
158 human coronavirus-EMC. *Nature* **495**, 251-254 (2013).
- 159 20. Okba, N. M. A. *et al.* Sensitive and Specific Detection of Low-Level Antibody
160 Responses in Mild Middle East Respiratory Syndrome Coronavirus Infections. *Emerg.*
161 *Infect. Dis.* **25**, 1868-1877 (2019).

Figure Legends

Fig.1 47D11 neutralizes SARS-CoV and SARS-CoV-2. a) Binding of 47D11 to HEK-293T cells expressing GFP-tagged spike proteins of SARS-CoV and SARS-CoV-2 detected by immunofluorescence assay. The human mAb 7.7G6 targeting the MERS-CoV S_{1B} spike domain was taken along as a negative control, cell nuclei in the overlay images are visualized with DAPI. b) Antibody-mediated neutralization of infection of luciferase-encoding VSV particles pseudotyped with spike proteins of SARS-CoV and SARS-CoV-2. Pseudotyped VSV particles pre-incubated with antibodies at indicated concentrations (see methods) were used to infect VeroE6 cells and luciferase activities in cell lysates were determined at 24 h post transduction to calculate infection (%) relative to non-antibody-treated controls. The average $\pm$ SD from at least two independent experiments performed is shown. Iso-CTRL: irrelevant isotype monoclonal antibody. c) Antibody-mediated neutralization of SARS-CoV and SARS-CoV-2 infection on VeroE6 cells. The experiment was performed with triplicate samples, the average $\pm$ SD is shown.

Fig.2 The neutralizing 47D11 monoclonal antibody binds the receptor binding domain of SARS-CoV and SARS-CoV-2 spike proteins without eliminating S_{1B}/ACE2 receptor interaction. a) ELISA binding curves of 47D11 to S_{ecto} (upper panel) or S_{1A} and S_{1B} (RBD) (lower panel) of SARS-S and SARS2-S coated at equimolar concentrations. The average $\pm$ SD from at least two independent experiments performed is shown. b) Interference of antibodies with binding of the S_{1B} of SARS-CoV and SARS-CoV-2 to cell surface ACE2-GFP analysed by flow cytometry. Prior to cell binding, S_{1B} was mixed with mAb (mAbs 47D11, 35F4, 43C6, 7.7G6, in H2L2 format) with indicated specificity in a mAb:S_{1B} molar ratio of 8:1 (see Suppl.Fig.4 for an extensive analysis using different mAb:S_{1B} molar ratio's). Cells are analysed for (ACE2-)GFP expression (x-axis) and S_{1B} binding (y-axis). Percentages of cells that scored negative, single positive, or double positive are shown in each quadrant. c) Divergence in surface residues in S_{1B} of SARS-CoV and SARS-CoV-2. Upper panel: Structure of the SARS-CoV spike protein S_{1B} RBD in complex with human ACE2 receptor (PDB: 2AJF)¹⁸. ACE2 (wheat color) is visualized in ribbon presentation. The S_{1B} core domain (blue) and subdomain (orange) are displayed in surface presentation using PyMOL, and are visualized with the same colors in the

linear diagram of the spike protein above, with positions of the S1 and S2 subunits, the S ectodomain (S_{ecto}), the S1 domains S1_{A-D} and the transmembrane domain (TM) indicated. Lower panel: Similar as panel above with surface residues on S1_B of SARS-CoV that are at variance with SARS-CoV-2 colored in white.

Acknowledgements

We thank dr. Yoshiharu Matsuura (Osaka University, Japan) for the provision of the luciferase-encoding VSV-G pseudotyped VSVΔG-luc virus, and Yongle Yang, Michael van der Reijden and Rick Janssens for technical support. We thank Christian Drosten (Charité Universitätsmedizin Berlin, Germany) for provision of the SARS-CoV-2 virus. This study was done within the framework of National Centre for One Health (NCOH) and the Innovative Medicines Initiative (IMI) Zoonotic Anticipation and Preparedness Initiative [ZAPI project; grant agreement no. 115760]. The mice used in this study were provided by Harbour Antibodies BV, a daughter company of Harbour Biomed (<http://www.harbourbiomed.com>). C. Wang was supported by a grant from the Chinese Scholarship Council (file number CSC201708620178).

Author Contributions

B.J.B. designed and coordinated the study. C.W., W.L., N.M.A.O., R.v.H. and D.D. conducted the experiments. D.D., B.L.H. and B.J.B. supervised part of the experiments. All authors contributed to the interpretations and conclusions presented. B.J.B. wrote the manuscript, B.L.H., F.J.M.K., A.D.M.E.O. and F.G. participated in editing the manuscript.

Materials and Methods

Expression and purification of coronavirus spike proteins. Coronavirus spike ectodomains (S_{ecto}) of SARS-CoV (residues 1–1,182; strain CUHK-W1; GenBank: AAP13567.1) and SARS-CoV-2 (residues 1–1,213; strain Wuhan-Hu-1; GenBank: QHD43416.1) were expressed transiently in HEK-293T cells with a C-terminal

trimerization motif and Strep-tag using the pCAGGS expression plasmid. Similarly, pCAGGS expression vectors encoding S1 or its subdomains of SARS-CoV (S1, residues 1-676; S1_A, residues 1-302; S1_B, residues, 325-533), and SARS-CoV-2 (S1, residues 1-682; S1_A, residues 1-294; S1_B, residues 329-538) C-terminally tagged with Fc domain of human or mouse IgG or Strep-tag were generated as described before¹⁹. Recombinant proteins were expressed transiently in HEK-293T cells and affinity purified from the culture supernatant by protein-A sepharose beads (GE Healthcare) or streptactin beads (IBA) purification. Purity and integrity of all purified recombinant proteins was checked by coomassie stained SDS-PAGE.

Generation of H2L2 mAbs. H2L2 mice were sequentially immunized in two weeks intervals with purified S_{ecto} of different CoVs in the following order: HCoV-OC43, SARS-CoV, MERS-CoV, HCoV-OC43, SARS-CoV and MERS-CoV. Antigens were injected at 20-25 µg/mouse using Stimune Adjuvant (Prionics) freshly prepared according to the manufacturer instruction for first injection, while boosting was done using Ribi (Sigma) adjuvant. Injections were done subcutaneously into the left and right groin each (50 µl) and 100 µl intraperitoneally. Four days after the last injection, spleen and lymph nodes are harvested, and hybridomas made by standard method using SP 2/0 myeloma cell line (ATCC#CRL-1581) as a fusion partner. Hybridomas were screened in antigen-specific ELISA and those selected for further development, subcloned and produced on a small scale (100 ml of medium). For this purpose, hybridomas are cultured in serum- and protein-free medium for hybridoma culturing (PFHM-II (1X), Gibco) with addition of non-essential amino acids 100X NEAA, Biowhittaker Lonza, Cat BE13-114E). H2L2 antibodies were purified from hybridoma culture supernatants using Protein-A affinity chromatography. Purified antibodies were stored at 4°C until use.

Production of human monoclonal antibody 47D11. For recombinant human mAb production, the cDNA's encoding the 47D11 H2L2 mAb variable regions of the heavy and light chains were cloned into expression plasmids containing the human IgG1 heavy chain and Ig kappa light chain constant regions, respectively (InvivoGen). Both plasmids contain the interleukin-2 signal sequence to enable efficient secretion of recombinant antibodies. Recombinant human 47D11 mAb and previously described Isotype-control (anti-Streptag mAb) or 7.7G6 mAb were produced in HEK-293T cells following transfection with pairs of the IgG1 heavy and light chain expression plasmids

according to protocols from InvivoGen. Human antibodies were purified from cell culture supernatants using Protein-A affinity chromatography. Purified antibodies were stored at 4°C until use.

Immunofluorescence microscopy. Antibody binding to cell surface spike proteins of SARS-CoV, SARS-CoV-2 and MERS-CoV was measured by immunofluorescence microscopy. HEK-293T cells seeded on glass slides were transfected with plasmids encoding SARS-S, SARS2-S or MERS-S - C-terminally fused to the green fluorescence protein (GFP) - using Lipofectamine 2000 (Invitrogen). Two days post transfection, cells were fixed by incubation with 2% paraformaldehyde in PBS for 20 min at room temperature and stained for nuclei with 4,6-diamidino-2-phenylindole (DAPI). Cells were subsequently incubated with mAbs at a concentration of 10 µg/ml for 1 h at room temperature, followed by incubation with Alexa Fluor 594 conjugated goat anti-human IgG antibodies (Invitrogen, Thermo Fisher Scientific) for 45 min at room temperature. The fluorescence images were recorded using a Leica Spell confocal microscope.

Flow cytometry-based receptor binding inhibition assay. Antibody interference of S1_B binding to human ACE2 receptor on the cell surface was measured by flow cytometry. HEK-293T cells were seeded at a density of 2.5×10⁵ cells per ml in a T75 flask. After reaching 70~80% confluency, cells were transfected with an expression plasmid encoding human ACE2 - C-terminally fused to the GFP - using Lipofectamine 2000 (Invitrogen). Two days post transfection, cells were dissociated by cell dissociation solution (Sigma-aldrich, Merck KGaA; cat. no. C5914). 2.5 µg/ml of human Fc tagged SARS-S1_B and SARS2-S1_B was preincubated with mAb at the indicated mAb:S1_B molar ratios for 1 hour on ice and subjected to flow cytometry. Single cell suspensions in FACS buffer were centrifuged at 400×g for 10 min. Cells were subsequently incubated with S1_B and mAb mixture for 1 h on ice, followed by incubation with Alexa Fluor 594 conjugated goat anti-human IgG antibodies (Invitrogen, Thermo Fisher Scientific) for 45 min at room temperature. Cells were subjected to flow cytometric analysis with a CytoFLEX Flow Cytometer (Beckman Coulter). The results were analysed by FlowJo (version 10).

Pseudotyped virus neutralization assay. Production of VSV pseudotyped with SARS-S and SARS2-S was performed as described previously with some adaptations¹¹. Briefly, HEK-293T cells were transfected with pCAGGS expression

vectors encoding SARS-S or SARS2-S carrying a 28- or 18-a.a. cytoplasmic tail truncation, respectively. One day post transfection, cells were infected with the VSV-G pseudotyped VSV Δ G bearing the firefly (*Photinus pyralis*) luciferase reporter gene. Twenty-four hours later, supernatants containing SARS-S/SARS2-S pseudotyped VSV particles were harvested and titrated on African green monkey kidney VeroE6 cells. In the virus neutralization assay, mAbs were serially diluted at two times the desired final concentration in DMEM supplemented with 1% fetal calf serum (Bodinco), 100 U/ml Penicillin and 100 μ g/ml Streptomycin. Diluted mAbs were incubated with an equal volume of pseudotyped VSV particles for 1 hour at room temperature, inoculated on confluent VeroE6 monolayers in 96-well plated, and further incubated at 37°C for 24 hours. Luciferase activity was measured on a Berthold Centro LB 960 plate luminometer using D-luciferin as a substrate (Promega). The percentage of infectivity was calculated as ratio of luciferase readout in the presence of mAbs normalized to luciferase readout in the absence of mAb. The half maximal inhibitory concentrations (IC₅₀) were determined using 4-parameter logistic regression (GraphPad Prism version 8).

Virus neutralization assay. Neutralization of authentic SARS-CoV and SARS-CoV-2 was performed using a plaque reduction neutralization test (PRNT) as described earlier, with some modifications²⁰. In brief, mAbs were two-fold serially diluted and mixed with SARS-CoV or SARS-CoV-2 for 1 hour. The mixture was then added to VeroE6 cells and incubated for 1 hr, after which the cells were washed and further incubated in medium for 8 hrs. The cells were then fixed and stained using a rabbit anti-SARS-CoV serum (Sino Biological) and a secondary peroxidase-labelled goat anti-rabbit IgG (Dako). The signal was developed using a precipitate forming TMB substrate (True Blue, KPL) and the number of infected cells per well were counted using the ImmunoSpot® Image analyzer (CTL Europe GmbH). The half maximal inhibitory concentrations (IC₅₀) were determined using 4-parameter logistic regression (GraphPad Prism version 8).

ELISA analysis of antibody binding to CoV spike antigens. NUNC Maxisorp plates (Thermo Scientific) were coated with equimolar antigen amounts at 4°C overnight. Plates were washed three times with Phosphate Saline Buffer (PBS) containing 0.05% Tween-20 and blocked with 3% Bovine Serum Albumin (BSA) in PBS containing 0.1% Tween-20 at room temperature for 2 hours. Four-folds serial dilutions of mAbs starting

at 10 µg/ml (diluted in blocking buffer) were added and plates were incubated for 1 hour at room temperature. Plates were washed three times and incubated with HRP-conjugated goat anti-human secondary antibody (ITK Southern Biotech) diluted 1:2000 in blocking buffer for 1 hour at room temperature. An HRP-conjugated anti-StrepMab (IBA, Cat.no: 2-1509-001) antibody was used to corroborate equimolar coating of the Strep-tagged spike antigens. HRP activity was measured at 450 nanometer using tetramethylbenzidine substrate (BioFX) and an ELISA plate reader (EL-808, Biotek). Half-maximum effective concentration (EC₅₀) binding values were calculated by non-linear regression analysis on the binding curves using GraphPad Prism (version 8).

Fig.1







Fig.2




Suppl.Fig.1

Hybridoma	SARS-S _{ecto}	SARS-S1	SARS-S1 _A	SARS2-S1
44B3	2.5	2.7	3.3	0.1
45E10	3.0	0.8	1.7	0.0
46F11	2.4	2.7	3.3	0.0
39F9	2.9	3.3	3.5	0.0
41A7	2.6	1.0	1.9	0.0
28 E3	2.4	2.3	3.2	0.0
34C10	1.3	1.0	1.9	0.0
16C10	2.4	0.6	1.7	0.1
14B1	2.6	2.9	3.3	0.1
30B1	0.6	0.5	1.1	0.0
28G10	1.0	1.3	2.6	0.0
28F6	2.4	2.9	3.0	0.0
40H10	1.2	0.7	1.9	0.0
39A4	1.7	1.5	2.8	0.0
37G1	1.3	0.9	1.7	0.0
44E11	2.8	3.3	3.5	0.1
19C1	1.9	0.4	1.2	0.1
58D2	2.6	2.8	3.4	0.1
14C1	2.8	1.2	2.6	0.0
45H1	2.3	3.1	3.6	0.0
24F5	3.3	3.4	3.6	0.0
52D9	1.5	1.6	2.3	1.3
45E6	2.4	2.6	3.3	0.0
47D11	3.4	3.0	0.0	1.5
47G10	2.6	2.8	0.1	0.0
48G1	3.3	3.4	0.1	0.0
49F1	1.8	2.0	0.0	1.3
43C6	3.1	3.4	0.1	0.1
22E10	3.2	3.4	0.1	0.0
28D11	2.7	3.1	0.1	0.0
28H3	2.8	1.8	0.0	0.0
25E7	3.1	3.3	0.1	0.1
22E8	1.2	1.2	0.1	0.0
35F4	3.2	3.6	0.1	0.0
43G5	3.2	3.3	0.1	0.1
47F8	1.4	1.4	0.0	0.0
43B4	3.2	3.3	0.1	0.0
49B10	1.1	0.6	0.0	0.2
51C11	1.9	1.9	0.0	0.0
36F6	1.7	2.7	0.1	0.3
65H8	3.2	3.3	0.1	0.1
65H9	1.6	1.7	0.1	2.5
48D5	3.3	3.5	0.1	0.0
35E2	2.5	3.3	0.2	0.0
44G3	2.4	2.8	0.1	0.0
9H9	1.8	0.1	0.0	0.1
25C3	3.0	0.1	0.1	0.1
29E6	1.1	0.1	0.1	0.0
43F11	2.8	0.1	0.1	0.0
47C4	1.5	0.0	0.1	0.0
13F11	3.0	0.0	0.0	0.0


ELISA reactivity hybr. sups		# hybr sups
anti-SARS-S1 _A		23
anti-SARS-S1 (but not binding S1 _A)		22
anti-SARS-S _{ecto} (but not binding S1)		6
Total		51

Suppl. Fig.1. ELISA cross-reactivity of antibody-containing supernatants of SARS-S H2L2 hybridomas towards SARS2-S1. SARS-S targeting hybridomas were developed by conventional hybridoma technology from immunized H2L2 transgenic mice (Harbour Biomed), as described before¹. These mice - carrying genes encoding the heavy and light chain human immunoglobulin repertoire - were sequentially immunized with 2-week intervals with trimeric spike protein ectodomains (S_{ecto}) of three human coronaviruses from the betacoronavirus genus in the following order: 1. HCoV-OC43-S_{ecto}, 2. SARS-CoV-S_{ecto}, 3. MERS-CoV-S_{ecto}, 4. HCoV-OC43-S_{ecto}, 5. SARS-CoV-S_{ecto}, 6. MERS-CoV-S_{ecto}. Four days after the last immunization, splenocytes and lymph node lymphocytes were harvested and hybridomas were generated. Antibodies in the cell supernatants were tested for ELISA-reactivity against SARS-S_{ecto}, SARS-S1, SARS-S1_A and SARS2-S1. Of the 51 hybridoma supernatants that reacted with SARS-S_{ecto} only, 23 reacted with SARS-S1_A, 22 with SARS-S1 but not SARS-S1_A, 6 with SARS-S_{ecto} but not SARS-S1. Four of the 51 SARS-S_{ecto} hybridoma supernatants reacted with SARS2-S1 (see column on the right). The table displays ELISA-signal intensities (OD_{450nm} values) of hybridoma supernatants for the different antigens.

1) Widjaja, I. et al. Towards a solution to MERS: protective human monoclonal antibodies targeting different domains and functions of the MERS-coronavirus spike glycoprotein. Emerg. Microbes Infect. 8, 516-530 (2019).




Suppl. Fig.2. ELISA binding curve of the anti-StrepMAb (IBA) antibody to Strep-tagged spike antigens to corroborate equimolar ELISA plate coating of SARS-S_{ecto} / SARS2-S_{ecto} (upper panel), SARS-S1_B / SARS2-S1_B (middle panel) and SARS-S1_A / SARS2-S1_A (lower panel) antigens used in Fig.2a.




Spike protein	K _D (nM)	K _{on} (M ⁻¹ sec ⁻¹)	K _{off} (sec ⁻¹)
SARS-Secto	0.745 (± 0.532)	3.75 (± 1.27) × 10 ⁴	3.42 (± 3.34) × 10 ⁻⁵
SARS2-Secto	10.8 (± 2.46)	9.77 (± 4.09) × 10 ³	9.85 (± 3.61) × 10 ⁻⁵
SARS-S1 _B	16.1 (± 13.3)	2.00 (± 0.961) × 10 ⁴	2.10 (± 0.239) × 10 ⁻⁴
SARS2-S1 _B	9.56 (± 2.68)	1.51 (± 0.285) × 10 ⁴	1.37 (± 0.0728) × 10 ⁻⁴

Suppl. Fig.3. Binding kinetics of 47D11 to the S ectodomain and S1_B of SARS-CoV and SARS-CoV-2. Binding kinetics of 47D11 to immobilized recombinant SARS-S_{ecto}, SARS2-S_{ecto}, SARS-S1_B and SARS2-S1_B was measured using biolayer interferometry at 25°C, as described previously²¹. Kinetic binding assay was performed by loading 47D11 mAb at optimal concentration (42 nM) on anti-human Fc biosensor for 10 mins. Antigen association step was performed by incubating the sensor with a range of concentrations of the recombinant spike ectodomain (1600-800-400-200-100-50-25 nM) for 10 min, followed by a dissociation step in PBS for 60 min. The kinetics constants were calculated using 1:1 Langmuir binding model on Fortebio Data Analysis 7.0 software.


SARS-S1_B – ACE2 receptor binding




Binding controls



SARS2-S1_B – ACE2 receptor binding




Suppl. Fig.4. 47D11 does not prevent binding of SARS-S1_B and SARS2-S1_B to ACE2-expressing cells. Human HEK-293T cells expressing human ACE2-GFP proteins (see Methods) were detached and fixed with 2% PFA, incubated with a fixed amount of human Fc-tagged S1_B domain of SARS-S or SARS2-S that was preincubated for 1h with mAb (mAbs 47D11, 35F4, 43C6, 7.7G6, in H2L2 format) at the indicated mAb:S1_B molar ratios, and analysed by flow cytometry using a Alexa Fluor 594-conjugated secondary antibody targeting the human Fc tag. Cells are analysed for GFP expression (x-axis, GFP signal) and antibody binding (y-axis, Alexa 594 signal). Percentages of cells that scored negative, single positive, or double positive are shown in each quadrant. Binding controls include PBS-treated cells (mock), treatment of cells with SARS-S1_B and SARS2-S1_B in the absence of antibody, and cells treated with antibodies only. The experiment was performed twice, data from a representative experiment are shown.




Suppl. Fig.5. ELISA-based receptor binding inhibition assay. The ELISA-based receptor binding inhibition assay was performed as described previously with some adaptations¹. Recombinant soluble human ACE2 was coated on NUNC Maxisorp plates (Thermo Scientific) at 4°C overnight. Plates were washed three times with PBS containing 0.05% Tween-20 and blocked with 3% BSA in PBS containing 0.1% Tween-20 at room temperature for 2 hours. Recombinant S_{ecto} and S1_B of SARS-S or SARS2-S (300 ng) and serially diluted mAbs (mAbs 47D11, 35F4, 43C6, 7.7G6, in H2L2 format) were mixed for 1h at RT, added to the plate for 1 hour at room temperature, after which the plates were washed three times. Binding to ACE2 was detected using HRP-conjugated StrepMAb (IBA) that recognizes the C-terminal Streptag on the S_{ecto} and S1_B proteins.

1) Widjaja, I. et al. Towards a solution to MERS: protective human monoclonal antibodies targeting different domains and functions of the MERS-coronavirus spike glycoprotein. *Emerg. Microbes Infect.* 8, 516-530 (2019).



Suppl. Fig.6. H2L2 monoclonal antibodies 35F4 and 43C6 neutralize SARS-CoV but not SARS-CoV-2. Antibody-mediated neutralization of infection of VSV particles pseudotyped with spike proteins of SARS-CoV (upper panel) and SARS-CoV-2 (lower panel) by the 35F4 and 43C6 H2L2 antibodies targeting SARS-S1 but not SARS2-S1 (see Suppl.Fig.1). An irrelevant antibody was taken along as a human IgG1 isotype control. Means $\pm$ SD of triplicates are shown.

Suppl.Fig.7



Suppl. Fig.7. Cell-cell fusion inhibition assay. The cell-cell-fusion inhibition assay was performed as described previously with some adaptations¹. VeroE6 cells were seeded with density of 10⁵ cells per ml. After reaching 70~80% confluency, cells were transfected with plasmids encoding full length SARS-S, SARS2-S and MERS-S – C-terminally fused to GFP - using Lipofectamine 2000 (Invitrogen). The furin recognition site in the SARS2-S was mutated (R⁶⁸²RAR to A⁶⁸²AAR) to inhibit cleavage of the protein by endogenous furin and allow trypsin-induced syncytia formation. Two days post transfection, cells were pretreated DMEM only or DMEM with 20 µg/ml mAbs for 1 h and subsequently treated with DMEM with 15 µg/ml trypsin (to activate the spike fusion function) in the absence or presence of 20 µg/ml mAbs (47D11 crossreactive SARS-S and SARS2-S, 35F4 reactive to SARS-S, 7.7G6 reactive to MERS-S). After incubation at 37°C for 2 hrs, the cells were fixed with 2% PFA in PBS for 20 min at room temperature and stained for nuclei with 4,6-diamidino-2-phenylindole (DAPI). Cells expressing the S-GFP proteins were detected by fluorescence microscopy and S-mediated cell-cell fusion was observed by the formation of (fluorescent) multi-nucleated syncytia. The fluorescence images were recorded using a Leica Spell confocal microscope. The experiment was performed twice, data from a representative experiment are shown.

1) Widjaja, I. et al. Towards a solution to MERS: protective human monoclonal antibodies targeting different domains and functions of the MERS-coronavirus spike glycoprotein. *Emerg. Microbes Infect.* 8, 516-530 (2019).

Suppl.Fig.8

SARS - RBD	323	CPFGEVFNATKFPSVYAWERKKISNCVADYSVLYNSTFFSTFKCYGVSATKLNDLCFSNV
SARS2 - RBD	438	CPFGEVFNATRFASVYAWNRRKRISNCVADYSVLYNASFSTFKCYGVSPTKLNDLCFTNV
		*****:.* *****:.* *****:*****: ***** *****:.*
SARS - RBD	383	YADSFVVKGDDVRQIAPGQTGVIADYNYKLPDDFMGCVLAWNTRNIDATSTGNYNYKYRY
SARS2 - RBD	396	YADSFVIRGDEVQRQIAPGQTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGGNYYLYRL
		*****:.* ***** ***** *****:*****:.*:.* ***** **
SARS - RBD	443	LRHGKLRPFERDISNVPFSPDGKPCTP-PALNCYWPLNDYGFYTTTGIGYQPYRVVLSF
SARS2 - RBD	456	FRKSNLKPFERDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYPYRVVLSF
		:*.:*:*****:.. ...**.:*****:.* *.*:*****
SARS - RBD	502	ELLNAPATVCGP
SARS2 - RBD	516	ELLHAPATVCGP
		:**

Suppl. Fig.8. Protein sequence alignment of the S1_B receptor binding domain (RBD) of the SARS-CoV and SARS-CoV-2 spike proteins by ClustalW. Numbering denotes the residue position in the full-length spike protein of SARS-CoV (Genbank: AAP13441.1) and SARS-CoV-2 (Genbank: QHD43416.1). Asterisks (*) indicated fully conserved residues, the colon symbol (:) indicates conservation between groups of very similar properties, and the period symbol (.) indicates conservation between groups of weakly similar properties. Sequences corresponding to the S1_B receptor binding core domain and the receptor binding subdomain are colored in blue and orange, respectively. The fourteen residues that are involved in binding of SARS-CoV S1_B to human ACE2 are highlighted in grey¹.

1) Li, F., Li, W., Farzan, M. & Harrison, S. C. Structure of SARS coronavirus spike receptor-binding domain complexed with receptor. Science **309**, 1864-1868 (2005).