

Wrong person, place and time: viral load and contact network structure predict SARS-CoV-2 transmission and super-spreading events

Ashish Goyal¹, Daniel B. Reeves¹, E. Fabian Cardozo-Ojeda¹, Joshua T. Schiffer^{1,2,3*†}, Bryan T. Mayer^{1†}

¹ Vaccine and Infectious Diseases Division, Fred Hutchinson Cancer Research Center

² Department of Medicine, University of Washington, Seattle

³ Clinical Research Division, Fred Hutchinson Cancer Research Center

† These authors contributed equally to the work.

Corresponding author: Joshua T. Schiffer, jschiffe@fredhutch.org

One Sentence Summary: We developed a coupled within-host and between-host mathematical model to identify viral shedding levels required for transmission of SARS-CoV-2 and influenza, and to explain why super-spreading events occur more commonly during SARS-CoV-2 infection.

Abstract

SARS-CoV-2 is difficult to contain because most transmissions occur during the pre-symptomatic phase of infection. Moreover, in contrast to influenza, while most SARS-CoV-2 infected people do not transmit the virus to anybody, a small percentage secondarily infect large numbers of people. We designed mathematical models of SARS-CoV-2 and influenza which link observed viral shedding patterns with key epidemiologic features of each virus, including distributions of the number of secondary cases attributed to each infected person (individual R_0) and the duration between symptom onset in the transmitter and secondarily infected person (serial interval). We identify that people with SARS-CoV-2 or influenza infections are usually contagious for fewer than two days congruent with peak viral load several days after infection, and that transmission is unlikely below a certain viral load. SARS-CoV-2 super-spreader events with over 10 secondary infections occur when an infected person is briefly shedding at a very high viral load and has a high concurrent number of exposed contacts. The higher predisposition of SARS-CoV-2 towards super-spreading events is not due to its 1-2 additional weeks of viral shedding relative to influenza. Rather, a person infected with SARS-CoV-2 exposes more people within equivalent physical contact networks than a person infected with influenza, likely due to aerosolization of virus. Our results support policies that limit crowd size in indoor spaces and provide viral load benchmarks for infection control and therapeutic interventions intended to prevent secondary transmission.

Introduction

The SARS-CoV-2 pandemic is an ongoing tragedy that has caused 700,000 deaths and massively disrupted the global economy. The pandemic is rapidly expanding in the United States and is re-emerging focally in many countries that had previous success in limiting its spread.¹

Two features have proven challenging in containing outbreaks. First, most transmissions occur during the pre-symptomatic phase of infection.² Underlying this observation is a highly variable incubation period, defined as time between infection and symptom onset, which often extends beyond an infected person's peak viral shedding.³

Second, there is substantial over-dispersion of the basic reproduction number (R_0) for an individual infected with SARS-CoV-2,⁴ meaning that most infected people do not transmit at all, while a minority may transmit to dozens of people, with the average, population R_0 achieving a high enough level (>1) to allow exponential growth of cases in the absence of an effective intervention.⁵ Approximately 10-20% of infected people account for 80% of SARS-CoV-2 transmissions.^{4,6} Super-spreader events, in which the duration of contact between a single transmitter and large number of secondarily infected people is often limited to hours, are well documented.^{7,8} This pattern is not evident for influenza which has more homogeneous individual transmissions numbers.^{9,10} Differing shedding kinetics between the two viruses might explain this distinction; SARS-CoV-2 is often present intermittently in the upper airways for many weeks,^{11,12} while influenza is rarely shed for more than a week.¹³ Alternatively, SARS-CoV-2 aerosolization may predispose to wider exposure networks given the presence of an infected person in a crowded indoor space.

Viral load is recognized as a strong determinant of transmission risk. For influenza, the dose of viral exposure is related to the probability of infection in human challenge studies,¹⁴ and early treatment reduces household transmission.^{15,16} Household shedding of human herpesvirus-6 is closely linked to subsequent infection in newborns,¹⁷ and infants shedding high levels of cytomegalovirus in the oropharynx predictably transmit the virus back to their mothers.¹⁸

The epidemiology of viral infections can also be perturbed by biomedical interventions that lower viral load at mucosal transmission surfaces. Reduction of genital herpes simplex virus-2 shedding with antiviral treatments decreases probability of transmission.¹⁹ Suppressive antiretroviral therapy (ART) for HIV virtually eliminates the possibility of partner-to-partner sexual transmission and has limited community transmission dramatically.^{20,21}

These concepts are relevant for SARS-CoV-2 infection and require urgent attention as the pandemic continues to wreak havoc. Early therapies that lower peak viral load may reduce the severity of COVID-19 but may also decrease the probability of transmission and of super-spreader events.²² Similarly, the effectiveness of policies such as limiting mass gatherings, and enforcing mask use can be directly evaluated by their ability to reduce exposure viral load and transmission risk.²³ Here we developed a transmission simulation framework to capture the contribution of viral load to observed epidemiologic transmission metrics for influenza and SARS-CoV-2 and used this approach to explain why SARS-CoV-2 is predisposed to super-spreading events.

Results

Overall approach. We designed a series of steps to estimate the viral load required for SARS-CoV-2 and influenza transmission, as well as conditions required to explain the observed over-dispersion of secondary infections (*individual R0*) and frequent super-spreader events associated with SARS-CoV-2 but not influenza. This process included within-host modeling of viral loads, simulations of exposures and possible transmissions based on various transmission dose response curves, testing of various parameter sets against epidemiologic data and exploratory analyses with the best fitting model (**Fig S1**).

Within-host mathematical model of SARS CoV-2 shedding. First, we used our previously developed within-host mathematical model (equations in the **Methods**),²⁴ to generate plausible viral load patterns in the upper airway of an infected person or *transmitter* who could potentially transmit the virus to others (**Fig 1, Fig S2a**). Briefly, the model captures observed upper airway viral kinetics from 25 people from four different countries.²⁵⁻²⁸ Key observed features include an early viral peak followed by a decelerating viral clearance phase, which in turn leads to a temporary plateau at a lower viral load, ultimately followed by rapid viral elimination. Our model captures these patterns by including a density dependent term for early infected cell elimination and a nonspecific acquired immune term for late infected cell elimination.

One limitation of our model is that only half of study participants provided longitudinal viral load data from the very early days of infection when COVID-19 is often asymptomatic. Therefore, the model's output is most reliable for later time points. In particular, we have somewhat limited information on viral expansion rate and duration of peak shedding. To impute

possible variability, we generated a set of heterogeneous shedding curves in which the viral upslope, the downslope of viral load after peak and the viral load during plateau phase were varied (**Fig S2b**). Overall, the model generated several distinct patterns of infection: rapid elimination after the initial peak, a prolonged plateau phase with a low viral load, and a prolonged plateau phase with higher viral load. We simulated the transmission model with and without imputed heterogeneity.

Transmission dose response curves. We defined an *exposure event* in very specific biologic terms as a discrete event consisting of sufficient contact in time and space between a transmitter and one or more uninfected persons (*exposure contacts*) to allow for the possibility of a successful transmission. We next designed hundreds of dose response curves which separately predict contagiousness (CD curves) and infectiousness (ID curves) at a certain viral dose given an exposure contact. *Contagiousness* is defined as the viral load dependent probability of passage of virus-laden droplets or airborne particles from the airways of a potential transmitter to the airway of an exposure contact. *Infectiousness* is defined as the viral load dependent probability of transmission given direct airway exposure to virus in an exposure contact. *Transmission risk* is the product of these two mechanistic probabilities derived from the ID and CD curves and results is a transmission dose (TD) response curve. Each CD or ID curve is defined by its ID50 (λ) or viral load at which contagion or infection probability is 50% (**Fig S2c**), as well as its slope (α) (**Fig S2d**).²⁹ The TD50 is defined as viral load at which there is 50% transmission probability. We assumed equivalent curves for contagiousness and infectiousness for model fitting purposes. We also considered a simpler model with only a single TD curve (for *infectiousness*) and obtained qualitatively similar results (**Supplement and Methods**). Our

model includes the possibility that increasing viral load is not a key determinant of transmission when $\alpha=0.01$ (**Fig 2d**).

Exposure contact rate simulations. We introduced heterogeneity of exposure contact rates among possible transmitters by randomly selecting from a gamma distribution defined by mean number of exposure contacts per day (θ) and a scaling factor (ρ) that controls daily variability (**Fig S3**).

Transmission simulations. For each defined exposure contact, viral load in the transmitter was sampled and transmission risk was then identified based on the product of the CD and ID curves, or the TD curve (**Fig S2e, f; Fig 1**). Based on these probabilities, we stochastically modeled whether a transmission occurred for each exposure contact. This process was repeated when there were multiple possible exposure events within a given discretized time interval and the total number of exposures and transmissions within that interval was calculated.

For each successful transmission, we assumed that it takes τ days for the first infected cell to produce virus. To inform simulated values of *serial interval* (SI or time between symptom onset in the secondarily infected and transmitter), we randomly selected the *incubation period* (IP), for both the transmitter and the newly infected person, from a gamma distribution based on existing data (**Fig S4a**).^{3,30} Incubation period was defined as time from infection to the time of the onset of symptoms, where the mean incubation for SARS-CoV-2 is 5.2 days compared to 2 days for influenza.^{3,9,30}

Model fitting. In order to identify the parameter set that best recapitulated the observed data, we then simulated several hundred thousands of parameter sets with ~250 possible TD curves defined by ID50 and CD50 (λ) and slope (α), along with ~180 combinations of the mean exposed contact rate per day (θ) and associated variance parameter (ρ), and values of $\tau \in [0.5, 1, 2, 3]$ days. We aimed to identify the parameter set that best recapitulated the following features of the observed epidemiologic and individual-level data for SARS-CoV-2: mean R_0 across individuals ($R_0 \in [1.4, 2.5]$),^{3,4,6,31,32} mean serial interval across individuals ($SI \in [4.0, 4.5]$),^{3,31,33} cumulative distribution functions of individual R_0 ,^{4,6,34-36} and cumulative distribution functions of serial intervals derived from SARS-CoV-2 transmission pair studies that were conducted early during the pandemic,³¹ prior to any confounding influence of social distancing measures. Here, we define *individual R_0* as the total number of secondary transmissions from the transmitter in a fully susceptible population (**Methods**). We further checked the closeness of the solved ID curve with the observed relationship between viral RNA and infectious virus levels from a longitudinal cohort of infected people.³⁷

Influenza modeling. Next, we performed equivalent analyses for influenza to explain the lower frequency of observed super-spreader events with this infection. Influenza viral kinetics were modelled using a previously data-validated model.³⁸ Incubation periods for influenza are lower and less variable than for SARS-CoV-2 and were randomly selected for each simulation of the model using a gamma distribution (**Fig S4b**).³⁹ We again fit the model to: mean R_0 across individuals ($R_0 \in [1.1, 1.5]$),⁴⁰⁻⁴² mean serial interval ($SI \in [2.9, 4.3]$),⁹ cumulative distribution functions of individual R_0 corresponding to the 2008-2009 influenza A H1N1 pandemic with

mean $R_0=1.26$ and dispersion parameter=2.36 in the negative binomial distribution, and cumulative distribution functions of serial intervals.^{9,10,40}

Model-predicted individual R_0 and serial intervals for SARS-CoV-2 infection. A single model parameter set ($[\alpha, \lambda, \tau, \theta, \rho] = [0.8, 10^7, 0.5, 4, 40]$) most closely reproduces empirically observed individual R_0 and serial interval histograms (**Fig 2a, c**) and cumulative distribution functions (**Fig 2b, d**). Despite assuming that each infected person sheds at a high viral load for a period of time (**Fig 1, Fig S2b**), the model captures the fact that ~75% of 10,000 simulated transmitters do not infect any other people and that each increase in the number of possible transmissions is associated with a decreasing probability (**Fig. 2a**).

SARS-CoV-2 viral load was recently measured with viral RNA levels and mapped to concurrent level of infectious virus by dividing by approximately 25.³⁷ We divided observed viral RNA levels at each exposure contact by 25, and noted that the modeled ID curve closely recapitulates predicted quantitative viral culture level (**Fig S5**).

The model also generates super-spreader events with 10,000 simulated transmissions (**Fig. 2b**). If super-spreaders are defined as those who produce at least 5 secondary infections, we estimate that ~10% of all infected people and ~35% of all transmitters are super-spreaders. If super-spreaders are defined as those who produce at least 10 secondary infections, we estimate that ~6% of all infected people and ~25% of all transmitters are super-spreaders. If super-spreaders are defined as those who produce at least 20 secondary infections, we estimate that ~2.5% of all infected people and ~10% of all transmitters are super-spreaders. If super-spreaders are defined as those producing ≥ 5 , ≥ 10 , or ≥ 20 secondary infections, the contribution to all secondary infections is estimated at ~85%, ~70%, or ~44%, respectively (**Table 1**).

The model also recapitulates the high variance of the serial interval observed within SARS-CoV-2 transmission pairs, including negative values observed in the data (**Fig 2c, d**). We next project *generation time*, defined as the period between when an individual becomes infected and when they transmit the virus, for all transmission pairs and identify that the mean serial interval (4.4 days) provides an accurate approximation of mean generation time. However, the variance of generation time is considerably lower and by definition does not include negative values. A majority of generation times fell between 4 and 7 days, compared to -5 to 12 days for the serial interval (**Fig 2e**).

Viral load thresholds for SARS-CoV-2 transmission. The optimized ID curve has an ID₅₀ of 10^7 viral RNA copies and a moderately steep slope (**Fig 3a**). The TD₅₀ for SARS-CoV-2 is slightly higher at $10^{7.5}$ viral RNA copies (**Fig 3a**). To assess the impact of these parameters on transmission, we performed simulations with 10,000 transmitters and concluded that transmission is very unlikely (~0.00005%) given an exposure to an infected person with an upper airway viral load of $<10^4$ SARS-CoV-2 RNA copies, and unlikely (~0.002%) given an exposure to an infected person with a viral load of $<10^5$ SARS-CoV-2 RNA copies. On the other hand, transmission is much more likely (39%) given an exposure to an infected person who is shedding $>10^7$ SARS-CoV-2 RNA copies, and 75% given an exposure to an infected person with a viral load of $>10^8$ SARS-CoV-2 RNA copies. We obtain similar results (not shown) when we solve our model using the assumption of homogeneous viral load trajectories as in **Fig S2a**.

Narrow duration of high infectivity during SARS-CoV-2 infection. We next plotted the probability of infection given an exposure to a transmitter. Under multiple shedding scenarios,

the window of high probability transmission is limited to time points around peak viral load, and some heterogeneity in regard to peak infectivity is noted between people (**Fig 3b-d**). In general, infected persons are likely to be most infectious (i.e., above TD50) for a ~0.5-1.0-day period between days 2 and 6 after infection. We therefore conclude that the observed wide variance in serial interval (**Fig 2c**) results primarily from the possibility of highly discrepant incubation periods between the transmitter and infected person, rather than wide variability in shedding patterns across transmitters.

Requirements for SARS CoV-2 super-spreader events. The solved value for exposed contact network heterogeneity (ρ) is 40 indicating high variability in day-to-day exposure contact rates (**Fig S3d**) with a high average number of exposed contacts per day ($\theta=4$). We generated a heatmap from our TD curve to identify conditions required for super-spreader events which included viral load exceeding 10^7 SARS CoV-2 RNA copies and a high number of daily exposure contacts per day. We observe an inflection point between 10^6 and 10^7 SARS CoV-2 RNA copies where large increases in the number of daily exposure contacts have a more limited impact on increasing the number of transmissions from a single person (**Fig 4a**). The exposure contact network occasionally results in days with ≥ 150 exposure contacts per day, which may allow an extremely high number of secondary infections from a single person (**Fig 4a**).

We next plotted transmission events simulated on a daily basis over 30 days since infection from 10,000 transmitters according to viral load at exposure and number of exposure contacts on that day (**Fig 4b**). Secondary transmissions to only 1-3 people occurred almost exclusively with daily numbers of exposure contacts below 10 with any exposure viral load exceeding 10^6 RNA copies or with higher numbers of exposure contacts per day and viral loads

exceeding 10^5 RNA copies. Massive super-spreader events with over 50 infected people almost always occurred at viral loads exceeding 10^7 RNA copies / day with high levels of concurrent exposure contacts (**Fig 4b**).

We next identified that over 50% of secondary infections were associated with a transmitter who has a high number of exposed contacts (11-100 per day) and a viral load exceeding 10^6 RNA copies (**Fig 4c**), which is the mechanistic underpinning of why ~70% of all secondary infections arose from transmitters who produced more than 10 secondary infections (**Table 1**).

Model predicted individual R_0 and serial intervals for influenza infection. A single model parameter set most closely reproduced empirically observed histograms and cumulative distribution functions for individual R_0 and serial intervals for influenza: $(\alpha, \lambda, \tau, \theta, \rho) = (0.7, 10^{5.5}, 0-0.5, 4, 1)$. ID50 values for influenza are lower than SARS CoV-2, but a direct comparison cannot be made because tissue culture infectious dose (TCID) has been more commonly used for measurements of influenza viral load, whereas viral RNA is used for SARS-CoV-2. Nevertheless, TCID is a closer measure of infectious virus and it is thus reasonable that ID50 based on TCID for influenza would be ~30-fold lower than ID50 based on total viral RNA (infectious and non-infectious virus) for SARS-CoV-2.³⁷

The other notable difference is a considerably lower ρ value for influenza (**Fig S3b**), denoting much less heterogeneity in the number of exposure contacts per person while the average daily exposure contact was the same for both viruses (4 per day). The model captures the fact that 40% of influenza infected people do not transmit to anyone else and that each increase in the number of individual transmissions is associated with a lower probability (**Fig. 5a**).

Relative to SARS-CoV-2, super-spreader events involving 5 or more people are predicted to be 5-fold less common overall and 10-fold less common among transmitters (~2% of all infected people and ~3% of transmitters) (**Fig. 5b, Table 1**). Super-spreaders defined as those infecting ≥ 5 individuals contribute to only ~10% to all transmissions (**Table 1**).

The model also recapitulates the lower variance of serial interval for influenza relative to SARS-CoV-2 (**Fig 5c, d**). We next identified that the mean and variance of the serial interval provide good approximations of the mean and variance for generation time. A majority of generation times fell between 2 and 6 days (**Fig 5e**).

Viral load thresholds for influenza transmission. Based on the optimized TD curve for influenza (**Fig 6a**), we next plotted the probability of infection given an exposure to an infected person. The TD50 for influenza is $10^{6.1}$ TCID/mL. Under various shedding scenarios, the window of high probability transmission is limited to time points around peak viral load (**Fig 6b-d**). In general, infected persons are likely to be most infectious (i.e., above TD50) for a ~0.5-1.0 day period.. The observed narrow variance in serial interval (**Fig 5c**) results primarily from the narrow range of incubation periods within the transmitter and secondarily infected person, as well as the limited variability in shedding patterns across transmitters.

Determinants of influenza individual R_0 . We generated a heat map from our TD curve to identify conditions governing influenza transmission to multiple people including viral load exceeding 10^6 influenza TCID and a high number of exposure contacts per day. The contact network never results in days with more than 15 exposure contacts per day, which severely limits

the possible number of transmissions from a single person relative to SARS-CoV-2 (**Fig 7a, S3b**).

We plotted transmission events simulated on a daily basis over 30 days since infection from 10,000 transmitters according to viral load at exposure and number of exposure contacts on that day (**Fig 7b**). Secondary transmissions to fewer than 5 people accounted for 90% of infections (**Table 1**) and occurred with fewer than 10 daily exposure contacts and exposure viral loads exceeding 10^4 TCID. Small scale super-spreader events with 5-10 infected people almost always occurred at viral loads exceeding 10^5 TCID with 5-10 concurrent exposure contacts (**Fig 7b**).

We next identified that over 50% of infections were associated with a transmitter who had fewer than 10 exposure contacts per day and a viral load exceeding $10^{4.5}$ TCID (**Fig 7c**), which is why no infected person ever transmitted to more than 10 other people (**Table 1**).

Differing exposed contact distributions, rather than viral kinetics, explain SARS CoV-2 super-spreader events. We sought to explain why SARS-CoV-2 has a more over-dispersed distribution of individual R_0 relative to influenza. To assess viral kinetics as a potential factor, we comparatively plotted transmission risk per exposure contact as a function of time since infection in 10,000 transmitters for each virus. The median per contact transmission risk is slightly higher for influenza; however, 75% and 95% transmission risks are marginally higher for SARS-CoV-2 compared to influenza with slightly higher peak transmission risk, and a longer tail of low transmission risk beyond 7 days (**Fig 8a**). The transmission risk was considerably higher for the 25% of simulated SARS-CoV-2 infections with the highest viral loads, suggesting that a substantial subset of infected people may be more pre-disposed to super-spreading. When plotted

as time since onset of symptoms the variability in transmission potential is considerably larger for persons with high SARS-CoV-2 viral load, owing to the variable incubation period of this virus (**Fig 8b**).

The median duration of shedding over infectivity thresholds was short and nearly equivalent for both viruses. For SARS-CoV-2 and influenza, median [range] time above ID10 was 2.7 [0, 7] and 2.4 [1.6, 3.7] days respectively; median time above ID25 was 1.7 [0, 3] and 1.5 [0, 2.2] days respectively; median time above ID50 was 0.8 [0, 1.3] and 0 [0, 1.3] days respectively; median time above ID75 was 0 [0, 0.4] and 0 [0, 0] days respectively; median time above ID90 was 0 [0, 0] and 0 [0, 0] days respectively. ID10, ID25 and ID50 values are more variable across SARS-CoV-2 simulations due to a minority of trajectories with prolonged moderate viral loads.

For SARS-CoV-2 and influenza, median [range] time above TD10 was 1.4 [0, 2.5] and 1.2 [0, 2.0] days respectively; median time above TD25 was 0.8 [0, 1.3] and 0.3 [0, 1.3] days respectively; median time above TD50 was 0 [0, 0.5] and 0 [0, 0.4] days respectively; median time above TD75 was 0 [0, 0] and 0 [0, 0] days respectively. TD10, TD25 and TD50 values are more variable across SARS-CoV-2 simulations due to a minority of trajectories with prolonged moderate viral loads (**Fig 8c**).

We next plotted the frequency of exposure contacts per day for both viruses and noted a higher frequency of days with no exposed contacts (**Fig 8d**), but also a higher frequency of days with more than 10 exposure contacts (**Fig 8e**) for SARS-CoV-2 relative to influenza, despite an equivalent mean number of daily exposure contacts. To confirm that this distribution drives the different observed distributions of individual R_0 values (**Fig 8f**), we simulated SARS-CoV-2 infection with an assumed $\rho=1$ and generated a distribution of individual R_0 similar to that of

influenza (**Fig S6a**). Similarly, we simulated influenza infection with an assumed $\rho=40$ and generated a distribution of individual R_0 similar to that of SARS-CoV-2 (**Fig S6b**). Under all scenarios, predicted distributions of serial interval (**Fig 8g, Fig S6**) and generation time (**Fig 8h, Fig S6**) were unchanged by shifts in the exposed contact network.

Projections of targeted physical distancing. Physical distancing is a strategy to decrease R_0 . We simulated a decrease in the contact rate uniformly across the population and noted a decrease in population R_0 (**Fig S7a**) as well the percent of infected people who will transmit (**Fig 7b**) and become super-spreaders (**Fig S7c-d**). An approximately 40% decrease in the average exposed contact rate decreased R_0 below 1 (**Fig S6a**). We further investigated whether lowering contact rate among larger groups only, in particular by banning exposure events with a high number of exposure contacts, could control the epidemic. We identify that limiting exposure contacts to no more than 5 per day is nearly equivalent to limiting exposure contacts altogether and that only a small decrease in mean exposure contact rate can achieve $R_0 < 1$ if exposure events with less than 20 contacts are eliminated (**Fig S8**).

Pre-symptomatic transmission and super-spreading risk. Much of the highest transmission risk for SARS-CoV-2 exists in the pre-symptomatic phase (**Fig 8b**) which explains why 62% of simulated transmissions occurred in the pre-symptomatic phase for SARS-CoV-2, compared to 10% for influenza. Similarly, 62% and 21% of SARS-CoV-2 and influenza super-spreader events with secondary transmissions ≥ 5 and 39% of SARS-CoV-2 super-spreader events with secondary transmissions $R_0 \geq 10$ fell in the pre-symptomatic period.

Discussion

Our results demonstrate that SARS-CoV-2 shedding kinetics are directly linked to the virus' most fundamental epidemiologic properties. First, we identify a transmission dose response curve which specifies that a nasal viral load below 10^5 RNA copies is unlikely to commonly result in transmission. For SARS-CoV-2, this threshold is consistent with the overall rarity of positive cultures at these levels.³⁷ We also predict a relatively steep TD curve such that transmission becomes much more likely when shedding exceeds 10^8 viral RNA copies and there is an exposure contact between an infected person and susceptible person. The amount of viral RNA can be roughly converted to an estimate of viral quantity by culture which approximates infectiousness. Our results therefore have relevance for dosing of SARS-CoV-2 in human challenge experiments that are being considered for vaccine trials.

While the duration of shedding for SARS-CoV-2 is often three weeks or longer,^{11,12} we predict that the duration of shedding above thresholds required for a moderate probability of transmission per contact is much shorter, often less than half a day, and is comparable to that of influenza. While transmission after the first week of infection is quite rare, our model is consistent with the observation that transmissions commonly occur during the pre-symptomatic phase of infection,² given the highly variable incubation period associated with SARS-CoV-2.

The observed high heterogeneity in serial interval is attributable almost entirely to the variable nature of the incubation period, rather than transmission occurring extremely late after infection. While our estimate for mean generation time is equivalent to that of mean serial interval, it is notable that the range of SARS-CoV-2 serial intervals is much wider than the range of generation times. This result is evident even though we built substantial heterogeneity into our viral shedding curves beyond that observed in the somewhat limited existing shedding data.

The finding of limited duration of SARS-CoV-2 infectivity has practical implications. First, considerable resources are being used in hospitals and skilled nursing facilities to isolate patients with persistent SARS-CoV-2 shedding. We propose that a low nasal viral load, particularly during late infection, need not justify full patient isolation procedures in the absence of aerosolizing procedures. This observation could save substantial hospital resources and valuable isolation beds during subsequent waves of infection. Similar considerations are relevant for employees wishing to return to work. Our results also suggest that time since first positive test may be predictive of lack of contagion, though more viral load kinetic studies will be needed to confirm the existing observation that viral loads after a week of infection are usually low and associated with negative viral cultures.³⁷ Finally, our conclusions are supportive of rapid, less sensitive assays which are more likely to detect infection at periods of contagion.⁴³

Many of these conclusions, including specific viral load thresholds for transmission, a steep dose response curve and a maximum 2-day duration of contagion within an infected individual are equally relevant for influenza infection. One important difference is that incubation periods for influenza are far less variable which means that at the individual level, the serial interval is much more likely to be predictive of the generation time.

Another finding is that SARS-CoV-2 super-spreading events are dependent on a large number of exposure contacts during the relatively narrow 1-2 days window during which a ~25% subset of infected people is shedding at extremely high levels above the TD50. Because we predict that super-spreader potential may be somewhat of a generalized property of infection, rather than a characteristic of a tiny subset of infected people, this result also has practical implications. A common experience during the pandemic has been early identification of a cluster of infected people within a specific confined environment such as a senior living home,

crowded work environment, athletic team, or restaurant. Our results demonstrate that newly diagnosed people within small clusters may be past the peak of their super-spreading potential. At this stage, many more infections have often been established and drastic quarantine procedures should be considered. Other undiagnosed, pre-symptomatic infected people may have super-spreader potential while the known infected person is no longer contagious, highlighting the importance of effective contact tracing.

At the prevention level, school opening and work opening strategies should focus on severely limiting the possible number of exposure contacts per day. Where large numbers of exposure contacts are unavoidable, mandatory masking policies, perhaps with N95 masks that may more significantly lower exposure viral loads should be considered.²³

Influenza infection is much less predisposed to super-spreader events than SARS-CoV-2. Yet, influenza shedding at levels above those required for a high probability of transmission occurs with only slightly lower frequency. Therefore, the markedly different probability of super-spreader events between the two viruses is unlikely to relate to different viral host kinetics, despite the fact that the overall duration of SARS-CoV-2 shedding exceeds duration of influenza shedding often by more than two weeks.

Rather, our analysis suggests that the exposure contact networks of SARS-CoV-2 transmitters are highly variable relative to those of influenza. One possible explanation underlying this finding is that SARS-CoV-2 is more predisposed to airborne transmission than influenza.⁴⁴ Here our precise definition of an exposure contact (sufficient contact between a transmitter and an uninfected person to potentially allow transmission) is of high relevance. Our result suggests that a SARS-CoV-2 infected person in a crowded, poorly ventilated room, may generate more exposure contacts than an influenza infected person in the same room, likely

based on wider dispersal and / or longer airborne survival of the virus. Thus, our results suggest a possible downstream quantitative effect of airborne transmission on SARS-CoV-2 epidemiology. Another possibly important variable is that pre-symptomatic transmission, which is a common feature of SARS-CoV-2 may predispose to multiple transmissions. This prediction reinforces current public health recommendation to avoid crowded indoor spaces with poor air recirculation.

On the other hand, a much higher proportion of SARS-CoV-2 infected people than influenza infected people do not transmit at all. This result lacks a clear mechanistic explanation but may imply that aerosolization occurs only in a subset of infected people. One theoretical explanation is that high viral load shedding in the pre-symptomatic phase is defined by lack of cough or sneeze leading to limited spatial diffusion of virus. Alternatively, it is also possible that a proportion of infected people never shed virus at high enough viral loads to allow efficient transmission. This possibility speaks to the need for more quantitative viral load data gathered during the initial stages of infection.

Age cohort structure differs between the two infections, with a lower proportion of observed pediatric infections for SARS-CoV-2. If adults have more high exposure events than children, then this could also explain super-spreader events. We are less enthusiastic about this hypothesis. First, SARS-CoV-2 super-spreader events have occurred in schools and camps and would likely be more common in the absence of widespread global school closures in high prevalence regions. Second, a sufficient proportion of influenza cases occur in adults to rule out the presence of frequent large super-spreading events in this population.

Our analysis has important limitations. First, exposure contacts were assumed to be homogeneous and we do not capture the volume of the exposing aerosol or droplet. For instance,

if a large-volume droplet contains ten times more viral particles than an aerosol droplet, then the exposure could be dictated by this volume as well as the viral load of the potential transmitter. It is possible that under rare circumstances with extremely high-volume exposures, even persons with extremely low viral loads may transmit. Second, based on the quality of available data, we fit our models for SARS-CoV-2 and influenza to viral RNA and viral culture respectively. Existing data suggest that kinetics of viral RNA and culture are similar during both infections, with culture having lower sensitivity to detect virus.³⁷ Third, our intra-host model of SARS-CoV-2 was fit to heterogeneous data with different sampling techniques and PCR assays.²⁴ Moreover, R_0 estimates have varied across the globe. Our estimates of TD50 are necessarily imprecise based on available data and should serve only as a conservative benchmark. Most importantly, we cannot rule out the possibility that a small minority of infected people shed at sufficient levels for transmission for much longer than has been observed to date. Finally, contagiousness could have different dose response dynamics than viral load dependent infectiousness and may require investigation in the future upon the availability of epidemiologically relevant additional data.

In conclusion, fundamental epidemiologic features of SARS-CoV-2 and influenza infections can be directly related to viral shedding patterns in the upper airway as well as the nature of exposure contact networks. We contend that this information should be leveraged for more nuanced public health practice in the next phase of the pandemic.

Methods

SARS-CoV-2 within-host model. To simulate SARS-CoV-2 shedding dynamics, we employed our previously-described viral infection model.²⁴ In this model, susceptible cells (S) after coming into contact with SARS-CoV-2 (V) become infected at rate βVS . The infected cells (I) produce new virus at a per-capita rate π . The model also includes the clearance of infected cells in two ways: (1) by an innate response with density dependent rate δI^k ; and (2) an acquired response with rate $\frac{mE^r}{E^r + \phi^r}$ mediated by SARS-CoV-2-specific effector cells (E). The clearance mediated by innate immunity depends on the infected cell density and is controlled by the exponent k . The Hill coefficient r parameterizes the nonlinearity of the second response and allows for rapid saturation of the killing. Parameter ϕ defines the effector cell level by which killing of infected cells by E is half maximal.

In the model, SARS-CoV-2-specific effector cells rise after 2 stages from precursors cells (M_1 and M_2). The first precursor cell compartment (M_1) proliferates in the presence of infection with rate ωIM_1 and differentiates into the effector cell at a per capita rate q during the next intermediate stage. Finally, effector cells die at rate δ_E . The model is expressed as a system of ordinary differential equations:

$$\begin{aligned}\frac{dS}{dt} &= -\beta VS \\ \frac{dI}{dt} &= \beta VS - \delta I^k I - m \frac{E^r}{E^r + \phi^r} I \\ \frac{dV}{dt} &= \pi I - \gamma V \\ \frac{dM_1}{dt} &= \omega I M_1 - q M_1 \\ \frac{dM_2}{dt} &= q(M_1 - M_2) \\ \frac{dE}{dt} &= q M_2 - \delta_E E\end{aligned}$$

471

472 We assumed $S(0) = 10^7$ cells/mL, $I(0) = 1$ cells/mL, $V(0) = \frac{\pi I(0)}{c}$ copies/mL, $M_1(0) = 1$,

473 $M_2(0) = 0$ and $E_0 = 0$.

474 When we introduce simulated heterogeneity in cases of SARS-CoV-2 (by increasing
475 the standard deviation of the random effects of parameters β by 20, δ by 2, k by 2 and π by 5 in
476 the original distribution from²⁴), some of the viral shedding curves suggest that viral shedding
477 could continue for long period (over 6 weeks). Indeed, while median viral shedding duration has
478 been estimated at 12-20 days, shedding for many months is also observed commonly.⁴⁵ We
479 assumed that viral loads after day 20 drop to a exposure-level viral load level (i.e., $V(0)$) as most
480 viral shedding observed after this point is transient and at an extremely low viral load.⁴⁶ The
481 population distribution of parameters to simulate artificial SARS-CoV-2 viral shedding dynamics
482 is provided in **Table S1**.

483

484 **Influenza within-host model.** To simulate viral shedding dynamics of influenza viral, we employ
485 a model³⁸ that is a simplified version of the viral dynamics model presented for SARS-CoV-2.
486 This model assumes $k = 0$ and $m = 0$ and can be expressed as a system of ordinary differential
487 equations:

$$\begin{aligned}\frac{dS}{dt} &= -\beta VS \\ \frac{dI}{dt} &= \beta VS - \delta I \\ \frac{dV}{dt} &= \pi I - \gamma V\end{aligned}$$

488 Following this model,³⁸ we assumed $S(0) = 4 \times 10^8$ cells/mL, $I(0) = 1$ cells/mL, $V(0) = \frac{\pi I(0)}{c}$
 489 copies/mL. To simulate artificial influenza viral shedding dynamics, we assumed the population
 490 distribution of parameters $\text{Log}_{10}(\beta)$, $\text{Log}_{10}(\pi)$, $\text{Log}_{10}(\gamma)$ and $\text{Log}_{10}(\delta)$ are -4.56 (0.17), -
 491 1.98 (0.14), 0.47 (0.03) and 0.60 (0.06), respectively.

492
 493 **Dose-response model.** For both viruses, to estimate the infectiousness $P_t[V(t)]$ (response) based
 494 on viral loads $V(t)$ (dose), we employed the function, $P_t[V(t)] = \frac{V(t)^\alpha}{\lambda^\alpha + V(t)^\alpha}$. Here, λ is the
 495 infectivity parameter that represents the viral load that corresponds to 50% infectiousness and
 496 50% contagiousness, and α is the Hill coefficient that controls the sharpness in the dose-response
 497 curve.

498
 499 **Transmission Model and Reproduction number.** Our transmission model assumes that only
 500 some contacts of an infected individual with viral load dependent infectiousness are physically
 501 exposed to the virus (defined as exposure contacts), that only some exposure contacts have virus
 502 passaged to their airways (contagiousness) and that only some exposed contacts with virus in
 503 their airways become secondarily infected (successful secondary infection). Contagiousness and
 504 infectiousness are then treated as viral load dependent multiplicative probabilities with
 505 transmission risk for a single exposure contact being the product. Contagiousness is considered
 506 to be viral load dependent based on the concept that a transmitter's dispersal cloud of virus is

more likely to prove contagious at higher viral load, which is entirely separate for considerations of viral infectivity within the airway once a virus contacts the surface of susceptible cells.

We next assume that the total exposed contacts within a time step (η_{Δ_t}) is gamma distributed, i.e. $\eta_{\Delta_t} \sim \Gamma\left(\frac{\theta}{\rho}, \rho\right) \Delta_t$, using the average daily contact rates (θ) and the dispersion parameter (ρ). To obtain the true number of exposure contacts with airway exposure to virus, we simply multiply the contagiousness of the transmitter with the total exposed contacts within a time step (i.e., $\zeta_t = \eta_{\Delta_t} P_t$).

Transmissions within a time step are simulated stochastically using time-dependent viral load to determine infectiousness (P_t). Successful transmission is modelled stochastically by drawing a random uniform variable ($U(0,1)$) and comparing it with infectiousness of the transmitter. In the case of successful transmission, the number of secondary infections within that time step (T_{Δ_t}) is obtained by the product of the infectiousness (P_t) and the number of exposure contacts drawn from the gamma distribution (ζ_t). In other words, the number of secondary infections for a time step is $T_{\Delta_t} = Ber(P_t) P_t \eta_{\Delta_t}$. If we disregard contagiousness by assuming $P_t = 1$ in ζ_t , we identify that there are little to no differences on overall results other than the emergent TD curve and optimal parameter set describing dose-response curve and exposed contact network, which no longer agrees as closely with in vitro probability of positive virus culture (**Fig S5**).³⁷

We obtain the number of secondary infections from a transmitter on a daily basis noting that viral load, and subsequent risk, does not change substantially within a day. We then summed up the number of secondary infections over 30 days since the time of exposure to obtain the individual reproduction number, i.e. $R_0 = \sum_{\Delta_t} T_{\Delta_t}$.

Serial interval and generation time. We further assume that upon successful infection, it takes τ days for the virus to move within-host, reach infection site and produce the first infected cell. To calculate serial interval (time between the onset of symptoms of transmitter and secondarily infected person), we sample the incubation period for both transmitter and secondarily infected person from a gamma distribution with a shape described in the **Fig S4**.^{3,30} In cases in which symptom onset in the newly infected person precedes symptom onset in the transmitter, the serial interval is negative; otherwise, serial interval is non-negative. Similarly, we calculate generation time as the difference between the time of infection of transmitter and the time of infection of secondarily infected person.

Fitting procedure. To estimate the values of unknown parameters in cases of SARS-CoV-2, we performed a grid search comprehensively exploring a total of ~500,000 combinations of 5 parameters taking the following values,

- (i) $\tau \in [0.5, 1, 2, 3]$ days,
- (ii) $\alpha \in [0.01, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 2.0, 3.0, 4.0, 5.0, 10.0]$
- (iii) $\lambda \in [10^0, 10^{0.5}, 10^{1.0} \dots, 10^8]$
- (iv) $\theta \in [0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 2.0, 3.0, 4.0, 5.0, 10.0, 20.0, 50.0]$.
- (v) $\rho \in [0.0001, 0.001, 0.01, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 2.0, 5.0, 10.0, 20.0, 30.0, 40.0, 50.0, 75.0, 100, 200, 500]$.

The parameter sets of $(\lambda, \tau, \alpha, \theta)$ were simulated for 1000 infected individuals to determine how well each set generates the summary statistics of mean R_0 , mean SI and the R_0 histograms by following a procedure explained in **Fig S1** and below:

Step A:

1. Simulate viral load $V(t)$ of 1,000 simulated infected individuals using **Eq. 1**
2. For each combination of $(\lambda, \tau, \alpha, \theta, \rho)$
 - a. For each time step Δ_t
 - i. Compute $P_t[V(t); \lambda, \alpha]$
 - ii. Draw $\eta_{\Delta_t} \sim \Gamma\left(\frac{\theta}{\rho}, \rho\right) \Delta_t$
 - iii. Calculate $T_{\Delta_t} = \text{Ber}(P_t) P_t \eta_{\Delta_t}$
 - b. Calculate $R_0 = \sum_{\Delta_t} T_{\Delta_t}$
 - i. Check if calculated mean R_0 is in the range:^{3,31}
 - c. Calculate Serial Interval based on τ and incubation period
 - i. Check if calculated SI is in the range in:^{3,31,33}

Step B:

1. If the parameter combination in Step A satisfy the criteria, then
 - i. Compute RSS for the obtained R_0 and histogram from:^{4,6,34,36} [Ref]

We visually checked whether our dose-response curve matched the observed probability of positive virus culture.³⁷ We assumed that viral loads derived from positive culture³⁷ can be considered equivalent to viral loads in the within-host model if divided by a positive integer. We found this positive integer to be 25 (**Fig S5**).

We performed a global sensitivity analysis to identify which parameter variability accounted for fit to different components of the data. Only narrow ranges of λ permitted close fit to the mean of R_0 and distribution functions of individual R_0 (**Fig S9**), while a specific value for α was necessary to fit to mean serial interval and distribution functions of individual R_0 (**Fig**

S9). Only narrow ranges of θ permitted close fit to the mean of R_0 and distribution functions of individual R_0 (**Fig S10**), while a specific value for ρ was necessary to fit to distribution functions of individual R_0 (**Fig S10**).

To obtain $TD50 (\lambda_T)$ based on $ID50 (\lambda)$, we use the relation

$$\frac{1}{\left(\left(\frac{10^\lambda}{V}\right)^\alpha + 1\right)^2} = \frac{1}{\left(\frac{10^{\lambda_T}}{V}\right)^{\alpha_T} + 1} = 0.5$$

From solving the second half ($\frac{1}{\left(\frac{10^{\lambda_T}}{V}\right)^{\alpha_T} + 1} = 0.5$), we get

$$V = 10^{\lambda_T}$$

Substituting $V = 10^{\lambda_T}$ in the first-half, we have

$$\frac{1}{\left(\left(\frac{10^\lambda}{10^{\lambda_T}}\right)^\alpha + 1\right)^2} = 0.5$$

$$\text{Or, } \left(\left(\frac{10^\lambda}{10^{\lambda_T}}\right)^\alpha + 1\right)^2 = 2$$

$$\text{Or, } \left(\frac{10^\lambda}{10^{\lambda_T}}\right)^\alpha = \sqrt{2} - 1$$

$$\text{Or, } 10^{\lambda_T \alpha} = \frac{10^{\lambda \alpha}}{\sqrt{2} - 1}$$

$$\text{Or, } \lambda_T = \lambda + \frac{0.38}{\alpha}$$

References

- 1 <https://coronavirus.jhu.edu/map.html>.
- 2 He, X. et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med* **26**, 672-675, doi:10.1038/s41591-020-0869-5 (2020).
- 3 Ganyani, T. et al. Estimating the generation interval for coronavirus disease (COVID-19) based on symptom onset data, March 2020. *Euro Surveill* **25**, doi:10.2807/1560-7917.ES.2020.25.17.2000257 (2020).
- 4 Endo, A., Centre for the Mathematical Modelling of Infectious Diseases COVID-19 Working Group, Abbott, S., Kucharski, A. & Funk, S. Estimating the overdispersion in COVID-19 transmission using outbreak sizes outside China. *Wellcome Open Res* **5**, doi:10.12688/wellcomeopenres.15842.3 (2020).
- 5 Lloyd-Smith, J. O., Schreiber, S. J., Kopp, P. E. & Getz, W. M. Superspreading and the effect of individual variation on disease emergence. *Nature* **438**, 355-359, doi:10.1038/nature04153 (2005).
- 6 Bi, Q. et al. Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. *Lancet Infect Dis*, doi:10.1016/S1473-3099(20)30287-5 (2020).
- 7 L., H., P., D., I., C. & al., e. High SARS-CoV-2 Attack Rate Following Exposure at a Choir Practice — Skagit County, Washington, March 2020. . *MMWR Morb Mortal Wkly Rep* **69:606–610**. (2020).
- 8 Park, S. Y. et al. Coronavirus Disease Outbreak in Call Center, South Korea. *Emerg Infect Dis* **26**, 1666-1670, doi:10.3201/eid2608.201274 (2020).

608 9 Cowling, B. J., Fang, V. J., Riley, S., Malik Peiris, J. S. & Leung, G. M. Estimation of
609 the serial interval of influenza. *Epidemiology* **20**, 344-347,
610 doi:10.1097/EDE.0b013e31819d1092 (2009).

611 10 Brugger, J. & Althaus, C. L. Transmission of and susceptibility to seasonal influenza in
612 Switzerland from 2003 to 2015. *Epidemics* **30**, 100373,
613 doi:10.1016/j.epidem.2019.100373 (2020).

614 11 Qi, L. et al. Factors associated with the duration of viral shedding in adults with COVID-
615 19 outside of Wuhan, China: a retrospective cohort study. *Int J Infect Dis* **96**, 531-537,
616 doi:10.1016/j.ijid.2020.05.045 (2020).

617 12 Cao, B. et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-
618 19. *N Engl J Med*, doi:10.1056/NEJMoa2001282 (2020).

619 13 Pawelek, K. A. et al. Modeling within-host dynamics of influenza virus infection
620 including immune responses. *PLoS Comput Biol* **8**, e1002588,
621 doi:10.1371/journal.pcbi.1002588 (2012).

622 14 Memoli, M. J. et al. Validation of the wild-type influenza A human challenge model
623 H1N1pdMIST: an A(H1N1)pdm09 dose-finding investigational new drug study. *Clin*
624 *Infect Dis* **60**, 693-702, doi:10.1093/cid/ciu924 (2015).

625 15 Pebody, R. G. et al. Use of antiviral drugs to reduce household transmission of pandemic
626 (H1N1) 2009, United Kingdom. *Emerg Infect Dis* **17**, 990-999,
627 doi:10.3201/eid1706.101161 (2011).

628 16 Goldstein, E. et al. Oseltamivir for treatment and prevention of pandemic influenza
629 A/H1N1 virus infection in households, Milwaukee, 2009. *BMC Infect Dis* **10**, 211,
630 doi:10.1186/1471-2334-10-211 (2010).

631 17 Mayer, B. T. et al. Estimating the Risk of Human Herpesvirus 6 and Cytomegalovirus
632 Transmission to Ugandan Infants from Viral Shedding in Saliva by Household Contacts.
633 Viruses **12**, doi:10.3390/v12020171 (2020).

634 18 Boucoiran, I. et al. Nonprimary Maternal Cytomegalovirus Infection After Viral
635 Shedding in Infants. Pediatr Infect Dis J **37**, 627-631,
636 doi:10.1097/INF.0000000000001877 (2018).

637 19 Corey, L. et al. Once-daily valacyclovir to reduce the risk of transmission of genital
638 herpes. N Engl J Med **350**, 11-20, doi:10.1056/NEJMoa035144 (2004).

639 20 Rodger, A. J. et al. Risk of HIV transmission through condomless sex in serodifferent gay
640 couples with the HIV-positive partner taking suppressive antiretroviral therapy
641 (PARTNER): final results of a multicentre, prospective, observational study. Lancet **393**,
642 2428-2438, doi:10.1016/S0140-6736(19)30418-0 (2019).

643 21 Cohen, M. S. et al. Antiretroviral Therapy for the Prevention of HIV-1 Transmission. N
644 Engl J Med **375**, 830-839, doi:10.1056/NEJMoa1600693 (2016).

645 22 Schiffer, J. T., Johnston, C., Wald, A. & Corey, L. An Early Test-and-Treat Strategy for
646 Severe Acute Respiratory Syndrome Coronavirus 2. Open Forum Infect Dis **7**, ofaa232,
647 doi:10.1093/ofid/ofaa232 (2020).

648 23 Leung, N. H. L. et al. Respiratory virus shedding in exhaled breath and efficacy of face
649 masks. Nat Med **26**, 676-680, doi:10.1038/s41591-020-0843-2 (2020).

650 24 Goyal, A., Cardozo-Ojeda, E. & Schiffer, J. Potency and timing of antiviral therapy as
651 determinants of duration of SARS CoV-2 shedding and intensity of inflammatory
652 response. medRxiv **2020.04.10.20061325**, doi:10.1101/2020.04.10.20061325 (2020).

- 25 Wölfel, R. et al. Virological assessment of hospitalized patients with COVID-2019.
Nature **581**, 465-469, doi:10.1038/s41586-020-2196-x (2020).
- 26 Lescure, F. X. et al. Clinical and virological data of the first cases of COVID-19 in
Europe: a case series. Lancet Infect Dis **20**, 697-706, doi:10.1016/S1473-3099(20)30200-
0 (2020).
- 27 Young, B. E. et al. Epidemiologic Features and Clinical Course of Patients Infected With
SARS-CoV-2 in Singapore. JAMA, doi:10.1001/jama.2020.3204 (2020).
- 28 Kim, J. Y. et al. Viral Load Kinetics of SARS-CoV-2 Infection in First Two Patients in
Korea. J Korean Med Sci **35**, e86, doi:10.3346/jkms.2020.35.e86 (2020).
- 29 Brouwer, A. F., Weir, M. H., Eisenberg, M. C., Meza, R. & Eisenberg, J. N. S. Dose-
response relationships for environmentally mediated infectious disease transmission
models. PLoS Comput Biol **13**, e1005481, doi:10.1371/journal.pcbi.1005481 (2017).
- 30 Lauer, S. A. et al. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From
Publicly Reported Confirmed Cases: Estimation and Application. Ann Intern Med **172**,
577-582, doi:10.7326/M20-0504 (2020).
- 31 Du, Z. et al. Serial Interval of COVID-19 among Publicly Reported Confirmed Cases.
Emerg Infect Dis **26**, 1341-1343, doi:10.3201/eid2606.200357 (2020).
- 32 World Health Organization. Statement on the meeting of the International Health
Regulations (2005) Emergency Committee regarding the outbreak of novel coronavirus
(2019-nCoV). (2020).
- 33 Nishiura, H., Linton, N. M. & Akhmetzhanov, A. R. Serial interval of novel coronavirus
(COVID-19) infections. Int J Infect Dis **93**, 284-286, doi:10.1016/j.ijid.2020.02.060
(2020).

676 34 Zhang, Y., Li, Y., Wang, L., Li, M. & Zhou, X. Evaluating Transmission Heterogeneity
677 and Super-Spreading Event of COVID-19 in a Metropolis of China. *Int J Environ Res*
678 *Public Health* **17**, doi:10.3390/ijerph17103705 (2020).

679 35 Dillon, A. et al. Clustering and superspreading potential of severe acute respiratory
680 syndrome coronavirus 2 (SARS-CoV-2) infections in Hong Kong. PREPRINT (Version
681 1) available at Research Square, doi:10.21203/rs.3.rs-29548/v1 (2020).

682 36 Miller, D. et al. Full genome viral sequences inform patterns of SARS-CoV-2 spread into
683 and within Israel. medRxiv, 2020.2005.2021.20104521,
684 doi:10.1101/2020.05.21.20104521 (2020).

685 37 van Kampen, J. J. A. et al. Shedding of infectious virus in hospitalized patients with
686 coronavirus disease-2019 (COVID-19): duration and key determinants. medRxiv,
687 2020.2006.2008.20125310, doi:10.1101/2020.06.08.20125310 (2020).

688 38 Baccam, P., Beauchemin, C., Macken, C. A., Hayden, F. G. & Perelson, A. S. Kinetics of
689 influenza A virus infection in humans. *J Virol* **80**, 7590-7599, doi:10.1128/JVI.01623-05
690 (2006).

691 39 Lessler, J. et al. Outbreak of 2009 pandemic influenza A (H1N1) at a New York City
692 school. *N Engl J Med* **361**, 2628-2636, doi:10.1056/NEJMoa0906089 (2009).

693 40 Opatowski, L. et al. Transmission characteristics of the 2009 H1N1 influenza pandemic:
694 comparison of 8 Southern hemisphere countries. *PLoS Pathog* **7**, e1002225,
695 doi:10.1371/journal.ppat.1002225 (2011).

696 41 Cowling, B. J. et al. The effective reproduction number of pandemic influenza:
697 prospective estimation. *Epidemiology* **21**, 842-846, doi:10.1097/EDE.0b013e3181f20977
698 (2010).

699 42 Roberts, M. G. & Nishiura, H. Early estimation of the reproduction number in the
700 presence of imported cases: pandemic influenza H1N1-2009 in New Zealand. PLoS One
701 6, e17835, doi:10.1371/journal.pone.0017835 (2011).

702 43 Larremore, D. B. et al. Test sensitivity is secondary to frequency and turnaround time for
703 COVID-19 surveillance. medRxiv, 2020.2006.2022.20136309,
704 doi:10.1101/2020.06.22.20136309 (2020).

705 44 van Doremalen, N. et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared
706 with SARS-CoV-1. N Engl J Med, doi:10.1056/NEJMc2004973 (2020).

707 45 Widders, A., Broom, A. & Broom, J. SARS-CoV-2: The viral shedding vs infectivity
708 dilemma. Infect Dis Health, doi:10.1016/j.idh.2020.05.002 (2020).

709 46 Huang, C.-G. et al. Relative COVID-19 viral persistence and antibody kinetics. medRxiv,
710 2020.2007.2001.20143917, doi:10.1101/2020.07.01.20143917 (2020).
711




Fig 1. SARS-CoV-2 and influenza transmission model schematic. In the above cartoon, the transmitter has 2 exposure events at discrete timepoints resulting in 7 total exposure contacts and 3 secondary infections. Transmission is more likely at the first exposure event due to higher exposure viral load. To model this process, the timing of exposure events and number of exposed contacts is governed by a random draw from a gamma distribution which allows for heterogeneity in number of exposed contacts per day (**Fig S3**). Viral load is sampled at the precise time of each exposure event. Probability of transmission is identified based on the product of two dose curves (**Fig S2C, D**) which capture contagiousness (probability of viral passage to an exposure contact's airway) and infectiousness (probability of transmission given viral presence in the airway). Incubation period (**Fig S4**) of the transmitter and secondarily infected person is an input into each simulation and is depicted graphically. Individual R_0 is an output of each simulation and is defined as the number of secondary infections generated by an infected individual. Serial interval is an output of each simulated transmission and is depicted graphically.




Fig 2. SARS-CoV-2 transmission model fit. **A.** Simulated and actual frequency histograms of individual R_0 values, **B.** Simulated and actual cumulative distribution of individual R_0 values. **C.** Simulated and actual frequency histograms of individual serial intervals, **D.** Simulated and actual cumulative distribution of individual serial intervals. **E.** Frequency distribution of simulated generation times.




Fig 3. SARS-CoV-2 transmission probability as a function of shedding. **A.** Optimal infectious dose (ID) response curve (infection risk = P_i) and transmission dose (TD) response curve (transmission risk = $P_t * P_i$) curves for SARS-CoV-2. Transmission probability is a product of two probabilities, contagiousness and infectiousness (**Fig 1**). **B-D.** Three simulated viral shedding curves. Heat maps represent risk of transmission at each shedding timepoint given an exposed contact with an uninfected person at that time.




Fig 4. Conditional requirements for SARS-CoV-2 superspreading events. **A.** Heatmap demonstrating the maximum number of feasible secondary infections per day from a transmitter given an exposure viral load on log10 scale (x-axis) and number of exposed contacts per day (y-axis). The exposed contact network allows a maximum of 150 exposed contacts per day (black dotted line) which is sufficient for multiple transmissions from a single person per day. **B.** 10,000 simulated transmitters followed for 30 days. The white space is a parameter space with no transmissions. Each dot represents the number of secondary transmissions from a transmitter per day. Input variables are log10 SARS-CoV-2 on the start of that day and number of contact exposures per day for the transmitter. There are 1,154,001 total exposure contacts and 15,992 total infections. **C.** 10,000 simulated infections with percent of infections due to exposure viral load binned in intervals of 0.5 intervals on log10 scale (x-axis) and number of exposed contacts (y-axis).




Fig 5. Influenza transmission model fit. **A.** Simulated and actual frequency histograms of individual R_0 values, **B.** Simulated and actual cumulative distribution of individual R_0 values. **C.** Simulated and actual frequency histograms of individual serial intervals, **D.** Simulated and actual cumulative distribution of individual serial intervals. **E.** Frequency distribution of simulated generation times.




Fig 6. Influenza transmission probability as a function of shedding.
A. Optimal infectious dose (ID) response curve (infection risk = P_i) and transmission dose (TD) response curve (transmission risk = $P_i * P_t$) curves for influenza. Transmission probability is a product of two probabilities, contagiousness and infectiousness (**Fig 1**). **B-D.** Three simulated viral shedding curves. Heat maps represent risk of transmission at each shedding timepoint given an exposed contact with an uninfected person at that time.




Fig 7. Conditional requirements for influenza super spreading events. **A.** Heatmap demonstrating the maximum number of secondary infections per day feasible from a transmitter given an exposure viral load on log10 scale (x-axis) and number of exposed contacts per day (y-axis). The exposed contact network allows a maximum of 15 exposed contacts per day (black dotted line) which is not sufficient for more than 15 transmissions from a single person per day. **B.** 10,000 simulated transmitters followed for 30 days. The white space is a parameter space with no transmissions. Each dot represents the number of secondary transmissions from a transmitter per day. Input variables are log10 influenza TCID on the start of that day and number of contact exposures per day for the transmitter. There are 1,239,984 total exposure contacts and 11,141 total infections. **C.** 10,000 simulated infections with percent of infections due to exposure viral load binned in intervals of 0.5 intervals on log10 scale (x-axis) and number of exposed contacts (y-axis).





Fig 8. Differing transmission contact distributions, rather than viral kinetics explain SARS CoV-2 super spreader events. **A.** Simulated transmission risk dynamics for 10,000 infected persons with SARS-CoV-2 and influenza. Solid line is median transmission risk. Dark, dotted line is transmission risk of 75th percentile viral loads, and light dotted line is transmission risk of 95th percentile viral loads. **B.** Same as **A** but plotted as transmission risk since onset of symptoms. Highest transmission risk for SARS-CoV-2 is pre-symptoms and for influenza is post symptoms. **C.** Boxplots of duration of time spent above TD10, TD25, TD50, TD75 and TD90 for 10,000 simulated SARS-CoV-2 and influenza shedding episodes. TD10, TD25, TD50, TD75 and TD90 are viral loads at which transmission probability is 10%, 25%, 50%, 75% and 90% respectively. The midlines are median values, boxes are interquartile ranges (IQR), and datapoints are outliers. Superimposed probability distributions of: **D & E.** number of transmission contacts per day, **F.** individual R_0 , **G.** serial interval and **H.** generation time for influenza and SARS-CoV-2.

A) Calculating Mean R_0 , Mean Serial Interval and histogram of R_0



B) Finding parameter sets




Fig S1. Mathematical model workflow.




Fig S2. Mathematical model of SARS-CoV-2 transmission dynamics. **A.** Simulated viral load shedding tracings of possible transmitters. **B.** Simulated viral load shedding with imputed heterogeneity. **C.** Simulated infection dose (ID) response curves with variance in infectivity (ID50) and **D.** dose response slopes. **E.** Simulated transmission dose (TD) response curves with variance in infectivity (TD50) and **F.** dose response slopes. The TD response curve is a product of the infection and contagion dose response curves.




Fig S3. Stochastic simulations of exposed contact frequency for varying dispersion (ρ). The average number of exposed contacts is 4 per day in each example with imputed daily heterogeneity based on an elevated value of ρ from a gamma distribution $\sim \Gamma(4/\rho, \rho)$.




Fig S4. Gamma distribution functions of incubation periods. A. SARS-CoV-2 (mean 5.2 days, shape parameter=3.45 and rate=0.66) and **B.** influenza (mean 2 days, shape parameter=6.25 and scale parameter=0.32).




Fig S5. Mathematical model recapitulation of relationship between SARS-CoV-2 viral load and viral culture. In a clinical study, quantitative viral culture was ~25-fold lower than viral RNA measurement by PCR (<https://www.medrxiv.org/content/10.1101/2020.06.08.20125310v1>). We identify high similarity between observed viral RNA level divided by 25 and model predicted infectiousness shown here with the ID curve..




Fig S6. Impact of changes in contact network heterogeneity on individual R0, serial interval, and generation time. A. SARS-CoV-2, and B. influenza. Lowering exposed contact network heterogeneity to levels observed with influenza decreases SARS-CoV-2 individual R0 over-dispersion. Increasing exposed contact network heterogeneity to levels observed with SARS-CoV-2 increases influenza R0 over-dispersion. Neither change impacts observed serial interval or estimate generation time.




Fig S7. Potential impact of population physical distancing on SARS-CoV2 epidemiology. **A.** Mean reproductive number **B.** Percent transmitters of all infected people **C.** Percent super-spreaders (individual $R_0 > 5$) of all infected people **D.** Percent super spreaders of all transmitters. Transmitters are defined as infected people who generate at least one secondary infection.




Fig S8. Potential impact of enhanced physical distancing only within high exposure contact networks on SARS-CoV-2 epidemiology. Simulations assume limitation of exposed contacts only among daily exposures of more than 5, 10, 20 or 50 people. Mean reproductive number decreases below one with only marginal decreases in overall rate of exposure contacts when contacts are limited to fewer than 20 people.




Fig S9. Sensitivity analysis of transmission curve parameter for model fit to SARS-CoV-2 data. Effects of varying transmission curve slope (x-axis) and TD50 for infectiousness (y-axis) on fit to **A.** Mean R0, **B.** Mean serial interval, **C.** Cumulative distribution function of individual R0, and **D.** Sum of Errors in A, B and C.




Fig S10. Sensitivity analysis of contact network structure for model fit to SARS-CoV-2 data. Effects of dispersion parameter (x-axis) and average exposed contacts per day (y-axis) on fit to **A.** Mean R0, **B.** Mean serial interval, **C.** Cumulative distribution function of individual R0, and **D.** Sum of Errors in A, B and C.

Super-spreader definitions	SARS-CoV-2			Influenza		
	All infected people	All transmitters	Contribution of super-spreaders to all transmissions	All infected people	All transmitters	Contribution of super-spreaders to all transmissions
Individual $R_0 \geq 5$	~10%	~35%	~85%	~2%	~3%	~10%
Individual $R_0 \geq 10$	~6%	~25%	~70%	~0%	~0%	~0%
Individual $R_0 \geq 20$	~2.5%	~10%	~44%	~0%	~0%	~0%

Table 1: Prevalence of super-spreaders among transmitters, and contribution of super-spreading events to all SARS-CoV-2 and influenza transmissions. Estimates are from 10,000 simulations.

$\text{Log}_{10}\beta$ (virions ⁻¹ day ⁻¹)	δ (day ⁻¹ cells ^{-k})	k (-)	$\text{Log}_{10}\pi$ (log ₁₀ day ⁻¹)	m (day ⁻¹ cells ⁻¹)	$\text{Log}_{10}\omega$ (day ⁻¹ cells ⁻¹)
-7.23	3.13	0.08	2.59	3.21	-4.55
0.2	0.02	0.02	0.05	0.33	0.01

Table S1: Population parameter estimates for simulated SARS-CoV-2 viral shedding dynamics. Parameters are from (doi: <https://doi.org/10.1101/2020.04.10.20061325>).¹³ The top row is the fixed effects (mean) and the bottom row is the standard deviation of the random effects. We also fixed $r=10$, $\delta E=1/\text{day}$, $q=2.4 \times 10^{-5}/\text{day}$ and $c=15/\text{day}$.