



Volcano geodesy using InSAR in 2020: the past and next decades

Michael P. Poland¹ · Howard A. Zebker²

Received: 14 September 2021 / Accepted: 8 January 2022

This is a U.S. government work and not under copyright protection in the U.S.; foreign copyright protection may apply 2022

Abstract

The study of volcano deformation has grown significantly through the year 2020 since the development of interferometric synthetic aperture radar (InSAR) in the 1990s. This relatively new data source, which provides evidence of changes in subsurface magma storage and pressure without the need for ground-based equipment, has matured during the past decade. It now provides a means to address previously inaccessible questions and offers input to increasingly complex models of magmatic processes. Here, we review how technological advances in InSAR during 2010–2020 have facilitated our ability to monitor and interpret volcanic processes, primarily through rapid and accurate observations of the changing surfaces at active volcanoes worldwide. Specifically, we examine how current systems achieve excellent resolution in time and space, provide global coverage, and generate products that are easy to use by non-specialists—factors that have often limited the practical study of volcanoes using radar measurements. We also look to the future, offering our perspective about how advancements in technology and data management in the decade to come will increase the value and accessibility of InSAR applied to the geodetic study of volcanoes and monitoring of hazardous volcanic processes. New developments will include the launch of additional satellites by both public space agencies and private companies, as well as implementation of algorithms for exploiting the growing volumes of data. To meet their full potential, these efforts will require coordination between data users and data providers so that the relevant imagery is acquired, made available to volcanologists in a timely fashion, and utilized to assess and mitigate volcanic hazards.

Keywords Volcano geodesy · Deformation · InSAR · Interferometric synthetic aperture radar

Introduction

Within a year of the introduction of interferometric synthetic aperture radar (InSAR) as a tool for mapping earthquake displacements in 1994 (Massonnet et al. 1994), the first interferograms of volcanic deformation were published—of apparent deflation at Mount Etna, Italy (Massonnet et al. 1995). Although subsequent analysis found that a significant component of the apparent deflation signal was due

to atmospheric path delay (Delacourt et al. 1998; Massonnet and Feigl 1998; Beauducel et al. 2000), the implications were clear: InSAR offered a means of assessing volcano deformation on a previously unimagined scale. It was suddenly possible to map deformation at volcanoes globally, regardless of the status of ground-based monitoring. There were still complications, of course; results are poor in heavily vegetated areas unless a long radar wavelength (especially L-band, versus the shorter C- and X-bands) is used. In addition, snow and ice cover confound the method, sensitivity to along-track (~ north–south) displacements is low, and data are not collected consistently over all volcanoes of interest. Nevertheless, InSAR was a game changer in volcanology, and application of the technique grew rapidly (Pinel et al. 2014). The discovery of many deforming volcanoes demonstrated how the measurements could be used to identify sources of potential future eruptions and has focused research and monitoring efforts on sites that otherwise may have been overlooked (e.g., Wicks et al. 2002; Dzurisin 2003; Pritchard and Simons 2004; Pritchard

Editorial responsibility: J.H. Fink

This paper constitutes part of a topical collection:
Looking Backwards and Forwards in Volcanology: A
Collection of Perspectives on the Trajectory of a Science

✉ Michael P. Poland
mpoland@usgs.gov

¹ U.S. Geological Survey - Cascades Volcano Observatory,
Vancouver, WA, USA

² Stanford University, Stanford, CA, USA

et al. 2018). An early and striking example was the recognition of widespread and rapid deformation of volcanoes in the Galápagos archipelago (Amelung et al. 2000), which prompted the installation of ground-based monitoring and fueled research into the dynamics of magma accumulation, dike emplacement, and eruption by generations of volcanologists (e.g., Jónsson et al. 1999; Chadwick et al. 2006; Geist et al. 2006; Jónsson 2009; Bagnardi et al. 2013; Davis et al. 2021). Importantly, these discoveries were not limited to remote unmonitored volcanoes. The recognition of uplift near South Sister volcano, Oregon, demonstrated the presence of an active magma body in an area previously thought to be dormant, despite a record of past ground-based monitoring (Wicks et al. 2002; Dzurisin et al. 2006, 2009; Lisowski et al. 2021).

Further SAR sensor development and the availability of large volumes of SAR data led to new insights into volcanic processes, especially when InSAR was combined with other datasets, like seismology, gas geochemistry, and physical volcanology (Kilbride et al. 2016). Used in response to volcanic crises, SAR-derived products including deformation, amplitude, interferometric coherence, and topography have contributed to successful forecasts of volcanic eruptions and are credited with saving thousands of lives (Pallister et al. 2013). We build on previous reviews (e.g., Massonnet and Sigmundsson 2000; Zebker et al. 2000; Pinel et al. 2014) and explore the evolution of satellite SAR and InSAR as applied to volcanology during the 2010s and explore the potential of the field in the 2020s. Future advances in SAR instruments and processing methods are inevitable, but exploiting these improvements will require commensurate investments in data handling and availability; otherwise, we risk underutilizing a vital resource for assessing and mitigating volcanic hazards around the world.

2010–2020: the decade that was

During the 2010s, use of InSAR in volcanology experienced an important transition, evolving from a tool used for research to one that could also be applied for timely monitoring of volcanic unrest and eruption. This evolution was fueled by developments in satellite capabilities as well as data availability and analysis methods. The most significant advance was the free availability of frequent and regular SAR data to the community—for the first time, monitoring agencies and research groups alike could rely on a steady flow of observations supporting both scientific research and hazards assessment and mitigation.

Sensor developments

The beginning of the 2010s saw a changing of the guard in terms of satellite SAR. The 2000s were dominated by the C-band ASAR instrument on the ENVISAT satellite and the L-band PALSAR instrument on the ALOS-1 satellite. Both satellites failed in 2011, leaving the C-band RADARSAT-2, which was launched in 2007, as the remaining source of long-term satellite SAR data. In the late 2000s, two X-band systems, TerraSAR-X and COSMO-SkyMed, were launched, and throughout the 2010s, they provided SAR imagery with spatial resolutions of a few meters (and in some cases less than a meter!). These data were used to highlight very localized deformation that would not otherwise have been detected by ground instruments or moderate-resolution SAR (Richter et al. 2013; Salzer et al. 2014), but they are only available at a limited number of volcanoes. By the middle of the decade, the L-band ALOS-2 satellite and C-band Sentinel-1 satellites had been launched. The major volcanological innovations offered by Sentinel-1 were frequent and consistent repeat observations of nearly all volcanoes on Earth, albeit at moderate (~ 15 m) spatial resolution. ALOS-2 provided a new source of L-band imagery, which had been absent since the failure of ALOS-1, at a range of spatial scales but with inconsistent and infrequent repeat intervals. Nevertheless, the combination of targeted high-resolution X-band data, broad systematic C-band coverage, and sporadic L-band acquisitions provided a suite of information that facilitated new applications in volcanology. It was during this decade that InSAR transitioned from a research application to a tool that could be applied to timely monitoring of volcanic hazards. For example, InSAR data acquired during dike emplacement events in Iceland (2014) and Hawai'i (2018) provided deformation data that revealed the staggering scope of magmatic intrusions before they erupted, triggering the months-long Holuhraun (Sigmundsson et al. 2015) and Kīlauea lower East Rift Zone (Neal et al. 2019; Fig. 1) eruptions, respectively.

Data availability and analysis

Two community efforts opened access to satellite SAR for volcanology in the 2010s. First, the Geohazard Supersites and Natural Laboratory (GSNL) initiative established a network of volcano “supersites” (listed at <https://geo-gsnl.org/>), where SAR images were freely available and could be used in combination with other volcanological data (Dumont et al. 2018; Dzurisin et al. 2019). Second, the Committee on Earth Observing Satellites (CEOS) Volcano Pilot and Demonstrator projects made available satellite SAR data first from Latin America, and then also from southwest Asia and Africa, allowing for detailed examination of volcanoes




Fig. 1 Cumulative deformation over 2018 at Hawai'i's Kilauea Volcano, at its summit and along its East Rift Zone, as observed by Sentinel-1 InSAR (left). Labeled white dots give locations of GPS stations used for reference to time series. Comparison of GPS (red dashed lines) and InSAR time series (green and blue dots based on

maximum 30-day temporal separation) show centimeter-level accuracy of InSAR time series when a tropospheric propagation correction (green dots) is applied (right). Figure and method adapted from Zebker (2021)

throughout that region and laying the groundwork for a better understanding of how satellite data in general might be used to monitor volcanoes (Pritchard et al. 2018; Reath et al. 2019). Although these initiatives cover only a small proportion of active volcanoes worldwide, they illustrate the potential of SAR data in volcano monitoring and research when utilized more fully than is possible via projects driven by single investigators or in response to isolated episodes of unrest or eruptive activity. The growing availability of InSAR results from volcanoes also led to an explosion (pun intended) in knowledge of the widespread nature of volcano deformation. In the 20 years between 1997 and 2017, the number of volcanoes known to have experienced some sort of surface deformation expanded from 44 to over 220 (Biggs and Pritchard 2017). Analyses of this growing database provided evidence of the strong connection between deformation and eruption (Biggs et al. 2014), the varied styles of magmatic and non-magmatic volcano deformation (Ebmeier et al. 2018), and the recognition that deformation served as a pre-eruptive phenomenon far more often and for a longer period than any other satellite dataset (Furtney et al. 2018).

The sheer volume of satellite SAR data created a new challenge: how could these data best be utilized? Several parallel developments emerged to aid data management. Multi-temporal analyses capitalized on growing computational power to generate spatially and temporally dense time series of ground deformation using persistent scattering (Hooper et al. 2012; Spaans and Hooper 2016) and small baseline subset analysis (Berardino et al. 2002). Implementation of automated processing engines (Lazecký et al. 2020) and artificial intelligence approaches enabled users to automatically scan many thousands interferograms

for indications of anomalous surface displacements (e.g., Anantrasirichai et al. 2018, 2019; Gaddes et al. 2018, 2019), increasing the likelihood that deformation of a volcano is detected. Corrections for atmospheric conditions also became widely available, aiding interpretation of InSAR results that might otherwise be ambiguous, especially on tall and steep volcanic edifices (Yu et al. 2018; Yip et al. 2019). In addition, the advent of user-friendly InSAR products yielded measurements in common coordinates rather than the radar-specific slant range and Doppler system, allowing non-specialists to take advantage of the powerful observing capabilities without requiring deep training in the details of radar technology (Zebker 2017).

2020–2030: the decade to be

The 2020s will see new satellite missions that provide additional capacity for InSAR monitoring of active volcanism, and missions developed by fully private enterprises will materially contribute to SAR data and capabilities. Exploiting the growing trove of SAR data for volcano monitoring and research will require coordination and cooperation among data providers and users, as well as continued development of resources that take advantage of all aspects of SAR and InSAR.

Sensor developments

The most anticipated technological advance in satellite SAR in the 2020s will undoubtedly be offered by NISAR (NASA-ISRO Synthetic Aperture Radar), an L- and

S-band satellite mission currently hoped for launch in 2023 but more likely to fly in 2024. NISAR will acquire repeat L-band (24-cm wavelength) data every 12 days over nearly all subaerial volcanoes on Earth, offering a swath width of 240 km, resolution of 3–10 m, single/dual/quad polarimetry, and open data access (Rosen et al. 2017). Additional S-band (12-cm wavelength) data will be available on a select group of potentially active volcanoes. Both of these longer wavelengths are less sensitive to surface decorrelation and will reveal deformation unseen by shorter-wavelength radars. Although the acquisition of L-band satellite SAR is not new—SeaSat, JERS, ALOS-1, ALOS-2, and SAOCOM all did or do carry L-band SAR sensors—the reliable worldwide coverage and free availability set NISAR apart. In fact, the massive quantity of data itself will pose a challenge, since the mission will generate more data than any other NASA mission by far (Jasper and Xaypraseuth 2017).

Sentinel-1 continuity through follow-on missions from the European Space Agency (ESA) will extend the current capability of consistent, global, freely accessible C-band SAR for 30 years, far outpacing NISAR's planned 3 years. It is highly likely that NISAR will last much longer than 3 years, however, and NASA is already planning NISAR's successor through a Surface Deformation and Change study. ESA will undoubtedly improve the capabilities of each new generation of Sentinel satellites, providing an ever-increasing ability to observe volcanoes through all stages of the eruption cycle. For example, ESA's proposed Earth Explorer 10 Harmony mission would combine with Sentinel-1 to provide repeat topography and better resolution of along-track displacements—both critical datasets for volcano surveillance (Kubaneck et al. 2021; López-Dekker et al. 2021).

We also expect private development of SAR capabilities to flourish during 2020–2030, following the precedent of entrepreneurial space launches. As of 2021, two commercial companies, ICEYE and Capella, are acquiring X-band SAR data of selected targets through constellations of smallsats that will eventually number in the tens of spacecraft. Both companies have offered data in support of volcanic crises, specifically for the 2021 eruptions of Soufrière, St. Vincent, and Nyiragongo, Democratic Republic of Congo. These and similar private companies will eventually achieve daily or better SAR coverage of any location on Earth. Such dense spatiotemporal coverage will represent a new era in satellite SAR monitoring of volcanic activity, provided that data are acquired and available to the volcanology community in advance of a crisis and on a consistent, instead of ad hoc, basis. The rapid response by international space agencies to unrest and eruption at Merapi volcano, Indonesia, in 2020–2021, provides an example of the value of such a response. High-resolution SAR data that would not otherwise have been available were freely provided and used to

identify localized but high-rate displacements on the flanks of the volcano (Fig. 2). This information was communicated to local volcanologists monitoring the eruption and aided their response to the crisis, offering important constraints on potential hazards, especially given the lack of monitoring instruments located on the upper flanks of the volcano and the danger in accessing those areas during the crisis.

With these developments in satellites and sensors, volcanologists must not become overly reliant on any single dataset but rather continue to take advantage of all satellite SAR data. The mix of available wavelengths, repeat intervals, and spatial resolutions that is currently and will become available through the 2020s offers the best opportunity for detecting the variety of potential surface changes that might precede, accompany, and follow volcanic eruptions.

Data availability and analysis

Three capabilities are needed from satellite SAR systems to aid with responses to volcanic unrest and eruptions: (1) flexible and responsive acquisition planning, including regular background observations that are appropriate for each volcano, (2) low-latency data delivery, and (3) free access. Current methods for tasking and accessing SAR data are heterogeneous, and no system is completely optimized for providing the “right” data at the “right” time of the “right” place and for the “right” price. Near-global and consistent dense temporal repeatability from Sentinel-like systems greatly reduces the need to specifically target volcanoes in advance of an eruption, but these broad views are not appropriate in all situations. Small island volcanoes, for example, require higher-resolution views than are available from current satellite missions with global background coverage. Targeting flexibility is therefore critical for fine-resolution, limited-coverage satellite systems that can best image small volcanic systems with localized activity.

Especially during a volcanic crisis, rapid delivery of satellite data is vital. Real- and near-real-time satellite monitoring of thermal and ash emissions are a cornerstone of detecting eruptions, especially at remote volcanoes (Poland et al. 2020), but all SAR data have a latency. As of the early 2020s, only Sentinel-1 and COSMO-SkyMed offer delivery of the full complex dataset within a few hours of acquisition, at least in ideal situations. Many other satellite SAR systems can provide at least amplitude data within hours in cases where the International Charter: Space and Major Disasters is invoked. In addition, only Sentinel-1 data are 100% openly accessible. Other satellite SAR datasets must be accessed via special initiatives, like the GSNL or CEOS projects (Pritchard et al. 2018), or via research agreements, although such proposals do not typically allow for open-ended response-style data usage. For satellite SAR to be




Fig. 2 TerraSAR-X staring-spotlight interferogram of Merapi volcano, Indonesia, spanning November 16–27, 2020. Upper image shows phase change overlain on amplitude. Zoomed frames show

only phase change and highlight multiple areas of localized high-rate deformation on the volcano’s west flank that could not be easily detected by limited ground-based datasets

fully integrated into volcano monitoring, they must be freely available to monitoring agencies.

The capabilities outlined above require coordination among the volcanology community and collaboration with space agencies and private companies. Thus far, international work on volcano monitoring with satellite SAR has been done by individual investigators, or on a “best effort” basis without specific funding through the GSNL and CEOS Volcano Pilot/Demonstrator projects—efforts that are neither sustainable nor robust and that do not meet the needs of the global volcanology community and of populations at

risk from volcanic eruptions. A formal coordination office, akin to that organized by the polar science community, for managing satellite SAR datasets could serve as a liaison between space agencies (private and public) and the scientists/institutions tasked with volcano monitoring and crisis response. This point-of-contact approach, ideally with some redundancy spread around the world to ensure rapid responses across time zones, would facilitate communication, ensure that conflicting acquisitions are not scheduled, provide a mechanism for managing datasets and user licenses that respects investments of the contributing space

agencies and private companies, and ensure feedback and accountability between data providers and users. A coordination office could also encourage capacity building at volcano observatories and monitoring agencies that might not be able to independently process, analyze, or interpret satellite SAR data.

Of course, commensurate developments will continue in analysis methods. Automated processing and anomaly detection are already a reality, thanks to advances in computing power and artificial intelligence approaches, but these methods are not yet implemented at the scale of global volcano monitoring. We expect that within the decade of the 2020s, operational use of such techniques will be implemented.

Satellite SAR volcano monitoring must also extend beyond interferometry. It has long been known that SAR data from volcanoes offer more information than merely phase changes caused by surface deformation—for example, providing insights into surface characteristics and volcanic activity from coherence (Zebker et al. 1996; Dietterich et al. 2012), backscatter (Pallister et al. 2013; Arnold et al. 2018), and topographic change (Kubanek et al. 2021; López-Dekker et al. 2021). Capitalizing on all aspects of the richness of SAR data is the only way to fully exploit their utility for monitoring volcanic unrest and eruptions and to best serve societal needs for volcanic hazards assessment and mitigation.

Conclusions

The 2010–2020 decade has seen great strides in our ability to measure, monitor, and understand volcanoes. Space-based InSAR permits worldwide observation of entire volcanic systems at fine resolution and offers direct measurements of how a surface deforms under stresses due to migration of magma or subsurface pressure changes. InSAR has matured from a limited research tool to a capability that is applied to monitoring and mitigating volcanic hazards.

The availability of free and comprehensive crustal deformation observations, most notably from Sentinel-1, has led to an explosion in new analysis methods that characterize change at the centimeter level, promising a chance at timely forecasts and the development of hazard warning tools. New SAR systems on the drawing board for the next decade and in today's production facilities will provide long-term continuous coverage at several wavelengths, with commercial and specialized SAR systems providing a range of imaging options that will facilitate flexible responses to threatening volcanic activity. For these data to be useful in volcanology, however, they must be accessible to scientists in a timely fashion—a task that will require robust communication, coordination, and collaboration between monitoring

agencies, research institutes, space agencies, and private companies.

Acknowledgements We are grateful for the comments from, and discussions with, Matt Pritchard, Susi Ebmeier, Larry Mastin, and Andy Hooper, which substantially improved the quality of this manuscript. Our thanks to numerous colleagues who have contributed to the Super-sites, CEOS Volcano Pilot/Demonstrator, NISAR, and other projects throughout the years, and whose research, discoveries, and insights helped to shape our perspective. Copernicus Sentinel data from 2018 shown in Figure 1 were processed and made available by the European Space Agency. TerraSAR-X data shown in Figure 2 were made available via the CEOS Volcano Demonstrator, © DLR 2020. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Author contribution The authors contributed equally to this work.

Funding Work was supported by the US Geological Survey and Stanford University.

Data availability Not applicable.

Code availability Not applicable.

Declarations

Competing interests The authors declare no competing interests.

References

- Amelung F, Jónsson S, Zebker H, Segall P (2000) Widespread uplift and 'trapdoor' faulting on Galápagos volcanoes observed with radar interferometry. *Nature* 407:993–996. <https://doi.org/10.1038/35039604>
- Anantrasirichai N, Biggs J, Albino F, Hill P, Bull D (2018) Application of machine learning to classification of volcanic deformation in routinely generated InSAR data. *J Geophys Res* 123:6592–6606. <https://doi.org/10.1029/2018JB015911>
- Anantrasirichai N, Biggs J, Albino F, Bull D (2019) The application of convolutional neural networks to detect slow, sustained deformation in InSAR time series. *Geophys Res Lett* 46:11,850–11,858. <https://doi.org/10.1029/2019GL084993>
- Arnold DW, Biggs J, Wadge G, Mothes P (2018) Using satellite radar amplitude imaging for monitoring syn-eruptive changes in surface morphology at an ice-capped stratovolcano. *Rem Sens Environ* 209:480–488. <https://doi.org/10.1016/j.rse.2018.02.040>
- Bagnardi M, Amelung F, Poland MP (2013) A new model for the growth of basaltic shields based on deformation of Fernandina volcano, Galápagos Islands. *Earth Planet Sci Lett* 377–378:358–366. <https://doi.org/10.1016/j.epsl.2013.07.016>
- Beauducel F, Briole P, Froger J-L (2000) Volcano-wide fringes in ERS synthetic aperture radar interferograms of Etna (1992–1998): deformation or tropospheric effect? *J Geophys Res* 105:16,391–16,402. <https://doi.org/10.1029/2000JB900095>
- Berardino P, Fornaro G, Lanari R, Sansosti E (2002) A new algorithm for surface deformation monitoring based on small baseline differential SAR interferograms. *IEEE Trans Geosci Rem Sens* 40:2375–2383. <https://doi.org/10.1109/TGRS.2002.803792>

- Biggs J, Pritchard ME (2017) Global volcano monitoring: what does it mean when volcanoes deform? *Elements* 13:17–22. <https://doi.org/10.2113/gselements.13.1.17>
- Biggs J, Ebmeier SK, Aspinall WP, Lu Z, Pritchard ME, Sparks RSJ, Mather TA (2014) Global link between deformation and volcanic eruption quantified by satellite imagery. *Nat Comm* 5:3471. <https://doi.org/10.1038/ncomms4471>
- Chadwick WW, Geist DJ, Jónsson S, Poland M, Johnson DJ, Meertens CM (2006) A volcano bursting at the seams; inflation, faulting, and eruption at Sierra Negra Volcano, Galápagos. *Geology* 34:1025–1028. <https://doi.org/10.1130/G22826A.1>
- Davis T, Bagnardi M, Lundgren P, Rivalta E (2021) Extreme curvature of shallow magma pathways controlled by competing stresses: insights from the 2018 Sierra Negra eruption. *Geophys Res Lett* 48:e2021GL093038. <https://doi.org/10.1029/2021GL093038>
- Delacourt C, Briole P, Achache J (1998) Tropospheric corrections of SAR interferograms with strong topography. *Appl Etna Geophys Res Lett* 25:2849–2852. <https://doi.org/10.1029/98GL02112>
- Dietterich HR, Poland MP, Schmidt DA, Cashman KV, Sherrod DR, Espinosa AT (2012) Tracking lava flow emplacement on the east rift zone of Kilauea, Hawai'i, with synthetic aperture radar coherence. *Geochem Geophys Geosys* 13. <https://doi.org/10.1029/2011GC004016>
- Dumont S, Sigmundsson F, Parks MM, Drouin VJP, Pedersen GBM, Jónsdóttir I, Höskuldsson Á, Hooper A, Spaans K, Bagnardi M, Gudmundsson MT, Barsotti S, Jónsdóttir K, Högnadóttir T, Magnússon E, Hjartardóttir ÁR, Dürig T, Rossi C, Oddsson B (2018) Integration of SAR data into monitoring of the 2014–2015 Holuhraun eruption, Iceland: Contribution of the Icelandic Volcanoes Supersite and the FutureVolc projects. *Front Earth Sci* 6:231. <https://doi.org/10.3389/feart.2018.00231>
- Dzurisin D (2003) A comprehensive approach to monitoring volcano deformation as a window on the eruption cycle. *Rev Geophys* 41:10. <https://doi.org/10.1029/2001RG000107>
- Dzurisin D, Lisowski M, Wicks CW Jr, Poland MP, Endo ET (2006) Geodetic observations and modeling of magmatic inflation at the Three Sisters volcanic center, central Oregon Cascade Range, USA. *J Volcanol Geotherm Res* 150:35–54. <https://doi.org/10.1016/j.jvolgeores.2005.07.011>
- Dzurisin D, Lisowski M, Wicks CW (2009) Continuing inflation at Three Sisters volcanic center, central Oregon Cascade Range, USA, from GPS, leveling, and InSAR observations. *Bull Volcanol* 71:1091–1110. <https://doi.org/10.1007/s00445-009-0296-4>
- Dzurisin D, Lu Z, Poland MP, Wicks CW (2019) Space-Based Imaging Radar Studies of U.S. Volcanoes. *Front Earth Sci* 6:249. <https://doi.org/10.3389/feart.2018.00249>
- Ebmeier SK, Andrews BJ, Araya MC, Arnold DWD, Biggs J, Cooper C, Cottrell E, Furtney M, Hickey J, Jay J, Lloyd R, Parker AL, Pritchard ME, Robertson E, Venzke E, Williamson JL (2018) Synthesis of global satellite observations of magmatic and volcanic deformation: implications for volcano monitoring & the lateral extent of magmatic domains. *J Appl Volcanol* 7:2. <https://doi.org/10.1186/s13617-018-0071-3>
- Furtney MA, Pritchard ME, Biggs J, Carn SA, Ebmeier SK, Jay JA, McCormick Kilbride BT, Reath KA (2018) Synthesizing multi-sensor, multi-satellite, multi-decadal datasets for global volcano monitoring. *J Volcanol Geotherm Res* 365:38–56. <https://doi.org/10.1016/j.jvolgeores.2018.10.002>
- Gaddes ME, Hooper A, Bagnardi M, Inman H, Albino F (2018) Blind signal separation methods for InSAR: the potential to automatically detect and monitor signals of volcanic deformation. *J Geophys Res* 123:10,226–10,251. <https://doi.org/10.1029/2018JB0161210>
- Gaddes ME, Hooper A, Bagnardi M (2019) Using machine learning to automatically detect volcanic unrest in a time series of interferograms. *J Geophys Res* 124:12,304–12,322. <https://doi.org/10.1029/2019JB017519>
- Geist D, Chadwick W, Johnson D (2006) Results from new GPS and gravity monitoring networks at Fernandina and Sierra Negra Volcanoes, Galápagos, 2000–2002. *J Volcanol Geotherm Res* 150:79–97. <https://doi.org/10.1016/j.jvolgeores.2005.07.003>
- Hooper A, Bekaert D, Spaans K, Arkan M (2012) Recent advances in SAR interferometry time series analysis for measuring crustal deformation. *Tectonophysics* 514–517:1–13. <https://doi.org/10.1016/j.tecto.2011.10.013>
- Jasper LEZ, Xaypraseuth P (2017) Data production on past and future NASA missions. 2017 IEEE Aerospace Conference. <https://doi.org/10.1109/AERO.2017.7943918>
- Jónsson S (2009) Stress interaction between magma accumulation and trapdoor faulting on Sierra Negra volcano, Galápagos. *Tectonophysics* 471:36–44. <https://doi.org/10.1016/j.tecto.2008.08.005>
- Jónsson S, Zebker H, Cervelli P, Segall P, Garbeil H, Mouginiis-Mark P, Rowland S (1999) A shallow-dipping dike fed the 1995 flank eruption at Fernandina volcano, Galápagos, observed by satellite radar interferometry. *Geophys Res Lett* 26:1077–1080. <https://doi.org/10.1029/1999GL900108>
- Kilbride BM, Edmonds M, Biggs J (2016) Observing eruptions of gas-rich compressible magmas from space. *Nat Comm* 7:13744. <https://doi.org/10.1038/ncomms13744>
- Kubanek J, Poland MP, Biggs J (2021) Applications of bistatic radar to volcano topography—a review of ten years of TanDEM-X. *IEEE J Sel Top Appl Earth Obs Rem Sens* 14:3282–3302. <https://doi.org/10.1109/JSTARS.2021.3055653>
- Lazecký M, Spaans K, González PJ, Maghsoudi Y, Morishita Y, Albino F, Elliott J, Greenall N, Hatton E, Hooper A, Juncu D, McDougall A, Walters RJ, Watson CS, Weiss JR, Wright TJ (2020) LiCSAR: An automatic InSAR tool for measuring and monitoring tectonic and volcanic activity. *Rem Sens* 12:2430. <https://doi.org/10.3390/rs12152430>
- Lisowski M, McCaffrey R, Wicks CW, Dzurisin D (2021) Geodetic constraints on a 25-year magmatic inflation episode near Three Sisters, central Oregon. *J Geophys Res*
- López-Dekker P, Biggs J, Chapron B, Hooper A, Käab A, Masina S, Mouginiot J, Nardelli BB, Pasquero C, Prats-Iraola P, Rampal P, Stroeve J, Rommenet B (2021) The Harmony mission: End of phase-0 science overview. 2021 IEEE International Geoscience and Remote Sensing Symposium IGARSS, pp 7752–7755. <https://doi.org/10.1109/IGARSS47720.2021.9554896>
- Massonnet D, Feigl KL (1998) Radar interferometry and its applications to changes in the earth's surface. *Rev Geophys* 36:441–500. <https://doi.org/10.1029/97RG03139>
- Massonnet D, Feigl K, Rossi M, Adragna F (1994) Radar interferometric mapping of deformation in the year after the Landers earthquake. *Nature* 369:227–230. <https://doi.org/10.1038/369227a0>
- Massonnet D, Briole P, Arnaud A (1995) Deflation of Mount Etna monitored by spaceborne radar interferometry. *Nature* 375:567–570. <https://doi.org/10.1038/375567a0>
- Massonnet D, Sigmundsson F (2000) Remote sensing of volcano deformation by radar interferometry from various satellites. In: Mouginiis-Mark PJ, Crisp JA, Fink JH (eds) *Remote Sensing of Active Volcanism*. *Geophys Mon*. 116:207–221. <https://doi.org/10.1029/GM116p0207>
- Neal CA, Brantley SR, Antolik L, Babb JL, Burgess M, Calles K, Cappos M, Chang JC, Conway S, Desmither L, Dotray P, Elias T, Fukunaga P, Fuke S, Johanson IA, Kamibayashi K, Kauahikaua J, Lee RL, Pekalib S, Miklius A, Million W, Moniz CJ, Nadeau PA, Okubo P, Parcheta C, Patrick MR, Shiro B, Swanson DA, Tollett W, Trusdell F, Younger EF, Zoeller MH, Montgomery-Brown EK, Anderson KR, Poland MP, Ball JL, Bard J, Coombs M, Dietterich HR, Kern C, Thelen WA, Cervelli PF, Orr T, Houghton BF, Gansecki C, Hazlett R, Lundgren P, Dieffenbach AK, Lerner

- AH, Waite G, Kelly P, Clor L, Werner C, Mulliken K, Fisher G, Damby D (2019) The 2018 rift eruption and summit collapse of Kīlauea Volcano. *Science* 363:367–374. <https://doi.org/10.1126/science.aav7046>
- Pallister JS, Schneider DJ, Griswold JP, Keeler RH, Burton WC, Noyles C, Newhall CG, Ratdomopurbo A (2013) Merapi 2010 eruption—Chronology and extrusion rates monitored with satellite radar and used in eruption forecasting. *J Volcanol Geotherm Res* 261:144–152. <https://doi.org/10.1016/j.jvolgeores.2012.07.012>
- Pinel V, Poland MP, Hooper A (2014) Volcanology: lessons learned from Synthetic Aperture Radar imagery. *J Volcanol Geotherm Res* 289:81–113. <https://doi.org/10.1016/j.jvolgeores.2014.10.010>
- Poland MP, Lopez T, Wright R, Pavolonis MJ (2020) Forecasting, detecting, and tracking volcanic eruptions from space. *Remote Sens Earth Sys Sci* 3:55–94. <https://doi.org/10.1007/s41976-020-00034-x>
- Pritchard ME, Simons M (2004) Surveying volcanic arcs with satellite radar interferometry: the Central Andes, Kamchatka and beyond. *GSA Today* 14:4–11. [https://doi.org/10.1130/1052-5173\(2004\)014%3c4:SVAWSR%3e2.0.CO;2](https://doi.org/10.1130/1052-5173(2004)014%3c4:SVAWSR%3e2.0.CO;2)
- Pritchard ME, Biggs J, Wauthier C, Sansosti E, Arnold DWD, Delgado F, Ebmeier SK, Henderson ST, Stephens K, Cooper C, Wnuk K, Amelung F, Aguilar V, Mothes P, Macedo O, Lara LE, Poland MP, Zoffoli S (2018) Towards coordinated regional multi-satellite InSAR volcano observations: results from the Latin America pilot project. *J App Volcanol* 7:5. <https://doi.org/10.1186/s13617-018-0074-0>
- Reath K, Pritchard M, Poland M, Delgado F, Carn S, Coppola D, Andrews B, Ebmeier SK, Rumpf E, Henderson S, Baker S, Lundgren P, Wright R, Biggs J, Lopez T, Wauthier C, Moruzzi S, Alcott A, Wessels R, Griswold J, Ogburn S, Loughlin S, Meyer F, Vaughan G, Bagnardi M (2019) Thermal, deformation, and degassing remote sensing time-series (A.D. 2000–2017) at the 47 most active volcanoes in Latin America: implications for volcanic systems. *J Geophys Res* 124:195–218. <https://doi.org/10.1029/2018JB016199>
- Richter N, Poland MP, Lundgren PR (2013) TerraSAR-X interferometry reveals small-scale deformation associated with the summit eruption of Kīlauea Volcano, Hawai‘i. *Geophys Res Lett* 40:1279–1283. <https://doi.org/10.1002/grl.50286>
- Rosen PA, Kim Y, Kumar R, Misra T, Bhan R, Sagi VR (2017) Global persistent SAR sampling with the NASA-ISRO SAR (NISAR) mission. 2017 IEEE Radar Conference (RadarConf), 8–12 May, Seattle. <https://doi.org/10.1109/RADAR.2017.7944237>
- Salzer JT, Nikkhoo M, Walter TR, Sudhaus H, Reyes-Dávila G, Bretón M, Arambula R (2014) Satellite radar data reveal short-term pre-explosive displacements and a complex conduit system at Volcán de Colima, Mexico. *Front Earth Sci* 2:12. <https://doi.org/10.3389/feart.2014.00012>
- Sigmundsson F, Hooper A, Hreinsdóttir S, Vogfjörð KS, Ófeigsson BG, Heimisson ER, Dumont S, Parks M, Spaans K, Gudmundsson GB, Drouin V, Árnadóttir T, Jónsdóttir K, Gudmundsson MT, Högnadóttir T, Fridriksdóttir HM, Hensch M, Einarsson P, Magnússon E, Samsonov S, Brandsdóttir B, White RS, Ágústsdóttir T, Greenfield T, Green RG, Hjartardóttir ÁR, Pedersen R, Bennett RA, Geirsson H, La Femina PC, Björnsson H, Pálsson F, Sturkell E, Bean CJ, Möllhoff M, Braiden AK, Eibl EPS (2015) Segmented lateral dyke growth in a rifting event at Bárðarbunga volcanic system, Iceland. *Nature* 517:191–195. <https://doi.org/10.1038/nature14111>
- Spaans K, Hooper A (2016) InSAR processing for volcano monitoring and other near-real time applications. *J Geophys Res* 121:2947–2960. <https://doi.org/10.1002/2015JB012752>
- Wicks CW Jr, Dzurisin D, Ingebritsen S, Thatcher W, Lu Z, Iverson J (2002) Magmatic activity beneath the quiescent Three Sisters volcanic center, central Oregon Cascade Range, USA. *Geophys Res Lett* 29:26-1-26-4. <https://doi.org/10.1029/2001GL014205>
- Yip STH, Biggs J, Albino F (2019) Re-evaluating volcanic deformation using atmospheric corrections: Implications for the magmatic system of Agung volcano, Indonesia. *Geophys Res Lett* 46:13,704-13,711. <https://doi.org/10.1029/2019GL085233>
- Yu C, Li Z, Penna NT, Crippa P (2018) Generic atmospheric correction model for Interferometric Synthetic Aperture Radar observations. *J Geophys Res* 123:9202–9222. <https://doi.org/10.1029/2017JB015305>
- Zebker HA (2017) User-friendly InSAR data products: Fast and simple timeseries processing. *IEEE Geosci Rem Sens Lett* 14:2122–2126. <https://doi.org/10.1109/LGRS.2017.2753580>
- Zebker H (2021) Accuracy of a model-free algorithm for temporal InSAR tropospheric correction. *Rem Sens* 13:409. <https://doi.org/10.3390/rs13030409>
- Zebker HA, Rosen P, Hensley S, Mouginiis-Mark PJ (1996) Analysis of active lava flows on Kīlauea Volcano, Hawaii, using SIR-C radar correlation measurements. *Geology* 24:495–498. [https://doi.org/10.1130/0091-7613\(1996\)024%3c0495:AOALFO%3e2.3.CO;2](https://doi.org/10.1130/0091-7613(1996)024%3c0495:AOALFO%3e2.3.CO;2)
- Zebker HA, Amelung F, Jonsson S (2000) Remote sensing of volcano surface and internal processes using radar interferometry. In: Mouginiis-Mark PJ, Crisp JA, Fink JH (eds) *Remote Sensing of Active Volcanism*. *Geophys Mon.* 116:205. <https://doi.org/10.1029/GM116p0179>