

Rev08

1 **Type of Article:** *Short Communication*

2

3 **Title:** NVX-CoV2373 vaccine protects cynomolgus macaque upper and lower airways
4 against SARS-CoV-2 challenge

5

6 **Authors and Affiliations:**

7 Mimi Guebre-Xabier¹ [#], Nita Patel¹ [#], Jing-Hui Tian¹, Bin Zhou¹, Sonia Maciejewski¹,
8 Kristal Lam¹, Alyse D. Portnoff¹, Michael J. Massare¹, Matthew B. Frieman², Pedro A.
9 Piedra³, Larry Ellingsworth¹, Gregory Glenn¹, and Gale Smith¹ *

10

11 ¹Novavax, Inc. 21 Firstfield Road, Gaithersburg, MD, 20878, USA. [mguebre-](mailto:mguebre-xabier@Novavax.com)
12 [xabier@Novavax.com](mailto:npatel@novavax.com) (M.G.X.), [npatel@novavax.com](mailto:jhtian@Novavax.com) (N.P.), jhtian@Novavax.com
13 (J.H.T.), bzhou@Novavax.com (B.Z.), smaciejewski@Novavax.com (S.M.),
14 klam@Novavax.com (K.L.), aportnoff@Novavax.com (A.D.P.),
15 mmassare@Novavax.com (M.J.M.), lellingsworth@novavax.com (L.E.),
16 gglenn@Novavax.com (G.G.), gsmith@Novavax.com (G.S.)

17 ²University of Maryland, School of Medicine, 685 West Baltimore St, Baltimore, MD
18 21201, USA. mfrieman@som.umaryland.edu (M.B.F.)

19 ³Department of Molecular Virology and Microbiology, and Pediatrics, Baylor College of
20 Medicine, Houston, Texas. ppiedra@bcm.edu (P.A.P.)

21

22 ***Correspondence:** Dr. Gale Smith, Novavax, Inc., 21 Firstfield Road, Gaithersburg, MD
23 20878 USA. +1-240-268-2000. gsmith@Novavax.com (G.S.)

24 [#]MGX and NP each contributed equally as co-lead authors.

Rev08

25 **Highlights**

26 • Full-length SARS-CoV-2 prefusion spike with Matrix-M1™ (NVX-CoV2373) vaccine.

27 • Induced hACE2 receptor blocking and neutralizing antibodies in macaques.

28 • Vaccine protected against SARS-CoV-2 replication in the nose and lungs.

29 • Absence of pulmonary pathology in NVX-CoV2373 vaccinated macaques.

30

Rev08

31 ABSTRACT

32 There is an urgent need for a safe and protective vaccine to control the global spread of
33 SARS-CoV-2 and prevent COVID-19. Here, we report the immunogenicity and
34 protective efficacy of a SARS-CoV-2 subunit vaccine (NVX-CoV2373) produced from
35 the full-length SARS-CoV-2 spike (S) glycoprotein stabilized in the prefusion
36 conformation. Cynomolgus macaques (*Macaca fascicularis*) immunized with NVX-
37 CoV2373 and the saponin-based Matrix-M adjuvant induced anti-S antibody that was
38 neutralizing and blocked binding to the human angiotensin-converting enzyme 2
39 (hACE2) receptor. Following intranasal and intratracheal challenge with SARS-CoV-2,
40 immunized macaques were protected against upper and lower infection and pulmonary
41 disease. These results support ongoing phase 1/2 clinical studies of the safety and
42 immunogenicity of NVX-CoV2327 vaccine (NCT04368988).

43 **Key words:** SARS-CoV-2, COVID-19, spike glycoprotein, NVX-CoV2373 nanoparticles,
44 Matrix-M1 adjuvant, nonhuman primate

Rev08

45 **1. INTRODUCTION**

46 There is an urgent need for a safe and effective severe acute respiratory syndrome
47 coronavirus 2 (SARS-CoV-2) vaccine to prevent coronavirus disease 2019 (Covid-19).
48 We have developed a recombinant nanoparticle vaccine constructed from the
49 full-length, wild-type SARS-CoV-2 spike glycoprotein (GenBank gene sequence
50 MN908947, nucleotides 21563-25384) optimized for the baculovirus-*Spodoptera*
51 *frugiperda* (Sf9) insect cell expression system [1]. In mice and nonhuman primates
52 (NHP), NVX-CoV2373 with a Matrix-M1 saponin-based adjuvant induced high titer anti-
53 spike IgG that blocks binding to the hACE2 receptor, neutralize wild type virus, and
54 protects mice against SARS-CoV-2 challenge with no evidence of vaccine-associated
55 enhanced respiratory disease. NVX-CoV2373 vaccine also induces polyfunctional CD4⁺
56 T-cell responses of IFN- γ , IL-2, and TNF- α biased towards a Th1 phenotype, and
57 generates antigen-specific germinal center B cells in the spleen. Safety and
58 immunogenicity NVX-CoV2327 vaccine is currently under evaluation in humans
59 (NCT04368988) and primary safety and immunogenicity outcomes described [2]. We
60 evaluate in the current study NVX-CoV2373 vaccine immunogenicity, induction of
61 receptor blocking, and neutralizing antibodies compared to levels in human COVID-19
62 convalescent sera. And in a nonhuman primate challenge model, protection against
63 upper and lower virus replication and pulmonary disease.

64 **2. MATERIALS AND METHODS**

65 **2.1. Cell lines, virus, antibody reagents, and receptors.**

Rev08

66 Vero E6 cells (ATCC, CRL-1586) were maintained in Minimal Eagles Medium (MEM)
67 supplemented with 10% fetal bovine serum, 1% glutamine and 1% penicillin and
68 streptomycin. The SARS-CoV-2 (WA-1, 2020) isolated was obtained from the Center for
69 Disease Control and stock virus prepared in Vero E6 cells. Histidine-tagged hACE2
70 receptor was purchased from Sino Biologics (Beijing, CN). Rabbit anti-SARS-CoV spike
71 protein was purchased from Biodefense and Emerging Infections Research Resources
72 Repository (BEI Resources, Manassas, VA).

73 2.2. *Recombinant SARS-CoV-2 spike protein*

74 NVX-CoV2327 was codon optimized synthetically produced from the full-length S
75 glycoprotein gene sequence (GenBank MN908947 nucleotides 21563-25384) for
76 expression in *Spodoptera frugiperda* (Sf9) cells (GenScript Piscataway, NJ, USA) as
77 described [1]. Briefly, the S1/S2 furin cleavage site 682-RRAR-685 was modified 682-
78 QQAQ-685 and two proline substitutions introduced at positions K986P and V987P (2P)
79 to stabilize the full-length SARS-CoV-2 S [3].

80 2.3. *Animal ethics*

81 The in-life portion of the study was conducted at BIOQUAL, Inc (Rockville, MD).
82 Female and male, > 3 years old at study initiation, cynomolgus macaques (*Macaca*
83 *fascicularis*) were obtained from Primgen, Inc (Hines, IL) and maintained at BIOQUAL,
84 Inc for the entire in-life portion of the study. BIOQUAL, Inc. is accredited by AAALACCC
85 International. Animals were maintained and treated according to the Institutional
86 Biosafety Committee guidelines and the study was pre-approved by the Institutional
87 Animal Care and Use Committee (IACUC). The study was conducted in accordance

Rev08

88 with the National Institutes of Health Guide for Care and Use of Laboratory Animals
89 (NIH publication 8023, Revised 1978).

90 *2.4. Cynomolgus macaque immunization*

91 Cynomolgus macaques >3 years old (n=4/group) at study initiation received 5 or
92 25 µg NVX-CoV2327 with 50 µg Matrix-M1 (Novavax AB, Uppsala, Sweden)
93 administered in 500 µL in the thigh muscle in two doses spaced 21 days apart. A
94 separate group was immunized with a fractional dose (2.5 µg) NVX-CoV2373 with 25 µg
95 Matrix-M1 in two doses spaced 21 days apart and a placebo group received formulation
96 buffer. Serum was collected before immunization on Day 0, Day 21 just prior to the
97 second immunization, and Day 33.

98 *Anti-spike (S) IgG ELISA*

99 Anti-SARS-CoV-2 spike (S) protein IgG ELISA titers were measured as described
100 [1]. Anti-S IgG EC₅₀ titers were calculated by 4-parameter fitting using SoftMax Pro
101 6.5.1 GxP software. Individual animal anti-S IgG EC₅₀ titers, group geometric mean
102 titers (GMT) were plotted using GraphPad Prism 7.05 software.

103 *2.5. Inhibition of hACE2 receptor binding and neutralization*

104 Antibodies that block binding of hACE2 receptor to the S-protein and neutralize in a
105 cytopathic effect assay (CPE) in Vero E6 cells were measured as described previously
106 as the serum titer that blocks 100% CPE [1]. Serum antibody titer at 50% binding
107 inhibition (IC₅₀) of hACE2 to SARS-CoV-2 S protein was determined in the SoftMax Pro
108 program. Individual animal hACE2 receptor inhibiting titers, mean titers, and SEM were
109 plotted using GraphPad Prism 7.05 software. Neutralizing antibody titers were

Rev08

110 determined as the dilution of serum that inhibited 100% of CPE (CPE₁₀₀) at 3 days post
111 infection of Vero E6 cells in a 96 well plate format.

112 2.6. *SARS-CoV-2 challenge procedure*

113 The virus challenge study was done at BIOQUAL, Inc. within a BSL-3 containment
114 facility. SARS-CoV-2 generated from isolate 2019-nCoV/USA-WA1/2020 was received
115 from BEI Resources (NR-52281; lot # 70033175) and expanded in Vero E6 cells for
116 challenge stock generation. Animals were sedated and challenged with a targeted total
117 dose of 1.04×10^4 pfu SARS-CoV-2 by intranasal (IN) and intratracheal (IT) in a volume
118 of 0.25 mL each route. BAL and nasal swabs were collected 2- and 4-days post
119 challenge. Necropsy was performed 7 days following challenge and lung tissues
120 collected for histopathology.

121 2.7. *RNA subgenomic RT-PCR*

122 The subgenomic viral mRNA (sgRNA) was measured in macaque bronchoalveolar
123 lavage (BAL) and nasal swabs collected 2- and 4-days post challenge using RT-PCR as
124 described [4]. To generate a standard curve, the SARS-CoV-2 E gene sgRNA was
125 cloned into a pcDNA3.1 expression plasmid. The insert was transcribed using an
126 AmpliCap-Max T7 High Yield Message Maker Kit (Cellscript, Madison, WI) to obtain
127 RNA for standards. Prior to RT-PCR, samples collected from challenged animals or
128 standards were reverse-transcribed using Superscript III VILO (Invitrogen) according to
129 the manufacturer's instructions. A Taqman custom gene expression assay
130 (ThermoFisher Scientific, Rockville, MD) was designed using the sequences targeting
131 the E gene sgRNA. Reactions were carried out on a Quant Studio 6 and 7 Flex Real-

Rev08

132 Time PCR System (Applied Biosystems, Foster City, CA) according to the
133 manufacturer's specifications. Standard curves were used to calculate sgRNA in copies
134 per mL. The quantitative assay was sensitive to 50 copies per mL.

135 *2.8. Human COVID-19 convalescent serum*

136 Convalescent serum samples (n=32) were provided by Dr. Pedro A Piedra (Baylor
137 College of Medicine, Houston, TX, USA). Samples were collected from COVID-19
138 individuals 18-79 years of age 4-6 weeks after testing positive for SARS CoV-2.
139 Symptoms ranged from asymptomatic, mild to moderate symptoms, to severe
140 symptoms requiring hospitalization. Sera were analyzed for anti-SARS-CoV-2 S IgG,
141 hACE2 receptor inhibition, and virus neutralizing antibody titers.

142 *2.9. Histopathology*

143 Animals were euthanized 7-days following SARS-CoV-2 challenge (Day 42) and
144 lung tissues collected. Tissue were prepared for histologic examination by Experimental
145 Pathology Laboratories, Inc. (EPL, Sterling, VA). The lungs were fixed with 10%
146 formalin, paraffin embedded, and sections stained with hematoxylin and eosin (H&E) for
147 histological examination. Slides were examined for total inflammation, periarteriolar, and
148 peribronchiolar inflammation and epithelial cell denuding.

149 **3. RESULTS**

150 *3.1. Immunogenicity of NVX-CoV2373 in nonhuman primates compared to COVID-19*
151 *convalescent human sera*

152 Macaques immunized with the prime/boost regimen of 2.5, 5, and 25 µg NVX-
153 CoV2373 with 25 µg in the low and 50 µg Matrix-M1 adjuvant in the two higher doses

Rev08

154 induced anti-S IgG (EC₅₀) antibodies at Day 21 after a single dose (GMT = 7,810,
155 22,386 and 21,472, respectively). Two weeks following a booster immunization, anti-S
156 IgG EC₅₀ titers increased to GMT EC₅₀ = 163,036, 335,017 and 469,739, respectively
157 (**Figure 1A**). In contrast, SARS-CoV-2 anti-S antibody in convalescent human sera was
158 6.9- to 14.2-fold less with at GMT EC₅₀ of 23,614 (**Figure 1B**). And, hACE2 receptor
159 inhibition titers of 649, 1,410, and 1,320 in 2.5, 5, and 25 µg NVX-CoV2373 dose groups
160 respectively were 5.2 – 11.2-fold higher than in convalescent sera (**Figure 1C**). Finally,
161 SARS-CoV-2 GMT neutralization antibody titers of 17,920 - 23,040 CPE₁₀₀ in
162 immunized macaques, were 7.9 – 10.1-fold higher than in convalescent sera (**Figure**
163 **1D**).

164 *3.2. Viral load in nasal swabs and BAL*

165 To evaluate the potential efficacy of NVX-CoV2373 vaccine, macaques were
166 challenged with SARS-CoV-2 virus in upper and lower airways. Macaques in the
167 placebo group had 9,131 sgRNA copies/mL in the BAL at 2 days post challenge and
168 remained elevated at day 4 except for one animal. In contrast, immunized animals had
169 no detectable sgRNA in BAL fluid other than one animal in the low dose group at day 2
170 which cleared replicating virus RNA by day 4 (**Figure 1E**). Half of the controls had ~4
171 log10 of virus sgRNA copies in nasal swabs and in contrast, no detectable sgRNA was
172 in the nose of NVX-CoV2373 vaccinated animals (**Figure 1F**).

173 *3.3. Lung pathology*

174 Lung tissues were collected from all animals at 7 days post challenge and sections
175 examined for pathologic changes within the upper and lower airways. Consistent with

Rev08

176 previous reports of SARS-CoV-2 infection in rhesus macaques [5-10] placebo control
177 animals had moderate to severe inflammation that involved the mucosa of the bronchi,
178 perivascular mononuclear infiltrate with mixed infiltrates of macrophages and
179 neutrophils within the alveoli. In contrast, there was little, or no inflammation observed in
180 the lungs of macaques immunized with NVX-CoV2373 vaccine 7 days post challenge
181 (**Figure 2**). These findings were consistent the absence of sgRNA in BAL fluids and
182 nasal swabs of vaccinated animals by day 4 post challenge (**Figure 1E and 1F**).

183 **Discussion**

184 Here, we report the immunogenicity and the protective efficacy of a prefusion,
185 stabilized, full-length SARS-CoV-2 S vaccine (NVX-CoV2373) in the cynomolgus
186 macaque model permissive to infection [11]. Prime and booster immunization with NVX-
187 CoV2373 vaccine with Matrix1-M adjuvant induced high levels of anti-S IgG and
188 antibodies that blocked SARS-CoV-2 spike protein binding to the hACE2 receptor and
189 neutralized the virus. Importantly, vaccinated nonhuman primates had little or no
190 detectable replicating virus (sgRNA) in either upper or lower respiratory tracks. These
191 results demonstrate a potential of NVX-CoV2373 vaccine to protect the lower
192 respiratory track against pulmonary disease and upper respiratory track against virus
193 replication thus helping to establish herd immunity and to halt the COVID-19 pandemic
194 and its devastating global impact.

195

Rev08

196 **ENDNOTES**

197 **Acknowledgments**

198 Funding for certain studies was provided by the Coalition for Epidemic Preparedness
199 Innovations (CEPI), PO Box 123, Torshov, 0412 Oslo, Norway.

200 **Author contributions**

201 MGX, GS, GG, JHT, ADP, MJM, MBF and LE contributed to conceptualization of
202 experiments, generation of data and analysis, and interpretation of the results. NP, JHT,
203 BZ, and SM performed experiments. MGX, NP and KL coordinated projects. GS, GG,
204 MBF, PAP, MGX, NP and LE contributed to drafting and making critical revisions with
205 the help of others.

206 **Competing interests**

207 Authors MGX, NP, JHT, BZ, SM, KL, ADP, MJM, GG, GS and LE are current or past
208 employees of Novavax, Inc., a for-profit organization, and these authors own stock or
209 hold stock options. These interests do not alter the authors adherence to policies on
210 sharing data and materials. MBF and PAP declare no competing interests.

211

Rev08

212 REFERENCES

213 1. Jing-Hui Tian, Nita Patel, Robert Haupt, Haixia Zhou, Stuart Weston, Holly
214 Hammond, et al. SARS-CoV-2 spike glycoprotein vaccine candidate NVX-CoV2373
215 elicits immunogenicity in baboons and protection in mice. bioRxiv 2020; doi:
216 <https://doi.org/10.1101/2020.06.29.178509>

217 2. Cheryl Keech, Gary Albert, Patricia Reed, Susan Neal, Joyce S. Plested, Mingzhu
218 Zhu et al. First-in-Human Trial of a SARS CoV 2 Recombinant Spike Protein
219 Nanoparticle Vaccine. medRxiv 2020; doi:
220 <https://doi.org/10.1101/2020.08.05.20168435>

221 3. Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, Graham BS, et
222 al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. Science
223 2020;367:1260-3. <https://doi:10.1126/science.abb2507>.

224 4. Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Müller MA, et al.
225 Virological assessment of hospitalized patients with COVID-2019. Nature
226 2020;581:465-9. <https://doi:10.1038/s41586-020-2196-x>.

227 5. Shan C, Yao YF, Yang XL, Zhou YW, Gao G, Peng Y, et al. Infection with novel
228 coronavirus (SARS-CoV-2) causes pneumonia in Rhesus macaques. Cell Res
229 2020;30:670-7. <https://doi:10.1038/s41422-020-0364-z>.

230 6. Chandrashekhar A, Liu J, Martinot AJ, McMahan K, Mercado NB, Peter L, et al.
231 SARS-CoV-2 infection protects against rechallenge in rhesus macaques. Science
232 2020;369:812-7. <https://doi:10.1126/science.abc4776>.

Rev08

233 7. Gao Q, Bao L, Mao H, Wang L, Xu K, Yang M, et al. Development of an inactivated
234 vaccine candidate for SARS-CoV-2. *Science* 2020;369:77-81.
235 <https://doi:10.1126/science.abc1932>.

236 8. Neeltje van Doremalen, Teresa Lambe, Alexandra Spencer, Sandra Belij-
237 Rammerstorfer, Jyothi N. Purushotham, Julia R. Port, et al. ChAdOx1 nCoV-19
238 vaccination prevents SARS-CoV-2 pneumonia in rhesus macaques. *bioRxiv*. 2020;
239 <https://doi:10.1101/2020.05.13.093195>.

240 9. Yu J, Tostanoski LH, Peter L, Mercado NB, McMahan K, Mahrokhan SH, et al. DNA
241 vaccine protection against SARS-CoV-2 in rhesus macaques. *Science*
242 2020;369:806-11. <https://doi:10.1126/science.abc6284> .

243 10. Kizzmekia S. Corbett, Barbara Flynn, Kathryn E. Foulds, Joseph R. Francica,
244 Seyhan Boyoglu-Barnum, Anne P. Werner, et al. Evaluation of the mRNA-1273
245 Vaccine against SARS-CoV-2 in Nonhuman Primates. *N Engl J Med* 2020;
246 <https://doi:10.1056/NEJMoa2024671>.


247 11. Rockx B, Kuiken T, Herfst S, Bestebroer T, Lamers MM, Oude Munnink BB, et al.
248 Comparative pathogenesis of COVID-19, MERS, and SARS in a nonhuman primate
249 model. *Science* 2020;368:1012-5. <https://doi:10.1126/science.abb7314>.

250

251

Rev08

252 **Figure 1.**



253

Rev08

254 **Figure 1.** Immunogenicity of NVX-CoV2373 vaccine in cynomolgus macaques. **(A)**

255 Groups of cynomolgus macaques (n = 4 per arm) were immunized weeks 0 and 3 with

256 2.5 μ g NVX-CoV2373 with 25 μ g Matrix-M1 or 5 μ g or 25 μ g NVX-CoV2373 with 50 μ g

257 Matrix-M1. Anti-spike EC₅₀ IgG titers were measured weeks 0, 1, 3, and 5. Lines

258 indicate anti-spike IgG titers for individual macaques in each group. **(B)** Anti-spike EC₅₀

259 IgG serum titers week 5 in NVX-CoV2373 immunized NHP compared to anti-S EC₅₀ IgG

260 titers in convalescent human sera. **(C)** ACE2 inhibition IC₅₀ serum titers week 5 NVX-

261 CoV2373 immunized macaques compared to ACE2 inhibition titers in convalescent

262 human sera, **(D)** Neutralization CPE₁₀₀ titers against wild type SARS-CoV-2 virus week

263 5 NVX-CoV2373 immunized macaques compared to neutralization CPE₁₀₀ titers in

264 convalescent human sera, **(E)** Subgenomic RNA copies in BAL fluid days 2 and 4 post

265 challenge SARS-CoV-2 virus in placebo and NVX-CoV2373 immunized macaques. **(F)**

266 Subgenomic RNA copies in nasal swab samples days 2 and 4 post challenge with

267 SARS-CoV-2 virus in placebo and NVX-CoV2373 immunized macaques. Dashed

268 horizontal line indicates the limit of detection (LOD). ConV: Convalescent serum. BAL:

269 bronchoalveolar lavage.

270

Rev08

271 **Figure 2.**




Figure 2. Representative histopathology of lungs from NVX-CoV2373 vaccinated cynomolgus macaques challenged with SARS-CoV-2 (WA1 strain). Histological findings of representative placebo treated animals included eosinophils expanding the mucosa of the bronchi, perivascular mononuclear infiltrate, with mixed macrophages and neutrophils within the alveoli at 7 days post infection. In the 5 µg dose group, one animal had mild to moderate perivascular infiltrate while other animals had no remarkable findings. There were no remarkable histological changes in the bronchial, vascular or alveoli in animals vaccinated with 50 µg NVX-CoV2373. There was no evidence of exacerbated lung inflammation in NVX-CoV2373 immunized animals.

272