

Macro-scale patterns in functional connectivity associated with ongoing thought patterns and dispositional traits

Samyogita Hardikar ¹, Brontë McKeown ^{2,15}, H. Lina Schaare ^{3, 4}, Ting Xu ⁵, Mark Edgar Lauckner ⁶, Sofie L. Valk ^{3, 4, 7}, Daniel S. Margulies ⁸, Adam Turnbull ^{9 10}, Boris Bernhardt ¹¹, Reinder Vos de Wael ¹¹, Arno Villringer ^{1, 12-14}, Jonathan Smallwood ¹⁵.

¹ Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

² Department of Psychology, University of York, York, YO10 5DD, United Kingdom

³ Otto Hahn Group Cognitive Neurogenetics, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

⁴ Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre Jülich, Jülich, Germany

⁵ Center for the Developing Brain, Child Mind Institute, New York, New York.

⁶ Max Planck Research Group: Adaptive Memory, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

⁷ Institute of Systems Neuroscience, Heinrich Heine University Düsseldorf, Düsseldorf, Germany

⁸ Frontlab, Institut du Cerveau et de la Moelle épinière, UPMC UMRS 1127, Inserm U 1127, CNRS UMR 7225 Paris, France

⁹ Department of Psychiatry and Behavioral Sciences, Stanford University, CA 94305, USA

¹⁰ Department of Brain and Cognitive Sciences, University of Rochester, NY 14627, USA

¹¹ McConnell Brain Imaging Centre, Montreal Neurological Institute and Hospital, McGill University, Montreal, Canada

¹² Day Clinic of Cognitive Neurology, Universitätsklinikum Leipzig, Leipzig, Germany.

¹³ MindBrainBody Institute, Berlin School of Mind and Brain, Humboldt-Universität zu Berlin, 10099 Berlin, Germany;

¹⁴ Center for Stroke Research Berlin (CSB), Charité - Universitätsmedizin Berlin, 10117 Berlin, Germany

¹⁵ Department of Psychology, Queens University, Kingston, Ontario, Canada.

Abstract

Complex macro-scale patterns of brain activity that emerge during periods of wakeful rest provide insight into the organisation of neural function, how these differentiate individuals based on their traits, and the neural basis of different types of self-generated thoughts. Although brain activity during wakeful rest is valuable for understanding important features of human cognition, its unconstrained nature makes it difficult to disentangle neural features related to personality traits from those related to the thoughts occurring at rest. Our study builds on recent perspectives from work on ongoing conscious thought that highlight the interactions between three brain networks - ventral and dorsal attention networks, as well as the default mode network. We combined measures of personality with state-of-the-art indices of ongoing thoughts at rest and brain imaging analysis, and explored whether this 'tri-partite' view can provide a framework within which to understand the contribution of states and traits to observed patterns of neural activity at rest. To capture macro-scale relationships between different brain systems, we calculated cortical gradients to describe brain organisation in a low dimensional space. Our analysis established that for more introverted individuals, regions of the ventral attention network were functionally more aligned to regions of the somatomotor system and the default mode network. At the same time, a pattern of detailed self-generated thought was associated with a decoupling of regions of dorsal attention from regions in the default mode network. Our study, therefore, establishes that interactions between attention systems and the default mode network are important influences on ongoing thought at rest and highlights the value of integrating contemporary perspectives on conscious experience when understanding patterns of brain activity at rest.

Introduction

Macro-scale patterns of brain activity at rest have the potential for elucidating the organisation of neural function, different types of psychiatric conditions (Cao et al., 2018; Koban et al., 2021), developmental changes including those during adolescence and old age (Dosenbach et al., 2010; Cui et al., 2020; Gratton et al., 2020; Wen et al., 2020), neurological disorders (Zhang et al., 2021) and are important for revealing the neural basis behind the landscape of self-generated experiences (Karapanagiotidis et al., 2019; McKeown et al., 2020; Kucyi et al., 2021). However, compared to controlled experimental conditions, interpreting neural activity recorded during resting-state functional magnetic resonance imaging (rs-fMRI) is challenging, partly because both trait-level aspects of the individual, and, the inherently complex and dynamic nature of ongoing experience at rest are both contributory factors to the observed brain activity (Smallwood et al., 2021b). It has recently been suggested that the meaning of different patterns of neural activity can be usefully constrained by pairing imaging data at rest with additional measures (Finn, 2021), for example by accounting for the patterns of thoughts individuals experience at rest (Karapanagiotidis et al., 2020, 2021; McKeown et al., 2020; Gonzalez-Castillo et al., 2021; Kucyi et al., 2021) or features of their personality (Hsu et al., 2018). While this methodological perspective is invaluable, we currently lack a theoretical framework within which to understand the brain-cognition relationships that these observations will establish. To address this gap in the literature, our study explores whether contemporary theories of the neural basis of ongoing conscious thought can provide a framework within which to interpret associations between macro-scale patterns of neural activity observed at rest, and measures of traits and ongoing experience.

Emerging views of how the brain supports patterns of ongoing conscious thought highlight interactions between three large scale networks (Menon, 2011; Huang et al., 2021; Smallwood et al., 2021b): the ventral attention network (VAN), the dorsal attention network (DAN) and the

default mode network (DMN)[⊕]. According to these ‘tri-partite’ network accounts, key hubs of the ventral attention network, such as the anterior insula and dorso-lateral prefrontal cortex, help gate access to conscious experience, influencing interactions between the DAN, which is more important for external mental content (Corbetta and Shulman, 2002), and the DMN which is important when states rely more on internal representations (Smallwood et al., 2021a). For example, Huang and colleagues established that activity levels in the anterior insula determine whether stimuli presented at perceptual threshold are consciously perceived (Huang et al., 2021), and this gating of external input emerged as a consequence of changes in the normal interactions between the DAN and DMN. They also found that disruptions to activity in the insula through anaesthesia resulted in reductions in self-generated mental imagery. Coming from a different perspective, Turnbull and colleagues (Turnbull et al., 2019b) used experience sampling during task performance to link patterns of neural activity to different features of ongoing thought. For example, they found activity in the dorsolateral prefrontal cortex (a member of the VAN according to a parcellation by Yeo, Krienen and colleagues, (Yeo et al., 2011)) was correlated with apparently contradictory patterns of ongoing thought – (i) self-generated episodic thoughts during periods of low demands, and (ii) patterns of detailed task focus when individuals were engaged in demanding external task. In the same study, neural activity within regions of the dorsal parietal cortex within the DAN, was exclusively reduced when participants engaged in self-generated thinking, highlighting a parallel neglect of external input seen by Huang and colleagues. Finally, in a second study, Turnbull and colleagues found that at rest, trait variance in the ability to focus on self-generated experience in laboratory situations with lower task demands is associated with decoupling of signals arising from the DAN and DMN (Turnbull et al., 2019). Summarising this emerging evidence, studies focused on understanding ongoing thought patterns from different perspectives converge on the view that regions of the VAN may be important for gating

[⊕] In this paper we refer to the networks using the taxonomy provided by Yeo, Krienen and colleagues (Yeo et al., 2011)

conscious access to different types of content by biasing interactions between the DAN and the DMN (Huang et al., 2021; Smallwood et al., 2021b).

Our current study explored whether this “tri-partite network” view of ongoing conscious thought derived from studies focused on understanding conscious experience, provides a useful organising framework for understanding the relation between observed brain activity at rest and patterns of cognition/ personality traits. We examined links between both (i) trait descriptions of individuals and (ii) patterns of ongoing thought. Our sample was a cohort of 144 participants who underwent a one-hour resting-state scan. Across this one-hour period, participants were interrupted on four occasions to answer a set of questions about their experiences at rest using multidimensional experience sampling (MDES), similar to a number of prior neuroimaging studies (Smallwood et al., 2016; Poerio et al., 2017; Wang et al., 2018). During a different session, the same participants also completed a battery of measures assessing features of their personality (such as the Big five (Costa and McCrae, 2008)) as well as subclinical/ psychiatric traits such as trait anxiety and depression (Zigmond and Snaith, 1983). Since our research question depends on understanding the hypothesised relationship between large scale networks (VAN, DMN and DAN), we used the Brainspace toolbox (Vos de Wael et al., 2020) to provide whole brain low dimensional representations of functional brain organisation, generating maps which represent the similarities and differences in the activity within different systems. We used R version 4.2.0 (R Core Team, 2021) to produce low dimensional representations of both traits and thoughts using principal component analysis (PCA). Using these two sets of data, we performed multiple regression to identify how brain network organisation varies with traits and states. In these analyses, the low dimensional representations of brain organisation were the dependent measures, and the components of traits and states were explanatory variables.

Methods

Data

The dataset used here is part of the MPI-Leipzig Mind-Brain-Body (MPILMBB) database (Mendes et al., 2019). The complete dataset consists of a battery of self-reported personality measures, measures of spontaneous thought, task data, and structural and resting-state functional MRI (one hour, divided into four adjacent 15-min sessions) from participants between 20 and 75 years of age. A detailed description of the participants, measures, and data acquisition protocol has been previously published along with the dataset (Mendes et al., 2019).

Participants

We limited our investigation to personality and thought self-reports, and rs-fMRI from participants under 50 years of age, who had complete data from at least three resting-state scans. The resulting sample included 144 participants (74 men, mean age= 26.77 years, SD= 4.03; 70 women, mean age = 26.93 years, SD = 5.55).

Resting state fMRI

The current sample includes fully pre-processed rs-fMRI data from 144 participants (four scans from 135 participants, and three scans from nine participants whose data were missing or incomplete). The rs-fMRI was acquired in axial orientation using T2*-weighted gradient-echo echo planar imaging (GE-EPI) sensitive to blood oxygen level-dependent (BOLD) contrast. Sequences were identical across the four runs, with the exception of alternating slice orientation and phase-encoding direction, to vary the spatial distribution of distortions and signal loss. Imaging and pre-processing parameters are described in detail in Mendes et al (Mendes et al., 2019).

Personality measures

To provide a broad description of individual traits we included data from the following 21 questionnaires:

Table 1. List of personality/ dispositional trait questionnaires

Abbreviation	Behavioural Measure
ACS	Attention Control Scale (Derryberry and Reed, 2002)
ASR	Adult Self Report (Achenbach and Rescorla, 2003)
BDI-II	Beck Depression Inventory -II (Beck et al., 1993)
BIS/BAS	Behavioural Inhibition and Approach System (Carver and White, 1994)
BP	Boredom Proneness Scale (Farmer and Sundberg, 1986)
ESS	Epworth Sleepiness Scale (Johns, 1991)
Gold-MSI	Goldsmiths Musical Sophistication Index (Müllensiefen et al., 2014)
HADS	Hospital Anxiety and Depression Scale (Zigmond and Snaith, 1983)
IAT	Internet Addiction Test (Young, 1998)
IMIS	Involuntary Musical Imagery Scale (Floridou et al., 2015)
MMI	Multimedia Multitasking Index (Ophir et al., 2009)
NEO PI-R	NEO Personality Inventory-Revised (Costa and McCrae, 2008)
PSSI	Personality Style and Disorder Inventory (Kuhl and Kazén, 2009)
SCS	Brief Self-Control Scale(Tangney et al., 2004)
SDS	Social Desirability Scale-17 (Crowne and Marlowe, 1960)
SES	Self-Esteem Scale (O'Malley and Bachman, 1979)
SD3	Short Dark Triad (Jones and Paulhus, 2014)
S-D-MW	Spontaneous and Deliberate Mind-Wandering (Carriere et al., 2013; Golchert et al.,
STAXI	State-Trait Anger Expression Inventory
TPS	Tuckman Procrastination Scale (Tuckman, 2016)
UPPS-P	UPPS-P Impulsive Behaviour Scale (Lynam et al., 2006; Schmidt et al., 2008)

Multi-dimensional experience sampling

Participants underwent a short MDES survey immediately after each 15 min rs-fMRI scan, which retrospectively measured various dimensions of spontaneous thought. The battery included 12 statements (Supplementary Table 1) which participants rated on a visual analog

scale with 5% response increments that go from 0% = “describes my thoughts not at all” to 100% = “describes my thoughts completely”. The current analysis sample includes MDES data for all available instances of rs-fMRI scans for each participant.

Analyses

Dimension reduction for questionnaire and MDES data

We performed two separate principal component analyses (PCA) to obtain low dimensional summaries of the 71 trait variables from 21 questionnaires, and 12 thought variables from MDES.

71 scores from the personality questionnaires of 144 participants were included and missing data (3.15% of including all variables) were imputed by the variable mean. PCA was performed on this matrix and five “trait” components (henceforth referred to as “traits”) were selected on the basis of eigenvalues >1 , using the Kaiser-Guttman criterion (Jolliffe, 2002) and their congruence with the previously well-established “Big Five” personality traits (Digman, 1990; Cobb-Clark and Schurer, 2012). For the MDES data, separate instances of responding for each participant were concatenated, resulting in a matrix with 576 observations of 12 variables. PCA was performed on this matrix, and five “thought” components (henceforth referred to as “thought patterns”) were selected on the basis of eigenvalues >1 . Varimax rotation was applied to both solutions to optimize the distinctiveness of each component. The five thought pattern scores were then averaged across the four scans, resulting in one score for each thought pattern for each participant, describing their location on a particular thought dimension.

Dimension reduction for whole-brain functional connectivity

Functional time-series for each participant was extracted using the Schaefer 400 parcellation (Schaefer et al., 2018) using the fully pre-processed data from all resting-state scans. The data from separate scans were concatenated, and a 400x400 connectivity matrix was calculated from the resulting time series for each participant using Pearson correlation. A

group connectivity matrix of the whole sample was calculated by averaging the 144 individual matrices.

In order to summarize whole brain connectivity in a low-dimensional space, we performed gradient analysis using the Brainspace toolbox (Vos de Wael et al., 2020). 10 macro-scale gradients were calculated for the group (Fig 2). First, we applied fisher's z transform to the group matrix, building an affinity matrix (kernel= normalized angle, sparsity= 0.9) and then decomposed it using PCA. We chose the PCA approach for gradient calculation, as Hong and colleagues (Hong et al., 2020) have shown that compared to non-linear decomposition methods, PCA provides better reliability and higher phenotypic predictive value for connectivity gradients. For ease of interpretation and comparability, group gradients were aligned to a subsample of the HCP dataset (Essen et al., 2013) included in BrainSpace. Finally, following McKeown et al. (McKeown et al., 2020) 10 gradients were calculated in order to maximize the gradient fit for all individuals during alignment. Individual gradients were calculated for each participant, aligned to the group-level gradients, resulting in a 400x10 matrix for each participant. 10 gradients were calculated to facilitate alignment across individuals irrespective of differences in rank order of individual gradients (to control for the possibility, for example, that the pattern summarised by group-level Gradient 2 is Gradient 4 for some participants). Subsequent regression analyses were limited to the first three group-aligned gradients, which have been relatively well-characterized in previous work (Margulies et al., 2016; McKeown et al., 2020; Turnbull et al., 2020). To visualize the functional axis captured by each gradient, we performed Neurosynth (Yarkoni et al., 2011) decoding on the group gradient maps (Fig 2). Further, we calculated the average gradient score for all parcels within each of the seven connectivity networks described by Yeo and colleagues (Yeo et al., 2011) (Fig 2).

Stability of thoughts-patterns and gradients

To quantify the stability of thoughts and connectivity gradients over the whole scanning period, intra-class correlation coefficients (ICC) were calculated for the thought patterns and following

Hong and colleagues (Hong et al., 2020), discriminability indices (Bridgeford et al., 2021) were calculated for whole gradients by treating the 4 scans and subsequent thought probes as separate instances. We used the two-way mixed effects model (i.e. type 3 ICC) used for quantifying test-retest reliability, where samples cannot be considered independent (Koo and Li, 2016). Only the 135 participants who had four full-length resting state scans were included in this analysis. As this analysis found reasonable levels of reliability (see below), the averages of the four separate thought scores were used as regressors in subsequent analysis. This allowed for both a robust measure of thought patterns over the whole testing period, and the inclusion of all 144 participants in the analysis.

Multiple Multivariate Regression

To investigate the relationship between individual differences in traits, thoughts, and macro-scale cortical gradients, we used multiple multivariate regression as implemented in the MATLAB SurfStat Toolbox (Worsley et al., 2009) [<http://www.math.mcgill.ca/keith/surfstat/>]. 400 separate linear models were estimated for 400 parcels, with the gradient scores from the first three gradients as the dependent variables, and with five trait scores (Fig 1A) and five thought scores (Fig 1B), as well as nuisance variables age, motion, and gender included as independent variables. The resulting significant effects from 400 parcels were corrected for False Discovery Rate (FDR, $q < .05$) (Storey, 2003) at the multivariate (three gradients) level. In order to test which gradient score was affected, follow-up univariate analyses were performed on the resulting parcels for each gradient separately, and effects were further Bonferroni corrected ($p_{\text{bonf}} < .025$) for the total number of comparisons performed for all parcels (including the analyses of all three gradients) for each variable. Additionally, to see how the trait components related to the thought components, we performed multiple multivariate regression with the thoughts as dependent variables and traits as independent variables.

Results

Traits and thought patterns

Application of PCA to the battery of personality questionnaires resulted in five “traits” (Fig 1 A, supp fig 1) with eigenvalues > 1 , explaining 48.4% of the variance. The five trait components, independent of the direction of loadings, largely map onto the “big five” personality factors: neuroticism, conscientiousness (positive loading on “procrastination” in our PCA result), extraversion (positive loading on “introversion” in our PCA result), agreeableness (positive loading on anti-social in our results) and openness to experience, respectively. Application of PCA to the MDES questions revealed five “thought patterns” (Fig. 1 B, supp fig 2) with eigenvalues > 1 , explaining 65.4% of the variance. Based on the most heavily loaded dimensions within each pattern, we named these: “modality” (image vs words), “positive episodic social”, “specific internal”, “self-relevant” and “prospective”.

Intraclass correlation – thoughts

Our first analysis established the reliability of thought components across the one hour of scanning. The five thought patterns showed low to moderate agreement between individual scores from single sessions (modality = 0.5856, positive episodic-social= 0.4531, specific internal= 0.5226, self-relevant = 0.5832, prospective= 0.3118), indicating a degree of variability between sessions. The average of all scores had high ICCs for the first four components (modality = 0.8497, positive episodic social = 0.7783, specific internal = 0.8141, self-relevant = 0.8484, prospective = 0.6444). The average scores from 4 sessions were used as regressors in subsequent analyses.


Next, we examined the relationship between the low dimensional representations of both personality and thoughts. Multiple multivariate regression using traits as predictor variables of thought patterns established that “negative affect” had a significant effect on thoughts ($5,134$) = 3.88, $p = 0.003$, partial $\eta^2 = 0.127$). Univariate follow-up showed that a high score on trait neuroticism was significantly associated with less “positive episodic social” thought (pattern 2;

$\beta = -0.229$, 95% CI = [-0.389 -0.07], $p = 0.005$, partial $\eta^2 = 0.055$) as well as greater “self-relevant” thought (pattern 4; $\beta = 0.229$, 95% CI = [0.066 0.391], $p = 0.006$, partial $\eta^2 = 0.053$), (Supplementary Fig. 3).

A. Trait Components

1. Negative affect

Anxious Depressed
Critical Items Avoidant
Borderline Anxiety Depression
Aggressive Behaviour
Somatic Complaints
Externalizing Inhibition
Neuroticism Dependent
Depressive Internalizing
Negative Urgency



2. Procrastination

Procrastination
Attention Problems Self Worth
Obsessive Compulsive
Adaptive Functioning (Edu)
Lack of Perseverance
Conscientiousness
Positive Urgency Self Control
Lack of Premeditation

4. Antisocial

Social Desirability
Anger Outward Antisocial
Anger Trait Narcissism
Agreeableness
Psychopathy Intrusive
Externalizing

3. Introversion

Withdrawn
Extraversion
Histrionic Schizoid
Rhapsodic
Paranoid
Adaptive Functioning (Friends)

5. Openness

Thought Problems
Movement (Music)
Openness
Schizotypal

B. Thought Patterns


1. Modality



2. Positive Episodic Social



3. Specific Internal



4. Self-relevant



5. Prospective



Figure 1. Principal components of traits and thoughts

A) First five trait components derived from PCA after varimax rotation are represented as word-clouds with negative loadings shown in blue, and positive loadings in red; the absolute loading value is represented by the font size of the item. In the bottom left panel, scree-plot showing the percentage of trait variance explained by the each of the first 10 components in grey, and the first five components after varimax rotation in red. **B)** Results of the application of PCA to the MDES data, depicted in the same way.

Macro-scale cortical gradients

The first three group level gradients are shown in Fig. 2 along with their Neurosynth meta-analytic associations and relationships to the Yeo-networks (Yeo et al., 2011) (7-network solution). The first gradient (G1) differentiates between visual regions at one end and DMN at the other. The second gradient (G2) describes the dissociation between somatomotor and visual cortices. The third gradient (G3) captures the segregation between different transmodal systems (the default mode network versus the fronto-parietal system). The three gradients are largely similar to the ones reported by Margulies et al. (Margulies et al., 2016) and subsequent literature (Hong et al., 2019; Paquola et al., 2019; Bethlehem et al., 2020; McKeown et al., 2020; Turnbull et al., 2020). Due to the difference in the decomposition provided by PCA vs. DM approach, the endpoints of G1 are different from those first reported by Margulies et al (Margulies et al., 2016) in that one end is anchored by the visual network alone, as opposed to visual and somatomotor, while in G2, the somatomotor network is separated from both the visual and default mode networks, as opposed to the visual network alone in the (Margulies et al., 2016) study. However, like those reported by Margulies and colleagues, the first two gradients together describe network-level connectivity space anchored at three ends by the visual, somatomotor and default mode network, respectively (Fig. 3). Stability of results irrespective of the decomposition approach is shown in supplementary Figure 4. Single gradients tended to be stable over the 4 sessions, with a discriminability index of 0.964 for Gradient 1, 0.918 for Gradient 2, and 0.983 for Gradient 3 over four adjacent scans from 135

participants. Discriminability indices are similar to those previously reported by (Hong et al., 2020).




Figure 2. Group-level gradients of functional connectivity

On the left are the first three group-averaged gradients, represented in left lateral and medial views. Regions with similar whole-brain connectivity profiles are shown in similar colours, with yellow and purple regions indicating most dissimilar connectivity patterns. Loading ranges and directions are arbitrary. In the middle, word clouds representing the top 10 positive (red) and negatively correlated (blue) Neurosynth decoding topic terms for each gradient map. On the right, radar-plots showing the Yeo-network profile of each group-level gradient depicted in the left column. Each radar-plot shows the mean gradient loadings for all parcels within the seven Yeo networks.




Figure 3. Comparison of group-level gradients to BrainSpace HCP template

The first scatterplot shows 400 parcel positions along G1 and G2 in the template calculated from the HCP subsample included in BrainSpace toolbox (Vos de Wael et al., 2020). The second scatterplot shows parcel positions in the group-level gradients G1 and G2 after Procrustes alignment to the HCP template. Parcels are color-coded according to their respective Yeo network. Yeo networks are shown as color-coded brain maps on the right.

Relationship between state- trait variability and cortical gradients

Having established low dimensional representations of thought, personality, and brain organisation, we next examined associations between different types of personality and ongoing thought experienced during the scan and our metrics of functional brain organisation. To this end, we performed a multiple multivariate regression with thoughts, traits, and nuisance variables (motion, age and gender) as independent variables, with whole brain functional organisation, as captured by the first three gradients, as dependent variables. In these analyses both trait “introversion” and a pattern describing “specific internal” thought showed significant effects at the multivariate level. Results from the univariate follow-up of effects within each gradient are shown in Fig. 4 and 5, and Table 2.

Trait Introversion (Fig 4)

Along the first gradient, a parcel within the right orbitofrontal cortex (within the executive control network, shown in orange) showed more similarity with transmodal regions for individuals high on introversion. Six parcels within the ventral attention network, including

anterior insula, operculum and cingulate cortex were closer to the somatomotor end along gradient two (shown in purple). The same regions showed lower scores along the third gradient in participants with higher introversion scores, indicating stronger integration with the default mode network. A parcel within posterior cingulate cortex (control) was also more segregated from the visual end of gradient two in participants with higher introversion scores.




Figure 4. Relationship between trait “Introversion” on the first three connectivity gradients

On the left, parcels within the first three gradients that show significant ($p_{\text{bonf}} < 0.025$) differences related to trait “introversion”, orange indicating regions within the “frontoparietal control network”, and violet indicating regions within the “ventral attention”. Scatter plots depict the relationship between individual scores for “introversion” thought (x-axis) and average

gradient score of all affected parcels (y-axis) within each gradient. Each datapoint is a participant. Both axes show standardized scores. Detailed results from individual parcels are reported in table 2. The right column shows Neurosynth decoding of ROI maps of affected parcels within each gradient, showing top ten positively correlated topic terms in red, and top ten negatively associated topic terms in blue.

Detailed internal cognition (Fig 5)

Relationships with patterns of detailed internal cognition were confined to the dorsal attention and visual networks. A region within the superior parietal lobule (DAN) had lower scores on the first gradient (more transmodal) and higher scores on the second gradient (less visual), indicating less similarity with visual regions whose ongoing experience was more “specific” and “internal”. Along the third gradient, higher “specific internal” thought scores were associated with greater separation between these regions and the default mode network. Finally, a parcel within the parahippocampal gyrus/ extrastriate (visual network) showed a broad spread along gradient one, with participants with higher “specific internal” thought scores falling on the transmodal/ DMN side and participants with lower scores (higher “surroundings”) closer to the visual system side.




Figure 5. Relationship between specific internal thought and the first three connectivity gradients

On the left, parcels within the first three gradients that show significant differences ($p_{\text{bonf}} < 0.025$) related to “specific internal” thought, green indicating regions within “dorsal attention network” (DAN), and purple indicating regions within the “visual network”. Scatter plots depict the relationship between individual scores for “specific internal” thought (x-axis) and average gradient score of all affected parcels (y-axis) within each gradient. Each datapoint is a participant. Both axes show standardized scores. Detailed results from individual parcels are reported in table 1. The right column shows Neurosynth decoding of ROI maps of affected parcels within each gradient, showing top ten positively correlated topic terms in red, and top ten negatively associated topic terms in blue.

Table 2. Relationships between first three connectivity gradients and introversion and specific internal thought

IV	DV	Yeo Network	Parcel	t (130)	p _{uncorr}	p _{bonf}
Introversion	G1	Control	OFC (R)	-4.2214	0.00002	0.0012
	G2	VAN	A Ins (L)	3.7888	0.00012	
			A Ins (R)	4.3110	0.00002	
			Fr Oper (R)	3.8081	0.00011	
			A Cing (R)	3.1879	0.00090	
		Control	P Cing (L)	4.0504	0.00004	
	G3	VAN	A Ins (L)	-3.9539	0.00006	
			A Ins (L)	-3.6784	0.00017	
			A Ins (R)	-4.2031	0.00002	
			Fr Oper (R)	-4.1767	0.00003	
			A Cing (R)	-3.9732	0.00006	
<hr/>						
Specific Internal						
Thought	G1	Visual	PHC/ ExStr	-2.9105	0.00212	0.0028
		DAN	SPL (L)	-4.5433	0.00001	
	G2	DAN	SPL (L)	4.1217	0.00003	
		DAN	SPL (R)	3.3542	0.00052	
	G3	DAN	SPL (L)	4.4548	0.00001	

IV = Independent Variable, **DV** = Dependent Variable, **G** = Gradient, **VAN** = Ventral Attention Network, **DAN** = Dorsal Attention Network, **OFC** = Orbitofrontal Cortex, **A** = Anterior, **P** = Posterior, **Fr** = Frontal, **Ins** = Insula, **L** = Left, **R** = Right, **Oper** = Operculum, **Cing** = Cingulate Cortex, **PHC** = Parahippocampal Cortex, **ExStr** = Extrastriate cortex, **SPL** = Superior Parietal Lobule. Results reported in the table are from univariate (single-gradient) follow-up tests for parcels showing a significant effect for each IV at the multivariate (3-gradient) level. Univariate tests are Bonferroni corrected for the total number of parcels (all 3 gradients) where tests were performed (21 parcels for Introversion, 9 for Specific Internal Thought).

Discussion

Our study investigates whether an emerging “tri-partite” perspective from contemporary views of ongoing conscious thought can provide a framework that can account for the relationship between dispositional traits, ongoing thought, and individual differences in large scale patterns of neural connectivity at rest. This tri-partite network account emphasises the roles of regions embedded within three large scale networks as important for ongoing experience: the ventral attention network, the dorsal attention network and the default mode network (Smallwood et al., 2021; Huang et al., 2021). We calculated macro-scale connectivity gradients from one hour of resting state fMRI for 144 participants. The variability of these gradients was then analysed as a function of self-reports of ongoing thought patterns (captured by the principal components of MDES) and personality traits (described by the principal components of a battery of personality and habit measures). Given the tendency of certain traits to be correlated with frequency of specific patterns of thought (e.g. depression level with intrusive and negative thought (Konu et al., 2021) we also looked for possible dependencies between trait and thought components through multivariate regression using traits as predictors.

Our analyses confirmed that both patterns of thought, and indices of traits, contribute to patterns of brain organisation in a manner that converges with the emerging tri-partite view of ongoing conscious thought. For example, it has been hypothesised that the ventral attention network (VAN) helps adjudicate between internal and external influences on ongoing thought (Smallwood et al., 2021, Huang et al., 2021) and we found that individuals who were high on dispositional “introversion” showed variation in anterior insula, overlying operculum and ACC: all regions making up part of the VAN. For more introverted people these regions showed greater alignment with somatomotor regions and less with visual cortex (gradient 2), and greater alignment with default mode network than the fronto parietal network (gradient 3). Notably, prior studies have found that regions of sensorimotor cortex are linked to deliberate mind-wandering (Golchert et al., 2017) and individuals who tend to generate patterns of episodic social cognition during periods of low task demands show greater temporal

correlation between the VAN and sensori-motor cortex (Turnbull et al., 2019a). Introversion reflects a predisposition towards internal subjective states rather than external objects (Jung, 1995), and so our analysis adds to an emerging literature (Golchert et al., 2017, Turnbull et al., 2019) that suggests that there may be an important relationship between sensorimotor cortex and the ventral attention network in patterns of internal thought.

Our study also highlighted that associations between observed patterns of neural organisation at rest and patterns of ongoing thought are also consistent with hypothesized tripartite view (Smallwood et al., 2021, Huang et al., 2021). For example, we found a pattern of detailed thinking during wakeful rest that was correlated with stronger decoupling between a region within dorsal parietal cortex from the visual network (as indexed by changes in both gradients 1 and 2), and greater separation from the default mode network (gradient 3). This region overlaps with a region within the DAN identified by (Turnbull et al., 2019b) in which brain activity was reduced during self-generated thoughts relative to external task focus, suggesting an important role for the DAN in external facing attention. Furthermore, using a technique known as “echoes” analysis (see Leech et al., 2012), Turnbull and colleagues (Turnbull et al., 2019b) established that individuals who engaged in self-generated thought during situations of low external demand, at rest showed greater separation between the dorsal attention network and lateral regions of the default mode network in a region of the dLPFC, also a member of the VAN. Thus our study confirms prior studies that highlight that greater segregation between the DAN and the DMN in the capacity to engage in thoughts that are less linked to the external environment. Together, therefore, the convergence between the current analysis and perspectives from research on conscious experience highlight a high degree of overlap in both the regions identified and the hypothesised functions. Our observations are important, therefore, because they help establish that, with the appropriate methodology (McKeown et al., 2020; Finn, 2021; Gonzalez-Castillo et al., 2021) neural accounts of conscious experience (Smallwood et al., 2021; Huang et al., 2021) provide an important valuable way to make sense of brain-cognition links observed at rest.

Second, our data provides evidence for the “decoupling” hypothesis of self-generated experience (Smallwood et al., 2013). This perspective emerged from observations that cortical processing of external inputs is reduced when individuals focus on internal self-generated thought (Smallwood et al., 2008; Kam et al., 2011; Baird et al., 2014), and assumes that this reduced processing of external input allows an internal train of thought to persist in a more detailed manner (p.524,Smallwood, 2013 (Smallwood, 2013b)). Our data is consistent with this view since both personality traits linked to internal focus (“introversion”), and patterns of detail experience that are not directed externally (“specific internal thought”), are linked to reductions in the similarity between neural activity in regions linked to attention and cognition, with regions of visual cortex. In this way our study provides novel insight into how the macro-scale functional patterns across the cortex support the emergence of detailed patterns of internal experience. Critically, in our study there was no external task from which thinking needed to be decoupled from, ruling out accounts of this process as a “lapse” in the normal upregulation of task relevant material needed for task completion (for discussion see Franklin et al., 2013, Smallwood 2013ab).

Although our study establishes how contemporary work on conscious experience can help understand patterns of brain organisation observed at rest and highlights how these approaches can be leveraged to understand the neural correlates of both an individuals’ traits, and their thoughts, there are nonetheless important questions that our study leaves open. For example, contemporary work on ongoing conscious thought highlights time and context as key variables necessary for understanding the neural correlates of different features of thinking. Since the aim of our study was to focus on the brain at rest, interpretations of our results should bear in mind that under different task conditions, neural correlates between thinking, personality and neural activity may be different. For example, prior studies have established that posterior elements of the default mode network can become integrated into task positive systems (Krieger-Redwood et al., 2016; Vatansever et al., 2017) and under demanding task conditions the default mode network is linked to patterns of task focused

cognition (Sormaz et al., 2018). Similarly, our analysis focused on ‘static’ indices of neural activity rather than dynamic measures. We chose to focus on static indices of neural activity because our prior studies have shown that brain-behaviour correlations can be relatively stable over time (Wang et al., 2020), and as we establish in our study these patterns show reasonable stability across a one hour session. Furthermore, our study, particularly with respect to the findings relating to DAN, map closely onto studies that use experience sampling to identify momentary correlations between neural activity and experience (Turnbull et al., 2019b, 2019a, 2020). Nonetheless, there are likely to be important dynamic features (Kucyi, 2018) of ongoing experience that our analysis of static brain organisation cannot capture. Recent discussions in cognitive neuroscience have highlighted the link between sample size and the reproducibility of brain-wide associations with behavioural phenotypes (Marek et al., 2022; Spisak et al., 2023). Our analysis of the trait and thought data alone revealed that “neuroticism” was related to high negative and episodic thoughts, however, we did not find any other significant relationships among traits and thought patterns. In the current data, neuroticism was the most prominent out of all five traits included in the analysis, accounting for 29% of the total variance explained by them. It is therefore likely that more extensive and robust correlations between thoughts patterns and other traits, as well as thoughts, traits and macroscale connectivity patterns would emerge with data sets with larger sample sizes. It is worth noting that mapping momentary changes between ongoing experience and neural activity will likely depend on a data set tailored to this question, in particular in which (i) experience sampling measures are collected more frequently as well as (ii) methodological advances that allow patterns of activity to be mapped without using temporal correlation. and that measure thinking across multiple contexts.

Author Contributions

Jonathan Smallwood, and Samyogita Hardikar conceived of this study. Bronte McKeown, Daniel Margulies, H. Lina Schaare, Reinder Vos de Wael, Mark Lauckner contributed analysis

tools, and Samyogita Hardikar performed the analysis. Samyogita Hardikar and Jonathan Smallwood wrote the main manuscript text and Samyogita Hardikar prepared the figures. Boris Bernhardt, Daniel Margulies, Sofie L Valk, Mark Lauckner, Adam Turnbull, Ting Xu, and Arno Villringer provided help with the conceptual and methodological framework for this study, as well as feedback on the manuscript. All authors reviewed the final manuscript.

References

Achenbach TM, Rescorla LA (2003) No TitleManual for the ASEBA adult forms & profiles. Research Center for Children, Youth, & Families. University of Vermont, Burlington, VT, USA.

Baird B, Smallwood J, Lutz A, Schooler JW (2014) The Decoupled Mind: Mind-wandering Disrupts Cortical Phase-locking to Perceptual Events. *J Cogn Neurosci* 26:2596–2607.

Beck A, Steer R, Brown G (1993) Beck Depression Inventory-II (BDI-II). ... Beck Depress Invent.

Bethlehem RAI, Paquola C, Seidlitz J, Ronan L, Bernhardt B, Consortium C-CCAN, Tsvetanov KA (2020) Dispersion of functional gradients across the adult lifespan. *Neuroimage* 222:117299 Available at: <https://linkinghub.elsevier.com/retrieve/pii/S1053811920307850> [Accessed November 3, 2021].

Bridgeford EW, Wang S, Yang Z, Wang Z, Xu T, Craddock C, Dey J, Kiar G, Gray-Roncal W, Colantuoni C, Douville C, Noble S, Priebe CE, Caffo B, Milham M, Zuo X-N, Reproducibility C for R and, Vogelstein JT (2021) Eliminating accidental deviations to minimize generalization error and maximize replicability: applications in connectomics and genomics. *bioRxiv*:802629 Available at: <https://www.biorxiv.org/content/10.1101/802629v8> [Accessed September 24, 2021].

Cao H et al. (2018) Cerebello-thalamo-cortical hyperconnectivity as a state-independent

functional neural signature for psychosis prediction and characterization. *Nat Commun*

9 Available at: [/pmc/articles/PMC6155100/](https://PMC6155100/) [Accessed March 22, 2022].

Carriere JSA, Seli P, Smilek D (2013) Wandering in both mind and body: Individual differences in mind wandering and inattention predict fidgeting. *Can J Exp Psychol Can Psychol expérimentale* 67:19–31 Available at: <http://doi.apa.org/getdoi.cfm?doi=10.1037/a0031438> [Accessed April 4, 2019].

Carver CS, White TL (1994) Behavioral Inhibition , Behavioral Activation , and Affective Responses to Impending Reward and Punishment : The BIS / BAS Scales. *J Pers Soc Psychol* 67:319–333.

Cobb-Clark DA, Schurer S (2012) The stability of big-five personality traits. *Econ Lett* 115:11–15.

Corbetta M, Shulman GL (2002) Control of goal-directed and stimulus-driven attention in the brain. *Nat Rev Neurosci* 3:201–215 Available at: <https://pubmed.ncbi.nlm.nih.gov/11994752/> [Accessed March 21, 2022].

Costa PT, McCrae RR (2008) The revised NEO personality inventory (NEO-PI-R). SAGE Handb Personal Theory Assess Vol 2 - Personal Meas Test:179–198.

Crowne DP, Marlowe D (1960) A new scale of social desirability independent of psychopathology. *J Consult Psychol* 24:349–354.

Cui Z et al. (2020) Individual Variation in Functional Topography of Association Networks in Youth. *Neuron* 106:340-353.e8.

Derryberry D, Reed MA (2002) Anxiety-related attentional biases and their regulation by attentional control. *J Abnorm Psychol* 111:225–236 Available at: [/record/2002-12652-002](https://doi.org/10.1037/0021-9010.111.2.225) [Accessed September 24, 2021].

Digman JM (1990) PERSONALITY STRUCTURE: EMERGENCE OF THE FIVE-FACTOR MODEL. *Annu Rev Psychol* 41:417–457 Available at: www.annualreviews.org

[Accessed October 10, 2023].

Dosenbach NUF, Nardos B, Cohen AL, Fair DA, Power JD, Church JA, Nelson SM, Wig GS, Vogel AC, Lessov-Schlaggar CN, Barnes KA, Dubis JW, Feczko E, Coalson RS, Pruett JR, Barch DM, Petersen SE, Schlaggar BL (2010) Prediction of individual brain maturity using fMRI. *Science* (80-) 329:1358–1361 Available at: <https://www.science.org/doi/abs/10.1126/science.1194144> [Accessed March 21, 2022].

Essen DC Van, Smith SM, Barch DM, Behrens TEJ, Yacoub E, Ugurbil K, Consortium for the W-MH (2013) The WU-Minn Human Connectome Project: An Overview. *Neuroimage* 80:62 Available at: [/pmc/articles/PMC3724347/](https://pmc/articles/PMC3724347/) [Accessed August 12, 2021].

Farmer R, Sundberg ND (1986) Boredom proneness--the development and correlates of a new scale. *J Pers Assess* 50:4–17 Available at: <https://europepmc.org/article/MED/3723312> [Accessed September 24, 2021].

Finn ES (2021) Is it time to put rest to rest? *Trends Cogn Sci* Available at: <https://linkinghub.elsevier.com/retrieve/pii/S1364661321002345> [Accessed October 14, 2021].

Floridou GA, Williamson VJ, Stewart L, Müllensiefen D (2015) The Involuntary Musical Imagery Scale (IMIS). *Psychomusicology Music Mind, Brain* 25:28–36 Available at: [/record/2015-22418-001](https://doi.org/10.1080/0938203X.2015.1002345) [Accessed September 24, 2021].

Golchert J, Smallwood J, Jefferies E, Seli P, Huntenburg JM, Liem F, Lauckner ME, Olschläger S, Bernhardt BC, Villringer A, Margulies DS (2017) Individual variation in intentionality in the mind-wandering state is reflected in the integration of the default-mode, fronto-parietal, and limbic networks. *Neuroimage* 146:226–235 Available at: <https://www.sciencedirect.com/science/article/pii/S1053811916306425?via%3Dihub> [Accessed July 31, 2018].

Gonzalez-Castillo J, Kam JWYY, Hoy CW, Bandettini PA (2021) How to Interpret Resting-State fMRI: Ask Your Participants. *J Neurosci* 41:1130–1141 Available at: <https://doi.org/10.1523/JNEUROSCI.1786-20.2020> [Accessed July 19, 2021].

Gratton C, Smith DM, Dorn M (2020) Digging Deeper to Chart the Landscape of Human Brain Development. *Neuron* 106:209–211 Available at: <https://doi.org/10.1016/j.neuron.2020.03.030> [Accessed February 15, 2022].

Hong SJ, de Wael RV, Bethlehem RAI, Lariviere S, Paquola C, Valk SL, Milham MP, Di Martino A, Margulies DS, Smallwood J, Bernhardt BC (2019) Atypical functional connectome hierarchy in autism. *Nat Commun* 10.

Hong SJ, Xu T, Nikolaidis A, Smallwood J, Margulies DS, Bernhardt B, Vogelstein J, Milham MP (2020) Toward a connectivity gradient-based framework for reproducible biomarker discovery. *Neuroimage* 223:117322.

Hsu WT, Rosenberg MD, Scheinost D, Constable RT, Chun MM (2018) Resting-state functional connectivity predicts neuroticism and extraversion in novel individuals. *Soc Cogn Affect Neurosci* 13:224–232 Available at: <https://academic.oup.com/scan/article/13/2/224/4823668> [Accessed March 22, 2022].

Huang Z, Tarnal V, Vlisides PE, Janke EL, McKinney AM, Picton P, Mashour GA, Hudetz AG (2021) Anterior insula regulates brain network transitions that gate conscious access. *Cell Rep* 35 Available at: <https://doi.org/10.1016/j.celrep.2021.109081> [Accessed May 27, 2021].

Johns MW (1991) A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep* 14:540–545 Available at: <https://europepmc.org/article/MED/1798888> [Accessed September 24, 2021].

Jolliffe IT (2002) Principal Component Analysis. New York: Springer-Verlag. Available at: <http://link.springer.com/10.1007/b98835> [Accessed February 15, 2022].

Jones DN, Paulhus DL (2014) Introducing the Short Dark Triad (SD3): A Brief Measure of Dark Personality Traits. *Assessment* 21:28–41 Available at: [/record/2014-10974-004](https://doi.org/10.1037/a0035744) [Accessed September 24, 2021].

Jung CG (1995) Memories, Dreams, Reflections. In: *Memories, Dreams, Reflections*, pp 414–415. London: Fontana Press.

Kam JWY, Dao E, Farley J, Kevin F, Smallwood J, Schooler JW, Handy TC (2011) Slow fluctuations in attentional control of sensory cortex. *J Cogn Neurosci* 23:460–470 Available at: <https://pubmed.ncbi.nlm.nih.gov/20146593/> [Accessed October 22, 2021].

Karapanagiotidis T, Jefferies E, Smallwood J (2021) Interactions between the neural correlates of dispositional internally directed thought and visual imagery. *Philos Trans R Soc B* 376:20190691 Available at: <https://royalsocietypublishing.org/doi/abs/10.1098/rstb.2019.0691> [Accessed August 11, 2021].

Karapanagiotidis T, Vidaurre D, Quinn AJ, Vatansever D, Poerio GL, Turnbull A, Ho NSP, Leech R, Bernhardt BC, Jefferies E, Margulies DS, Nichols TE, Woolrich MW, Smallwood J (2020) The psychological correlates of distinct neural states occurring during wakeful rest. *Sci Rep* 10 Available at: [/pmc/articles/PMC7712889/](https://pmc/articles/PMC7712889/) [Accessed August 11, 2021].

Karapanagiotidis T, Vidaurre D, Quinn AJ, Vatansever D, Poerio GL, Turnbull A, Leech R, Bernhardt B, Jefferies E, Margulies DS, Nichols TE, Woolrich MW, Smallwood J (2019) Emergence of neural dynamics within a co-ordinate space of large-scale neural hierarchies. *bioRxiv*:2019.12.21.885772 Available at: <https://www.biorxiv.org/content/10.1101/2019.12.21.885772v1> [Accessed January 20, 2020].

Koban L, Gianaros PJ, Kober H, Wager TD (2021) The self in context: brain systems linking mental and physical health. *Nat Rev Neurosci* 2021 22:309–322 Available at:

<https://www.nature.com/articles/s41583-021-00446-8> [Accessed March 10, 2022].

Konu D, McKeown B, Turnbull A, Siu Ping Ho N, Karapanagiotidis T, Vanderwal T, McCall C, Tipper SP, Jefferies E, Smallwood J (2021) Exploring patterns of ongoing thought under naturalistic and conventional task-based conditions. *Conscious Cogn* 93:103139.

Koo TK, Li MY (2016) A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *J Chiropr Med* 15:155 Available at: [/pmc/articles/PMC4913118/](https://pmc/articles/PMC4913118/) [Accessed August 11, 2021].

Krieger-Redwood K, Jefferies E, Karapanagiotidis T, Seymour R, Nunes A, Ang JWA, Majernikova V, Mollo G, Smallwood J (2016) Down but not out in posterior cingulate cortex: Deactivation yet functional coupling with prefrontal cortex during demanding semantic cognition. *Neuroimage* 141:366–377 Available at: <https://pubmed.ncbi.nlm.nih.gov/27485753/> [Accessed September 24, 2021].

Kucyi A (2018) Just a thought: How mind-wandering is represented in dynamic brain connectivity. *Neuroimage* 180:505–514.

Kucyi A, Esterman M, Capella J, Green A, Uchida M, Biederman J, Gabrieli JDE, Valera EM, Whitfield-Gabrieli S (2021) Prediction of stimulus-independent and task-unrelated thought from functional brain networks. *Nat Commun* 12:1–17 Available at: <https://doi.org/10.1038/s41467-021-22027-0> [Accessed May 13, 2021].

Kuhl J, Kazén M (2009) Persönlichkeits-Stil-und-Störungs-Inventar (PSSI): Manual (Hogrefe, ed).

Leech R, Braga R, Sharp DJ (2012) Echoes of the Brain within the Posterior Cingulate Cortex. *J Neurosci* 32:215–222 Available at: <https://www.jneurosci.org/content/32/1/215> [Accessed September 24, 2021].

Lynam DR, Smith GT, Whiteside SP, Cyders MA (2006) The UPPS-P: Assessing five personality pathways to impulsive behavior. West Lafayette, Purdue Univ 10.

Marek S et al. (2022) Reproducible brain-wide association studies require thousands of individuals. *Nature* 603:654 Available at: [/pmc/articles/PMC8991999/](https://PMC8991999/) [Accessed December 14, 2022].

Margulies DS, Ghosh SS, Goulas A, Falkiewicz M, Huntenburg JM, Langs G, Bezgin G, Eickhoff SB, Castellanos FX, Petrides M, Jefferies E, Smallwood J (2016) Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proc Natl Acad Sci U S A* 113:12574–12579 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27791099> [Accessed December 11, 2019].

McKeown B, Strawson WH, Wang HT, Karapanagiotidis T, Vos de Wael R, Benkarim O, Turnbull A, Margulies D, Jefferies E, McCall C, Bernhardt B, Smallwood J (2020) The relationship between individual variation in macroscale functional gradients and distinct aspects of ongoing thought. *Neuroimage* 220:117072.

Mendes N et al. (2019) A functional connectome phenotyping dataset including cognitive state and personality measures. *Sci Data* 6:180307 Available at: <http://www.nature.com/articles/sdata2018307> [Accessed April 4, 2019].

Menon V (2011) Large-scale brain networks and psychopathology: a unifying triple network model. *Trends Cogn Sci* 15:483–506 Available at: <https://pubmed.ncbi.nlm.nih.gov.browser.cbs.mpg.de/21908230/> [Accessed March 10, 2022].

Müllensiefen D, Gingras B, Musil J, Stewart L (2014) Measuring the facets of musicality: The Goldsmiths Musical Sophistication Index (Gold-MSI). *Pers Individ Dif* 60:S35.

O'Malley PM, Bachman JG (1979) Self-esteem and education: Sex and cohort comparisons among high school seniors. *J Pers Soc Psychol* 37:1153–1159 Available at: [/record/1980-33361-001](https://record/1980-33361-001) [Accessed September 24, 2021].

Ophir E, Nass C, Wagner AD (2009) Cognitive control in media multitaskers. *Proc Natl Acad Sci* 106:15583–15587 Available at: <https://www.pnas.org/content/106/37/15583>

[Accessed September 24, 2021].

Paquola C, Vos De Wael R, Wagstyl K, Bethlehem RAI, Hong S-J, Seidlitz J, Bullmore ET, Evans AC, Misic B, Margulies DS, Smallwood J, Bernhardt BC (2019) Microstructural and functional gradients are increasingly dissociated in transmodal cortices. Kennedy H, ed. PLoS Biol 17:e3000284 Available at:
<http://dx.plos.org/10.1371/journal.pbio.3000284> [Accessed November 12, 2019].

Poerio GL, Sormaz M, Wang H-T, Margulies D, Jefferies E, Smallwood J (2017) The role of the default mode network in component processes underlying the wandering mind. Soc Cogn Affect Neurosci 12:1047–1062 Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/28402561> [Accessed July 31, 2018].

R Core Team (2021) R: A language and environment for statistical computing. Available at:
<https://www.r-project.org/>.

Schaefer A, Kong R, Gordon EM, Laumann TO, Zuo X-N, Holmes AJ, Eickhoff SB, Yeo BTT (2018) Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. Cereb Cortex 28:3095–3114 Available at: <https://academic-oup-com.browser.cbs.mpg.de/cercor/article/28/9/3095/3978804> [Accessed August 11, 2021].

Schmidt RE, Gay P, D'Acremont M, Van Der Linden M (2008) A German adaptation of the upps impulsive behavior scale: Psychometric properties and factor structure. Swiss J Psychol 67:107–112 Available at:
<http://psycontent.metapress.com/openurl.asp?genre=article&id=doi:10.1024/1421-0185.67.2.107> [Accessed June 5, 2014].

Smallwood J (2013a) Distinguishing how from why the mind wanders: A process-occurrence framework for self-generated mental activity. Psychol Bull 139:519–535.

Smallwood J (2013b) Searching for the elements of thought: Reply to franklin, mrazek,

broadway, and schooler (2013). *Psychol Bull* 139:542–547.

Smallwood J, Beach E, Schooler JW, Handy TC (2008) Going AWOL in the Brain: Mind Wandering Reduces Cortical Analysis of External Events. *J Cogn Neurosci* 20:458–469.

Smallwood J, Bernhardt BC, Leech R, Bzdok D, Jefferies E, Margulies DS (2021a) The default mode network in cognition: a topographical perspective. *Nat Rev Neurosci* 2021:1–11 Available at: <https://www-nature-com.browser.cbs.mpg.de/articles/s41583-021-00474-4> [Accessed July 7, 2021].

Smallwood J, Karapanagiotidis T, Ruby F, Medea B, de Caso I, Konishi M, Wang H-T, Hallam G, Margulies DS, Jefferies E (2016) Representing Representation: Integration between the Temporal Lobe and the Posterior Cingulate Influences the Content and Form of Spontaneous Thought Biagini G, ed. *PLoS One* 11:e0152272 Available at: <http://dx.plos.org/10.1371/journal.pone.0152272> [Accessed May 28, 2018].

Smallwood J, Turnbull A, Wang H ting, Ho NSP, Poerio GL, Karapanagiotidis T, Konu D, McKeown B, Zhang M, Murphy C, Vatansever D, Bzdok D, Konishi M, Leech R, Seli P, Schooler JW, Bernhardt B, Margulies DS, Jefferies E (2021b) The neural correlates of ongoing conscious thought. *iScience* 24:102132.

Sormaz M, Murphy C, Wang H, Hymers M, Karapanagiotidis T, Poerio G, Margulies DS, Jefferies E, Smallwood J (2018) Default mode network can support the level of detail in experience during active task states. *Proc Natl Acad Sci U S A* 115:9318–9323 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/30150393> [Accessed April 17, 2019].

Spisak T, Bingel U, Wager TD (2023) Multivariate BWAS can be replicable with moderate sample sizes. *Nature* 615:E4–E7 Available at: <https://www.nature.com/articles/s41586-023-05745-x> [Accessed March 31, 2023].

Storey JD (2003) The positive false discovery rate: a Bayesian interpretation and the q-

value. <https://doi.org/10.1214/aos/1074290335> 31:2013–2035 Available at: <https://projecteuclid.org/journals/annals-of-statistics/volume-31/issue-6/The-positive-false-discovery-rate--a-Bayesian-interpretation-and/10.1214/aos/1074290335.full> [Accessed November 8, 2021].

Tangney JP, Baumeister RF, Boone AL (2004) High self-control predicts good adjustment, less pathology, better grades, and interpersonal success. *J Pers* 72:271–324 Available at: <https://pubmed.ncbi.nlm.nih.gov/15016066/> [Accessed September 24, 2021].

Tuckman BW (2016) The Development and Concurrent Validity of the Procrastination Scale: <http://dx.doi.org/10.1177/0013164491512022> 51:473–480 Available at: <https://journals.sagepub.com/doi/10.1177/0013164491512022> [Accessed September 24, 2021].

Turnbull A, Karapanagiotidis T, Wang HT, Bernhardt BC, Leech R, Margulies D, Schooler J, Jefferies E, Smallwood J (2020) Reductions in task positive neural systems occur with the passage of time and are associated with changes in ongoing thought. *Sci Rep* 10 Available at: [/pmc/articles/PMC7303126/](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7303126/)?report=abstract [Accessed November 11, 2020].

Turnbull A, Wang H-T, Schooler JW, Jefferies E, Margulies DS, Smallwood J (2019a) The ebb and flow of attention: Between-subject variation in intrinsic connectivity and cognition associated with the dynamics of ongoing experience. *Neuroimage* 185:286–299 Available at: <https://www.sciencedirect.com/science/article/pii/S1053811918319414?via%3Dihub> [Accessed January 18, 2019].

Turnbull A, Wang H, Murphy C, Ho NSP, Wang X, Sormaz M, Karapanagiotidis T, Leech RM, Bernhardt B, Margulies DS, Vatansever D, Jefferies E, Smallwood J (2019b) Left dorsolateral prefrontal cortex supports context-dependent prioritisation of off-task thought. *Nat Commun* 10:3816 Available at: <http://www.nature.com/articles/s41467-019-11030-0>

019-11764-y [Accessed November 12, 2019].

Vatansever D, Menon DK, Stamatakis EA (2017) Default mode contributions to automated information processing. *Proc Natl Acad Sci U S A* 114:12821–12826 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/29078345> [Accessed April 29, 2019].

Vos de Wael R, Benkarim O, Paquola C, Lariviere S, Royer J, Tavakol S, Xu T, Hong S-J, Langs G, Valk S, Misic B, Milham M, Margulies D, Smallwood J, Bernhardt BC (2020) BrainSpace: a toolbox for the analysis of macroscale gradients in neuroimaging and connectomics datasets. *Commun Biol* 2020 31 3:1–10 Available at: <https://www.nature.com/articles/s42003-020-0794-7> [Accessed November 3, 2021].

Wang H-T, Ho NSP, Bzdok D, Bernhardt BC, Margulies DS, Jefferies E, Smallwood J (2020) Neurocognitive patterns dissociating semantic processing from executive control are linked to more detailed off-task mental time travel. *Sci Reports* 2020 101 10:1–14 Available at: <https://www.nature.com/articles/s41598-020-67605-2> [Accessed September 24, 2021].

Wang H-TT, Poerio G, Murphy C, Bzdok D, Jefferies E, Smallwood J (2018) Dimensions of Experience: Exploring the Heterogeneity of the Wandering Mind. *Psychol Sci* 29:56–71 Available at: <http://www.ncbi.nlm.nih.gov/pubmed/29131720> [Accessed July 31, 2018].

Wen X, He H, Dong L, Chen J, Yang J, Guo H, Luo C, Yao D (2020) Alterations of local functional connectivity in lifespan: A resting-state fMRI study. *Brain Behav* 10:e01652 Available at: <https://onlinelibrary.wiley.com/doi/full/10.1002/brb3.1652> [Accessed March 21, 2022].

Worsley KJ, Taylor JE, Carbonell F, Chung MK, Duerden E, Bernhardt B, Lyttelton O, Boucher M, Evans AC (2009) SurfStat: A Matlab toolbox for the statistical analysis of univariate and multivariate surface and volumetric data using linear mixed effects models and random field theory. *Neuroimage* 47 Available at: <http://www.stat.uchicago.edu/~worsley/surfstat/poster.htm> [Accessed August 11, 2021].

Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD, T Y, RA P, TE N, DC VE, TD W (2011) Large-scale automated synthesis of human functional neuroimaging data.

8:665–670 Available at: <https://pubmed.ncbi.nlm.nih.gov/browser.cbs.mpg.de/21706013/> [Accessed August 11, 2021].

Yeo BTT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, Roffman JL, Smoller JW, Zöllei L, Polimeni JR, Fisch B, Liu H, Buckner RL (2011) The organization

of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol* 106:1125–1165 Available at:

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3174820/> [Accessed March 30, 2021].

Young KS (1998) Internet Addiction: The Emergence of a New Clinical Disorder.

CyberPsychology Behav 1:237–244 Available at:

<https://www.liebertpub.com/doi/abs/10.1089/cpb.1998.1.237> [Accessed September 24, 2021].

Zhang J, Kucyi A, Raya J, Nielsen AN, Nomi JS, Damoiseaux JS, Greene DJ, Horovitz SG,

Uddin LQ, Whitfield-Gabrieli S (2021) What have we really learned from functional connectivity in clinical populations? *Neuroimage* 242:118466 Available at:

<https://linkinghub.elsevier.com/retrieve/pii/S1053811921007394> [Accessed August 25, 2021].


Zigmond AS, Snaith RP (1983) The hospital anxiety and depression scale. *Acta Psychiatr*

Scand 67:361–370 Available at: <https://pubmed.ncbi.nlm.nih.gov/6880820/> [Accessed September 24, 2021].


Supplementary tables and figures

Supplementary Table 1. Multidimensional Experience Sampling (MDES) Statements


Dimension	Statement
Positive	“My thoughts were positive”
Negative	“My thoughts were negative.”
Future	“I thought about future events.”
Past	“I thought about past events.”
Myself	“I thought about myself.”
People	“I thought about other people.”
Surroundings	“I thought about my present environment/ surrounding.”
Wakeful	“I was completely awake.”
Images	“My thoughts were in the form of images.”
Words	“My thoughts were in the form of words”
Specific	“My thoughts were more specific than vague”
Intrusive	“My thoughts were intrusive”



Supplementary figure 1. Heatmap showing variable component loadings for the first 5 principal components derived from trait questionnaires



Supplementary figure 2. Heatmap showing variable component loadings for the first 5 principal components derived from MDES



Supplementary figure 3. Scatter-plots showing the relationship between trait “Negative affect”, and “Positive Episodic Social” and “Self-relevant” thought.