

1 **Concentration and quantification of *Tilapia tilapinevirus* from water using a**  
2 **simple iron flocculation coupled with probe-based RT-qPCR**

3

4 Suwimon Taengphu<sup>1</sup>, Pattanapon Kayansamruaj<sup>2</sup>, Yasuhiko Kawato<sup>3</sup>, Jerome Delamare-  
5 Deboutteville<sup>4</sup>, Chadag Vishnumurthy Mohan<sup>4</sup>, Ha Thanh Dong<sup>5\*</sup>, Saengchan Senapin<sup>1,6\*</sup>

6

7 <sup>1</sup> Fish Health Platform, Center of Excellence for Shrimp Molecular Biology and Biotechnology,  
8 (Centex Shrimp), Faculty of Science, Mahidol University, Bangkok, Thailand

9 <sup>2</sup> Center of Excellence in Aquatic Animal Health Management, Faculty of Fisheries, Kasetsart  
10 University, Bangkok, Thailand

11 <sup>3</sup> Nansei Main Station, National Research Institute of Aquaculture, Japan Fisheries Research and  
12 Education Agency, Minami-Ise, Mie 516-0193, Japan

13 <sup>4</sup> WorldFish, Penang, Malaysia

14 <sup>5</sup> School of Environment, Resources and Development, Asian Institute of Technology, Pathum  
15 Thani, Thailand

16 <sup>6</sup> National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and  
17 Technology Development Agency (NSTDA), Pathum Thani, Thailand

18

19 Corresponding Authors:

20 Saengchan Senapin

21 Email address: saengchan@biotec.or.th

22 Ha Thanh Dong

23 Email address: htdong@ait.ac.th

24 **Abstract**

25 *Tilapia tilapinevirus* (also known as tilapia lake virus, TiLV) is an important virus responsible  
26 for die-off of farmed tilapia globally. Detection and quantification of the virus from  
27 environmental DNA/RNA (eDNA/eRNA) using pond water represents a potential, noninvasive  
28 routine approach for pathogen monitoring and early disease forecasting in aquaculture systems.  
29 Here, we report a simple iron flocculation method for viral concentration from water combined  
30 with a newly developed hydrolysis probe quantitative RT-qPCR method for detection and  
31 quantification of TiLV. The RT-qPCR method targeting a conserved region of TiLV genome  
32 segment 9 has a detection limit of 10 viral copies per  $\mu\text{L}$  of template. The method had a 100%  
33 analytical specificity and sensitivity for TiLV. The optimized iron flocculation method was able  
34 to recover  $16.11 \pm 3.3\%$  of virus from water samples spiked with viral cultures. During disease  
35 outbreak cases from an open-caged system and a closed hatchery system, both tilapia and water  
36 samples were collected for detection and quantification of TiLV. The results revealed that TiLV  
37 was detected from both clinically sick fish and asymptomatic fish. Most importantly, the virus  
38 was successfully detected from water samples collected from different locations in the affected  
39 farms e.g. river water samples from affected cages ( $8.50 \times 10^2$  to  $2.79 \times 10^4$  copies/L) and fish-  
40 rearing water samples, sewage, and reservoir ( $4.29 \times 10^2$  to  $3.53 \times 10^3$  copies/L) from affected  
41 and unaffected ponds of the hatchery. In summary, this study suggests that the eRNA detection  
42 system using iron flocculation coupled with probe based-RT-qPCR is feasible for concentration  
43 and quantification of TiLV from water. This approach might be useful for noninvasive  
44 monitoring of TiLV in tilapia aquaculture systems and facilitating appropriate decisions on  
45 biosecurity interventions needed.

46

47 **Introduction**

48 *Tilapia tilapinevirus* (commonly called tilapia lake virus, TiLV) is a novel and only virus in a  
49 new genus *Tilapinevirus* under the family *Amnooviridae* (International Committee on  
50 Taxonomy of Viruses. 2019). Since its first discovery in 2014, the virus had significant impacts  
51 on tilapia aquaculture worldwide (Eyngor et al. 2014; Ferguson et al. 2014; Jansen et al. 2019).  
52 TiLV is an RNA virus with a 10 segmented negative sense single stranded genome of  
53 approximately 10.323 kb in size (Bacharach et al. 2016). Disease caused by TiLV usually results  
54 in cumulative mortality from 20 to 90% (Behera et al. 2018; Dong et al. 2017a; Eyngor et al.

55 2014; Ferguson et al. 2014; Surachetpong et al. 2017). So far, there are 16 countries that  
56 reportedly confirmed detection of TiLV (Jansen et al. 2019; Surachetpong et al. 2020), but a  
57 wider geographical spread has been hypothesized due to active movements of live tilapia with  
58 other countries (Dong et al. 2017b). Waterborne spread of TiLV might also contribute to  
59 pathogen dissemination to new areas as well as transmission to other fish species  
60 (Chiamkunakorn et al. 2019; Eyangor et al. 2014; Jaewwimol et al. 2018; Piamsomboon &  
61 Wongtavatchai 2021). Experimental evidences have already demonstrated that TiLV is both  
62 horizontally and vertically transmitted (Dong et al. 2020; Eyangor et al. 2014; Jaewwimol et al.  
63 2018; Yamkasem et al. 2019).

64 With respect to waterborne transmission of fish pathogens, several studies employed various  
65 viral concentration methods from water for pathogen detection (For example, Haramoto et al.  
66 (2007); Kawato et al. (2016); Minamoto et al. (2009); Nishi et al. (2016)). The concept is one of  
67 the applications of environmental DNA (eDNA) which is nucleic acids extracted from  
68 environmental samples such as water, soil, and feces (Bass et al. 2015; Gomes et al. 2017). The  
69 eDNA gives advantages in disease monitoring, control measure design, risk factor analysis and  
70 studies of viral survival nature (example review in Oidtmann et al. (2018)). The work described  
71 by Kawato et al. (2016) used an iron flocculation method to concentrate red sea bream iridovirus  
72 (RSIV) in a challenge model with Japanese amberjack (*Seriola quinqueradiata*). Results from  
73 that study showed that detection by qPCR of RSIV from fish-rearing water samples peaked more  
74 than five days before fish mortality occurred, suggesting potential benefit of using iron  
75 flocculation method for disease forecast. Others studies used a cation-coated filter method to  
76 detect DNAs of cyprinid herpesvirus 3 (CyHV-3) (also known as koi herpesvirus, KHV) from  
77 concentrated river water samples three to four months before mass mortalities events occurred in  
78 wild carp in Japan (Haramoto et al. 2007; Minamoto et al. 2009). Additionally, the virus was still  
79 detectable in river water for at least three months after the outbreaks (Minamoto et al. 2009).  
80 These findings helped local authorities and farmers to make rapid decisions for emergency  
81 harvest, biosecurity implementation, follow appropriate disinfection procedures and fallowing  
82 periods.

83 Several molecular methods have been developed for detection of TiLV including RT-PCR  
84 (Eyangor et al. 2014), nested and semi-nested PCR (Dong et al. 2017a; Kembou Tsofack et al.  
85 2017; Taengphu et al. 2020), RT-qPCR (Tattiyapong et al. 2018; Waiyamitra et al. 2018), loop-

86 mediated isothermal amplification (LAMP) (Kampeera et al. 2021; Phusantisampan et al. 2019;  
87 Yin et al. 2019) and Nanopore-based PCR amplicon approach (Delamare-Deboutteville et al.  
88 2021). However, all of these methods target fish tissue specimens for diagnosis, none of which  
89 reported any application for TiLV detection from environmental water samples. Previous probe-  
90 based RT-qPCR methods developed to detect TiLV from tilapia clinical samples with detection  
91 limits of  $2.7 \times 10^4$  or ~70,000 copies (Kembou Tsofack et al. 2017; Waiyamitra et al. 2018) might  
92 not be sensitive enough to detect low viral loads of TiLV in environmental water samples. Based  
93 on publicly available TiLV genomic sequence data (Ahsan et al. 2020; Chaput et al. 2020;  
94 Debnath et al. 2020; Pulido et al. 2019; Subramaniam et al. 2019; Thawornwattana et al. 2021),  
95 we developed a new probe-based RT-qPCR assay targeting TiLV genomic segment 9 and  
96 applied to detect TiLV not only from fish tissues but also from environmental RNA (eRNA)  
97 concentrated from water samples. A simple iron flocculation method for concentration of TiLV  
98 from fish-rearing water samples coupled with our new RT-qPCR assay to detect and quantify  
99 TiLV eRNA was described in the present study.

100

## 101 **Materials & Methods**

### 102 **Development of a new probe-based quantitative RT-qPCR method for TiLV**

#### 103 ***Primer & probe design and establishment of PCR conditions***

104 A new hydrolysis probe-based RT-qPCR method was developed and optimized for detection and  
105 quantification of TiLV. Out of the 10 segments of the TiLV genome, segment 9 was reported to  
106 have relatively high identity (97.44 - 99.15%) among various TiLV isolates (Pulido et al. 2019).  
107 Primers and probe were thus designed based on conserved regions of TiLV genome segment 9  
108 following multiple sequence alignments of all available sequences (n=25 or 27) retrieved from  
109 the GenBank database at NCBI as of June 2021 (Fig. S1). Primer Seg9-TaqMan-F (5'-CTA  
110 GAC AAT GTT TTC GAT CCA G-3') had a 100% perfect match with all retrieved 27  
111 sequences while primer Seg9-TaqMan-R (5'-TTC TGT GTC AGT AAT CTT GAC AG-3') and  
112 probe (5'-6-FAM-TGC CGC CGC AGC ACA AGC TCC A-BHQ-1-3') had one mismatch  
113 nucleotide from 25 and 27 available sequences, respectively (Fig. S1). The final composition of  
114 the optimized TiLV RT-qPCR 20  $\mu$ L reaction consists of 1X master mix (qScript XLT 1-Step  
115 RT-qPCR ToughMix Low ROX buffer) (Quanta Bio), 1.5-2  $\mu$ L ( $\leq$ 300 ng) of RNA template, 450  
116 nM of each forward and reverse primers, and 150 nM of Seg9-TaqMan-Probe. Size of the

117 amplified product is expected at 137 bp. Cycling conditions include a reverse transcription step  
118 at 50 °C for 10 min, then an initial denaturation step at 95 °C for 1 min followed by 40 cycles of  
119 95 °C for 10 s and 58 °C for 30 s. RT-qPCR amplification was carried out using Bio-Rad CFX  
120 Connect Real-Time PCR machine. Positive control plasmid (pSeg9-351) was previously  
121 constructed by inserting a 351 bp-TiLV segment 9 open reading frame (ORF) into pGEM T-easy  
122 vector (Promega) as reported earlier (Thawornwattana et al. 2021).

123 ***Analytical specificity and sensitivity tests***

124 Specificity of the Seg9-targeted RT-qPCR was tested with RNA extracted (150 ng/reaction) from  
125 clinically healthy tilapia, 15 common fish bacterial pathogens, and fish tissues infected with  
126 nervous necrosis virus (NNV), infectious spleen and kidney necrosis virus (ISKNV), or scale  
127 drop disease virus (SDDV) (Table S1). Detection limit of the method was investigated using 10-  
128 fold serial dilutions of pSeg9-351 plasmid template from  $10^6$  to 1 copies/ $\mu$ L template. The assays  
129 were performed in duplicate. Calculation of viral copy numbers was performed using standard  
130 curves prepared by plotting the  $\log_{10}$  of serial plasmid dilutions versus quantification cycle (Cq)  
131 values.

132 ***Diagnostic specificity and sensitivity of the assay***

133 We assessed the Seg9 RT-qPCR assay against RNA extracted from 65 samples held in our  
134 laboratory. Forty-four samples originated from known TiLV outbreaks and 21 from known non-  
135 diseased samples (healthy tilapia). Diagnostic test results were obtained using semi-nested RT-  
136 PCR methods as described before (Dong et al. 2017a; Taengphu et al. 2020). Analytical  
137 specificity and sensitivity of the assay were calculated according to formulas described by  
138 Martin (1984) as:

- 139 • Sensitivity % = [number of true positive samples / (number of true positive samples +  
140 number of false negative samples)]  $\times$  100
- 141 • Specificity % = [number of true negative samples / (number of true negative samples +  
142 number of false positive samples)]  $\times$  100

143 **Optimization for viral concentration protocol**

144 ***Virus preparation***

145 Viral stock used in this study was isolated from TiLV-infected Nile tilapia using E-11 cell line, ,  
146 a clone of the cell line SSN-1 derived from whole fry tissue of snakehead fish (Sigma-Aldrich  
147 cat no. 01110916-1VL). The virus was propagated as described in Dong et al. (2020). Briefly,

148 200  $\mu$ L of TiLV stock ( $\sim 10^8$  copies/mL) was added into a 75 mL cell culture flask containing a  
149 monolayer of E-11 cell and 5 mL of L15 medium (Leibovitz), incubated at 25 °C for 5 days. The  
150 culture supernatant containing viral particles was collected after centrifugation at 15,000 x g for  
151 10 min at 4 °C. The viral stock was kept in aliquots of 1 mL at -80 °C until used.

152 ***Iron flocculation***

153 Viral concentration using iron flocculation method was performed using the protocol previously  
154 described by Kawato et al. (2016) with some modifications. Workflow of this method is  
155 illustrated in Fig. 1. Briefly, 100  $\mu$ L ( $\sim 10^7$ - $10^8$  copies) of TiLV viral stock was added into 500  
156 mL of sterile water that contained 1% marine salt and 36  $\mu$ M ferric chloride. The suspension was  
157 stirred at room temperature for 1 h before being mechanically filtered through a 0.4- $\mu$ m pore size  
158 polycarbonate filter (Advantec) with a vacuum pump connected to a filter holder KG-47  
159 (Advantec) under < 15 psi pressure. The flocculate-trapped filters were either directly subjected  
160 to nucleic acid extraction or resuspended with oxalate-EDTA buffer (John et al. 2011) prior to  
161 nucleic acid extraction using Patho Gene-spin DNA/RNA extraction kit (iNtRON). Experiments  
162 were carried out in two to four replicates. Viral concentration, percentage (%) recovery and fold  
163 reduction of the virus copies were calculated from Cq values after flocculation compared to that  
164 of the starting viral stock.

165 **Detection of TiLV from fish and pond water sources during disease outbreaks**

166 During 2020-2021, two disease outbreaks were reported to our laboratory. One occurred in an  
167 open-caged system (juvenile hybrid red tilapia, *Oreochromis* sp.) and the other in a closed  
168 hatchery system (earthen ponds, Nile tilapia, *O. niloticus*). The fish experienced abnormal  
169 mortalities with clinical symptoms of disease resembling those caused by TiLV, e.g. darkened  
170 body (Nile tilapia), pale color and reddish opercula (red hybrid tilapia), abdominal distension,  
171 and exophthalmia. In the first outbreak, we received fish specimens and water samples collected  
172 from four cages namely A, B, C and D with two-three fish and two bottles of 500 mL water  
173 samples from each cage. The samples were kept on ice during transportation and shipped to our  
174 laboratory within 24 h. In the latter outbreak, internal organs from both sick and healthy looking  
175 tilapia from different ponds as well as snails and sludge were collected and preserved in Trizol  
176 reagent (Invitrogen) by a hatchery veterinarian and sent to our laboratory. Water (500 mL/bottle)  
177 from fish ponds, reservoir, and sewage (outgoing waste water from ponds) was also collected  
178 from this hatchery.

179 Fish specimens were subjected to RNA extraction while water samples were centrifuged (5,000 x  
180 g for 5 min) to remove suspended matters before subjected to iron flocculation and subsequent  
181 nucleic acid extraction by Patho Gen-spin column kit. Viral detection and quantification were  
182 then performed to investigate the presence of TiLV by the established Seg 9 RT-qPCR assay  
183 described above. Plasmid template pSeg9-351 was used in a positive control reaction while  
184 nuclease-free water was used for negative control.

185

## 186 **Results**

### 187 **A new probe-based RT-qPCR method for detection and quantification of TiLV**

188 The Seg9 RT-qPCR method developed in this study had a detection limit (sensitivity) of 10  
189 copies/µL template with mean  $C_q \pm SD$  values of the detection limit at  $38.24 \pm 0.09$  (Fig. 2a).  
190 Hence, samples with a  $C_q$  value  $\geq 38.24$  were considered TiLV negative or under the limit of  
191 this detection method. Amplification efficiency (E) of the established RT-qPCR was 94.0% with  
192  $R^2$  of 0.998 (Fig. 2b). Analytical specificity test revealed that the method was highly specific to  
193 TiLV only since no amplifications were found when the method was assayed with RNA  
194 templates extracted from three other viruses, 15 bacterial species, and healthy tilapia (Fig. 2c,  
195 Table S1). The method had 100% diagnostic specificity and 100% diagnostic sensitivity when  
196 assayed with previously diagnosed TiLV infected and non-infected fish samples ( $n = 65$  with  $C_q$   
197 value ranges 13.02 – 34.85) (Table 1).

### 198 **Conditions for viral concentration and percentage recovery**

199 Percentage recovery of TiLV after iron flocculation but without suspension of the membrane  
200 filter in oxalate-EDTA buffer was only  $2.04 \pm 0.5\%$  ( $n=2$ ), which corresponded to a  $50.55 \pm$   
201 12.2-fold reduction in the viral concentration compared to the original viral stock (Table 2). This  
202 was significantly improved with an additional suspension step of the flocculate-trapped filters  
203 into oxalate-EDTA buffer prior to RNA extraction. The percentage recovery of TiLV increased  
204 to  $16.11 \pm 3.3\%$  ( $n=4$ ), which is equivalent to a  $6.38 \pm 1.1$ -fold reduction in viral concentration  
205 after iron flocculation (Table 2). Figure 2d showed representative results of viral quantification  
206 using Seg 9 RT-qPCR assays of TiLV from water after iron flocculation with the resuspension  
207 step.

### 208 **Virus quantification from tilapia and different water sources during disease outbreaks**

209 The results of TiLV detection and quantification from fish tissues and water samples are shown  
210 in Tables 3 and 4. In the first disease outbreak (open-cages), TiLV was detected from both fish  
211 and water samples from all four cages (A-D) (Table 3). Fish samples had Cq values ranging from  
212 12.40 to 36.22, equivalent to  $3.98 \times 10^8$  to  $5.6 \times 10^1$  viral copies/150 ng RNA template,  
213 respectively (Table 3, Fig. 2e). Interestingly, eight water samples collected from four cages had a  
214 similar viral load ranging from  $8.50 \times 10^2$  to  $3.40 \times 10^4$  copies/L (Cq 31.19 - 36.76) (Table 3,  
215 Fig. 2f).

216 In the second disease event (earthen ponds), samples were collected from eight ponds; one had  
217 unusually mortality, five showed no sign of disease, one was a sewage pond and one a reservoir  
218 pond (Table 4). In the affected fingerling pond C1, TiLV was detected from five diseased fish  
219 ( $9.53 \times 10^7$  to  $1.17 \times 10^9$  copies/150 ng RNA template), one asymptomatic fish ( $3.80 \times 10^3$   
220 copies/150 ng RNA template), and water sample from one location ( $8.41 \times 10^3$  copies/L) (Table  
221 4). TiLV was undetectable from snail and sludge samples originating from pond C1. TiLV  
222 investigation from the remaining 7 other ponds revealed that TiLV was also detectable—but in  
223 relatively low viral loads from some asymptomatic fish (both fingerling and brood fish) and  
224 water from culture ponds as well as water from the reservoir and sewage ponds that were  
225 collected during the disease event (Table 4).

226

## 227 **Discussion**

228 Methods to concentrate and recover viral particles from environmental water samples have been  
229 long applied in human health studies especially with waterborne diseases caused by enteric  
230 viruses (example review in Cashdollar & Wymer (2013); Haramoto et al. (2018)). It has later  
231 become an essential process for aquatic environment research (Jacquet et al. 2010). Several  
232 techniques have been used for viral concentration from aquatic environment, including  
233 coagulation/flocculation, filtration/ultrafiltration, and centrifugation/ultracentrifugation  
234 (Cashdollar & Wymer 2013; Ikner et al. 2012). Our present study employed an iron flocculation  
235 method which was initially described for virus removal from freshwater (Chang et al. 1958) and  
236 virus concentration from marine water (John et al. 2011). It was later adapted to detect and  
237 quantify two fish viruses: nervous necrosis virus (NNV) (an RNA virus) and red sea bream  
238 iridovirus (RSIV) (a DNA virus) that were experimentally spiked in fish-rearing water (Kawato  
239 et al. 2016; Nishi et al. 2016). The recovery rate was estimated by qPCR and yielded >50 and

240 >80% for NNV and RSIV, respectively. In this study, while the recovery rate of TiLV (an RNA  
241 virus) from spiked-water was considerably lower ( $16.11 \pm 3.3\%$ ), it is in a similar range of  
242 practical methods used for concentrating and detecting human viruses from water environments  
243 (Haramoto et al. 2018). For example, murine norovirus-1 (MNV-1) used as a viral model in viral  
244 concentration assay of human enteric viruses was recovered from spiked-water at 5.8–21.9%  
245 using the electronegative hydroxyapatite (HA)-filtration combined with polyethylene glycol  
246 (PEG) concentration method. The protocol was then used for detection of human noroviruses  
247 (NoV) and hepatitis A virus (HAV) in all water types (De Keuckelaere et al. 2013). More  
248 recently, researchers used porcine coronavirus (porcine epidemic diarrhea virus, PEDV) and  
249 mengovirus (MgV) as model viruses to concentrate severe acute respiratory syndrome  
250 coronavirus 2 (SARS-CoV-2) from water samples (Randazzo et al. 2020). By using an aluminum  
251 hydroxide adsorption-precipitation concentration method, PEDV and MgV spiked in water were  
252 recovered at 3.3-11.0%. The method can then be applied to detect SARS-CoV-2 RNA in  
253 untreated wastewater samples of  $\sim 10^{5.4}$  genomic copies/L (Randazzo et al. 2020).

254

255 Despite a low recovery rate from water samples in this study, we confirmed the usefulness of the  
256 iron flocculation and RT-qPCR approach to concentrate and determine the concentration of  
257 TiLV from fish-rearing water and other water sources from two aquaculture production systems  
258 during disease outbreaks. The inherent nature of DNA and RNA viruses and their ability to  
259 survive outside their hosts may also contribute to those differences observed in recovery rates  
260 (Cashdollar & Wymer 2013; Pinon & Viallette 2018). Other viral concentration techniques using  
261 different coagulant/flocculant chemicals as well as more efficient RNA extraction methods  
262 should be tested for further improvement of TiLV recovery from water.

263

264 After the viral concentration and recovery process, downstream viral detection methods include  
265 cell culture methods, PCR-based assays, and viral metagenomics analysis (example review in  
266 Haramoto et al. (2018)). Here, we employed RT-qPCR technique for detection and quantification  
267 of TiLV, although the detected amounts did not represent the viral viability. Using all TiLV  
268 genomic sequences publicly available, we designed a new set of conserved primers and probe  
269 targeting the viral genomic segment 9. The newly established RT-qPCR protocol was highly  
270 specific to TiLV and did not cross-amplify RNA extracted from other common bacterial and

271 viral aquatic pathogens. The method is very sensitive as it can detect as low as 10 viral copies  
272 per  $\mu\text{L}$  of template, >2,700 times more sensitive than previous probe-based RT-qPCR methods  
273 (Kembou Tsofack et al. 2017; Waiyamitra et al. 2018), reflecting high specificity of the newly  
274 designed primers and probe. Our RT-qPCR method has 100% diagnostic specificity and  
275 sensitivity in agreement with previous results (n=65) obtained using semi-nested RT-PCR  
276 protocols (Dong et al. 2017a; Taengphu et al. 2020). Increased number of sample sizes with  
277 diverse geographical sources may be required for further investigation. Most importantly, this  
278 new Seg 9 RT-qPCR assay was able to detect and quantify TiLV load from various types of field  
279 samples, including clinically sick fish, asymptomatic fish, and water samples, as opposed to  
280 other molecular diagnostic methods optimized solely for fish specimens.

281

282 The viral loads from water samples collected during the two disease events were approximately  
283  $\sim 10^3$  viral copies/L (earthen ponds) and  $\sim 10^4$  viral copies/L (open-cages), but in reality, these  
284 concentrations might be significantly higher due to substantial losses during the concentration  
285 and recovery process. Higher viral loads observed in some of the water samples collected during  
286 the disease outbreak were probably due to active shedding of the virus from diseased fish into the  
287 environment, and might be an additional evidence of the waterborne transmission nature of TiLV  
288 reported previously (Eyngor et al. 2014; Yamkasem et al. 2019). Potential application for TiLV  
289 outbreak forecasting should be further investigated by experimental infection to monitor viral  
290 loads in water in relation to fish morbidity and mortality as previously described for other fish  
291 pathogens (Haramoto et al. 2007; Kawato et al. 2016; Minamoto et al. 2009; Nishi et al. 2016).

292

## 293 **Conclusions**

294 In summary, the viral concentration method by iron flocculation used in concert with a newly  
295 developed probe-based RT-qPCR was not only successful for detection and quantification of  
296 TiLV from water in diseased pond/cages, but also from unaffected ponds, reservoir, and sewage  
297 water. This method, apart from its potential practical use for future monitoring programs of TiLV  
298 viral load in water samples from various culturing units, our approach could become useful to  
299 detect possible TiLV contamination from incoming and outgoing waste water as well as to test  
300 the systems after disinfection treatments. Such application will support health professionals and

301 farmers to design appropriate biosecurity interventions to reduce the loss caused by TiLV in  
302 tilapia farms and hatcheries.

303

304 **Acknowledgements**

305 This study was financially funded by the CGIAR Research Program on Fish Agri-Food Systems  
306 (FISH) led by WorldFish. The authors would like to thank K. Pimsannil, W. Meemetta and Ms.  
307 Thu Thao Mai for their skilled technical assistance.

308 **References**

309 Ahasan MS, Keleher W, Giray C, Perry B, Surachetpong W, Nicholson P, Al-Hussinee L,  
310 Subramaniam K, and Waltzek TB. 2020. Genomic Characterization of Tilapia Lake Virus  
311 Isolates Recovered from Moribund Nile Tilapia (*Oreochromis niloticus*) on a Farm in the  
312 United States. *Microbiol Resour Announc* 9:e01368-01319. 10.1128/MRA.01368-19

313 Bacharach E, Mishra N, Briese T, Zody MC, Kembou Tsofack JE, Zamostiano R, Berkowitz A,  
314 Ng J, Nitido A, Corvelo A, Toussaint NC, Abel Nielsen SC, Hornig M, Del Pozo J,  
315 Bloom T, Ferguson H, Eldar A, and Lipkin WI. 2016. Characterization of a Novel  
316 Orthomyxo-like Virus Causing Mass Die-Offs of Tilapia. *mBio* 7:e00431-00416.  
317 10.1128/mBio.00431-16

318 Bass D, Stentiford GD, Littlewood DTJ, and Hartikainen H. 2015. Diverse Applications of  
319 Environmental DNA Methods in Parasitology. *Trends Parasitol* 31:499-513.  
320 10.1016/j.pt.2015.06.013

321 Behera BK, Pradhan PK, Swaminathan TR, Sood N, Paria P, Das A, Verma DK, Kumar R,  
322 Yadav MK, Dev AK, Parida PK, Das BK, Lal KK, and Jena JK. 2018. Emergence of  
323 Tilapia Lake Virus associated with mortalities of farmed Nile Tilapia *Oreochromis*  
324 *niloticus* (Linnaeus 1758) in India. *Aquac Res* 484:168-174.  
325 10.1016/j.aquaculture.2017.11.025

326 Cashdollar JL, and Wymer L. 2013. Methods for primary concentration of viruses from water  
327 samples: a review and meta-analysis of recent studies. *J Appl Microbiol* 115:1-11.  
328 10.1111/jam.12143

329 Chang SL, Stevenson RE, Bryant AR, Woodward RL, and Kabler PW. 1958. Removal of  
330 Coxsackie and bacterial viruses in water by flocculation. II. Removal of Coxsackie and  
331 bacterial viruses and the native bacteria in raw Ohio River water by flocculation with  
332 aluminum sulfate and ferric chloride. *Am J Public Health*  
333 48:159-169. 10.2105/ajph.48.2.159.

334 Chaput DL, Bass D, Alam MM, Hasan NA, Stentiford GD, Aerle RV, Moore K, Bignell JP,  
335 Haque MM, and Tyler CR. 2020. The Segment Matters: Probable Reassortment of  
336 Tilapia Lake Virus (TiLV) Complicates Phylogenetic Analysis and Inference of  
337 Geographical Origin of New Isolate from Bangladesh. *Viruses* 12:258.  
338 10.3390/v12030258

339 Chiamkunakorn C, Machimbirike VI, Senapin S, Khunrae P, Dong HT, and Rattanarojpong T.  
340 2019. Blood and liver biopsy for the non-destructive screening of tilapia lake virus. *J  
341 Fish Dis* 42:1629-1636. 10.1111/jfd.13076

342 De Keuckelaere A, Baert L, Duarte A, Stals A, and Uyttendaele M. 2013. Evaluation of viral  
343 concentration methods from irrigation and processing water. *J Virol Methods* 187:294-  
344 303. 10.1016/j.jviromet.2012.11.028

345 Debnath PP, Delamare-Deboutteville J, Jansen MD, Phiwsaiya K, Dalia A, Hasan MA, Senapin  
346 S, Mohan CV, Dong HT, and Rodkhum C. 2020. Two-year surveillance of tilapia lake  
347 virus (TiLV) reveals its wide circulation in tilapia farms and hatcheries from multiple  
348 districts of Bangladesh. *J Fish Dis* 43:1381-1389. 10.1111/jfd.13235

349 Delamare-Deboutteville J, Taengphu S, Gan HM, Kayansamruaj P, Debnath PP, Barnes A,  
350 Wilkinson S, Kawasaki M, Vishnumurthy Mohan C, Senapin S, and Dong HT. 2021.  
351 Rapid genotyping of tilapia lake virus (TiLV) using Nanopore sequencing. *J Fish Dis*:1-  
352 12. 10.1111/jfd.13467

353 Dong HT, Ataguba GA, Khunrae P, Rattanarojpong T, and Senapin S. 2017b. Evidence of TiLV  
354 infection in tilapia hatcheries from 2012 to 2017 reveals probable global spread of the  
355 disease. *Aquac Res* 479:579-583. 10.1016/j.aquaculture.2017.06.035

356 Dong HT, Senapin S, Gangnonngiw W, Nguyen VV, Rodkhum C, Debnath PP, Delamare-  
357 Deboutteville J, and Mohan CV. 2020. Experimental infection reveals transmission of  
358 tilapia lake virus (TiLV) from tilapia broodstock to their reproductive organs and  
359 fertilized eggs. *Aquac Res* 515. 10.1016/j.aquaculture.2019.734541

360 Dong HT, Siriroob S, Meemetta W, Santimanawong W, Gangnonngiw W, Pirarat N, Khunrae P,  
361 Rattanarojpong T, Vanichviriyakit R, and Senapin S. 2017a. Emergence of tilapia lake  
362 virus in Thailand and an alternative semi-nested RT-PCR for detection. *Aquac Res*  
363 476:111-118. 10.1016/j.aquaculture.2017.04.019

364 Eyangor M, Zamostiano R, Kembou Tsofack JE, Berkowitz A, Bercovier H, Tinman S, Lev M,  
365 Hurvitz A, Galeotti M, Bacharach E, and Eldar A. 2014. Identification of a novel RNA  
366 virus lethal to tilapia. *J Clin Microbiol* 52:4137-4146. 10.1128/JCM.00827-14

367 Ferguson HW, Kabuusu R, Beltran S, Reyes E, Lince JA, and del Pozo J. 2014. Syncytial  
368 hepatitis of farmed tilapia, *Oreochromis niloticus* (L.): a case report. *J Fish Dis* 37:583-  
369 589. 10.1111/jfd.12142

370 Gomes GB, Hutson KS, Domingos JA, Chung C, Hayward S, Miller TL, and Jerry DR. 2017.  
371 Use of environmental DNA (eDNA) and water quality data to predict protozoan parasites  
372 outbreaks in fish farms. *Aquac Res* 479:467-473. 10.1016/j.aquaculture.2017.06.021

373 Haramoto E, Kitajima M, Hata A, Torrey JR, Masago Y, Sano D, and Katayama H. 2018. A  
374 review on recent progress in the detection methods and prevalence of human enteric  
375 viruses in water. *Water Res* 135:168-186. 10.1016/j.watres.2018.02.004

376 Haramoto E, Kitajima M, Katayama H, and Ohgaki S. 2007. Detection of koi herpesvirus DNA  
377 in river water in Japan. *J Fish Dis* 30:59-61. 10.1111/j.1365-2761.2007.00778.x.

378 Ikner LA, Gerba CP, and Bright KR. 2012. Concentration and recovery of viruses from water: a  
379 comprehensive review. *Food Environ Virol* 4:41-67. 10.1007/s12560-012-9080-2

380 International Committee on Taxonomy of Viruses. 2019. Taxonomy, Virus Taxonomy: 2018b  
381 Release. Available at <https://talk.ictvonline.org/taxonomy/>.

382 Jacquet S, Miki T, Noble R, Peduzzi P, and Wilhelm S. 2010. Viruses in aquatic ecosystems:  
383 important advancements of the last 20 years and prospects for the future in the field of  
384 microbial oceanography and limnology. *Adv Oceanogr Limnol* 1:97-141.  
385 10.1080/19475721003743843

386 Jaemwimol P, Rawiwan P, Tattiayapong P, Saengnual P, Kamlangdee A, and Surachetpong W.  
387 2018. Susceptibility of important warm water fish species to tilapia lake virus (TiLV)  
388 infection. *Aquac Res* 497:462-468. 10.1016/j.aquaculture.2018.08.028

389 Jansen MD, Dong HT, and Mohan CV. 2019. Tilapia lake virus: a threat to the global tilapia  
390 industry? *Rev Aquacult* 11:725-739. 10.1111/raq.12254

391 John SG, Mendez CB, Deng L, Poulos B, Kauffman AK, Kern S, Brum J, Polz MF, Boyle EA,  
392 and Sullivan MB. 2011. A simple and efficient method for concentration of ocean viruses  
393 by chemical flocculation. *Env Microbiol Rep* 3:195-202. 10.1111/j.1758-  
394 2229.2010.00208.x

395 Kampeera J, Dangtip S, Suvannakad R, Khumwan P, Senapin S, and Kiatpathomchai W. 2021.  
396 Reverse transcription loop-mediated isothermal amplification (RT-LAMP) combined  
397 with colorimetric gold nanoparticle (AuNP) probe assay for visual detection of tilapia  
398 lake virus (TiLV) in Nile and red hybrid tilapia. *J Fish Dis* 00:1-13. 10.1111/jfd.13482

399 Kawato Y, Ito T, Kamaishi T, Fujiwara A, Ototake M, Nakai T, and Nakajima K. 2016.  
400 Development of red sea bream iridovirus concentration method in seawater by iron  
401 flocculation. *Aquac Res* 450:308-312. 10.1016/j.aquaculture.2015.08.016

402 Kembou Tsofack JE, Zamostiano R, Watted S, Berkowitz A, Rosenbluth E, Mishra N, Briese T,  
403 Lipkin WI, Kabuusu RM, Ferguson H, Del Pozo J, Eldar A, and Bacharach E. 2017.  
404 Detection of Tilapia Lake Virus in Clinical Samples by Culturing and Nested Reverse  
405 Transcription-PCR. *J Clin Microbiol* 55:759-767. 10.1128/JCM.01808-16

406 Martin SW. 1984. Estimating disease prevalence and the interpretation of screening. *Prev Vet  
407 Med* 2:463-472. 10.1016/0167-5877(84)90091-6

408 Minamoto T, Honjo MN, Uchii K, Yamanaka H, Suzuki AA, Kohmatsu Y, Iida T, and Kawabata  
409 Z. 2009. Detection of cyprinid herpesvirus 3 DNA in river water during and after an  
410 outbreak. *Vet Microbiol* 135:261-266. 10.1016/j.vetmic.2008.09.081

411 Nishi S, Yamashita H, Kawato Y, and Nakai T. 2016. Cell Culture Isolation of Piscine  
412 Nodavirus (Betanodavirus) in Fish-Rearing Seawater. *Appl Environ Microbiol* 82:2537-  
413 2544. 10.1128/AEM.03834-15

414 Oidtmann B, Dixon P, Way K, Joiner C, and Bayley AE. 2018. Risk of waterborne virus spread -  
415 review of survival of relevant fish and crustacean viruses in the aquatic environment and  
416 implications for control measures. *Rev Aquacult* 10:641-669. 10.1111/raq.12192

417 Phusantisampan T, Tattiyapong P, Mutrakulcharoen P, Sriariyanun M, and Surachetpong W.  
418 2019. Rapid detection of tilapia lake virus using a one-step reverse transcription loop-  
419 mediated isothermal amplification assay. *Aquac Res*  
420 507:35-39. 10.1016/j.aquaculture.2019.04.015

421 Piamsomboon P, and Wongtavatchai J. 2021. Detection of Tilapia Lake Virus (TiLV) in Healthy  
422 Fish from the Pre-Existing Disease Environment Using Different RT-PCR Methods. *Turk  
423 J Fish Aquat Sci* 21:205-209. 10.4194/1303-2712-v21\_4\_05

424 Pinon A, and Viallette M. 2018. Survival of Viruses in Water. *Intervirology* 61:214-222.  
425 10.1159/000484899

426 Pulido LLH, Mora CM, Hung AL, Dong HT, and Senapin S. 2019. Tilapia lake virus (TiLV)  
427 from Peru is genetically close to the Israeli isolates. *Aquac Res* 510:61-65.  
428 10.1016/j.aquaculture.2019.04.058

429 Randazzo W, Truchado P, Cuevas-Ferrando E, Simon P, Allende A, and Sanchez G. 2020.  
430 SARS-CoV-2 RNA in wastewater anticipated COVID-19 occurrence in a low prevalence  
431 area. *Water Res* 181:115942. 10.1016/j.watres.2020.115942

432 Subramaniam K, Ferguson HW, Kabuusu R, and Waltzek TB. 2019. Genome sequence of tilapia  
433 lake virus associated with syncytial hepatitis of tilapia in an Ecuadorian aquaculture  
434 facility. *Microbiol Resour Announc* 8:e00084-00019. 10.1128/MRA.00084-19

435 Surachetpong W, Janetanakit T, Nonthabenjawan N, Tattiyapong P, Sirikanchana K, and  
436 Amonsin A. 2017. Outbreaks of Tilapia Lake Virus Infection, Thailand, 2015-2016.  
437 *Emerg Infect Dis* 23:1031-1033. 10.3201/eid2306.161278

438 Surachetpong W, Roy SRK, and Nicholson P. 2020. Tilapia lake virus: The story so far. *J Fish  
439 Dis* 43:1115-1132. 10.1111/jfd.13237

440 Taengphu S, Sangsuriya P, Phiwsaiya K, Debnath PP, Delamare-Deboutteville J, Mohan CV,  
441 Dong HT, and Senapin S. 2020. Genetic diversity of tilapia lake virus genome segment 1  
442 from 2011 to 2019 and a newly validated semi-nested RT-PCR method. *Aquac Res* 526.  
443 10.1016/j.aquaculture.2020.735423

444 Tattiayapong P, Sirikanchana K, and Surachetpong W. 2018. Development and validation of a  
445 reverse transcription quantitative polymerase chain reaction for tilapia lake virus  
446 detection in clinical samples and experimentally challenged fish. *J Fish Dis* 41:255-261.  
447 10.1111/jfd.12708

448 Thawornwattana Y, Dong HT, Phiwsaiya K, Sangsuriya P, Senapin S, and Aiewsakun P. 2021.  
449 Tilapia lake virus (TiLV): Genomic epidemiology and its early origin. *Transbound*  
450 *Emerg Dis* 68:435-444. 10.1111/tbed.13693

451 Waiyamitra P, Tattiayapong P, Sirikanchana K, Mongkolsuk S, Nicholson P, and Surachetpong  
452 W. 2018. A TaqMan RT-qPCR assay for tilapia lake virus (TiLV) detection in tilapia.  
453 *Aquac Res* 497:184-188. 10.1016/j.aquaculture.2018.07.060

454 Yamkasem J, Tattiayapong P, Kamlangdee A, and Surachetpong W. 2019. Evidence of potential  
455 vertical transmission of tilapia lake virus. *J Fish Dis* 42:1293-1300. 10.1111/jfd.13050

456 Yin J, Wang Q, Wang Y, Li Y, Zeng W, Wu J, Ren Y, Tang Y, Gao C, Hu H, and Bergmann  
457 SM. 2019. Development of a simple and rapid reverse transcription-loopmediated  
458 isothermal amplification (RT-LAMP) assay for sensitive detection of tilapia lake virus. *J*  
459 *Fish Dis* 42:817-824. 10.1111/jfd.12983

460 **Tables and Figures**

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486


487



488 **Figure 1:** Workflow of TiLV flocculation, concentration and quantification used in this study.

489 An iron flocculation method was used to concentrate viruses from water (a). The water  
490 suspension containing the virus was filtered through a 0.4- $\mu$ m pore size polycarbonate membrane  
491 filter with a vacuum pressure pump (b-c). The flocculate-trapped filter (d) was then resuspended  
492 in oxalate-EDTA buffer (e) prior to nucleic acid extraction (f) and TiLV quantification (g).

493  
494  
495  
496  
497  
498  
499



500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511

512 **Figure 2:** Performance of the newly established probe-based RT-qPCR detection of TiLV  
513 genomic segment 9. a) Analytical sensitivity assay determined using serial dilutions of plasmid  
514 DNA containing a 351-bp TiLV segment 9 insert. Amplification results were from two technical  
515 replicate tests. b) A standard curve was derived from the assays in (a) showing an amplification  
516 efficiency (E) of 94.0%. c) Analytical specificity test of the RT-qPCR protocol against RNAs  
517 extracted from common pathogens of fish and healthy looking tilapia as listed in Table S1. d)  
518 TiLV quantification from template extracted from stock virus (S) and flocculate-trapped filters  
519 (F) with resuspension step using two replicates. e) TiLV quantification from fish samples  
520 collected from an outbreak open cage. f) TiLV quantification from water samples collected from  
521 an outbreak open cage. P, positive control; N, no template control; RFU, relative fluorescence  
522 units.

523 **Table 1:** Diagnostic specificity and sensitivity of the Seg9 probe-based RT-qPCR method

Test results	Diseased samples (n=44)	Non-diseased samples (n=21)
Positive (+)	True positive 44	False positive 0
Negative (-)	False negative 0	True negative 21
Diagnostic sensitivity (%)	100	
Diagnostic specificity (%)	100	

524

**Table 2:** Percentage (%) recovery of viruses from water using different conditions

Sample type	Before and after flocculation	Suspension step	Total viral copy number	% recovery	Fold reduction
Water spiked with TiLV culture	Before (viral stock)		$3.92 \times 10^8$		
	After (Rep.1)	No	$9.34 \times 10^6$	2.38	41.93
	After (Rep.2)	No	$6.62 \times 10^6$	1.69	59.18
	<b>Mean <math>\pm</math> SD</b>			<b>2.04 <math>\pm</math> 0.5</b>	<b>50.55 <math>\pm</math> 12.2</b>
	Before (viral stock 1)		$1.27 \times 10^8$		
	After (Rep.1)	Yes	$2.67 \times 10^7$	21.08	4.74
	Before (viral stock 2)		$3.21 \times 10^7$		
	After (Rep.2)	Yes	$4.67 \times 10^6$	14.55	6.87
	Before (viral stock 3)*		$4.16 \times 10^7$		
	After (Rep.3)*	Yes	$5.85 \times 10^6$	14.07	7.10
	Before (viral stock 4)*		$3.07 \times 10^7$		
	After (Rep.4)*	Yes	$4.52 \times 10^6$	14.74	6.78
	<b>Mean <math>\pm</math> SD</b>			<b>16.11 <math>\pm</math> 3.3</b>	<b>6.38 <math>\pm</math> 1.1</b>

Rep, replicate; \* denotes experiments where qPCR results were shown in Fig. 2d.

**Table 3:** Quantification of TiLV from fish and water during an outbreak in open-cages

Cage	Samples	Cq	TiLV load*	Interpretation
A	Diseased fish A1-1 (liver + spleen)	13.02	$2.64 \times 10^8$	+
	Diseased fish A1-2 (liver + spleen)	30.69	$2.18 \times 10^3$	+
	Diseased fish A1-3 (liver + spleen)	13.11	$2.49 \times 10^8$	+
	Water sample A1	36.76	$8.50 \times 10^2$	+
B	Water sample A2	31.95	$2.06 \times 10^4$	+
	Diseased fish B1-1 (liver + spleen)	14.35	$1.10 \times 10^8$	+
	Diseased fish B1-2 (liver + spleen)	17.49	$1.37 \times 10^7$	+
	Diseased fish B1-3 (liver + spleen)	13.13	$2.46 \times 10^8$	+
C	Water sample B1	32.54	$1.39 \times 10^4$	+
	Water sample B2	31.60	$2.59 \times 10^4$	+
	Diseased fish C1-1 (liver + spleen)	14.76	$8.34 \times 10^7$	+
	Diseased fish C1-2 (liver + spleen)	13.87	$1.50 \times 10^8$	+
D	Water sample C1	32.71	$1.24 \times 10^4$	+
	Water sample C2	31.49	$2.79 \times 10^4$	+
	Diseased fish D1-1 (liver + spleen)	36.22	$5.6 \times 10^1$	+
	Diseased fish D1-2 (liver + spleen)	12.40	$3.98 \times 10^8$	+
D	Diseased fish D1-3 (liver + spleen)	18.67	$6.26 \times 10^6$	+
	Water sample D1	35.90	$1.50 \times 10^4$	+
	Water sample D2	31.19	$3.40 \times 10^4$	+


Gray highlights water samples; \*viral copy (per reaction for 150 ng fish extracted RNA & per L of water sample); +, detected.

**Table 4:** Quantification of TiLV from fish and pond water during an outbreak in earthen closed-ponds

Pond		Samples	Cq	TiLV load*	Interpretation
Fingerling pond C1 (affected pond)	Fish	Diseased F1 (liver + spleen)	12.42	$3.93 \times 10^8$	+
		Diseased F2 (liver + spleen)	14.56	$9.53 \times 10^7$	+
		Diseased F3 (liver + spleen)	12.11	$4.83 \times 10^8$	+
		Diseased F4 (liver + spleen)	10.77	$1.17 \times 10^9$	+
		Diseased F5 (liver)	13.46	$4.17 \times 10^8$	+
	Water	Normal looking (whole fish)	29.85	$3.80 \times 10^3$	+
		Location 1	39.73	-	-
	Snail Sludge	Location 2	33.30	$8.41 \times 10^3$	+
		Pooled sample	-	-	-
		Pooled sample 1	-	-	-
		Pooled sample 2	-	-	-
Fingerling pond C2	Fish	Normal looking F1 (whole fish)	-	-	-
		Normal looking F2 (whole fish)	32.88	$5.11 \times 10^2$	+
	Water	Location 1	34.66	$3.42 \times 10^3$	+
		Location 2	39.76	-	-
Fingerling C3	Fish	Normal looking F1 (whole fish)	37.34	$2.6 \times 10^1$	
		Normal looking F2 (whole fish)	-	-	-
	Water	Location 1	-	-	-
		Location 2	-	-	-
Broodstock pond B1	Fish	Female brood 1, normal looking <sup>#</sup>	37.08	$3.10 \times 10^1$	
		Female brood 2, normal looking <sup>#</sup>	35.42	$9.50 \times 10^1$	
		Male brood 1, normal looking <sup>#</sup>	38.28	-	-
		Male brood 2, normal looking <sup>#</sup>	36.18	$5.70 \times 10^1$	
Broodstock pond B2	Water	Location 1	37.79	$4.29 \times 10^2$	+
		Location 1	-	-	-
Broodstock pond B3	Water	Location 2	-	-	-
		Location 1	-	-	-
Sewage	Water	Location 2	-	-	-
		Location 1	34.61	$3.53 \times 10^3$	+
Reservoir	Water	Location 2	-	-	-
		Location 1	-	-	-
		Location 2	37.78	$4.32 \times 10^2$	+

Gray highlights water samples; \*viral copy (per reaction for 150 ng fish extracted RNA & per L of water sample); #, liver, kidney, spleen, gill, gonad; -, not detected; +, detected; C2, C3, B1-B3 apparently healthy ponds with no signs of disease

## Supplementary data



**Figure S1.** Nucleotide sequence alignments of TiLV segment 9 sequences (n=27) retrieved from the GenBank database at NCBI. Accession numbers and viral isolate names of all 27 sequences are shown on the left panel. Position of primers and probe used in the newly developed RT-qPCR assay are marked. Numbers denote nucleotide positions to the putative coding region.

**Table S1:** Sample used for evaluation of analytical specificity and sensitivity of the probe-based RT-qPCR method

Samples	Host	Sample type	RT-qPCR result (Cq)	ND, not dete ctab le
Clinically healthy tilapia	NA	RNA	ND	
NNV-infected tissue	Grouper	RNA	ND	
ISKNV-infected tissue	Asian sea bass	RNA	ND	
SDDV-infected tissue	Asian sea bass	RNA	ND	
<i>Streptococcus agalactiae</i>	Nile tilapia	RNA	ND	
<i>Streptococcus iniae</i>	Asian sea bass	RNA	ND	
<i>Edwardsiella ictaluri</i>	Striped catfish	RNA	ND	
<i>Edwardsiella tarda</i>	Nile tilapia	RNA	ND	
<i>Flavobacterium columnare</i>	Asian sea bass	RNA	ND	
<i>Francisella orientalis</i>	Hybrid red tilapia	RNA	ND	
<i>Aeromonas hydrophila</i>	Tilapia	RNA	ND	
<i>Aeromonas veronii</i>	Nile tilapia	RNA	ND	
<i>Aeromonas dhakensis</i>	Hybrid red tilapia	RNA	ND	
<i>Aeromonas caviae</i>	Nile tilapia	RNA	ND	
<i>Aeromonas jandaei</i>	Nile tilapia	RNA	ND	
<i>Plesiomonas shigelloides</i>	Nile tilapia	RNA	ND	
<i>Chryseobacterium</i> sp.	Nile tilapia	RNA	ND	
<i>Vogesella</i> sp.	Nile tilapia	RNA	ND	
<i>Vibrio cholerae</i>	Nile tilapia	RNA	ND	