

An ancient truncated duplication of the anti-Mullerian hormone receptor type 2 gene is a potential conserved master sex determinant in the Pangasiidae catfish family

Short running title: An ancient conserved sex determinant in Pangasiids

Ming Wen^{1,2}, Qiaowei Pan^{2,3}, Elodie Jouanno², Jerome Montfort², Margot Zahm⁴, Cédric Cabau⁵, Christophe Klopp^{4,5}, Carole Iampietro⁶, Céline Roques⁶, Olivier Bouchez⁶, Adrien Castinel⁶, Cécile Donnadieu⁶, Hugues Parrinello⁷, Charles Poncet⁸, Elodie Belmonte⁸, Véronique Gautier⁸, Jean-Christophe Avarre⁹, Remi Dugue⁹, Rudhy Gustiano¹⁰, Trần Thị Thúy Hà¹¹, Marc Campet¹², Kednapat Sriphairoj¹³, Josiane Ribolli¹⁴, Fernanda L. de Almeida¹⁵, Thomas Desvignes¹⁶, John H., Postlethwait¹⁶, Christabel Floi Bucac^{3,17}, Marc Robinson-Rechavi^{3,17}, Julien Bobe², Amaury Herpin², Yann Guiguen^{2*}

AFFILIATIONS

¹ State Key Laboratory of Developmental Biology of Freshwater Fish, College of Life Science, Hunan Normal University, Changsha, China.

² INRAE, LPGP, 35000 Rennes, France.

³ Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland.

⁴ Plate-forme bio-informatique Genotoul, Mathématiques et Informatique Appliquées de Toulouse, INRAE, Castanet Tolosan, France.

⁵ SIGENAE, GenPhySE, Université de Toulouse, INRAE, ENVT, Castanet Tolosan, France.

⁶ INRAE, US 1426, GeT-PlaGe, Genotoul, Castanet-Tolosan, France.

⁷ Montpellier GenomiX (MGX), c/o Institut de Génomique Fonctionnelle, 141 rue de la Cardonille, 34094, Montpellier Cedex 05, France.

⁸ GDEC Gentyane, INRAE, Université Clermont Auvergne, Clermont-Ferrand, France.

⁹ ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France.

¹⁰ Research Institute of Freshwater Fisheries (CRIFI-RIFF), Instalasi Penelitian Perikanan Air Tawar, Jalan Ragunan-Pasar Minggu, P.O. Box 7220/jkspm, Jakarta 12540, Indonesia.

¹¹ Research Institute for Aquaculture No.1. Dinh Bang, Tu Son, Bac Ninh, Viet Nam.

¹² Neovia Asia, HCM city Vietnam.

¹³ Faculty of Natural Resources and Agro-Industry, Kasetsart University Chalermphrakiat Sakon Nakhon Province Campus, Sakon Nakhon, Thailand.

34 ¹⁴ Laboratório de Biologia e Cultivo de Peixes de Água Doce, Universidade Federal de Santa
35 Catarina, Florianópolis, SC, Brasil.

36 ¹⁵ Embrapa Amazônia Ocidental, Manaus, Amazonas, Brasil.

37 ¹⁶ Institute of Neuroscience, University of Oregon, Eugene OR 97403, USA.

38 ¹⁷ SIB Swiss Institute of Bioinformatics, 1015 Lausanne, Switzerland.

39

40 * Corresponding author: Yann Guiguen (yann.guiguen@inrae.fr)

41

42 **Key words:** Pangasiid catfishes, *amhr2*, sex determination, male genome assembly, evolution

43

ABSTRACT

The evolution of sex determination (SD) mechanisms in teleost fishes is amazingly dynamic, as reflected by the variety of different master sex-determining genes identified, even sometimes among closely related species. Pangasiids are a group of economically important catfishes in many South-Asian countries, but little is known about their sex determination system. Here, we generated novel genomic resources for 12 Pangasiid species and provided a first characterization of their SD system. Based on an Oxford Nanopore long-read chromosome-scale high quality genome assembly of the striped catfish *Pangasianodon hypophthalmus*, we identified a duplication of the anti-Müllerian hormone receptor type II gene (*amhr2*), which was further characterized as being sex-linked in males and expressed only in testicular samples. These first results point to a male-specific duplication on the Y chromosome (*amhr2by*) of the autosomal *amhr2a*. Sequence annotation revealed that the *P. hypophthalmus* *Amhr2by* is truncated in its N-terminal domain, lacking the cysteine-rich extracellular part of the receptor that is crucial for ligand binding, suggesting a potential route for its neofunctionalization. Short-read genome sequencing and reference-guided assembly of 11 additional Pangasiid species, along with sex-linkage studies, revealed that this truncated *amhr2by* duplication is also conserved as a male-specific gene in many Pangasiids. Reconstructions of the *amhr2* phylogeny suggested that *amhr2by* arose from an ancient duplication / insertion event at the root of the Siluroidei radiation that is dated around 100 million years ago. Altogether these results bring multiple lines of evidence supporting that *amhr2by* is an ancient and conserved master sex-determining gene in Pangasiid catfishes, a finding that highlights the recurrent usage of the transforming growth factor β pathway in teleost sex determination and brings another empirical case towards the understanding of the dynamics or stability of sex determination systems.

INTRODUCTION

Catfishes (Order Siluriformes) with approximately 4,000 species (Sullivan, Lundberg, & Hardman, 2006) are economically and ecologically important fish worldwide. Among catfishes, the Pangasiid family (Pangasiidae) is recognized as a monophyletic group including four extant genera, i.e., *Helicophagus*, *Pangasianodon*, *Pangasius* and *Pteropangasius* (Pouyaud, Gustiano, & Teugels, 2016). These species have a wide range of habitats both in fresh and brackish water across southern Asia, from Pakistan to Borneo (Roberts & Vidthayanon, 1991). Many Pangasiids, because of their rapid growth rate, are also important aquaculture species, such as *Pangasius bocourti*, *Pangasius*

77 *djambal* and *Pangasianodon hypophthalmus* (Lazard, Cacot, Slembrouck, & Legendre, 2009). *P.*
78 *hypophthalmus*, for example, has become a major aquaculture species extensively farmed in many
79 Asian countries (Anka, Faruk, Hasan, & Azad, 2014; Na-Nakorn & Moeikum, 2009, p.; Phuong &
80 Oanh, 2010; Singh & Lakra, 2012) and has even been recently introduced in the Brazilian finfish
81 aquaculture.

82 Sex determination (SD) mechanisms have not been investigated in detail in Pangasiid catfishes, but
83 genetic sex-linked markers that could facilitate broodstock management for aquaculture or
84 conservation purposes, have been searched without success in both *P. hypophthalmus* and *P. gigas*
85 (Sriphairoj, Na-Nakorn, Brunelli, & Thorgaard, 2007). SD in vertebrates can rely on genetic (GSD
86 for genetic SD), environmental (ESD for environmental SD) or a combination of both genetic and
87 environmental factors (such as thermal effects on GSD = GSD+TE) (Baroiller, D'Cotta, & Saillant,
88 2009; Kobayashi, Nagahama, & Nakamura, 2013; Ospina-Alvarez & Piferrer, 2008). In teleost fishes,
89 SD has been found to be extremely plastic with both GSD, ESD and GSD+TE systems. In addition,
90 teleosts exhibit a wide range of GSD systems, with both classical male (XX/XY) and female
91 heterogamety (ZZ/ZW), but also more complex GSD systems relying on polygenic SD with or
92 without multiple sex chromosomes (Devlin & Nagahama, 2002; Mank & Avise, 2009; Moore &
93 Roberts, 2013). These transitions or turnovers of different GSD systems have been found in closely
94 related species belonging to the same genus (Takehana, Hamaguchi, & Sakaizumi, 2008) and even
95 across populations of the same species (Kallman, 1973). A similar high turnover has also been found
96 for master sex determining (MSD) genes at the top of the genetic sex determination cascade (Matsuda
97 et al., 2002; Myosho et al., 2012; Nanda et al., 2002; Q. Pan et al., 2016; Takehana et al., 2014). Many
98 of these fish MSD genes belong to the “usual suspect” category (Herpin & Scharl, 2015) because
99 they derived from key genes regulating the gonadal sex differentiation network. These “usual
100 suspect” MSD genes currently belong to a few gene families, like the *Dmrt* (Chen et al., 2014;
101 Matsuda et al., 2002; Nanda et al., 2002), *Sox* (Takehana et al., 2014), steroid-pathway (Koyama et
102 al., 2019; Purcell et al., 2018) and Transforming Growth Factor beta (TGFβ) families (Q. Pan et al.,
103 2021), which have been independently and recurrently used to generate new MSD genes. The greatest
104 diversity of MSD genes is found within the TGFβ family with the anti-mullerian hormone, *amh*
105 (Hattori et al., 2012; M. Li et al., 2015; Q. Pan et al., 2019), the gonadal soma derived factor, *gsdf*
106 (Myosho et al., 2012; Rondeau et al., 2013), or the growth/differentiation factor 6, *gdf6* (Imarazene
107 et al., 2021; Reichwald et al., 2015) genes, but also TGF-β type II and type I receptors with the anti-
108 Mullerian hormone receptor type 2, *amhr2* (Feron et al., 2020; Kamiya et al., 2012) and the bone
109 morphogenetic protein receptor, type IBb, *bmpr1bb* (Rafati et al., 2020) genes. However, a few
110 exceptions to the “usual suspects” rule have also been identified with, for instance, the conserved

salmonid MSD *sdY* gene that evolved from an immunity-related gene (Bertho, Herpin, Scharl, & Guiguen, 2021; Yano et al., 2012, 2013). Based on a chromosome-scale high quality genome assembly, and previously published whole-organ transcriptomic data (Pasquier et al., 2016) of *Pangasianodon hypophthalmus*, we identified a male-specific duplication of *amhr2* (*amhr2by*) in that species. This potential Y chromosome-specific *amhr2by* encodes an N-terminal truncated protein that lacks the cysteine-rich extracellular part of the receptor, which is key for proper Amh ligand binding. Sex-linkage studies and genome sequencing of 11 additional Pangasiid species show that *amhr2by* is conserved as a male-specific gene in at least four Pangasiid species, stemming from a single ancient duplication / insertion event at the root of the Siluroidei suborder radiation that is dated around 100 million years ago (Kappas, Vittas, Pantzartzi, Drosopoulou, & Scouras, 2016). Together, these results bring multiple lines of evidence supporting the hypothesis that *amhr2by* is potentially an ancient and conserved master sex-determining gene in Pangasiid catfishes and highlight the recurrent usage of the transforming growth factor β pathway in teleost sex determination.

MATERIAL AND METHODS

Samples collection

For high-quality genome reference sequencing, a single *P. hypophthalmus* male was sampled from captive broodstock populations originating from Indonesia and maintained in the experimental facilities of ISEM (Institut des Sciences de l'Evolution de Montpellier, France). High molecular weight (HMW) genomic DNA (gDNA) was extracted from a 0.5-ml blood sample stored in a TNES-Urea lysis buffer (TNES-Urea: 4 M urea; 10 mM Tris-HCl, pH 7.5; 125 mM NaCl; 10 mM EDTA; 1% SDS). HMW gDNA was then purified using a slightly-modified phenol-chloroform extraction (Q. Pan et al., 2019). For the chromosome contact map (Hi-C), 1.5 ml of blood was taken from the same animal and slowly cryopreserved with 15 % dimethyl sulfoxide (DMSO) in a Mr. Frosty Freezing Container (Thermo Scientific) at -80°C. For sex-linkage analyses and short-read genome sequencing, fin clips were sampled and stored in 90% ethanol. *P. djambal* fin clips were sampled from captive broodstock populations originating from Indonesia and maintained in the experimental facilities of ISEM. *P. gigas* fin clips were sampled on broodstock populations kept for a restocking program in Thailand. *P. bocourti* and *P. conchophilus* fin clips were sampled at market places in Vietnam. *P. elongatus*, *P. siamensis*, *P. macronema*, *P. larnaudii*, *P. mekongensis*, and *P. krempfi*

were wild samples collected in Vietnam. *P. sanitwongsei* fin clip samples were obtained through the aquaculture trade and their precise origin is unknown.

Chromosome-scale genome sequencing and assembly of *P. hypophthalmus*

Oxford Nanopore sequencing

All library preparations and sequencing were performed using Oxford Nanopore Ligation Sequencing Kits SQK-LSK108 and SQK-LSK109 according to the manufacturer's instructions (Oxford Nanopore Technologies). For the SQK-LSK108 sequencing Kit, 90 µg of DNA was purified then sheared to 20 kb fragments using the megaruptor1 system (Diagenode). For each library, a DNA-damage repair step was performed on 5 µg of DNA. Then an END-repair-dA-tail step was performed for adapter ligation. Libraries were loaded onto nine R9.4.1 flowcells and sequenced on a GridION instrument at a concentration of 0.1 pmol for 48 h. For the SQK-LSK109 sequencing Kit, 10 µg of DNA was purified then sheared to 20 kb fragments using the megaruptor1 system (Diagenode). For this library, a one-step DNA-damage repair + END-repair-dA-tail procedure was performed on 2 µg of DNA. Adapters were then ligated to DNAs in the library. The library was loaded onto one R9.4.1 flowcell and sequenced on a GridION instrument at a concentration of 0.08 pmol for 48h.

10X Genomics sequencing

The Chromium library was prepared according to 10X Genomics' protocols using the Genome Reagent Kit v2. The library was prepared from 10 ng of high molecular weight (HMW) gDNA. Briefly, in the microfluidic Genome Chip, a library of Genome Gel Beads, was combined with HMW template gDNA in master mix and partitioning oil to create Gel Bead-In-EMulsions (GEMs) in the Chromium apparatus. Each Gel Bead was then functionalized with millions of copies of a 10x™ barcoded primer. Dissolution of the Genome Gel Bead in the GEM released primers containing (i) an Illumina R1 sequence (Read 1 sequencing primer), (ii) a 16 bp 10x Barcode, and (iii) a 6 bp random primer sequence. The R1 sequence and the 10x™ barcode were added to the molecules during the GEM incubation. P5 and P7 primers, R2 sequence, and Sample Index were added during library construction. 10 cycles of PCR were applied to amplify the library. The library was sequenced on an Illumina HiSeq3000 using a paired-end format with read length of 150 bp with the Illumina HiSeq3000 sequencing kits.

Hi-C sequencing

174 Hi-C library generation was carried out according to a protocol adapted from Rao et al. 2014 (Foissac
175 et al., 2019). The blood sample was spun down, and the cell pellet was resuspended and fixed in 1%
176 formaldehyde. Five million cells were processed for the Hi-C library. After overnight digestion with
177 HindIII (NEB), DNA ends were labeled with Biotin-14-DCTP (Invitrogen) using the klenow (NEB)
178 and religated. In total, 1.4 µg of DNA was sheared to an average size of 550 bp (Covaris). Biotinylated
179 DNA fragments were pulled down using M280 Streptavidin Dynabeads (Invitrogen) and ligated to
180 PE adaptors (Illumina). The Hi-C library was amplified using PE primers (Illumina) with 10 PCR
181 amplification cycles. The library was sequenced using a HiSeq3000 (Illumina, California, USA) in
182 150 bp paired-end format.

183

184 **Genome assembly**

185

186 GridION data were trimmed using Porechop v0.2.1 (<https://github.com/rrwick/Porechop>) and filtered
187 using NanoFilt v2.2.0 (De Coster, D'Hert, Schultz, Cruts, & Van Broeckhoven, 2018) with the
188 parameters -l 3000 and -q 7. A *de novo* assembly was constructed with SmartDeNovo (Ruan,
189 2015/2019), Wtdbg2 v2.1 (Ruan & Li, 2020) and flye v2.3.7 (Kolmogorov, Yuan, Lin, & Pevzner,
190 2019), each with default parameters. The resulting assembly metrics were compared, and the draft
191 assembly with the best metrics generated by SmartDeNovo was kept and used as reference. This
192 assembly was then further corrected using long reads. After mapping the trimmed and filtered
193 GridION reads with minimap2 v2.11 (H. Li, 2018) with parameter -x map-ont, the assembly was
194 polished using Racon (Vaser, Sović, Nagarajan, & Šikić, 2017) v1.3.1 with default parameters for
195 three rounds. The assembly was then corrected using short reads. After mapping 10X short reads with
196 Long Ranger v2.1.1, Pilon (Walker et al., 2014) v1.22 was run with parameters --fix bases,gaps --
197 changes. Again, three rounds of these short reads polishing were performed. The final polished
198 genome assembly was then scaffolded using Hi-C information. Reads were aligned to the draft
199 genome using Juicer (Durand, Shamim, et al., 2016) with default parameters. A candidate assembly
200 was then generated with 3D de novo assembly (3D-DNA) pipeline (Dudchenko et al., 2017) with the
201 -r 0 parameter. The candidate assembly was manually reviewed using Juicebox (Durand, Robinson,
202 et al., 2016) assembly tools. Gaps in this chromosome scaled assembly were filled using GapCloser
203 (https://github.com/CAFS-bioinformatics/LR_Gapcloser) v1.1 with default parameters. Reads used
204 to fill these gaps were GridION and PromethION reads filtered with NanoFilt and then corrected with
205 Canu (Koren et al., 2017) v1.6 using parameters --correct genomeSize = 753m --nanopore-raw. The
206 assembly was then corrected one last time using short reads polishing pipeline.

Genome analysis and protein-coding gene annotation

K-mer-based estimation of the genome size was carried out with GenomeScope (Vurtture et al., 2017) v2.0. 10X reads were processed with Jellyfish v1.1.11 (Marçais & Kingsford, 2011) to count 21-mer with a max k-mer coverage of 10,000 and 1,000,000. BUSCO (Simão, Waterhouse, Ioannidis, Kriventseva, & Zdobnov, 2015) v3.0.2 was run with parameters –species zebrafish and –limit 10 on the single-copy orthologous gene library from the actinopterygii lineage. The first annotation step was to identify repetitive content using RepeatMasker v4.0.7 (<https://www.repeatmasker.org/>), Dust (Morgulis, Gertz, Schäffer, & Agarwala, 2006), and TRF v4.09 (Benson, 1999). A species-specific *de novo* repeat library was built with RepeatModeler v1.0.11 (<http://www.repeatmasker.org/RepeatModeler/>) and repeated regions were located using RepeatMasker with the *de novo* and *Danio rerio* libraries. Bedtools v2.26.0 (Quinlan & Hall, 2010) was used to merge repeated regions identified with the three tools and to soft mask the genome. The Maker3 genome annotation pipeline v3.01.02-beta (Holt & Yandell, 2011) combined annotations and evidence from three approaches: similarity with fish proteins, assembled transcripts, and *de novo* gene predictions. Protein sequences from 11 fish species (*Astyanax mexicanus*, *Danio rerio*, *Gadus morhua*, *Gasterosteus aculeatus*, *Lepisosteus oculatus*, *Oreochromis niloticus*, *Oryzias latipes*, *Poecilia formosa*, *Takifugu rubripes*, *Tetraodon nigroviridis*, *Xiphophorus maculatus*) found in Ensembl were aligned to the masked genome using Exonerate v2.4 (Slater & Birney, 2005). RNA-Seq reads of *P. hypophthalmus* (NCBI BioProject PRJNA256973) from the PhyloFish project (Pasquier et al., 2016) were used for genome annotation and aligned to the chromosomal assembly using STAR v2.5.1b (Dobin et al., 2013) with outWigType and outWigStrand options to output signal wiggle files. Cufflinks v2.2.1 (Trapnell et al., 2010) was used to assemble the transcripts that were used as RNA-seq evidence. Braker v2.0.4 (Hoff, Lange, Lomsadze, Borodovsky, & Stanke, 2016) provided *de novo* gene models with wiggle files provided by STAR as hint files for GeneMark (Hoff et al., 2016) and Augustus (Stanke et al., 2006) training. The best supported transcript for each gene was chosen using the quality metric called Annotation Edit Distance (AED) (Eilbeck, Moore, Holt, & Yandell, 2009).

miRNA gene and mature miRNA annotation

Small RNA Illumina sequencing libraries were prepared using the NEXTflex Small RNA-Seq Kit v3 (PerkinElmer) following the manufacturer's instructions and starting with the same total RNA extracts as for the Phylofish project (Pasquier et al., 2016). Total RNA was extracted using Trizol reagent (Euromedex, France) according to the manufacturer's instructions. Libraries were sequenced

on an Illumina HiSeq 2500 sequencer and raw reads were pre-processed using CUTADAPT version 3.4 (Martin, 2011). All eight adult organ libraries (brain, gills, heart ventricle, skeletal muscle, intestine, liver, ovary and testis) were simultaneously analyzed using *Prost!* (Thomas Desvignes, Batzel, Sydes, Eames, & Postlethwait, 2019) selecting for read length 17 to 25 nucleotides and with a minimum of five identical reads. Reads were then aligned to the species' reference genome using bbmapskimmer.sh version 37.85 of the BBMap suite (<https://sourceforge.net/projects/bbmap/>). Gene and mature miRNA annotations were performed as previously described (Thomas Desvignes et al., 2019) based on established miRNA gene orthologies among ray-finned fish species (Thomas Desvignes, Sydes, Montfort, Bobe, & Postlethwait, 2021) and using previously published miRNA annotations in spotted gar, zebrafish, three-spined stickleback, Japanese medaka, shortfin molly and blackfin icefish as reference (Braasch et al., 2016; Thomas Desvignes et al., 2019, 2021; Kelley et al., 2021; B.-M. Kim et al., 2019). miRNA and isomiR nomenclature follow the rules established for zebrafish (T. Desvignes et al., 2015).

Short-read sequencing and genome-guided assemblies of other Pangasiids

Short-read sequencing

The *P. gigas* and *P. djambal* genomes were sequenced using an Illumina 2x250 bp format. DNA library construction was performed according to the manufacturer's instruction using the Truseq DNA nano library prep kit (Illumina). Briefly, gDNA was quantified using the HS dsDNA Assay kit on the Qubit (Invitrogen). 200 ng of gDNA were sonicated on a Bioruptor (Diagenode). Sonicated gDNA was end repaired and size selected on magnetic beads aiming for fragments of an average size of 550 pb. Selected fragments were adenylated on their 3' ends before ligation of Illumina's indexed adapters. The library was amplified using 8 PCR cycles and verified on a Fragment Analyzer using the HS NGS fragment kit (Agilent). The library was quantified by qPCR using the KAPA Library quantification kit (Roche, ref. KK4824) and sequenced on half a lane of HiSeq2500 in paired end 2x250nt using the clustering and SBS rapid kit following the manufacturer's instructions. All other species were sequenced using an Illumina 2x150 bp strategy according to Illumina's protocols using the Illumina TruSeq Nano DNA HT Library Prep Kit. Briefly, DNA was fragmented by sonication, size selection was performed using SPB beads (kit beads) and adaptors were ligated to be sequenced. Library quality was assessed using an Advanced Analytical Fragment Analyzer and libraries were quantified by qPCR using the Kapa Library Quantification Kit. DNA-seq experiments were performed on one Illumina NovaSeq S4 lane using a paired-end read length of 2x150 bp with the Illumina NovaSeq6000 Reagent Kits.

Assembly and annotation

The *P. gigas* and *P. djambal* genomes were assembled from 2x250 bp short reads using the DiscovarDeNovo assembler (<https://github.com/bayolau/discovardenovo/>) with default parameters. For *P. sanitwongsei*, *P. conchophilus*, *P. bocourti*, *P. larnaudii*, *P. mekongensis*, and *P. krempfi*, 2x150 bp reads were assembled using SPADes v.3.11.1 (Bankevich et al., 2012) and then purged using purge_dups (Guan et al., 2020). The *P. elongatus*, *P. macronema* and *P. siamensis* 2x150 bp short reads were assembled with SPADes v.3.14.1 instead of v.3.11.1 because of a higher individual genome heterozygosity (> 1%), followed by a more stringent purge with Redundans v0.14a (Pryszcz & Gabaldón, 2016). All these species were then assembled into pseudo-chromosomes using a reference-guided strategy and the “query assembled as reference” function from DGenies v1.2.0 (Cabanettes & Klopp, 2018), and the GENO_Phyp_1.0 *P. hypophthalmus* assembly used as a reference. Genes from the NCBI annotation of GENO_Phyp_1.0 were then mapped to chromosome-scale assemblies using Liftoff (Shumate & Salzberg, 2021).

Species and gene phylogenies

Whole-genome species phylogeny analysis was carried out with protein gene annotation from our 12 Pangasidae species combined with protein sequences from *Ictalurus punctatus* (siluriformes) as a Pangasidae outgroup species. Outgroup species protein sequences were retrieved from Ensembl release 103 (Howe et al., 2021). Orthogroups were identified using OrthoFinder (Emms & Kelly, 2019), followed by multiple sequence alignment of concatenated one-to-one orthologs (n = 8151) using MAFFT version 7.475 (Katoh & Standley, 2013). Species tree inference was performed via IQ-TREE 2 (Minh et al., 2020), the latter using a standard non-parametric bootstrap (r = 100).

Gene and protein phylogenetic reconstructions were performed on all *amhr2*/Amhr2 homologous sequences from 28 catfish species along with *amhr2* sequences from *Astyanax mexicanus* (characiformes) and *Electrophorus electricus* (gymnotiformes) as siluriformes outgroups (. Full-length CDS were predicted based on their genomic and protein sequence annotation or retrieved from GenBank (see Table S2 and multi-fasta files of these sequences are publicly available at <https://doi.org/10.15454/M3HYAX>). To verify the tree topology of *amhr2*/Amhr2 homologs, besides complete protein and cDNA sequences, we also constructed phylogenetic trees with only the first and second codons of the coding sequences (Lemey, 2009). All putative CDS and protein sequences were

then aligned using MAFFT (version 7.450) (Katoh & Standley, 2013). Residue-wise confidence scores were computed with GUIDANCE 2 (Sela, Ashkenazy, Katoh, & Pupko, 2015), and only well-aligned residues with confidence scores above 0.99 were retained. Phylogenetic relationships among the *amhr2* sequences were inferred with both maximum-likelihood implemented in IQ-TREE (version 1.6.7) (Minh et al., 2020), and Bayesian methods implemented in Phylobayes (version 4.1) (Lartillot, Lepage, & Blanquart, 2009). More precisely, alignment files from either full-length cDNA, third-codon-removed cDNA, or full-length proteins were used for model selection and tree inference with IQ-TREE (version 1.6.7) (Minh et al., 2020) with 1000 bootstraps and the 1000 SH-like approximate likelihood ratio test for robustness. The same alignment files were run in a Bayesian framework with Phylobayes (version 4.1) (Lartillot et al., 2009) using the CAT-GTR model with default parameters, and two chains were run in parallel for approximately 2000 cycles with the first 500 cycles discarded as burnt-in until the average standard deviation of split frequencies remained \leq 0.001. The resulting phylogenies were visualized with Figtree (version 1.44).

Selection analysis on *amhr2* sequences

Selection analysis was performed on the *amhr2* phylogeny using Godon (Davydov, Salamin, & Robinson-Rechavi, 2019). Analyses were performed separately for (a) exons conserved in both *amhr2a* and *amhr2by* (“conserved exons”) and (b) the exon region found only in *amhr2a* (“first exons”). Three codon models were used: M8 (Yang, Nielsen, Goldman, & Pedersen, 2000), M8 with codon gamma rate variation (Davydov et al., 2019), and the branch-site model (Zhang, Nielsen, & Yang, 2005) (conserved exons only). For the branch-site model, the branch leading to the *amhr2by* clade was set as the foreground branch.

Transcriptome analyses

Reads from *P. hypophthalmus* adult organs and embryos (Pasquier et al., 2016) were mapped on the complete *P. hypophthalmus* reference transcriptome using bwa mem version 0.7.17 (H. Li, 2013). Unique mapped reads were then filtered and a raw count matrix was generated with htseq-count (Anders, Pyl, & Huber, 2015) and normalized using DESeq2 (Love, Huber, & Anders, 2014). Genes of interest were extracted from this complete transcriptome dataset and missing values were replaced by a minimal value (0.1) in the normalized raw count matrix. Hierarchical classification was carried out after log transformation and gene median centering using the cluster 3.0 software (de Hoon, Imoto, Nolan, & Miyano, 2004) with an uncentered correlation similarity metric and an average linkage clustering method.

Read-coverage analyses around the *amhr2a* and *amhr2by* loci in Pangasiids

To assess whether *amhr2by* is a potential Y specific gene in species for which whole genome sequencing was only obtained from one sample, we computed the read coverage throughout the genome and extracted the read coverage information around the *amhr2a* and *amhr2by* loci. In *P. hypophthalmus*, ONT reads were mapped on its own genome assembly using minimap version 2.11 (H. Li, 2018). In other Pangasiids, Illumina pair-end reads were mapped onto the *P. hypophthalmus* genome assembly using bwa version 0.7.17 (H. Li, 2013), indexed using samtools version 1.8 (H. Li et al., 2009) and sorted by PICARD SortSam. Then a pileup file was generated using samtools mpileup (H. Li et al., 2009) with per-base alignment quality disabled and (-B). Subsequently, a sync file containing the nucleotide composition for each position in the reference was created from the pileup file using popoolation mpileup2sync version 1.201 with a min quality of 20 (-min-qual 20) (Kofler, Pandey, & Schlötterer, 2011). Read depth was then calculated in a 10 kb non-overlapping window using PSASS (version 2.0.0, doi:10.5281/zenodo.2615936).

Primer design

P. hypophthalmus amhr2a and *amhr2by* genes were aligned with bioedit version 7.0.5.3 and specific primers were designed based on this alignment to select highly divergent positions for each paralog. Selected primer sequences were forward: 5'-GGAGTCTATAAACCCGTGGTAGC-3', and reverse: 5'-CTATGTCACGCTGAACCTCCAGTGT-3' (expected amplicon size: 153 bp) for the *amhr2by* gene and forward: 5'-GGAGTCTATAAGCCAGCGGTGGCT-3', and reverse: 5'-CTATGCCAGAATAACCCTGCAATGC-3' (expected amplicon size: 142 bp) for the *amhr2a* gene.

DNA extraction for PCR sex genotyping

DNA from fin clips was extracted using a Chelex-based extraction method. Briefly, a piece of fin clip from each sample was placed into a PCR tube, and then 150 µl 5% Chelex and 20 µl 1 mg/ml proteinase K were added to each tube. Tubes were then vortexed and quickly spun down. After that, samples were incubated for 2 h at 56°C followed by boiling 10 min at 99°C. DNA was then centrifuged at 7500 g for 5 min and diluted to 1:2 with double distilled water. Genotyping PCR reactions were run in 12.5 µl with 1.25 µl JumpStart PCR buffer 10X, 0.125 µl 25 mM dNTP, 0.25 µl 10 µM forward and reverse primers, 8.5 µl ddH₂O and 2 µl DNA. PCR cycling conditions were: 95°C for 3 min as initial denaturation, then 35 cycles for amplification with denaturation at 95°C for 30 s, annealing at 52°C for 30 s and extension at 72°C for 30 s, and finally another more extension at 72°C for 30 s and hold at 4°C.

RESULTS

A high-quality chromosome-scale genome assembly of *P. hypophthalmus*

A high-quality reference genome of a male *P. hypophthalmus* was sequenced using a combination of 10X Linked-Reads, Oxford Nanopore long reads and a chromosome contact map (Hi-C). Its genome size based on the kmer linked-reads distribution was estimated around 810 Mb including, respectively 65% and 35 % of unique and repeated sequences. The heterozygosity level of this *P. hypophthalmus* genome was estimated at around 1.2 %. The integration of all sequencing data provided a genome assembly size of 760 Mb (93% of the kmer estimated size), containing 612 contigs, a scaffold N50 of 26.4 Mb (Table 1) and 99.2% of all sequences anchored onto 30 chromosomes after Hi-C integration (see assembly metrics and comparison with other genome assemblies in Table 1). Combining *de novo* gene predictions, homology to teleost proteins, and evidence from transcripts, 25,076 protein-coding genes were annotated in our male *P. hypophthalmus* reference genome using our in-house genome annotation protocol. Because our *P. hypophthalmus* genome assembly has been derived by NCBI to produce a Reference Sequence (RefSeq) record (GCF_009078355.1) and was annotated by the NCBI Eukaryotic Genome Annotation Pipeline, the NCBI annotation will be used thereafter as reference in the following text. In addition to protein-coding genes, 323 microRNA genes (miRNAs) and 389 mature miRNAs were annotated using Illumina small-RNA sequencing data from a panel of eight organs. Gene and mature miRNA annotations as well as analyzed expression patterns are publicly available on FishmiRNA (<http://www.fishmirna.org/>) (Thomas Desvignes et al., 2022). This genome-wide miRNA annotation represents the first exhaustive miRNA annotation available for a Pangasiid species.

Characterization of a male-specific *amhr2* duplication in *P. hypophthalmus*

Because many teleost MSD genes evolved from the duplication of an autosomal “usual suspect” gene, we first searched for potential duplicates of *dmrt1*, *amh*, *amhr2*, *sox3*, *gsdf* and *gdf6* genes in the *P. hypophthalmus* genome assemblies. We found no gene duplication for *dmrt1*, *amh*, *sox3*, *gsdf* and *gdf6* (*gdf6a* and *gdf6b*), but two *amhr2* homologs were found in the two male *P. hypophthalmus* assemblies (GENO_Phyp_1.0 and VN_pangasius) while only one *amhr2* gene was detected in the female *P. hypophthalmus* ASM1680104v1 assembly. In the GENO_Phyp_1.0 *P. hypophthalmus* assembly, these two *amhr2* homologs, i.e., LOC113540131 (annotated as bone morphogenetic protein receptor type-2-like) and LOC113533735 (annotated as anti-Mullerian hormone type-2 receptor-like) are located respectively on chromosome 4 (Chr04:32,081,919-32,105,291) and 10

(Chr10: 26,334,822-26,348,340). The single *amhr2* locus found in the female ASM1680104v1 assembly (in ASM1680104v1 Chr04), is on chromosome 4 and shares 99% identity over 13.5 kb (100% overlap) with LOC113533735, and 87% identity on only 3% overlapping regions with LOC113540131. Using primers (see Materials and Methods) designed to amplify specifically either LOC113540131 or LOC113533735, we genotyped *P. hypophthalmus* males (N=12) and females (N=11) and found that LOC113540131 is significantly linked with maleness ($p = 7.12 \times 10^{-5}$) with a single positive outlier among 11 phenotypic females (see Table 2). In contrast, LOC113533735 was detected in all males and females (Fig. 1). These genotyping results, along with the absence of LOC113540131 in the female ASM1680104v1 assembly, strongly support the hypothesis that LOC113540131 is a Y-specific male-specific gene. We thus called the LOC113540131 gene, *amhr2by*, as the male-specific Y chromosome paralog of the autosomal LOC113533735 gene named *amhr2a*.

Comparison of *P. hypophthalmus amhr2by* and *amhr2a* and their inferred proteins

Overall, the predicted structure of the autosomal *P. hypophthalmus amhr2a* and the canonical vertebrate *Amhr2* are similar with the same number of introns and exons. The mVISTA (Frazer, Pachter, Poliakov, Rubin, & Dubchak, 2004) alignments of *P. hypophthalmus amhr2a* and *amhr2by* genes along with their CDS (Fig. 2A), show that these two genes display some sequence identity only within their shared exons, with no significant homology detected in their intronic, 3'UTR, and 5'UTR sequences (Fig. 2A). In addition, the *amhr2by* gene is lacking the first two exons of *amhr2a*, and the third *amhr2by* exon is also truncated. The *amhr2by* and *amhr2a* CDS share 78.78% identity on 1,164 bp of overlapping sequences (78% of the *amhr2a* CDS that is 1,455 bp long). Correspondingly, the two deduced proteins share 70.32% identity over 380 overlapping amino-acids, and Amhr2by lacks 112 amino-acids at its N-terminal extremity corresponding to two first exons and part of exon 3 of Amhr2a. (Fig. 2B and 2C). Hence, the *P. hypophthalmus* Amhr2by translates as an N-terminal-truncated type II receptor lacking its whole extra-cellular domain mediating ligand binding, while overall the remaining of the other functional domains (transmembrane and serine-threonine kinase domain) remain similar between Amhr2a and Amhr2by (Fig. 2B and 2C).

Expression of *amhr2by* and *amhr2a* in *P. hypophthalmus* adult tissues

Using *P. hypophthalmus* RNA-Seq from the PhyloFish database (Pasquier et al., 2016), we examined the organ expression of *amhr2a* and *amhr2by* along with a series of SD genes previously identified in other teleosts, i.e., *amh*, *dmrt1*, *gsdf*, *gdf6a*, *gdf6b* and *sox3*. Among these genes, *amh*, *dmrt1*, and *gsdf* display predominant expression in the adult testis and / or ovary with a much lower expression

in the eight additional somatic organs examined or in embryos (Fig. 3A). The two *amhr2* genes also have a gonadal-predominant expression pattern with *amhr2a* being expressed in both ovary and testis while *amhr2by* being strictly expressed in the testis as expected for a Y chromosome sex determination gene (Fig. 3B). The two *gdf6* paralogs (*gdf6a*, *gdf6b*) and *sox3* have no expression or a low expression in gonads and are more expressed in embryos for *sox3* and *gdf6a* or in bones and brain for *sox3*.

Sex-linkage of *amhr2by* in Pangasiids

To explore the evolution of *amhr2by* in Pangasiids, we obtained gDNA samples from 11 additional Pangasiid species with at least some specimens being phenotypically sexed for four of these species (Table S2). Samples from fish that were phenotypically sexed (i.e., *Pangasianodon gigas*, *Pangasius djambal*, *Pangasius conchophilus*, and *Pangasius bocourti*) were PCR genotyped to explore the potential conservation of *amhr2by* male sex-linkage in Pangasiids. In three of these species, *amhr2by* was found to be significantly associated with male phenotype ($p < 8.528e^{-04}$) (Table 2), except in *P. gigas*, the association was not significant ($p = 0.3865$) due to the combination of low sample size (3 males and 3 females) and the presence of one female outlier (Table 2). To complement this genotyping information, one male individual of *P. gigas*, *P. djambal*, *P. conchophilus*, and *P. bocourti* and one individual of unknown sex for *P. elongatus*, *P. siamensis*, *P. sanitwongsei*, *P. macronema*, *P. larraudii*, *P. mekongensis* and *P. krempfi* were sequenced using Illumina short-read strategies. These genomic short-read sequences were assembled and anchored using a reference-guided strategy (Lischer & Shimizu, 2017) on the *P. hypophthalmus* chromosome assembly, and the NCBI gene annotation of GENO_Phyp_1.0 was lifted over to these assemblies (see genome and annotation metrics in Table S1). The *amhr2a* genes were extracted from all these guided assemblies, and *amhr2by* homologs were extracted from the four male assemblies, i.e., *P. gigas*, *P. djambal*, *P. conchophilus*, and *P. bocourti* as well as from the unknown sex assemblies of *P. sanitwongsei*, and *P. krempfi*. To better explore sex-linkage in species for which we only sequenced a single individual, read coverage was explored around the *amhr2a* and *amhr2by* loci using the *P. hypophthalmus* as reference genome (Fig. S1). Under the hypothesis that *amhr2by* is also a male-specific Y chromosomal gene in additional Pangasiids, we expected a half coverage around *amhr2by* in males (hemizygous in XY) and an average read coverage around the autosomal *amhr2a*. In agreement with that hypothesis, a half coverage was found around the *amhr2by* locus for all species in which *amhr2by* was identified i.e., the male individuals of *P. hypophthalmus*, *P. gigas*, *P. djambal*, *P. conchophilus*, and *P. bocourti* and individuals of unknown sex in *P. sanitwongsei*, and *P. krempfi*. This result supports hemizyosity of *amhr2by* in these species as expected for a Y chromosomal gene. In other

species, i.e., *P. elongatus*, *P. siamensis*, *P. macronema*, *P. larnaudii*, and *P. mekongensis*, no conclusion can be drawn because the absence of finding *amhr2by* in these individuals could be because they are XX females without a Y chromosome and an *amhr2by* gene, or these species may have lost *amhr2by* as a Y chromosome gene.

Evolution of *amhr2* in Siluriformes

These whole-genome annotations were combined with protein sequences from channel catfish, *Ictalurus punctatus* (Siluriformes, Ictaluridae) used as a Pangasiid outgroup, and 8151 groups of one-to-one orthologs were used after concatenation to construct a whole-genome species tree inference (Fig. 4). In addition, all Pangasiids *amhr2* sequences deduced from our genomic resources were used for phylogenetic analyses with other available catfish *amhr2* genes (Table S2), along with *amhr2* from a gymnotiform (*Electrophorus electricus*) and a characiform (*Astyanax mexicanus*) as the closest species outgroups to the Siluriformes order. The topologies of all trees, i.e., using maximum-likelihood and Bayesian methods on proteins, CDS, and CDS with third codons removed (see Materials and Methods), were all congruent in showing that most of the *amhr2* from the sub-order Siluroidei (Sullivan et al., 2006) cluster with the Pangasiid *amhr2a*, and that outside the Pangasiid family, only a single species (*Pimelodus maculatus*, Pimelodidae) has an *amhr2* duplication clustering with the *amhr2by* sequences (Fig. 5, Fig. S2). Within the Siluriformes, a single *amhr2* in *Corydoras* sp (Callichthyidae, Loricarioidei) roots the *amhr2a* and *amhr2b* duplications (Fig. 5, Fig. S2), suggesting that *amhr2b* (*P. maculatus*) and *amhr2by* (Pangasiids) arose from an ancient duplication / insertion event at the root of the Siluroidei radiation that is dated around 100 million years ago (Kappas et al., 2016). We also searched for selection acting on the Pangasiid *amhr2* sequences, but detected no statistically significant signal of positive selection (Table 3) for either all exons conserved in both *amhr2a* and *amhr2by* (“conserved exons”) or for the exon region found only in *amhr2a* (“first exons”).

DISCUSSION

The Pangasiid family contains both important aquaculture species (Lazard et al., 2009) and key ecological catfish species (Eva et al., 2016) in many south Asian countries. Here, we present a reference genome for striped catfish, *Pangasianodon hypophthalmus*, and provide an additional high-quality genomic resource combining long-read sequencing and a chromosomal assembly for this species. This *de novo* genome (GENO_Phyph_1.0, GCA_009078355.1) was assembled into 30 large scaffolds that most likely correspond to the 30 chromosomes reported previously in cytological

studies (Sreeputhorn et al., 2017). This assembly also improves the metrics of the previously publicly available male assembly VN_pangasius (GCA_003671635.1) that was not anchored on chromosomes (O. T. P. Kim et al., 2018), and is comparable in terms of assembly metrics to the newest female ASM1680104v1 (GCA_016801045.1) chromosome-anchored assembly (Z. Gao et al., 2021). In addition to this *P. hypophthalmus* chromosome-anchored assembly, we also provided short-read genome sequencing for eleven additional Pangasiid species belonging to the genera *Pangasianodon* (1 additional species) and *Pangasius* (10 additional species). These short-read assemblies have been anchored and annotated on our reference *P. hypophthalmus* genome assembly and now present a large public set of genomic resources for the Pangasiid family.

Phylogenetic relationships within Siluriformes are still debated with no consensus for clear placement of some families within this order (Kappas et al., 2016; Sullivan et al., 2006). But at a broader scale, it is generally accepted that the sub-order Loricarioidei (defined also as a super-family) containing the armored catfish families (Callichthyids and Loricariids) is the earliest-diverging Siluriformes clade with the Diplomystoidei sub-order being the sister group to the remaining Siluroidei sub-order (Kappas et al., 2016; Sullivan et al., 2006). Pangasiids belong to the Siluroidei sub-order and have been characterized as the sister group to either Ictaluridae and Cranoglanididae (Kappas et al., 2016) or Schilbidae (Villela et al., 2017). Their phylogeny has been explored using both mitochondrial and nuclear makers (Karinthanyakit & Jondeung, 2012; Pouyaud et al., 2016). Here, using a phylogenomic approach (Delsuc, Brinkmann, & Philippe, 2005), we were able to determine the precise phylogenetic relationships among the 12 Pangasiid species for which we produced genome sequencing. Our results confirmed the basal position of the *Pangasianodon* genus as already described (Karinthanyakit & Jondeung, 2012; Na-Nakorn et al., 2006; Pouyaud et al., 2016) and, although we did not sequence any *Helicophagus* or *Pseudolais* genera, results allowed us to resolve the taxonomic positions of several *Pangasius* species (Karinthanyakit & Jondeung, 2012).

The molecular basis of genetic sex determination has been explored in only a few catfishes, with reports on the identification of male sex-specific sequences supporting a XX/XY sex determination system in *Pseudobagrus ussuriensis* (Z.-J. Pan, Li, Zhou, Qiang, & Gui, 2015) and *Pelteobagrus (Tachysurus) fulvidraco* (Dan, Mei, Wang, & Gui, 2013; Wang, Mao, Chen, Liu, & Gui, 2009) from the Bagridae family, and in *Clarias gariepinus* from the Clariidae family (Kovács, Egedi, Bártfai, & Orbán, 2000). In the Ictalurid channel catfish, *Ictalurus punctatus*, based on whole genome sequencing of a YY individual and genome-wide analyses, an isoform of the breast cancer anti-resistance 1 (*bcar1*) gene has been characterized as the male master sex determining gene (Bao et al.,

2019). In Pangasiids, genetic sex-markers have been searched without success in *P. hypophthalmus* and *P. gigas* (Sriphairoj et al., 2007). In our study, based on chromosome-scale genome assemblies of many Pangasiid species, transcriptomic data (Pasquier et al., 2016), and sex-linkage analyses we identified a male-specific duplication of the *amhr2* (*amhr2by*) gene as a potentially conserved male master sex determining gene in that fish family. The role of *Amhr2* as a master sex determining gene has been functionally characterized in the tiger pufferfish, *Takifugu rubripes* and Ayu, *Plecoglossus altivelis* (Kamiya et al., 2012; Nakamoto et al., 2021) and strongly suggested by sex-linkage information in common seadragon, *Phyllopteryx taeniolatus*, alligator pipefish, *Syngnathoides biaculeatus* (Qu et al., 2021), other species of pufferfishes (Duan et al., 2021; F.-X. Gao et al., 2020; Kamiya et al., 2012) and yellow perch, *Perca flavescens* (Feron et al., 2020). In addition, the anti-Mullerian hormone, Amh, which is the cognate ligand of AmhR2, has also been demonstrated or suggested as a master sex determining gene in a few fish species (Hattori et al., 2012; M. Li et al., 2015; Q. Pan et al., 2019, 2021; Song et al., 2021). Our results thus provide a new example of the repeated and independent recruitment of Amh and TGF β pathway members in fish genetic sex determination (Q. Pan et al., 2021). Although formal proof that this *amhr2by* gene is a conserved master sex determining gene in Pangasiids will require additional gene expression analyses and functional demonstrations, our results have application as a useful marker for sex control in many Pangasiid species in aquaculture. Sex dimorphic growth is often one of the main reasons for breeding all-male or all-female populations for aquaculture purposes. In Pangasiids, females have a faster growth rate in *P. djambal* above 3 kg, probably linked with the early maturation of males (Legendre et al., 2000). In contrast, weight gain was better in males compared to females in *P. bocourti* (Meng-Umphang, 2009). In addition, our results will also allow better management of breeders used for restocking in the large and endangered Mekong Giant Catfish, *P. gigas*, because maturation takes as long as 16-20 years in this species (Sriphairoj et al., 2007).

Our results on Pangasiid sex determination also raise interesting questions on Amhr2 structure and evolution. For instance, the N-terminal truncation of all the Pangasiid Amhr2by proteins is intriguing because this N-terminal part of the TGF β type II receptors encodes the complete extracellular ligand-binding domain that is known to be crucial for ligand binding specificity (Hart et al., 2021). N-terminal truncations of TGF β receptors acting as sex-determining genes have been already reported for Amhr2 in yellow perch (Feron et al., 2020) and common seadragon (Qu et al., 2021), and for Bmpr1b in the Atlantic herring, *Clupea harengus* (Rafati et al., 2020). In the Atlantic herring, the N-terminal truncated Bmpr1bby protein lacks the canonical TGF β receptor extracellular domains, but has maintained its ability to propagate a specific intracellular signal through kinase activity and Smad

578 protein phosphorylation (Rafati et al., 2020). Together, these studies suggest that some TGF β
 579 receptors truncated in their N-terminal extracellular ligand-binding domain can still trigger a
 580 biological response independent from any ligand activation. The fact that convergently, many fish
 581 master sex determining genes encoding a TGF β receptor with a similar N-terminal truncation,
 582 suggests that such a ligand-independent action is probably an important step that could have been
 583 selected independently to allow an autonomous action of the master sex determining gene. A second
 584 interesting and unexpected result from our study is that the duplication of *amhr2* genes that gave birth
 585 to the Pangasiids *amhr2by* gene is potentially ancient and so is likely to still be present in additional
 586 catfish species outside the Pangasidae family. This result is well supported by the topologies of our
 587 *amhr2* phylogenetic gene trees that place the origin of this duplication at the root of the Siluroidei
 588 sub-order that is dated around 100 Mya (Kappas et al., 2016). We also found one example of an
 589 *amhr2b* that is retained in the *Pimelodus maculatus* (Pimelodidae family) genome, although we do
 590 not know if this gene is also sex-linked in this species. But surprisingly no other *amhr2* duplication
 591 has been reported yet in other catfish species. Gains and losses of master sex determining genes have
 592 been already described such as in Esociformes in which some species have completely lost the *amh*
 593 duplication (*amhby*) that is a master sex determining gene in other closely related species from the
 594 same family (Q. Pan et al., 2019, 2021). Such complete gene losses can also be expected in catfishes
 595 like for instance in the channel catfish that relies on the *bcar1* gene as master sex determining gene
 596 (Bao et al., 2019), with no remains of an *amhr2* gene duplication. This situation is also probably the
 597 case for additional catfish species in which we did not find any *amhr2* duplication in male genome
 598 assemblies like in the Ictaluridae, *Ameiurus melas*, the Clariidae, *Clarias magur*, and the
 599 Auchenipteridae, *Ageneiosus marmoratus*. But if *amhr2b* is also male-specific as in Pangasiids, the
 600 question remains open for the additional catfish species where only female genome assemblies are
 601 currently available, such as in the Sisoridae, Siluridae and Bagridae families. A more extensive search
 602 for a potential duplication of *amhr2* genes in additional Siluroidei catfishes would be needed to better
 603 understand the fate of the *amhr2b* gene and whether it remains a master sex determining gene like in
 604 the Pangasiid family.

605
 606 Together our results bring multiple lines of evidence supporting the hypothesis that the conserved
 607 Pangasiid *amhr2by* is a potential sex determining gene that stemmed from an ancient duplication
 608 common to all Siluroidei catfishes. Our results highlight the recurrent usage of the TGF β pathway in
 609 teleost sex determination (Q. Pan et al., 2021) and the potential functional innovation through protein
 610 truncation. Furthermore, our results showcase the less considered long-term stability of sex
 611 determination gene in teleosts, a group that often receives attention for its dynamic evolution of sex

determination systems.

DATA AVAILABILITY

The Whole Genome Shotgun project of *P. hypophthalmus*, is available in the Sequence Read Archive (SRA), under BioProject reference PRJNA547555 with 10X genomics and Hi-C Illumina sequencing data is available in SRA under accession number SRX6071341 and SRX6071345 and Oxford Nanopore long reads data under SRA accession numbers SRX6071342 to SRX6071344 and SRX6071346 to SRX6071355. *P. hypophthalmus* small RNA-Seq sequences are available in SRA under Bioproject PRJNA256963. *P. gigas* and *P. djambal* genomes assembled with a *P. hypophthalmus* reference-guided strategy have been submitted to SRA under the respective BioProjects PRJNA593917 and PRJNA605300. All other Pangasiidae genomes assembled with a *P. hypophthalmus* reference-guided strategy without their genome annotations are available in SRA under BioProject PRJNA795327, and their genome assemblies plus their annotations are available in the omics dataverse (Open source research data repository) server with the following DOI (<https://doi.org/10.15454/M3HYAX>). *Pangasius siamensis* has been considered by NCBI curators as a *P. macronema* synonym and its genome is then recorded in NCBI with *P. macronema* as a Biosample species name, with sample name PaSia (for *Pangasius siamensis*) under accession BioSample number SAMN24707637.

BENEFIT-SHARING STATEMENT

A research collaboration was developed with scientists from the countries providing genetic samples (KS in Thailand, GR in Indonesia, TTTH in Vietnam, and JR and FLA in Brazil), all collaborators are included as co-authors, the results of research have been shared with the provider communities, and the research addresses a priority concern, in this case the conservation of organisms being studied. More broadly, our group is committed to international scientific partnerships, as well as institutional capacity building.

ACKNOWLEDGEMENTS

This project was supported by funds from the “Agence Nationale de la Recherche” (PhyloSex project, grant No. ANR-13-ISV7-0005; GenoFish project, grant No. ANR-16-CE12-0035) and R35 GM139635 and R01 OD011116 from the National Institutes of Health, USA. The GeT-PlaGe, Gentyane and Montpellier GenomiX (MGX) facilities were supported by France Génomique National infrastructure, funded as part of “Investissement d’avenir” program managed by Agence Nationale pour la Recherche (grant No. ANR-10-INBS-09). MRR was supported by the Swiss

646 National Science Foundation (grant 173048). We are grateful to the Genotoul bioinformatics platform
647 Toulouse Midi-Pyrenees (Bioinfo Genotoul) for providing computing and/or storage resources.

648

649 REFERENCES

- 650 Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq—A Python framework to work with high-throughput
651 sequencing data. *Bioinformatics (Oxford, England)*, 31(2), 166–169. doi:
652 10.1093/bioinformatics/btu638
- 653 Anka, I. Z., Faruk, M., Hasan, M. M., & Azad, M. (2014). *Environmental Issues of Emerging Pangas (*
654 *Pangasianodon hypophthalmus) Farming in Bangladesh*. doi: 10.3329/PA.V24I1-2.19118
- 655 Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., ... Pevzner, P. A. (2012).
656 SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing.
657 *Journal of Computational Biology*, 19(5), 455–477. doi: 10.1089/cmb.2012.0021
- 658 Bao, L., Tian, C., Liu, S., Zhang, Y., Elawad, A., Yuan, Z., ... Liu, Z. (2019). The Y chromosome sequence
659 of the channel catfish suggests novel sex determination mechanisms in teleost fish. *BMC Biology*,
660 17(1), 6. doi: 10.1186/s12915-019-0627-7
- 661 Baroiller, J. F., D'Cotta, H., & Saillant, E. (2009). Environmental effects on fish sex determination and
662 differentiation. *Sexual Development: Genetics, Molecular Biology, Evolution, Endocrinology,*
663 *Embryology, and Pathology of Sex Determination and Differentiation*, 3(2–3), 118–135. doi:
664 10.1159/000223077
- 665 Benson, G. (1999). Tandem repeats finder: A program to analyze DNA sequences. *Nucleic Acids Research*,
666 27(2), 573–580.
- 667 Bertho, S., Herpin, A., Scharl, M., & Guiguen, Y. (2021). Lessons from an unusual vertebrate sex-determining
668 gene. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*,
669 376(1832), 20200092. doi: 10.1098/rstb.2020.0092
- 670 Braasch, I., Gehrke, A. R., Smith, J. J., Kawasaki, K., Manousaki, T., Pasquier, J., ... Postlethwait, J. H. (2016).
671 The spotted gar genome illuminates vertebrate evolution and facilitates human-teleost comparisons.
672 *Nature Genetics*, 48(4), 427–437. doi: 10.1038/ng.3526
- 673 Cabanettes, F., & Klopp, C. (2018). D-GENIES: Dot plot large genomes in an interactive, efficient and simple
674 way. *PeerJ*, 6, e4958. doi: 10.7717/peerj.4958
- 675 Chen, S., Zhang, G., Shao, C., Huang, Q., Liu, G., Zhang, P., ... Wang, J. (2014). Whole-genome sequence of
676 a flatfish provides insights into ZW sex chromosome evolution and adaptation to a benthic lifestyle.
677 *Nature Genetics*, 46(3), 253–260. doi: 10.1038/ng.2890
- 678 Dan, C., Mei, J., Wang, D., & Gui, J.-F. (2013). Genetic Differentiation and Efficient Sex-specific Marker
679 Development of a Pair of Y- and X-linked Markers in Yellow Catfish. *International Journal of*
680 *Biological Sciences*, 9(10), 1043–1049. doi: 10.7150/ijbs.7203
- 681 Davydov, I. I., Salamin, N., & Robinson-Rechavi, M. (2019). Large-Scale Comparative Analysis of Codon
682 Models Accounting for Protein and Nucleotide Selection. *Molecular Biology and Evolution*, 36(6),
683 1316–1332. doi: 10.1093/molbev/msz048
- 684 De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M., & Van Broeckhoven, C. (2018). NanoPack: Visualizing
685 and processing long-read sequencing data. *Bioinformatics*, 34(15), 2666–2669. doi:
686 10.1093/bioinformatics/bty149
- 687 de Hoon, M. J. L., Imoto, S., Nolan, J., & Miyano, S. (2004). Open source clustering software. *Bioinformatics*
688 *(Oxford, England)*, 20(9), 1453–1454. doi: 10.1093/bioinformatics/bth078
- 689 Delsuc, F., Brinkmann, H., & Philippe, H. (2005). Phylogenomics and the reconstruction of the tree of life.
690 *Nature Reviews. Genetics*, 6(5), 361–375. doi: 10.1038/nrg1603
- 691 Desvignes, T., Batzel, P., Berezikov, E., Eilbeck, K., Eppig, J. T., McAndrews, M. S., ... Postlethwait, J. H.
692 (2015). miRNA Nomenclature: A View Incorporating Genetic Origins, Biosynthetic Pathways, and
693 Sequence Variants. *Trends in Genetics: TIG*, 31(11), 613–626. doi: 10.1016/j.tig.2015.09.002
- 694 Desvignes, Thomas, Bardou, P., Montfort, J., Sydes, J., Guyomar, C., George, S., ... Bobe, J. (2022).
695 FishmiRNA: An evolutionarily supported microRNA annotation and expression database for ray-
696 finned fishes. *Molecular Biology and Evolution*, msac004. doi: 10.1093/molbev/msac004
- 697 Desvignes, Thomas, Batzel, P., Sydes, J., Eames, B. F., & Postlethwait, J. H. (2019). miRNA analysis with

698 Prost! Reveals evolutionary conservation of organ-enriched expression and post-transcriptional
699 modifications in three-spined stickleback and zebrafish. *Scientific Reports*, 9(1), 3913. doi:
700 10.1038/s41598-019-40361-8

701 Desvignes, Thomas, Sydes, J., Montfort, J., Bobe, J., & Postlethwait, J. H. (2021). Evolution after Whole-
702 Genome Duplication: Teleost MicroRNAs. *Molecular Biology and Evolution*, 38(8), 3308–3331. doi:
703 10.1093/molbev/msab105

704 Devlin, R. H., & Nagahama, Y. (2002). Sex determination and sex differentiation in fish: An overview of
705 genetic, physiological, and environmental influences. *Aquaculture*, 208(3), 191–364. doi:
706 10.1016/S0044-8486(02)00057-1

707 Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., ... Gingeras, T. R. (2013). STAR:
708 Ultrafast universal RNA-seq aligner. *Bioinformatics (Oxford, England)*, 29(1), 15–21. doi:
709 10.1093/bioinformatics/bts635

710 Duan, W., Gao, F.-X., Chen, Z., Gao, Y., Gui, J.-F., Zhao, Z., & Shi, Y. (2021). A sex-linked SNP mutation
711 in amhr2 is responsible for male differentiation in obscure puffer (Takifugu obscurus). *Molecular
712 Biology Reports*, 48(8), 6035–6046. doi: 10.1007/s11033-021-06606-4

713 Dudchenko, O., Batra, S. S., Omer, A. D., Nyquist, S. K., Hoeger, M., Durand, N. C., ... Aiden, E. L. (2017).
714 De novo assembly of the Aedes aegypti genome using Hi-C yields chromosome-length scaffolds.
715 *Science (New York, N.Y.)*, 356(6333), 92–95. doi: 10.1126/science.aal3327

716 Durand, N. C., Robinson, J. T., Shamim, M. S., Machol, I., Mesirov, J. P., Lander, E. S., & Aiden, E. L. (2016).
717 Juicebox Provides a Visualization System for Hi-C Contact Maps with Unlimited Zoom. *Cell Systems*,
718 3(1), 99–101. doi: 10.1016/j.cels.2015.07.012

719 Durand, N. C., Shamim, M. S., Machol, I., Rao, S. S. P., Huntley, M. H., Lander, E. S., & Aiden, E. L. (2016).
720 Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. *Cell Systems*,
721 3(1), 95–98. doi: 10.1016/j.cels.2016.07.002

722 Eilbeck, K., Moore, B., Holt, C., & Yandell, M. (2009). Quantitative measures for the management and
723 comparison of annotated genomes. *BMC Bioinformatics*, 10, 67. doi: 10.1186/1471-2105-10-67

724 Emms, D. M., & Kelly, S. (2019). OrthoFinder: Phylogenetic orthology inference for comparative genomics.
725 *Genome Biology*, 20(1), 238. doi: 10.1186/s13059-019-1832-y

726 Eva, B., Harmony, P., Thomas, G., Francois, G., Alice, V., Claude, M., & Tony, D. (2016). Trails of river
727 monsters: Detecting critically endangered Mekong giant catfish Pangasianodon gigas using
728 environmental DNA. *Global Ecology and Conservation*, 7, 148–156. doi:
729 10.1016/j.gecco.2016.06.007

730 Feron, R., Zahm, M., Cabau, C., Klopp, C., Roques, C., Bouchez, O., ... Guiguen, Y. (2020). Characterization
731 of a Y-specific duplication/insertion of the anti-Mullerian hormone type II receptor gene based on a
732 chromosome-scale genome assembly of yellow perch, *Perca flavescens*. *Molecular Ecology
733 Resources*, 20(2), 531–543. doi: 10.1111/1755-0998.13133

734 Foissac, S., Djebali, S., Munyard, K., Vialaneix, N., Rau, A., Muret, K., ... Giuffra, E. (2019). Multi-species
735 annotation of transcriptome and chromatin structure in domesticated animals. *BMC Biology*, 17(1),
736 108. doi: 10.1186/s12915-019-0726-5

737 Frazer, K. A., Pachter, L., Poliakov, A., Rubin, E. M., & Dubchak, I. (2004). VISTA: Computational tools for
738 comparative genomics. *Nucleic Acids Research*, 32(Web Server issue), W273-279. doi:
739 10.1093/nar/gkh458

740 Gao, F.-X., Shi, Y., Duan, W., Lu, W.-J., Huang, W., Zhang, X.-J., ... Gui, J.-F. (2020). A rapid and reliable
741 method for identifying genetic sex in obscure pufferfish (Takifugu obscurus). *Aquaculture*, 519,
742 734749. doi: 10.1016/j.aquaculture.2019.734749

743 Gao, Z., You, X., Zhang, X., Chen, J., Xu, T., Huang, Y., ... Shi, Q. (2021). A chromosome-level genome
744 assembly of the striped catfish (Pangasianodon hypophthalmus). *Genomics*, 113(5), 3349–3356. doi:
745 10.1016/j.ygeno.2021.07.026

746 Guan, D., McCarthy, S. A., Wood, J., Howe, K., Wang, Y., & Durbin, R. (2020). Identifying and removing
747 haplotypic duplication in primary genome assemblies. *Bioinformatics*, 36(9), 2896–2898. doi:
748 10.1093/bioinformatics/btaa025

749 Hart, K. N., Stocker, W. A., Nagykerly, N. G., Walton, K. L., Harrison, C. A., Donahoe, P. K., ... Thompson,
750 T. B. (2021). Structure of AMH bound to AMHR2 provides insight into a unique signaling pair in the
751 TGF- β family. *Proceedings of the National Academy of Sciences of the United States of America*,
752 118(26), e2104809118. doi: 10.1073/pnas.2104809118

- 753 Hattori, R. S., Murai, Y., Oura, M., Masuda, S., Majhi, S. K., Sakamoto, T., ... Strüssmann, C. A. (2012). A
754 Y-linked anti-Müllerian hormone duplication takes over a critical role in sex determination.
755 *Proceedings of the National Academy of Sciences of the United States of America*, 109(8), 2955–2959.
756 doi: 10.1073/pnas.1018392109
- 757 Herpin, A., & Scharl, M. (2015). Plasticity of gene-regulatory networks controlling sex determination: Of
758 masters, slaves, usual suspects, newcomers, and usurpators. *EMBO Reports*, 16(10), 1260–1274. doi:
759 10.15252/embr.201540667
- 760 Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M., & Stanke, M. (2016). BRAKER1: Unsupervised RNA-
761 Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS. *Bioinformatics (Oxford, England)*, 32(5), 767–769. doi: 10.1093/bioinformatics/btv661
- 762 Holt, C., & Yandell, M. (2011). MAKER2: An annotation pipeline and genome-database management tool for
763 second-generation genome projects. *BMC Bioinformatics*, 12(1), 491. doi: 10.1186/1471-2105-12-491
- 764 Howe, K. L., Achuthan, P., Allen, J., Allen, J., Alvarez-Jarreta, J., Amode, M. R., ... Flicek, P. (2021). Ensembl
765 2021. *Nucleic Acids Research*, 49(D1), D884–D891. doi: 10.1093/nar/gkaa942
- 766 Imarazene, B., Du, K., Beille, S., Jouanno, E., Feron, R., Pan, Q., ... Guiguen, Y. (2021). A Supernumerary
767 “B-Sex” Chromosome Drives Male Sex Determination in the Pachón Cavefish, *Astyanax mexicanus*
768 (SSRN Scholarly Paper No. ID 3875774). Rochester, NY: Social Science Research Network. doi:
769 10.2139/ssrn.3875774
- 770 Kallman, K. D. (1973). The Sex-Determining Mechanism of the Platyfish, *Xiphophorus maculatus*. In J. H.
771 Schröder (Ed.), *Genetics and Mutagenesis of Fish* (pp. 19–28). Berlin, Heidelberg: Springer. doi:
772 10.1007/978-3-642-65700-9_2
- 773 Kamiya, T., Kai, W., Tasumi, S., Oka, A., Matsunaga, T., Mizuno, N., ... Kikuchi, K. (2012). A trans-species
774 missense SNP in *Amhr2* is associated with sex determination in the tiger pufferfish, *Takifugu rubripes*
775 (*fugu*). *PLoS Genetics*, 8(7), e1002798. doi: 10.1371/journal.pgen.1002798
- 776 Kappas, I., Vittas, S., Pantzartzi, C. N., Drosopoulou, E., & Scouras, Z. G. (2016). A Time-Calibrated
777 Mitogenome Phylogeny of Catfish (Teleostei: Siluriformes). *PLOS ONE*, 11(12), e0166988. doi:
778 10.1371/journal.pone.0166988
- 779 Karinthanyakit, W., & Jondeung, A. (2012). Molecular phylogenetic relationships of pangasiid and schilbid
780 catfishes in Thailand. *Journal of Fish Biology*, 80(7), 2549–2570. doi: 10.1111/j.1095-
781 8649.2012.03303.x
- 782 Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: Improvements
783 in performance and usability. *Molecular Biology and Evolution*, 30(4), 772–780. doi:
784 10.1093/molbev/mst010
- 785 Kelley, J. L., Desvignes, T., McGowan, K. L., Perez, M., Rodriguez, L. A., Brown, A. P., ... Tobler, M. (2021).
786 MicroRNA expression variation as a potential molecular mechanism contributing to adaptation to
787 hydrogen sulphide. *Journal of Evolutionary Biology*, 34(6), 977–988. doi: 10.1111/jeb.13727
- 788 Kim, B.-M., Amores, A., Kang, S., Ahn, D.-H., Kim, J.-H., Kim, I.-C., ... Park, H. (2019). Antarctic blackfin
789 icefish genome reveals adaptations to extreme environments. *Nature Ecology & Evolution*, 3(3), 469–
790 478. doi: 10.1038/s41559-019-0812-7
- 791 Kim, O. T. P., Nguyen, P. T., Shoguchi, E., Hisata, K., Vo, T. T. B., Inoue, J., ... Satoh, N. (2018). A draft
792 genome of the striped catfish, *Pangasianodon hypophthalmus*, for comparative analysis of genes
793 relevant to development and a resource for aquaculture improvement. *BMC Genomics*, 19(1), 733.
794 doi: 10.1186/s12864-018-5079-x
- 795 Kobayashi, Y., Nagahama, Y., & Nakamura, M. (2013). Diversity and plasticity of sex determination and
796 differentiation in fishes. *Sexual Development: Genetics, Molecular Biology, Evolution, Endocrinology, Embryology, and Pathology of Sex Determination and Differentiation*, 7(1–3), 115–
797 125. doi: 10.1159/000342009
- 798 Kofler, R., Pandey, R. V., & Schlötterer, C. (2011). PoPoolation2: Identifying differentiation between
799 populations using sequencing of pooled DNA samples (Pool-Seq). *Bioinformatics (Oxford, England)*,
800 27(24), 3435–3436. doi: 10.1093/bioinformatics/btr589
- 801 Kolmogorov, M., Yuan, J., Lin, Y., & Pevzner, P. A. (2019). Assembly of long, error-prone reads using repeat
802 graphs. *Nature Biotechnology*, 37(5), 540–546. doi: 10.1038/s41587-019-0072-8
- 803 Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., & Phillippy, A. M. (2017). Canu: Scalable
804 and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome*
805 *Research*, 27(5), 722–736. doi: 10.1101/gr.215087.116
- 806
- 807

- 808 Kovács, B., Egedi, S., Bártfai, R., & Orbán, L. (2000). Male-specific DNA markers from African catfish
809 (Clarias gariepinus). *Genetica*, 110(3), 267–276. doi: 10.1023/A:1012739318941
- 810 Koyama, T., Nakamoto, M., Morishima, K., Yamashita, R., Yamashita, T., Sasaki, K., ... Sakamoto, T. (2019).
811 A SNP in a Steroidogenic Enzyme Is Associated with Phenotypic Sex in Seriola Fishes. *Current*
812 *Biology*, 29(11), 1901–1909.e8. doi: 10.1016/j.cub.2019.04.069
- 813 Lartillot, N., Lepage, T., & Blanquart, S. (2009). PhyloBayes 3: A Bayesian software package for phylogenetic
814 reconstruction and molecular dating. *Bioinformatics*, 25(17), 2286–2288. doi:
815 10.1093/bioinformatics/btp368
- 816 Lazard, J., Cacot, P., Slembrouck, J., & Legendre, M. (2009). La pisciculture des Pangasiidae. *Cahiers*
817 *Agricultures*, 18(2–3), 164–173 (1). doi: 10.1684/agr.2009.0284
- 818 Legendre, M., Pouyaud, L., Slembrouck, J., Gustiano, R., Kristanto, A. H., Subagja, J., ... Maskeer. (2000).
819 Pangasius djambal: A new candidate species for fish culture in Indonesia. (Horizon (IRD)).
- 820 Lemey, P. (2009). *The Phylogenetic Handbook: A Practical Approach to Phylogenetic Analysis and*
821 *Hypothesis Testing* (\$ {number}nd édition). Cambridge, UK ; New York: Cambridge University Press.
- 822 Li, H. (2013). *Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM*. Retrieved
823 from <https://arxiv.org/abs/1303.3997v2>
- 824 Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics (Oxford, England)*,
825 34(18), 3094–3100. doi: 10.1093/bioinformatics/bty191
- 826 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... 1000 Genome Project Data Processing
827 Subgroup. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics (Oxford,*
828 *England)*, 25(16), 2078–2079. doi: 10.1093/bioinformatics/btp352
- 829 Li, M., Sun, Y., Zhao, J., Shi, H., Zeng, S., Ye, K., ... Wang, D. (2015). A Tandem Duplicate of Anti-Müllerian
830 Hormone with a Missense SNP on the Y Chromosome Is Essential for Male Sex Determination in Nile
831 Tilapia, Oreochromis niloticus. *PLOS Genetics*, 11(11), e1005678. doi:
832 10.1371/journal.pgen.1005678
- 833 Lischer, H. E. L., & Shimizu, K. K. (2017). Reference-guided de novo assembly approach improves genome
834 reconstruction for related species. *BMC Bioinformatics*, 18(1), 474. doi: 10.1186/s12859-017-1911-6
- 835 Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-
836 seq data with DESeq2. *Genome Biology*, 15(12), 550. doi: 10.1186/s13059-014-0550-8
- 837 Mank, J. E., & Avise, J. C. (2009). Evolutionary diversity and turn-over of sex determination in teleost fishes.
838 *Sexual Development: Genetics, Molecular Biology, Evolution, Endocrinology, Embryology, and*
839 *Pathology of Sex Determination and Differentiation*, 3(2–3), 60–67. doi: 10.1159/000223071
- 840 Marçais, G., & Kingsford, C. (2011). A fast, lock-free approach for efficient parallel counting of occurrences
841 of k-mers. *Bioinformatics (Oxford, England)*, 27(6), 764–770. doi: 10.1093/bioinformatics/btr011
- 842 Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
843 *EMBnet.Journal*, 17(1), 10–12. doi: 10.14806/ej.17.1.200
- 844 Matsuda, M., Nagahama, Y., Shinomiya, A., Sato, T., Matsuda, C., Kobayashi, T., ... Sakaizumi, M. (2002).
845 DMY is a Y-specific DM-domain gene required for male development in the medaka fish. *Nature*,
846 417(6888), 559–563. doi: 10.1038/nature751
- 847 Meng-Umphang, K. (2009). Growth Performance, Sex Hormone Levels and Maturation Ability of Pla Pho
848 (Pangasius bocourti) Fed with Spirulina Supplementary Pellet and Hormone Application. *Int. J. Agric.*
849 *Biol.*, 11(4), 5.
- 850 Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler, A., & Lanfear,
851 R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic
852 Era. *Molecular Biology and Evolution*, 37(5), 1530–1534. doi: 10.1093/molbev/msaa015
- 853 Moore, E. C., & Roberts, R. B. (2013). Polygenic sex determination. *Current Biology: CB*, 23(12), R510–512.
854 doi: 10.1016/j.cub.2013.04.004
- 855 Morgulis, A., Gertz, E. M., Schäffer, A. A., & Agarwala, R. (2006). A fast and symmetric DUST
856 implementation to mask low-complexity DNA sequences. *Journal of Computational Biology: A*
857 *Journal of Computational Molecular Cell Biology*, 13(5), 1028–1040. doi: 10.1089/cmb.2006.13.1028
- 858 Myosho, T., Otake, H., Masuyama, H., Matsuda, M., Kuroki, Y., Fujiyama, A., ... Sakaizumi, M. (2012).
859 Tracing the emergence of a novel sex-determining gene in medaka, Oryzias luzonensis. *Genetics*,
860 191(1), 163–170. doi: 10.1534/genetics.111.137497
- 861 Nakamoto, M., Uchino, T., Koshimizu, E., Kuchiishi, Y., Sekiguchi, R., Wang, L., ... Sakamoto, T. (2021). A
862 Y-linked anti-Müllerian hormone type-II receptor is the sex-determining gene in ayu, Plecoglossus

- altivelis. *PLoS Genetics*, 17(8), e1009705. doi: 10.1371/journal.pgen.1009705
- Na-Nakorn, U., & Moeikum, T. (2009). Genetic diversity of domesticated stocks of striped catfish, *Pangasianodon hypophthalmus* (Sauvage 1878), in Thailand: Relevance to broodstock management regimes. *Aquaculture*, 297(1/4), 70–77.
- Na-Nakorn, U., Sukmanom, S., Nakajima, M., Taniguchi, N., Kamonrat, W., Poompuang, S., & Nguyen, T. T. T. (2006). MtDNA diversity of the critically endangered Mekong giant catfish (*Pangasianodon gigas* Chevey, 1913) and closely related species: Implications for conservation. *Animal Conservation*, 9(4), 483–494. doi: 10.1111/j.1469-1795.2006.00064.x
- Nanda, I., Kondo, M., Hornung, U., Asakawa, S., Winkler, C., Shimizu, A., ... Scharl, M. (2002). A duplicated copy of DMRT1 in the sex-determining region of the Y chromosome of the medaka, *Oryzias latipes*. *Proceedings of the National Academy of Sciences of the United States of America*, 99(18), 11778–11783. doi: 10.1073/pnas.182314699
- Ospina-Alvarez, N., & Piferrer, F. (2008). Temperature-dependent sex determination in fish revisited: Prevalence, a single sex ratio response pattern, and possible effects of climate change. *PloS One*, 3(7), e2837. doi: 10.1371/journal.pone.0002837
- Pan, Q., Anderson, J., Bertho, S., Herpin, A., Wilson, C., Postlethwait, J. H., ... Guiguen, Y. (2016). Vertebrate sex-determining genes play musical chairs. *Comptes Rendus Biologies*, 339(7–8), 258–262. doi: 10.1016/j.crv.2016.05.010
- Pan, Q., Feron, R., Yano, A., Guyomard, R., Jouanno, E., Vigouroux, E., ... Guiguen, Y. (2019). Identification of the master sex determining gene in Northern pike (*Esox lucius*) reveals restricted sex chromosome differentiation. *PLOS Genetics*, 15(8), e1008013. doi: 10.1371/journal.pgen.1008013
- Pan, Q., Kay, T., Depincé, A., Adolphi, M., Scharl, M., Guiguen, Y., & Herpin, A. (2021). Evolution of master sex determiners: TGF- β signalling pathways at regulatory crossroads. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 376(1832), 20200091. doi: 10.1098/rstb.2020.0091
- Pan, Z.-J., Li, X.-Y., Zhou, F.-J., Qiang, X.-G., & Gui, J.-F. (2015). Identification of Sex-Specific Markers Reveals Male Heterogametic Sex Determination in *Pseudobagrus ussuriensis*. *Marine Biotechnology (New York, N.Y.)*, 17(4), 441–451. doi: 10.1007/s10126-015-9631-2
- Pasquier, J., Cabau, C., Nguyen, T., Jouanno, E., Severac, D., Braasch, I., ... Bobe, J. (2016). Gene evolution and gene expression after whole genome duplication in fish: The PhyloFish database. *BMC Genomics*, 17, 368. doi: 10.1186/s12864-016-2709-z
- Phuong, N. T., & Oanh, D. T. H. (2010). Striped Catfish Aquaculture in Vietnam: A Decade of Unprecedented Development. In S. S. De Silva & F. B. Davy (Eds.), *Success Stories in Asian Aquaculture* (pp. 131–147). Dordrecht: Springer Netherlands. doi: 10.1007/978-90-481-3087-0_7
- Pouyaud, L., Gustiano, R., & Teugels, G. G. (2016). contribution to the phylogeny of the Pangasiidae based on mitochondrial 12S rDNA. *Indonesian Journal of Agricultural Science*, 5(2), 4562. doi: 10.21082/ijas.v5n2.2004.p4562
- Pryszcz, L. P., & Gabaldón, T. (2016). Redundans: An assembly pipeline for highly heterozygous genomes. *Nucleic Acids Research*, 44(12), e113–e113. doi: 10.1093/nar/gkw294
- Purcell, C. M., Seetharam, A. S., Snodgrass, O., Ortega-García, S., Hyde, J. R., & Severin, A. J. (2018). Insights into teleost sex determination from the *Seriola dorsalis* genome assembly. *BMC Genomics*, 19(1), 31. doi: 10.1186/s12864-017-4403-1
- Qu, M., Liu, Y., Zhang, Y., Wan, S., Ravi, V., Qin, G., ... Lin, Q. (2021). Seadragon genome analysis provides insights into its phenotype and sex determination locus. *Science Advances*, 7(34), eabg5196. doi: 10.1126/sciadv.abg5196
- Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics (Oxford, England)*, 26(6), 841–842. doi: 10.1093/bioinformatics/btq033
- Rafati, N., Chen, J., Herpin, A., Pettersson, M. E., Han, F., Feng, C., ... Andersson, L. (2020). Reconstruction of the birth of a male sex chromosome present in Atlantic herring. *Proceedings of the National Academy of Sciences*, 117(39), 24359–24368. doi: 10.1073/pnas.2009925117
- Reichwald, K., Petzold, A., Koch, P., Downie, B. R., Hartmann, N., Pietsch, S., ... Platzter, M. (2015). Insights into Sex Chromosome Evolution and Aging from the Genome of a Short-Lived Fish. *Cell*, 163(6), 1527–1538. doi: 10.1016/j.cell.2015.10.071
- Roberts, T. R., & Vidhayanon, C. (1991). Systematic Revision of the Asian Catfish Family Pangasiidae, with Biological Observations and Descriptions of Three New Species. *Proceedings of the Academy of*

- 918 *Natural Sciences of Philadelphia*, 143, 97–143.
- 919 Rondeau, E. B., Messmer, A. M., Sanderson, D. S., Jantzen, S. G., von Schalburg, K. R., Minkley, D. R., ...
 920 Koop, B. F. (2013). Genomics of sablefish (*Anoplopoma fimbria*): Expressed genes, mitochondrial
 921 phylogeny, linkage map and identification of a putative sex gene. *BMC Genomics*, 14(1), 452. doi:
 922 10.1186/1471-2164-14-452
- 923 Ruan, J. (2019). *Ultra-fast de novo assembler using long noisy reads: Ruanjue/smartdenovo* [C]. Retrieved
 924 from <https://github.com/ruanjue/smartdenovo> (Original work published 2015)
- 925 Ruan, J., & Li, H. (2020). Fast and accurate long-read assembly with wtdbg2. *Nature Methods*, 17(2), 155–
 926 158. doi: 10.1038/s41592-019-0669-3
- 927 Sela, I., Ashkenazy, H., Katoh, K., & Pupko, T. (2015). GUIDANCE2: Accurate detection of unreliable
 928 alignment regions accounting for the uncertainty of multiple parameters. *Nucleic Acids Research*,
 929 43(W1), W7-14. doi: 10.1093/nar/gkv318
- 930 Shumate, A., & Salzberg, S. L. (2021). Liftoff: Accurate mapping of gene annotations. *Bioinformatics*, 37(12),
 931 1639–1643. doi: 10.1093/bioinformatics/btaa1016
- 932 Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO:
 933 Assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*,
 934 31(19), 3210–3212. doi: 10.1093/bioinformatics/btv351
- 935 Singh, A. K., & Lakra, W. S. (2012). Culture of Pangasianodon hypophthalmus into India: Impacts and present
 936 scenario. *Pakistan Journal of Biological Sciences: PJBS*, 15(1), 19–26. doi: 10.3923/pjbs.2012.19.26
- 937 Slater, G. S. C., & Birney, E. (2005). Automated generation of heuristics for biological sequence comparison.
 938 *BMC Bioinformatics*, 6, 31. doi: 10.1186/1471-2105-6-31
- 939 Song, W., Xie, Y., Sun, M., Li, X., Fitzpatrick, C. K., Vaux, F., ... He, Y. (2021). A duplicated amh is the
 940 master sex-determining gene for *Sebastes rockfish* in the Northwest Pacific. *Open Biology*, 11(7),
 941 210063. doi: 10.1098/rsob.210063
- 942 Sreeputhorn, K., Mangumphan, K., Muanphet, B., Tanomtong, A., Supiwong, W., & Kaewmad, P. (2017).
 943 *The First Report on Chromosome Analysis of F1 Hybrid Catfish: Mekong Giant Catfish (*
 944 *Pangasianodon gigas**) × Striped Catfish (**Pangasianodon hypophthalmus**) and Spot Pangasius (**Pangasius*
 945 *larnaudii**) × Pangasianodon hypophthalmus (Siluriformes, Pangasiidae)*. doi:
 946 10.1508/CYTOLOGIA.82.457
- 947 Sriphairoj, K., Na-Nakorn, U., Brunelli, J. P., & Thorgaard, G. H. (2007). No AFLP sex-specific markers
 948 detected in *Pangasianodon gigas* and *P. hypophthalmus*. *Aquaculture*, 273(4), 739–743. doi:
 949 10.1016/j.aquaculture.2007.09.018
- 950 Stanke, M., Keller, O., Gunduz, I., Hayes, A., Waack, S., & Morgenstern, B. (2006). AUGUSTUS: Ab initio
 951 prediction of alternative transcripts. *Nucleic Acids Research*, 34(Web Server issue), W435-439. doi:
 952 10.1093/nar/gkl200
- 953 Sullivan, J. P., Lundberg, J. G., & Hardman, M. (2006). A phylogenetic analysis of the major groups of
 954 catfishes (Teleostei: Siluriformes) using rag1 and rag2 nuclear gene sequences. *Molecular*
 955 *Phylogenetics and Evolution*, 41(3), 636–662. doi: 10.1016/j.ympev.2006.05.044
- 956 Takehana, Y., Hamaguchi, S., & Sakaizumi, M. (2008). Different origins of ZZ/ZW sex chromosomes in
 957 closely related medaka fishes, *Oryzias javanicus* and *O. hubbsi*. *Chromosome Research: An*
 958 *International Journal on the Molecular, Supramolecular and Evolutionary Aspects of Chromosome*
 959 *Biology*, 16(5), 801–811. doi: 10.1007/s10577-008-1227-5
- 960 Takehana, Y., Matsuda, M., Myosho, T., Suster, M. L., Kawakami, K., Shin-I, T., ... Naruse, K. (2014). Co-
 961 option of Sox3 as the male-determining factor on the Y chromosome in the fish *Oryzias dancena*.
 962 *Nature Communications*, 5, 4157. doi: 10.1038/ncomms5157
- 963 Trapnell, C., Williams, B. A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M. J., ... Pachter, L. (2010).
 964 Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform
 965 switching during cell differentiation. *Nature Biotechnology*, 28(5), 511–515. doi: 10.1038/nbt.1621
- 966 Vaser, R., Sović, I., Nagarajan, N., & Šikić, M. (2017). Fast and accurate de novo genome assembly from long
 967 uncorrected reads. *Genome Research*, 27(5), 737–746. doi: 10.1101/gr.214270.116
- 968 Villela, L. C. V., Alves, A. L., Varela, E. S., Yamagishi, M. E. B., Giachetto, P. F., da Silva, N. M. A., ...
 969 Caetano, A. R. (2017). Complete mitochondrial genome from South American catfish
 970 *Pseudoplatystoma reticulatum* (Eigenmann & Eigenmann) and its impact in Siluriformes phylogenetic
 971 tree. *Genetica*, 145(1), 51–66. doi: 10.1007/s10709-016-9945-7
- 972 Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H., Gurtowski, J., & Schatz, M. C.

- (2017). GenomeScope: Fast reference-free genome profiling from short reads. *Bioinformatics (Oxford, England)*, 33(14), 2202–2204. doi: 10.1093/bioinformatics/btx153
- Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., ... Earl, A. M. (2014). Pilon: An integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PloS One*, 9(11), e112963. doi: 10.1371/journal.pone.0112963
- Wang, D., Mao, H.-L., Chen, H.-X., Liu, H.-Q., & Gui, J.-F. (2009). Isolation of Y- and X-linked SCAR markers in yellow catfish and application in the production of all-male populations. *Animal Genetics*, 40(6), 978–981. doi: 10.1111/j.1365-2052.2009.01941.x
- Yang, Z., Nielsen, R., Goldman, N., & Pedersen, A. M. (2000). Codon-substitution models for heterogeneous selection pressure at amino acid sites. *Genetics*, 155(1), 431–449. doi: 10.1093/genetics/155.1.431
- Yano, A., Guyomard, R., Nicol, B., Jouanno, E., Quillet, E., Klopp, C., ... Guiguen, Y. (2012). An Immune-Related Gene Evolved into the Master Sex-Determining Gene in Rainbow Trout, *Oncorhynchus mykiss*. *Current Biology*, 22(15), 1423–1428. doi: 10.1016/j.cub.2012.05.045
- Yano, A., Nicol, B., Jouanno, E., Quillet, E., Fostier, A., Guyomard, R., & Guiguen, Y. (2013). The sexually dimorphic on the Y-chromosome gene (sdY) is a conserved male-specific Y-chromosome sequence in many salmonids. *Evolutionary Applications*, 6(3), 486–496. doi: 10.1111/eva.12032
- Zhang, J., Nielsen, R., & Yang, Z. (2005). Evaluation of an improved branch-site likelihood method for detecting positive selection at the molecular level. *Molecular Biology and Evolution*, 22(12), 2472–2479. doi: 10.1093/molbev/msi237

AUTHOR CONTRIBUTIONS

YG, and JHP designed the project. JCA, RD, MC, TTTH, RG, KS, JR and FLA collected the samples, EJ, MW, CI, AC, CR, OB, SV, CL, CP, EB, VG and HA extracted the gDNA, made the genomic libraries and sequenced them. CC, CK, MZ, MW, QP and YG processed the genome assemblies and / or analyzed the results. TD, JM and JB processed and analyzed the small RNA sequencing data for miRNA analysis. CFB, MW, QP and MRR performed phylogenetic analyses. CFB and MRR performed the selection analysis. MW, JHP, CC, CK, CR, QP and YG wrote the manuscript with inputs from all other coauthors. JHP, CD, JB and YG, supervised the project administration and raised funding. All the authors read and approved the final manuscript.

COMPETING INTERESTS

All authors declare no competing interests.

FIGURES




Figure 1. Sex genotyping in *P. hypophthalmus*. The *amhr2a* sequence (upper panel) is PCR amplified in both male and female samples, while the *amh2by* sequence (bottom panel) is only amplified in male samples, indicating that *amhr2by* is male-specific i.e., Y-chromosome linked.





Figure 2. Structure of *amhr2a* and *amhr2by* and deduced proteins in *P. hypophthalmus*. (A) Identity plot of the alignment of the autosomal *amhr2a* with the Y-linked *amhr2by* sequences. Exons (E) of both *amhr2* genes are depicted with gray boxes. (B) Clustal W alignment of Amhr2a and Amhr2by proteins. Identical amino acids are shaded in gray and conserved cysteines in the extracellular domain of Amhr2a are highlighted in yellow. The different domains (signal peptide, cysteine knot and transmembrane domain) of the receptors are boxed. Intron-exon boundaries are boxed in black for both receptors. (C) Schematic representation of *P. hypophthalmus* autosomal Amhr2a and Y-linked Amhr2bY proteins showing the architecture of Amh receptors and the correspondence between exons of Amhr2a and Amhr2by, highlighting the absence of the entire extracellular domain in the truncated Amhr2bY.






Figure 4: Whole-genome-based phylogenetic tree of all sequenced Pangasiid species. Maximum-likelihood phylogeny of 12 Pangasiidae species with *Ictalurus punctatus* (siluriformes) as a Pangasiidae outgroup, based on alignment of concatenated protein sequences. Branch length scale corresponds to 0.02 amino acid substitutions per site. Support values at each node are proportions of 100 standard non-parametric bootstrap replicates.






Figure 5. Phylogeny of *amhr2* in catfishes reveals an ancient *amhr2a* / *amhr2b* duplication in Siluroidei. Maximum-likelihood phylogeny of *amhr2* coding sequences (see Supplementary Figure 1 for other phylogenetic approaches) from 28 catfish species with *amhr2* coding sequences from *Astyanax mexicanus* (Characiformes) and *Electrophorus electricus* (Gymnotiformes) as Siluriformes outgroups. Family and suborders are given for all catfish species on the right panel of the figure. The *amhr2b* cluster including the *amhr2b* of Pangasiids is shaded in purple, the *amhr2a* cluster shaded in red, and the *Corydorasp amhr2* pre-duplication is shaded in yellow. The branch length scale representing the number of substitutions per site is given at the root of the Siluriformes tree. Bootstrap values are given only for values over 80 and are inserted in a white circle at key nodes for the Siluroidei *amhr2* duplication.



Supplementary Figure 1. Relative genome read coverage around the *amhr2a* and *amhr2by* loci in 12 Pangasiids supports hemizyosity of *amhr2by* in some species. (A-L) Relative average read coverage was deduced from each species short-read remapping on the *P. hypophthalmus* genome reference and is shown in blue for the autosomal *amhr2a* locus and in red for the *amhr2by* locus (left and right side respectively of each species panel). Half read coverage compared to genome average was detected around the *amhr2by* locus compared to the *amhr2a* locus in the male genomes of *P. hypophthalmus*, *P. gigas*, *P. djambal*, *P. conchophilus*, and *P. bocourti* (A-E) and in the unknown sex genomes of *P. sanitwongsei*, and *P. krempfi* (F-G), supporting hemizyosity of *amhr2by* in these species as it would be expected for a Y chromosomal gene. In *P. elongatus*, *P. siamensis*, *P. macronema*, *P. larnaudii*, and *P. mekongensis* (H-L) no reads were significantly remapped on the *P. hypophthalmus* *amhr2by* locus either because these sequenced individuals are XX females without a Y chromosome *amhr2by* gene, or because these species have lost *amhr2by* as a Y chromosome gene.



Supplementary Figure 2. Phylogenies of Amhr2 / amhr2 in catfishes support an ancient amhr2a / amhr2b duplication in Siluroidei. Maximum-likelihood (A, B, C) and Bayesian (D, E, F) phylogenies of Amhr2 proteins (A, D), amhr2 coding (CDS) sequences (B, E) and amhr2 CDS sequences with the third codon removed (C, F) from 28 catfish species with sequences from *Astyanax mexicanus* (Characiformes) and *Electrophorus electricus* (Gymnotiformes) as Siluriformes outgroups. The amhr2b cluster including the amhr2by of Pangasiids is shaded in purple, the amhr2a cluster shaded in red, and the *Corydoras sp* amhr2 pre-duplication is shaded in yellow. The branch length scale representing the number of substitutions per site is given below each tree. Bootstrap values are given only for values over 80 except at key nodes for the Siluroidei amhr2 duplication (white circles).

1090 TABLES

1091 **Table 1: Comparison of our *P. hypophthalmus* reference genome assembly metrics (our study)**
1092 **with the other *P. hypophthalmus* available assemblies.**

Assemblies	GCA_003671635.1	Our study	GCA_016801045.1
Release date	05/04/2018	10/22/2019	14/10/2020
Sex of the sequenced individual	male	male	female
Total sequence length	715.8 Mb	758.9 Mb	742.5 Mb
Total ungapped length	696.5 Mb	758.8 Mb	742.3 Mb
Number of contigs	23,34	612	808
Contig N50	0.06 Mb	16.19 Mb	3.48 Mb
Contig L50	3,254	18	63
Total number of chromosomes	N.A	30	30
Number of component sequences (WGS or clone)	568	150	402

1093 N.A = Not Applicable

1094 **Table 2: Sex-linkage of *amhr2by* in five different Pangasiid species.** Associations between
1095 *amhr2by* specific PCR amplifications and sex phenotypes are provided for both males and females
1096 (number of positive individuals for *amhr2by*/total number of individuals) along with the p value of
1097 association with sex that was calculated for each species based on the Pearson's Chi-square test with
1098 Yates' continuity correction.

Species	males	females	p value
<i>Pangasianodon hypophthalmus</i>	12/12	1/11	7.12e-05
<i>Pangasianodon gigas</i>	3/3	1/3	0.3865*
<i>Pangasius bocourti</i>	12/12	1/20	8.411e-07
<i>Pangasius conchophilus</i>	22/22	0/10	1.559e-07
<i>Pangasius djambal</i>	6/6	0/9	8.528e-04

1099 * non-significant association with sex

1100 **Table 3: Positive selection analyses reveal no significant signal of positive selection on Pangasiid**
1101 ***amhr2*.** P-values were computed using a chi-square distribution with 1 degree of freedom. None of
1102 the p-values passed a Bonferroni corrected limit of significance: $0.05/3 = 0.0167$. DlnL = difference
1103 in log-likelihood between models with and without positive selection; likelihood ratio test statistic.

Model	Conserved exons		First exons	
	DlnL	p-value	DlnL	p-value
M8 gamma	0.0000000	0.5000000	3.37482	3.309990e-02
Branch-site gamma	0.2976536	0.2926786	-	-

1104

Supplementary Table 1: Genome assembly characteristics and annotation metrics of 12 Pangasiid species.

Species	Sex	Sequencing/assembly	Guided assembly	N	G.S (Gb)	Max (Mb)	N50 (Mb)	L50	% Chr	Annotation	Buscos (C)	Buscos (S)	Buscos (D)	Buscos (F)	Buscos (M)
<i>Pangasianodon hypophthalmus</i>	male	ONT, 10X, Hi-C Smartdenovo/lonranger/juicer	N.A	612	0.759	35.6	26.16	13	99.2	de novo / GenBank	96.6	95.6	1.0	1.0	2.4
<i>Pangasianodon gigas</i>	male	Illumina 2x 250 bp Discovar de novo	Dgenies	283151	0.841	35.47	26.7	12	89.0	de novo / in house					
<i>Pangasius djambal</i>	male	Illumina 2x 250 bp Discovar de novo	Dgenies	415588	0.867	34.66	28.07	11	82.7	de novo / in house					
<i>Pangasius conchophilus</i>	male	Illumina 2x150 bp SPADes v.3.11.1 /purge_dup	Dgenies	937794	0.815	30.1	17.57	18	73.0	Lifted from <i>P. hypophthalmus</i>	81.9	80.7	1.2	6.2	11.9
<i>Pangasius bocourti</i>	male	Illumina 2x150 bp SPADes v.3.11.1 /purge_dup	Dgenies	740730	0.780	33.3	24.19	14	86.7	Lifted from <i>P. hypophthalmus</i>	95.5	94.6	0.9	1.4	3.1
<i>Pangasius elongatus</i>	U	Illumina 2x150 bp SPADes v.3.14.1 /redundans v0.14a	Dgenies	126560	0.712	31.0	22.43	14	87.6	Lifted from <i>P. hypophthalmus</i>	90.7	89.8	0.9	3.1	6.2
<i>Pangasius siamensis</i>	U	Illumina 2x150 bp SPADes v.3.14.1 /redundans v0.14a	Dgenies	53159	0.685	32.4	23.59	13	95.4	Lifted from <i>P. hypophthalmus</i>	93.0	92.0	1.0	2.1	4.9
<i>Pangasius sanitwongsei</i>	U	Illumina 2x150 bp SPADes v.3.11.1 /purge_dup	Dgenies	387468	0.743	34.1	24.55	14	92.4	Lifted from <i>P. hypophthalmus</i>	96.3	95.6	0.7	1.1	2.6
<i>Pangasius macronema</i>	U	Illumina 2x150 bp SPADes v.3.14.1 /redundans v0.14a	Dgenies	42101	0.683	32.4	23.73	13	95.9	Lifted from <i>P. hypophthalmus</i>	93.2	91.9	1.3	2.3	4.5
<i>Pangasius larnaudii</i>	U	Illumina 2x150 bp SPADes v.3.11.1 /purge_dup	Dgenies	375123	0.733	33.0	23.84	14	91.3	Lifted from <i>P. hypophthalmus</i>	95.6	94.8	0.8	1.5	2.9
<i>Pangasius mekongensis</i>	U	Illumina 2x150 bp SPADes v.3.11.1 /purge_dup	Dgenies	443231	0.766	33.7	24.28	14	88.8	Lifted from <i>P. hypophthalmus</i>	93.6	92.8	0.8	2.5	3.9
<i>Pangasius krempfi</i>	U	Illumina 2x150 bp SPADes v.3.11.1 /purge_dup	Dgenies	448669	0.739	33.5	24.31	14	91.3	Lifted from <i>P. hypophthalmus</i>	95.1	94.2	0.9	1.8	3.1

Sex = phenotypic sex of the animal sequenced (U= unknown), N= Number of contigs, G.S = genome assembly size (kb), Max = size of the longest scaffold, N50 = scaffold N50 (Mb), L50 = scaffold L50, % Chr = percentage of the assembly in chromosomes, Buscos (V4, in genome mode with actinopterygii lineage) score in percentage (C = Complete, S = Single copy, D = Duplicated, F = Fragmented, M = Missing). N.A = Not Applicable.

Supplementary Table 2: Origin of the catfish *amhr2* sequences used for phylogenetic analyses.

Species	Family	Sub-order	order	Sex	Gene	Source	Sequences deduced from
<i>Pangasianodon hypophthalmus</i>	Pangasiidae	Siluroidei	Siluriformes	male	<i>amhr2a</i> <i>amhr2by</i>	This study	Genome annotation
<i>Pangasianodon gigas</i>	Pangasiidae	Siluroidei	Siluriformes	male	<i>amhr2a</i> <i>amhr2by</i>	This study	Genome annotation
<i>Pangasius djambal</i>	Pangasiidae	Siluroidei	Siluriformes	male	<i>amhr2a</i> <i>amhr2by</i>	This study	Genome annotation
<i>Pangasius conchophilus</i>	Pangasiidae	Siluroidei	Siluriformes	male	<i>amhr2a</i> <i>amhr2by</i>	This study	Genome annotation
<i>Pangasius bocourti</i>	Pangasiidae	Siluroidei	Siluriformes	male	<i>amhr2a</i> <i>amhr2by</i>	This study	Genome annotation
<i>Pangasius elongatus</i>	Pangasiidae	Siluroidei	Siluriformes	unknown	<i>amhr2a</i>	This study	Genome annotation
<i>Pangasius siamensis</i>	Pangasiidae	Siluroidei	Siluriformes	unknown	<i>amhr2a</i>	This study	Genome annotation
<i>Pangasius sanitwongsei</i>	Pangasiidae	Siluroidei	Siluriformes	unknown	<i>amhr2a</i> <i>amhr2by</i>	This study	Genome annotation
<i>Pangasius macronema</i>	Pangasiidae	Siluroidei	Siluriformes	unknown	<i>amhr2a</i>	This study	Genome annotation
<i>Pangasius larnaudii</i>	Pangasiidae	Siluroidei	Siluriformes	unknown	<i>amhr2a</i>	This study	Genome annotation
<i>Pangasius mekongensis</i>	Pangasiidae	Siluroidei	Siluriformes	unknown	<i>amhr2a</i>	This study	Genome annotation
<i>Pangasius krempfi</i>	Pangasiidae	Siluroidei	Siluriformes	unknown	<i>amhr2a</i> <i>amhr2by</i>	This study	Genome annotation
<i>Clarias batrachus</i>	Clariidae	Siluroidei	Siluriformes	unknown	<i>amhr2a</i>	GCA_003987875.1	inferred from genome assembly
<i>Clarias macrocephalus</i>	Clariidae	Siluroidei	Siluriformes	female	<i>amhr2a</i>	GCA_011419295.1	inferred from genome assembly
<i>Clarias magur</i>	Clariidae	Siluroidei	Siluriformes	male	<i>amhr2a</i>	GCA_013621035.1	inferred from genome assembly
<i>Bagarius yarrelli</i>	Sisoridae	Siluroidei	Siluriformes	female	<i>amhr2a</i>	GCA_005784505.1	inferred from genome assembly
<i>Glyptosternon maculatum</i>	Sisoridae	Siluroidei	Siluriformes	female	<i>amhr2a</i>	http://gigadb.org/dataset/view/id/100489	inferred from genome assembly
<i>Silurus glanis</i>	Siluridae	Siluroidei	Siluriformes	female	<i>amhr2a</i>	GCA_014706435.1	inferred from genome assembly
<i>Silurus meridionalis</i>	Siluridae	Siluroidei	Siluriformes	female	<i>amhr2a</i>	GCA_014805685.1	KAF7704051.1
<i>Ompok bimaculatus</i>	Siluridae	Siluroidei	Siluriformes	unknown	<i>amhr2a</i>	GCA_009108245.1	inferred from genome assembly
<i>Hemibagrus wyckioides</i>	Bagridae	Siluroidei	Siluriformes	female	<i>amhr2a</i>	GCA_019097595.1	KAG7327988.1
<i>Tachysurus fulvidraco</i>	Bagridae	Siluroidei	Siluriformes	female	<i>amhr2a</i>	GCF_003724035.1	XP_027015428.1
<i>Ageneiosus marmoratus</i>	Auchenipteridae	Siluroidei	Siluriformes	male	<i>amhr2a</i>	GCA_003347165.1	inferred from genome assembly
<i>Heteropneustes fossilis</i>	Heteropneustidae	Siluroidei	Siluriformes	male	<i>amhr2a</i>	unpublished	Inferred from transcriptome information
<i>Ameiurus melas</i>	Ictaluridae	Siluroidei	Siluriformes	male	<i>amhr2a</i>	GCA_012411365.1	KAF4083677.1
<i>Ictalurus punctatus</i>	Ictaluridae	Siluroidei	Siluriformes	female	<i>amhr2a</i>	GCF_001660625.1	XP_017331275.1
<i>Pimelodus maculatus</i>	Pimelodidae	Siluroidei	Siluriformes	male	<i>amhr2a</i> <i>amhr2b</i>	unpublished	inferred from genome assembly
<i>Corydoras sp C115</i>	Callichthyidae	Loricarioidei	Siluriformes	unknown	<i>amhr2</i>	GCA_019802505.1	inferred from genome assembly
<i>Electrophorus electricus</i>	Gymnotidae	NR	Gymnotiformes	unknown	<i>amhr2</i>	GCF_013358815.1	XP_035376390.1
<i>Astyanax mexicanus</i>	Stethaprioninae	NR	Characiformes	female	<i>amhr2</i>	GCF_000372685.2	XP_022538368.1

NR : not relevant