

Towards understanding diversity, endemism and global change vulnerability of soil fungi

Leho Tedersoo^{1*}, Vladimir Mikryukov², Alexander Zizka³, Mohammad Bahram⁴, Niloufar Hagh-Doust², Sten Anslan², Oleh Prylutskyi⁵, Manuel Delgado-Baquerizo⁶, Fernando T. Maestre⁷, Jaan Pärn², Maarja Öpik², Mari Moora², Martin Zobel², Mikk Espenberg², Ülo Mander², Abdul Nasir Khalid⁸, Adriana Corrales⁹, Ahto Agan¹⁰, Aída-M. Vasco-Palacios¹¹, Alessandro Saitta¹², Andrea C. Rinaldi¹³, Annemieke Verbeken¹⁴, Bobby P. Sulistyo¹⁵, Boris Tamgnoue¹⁶, Brendan Furneaux¹⁷, Camila Duarte Ritter¹⁸, Casper Nyamukondiwa¹⁹, Cathy Sharp²⁰, César Marín²¹, Daniyal Gohar¹, Darta Klavina²², Dipon Sharmah²³, Dong Qin Dai²⁴, Eduardo Nouhra²⁵, Elisabeth Machteld Biersma²⁶, Elisabeth Rähn¹⁰, Erin K. Cameron²⁷, Eske De Crop¹⁴, Eveli Otsing¹, Evgeny A. Davydov²⁸, Felipe E. Albornoz²⁹, Francis Q. Brearley³⁰, Franz Buegger³¹, Geoffrey Zahn³², Gregory Bonito³³, Inga Hiiesalu², Isabel C. Barrio³⁴, Jacob Heilmann-Clausen³⁵, Jelena Ankuda³⁶, John Y. Kupagme¹, Jose G. Maciá-Vicente³⁷, Joseph Djeugap Fovo¹⁶, József Geml³⁸, Juha M. Alatalo³⁹, Julieta Alvarez-Manjarrez⁴⁰, Kadri Pöldmaa², Kadri Runnel², Kalev Adamson¹⁰, Kari Anne Bråthen⁴¹, Karin Pritsch³¹, Kassim I. Tchan⁴², Kęstutis Armolaitis³⁶, Kevin D. Hyde⁴³, Kevin K. Newsham⁴⁴, Kristel Panksep⁴⁵, Adebola A. Lateef⁴⁶, Liis Tiirmann¹, Linda Hansson⁴⁷, Louis J. Lamit^{48,49}, Malka Saba⁵⁰, Maria Tuomi⁴¹, Marieka Gryzenhout⁵¹, Marijn Bauters⁵², Meike Piepenbring⁵³, Nalin Wijayawardene²⁴, Nourou S. Yorou⁴², Olavi Kurina⁵⁴, Peter E. Mortimer⁵⁵, Peter Meidl⁵⁶, Petr Kohout⁵⁷, R. Henrik Nilsson⁵⁸, Rasmus Puusepp¹, Rein Drenkhan¹⁰, Roberto Garibay-Orijel⁵⁹, Roberto Godoy⁶⁰, Saad Alkahtani⁶¹, Saleh Rahimlou¹, Sergey V. Dudov⁶², Sergei Pölme¹, Soumya Ghosh⁵¹, Sunil Mundra⁶³, Talaat Ahmed³⁹, Tarquin Netherway⁴, Terry W. Henkel⁶⁴, Tomas Roslin⁴, Vincent Nteziryayo⁶⁵, Vladimir E. Fedosov⁶², Vladimir G. Onipchenko⁶², W. A. Erandi Yasanthika⁴³, Young Woon Lim⁶⁶, Nadejda A. Soudzilovskaia⁶⁷, Alexandre Antonelli⁶⁸, Urmas Kõljalg², Kessy Abarenkov⁶⁹

¹Center of Mycology and Microbiology, University of Tartu, Tartu, Estonia

²Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

³Department of Biology, Philipps-University, Marburg, Germany

⁴Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

⁵Department of Mycology and Plant Resistance, School of Biology, V.N. Karazin Kharkiv National University, Kharkiv, Ukraine

⁶Instituto de Recursos Naturales y Agrobiología de Sevilla (IRNAS), CSIC, Sevilla, Spain

⁷Instituto Multidisciplinar para el Estudio del Medio ‘Ramón Margalef’ and Departamento de Ecología, Universidad de Alicante; 03690, Alicante, Spain

⁸Institute of Botany, University of the Punjab, Lahore, Pakistan

⁹Centro de Investigaciones en Microbiología y Biotecnología-UR (CIMBIUR), Universidad del Rosario, Bogotá, Colombia

- 37 ¹⁰Institute of Forestry and Engineering, Estonian University of Life Sciences, Tartu, Estonia
- 38 ¹¹Escuela de Microbiología, Universidad de Antioquia, Medellín, Antioquia, Colombia
- 39 ¹²Department of Agricultural, Food and Forest Sciences, University of Palermo, Palermo, Italy
- 40 ¹³Department of Biomedical Sciences, University of Cagliari, Cagliari, Italy
- 41 ¹⁴Department Biology, Ghent University, Ghent, Belgium
- 42 ¹⁵Department of Biomedicine, Indonesia International Institute for Life Sciences, Jakarta, Indonesia
- 43 ¹⁶Department of Crop Science, University of Dschang, Dschang, Cameroon
- 44 ¹⁷Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden
- 45 ¹⁸Departamento de Zootecnia, Universidade Federal do Paraná, Curitiba, PR, Brazil
- 46 ¹⁹Department of Biological Sciences and Biotechnology, Botswana International University of
47 Science and Technology, Palapye, Botswana
- 48 ²⁰Natural History Museum of Zimbabwe, Bulawayo, Zimbabwe
- 49 ²¹Centro de Investigación e Innovación para el Cambio Climático (CiiCC), Universidad Santo Tomás,
50 Santiago, Chile
- 51 ²²Latvian State Forest Research Institute Silava, Salaspils, Latvia
- 52 ²³Department of Botany, Jawaharlal Nehru Rajkeeya Mahavidyalaya, Pondicherry University, Port
53 Blair, India
- 54 ²⁴College of Biological Resource and Food Engineering, Qujing Normal University, Qujing, Yunnan,
55 China
- 56 ²⁵Instituto Multidisciplinario de Biología Vegetal (CONICET), Universidad Nacional de Córdoba,
57 Córdoba, Argentina
- 58 ²⁶Natural History Museum of Denmark, Copenhagen, Denmark
- 59 ²⁷Department of Environmental Science, Saint Mary's University, Halifax, Nova Scotia, Canada
- 60 ²⁸Altai State University, Barnaul, Russia
- 61 ²⁹CSIRO Land and Water, Wembley, WA, Australia
- 62 ³⁰Department of Natural Sciences, Manchester Metropolitan University, Manchester, UK
- 63 ³¹Helmholtz Zentrum München, Neuherberg, Germany
- 64 ³²Utah Valley University, Orem UT, USA
- 65 ³³Plant, Soil and Microbial Sciences, Michigan State University, East Lansing MI, USA
- 66 ³⁴Faculty of Natural and Environmental Sciences, Agricultural University of Iceland, Hvanneyri,
67 Iceland
- 68 ³⁵Center for Macroecology, Evolution and Climate, University of Copenhagen, Copenhagen,
69 Denmark
- 70 ³⁶Department of Silviculture and Ecology, Institute of Forestry of Lithuanian Research Centre for
71 Agriculture and Forestry (LAMMC). Girionys, Lithuania
- 72 ³⁷Plant Ecology and Nature Conservation, Wageningen University & Research, Wageningen, The
73 Netherlands

- 74 ³⁸ELKH-EKKE Lendület Environmental Microbiome Research Group, Eszterházy Károly Catholic
75 University, Eger, Hungary
- 76 ³⁹Environmental Science Center, Qatar University, Doha, Qatar
- 77 ⁴⁰Biology Department, Stanford University, Stanford CA, USA
- 78 ⁴¹Department of Arctic and Marine Biology, The Arctic University of Norway, Tromsø, Norway
- 79 ⁴²Research Unit Tropical Mycology and Plants-Soil Fungi Interactions, University of Parakou,
80 Parakou, Benin
- 81 ⁴³Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai, Thailand
- 82 ⁴⁴NERC British Antarctic Survey, High Cross, Cambridge, UK
- 83 ⁴⁵Chair of Hydrobiology and Fishery, Estonian University of Life Sciences, Tartu, Estonia
- 84 ⁴⁶Department of Plant Biology, University of Ilorin, Ilorin, Nigeria
- 85 ⁴⁷Gothenburg Centre for Sustainable Development, Gothenburg, Sweden
- 86 ⁴⁸Department of Biology, Syracuse University, Syracuse NY, USA
- 87 ⁴⁹Department of Environmental and Forest Biology, State University of New York College of
88 Environmental Science and Forestry, Syracuse NY, USA
- 89 ⁵⁰Department of Plant Sciences, Quaid-i-Azam University, Islamabad, Pakistan
- 90 ⁵¹Department of Genetics, University of the Free State, Bloemfontein, South Africa
- 91 ⁵²Department of Environment, Ghent University, Ghent, Belgium
- 92 ⁵³Mycology Working Group, Goethe University Frankfurt am Main, Frankfurt am Main, Germany
- 93 ⁵⁴Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, Tartu,
94 Estonia
- 95 ⁵⁵Center For Mountain Futures, Kunming Institute of Botany, Chinese Academy of Sciences,
96 Kunming, China
- 97 ⁵⁶Freie Universität Berlin, Institut für Biologie, Berlin, Germany
- 98 ⁵⁷Institute of Microbiology, Czech Academy of Sciences, Prague, Czech Republic
- 99 ⁵⁸Gothenburg Global Biodiversity Centre, University of Gothenburg, Gothenburg, Sweden
- 100 ⁵⁹Instituto de Biología, Universidad Nacional Autónoma de México, Ciudad de México, México
- 101 ⁶⁰Instituto Ciencias Ambientales y Evolutivas, Universidad Austral de Chile, Valdivia, Chile
- 102 ⁶¹College of Science, King Saud University, Riyadh, Saudi Arabia
- 103 ⁶²Department of Ecology and Plant Geography, Moscow Lomonosov State University, Moscow,
104 Russia
- 105 ⁶³Department of Biology, College of Science, United Arab Emirates University, Abu Dhabi, UAE
- 106 ⁶⁴Department of Biological Sciences, California State Polytechnic University, Arcata CA, USA
- 107 ⁶⁵Department of Food Science and Technology, University of Burundi, Bujumbura, Burundi
- 108 ⁶⁶School of Biological Sciences and Institute of Microbiology, Seoul National University, Seoul,
109 Korea
- 110 ⁶⁷Centre for Environmental Sciences, Hasselt University, Hasselt, Belgium

⁶⁸Royal Botanic Gardens, Kew, Richmond, United Kingdom

⁶⁹University of Tartu Natural History Museum, Tartu, Estonia

*Corresponding author. Email: leho.tedersoo@ut.ee; tel. +372 56654986

Keywords: biodiversity, climate change, global change vulnerability, conservation priorities, global maps, biogeography, mycorrhizal fungi, pathogens, saprotrophs

Summary

Fungi play pivotal roles in ecosystem functioning, but little is known about their global patterns of diversity, endemism, vulnerability to global change drivers and conservation priority areas. We applied the high-resolution PacBio sequencing technique to identify fungi based on a long DNA marker that revealed a high proportion of hitherto unknown fungal taxa. We used a Global Soil Mycobiome consortium dataset to test relative performance of various sequencing depth standardization methods (calculation of residuals, exclusion of singletons, traditional and SRS rarefaction, use of Shannon index of diversity) to find optimal protocols for statistical analyses. Altogether, we used six global surveys to infer these patterns for soil-inhabiting fungi and their functional groups. We found that residuals of log-transformed richness (including singletons) against log-transformed sequencing depth yields significantly better model estimates compared with most other standardization methods. With respect to global patterns, fungal functional groups differed in the patterns of diversity, endemism and vulnerability to main global change predictors. Unlike α -diversity, endemism and global-change vulnerability of fungi and most functional groups were greatest in the tropics. Fungi are vulnerable mostly to drought, heat, and land cover change. Fungal conservation areas of highest priority include wetlands and moist tropical ecosystems.

Introduction

Human activities affect nearly all habitats through changes in climate and land-use, which in turn alter vegetation cover and composition. These changes negatively impact many species that have narrow environmental tolerances and limited dispersal capacity across anthropogenic landscapes (Schulte to Bühne et al. 2020). Anthropogenic impacts most strongly affect endemic species – i.e., taxa with small distribution ranges and narrow ecological niches (Brook et al. 2008). Diversity of endemic plants and animals is higher in areas characterized by historical stability, high precipitation, environmental heterogeneity, and insularity. Unfortunately, these areas usually coincide with major human degradations of the environment (Kier et al. 2009; Stein et al. 2014; Sandel et al. 2020).

Unlike the situation with plants and animals, global patterns of fungal diversity, endemism and vulnerability to environmental change remain virtually unknown (Cameron et al. 2019; Guerra et al. 2021b; but see Talbot et al. 2014; Davison et al. 2015). This is alarming, given the fundamental roles that fungi play in carbon and nutrient cycling processes (Wardle & Lindahl 2014; Crowther et al. 2019). Comparative studies have indicated that aboveground and belowground biodiversity are driven by different environmental predictors at local and global scales (Cameron et al. 2019; Le Provost et al. 2021). This suggests differential responses of macro- and microorganisms to land use and climate changes (Guerra et al. 2021b). As for plants and animals, soil fungal communities are likely vulnerable to global change drivers. For instance, high-temperature stress (Malcolm et al. 2008; Barcenas-Moreno et al. 2009; Morgado et al. 2015; Misiak et al. 2021) and prolonged drought (Schmidt et al. 2017; de Vries et al. 2018) can alter fungal growth, functionality and community composition. Likewise, changes in land use that result in habitat fragmentation may lead to shifts in prevalence of pathogenic, mutualistic, and free-living fungal groups (Brinkmann et al. 2019; Makiola et al. 2019; Le Provost et al. 2021; Rodriguez-Ramos et al. 2021).

While thousands of plant and animal species are listed as threatened on the IUCN global Red List, only 262 out of an estimated 2.2-3.8 million fungal species (Hawksworth & Lücking 2017) have been listed as such. The majority of these are from high-income countries in temperate regions (IUCN 2021) and are from fungal groups that make conspicuous macroscopic fruiting bodies (Cui et al. 2021). However, the vast majority of fungi produce no or inconspicuous fruiting bodies and are therefore hard to survey, which has hampered their conservation assessment (Gonçalves et al. 2021).

Here we used the most advanced high-resolution sequencing technology to globally survey soil fungal diversity and assess their endemism and vulnerability to global change. We hypothesized that i) the endemism of fungi is relatively higher in the tropics due to greater regional climatic stability; and ii) vulnerability of fungi to global change is highest in habitats experiencing the strongest global warming effects (polar regions) and intensive land use (dry tropics). We predicted that because of their intimate associations with other organisms, endemism and vulnerability patterns are more evident for macrofungi and biotrophic groups compared with saprotrophic microfungal groups. We then propose global conservation priorities for these ecologically pivotal fungi.

Results and Discussion

Fungal diversity

We used the recently generated Global Soil Mycobiome consortium dataset (GSMc; 3,200 plots, Tedersoo et al. 2021b) along with data from five other global soil surveys (**Fig. 1**; see methods) and international nucleotide sequence databases to determine the diversity and endemism of fungal

functional groups – viz. arbuscular mycorrhizal (AM) fungi, ectomycorrhizal (EcM) fungi, non-EcM Agaricomycetes (mostly saprotrophic macrofungi), molds, pathogens, opportunistic human parasites (OHPs, mostly thermophilic saprotrophs), early-diverging unicellular lineages (mostly chytrids, aphelids, and rozellids), and yeasts. Compared to previous meta-analytical approaches (e.g. Vetrovsky et al. 2019), our cumulative data comprise the largest available globally standardized database based on directly comparable soil sampling and long-read molecular analysis protocols. Collectively, all datasets yielded 20,182,427 fungal reads composed of 905,841 ‘species’ – operational taxonomic units (OTUs), each defined as <98% sequence similarity of the rRNA ITS barcode from all other OTUs. The genera *Tomentella* (Basidiomycota), *Penicillium* (Ascomycota), and *Mortierella* (Mortierellomycota) were the most species-rich (**Fig. 1**).

We combined machine-learning and general linear modeling (GLM) approaches to find the best predictors of fungal species richness and the Shannon index of diversity for settling the contrasting results obtained from previous global studies (Tedersoo et al. 2014; Egidi et al. 2019; Vetrovsky et al. 2019). At the site scale (α -diversity), the best supported results were obtained for residuals of logarithm-transformed richness accounting for sequencing depth (**Fig. 2**). Since the datasets were retrieved using different sampling design and therefore differed strongly in the inferred richness (**Fig. 3**), we focused mainly on analyses of the largest, GSMc dataset. Total fungal richness had a broadly unimodal relationship with soil pH ($R^2_{\text{adj}}=0.133$) and responded positively to vegetation age ($R^2_{\text{adj}}=0.045$; **Fig. 4**). Deserts and Antarctic habitats supported the lowest richness among all biomes (**Fig. 4**). We validated the results using several datasets, in which fungal richness had a unimodal relationship with soil pH and positive response to mean annual precipitation (MAP)(**Fig. 5**). Across datasets, fungal γ -diversity at the ecoregion level was best explained by average MAP ($R^2_{\text{adj}}=0.179$; **Fig. 6**). The differences in richness trends between α -diversity and γ -diversity indicate that high precipitation favors niche differentiation at the regional scale, as reflected by higher turnover between sites (i.e. increasing α - to γ -diversity).

Our results thus update previous patterns of α -diversity decrease (Tedersoo et al. 2014) or increase (Vetrovsky et al. 2019) at high latitudes and confirm relatively lower fungal diversity in Antarctica. The latter pattern has been/can be ascribed to low plant diversity and coverage (Newsham et al. 2016). The more prominent latitudinal gradient in γ -diversity reflects a greater positive effect of MAP on the regional fungal species pool. Disregarding Antarctica, the lack of a global α -diversity latitudinal gradient in fungi is unique among terrestrial organisms (Kinlock et al. 2018). By comparison, the γ -diversity patterns detected resemble those found for soil fauna (Aslani et al. 2022; Potapov et al. 2022) and protistan parasites (Oliverio et al. 2020), all which show slight richness peaks in tropical latitudes. The distinctly weaker latitudinal diversity gradients of soil organisms compared with most aquatic and terrestrial macro-organisms may be related to indirect effects of temperature-related climatic variables as well as soil pH and C/N ratio as main drivers of soil habitat quality. The

differences may also be related to higher dispersal capacity of soil organisms who have microscopic body sizes or dispersal propagules (Soininen et al. 2013; Aslani et al. 2022).

Fungal endemism

To estimate relative endemism among the world's ecoregions (**Fig. 7; Table 1; see methods**), we combined indices of community similarity, uniqueness, and species ranges into an overall endemism index (see Methods). Five metrics were combined, including the number and proportion of endemic species, mean maximum geographical range of species, Jaccard index, and beta-sim index (**Box 1**). We found that endemism of all fungi peaked in moist tropical biomes and it was positively related to mean annual air temperature (MAT; $R^2_{adj}=0.277$; **Fig. 8**) and soil acidity ($R^2_{adj}=0.108$; **Table 2**).

While endemism patterns of non-EcM Agaricomycetes and AM fungi were similar to those shown for all fungi, different patterns were found for other functional groups. Endemism of EcM fungi was related to high mean annual precipitation (MAP) ($R^2_{adj}=0.147$). Molds, pathogens and yeasts showed multiple endemism hotspots. Molds ($R^2_{adj}=0.199$) and pathogens ($R^2_{adj}=0.105$) had relatively greater endemism in strongly acidic or alkaline soils, indicating that extreme soil conditions may support unique soil biota, with limited effective dispersal across edaphically extreme habitats. Human footprint (see Methods) had a weak negative effect on endemism of all fungi ($R^2_{adj}=0.018$), pathogens ($R^2_{adj}=0.015$), and OHPs ($R^2_{adj}=0.056$), suggesting that anthropogenic habitat loss or homogenization may affect endemic species (Finderup Nielsen et al. 2019). European ecoregions had the lowest endemism for all fungi ($R^2_{adj}=0.065$), pathogens ($R^2_{adj}=0.086$) and unicellular fungi ($R^2_{adj}=0.035$) compared with those of other areas. Averaged current aerial bioclimatic variables better explained endemism compared with the ranges of those variables or bioclimatic variables of soil and last glacial maximum (LGM). Climate change since the LGM had a weak positive effect on endemism of molds (mean diurnal range and overall climate change: $R^2_{adj}=0.073$) and OHPs (isothermality and mean diurnal range: $R^2_{adj}=0.056$) but not other groups.

We found that patterns in fungal endemism were relatively consistent among the five individual endemism indices and that they resemble endemism patterns of vascular plants and animals, which exhibit major hotspots in wet tropical habitats (Kier et al. 2009; Barlow et al. 2018). However, endemism patterns in fungi were somewhat weaker, which may reflect the greater long-distance dispersal capacity of fungal spores relative to propagules of plants and animals (Golan & Pringle 2017). In terms of the greater macroorganism richness and endemism found in the tropics, the literature abounds with hypotheses, including narrower niche breadth, more asymmetric interactions (i.e., greater specialization), climatic stability, and more rapid evolution due to environmental energy (Vazquez & Stevens 2004; Brown 2013) in the tropics than elsewhere. Negligible effects of the LGM suggest that

climatic stability is not an important driver of fungal endemism, a pattern that contrasts with those of plants and animals (Rosauer & Jetz 2015). The greater phylogenetic diversity of fungi noted for the tropics (e.g. Tedersoo et al. 2018) may in part reflect tropical origins for many lineages, as well as the radiation and rapid speciation of a limited number of EcM fungal genera into higher latitude areas (Kennedy et al. 2012; Sanchez-Ramirez et al. 2015). On a global scale, plant diversity does not appear to be causally related to fungal diversity (Tedersoo et al. 2014), but there is some evidence for stronger mutualistic plant-fungal interactions related to high rainfall (Pöhlme et al. 2018). Pathogenic interactions warrant further research in this respect, given their major importance as regulators of plant diversity (Chen et al. 2019). Tropical soil fungi have relatively greater dispersal limitations (Bahram et al. 2013) and narrower distribution ranges (Tedersoo et al. 2014), suggesting that high local diversity may contribute to greater regional-scale endemism.

Vulnerability of fungi to global change drivers

Communities with many species at their environmental niche limits may be particularly vulnerable to local extinctions (Watson et al. 2013; Smith et al. 2020b). Thus, we evaluated the relative vulnerability of soil fungal functional groups by estimating the percentage of species occurring at their upper niche limits to three major global change drivers – land use (land cover change), heat (maximum monthly temperature), and drought (lowest quarterly precipitation). We projected to the year 2070 relative to the 2015 baseline, using the average vulnerability index (Smith et al. 2020b), land use extrapolations of the LUH2 global dataset (Hurtt et al. 2020), and climatic extrapolations based on the CCS8.5 scenario (Karger et al. 2021). For all fungi taken together, predicted vulnerability to heat (best predictor: maximum monthly temperature; $R^2_{adj}=0.583$) and drought (precipitation seasonality; $R^2_{adj}=0.456$) were the greatest in the tropical and subtropical latitudes. Vulnerability to land use change (isothermality; $R^2_{adj}=0.145$) peaked in the tropics. The overall additive global change vulnerability was thus the highest in densely populated tropical and subtropical regions. Fungal functional groups had similar vulnerability patterns, which were mostly related to temperature. Among fungal groups, average vulnerability scores were highest for AM and EcM symbionts and unicellular fungi, but these scores differed only slightly across the global change drivers (**Fig. 9**). The actual vulnerability is probably underestimated for biotrophic pathogens and EcM fungi, because these groups associate with a limited number of plant species and are sometimes host-specific (Kennedy et al. 2015). Therefore, the loss of one of the few key symbiotic partners may greatly reduce the biotic niche of specialist fungi.

Patterns of vulnerability in fungi are somewhat similar to those of terrestrial plants and animals, where vulnerability peaks in drylands prone to desertification (Warren et al. 2013), arctic/alpine areas (cold-adapted species), and regions with dense human populations (Watson et al. 2013). The relatively low vulnerability to heat in tundra-inhabiting fungi can be explained by their relatively high

temperature optima (Maynard et al. 2019; but see Misiak et al. 2021), acclimation (Romero-Olivares et al. 2017), and poleward migration potential, despite relatively greater predicted warming in Arctic ecosystems. Above certain tolerance thresholds, soil organisms may be physiologically constrained by increasing soil temperature and evaporation, lower soil water potentials, and loss of oxygen due to greater respiration and faster decomposition, which result in hampered soil functioning and ecosystem multifunctionality (Delgado-Baquerizo et al. 2017). Open areas are predicted to increase due to climate change and human activities. This will further expose soil to solar radiation and result in the loss of fungal plant hosts. While here we calculated average vulnerabilities by adding up the effects of individual drivers, global change impacts tend to be synergistic (Rillig et al. 2019), so actual vulnerabilities may be much higher.

Implications for conservation

Most fungi and soil organisms do not enjoy the protection and conservation measures that are afforded to more “charismatic” animals and plants (Ducarme et al. 2013). Nonetheless, fungi and other soil biota are pivotal to soil health, nutrient cycling, water storage, food security, and many other ecosystem services. Their biodiversity should hence be brought to the center stage of global sustainability thinking and conservation planning. For example, these organisms should be factored in when selecting protected areas otherwise based on plant and animal conservation (Guerra et al. 2021a). The fact that many EcM and plant pathogenic fungal species are associated with specific host plants indicates that on the local scale it is not only the narrowly-distributed species but also unique biotic associations that require focused conservation measures. From the fungal perspective, it is particularly important to protect plant species that act as hubs in modules of biotic interaction networks, because these hub species typically associate with multiple, distinct fungal partners (Pölme et al. 2018). In other cases, certain unique plant species or higher taxonomic groups should be prioritized. For example, in southern South America, the drought-sensitive tree family Nothofagaceae is the only group known to support EcM fungi that are endemic to this area (Godoy & Marin 2019)

Although the vulnerability to environmental change differed among fungal groups, their overall global patterns were similar. This suggests that broad habitat conservation measures may work for most fungal groups, including macroscopic non-EcM Agaricomycetes and EcM fungi as well as more cryptic pathogens and other groups. To accomplish this, fungi need to be incorporated into conservation frameworks (Gonçalves et al. 2021). Actions to fill existing information gaps at the local and global levels must also be taken, and global-scale surveys should take into account the soil biodiversity assessments, complementing the traditional collections-based assessment with metabarcoding of environmental DNA. This applies to national conservation evaluation programs and engagement in global policy-making initiatives, such as the System of Environmental Economic

Accounting of the United Nations, World Biodiversity Forum, and Post-2020 Global Biodiversity Framework. Furthermore, promoting the red-listing of endangered fungal species at the national and global levels is critical (FAO 2020; IUCN 2021). Fungi need active and specific inclusion in national and global conservation policies and strategies, not just passive and implicit protection.

Our study provides evidence that soil fungi may be highly vulnerable to global change, which needs to be considered when planning how to preserve these key organisms in a changing world. As with plants and animals, fungi appear to be environmentally sensitive due to the strong impacts that land cover change, low moisture, and high temperatures have on taxonomic and functional composition (Brinkmann et al. 2019; Makiola et al. 2019; this study). The endemism of fungi is highest in tropical forest biomes (Kier et al. 2009; this study), so conservation measures advocated for tropical plants and animals (Brooks et al. 2006; Barlow et al. 2018) are likely to conserve fungi. Tropical forests are under continued threat from deforestation and degradation driven by expanding agriculture, extractive industries, and infrastructural projects (Bebbington et al. 2018). Conservation of herbaceous wetlands, tropical rainforests and tropical woodlands is supported by our global fungal conservation priority map that accounts for endemism, vulnerability, and γ -diversity (**Fig. 10**). Additionally, given the importance of soil pH for soil microbial diversity and composition, it is essential to prioritize areas with high pedodiversity or mixed landscapes including bogs, various forest types, and grasslands. As a crucial measure, desertification and loss of soil organic matter needs to be controlled by reducing the conversion of primary forest to crops and pasture (Smith et al. 2020a). This is important not only to prevent land degradation processes from impairing bacterial and fungal diversity, but also to sustain the capacity of drylands to provide essential functions and services, such as soil fertility, carbon storage, and food production for more than one billion people (Sivakumar 2007; Delgado-Baquerizo et al. 2018).

Conclusions

In conclusion, soil fungi show strong endemism patterns, which differ by functional groups and are driven by both climatic and edaphic factors. Fungal groups also differ strongly in their relative vulnerability scores to global change, which peak in heavily populated tropical dryland areas. Unfortunately, these are the very areas most prone to further land degradation and desertification. Fungal endemism and vulnerability patterns only partly mirror those of vascular plants and animals, which may be ascribed to their more efficient dispersal mechanisms. Global conservation efforts should include fungal biodiversity surveys alongside assessments of soil health, below-aboveground feedbacks, and areas of highest conservation priority, to secure the protection of habitats. Even more, they should include the monitoring of regional fungal communities over time, to pick relevant

changes and to provide early warning signals of impending change. What we do not know, we cannot efficiently manage or protect.

Methods

Datasets

To study fungal endemism and vulnerability to global change, we combined data from the Global Soil Mycobiome consortium (GSMc) open dataset (Teder et al. 2021b) with materials from five other global soil biological surveys (**Fig. 1**) – BIODESERT (Maestre et al. 2022), MUSGONET (including the natural sites in Delgado-Baquerizo et al. 2021), CLIMIFUN (Bastida et al. 2021), GlobalAM (Davison et al. 2021), GlobalWetlands (Bahram et al. 2022) as well as Sanger sequence data from soil-inhabiting fungi obtained from the UNITE database (Nilsson et al. 2019) covering GenBank. We obtained the DNA from all five surveys and performed new DNA metabarcoding analyses following the protocols outlined for the GSMc dataset (Teder et al. 2021b).

All datasets comprised information on geographical coordinates and soil pH. Based on geographical coordinates, we assigned the following climatic and land cover metadata to the samples: i) CHELSA v2.1 bioclimatic variables for the period 1981–2010 (Karger et al., 2020), ii) CHELSA-TraCE21k v1.0. for the LGM (Karger et al. 2021), and iii) CHELSA v2.1 climate extrapolations for the year 2070 following the RCP8.5 global warming scenario with SSP5 socioeconomic conditions and the GFDL-ESM4 global circulation model (Karger et al., 2020); iv) normalized difference vegetation index (NDVI; Filipponi et al. 2018); v) SoilGrids v.2 soil pH from 0–5 cm depth (Poggio et al., 2021); vi) land cover type using Copernicus classification v.3 (Buchhorn et al. 2020) for the year 2015; and vii) human footprint index based on the Land-Use Harmonization (LUH2; Hurtt et al., 2020) or the year 2015 and 2070 extrapolation. Based on original descriptions of vegetation (age, cover, relative abundance of species, fire history) or remote sensing data (Google Maps), samples were assigned to biomes (Olson et al. 2001) and land cover types. Based on Z-transformed differences in present and LGM bioclimatic variables, we calculated for each sample an averaged LGM climate change index. Further, for each sample we estimated the human footprint index as the cumulative sum of land-use state transitions, with the year 1960 used as a baseline.

Bioinformatics

To infer fungal species and taxonomy, we used a long-read sequencing approach involving the ribosomal RNA 18S gene V9 subregion, ITS1 spacer, 5.8S gene, and ITS2 spacer to enhance

taxonomic resolution and accuracy. We used degenerate, universal eukaryotic primers to cover as many divergent taxa within the fungi and micro-eukaryotes as possible (Tedersoo et al. 2021a). The amplicon samples were prepared in 82 PacBio SMRTbell sequencing libraries and sequenced on 48 PacBio Sequel 8M SMRT cells. The obtained reads were quality-filtered, demultiplexed to samples, trimmed to include only the full-length ITS region, and clustered to operational taxonomic units (conditionally termed as species) at 98% sequence similarity, which roughly corresponds to species-level divergence. Taxonomy was assigned based on information from the 10 best BLASTn matches against the UNITE 9.1 beta dataset (<https://doi.org/10.15156/BIO/1444285>). The resulting species-by-sample matrices were manually checked library-wise for external and cross-contamination and rates of index switching artifacts. We excluded several samples for which we suspected contamination, and removed rare occurrences of dominant species using the following thresholds: abundances = 1 for species with total abundance of >99 and abundances = 2 for species with total abundance of >999.

Based on FungalTraits 1.3 (Pöhlme et al. 2020), species belonging to the kingdom Fungi were assigned to functional groups based on ecological or physiological characters: i) AM fungi (including all Glomeromycota but excluding all Endogonomycetes, because there is not enough information to distinguish AM species from free-living species); ii) EcM fungi (excluding dubious lineages); iii) non-EcM Agaricomycetes (mostly saprotrophic fungi with macroscopic fruiting bodies); iv) molds (including Mortierellales, Mucorales, Umbelopsidales, and Aspergillaceae and Trichocomaceae of Eurotiales and Trichoderma of Hypocreales); v) putative pathogens (including plant, animal and fungal pathogens as primary or secondary lifestyles); vi) OHPs (excluding Mortierellales); vii) yeasts (excluding dimorphic yeasts); and viii) other unicellular (non-yeast) fungi (including chytrids, aphids, rozellids, and other early-diverging fungal lineages). Other groups such as lichen-forming fungi were not considered, owing to their relative infrequency in soil across samples and ecoregions. Among these groups, mostly non-EcM Agaricomycetes and EcM fungi comprise many red-listed species of conspicuous conch-shaped, resupinate, or stipitate fruiting bodies and are hence considered to be of relatively higher conservation interest (Cao et al. 2021; IUCN 2021).

Fungal diversity

To assess patterns in global fungal α -diversity, we first calculated the residuals of logarithmically-transformed fungal richness and richness of major functional groups by performing linear regression against the logarithm of sequencing depth. We also compared other approaches such as residuals from untransformed richness against square-root-transformed and log-transformed sequencing depth (Tedersoo et al. 2014), exclusion of singletons, Shannon index of diversity, traditional rarefaction (depth, 500 reads), and SRS-rarefaction (Beule & Karlovsky 2020) to 500 (minimum) or 3894 (median) reads (**Fig. 2**). Because the approach including singletons and log-log transformation for

selecting residuals resulted in best-supported models (**Fig. 2**), we chose this approach for further analyses. Because the sampling protocols differed in sampling area size, number of subsamples and DNA extraction protocols, we included only the GSMc dataset in analyses of fungal richness and composition. The GSMc dataset furthermore featured information about vegetation (age, proportion of dominant plant taxa and mycorrhiza types, fire history) soil properties (C, N, P, K, Ca, Mg concentration), and sampling date. Based on geographical coordinates and sampling dates, we calculated geographic and temporal eigenvectors using the *adespatial* package of R (Dray et al. 2018). Phylogenetic eigenvectors were calculated for woody plant species composition by mapping the taxa to a multigene vascular plant phylogram (Qian & Jin 2016).

Because of metadata availability and comparability, we exclusively used the GSMc dataset (3200 composite samples by 722,682 species) to estimate the best predictors of fungal α -diversity and composition and to update global fungal diversity distribution maps. We first calculated the residuals of logarithmically-transformed fungal richness and richness of major functional groups by performing linear regression against the logarithm of sequencing depth. Besides richness residuals and Shannon index of diversity, we calculated fungal phylogenetic diversity (PD), mean phylogenetic distance (MPD), and mean neighbor taxonomic distance (MNTD) indices based on a classification tree (Tedersoo et al. 2018) using the *PhyloMeasures* package of R (Tsirogianis & Sandel 2016). For predictors, we included biome and continent (dummy variables), bioclimatic, edaphic, and vegetation-related variables, as well as eigenvectors of woody plant phylogenetic composition, spatial and temporal distance. Using random forest, we retrieved 20 candidate variables for GLM modeling. In GLM modeling, we included quadratic terms to account for non-linearity. To avoid an excessive number of predictors, only significant variables ($P < 0.001$; $R^2 > 0.020$) were kept in the final models. We also performed additional correlation analyses to illustrate the latitudinal gradient of α -diversity and γ -diversity. For generating fungal diversity maps, we performed additional analyses using bioclimatic variables, database $pH_{H2O(0-5\text{ cm})}$, human footprint index and Copernicus land use categories and their interactions with continuous predictors as described for vulnerability maps. The GSMc dataset-based α -diversity analyses were validated by performing similar analyses using other datasets. Richness residuals were calculated separately for these data and all datasets were subjected to model selection for soil pH and bioclimatic variables, because information about other predictors was inconsistent. We also modeled various versions of soil pH including the original measurements of pH_{KCl} as well as pH_{H2O} extrapolations for soil depths of 0-5 cm, 0-30 cm, and 15-30 cm. Since the pH_{KCl} fit significantly better into global models (**Fig. 2B**) and pH_{H2O} extrapolations were missing for smaller islands and Antarctic habitats, we used pH_{KCl} in subsequent analyses.

For γ -diversity, logarithmically-transformed cumulative ecoregion (see below) species richness was subjected to model selection against the logarithmically-transformed number of samples and sequencing depth to calculate residual richness. Residual richness and endemism indices of all fungi

and functional groups were subjected to random forest machine learning analysis to pre-select ten most important variables for GLM.

Endemicity

To infer endemicity patterns in fungi, samples (including data obtained from UNITE, covering International Nucleotide Sequence Databases entries) were assigned to ecological regions (Olson et al. 2001) based on their geographical coordinates, allowing a 10-km of buffer zone between a terrestrial ecological region and water (due to low resolution of the map layers in shore areas). Based on climatic and floristic similarities, the ecological regions were further aggregated into larger areas or split into smaller, geographically distinct units, which we refer to as ecoregions (**Fig. 7**). Each of 174 ecoregions comprised 1 to 45 soil samples, with surplus samples excluded randomly. Five indices of endemism – viz., the number of endemic species (weight = 16.7%), proportion of endemic species (weight = 16.7%), mean maximum geographical range of taxa (weight = 33.3%), Jaccard index (weight = 16.7%), and beta-sim index (weight = 16.7%) – were selected for calculating the averaged endemicity index based on the community matrix (Crisp et al. 2001; Villéger & Brosse 2012; **Box 1**). The former two and latter two indices reflect similar aspects of endemicity and were therefore downweighted. To account for differences in sampling intensity, we calculated residuals for the numbers of all species and endemic species by regressing these against the logarithmically-transformed number of samples and sequencing depth.

Of the indices used, only the number and proportion of endemic taxa were significantly positively correlated with species richness (all fungi: $r=0.707$ and $r=0.212$, respectively), whereas others had no significant correlation. Furthermore, species richness was not included among the best predictors of averaged endemicity, indicating that these metrics are independent. Endemicity indices were calculated using the betapart package v.1.5.4 (Baselga & Orme 2012) of R v.4.1.10 (R Core Team 2022). Endemicity indices of all fungi and functional groups were subjected to random forest machine learning analysis to pre-select the ten most important variables for GLM. We used the variance (as coefficient of variation) and averaged values of bioclimatic variables, area, latitude, longitude, altitude, and soil pH as well as continents (dummy variables) to explain endemicity. GLMs were fitted using second-order polynomial terms for continuous variables. Only significant variables ($P<0.050$; $r^2>0.020$) were kept in the final models. Based on the predictions revealed by GLMs, endemicity maps were constructed using the sf v.1.0-5 (Pebesma 2018) package of R.

488 *Global change vulnerability*

489 The vulnerability of soil fungal groups was estimated relative to three global change drivers – heat
490 (maximum monthly temperature), drought (negative of inverse hyperbolic sine-transformed
491 precipitation in the driest quarter), and land cover change – for the year 2070 (relative to 2015
492 baseline) using the community-mean percentile vulnerability index (V_2 ; Smith et al. 2020b). This
493 index is based on averaging percentiles of all species at a given global change driver value.

$$v_{2i} = \left(\frac{\sum_{j=1}^n a_{ij} F_j(x_i)}{\sum_{j=1}^n a_{ij}} \right) \times 100$$

494

495 where a_{ij} is the presence (0 or 1) of species j in site i , $F_j(x_i)$ is the percentile of species j given site
496 parameter value x_i , x_i is the parameter value of site i , and n is the total number of species observed.

497 Precipitation in the driest quarter was selected as a proxy for drought, because bioclimatic variables
498 cover larger areas (including islands) and offer greater resolution compared with other measures of
499 soil water content and indicators of drought. The vulnerability scores were calculated for each soil
500 sample using `vuln v.0.0.05` (Smith et al. 2020b) package of R. We also constructed the average
501 vulnerability score by equally weighting all components. The vulnerability scores were unrelated to
502 sequencing depth and sample size. We performed a similar random forest and GLM modeling
503 exercise for determining the main predictors of vulnerability as described above, but allowed
504 interaction terms between categorical and continuous predictors and used a more relaxed threshold for
505 keeping variables in the model ($P < 0.001$; $R^2 > 0.01$) due to greater sample size. To construct
506 vulnerability maps we used a regression-kriging approach (Hengl & MacMillan 2019). To predict
507 vulnerability scores for each global driver and estimate their prediction uncertainty, thin plate splines
508 (basis dimensionality = 3) were fitted using a generalized additive model (GAM) with `mgcv v.1.8-38`
509 (Wood 2011) package. To incorporate the spatial autocorrelation signal, we calculated residuals at the
510 sampling sites and used inverse distance weighting (IDW) to interpolate residuals beyond the
511 sampling sites. To obtain final vulnerability predictions, interpolated residuals were added to the
512 results based on the predicted regression part. By using the relative vulnerability values, we also
513 prepared the map of fungal vulnerability ascribed to each of the three components. Vulnerability maps
514 were visualized using the `raster v.3.5-9` (Hijmans 2021) package of R.

515 The maps for conservation priorities were calculated for all fungi using sampling points used in
516 vulnerability analyses, except points corresponding to cropland and urban and village land cover. For
517 each sampling point, the respective average endemism, γ -diversity, and vulnerability scores were z-
518 transformed, followed by adding a constant (5, to exclude negative values), multiplied (to downweigh

areas with any low values), and used in a regression-kriging approach (**Table 3**) as described for vulnerability.

Based on methods comparison, we conclude that inclusion of singletons may increase richness model performance at relatively low sequencing depth in high-quality, third-generation sequencing datasets. Furthermore, residuals of log-transformed richness against log-transformed sequencing depth yields significantly better model estimates compared with most other standardization methods. Richness data from studies with different sampling designs must not be pooled in a common analysis unless the factor “study” or sampling attributes (e.g. number of subsamples, volume of samples) are accounted for. Use of original soil pH data strongly outperforms extrapolated data; therefore, soil pH extrapolations should not be used for testing relative effects of edaphic and climatic and other properties. This probably applies to other edaphic and vegetation-related values that greatly vary on a local scale.

Acknowledgements

The bulk of the funding is derived from the Estonian Science Foundation (grants PRG632, PRG1170, PSG136, MOBTP198), Norway-Baltic financial mechanism (grant EMP442) and Novo Nordisk Fonden (NNF20OC0059948). LT, VM, AZ, MB, NAS, AA, UK, and KA designed the study; VM, LT, AZ, MB, NH-D, SA, and OP analyzed data; LT, MD-B, FTM, JP, MÖ, MM, MZ, ME, and ÜM contributed DNA extracts from global surveys; other authors contributed materials, data, and/or chemical analyses; LT wrote the first draft and all authors contributed to the writing of the article. The authors confirm that there are no competing interests. Data for this paper have been deposited to the PlutoF data repository (GSMc data: <https://doi.org/10.15156/BIO/2263453>). Representative sequences of identical reads per sample are available from the UNITE database. The scripts used for the bioinformatic analysis are available at GitHub: <https://github.com/Mycology-Microbiology-Center/GSMc>

References

- Aslani F, Geisen S, Ning D, Tedersoo L, Bahram M. 2021. Towards revealing the global diversity and community assembly of soil eukaryotes. *Ecology Letters* 25: 65-76.
- Bahram M, Kõljalg U, Courty P-E, Diedhiou A, Kjoller R, Põlme S, Ryberg M, Veldre V, Tedersoo L. 2013. The distance-decay of similarity in communities of ectomycorrhizal fungi in different ecosystems and scales. *Journal of Ecology* 101: 1335–1344.
- Bahram M, Espenberg M, Pärn J, Lehtovirta-Morley L, Anslan S, Kasak K, Kõljalg U, Liira J, Maddison M, Moora M, Niinemets Ü, Öpik M, Pärtel M, Soosaar K, Zobel M, Hildebrand F, Tedersoo L, Mander Ü. 2022. Structure and function of the soil microbiome underlying N₂O emissions from global wetlands. *Nature Communications* 13:1430.
- Barcenas-Moreno GE, Gomez-Brandon MA, Rousk J, Bååth E. 2009. Adaptation of soil microbial communities to temperature: comparison of fungi and bacteria in a laboratory experiment. *Global Change Biology* 15: 2950-2957.
- Barlow J, França F, Gardner TA, Hicks CC, Lennox GD, Berenguer E, Castello L, Economo EP, Ferreira J, Guénard B, Leal CG, Isaac V, Lees A, Parr C, Wilson S, Young P, Graham N. 2018. The future of hyperdiverse tropical ecosystems. *Nature* 559: 517-526.
- Baselga A, Orme CDL. 2012. betapart: an R package for the study of beta diversity. *Methods in Ecology and Evolution* 3: 808-812.

- Bastida F, Eldridge DJ, García C, Png GK, Bardgett RD, Delgado-Baquerizo M. 2021. Soil microbial diversity–biomass relationships are driven by soil carbon content across global biomes. *The ISME Journal* 15: 2081–2091.
- Bebbington AJ, Bebbington DH, Sauls LA, Rogan J, Agrawal S, Gamboa C, Imhof A, Johnson K, Rosa H, Royo A, Toumbourou T. 2018. Resource extraction and infrastructure threaten forest cover and community rights. *Proceedings of the National Academy of Sciences* 115: 13164-13173.
- Beule L, Karlovsky P. 2020. Improved normalization of species count data in ecology by scaling with ranked subsampling (SRS): application to microbial communities. *PeerJ* 8:e9593.
- Brinkmann N, Schneider D, Sahner J, Ballauff J, Edy N, Barus H, Irawan B, Budi SW, Qaim M, Daniel R, Polle A. 2019. Intensive tropical land use massively shifts soil fungal communities. *Scientific Reports* 9:3403.
- Brook BW, Sodhi NS, Bradshaw CJ. 2008. Synergies among extinction drivers under global change. *Trends in Ecology & Evolution* 23: 453-460.
- Brooks TM, Mittermeier RA, Da Fonseca GA, Gerlach J, Hoffmann M, Lamoreux JF, Mittermeier CG, Pilgrim JD, Rodrigues AS. 2006. Global biodiversity conservation priorities. *Science* 313: 58-61.
- Brown JH. 2013. Why are there so many species in the tropics? *Journal of Biogeography* 41: 8-22.
- Buchhorn M, Lesiv M, Tsendbazar NE, Herold M, Bertels L, Smets B. 2020. Copernicus global land cover layers - collection 2. *Remote Sensing* 12:1044.
- Cameron EK, Martins IS, Lavelle P, Mathieu J, Tedersoo L, Bahram M, Gottschall F, Guerra CA, Hines J, Patoine G, Siebert J. 2019. Global mismatches in aboveground and belowground biodiversity. *Conservation Biology* 33: 1187-1192.
- Cao Y, Wu G, Yu D. 2021. Include macrofungi in biodiversity targets. *Science* 372: 1160-1160.
- Chen L, Swenson NG, Ji N, Mi X, Ren H, Guo L, Ma K. 2019. Differential soil fungus accumulation and density dependence of trees in a subtropical forest. *Science* 366: 124-128.
- Crisp MD, Laffan S, Linder HP, Monro A. 2001. Endemism in the Australian flora. *Journal of Biogeography* 28: 183-198.
- Crowther TW, Van den Hoogen J, Wan J, Mayes MA, Keiser AD, Mo L, Averill C, Maynard DS. 2019. The global soil community and its influence on biogeochemistry. *Science* 365:eaav0550.
- Davison J, Moora M, Semchenko M, Adenan SB, Ahmed T, Akhmetzhanova AA, Alatalo JM, Al-Quraishy S, Andriyanova E, Anslan S. 2021. Temperature and pH define the realised niche space of arbuscular mycorrhizal fungi. *New Phytologist* 231: 763-776.
- Davison J, Moora M, Öpik M, Adholeya A, Ainsaar L, Ba A, Burla S, Diedhiou AG, Hiiesalu I, Jairus T, Johnson NC, Kane A, Koorem K, Kochar M, Ndiaye C, Pärtel M, Reier Ü, Saks Ü, Singh R, Vasar M, Zobel M. 2015. Global assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism. *Science* 349: 970-973.
- de Vries FT, Griffiths RI, Bailey M, Craig H, Girlanda M, Gweon HS, Hallin S, Kaisermann A, Keith AM, Kretschmar M, Lemanceau P, Bardgett RD. 2018. Soil bacterial networks are less stable under drought than fungal networks. *Nature Communications* 9:3033.
- Delgado-Baquerizo M, Eldridge DJ, Liu YR, Sokoya B, Wang JT, Hu HW, He JZ, Bastida F, Moreno JL, Bamigboye AR, Blanco-Pastor JL, Singh BK. 2021. Global homogenization of the structure and function in the soil microbiome of urban greenspaces. *Science Advances* 7:eabg5809.
- Delgado-Baquerizo M, Eldridge DJ, Ochoa V, Gozalo B, Singh BK, Maestre FT. 2017. Soil microbial communities drive the resistance of ecosystem multifunctionality to global change in drylands across the globe. *Ecology Letters* 20: 1295-1305.
- Delgado-Baquerizo M, Eldridge DJ, Travers SK, Val J, Oliver I, Bissett A. 2018. Effects of climate legacies on above-and below-ground community assembly. *Global Change Biology* 24: 4330–4339.
- Dray S, Blanchet G, Borcard D, Guenard G, Jombart T, Larocque G, Legendre P, Madi N, Wagner HH, Dray MS. 2018. Package ‘adespatial’.
<https://cran.microsoft.com/web/packages/adespatial/adespatial.pdf>
- Ducarme F, Luque GM, Courchamp F. 2013. What are “charismatic species” for conservation biologists. *BioSciences Master Reviews* 10: 1-8.
- Egidi E, Delgado-Baquerizo M, Plett JM, Wang J, Eldridge DJ, Bardgett RD, Maestre FT, Singh BK. 2019. A few Ascomycota taxa dominate soil fungal communities worldwide. *Nature Communications* 10:2369.

FAO. 2020. State of knowledge of soil biodiversity: status, challenges and potentialities. FAO, Rome. <https://doi.org/10.4060/cb1928en>

Filipponi F, Valentini E, Nguyen Xuan A, Guerra CA, Wolf F, Andrzejak M, Taramelli A. 2018. Global MODIS fraction of green vegetation cover for monitoring abrupt and gradual vegetation changes. *Remote Sensing* 10:653.

Finderup Nielsen T, Sand-Jensen K, Dornelas M, Bruun HH. 2019. More is less: net gain in species richness, but biotic homogenization over 140 years. *Ecology Letters* 22: 1650-1657.

Godoy R, Marin C. 2019. Mycorrhizal studies in temperate rainforests of Southern Chile. In: Pagano MC, Lugo MA (eds). *Mycorrhizal Fungi in South America*. (pp. 315-341). Springer, Cham.

Golan JJ, Pringle A. 2017. Long-distance dispersal of fungi. *Microbiology Spectrum* 5: 1-24.

Gonçalves SC, Haelewaters D, Furci G, Mueller GM. 2021. Include all fungi in biodiversity goals. *Science* 373: 403.

Guerra CA, Bardgett RD, Caon L, Crowther TW, Delgado-Baquerizo M, Montanarella L, Navarro LM, Orgiazzi A, Singh BK. 2021a. Tracking, targeting, and conserving soil biodiversity. *Science* 371: 239-241.

Guerra CA, Delgado-Baquerizo M, Duarte E, Marigiliano O, Görgen C, Maestre FT, Eisenhauer N. 2021b. Global projections of the soil microbiome in the Anthropocene. *Global Ecology and Biogeography* 30: 987-999.

Hawksworth DL, Lücking R. 2017. Fungal diversity revisited: 2.2 to 3.8 million species. *Microbiology Spectrum* 1: 79-95.

Hengl T, MacMillan RA. 2019. Predictive Soil Mapping with R. OpenGeoHub Foundation, Wageningen. www.soilmapper.org.

Hijmans RJ. 2021. raster: Geographic Data Analysis and Modeling. R package version 3.5-9. <https://CRAN.R-project.org/package=raster>

Hurt GC, Chini L, Sahajpal R, Frolking S, Bodirsky BL, Calvin K, Doelman JC, Fisk J, Fujimori S, Klein Goldewijk K, Hasegawa T. 2020. Harmonization of global land use change and management for the period 850–2100 (LUH2) for CMIP6. *Geoscientific Model Development* 13: 5425-5464.

IUCN 2021. The IUCN Red List of Threatened Species. Version 2021-2. International Union for Conservation of Nature, Gland.

Karger DN, Nobis MP, Normand S, Graham CH, Zimmermann NE. 2021. CHELSA-TraCE21k v1. 0. Downscaled transient temperature and precipitation data since the last glacial maximum. *Climate of the Past Discussions* 2021:30.

Karger DN, Schmatz DR, Dettling G, Zimmermann NE. 2020. High-resolution monthly precipitation and temperature time series from 2006 to 2100. *Scientific Data* 7:248.

Kennedy PG, Matheny PB, Ryberg KM, Henkel TW, Uehling JK, Smith ME. 2012. Scaling up: examining the macroecology of ectomycorrhizal fungi. *Molecular Ecology* 21: 4151-4154.

Kennedy PG, Walker JKM, Bogar LM. 2015. Interspecific mycorrhizal networks and non-networking hosts: exploring the ecology of host genus *Alnus*. *Ecological Studies* 224: 227-254.

Kier G, Kreft H, Lee TM, Jetz W, Ibsch PL, Nowicki C, Mutke J, Barthlott W. 2009. A global assessment of endemism and species richness across island and mainland regions. *Proceedings of the National Academy of Sciences USA* 106: 9322-9327.

Kinlock NL, Prowant L, Herstoff EM, Foley CM, Akin-Fajiye M, Bender N, Umarani M, Ryu HY, Şen B, Gurevitch J. 2018. Explaining global variation in the latitudinal diversity gradient: Meta-analysis confirms known patterns and uncovers new ones. *Global Ecology and Biogeography* 27: 125-141.

Le Provost G, Thiele J, Westphal C, Penone C, Allan E, Neyret M, Van Der Plas F, Ayasse M, Bardgett RD, Birkhofer K, Boch S. 2021. Contrasting responses of above-and belowground diversity to multiple components of land-use intensity. *Nature Communications* 12:3918.

Makiola A, Dickie IA, Holdaway RJ, Wood JR, Orwin KH, Glare TR. 2019. Land use is a determinant of plant pathogen alpha-but not beta-diversity. *Molecular Ecology* 28: 3786-98.

Maestre FT, Eldridge DJ, Gross N, Le Bagousse-Pinguet Y, Saiz H, Gozalo B, Ochoa V, Gaitán JJ. 2022. The BIODESERT survey: Assessing the impacts of grazing on the structure and functioning of global drylands. *Web Ecology*, doi: 10.5194/we-21-1-2021.

Malcolm GM, Lopez-Gutierrez JC, Koide RT, Eissenstat DM. 2008. Acclimation to temperature and temperature sensitivity of metabolism by ectomycorrhizal fungi. *Global Change Biology* 14: 1169-1180.

Maynard DS, Bradford MA, Covey KR, Lindner D, Glaeser J, Talbert DA, Tinker PJ, Walker DM, Crowther TW. 2019. Consistent trade-offs in fungal trait expression across broad spatial scales. *Nature Microbiology* 4:846.

Misiak M, Goodall-Copestake WP, Sparks TH, Worland MR, Boddy L, Magan N, Convey P, Hopkins DW, Newsham KK. 2021. Inhibitory effects of climate change on the growth and extracellular enzyme activities of a widespread Antarctic soil fungus. *Global Change Biology* 27: 1111-1125.

Newsham KK, Hopkins DW, Carvalhais LC, Fretwell PT, Rushton SP, O'Donnell AG, Dennis PG. 2016. Relationship between soil fungal diversity and temperature in the maritime Antarctic. *Nature Climate Change* 6: 182-186.

Nilsson RH, Larsson K-H, Taylor AFS, Bengtsson-Palme J, Jeppesen TS, Schigel D, Kennedy P, Picard K, Glöckner FO, Tedersoo L, Saar I, Kõljalg U, Abarenkov K. 2019. The UNITE database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Research* 47: D259-D264.

Pebesma E. 2018. Simple Features for R: Standardized Support for Spatial Vector Data. *The R Journal* 10: 439-446.

Oliverio AM, Geisen S, Delgado-Baquerizo M, Maestre FT, Turner BL, Fierer N. 2020. The global-scale distributions of soil protists and their contributions to belowground systems. *Science Advances* 6:eaax8787.

Olson DM, Dinerstein E, Wikramanayake ED, Burgess ND, Powell GWN, Underwood EC, Damico JA, Itoua I, Strand HE, Morrison JC, Loucks CJ, Allnutt TF, Ricketts TH, Kura Y, Lamoreux JF, Wettengel WW, Hedao P, Kassem KR. 2001. Terrestrial ecoregions of the World: a new map of life on earth. *Bioscience* 51: 933-938.

Poggio L, de Sousa LM, Batjes NH, Heuvelink G, Kempen B, Ribeiro E, Rossiter D. 2021. SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty. *Soil* 7: 217-240.

Pölmé S, Abarenkov K, Nilsson RH, Lindahl BD, Clemmensen KE, Kauserud H, Nguyen N, Kjoller R, Bates ST, Baldrian P. FungalTraits: a user-friendly traits database of fungi and fungus-like stramenopiles. *Fungal Diversity* 105: 1-16.

Pölmé S, Bahram M, Jacquemyn H, Kennedy P, Kohout P, Moora M, Oja J, Öpik M, Pecoraro L. 2018. Host preference and network properties in biotrophic plant-fungal associations. *New Phytologist* 217: 1230-1239.

Potapov AM, Guerra CA, van den Hoogen J, Babenko A, Bellini BC, Berg MP, Chown SL, Deharveng L, Kova L, Kuznetsova NA, Ponge JF. 2022. Globally invariant metabolism but density-diversity mismatch in springtails. *bioRxiv* 2022:475345.

Qian H, Jin Y. 2016. An updated megaphylogeny of plants, a tool for generating plant phylogenies and an analysis of phylogenetic community structure. *Journal of Plant Ecology* 9: 233-239.

R Core Team. 2021. R: a language and environment for statistical computing. <http://www.R-project.org/>.

Rillig MC, Ryo M, Lehmann A, Aguilar-Trigueros CA, Buchert S, Wulf A, Iwasaki A, Roy J, Yang G. 2019. The role of multiple global change factors in driving soil functions and microbial biodiversity. *Science* 366: 886-890.

Rodriguez-Ramos JC, Cale JA, Cahill Jr, JF, Simard SW, Karst J, Erbilgin N. 2021. Changes in soil fungal community composition depend on functional group and forest disturbance type. *New Phytologist* 229: 1105-1117.

Romero-Olivares AL, Allison SD, Treseder KK. 2017. Soil microbes and their response to experimental warming over time: a meta-analysis of field studies. *Soil Biology and Biochemistry* 107: 32-40.

Rosauer DF, Jetz W. 2015. Phylogenetic endemism in terrestrial mammals. *Global Ecology and Biogeography* 24: 168-179.

Sanchez-Ramirez S, Etienne RS, Moncalvo J-M. 2015. High speciation rate at temperate latitudes explains unusual diversity gradients in a clade of ectomycorrhizal fungi. *Evolution* 8: 2196-2209.

Sandel B, Weigelt P, Kreft H, Keppel G, van der Sande MT, Levin S, Smith S, Craven D, Knight TM. 2020. Current climate, isolation and history drive global patterns of tree phylogenetic endemism. *Global Ecology and Biogeography* 29: 4-15.

- Schmidt PA, Schmitt I, Otte J, Bandow C, Römbke J, Bálint M, Rolshausen G. 2017. Season-long experimental drought alters fungal community composition but not diversity in a grassland soil. *Microbial Ecology* 75: 468-478.
- Schulte to Bühne H, Tobias JA, Durant SM, Pettoirelli N. 2020. Improving predictions of climate change–land use change interactions. *Trends in Ecology & Evolution* 36: 29-38
- Sivakumar MV. 2007. Interactions between climate and desertification. *Agricultural and Forest Meteorology* 142: 143-155.
- Smith P, Calvin K, Nkem J, Campbell D, Cherubini F, Grassi G, Korotkov V, Le Hoang A, Lwasa S, McElwee P, Nkonya E. 2020a. Which practices co-deliver food security, climate change mitigation and adaptation, and combat land degradation and desertification? *Global Change Biology* 26: 1532-1575.
- Smith RJ, Jovan S, McCune B. 2020b. Climatic niche limits and community-level vulnerability of obligate symbioses. *Journal of Biogeography* 47: 382-395.
- Soininen J, Korhonen JJ, Luoto M. 2013. Stochastic species distributions are driven by organism size. *Ecology* 94: 660–670.
- Stein A, Gerstner K, Kreft H. 2014. Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales. *Ecology Letters* 17: 866-880.
- Talbot JM, Bruns TD, Taylor JW, Smith DP, Branco S, Glassman SI, Erlandson S, Vilgalys R, Laio H-L, Smith ME, Peay KG. 2014. Endemism and functional convergence across the North American mycobiome. *Proceedings of the National Academy of Sciences USA* 111: 6341–6346.
- Tedersoo L, Albertsen M, Anslan S, Callahan B. 2021a. Perspectives and benefits of high-throughput long-read sequencing in microbial ecology. *Applied and Environmental Microbiology* 87:e00626-21.
- Tedersoo L, Bahram M, Pölme S, Kõljalg U, Yorou NS, Wijesundera R, Villarreal-Ruiz L, Vasco-Palacios A, Quang Thu P, Suija A, Smith ME, Sharp C, Saluveer E, Saitta A, Ratkowsky D, Pritsch K, Riit T, Pöldmaa K, Piepenbring M, Phosri C, Peterson M, Parts K, Pärtel K, Otsing E, Nouhra E, Njouonkou AL, Nilsson RH, Morgado LN, Mayor J, May TW, Kohout P, Hosaka K, Hiiesalu I, Henkel TW, Harend H, Guo L, Greslebin A, Grelet G, Geml J, Gates G, Dunstan W, Dunk C, Drenkhan R, Dearnaley J, De Kesel A, Dang T, Chen X, Buegger F, Brearley FQ, Bonito G, Anslan S, Abell S, Abarenkov K. 2014. Global diversity and geography of soil fungi. *Science* 346: 1078.
- Tedersoo L, Mikryukov V, Anslan S, Bahram M, Khalid AN, Corrales A, Agan A, Vasco-Palacios AM, Saitta A, Antonelli A, Rinaldi AC. 2021b. The Global Soil Mycobiome consortium dataset for boosting fungal diversity research. *Fungal Diversity* 111: 573-588.
- Tedersoo L, Sánchez-Ramírez S, Kõljalg U, Bahram M, Döring M, Schigel D, May T, Ryberg M, Abarenkov K. 2018. High-level classification of the Fungi and a tool for evolutionary ecological analyses. *Fungal Diversity* 90: 135–159.
- Tsirogiannis C, Sandel B. 2016. PhyloMeasures: a package for computing phylogenetic biodiversity measures and their statistical moments. *Ecography* 39: 709-714.
- Vazquez DP, Stevens RD. 2004. The latitudinal gradient in niche breadth: concepts and evidence. *The American Naturalist* 164: E1-E9.
- Vetrovsky T, Kohout P, Kopecký M, Machac A, Man M, Bahnmann BD, Brabcová V, Choi J, Meszárošová L, Human ZR, Lepinay C, Baldrian P. 2019. A meta-analysis of global fungal distribution reveals climate-driven patterns. *Nature Communications* 10: 1-9.
- Villéger S, Brosse S. 2012. Measuring changes in taxonomic dissimilarity following species introductions and extirpations. *Ecological Indicators* 18: 552-558.
- Wardle DA, Lindahl BD. 2014. Disentangling global soil fungal diversity. *Science* 346: 1052-1053.
- Warren R, VanDerWal J, Price J, Welbergen JA, Atkinson I, Ramirez-Villegas J, Osborn TJ, Jarvis A, Shoo LP, Williams SE, Lowe J. 2013. Quantifying the benefit of early climate change mitigation in avoiding biodiversity loss. *Nature Climate Change* 3: 678-682.
- Watson JE, Iwamura T, Butt N. 2013. Mapping vulnerability and conservation adaptation strategies under climate change. *Nature Climate Change* 3: 989-994.
- Wood SN. 2011. Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *Journal of the Royal Statistical Society B* 73: 3-36.

Figures




Fig. 1. Distribution of samples and fungal species across datasets. (A) Global sampling map, with different symbols representing different datasets; (B) species distribution of fungi among datasets, with the proportion of unique and shared species indicated in the diagram; (C) Krona chart indicating taxonomic distribution of fungal species (interactive chart can be browsed at <https://plutof.ut.ee/#/doi/10.15156/BIO/2483900>); (D) species richness and total read abundance of the top 10 most diverse fungal genera.




Fig. 2. Comparison of (A) richness proxies (use of log-transformation, residuals of sequencing depth, SRS or simple rarefaction) and (B) measures of soil pH on analytical performance. Relative goodness was estimated based on the determination coefficients of the best models (A) or pH-only models (B). In left panels, significant among-group differences are indicated with different letters based on Tukey Posthoc tests; bars, means; whiskers, SE. Soil pH_{KCl} were determined experimentally, whereas pH_{H₂O} were obtained from Poggio et al. (2021).




Fig. 3. Relative 'species' accumulation curves (A-C), sequencing depth (D) and 'species' richness (E-F) across four datasets. (A) The log-log relationship between the number of reads and 'species' richness that was used for calculation of residuals and further analyses; (B-C) Relatively lower performance of log-linear relationships of log-transformed and square-root-transformed sequencing depth; (D) Initial differences in sequencing depth among datasets; (E-F) Fungal 'species' richness differences relative to the average in the raw data (E) and residuals of the log-log regression analysis (F). In D-F, boxes indicate standard errors around the mean and whiskers indicate 95% confidence intervals; letters above whiskers indicate statistically significant differences among datasets (using log-transformed data for D-E). These analyses indicate that the log-log transformation for calculating residuals is relatively more robust compared with other methods and that richness estimates from studies with different methods cannot be directly compared.




Fig. 4. Response of α -diversity of all fungi to soil pH, vegetation age, mean annual precipitation, latitude, biomes, and land cover categories. For continuous predictors, black lines indicate linear and polynomial fits and red lines indicate lowess fits. For categorical predictors, boxes represent standard error around the mean (central line), whiskers depict 95% CI and letters above boxes indicate statistically significant different groups ($P < 0.001$).




Fig. 5. Comparison of α -diversity patterns in all fungi across four most inclusive datasets: (A) best models (only bioclimatic variables and soil pH were included in model selection), (B) lowess regression curves for the best-fitting climatic predictor, and (C) lowess regression curves for soil pH.




Fig. 6. The effect of average mean annual precipitation on γ -diversity of fungi at the ecoregion scale. Black line, best quadratic fit; red line, lowess curve.




Fig. 7. Distribution of 174 ecoregions used in endemicity analyses. Ecoregions excluded from the analyses due to the lack of data are indicated in gray. Their explanation is given in Table 1.




Fig. 8. The effect of average mean annual precipitation on endemicity of fungi at the ecoregion scale. Black line, best quadratic fit; red line, lowess curve.




Fig. 9. Vulnerability of fungi and functional groups to global change drivers. Different letters indicate statistically significant ($P < 0.001$) differences among functional groups (a-e) and among global change drivers within functional groups (x-z).




Fig. 10. Relationships between conservation priority areas with mean annual temperature (A) and Copernicus land cover types (B). In A, black and red lines indicate best-fitting linear and lowess functions, respectively. In B, central lines and whiskers indicate mean and standard errors, respectively; letters above whiskers indicate statistically significant differences among land cover types.

Calculation of endemism indices

To estimate endemism at the ecoregion level, we calculated five endemism indices (E_1 - E_5). Weighted endemism indices were calculated based on these five indices using z-transformation of residuals in regression analyses against sequencing and sampling depth.

Five endemism indices

E_1 Number of endemic species	E_1	
E_2 Proportion of endemic species	$E_2 = \frac{E_1}{S}$	S , total species richness in ecoregion
E_3 Mean maximum range of species (average weighted endemism; Crisp et al. 2001)	$E_3 = -\frac{\sum_i d_{ri}}{S}$	d_{ri} , range of species i
E_4 Jaccard index of similarity (Jaccard 1912)	$E_4(C_1, C_2) = \frac{C_1 \cap C_2}{C_1 \cup C_2}$	C , ecoregion communities
E_5 β_{SIM} index of similarity (multiple-site Simpson-based community similarity index, Baselga et al. 2010)	$E_5 = \frac{\sum_{i < j} \min(b_{ij}, b_{ji})}{\sum_i (S_i - S_i) + \sum_{i < j} \min(b_{ij}, b_{ji})}$	S_i , number of species in site i S_j , total number of species b_{ij} and b_{ji} , number of species exclusive to sites i and j , respectively

Accounting for sampling and sequencing depth for endemism indices

E_{nR} residuals of n endemism index	$E_{nR} = E_n - \hat{E}_n$ $\hat{E}_n = \beta_1 \log(L_1 + L_2) + \beta_2 \log(R_1 + R_2) + \beta_3 \log(L_1) + \beta_4 \log(L_2) + \beta_5 \log(R_1) + \beta_6 \log(R_2) + \beta_7 (L_1 + L_2) + \beta_8 (R_1 + R_2) + \beta_9 L_1 + \beta_{10} L_2 + \beta_{11} R_1 + \beta_{12} R_2 + \varepsilon$	L_1 , number of soil sampling localities L_2 , number of UNITE localities R_1 , number of reads of soil samples R_2 , number of UNITE Sanger reads
--	--	---

Residuals are calculated based on the best-fitting model.

Standardizing residuals of endemism indices

E_{nRz} Z-transformed E_{nR}	$E_{nRz} = \frac{E_{nR} - \mu}{\sigma}$	μ , mean E_{nR} σ , standard deviation of E_{nR}
----------------------------------	---	--

Weighting standardized endemism indices

E_{ave} weighted E_{nRz}	$E_{ave} = \frac{E_{1Rz} + E_{2Rz} + 2E_{3Rz} + E_{4Rz} + E_{5Rz}}{6}$
------------------------------	--

E_3 receives double weight, because E_1 and E_2 as well as E_3 and E_4 are calculated on a similar basis.

Correlation matrix of endemism indices

	E_{ave}	E_{1Rz}	E_{2Rz}	E_{3Rz}	E_{4Rz}	E_{5Rz}
E_{ave}	1.000	0.767	0.805	0.554	0.754	0.816
E_{1Rz}	0.767	1.000	0.806	0.188	0.452	0.540
E_{2Rz}	0.805	0.806	1.000	0.072	0.788	0.745
E_{3Rz}	0.554	0.188	0.072	1.000	0.169	0.270
E_{4Rz}	0.754	0.452	0.788	0.169	1.000	0.782
E_{5Rz}	0.816	0.540	0.745	0.270	0.782	1.000

Correlation matrix of average endemism of fungal groups

	1	2	3	4	5	6	7	8	9
all fungi (1)	1.000	0.797	0.591	0.517	0.568	0.628	0.675	0.619	0.523
non-EcM Agaricomycetes (2)	0.797	1.000	0.790	0.398	0.461	0.470	0.460	0.595	0.532
AM fungi (3)	0.591	0.790	1.000	0.280	0.375	0.232	0.233	0.578	0.461
EcM fungi (4)	0.517	0.398	0.280	1.000	0.234	0.268	0.250	0.270	0.176
molds (5)	0.568	0.461	0.375	0.234	1.000	0.600	0.413	0.364	0.513
OHP (6)	0.628	0.470	0.232	0.268	0.600	1.000	0.809	0.159	0.354
pathogens (7)	0.675	0.460	0.233	0.250	0.413	0.809	1.000	0.264	0.331
unicellular fungi (8)	0.619	0.595	0.578	0.270	0.364	0.159	0.264	1.000	0.463
yeasts (9)	0.523	0.532	0.461	0.176	0.513	0.354	0.331	0.463	1.000

References

Baselga, A. Partitioning the turnover and nestedness components of beta diversity. *Glob. Ecol. Biogeogr.* **19**, 134-143 (2010).
Jaccard, P. The distribution of the flora in the alpine zone. *New Phytol.* **11**, 37-50 (1912).

Box 1. Calculation of endemism indices and correlation among standardized indices for all fungi and among functional groups.

864 **Table 1. The ecoregions used in endemicity analyses.**

Ecoregion number	Putative name
1	Alaskan forests
2	Alaskan tundra
3	Albertine Rift montane forests
4	Yunga
5	Alps conifer and mixed forests
6	Altai steppes and semideserts
7	Anatolian forests
8	Andaman Islands rain forests
9	Yucatan forests
10	Antarctic desert
11	Appalachian mixed forests
12	Appennine forests
13	Arabic drylands
14	Argentina savanna
15	Australia Central drylands
16	Australia E subtropical woodlands
17	Australia NE savannas
18	Australia NW savannas
19	Australia SE shrublands and savannas
20	Australia SE temperate forests
21	Australia SW woodlands
22	Australia W savannas
23	Australia W shrublands
24	Balkan forests
25	Baltic mixed forests
26	Benelux Atlantic mixed forests
27	Borneo-Java rain forests
28	Botswana woodlands
29	Brazil S forests
30	Burman forests
31	Caatinga
32	California Sierra Nevada forests
33	Cameroon W forests
34	Canada N boreal forests
35	Canada NE boreal forests
36	Canada NW boreal forests
37	Canary Islands dry woodlands and forests
38	Cape fynbos
39	Cape karoo
40	Cape thickets
41	Cape Verde Islands dry forests
42	Caucasia
43	Central American forests
44	Central Asian drylands

- 45 Central Atlantic rain forests
- 46 Central Congolian forests
- 47 Central European mixed forests
- 48 Central Siberian tundra
- 49 Central Sudanian savanna
- 50 Cerrado
- 51 Chaco
- 52 Chilean matorral
- 53 China Central forests
- 54 China E forests
- 55 Colombia montane forests
- 56 Zimbabwe-Mozambique woodlands
- 57 Colombia N forests
- 58 Colombia SW dry forests
- 59 Colombia W forests
- 60 Crete Mediterranean forests
- 61 Cuba forests
- 62 Czech mixed forests
- 63 Da Hinggan-Dzhagdy Mountains conifer forests
- 64 East African shrublands
- 65 East African woodlands
- 66 East Asian drylands
- 67 East European forest steppe
- 68 East Guinean forests
- 69 East Himalayan forests
- 70 East Siberia forest and mountain tundra
- 71 West Amazon forests
- 72 Eastern South African woodlands
- 73 Estonian Sarmatic mixed forests
- 74 Ethiopian montane woodlands
- 75 Fiji forests
- 76 Finland taiga
- 77 Florida forests
- 78 France Atlantic mixed forests
- 79 Great Britain Forests
- 80 Great Lakes forests
- 81 Zambia woodlands
- 82 Greenland tundra
- 83 Hawaii forests
- 84 Highveld grasslands
- 85 Iceland boreal birch forests and alpine tundra
- 86 Zagros Mountains forest steppe
- 87 Isthmian forests
- 88 Japan forests
- 89 Kalahari xeric savanna
- 90 Victoria Basin forest-savanna mosaic
- 91 Karakoram drylands

- 92 Western Ghats forests
- 93 Korean forests
- 94 Latvian Sarmatic mixed forests
- 95 Lesser Antilles forests
- 96 Madagascar woodlands
- 97 Madeira evergreen forests
- 98 Magellanic subpolar forests
- 99 Manchurian forests
- 100 Mascarene forests
- 101 Western European broadleaf forests
- 102 Mexico NE shrublands and forests
- 103 Mexico NW deserts and montane forests
- 104 Mexico S dry forests
- 105 Mexico SW lowland forests
- 106 Mexico SW montane forests
- 107 Mongolian deserts
- 108 Mongolian steppe
- 109 Namib drylands
- 110 New Caledonia forests
- 111 New Guinea forests
- 112 New Zealand forests
- 113 North African woodlands
- 114 North Amazon forests
- 115 North Atlantic rain forests
- 116 North India moist forests
- 117 Northeast Siberian taiga
- 118 Northeast Siberian tundra
- 119 Northern Indochina subtropical forests
- 120 Northern Zanzibar-Inhambane coastal forest mosaic
- 121 Western Congolian forests
- 122 West Sudanian savanna
- 123 Oman drylands
- 124 Pacific deserts
- 125 Pannonian forests
- 126 Pantelleria-Lampedusa forests
- 127 Patagonian steppe
- 128 Peninsular Malaysian rain forests
- 129 West Siberian forests
- 130 Polynesian forests
- 131 Portugal woodlands
- 132 Puerto Rico forests
- 133 Puna
- 134 Queensland tropical rain forests
- 135 Russian Sarmatic mixed forests
- 136 Russian taiga
- 137 Sahelian drylands
- 138 Sardinian-Corsican forests

- 139 Scandinavian coastal conifer forests
 - 140 Scandinavian Montane Birch forest and grasslands
 - 141 Scandinavian Sarmatic mixed forests
 - 142 Scandinavian taiga
 - 143 Sicilian forests
 - 144 South Asian drylands
 - 145 South Atlantic rain forests
 - 146 South Caspian forests
 - 147 South Siberian steppe
 - 148 Southeast African bushveld
 - 149 Southeast Amazon forests
 - 150 Southern Indochina tropical forests
 - 151 Spain woodlands
 - 152 Sri Lanka forests
 - 153 Subantarctic islands tundra
 - 154 West Himalayan forests
 - 155 Svalbard Arctic desert
 - 156 Taiwan forests
 - 157 Tarim basin drylands
 - 158 Tasmanian forests
 - 159 Tenasserim-South Thailand semi-evergreen rain forests
 - 160 Tibetan steppe
 - 161 Trans-Baikal forests
 - 162 Transylvanian forests
 - 163 West Guinean forests
 - 164 USA California chaparral
 - 165 USA Central forests and grasslands
 - 166 USA NE hemiboreal forests
 - 167 USA SE forests
 - 168 USA Southern Rockies and steppe
 - 169 USA western steppe and montane forests
 - 170 USA-Canada Pacific forests and grasslands
 - 171 USA-Canada Rockies and central forests and grasslands
 - 172 Ussuri broadleaf and mixed forests
 - 173 Valdivian temperate forests
 - 174 Venezuela woodlands
-

865

866

867 **Table 2. The best predictors of endemism indices in ecoregions for all fungi.**

	DF	Sum of squares	Mean squares	F-value	R ² _{adj}	P-value	Trend
MAT _{mean}	2	27.2	13.6	97.4	0.277	<0.001	U-shaped
soil pH _{mean}	1	10.6	10.6	38.2	0.108	<0.001	negative
subcontinent: Europe	1	6.5	6.5	23.5	0.065	<0.001	negative
human footprint index	1	2.1	2.1	7.5	0.018	0.007	negative
error	168	48.3					

868

869 **Table 3. The best models of conservation priority co-kriging maps for fungi. All P-values are <0.001.**

	Df	Sum of squares	Mean squares	F-value	R ² _{adj}	Trend
air MAT ¹	2	2133	1066	1426	0.266	positive
MPWM	1	992	992	663	0.134	positive
climate isothermality	1	169	169	113	0.023	positive
MAP	2	167	84	112	0.022	positive
Land cover type ²	7	128	18	12	0.015	
Land cover type x MAP	7	121	17	12	0.014	
error	2477	3705				

¹Abbreviations: MAP, mean annual precipitation; MAT, mean annual temperature; MPWM, mean precipitation of wettest month;

²Copernicus land cover types: broadleaf forest, mixed forest, coniferous forest, shrubland, tundra, grassland, desert, wetland (note that cropland and urban and village biomes were excluded).

870