

Mitochondrial phospholipid import mediated by VDAC, a dimeric beta barrel scramblase

Helene Jahn¹, Ladislav Bartoš^{2,3}, Joost C. M. Holthuis⁴, Robert Vácha^{2,3,*}, Anant K. Menon^{1,*}

¹Department of Biochemistry, Weill Cornell Medical College; New York, NY 10065, USA

²CEITEC - Central European Institute of Technology, Masaryk University; Kamenice 5, 62500 Brno, Czech Republic

³National Centre for Biomolecular Research, Faculty of Science, Masaryk University; Kamenice 5, 62500 Brno, Czech Republic

⁴Department of Molecular Cell Biology, University of Osnabrück; Osnabrück 49076, Germany

*Corresponding authors: E-mails: robert.vacha@muni.cz, akm2003@med.cornell.edu

Character count: 64,346 (main text: 45,370; References: 12,237; Main figure legends: 6,088)

ABSTRACT

Mitochondria depend on phospholipids supplied by the endoplasmic reticulum, but how these lipids cross the outer membrane to enter mitochondria is unclear. We identified the voltage-dependent ion channel (VDAC) as a scramblase that facilitates lipid entry. Our analyses show that phospholipid transport across the outer membrane is impaired in yeast mitochondria lacking VDAC homologs. On reconstitution into membrane vesicles, dimers of human VDAC1 and VDAC2 facilitate rapid scrambling, whereas VDAC1 monomers are less active. Molecular dynamics simulations of VDAC1 dimers support rapid scrambling at a dimer interface where polar residues induce large water defects and bilayer thinning. Thus, VDAC, a member of a superfamily of beta barrel proteins, moonlights as a new class of phospholipid scramblase operating by an unprecedented mechanism.

Key words: VDAC, scramblase, phospholipid, dimer, mitochondria

INTRODUCTION

The double membrane of mitochondria is composed of phospholipids which are supplied by the endoplasmic reticulum (ER) or assembled *in situ* from ER-derived phospholipid precursors (Horvath & Daum, 2013; Tamura *et al.*, 2020; Tatsuta & Langer, 2017). For example, cardiolipin, the signature lipid of mitochondria, is synthesized on the matrix side of the inner mitochondrial membrane (IMM) from the ER-derived phospholipid phosphatidic acid (PA), and subsequently remodeled in the inter-membrane space (IMS) by enzymatic exchange of acyl chains with ER-derived phosphatidylcholine (PC) (Horvath & Daum, 2013; Schlame & Greenberg, 2017; Tamura *et al.*, 2020; Tatsuta & Langer, 2017). This poses a considerable lipid trafficking problem, as, after delivery to the cytoplasmic face of the outer mitochondrial membrane (OMM) by non-vesicular mechanisms (Reinisch & Prinz, 2021), PA, PC, and other phospholipids must cross the barrier of the OMM before moving on to the IMM (Fig. 1A). Previous work suggested that phospholipids are scrambled across the OMM, i.e., flip-flopped across the OMM by an ATP-independent mechanism (Dolis *et al.*, 1996; Janssen *et al.*, 1999; Jasinska *et al.*, 1993; Lampl *et al.*, 1994; Pomorski & Menon, 2016), but the identity of the OMM scramblase(s) is not known.

The Voltage Dependent Anion Channel (VDAC) is an abundant, multi-functional OMM β -barrel protein with three isoforms (VDAC1-3) in mammals and two (Por1-2) in budding yeast (Bergdoll, 2017; De Pinto, 2021; Di Rosa *et al.*, 2021; Raghavan *et al.*, 2012; Shoshan-Barmatz *et al.*, 2015; Zeth & Zachariae, 2018). It forms the pore through which ATP and other metabolites cross the OMM and plays a key role in apoptosis (Bergdoll, 2017; De Pinto, 2021; Di Rosa *et al.*, 2021; Raghavan *et al.*, 2012; Shoshan-Barmatz *et al.*, 2015; Zeth & Zachariae, 2018). The different isoforms also play non-redundant roles in cells (Bergdoll, 2017; Messina *et al.*, 2012; Raghavan *et al.*, 2012). Previous molecular dynamics (MD) simulations of mouse VDAC1 (Villinger *et al.*, 2010) indicated membrane thinning near an outward-facing glutamate residue (E73), situated midway along the β -barrel. This thinning was observed to promote phospholipid flip-flop (Villinger *et al.*, 2010), with headgroups transiently approaching E73 as the lipids crossed from one side of the bilayer to the other. Using coarse-grained MD (CGMD) simulations, we confirmed that the surface of VDAC1 provides a low energy path for phospholipid transit across

the bilayer, populated by membrane-facing polar residues and centered on E73 (Fig. S1). These observations suggested that VDAC1 - and perhaps other VDAC isoforms - may be able to facilitate phospholipid rapid flip-flop, thereby providing the scramblase activity needed to move lipids across the OMM. Here we show that both VDAC1 and VDAC2 are phospholipid scramblases. However, the hypothesized pathway centered on E73 appears not to play an important role as fast scrambling requires dimerization, such that phospholipids transit between the leaflets of the bilayer at a dimer interface where polar residues induce large water defects and bilayer thinning.

RESULTS

A role for VDAC in phospholipid entry into mitochondria. As an initial test of the possibility that VDAC plays a role in scrambling phospholipids across the OMM, we assayed the ability of isolated yeast mitochondria to convert exogenously supplied phosphatidylserine (PS) to phosphatidylethanolamine (PE), a process requiring transport of PS across the OMM to the site of PS decarboxylase (Psd1) in the IMS (Fig. S2A) (Tatsuta & Langer, 2017). Mitochondria were isolated from wild type yeast as well as strains lacking the VDAC homologs Por1 and/or Por2, adjusted to equal protein concentration (reflected in immunoblots showing similar levels of Psd1 and Tom70 (Fig. S2B)), and assayed for transport-coupled PS decarboxylation (Fig. S2C). We used the short-chain PS analog NBD-PS for these experiments because its slight water-solubility allowed its delivery to the assay without the need of solvents that might disrupt OMM integrity, and because its NBD fluorophore enabled easy visualization of its conversion to NBD-PE on thin layer chromatograms (Fig. S2C). Compared with wild type mitochondria, we found that the rate of PE production was ~1.5 fold lower in mitochondria lacking Por1 and Por2 (Fig. S2D,E), consistent with the idea that efficient trans-bilayer movement of PS across the OMM depends on VDAC.

VDAC2 scrambles phospholipids in reconstituted vesicles. We next directly evaluated the ability of human VDAC proteins to scramble phospholipids using a previously described assay (Wang *et al*, 2018)(Fig. 1B). We purified correctly folded human VDAC2 after producing it recombinantly in *E. coli* (Fig. S3) and reconstituted it into large unilamellar vesicles composed mainly of PC with a trace quantity of [³H]inositol-labeled phosphatidylinositol ([³H]PI). Protein-free vesicles were prepared in parallel. On adding bacterial PI-specific phospholipase C (PI-PLC) to the protein-free vesicles we observed rapid hydrolysis of ~50% of the [³H]PI, corresponding to those molecules residing in the outer leaflet of the vesicles and therefore accessible to the enzyme. However, in samples reconstituted with VDAC2, the extent of hydrolysis increased to ~75% indicating flipping of [³H]PI molecules from the inner to the outer leaflet in a large fraction of the vesicles, at a rate similar to or greater than that of the hydrolysis reaction. As flipping was observed in the absence of any metabolic energy supplied to the system, this result indicates that VDAC2 is a scramblase (Fig. 1C).

We further characterized lipid scrambling by VDAC2 using fluorescence-based methods (Chang *et al*, 2004; Kubelt *et al*, 2002; Menon *et al*, 2011). VDAC2 was reconstituted into vesicles containing a trace quantity of 1-myristoyl-2-C6-NBD-PC, a fluorescent PC analog. As VDAC2 is a channel/pore, reconstitution efficiency was assessed using dithionite-mediated

bleaching of NBD-PC (Fig. 1D). All NBD-PC molecules are expected to be bleached in a vesicle that contains a VDAC2 channel, which permits dithionite entry, whereas only those molecules in the outer leaflet will be bleached in vesicles lacking a channel. Fig. 1E shows that >90% of the NBD-PC is bleached, indicating efficient reconstitution of VDAC2 (~53% bleaching was observed in protein-free vesicles, as expected for 150 nm-diameter vesicles). Next, scramblase activity was assayed using fatty acid-free bovine serum albumin (BSA), which extracts NBD-PC molecules from the outer leaflet of the vesicles, resulting in a decrease in fluorescence as the quantum efficiency of NBD-PC complexed to BSA is ~60% lower than that of NBD-PC in the membrane (Chang *et al.*, 2004; Kubelt *et al.*, 2002; Menon *et al.*, 2011)(Fig. 1F). Thus, quantitative extraction of outer leaflet NBD-PC from protein-free liposomes causes ~30% reduction in fluorescence as seen in Fig. 1G. For vesicles reconstituted with VDAC2, the observed fluorescence reduction is ~55% (Fig. 1G), the extent predicted if ~90% of the vesicles include a scramblase, consistent with the reconstitution efficiency deduced from the channel assay. Curve fitting revealed that fluorescence loss on BSA treatment of VDAC2-containing vesicles is characterized by a rapid phase (half-time ~5-10 s, also seen in protein-free samples) corresponding to the extraction of NBD-PC located initially in the outer leaflet, followed by a slower phase (half-time ~50 s) which we attribute to trans-bilayer movement. These data reaffirm that VDAC2 is a scramblase, capable of transporting both anionic (PI) and zwitterionic (PC) phospholipids.

To elucidate the scrambling mechanism, we tested the effect of disrupting the predicted lipid transit pathway (Fig. S1) by substituting the membrane-facing glutamate residue (E84 in VDAC2) with leucine. We found, unexpectedly, that the E84L mutation had no detectable impact on the ability of VDAC2 to scramble NBD-PC (Fig. S4), suggesting that the hypothesized pathway is not the principal avenue for lipid scrambling.

VDAC dimers support fast scrambling. Whereas channel activity is intrinsic to a VDAC monomer (Bergdoll, 2017; Zeth & Zachariae, 2018), we considered the possibility that scramblase activity may depend on VDAC's quaternary structure. VDAC is known to dimerize and oligomerize, both in the OMM as visualized by atomic force microscopy (Goncalves *et al.*, 2008; Hoogenboom *et al.*, 2007), as well as in detergent solution and after reconstitution into vesicles (Bayrhuber *et al.*, 2008; Bergdoll *et al.*, 2018; Betaneli *et al.*, 2012; Geula *et al.*, 2012; Schredelseker *et al.*, 2014; Shoshan-Barmatz *et al.*, 2013; Zalk *et al.*, 2005). Various factors, including pH, regulate the oligomeric behavior of VDAC (Bergdoll *et al.*, 2018; Betaneli *et al.*, 2012; Kim *et al.*, 2019; Schredelseker *et al.*, 2014; Shoshan-Barmatz *et al.*, 2013; Zalk *et al.*, 2005). To establish the functional oligomeric state of VDAC, we determined vesicle occupancy statistics in our reconstituted system (Cliff *et al.*, 2020; Goren *et al.*, 2014; Ploier *et al.*, 2016). We varied the protein/phospholipid ratio (PPR) of the preparation by reconstituting different amounts of VDAC2 into a fixed quantity of lipid and used the channel assay to determine the fraction of vesicles, at each ratio, that had been functionalized with a VDAC2 channel. The same analyses were also performed using purified human VDAC1 (Fig. S3). As shown in Fig. 2A,B, both VDAC2 and VDAC1 readily functionalize vesicles with channels, with Poisson analysis indicating that they reconstitute as higher order structures that we estimate to be decamers (Fig. 2C). This is likely because the proteins are intrinsically multimeric as purified (Fig. S5) and/or multimerize during detergent removal *en route* to reconstitution as seen in other instances (Betaneli *et al.*, 2012; Goren *et al.*, 2014; Ploier *et al.*, 2016).

The same vesicle preparations were next taken for scramblase assays. VDAC2 showed robust scramblase activity as expected, the fraction of functionalized vesicles scaling with the protein/phospholipid ratio as seen for the channel assays (Fig. 2D). However, VDAC1 was surprisingly less potent (Fig. 2E). To evaluate this difference quantitatively, we graphed the fraction of scramblase-active vesicles versus the fraction of channel-active vesicles for both isoforms (Fig. 2F). For VDAC2, these fractions were highly correlated, with ~85% of the channel active vesicles displaying scramblase activity over the range of the protein/phospholipid ratios that we tested. However, for VDAC1, scramblase activity was essentially undetectable on the timescale of the assay until sufficient protein was reconstituted to populate about a third of the vesicles with channels. As the fraction of channel-active vesicles increased beyond ~0.3, the fraction of scramblase-active vesicles also increased but less efficiently (correlation ~70%) than in the case of VDAC2 (Fig. 2F).

To understand the molecular basis for the difference in the behavior of these otherwise highly similar isoforms of VDAC (75% identity, 91% positive match (Messina *et al.*, 2012; Schredelseker *et al.*, 2014)), we used chemical crosslinking to probe their quaternary structure after reconstitution. VDAC1- and VDAC2-containing vesicles were treated with EGS, an amine-reactive crosslinker which has been previously used to determine the oligomeric nature of VDAC (Geula *et al.*, 2012; Zalk *et al.*, 2005)(Fig. S7A). Immunoblotting revealed that in the absence of crosslinker, both proteins were detected mainly as monomers, with a small fraction of VDAC2 forming SDS-resistant dimers. However, the crosslinker efficiently captured dimers and multimers of VDAC2 whereas VDAC1 was recovered mainly in monomeric form (Fig. 2G). We conclude that despite their similar incorporation into vesicles (Fig. 2C), VDAC2 molecules remain associated in the membrane such that complexes can be captured by EGS crosslinking, whereas VDAC1 molecules dissociate, thereby losing their ability to facilitate fast scrambling (assays run over an extended time frame reveal that VDAC1 scrambles lipids slowly, with a long half-time of ~4 h (Fig. S6)). These data suggest that a dimer or higher order multimer of VDAC is necessary for rapid scrambling.

VDAC1 dimers could be formed by crosslinking providing an opportunity to determine whether dimerization is sufficient for scrambling. We identified conditions in which we could crosslink VDAC1 efficiently with EGS after purifying it in LDAO detergent. Immunoblot analysis revealed that the crosslinked sample, VDAC1^x, consisted principally of dimers, with a small fraction of higher order multimers (Fig. 3A). The EGS-captured dimers likely correspond to VDAC1 protomers oriented in parallel, interacting via strands β 1,2,18,19 (Fig. S7C,D), similar to the common dimeric structure reported for human and rat VDAC1 (Bayrhuber *et al.*, 2008; Geula *et al.*, 2012), and zebrafish VDAC2 (Schredelseker *et al.*, 2014). Importantly, the parallel orientation of the VDAC1 protomers within the dimer structure is supported by spectroscopic analyses of LDAO-solubilized mouse VDAC1 (Bergdoll *et al.*, 2018) and structural studies of human VDAC1 purified in LDAO (Bayrhuber *et al.*, 2008).

Reconstitution of VDAC1^x and mock-treated VDAC1 showed that they both functionalized vesicles efficiently as evidenced by channel activity (Figure 3B). However, whereas the mock-treated protein was barely active as a scramblase, VDAC1^x showed robust scrambling activity when reconstituted at the same protein/phospholipid ratio (Fig. 3C). Quantification of the results (Fig. 3D) showed that ~90% of channel-active vesicles populated with VDAC1^x are also scramblase active, whereas this number fell to ~30% in the case of reconstituted VDAC

monomers. We note that VDAC1^x scrambles lipids ~2-fold more slowly than native dimers/multimers of VDAC2 (Fig. 3C and 1G), perhaps because of the physical constraints imposed by crosslinking. To verify our results with EGS, we used DSP, a cleavable crosslinker containing a DTT-susceptible disulfide bond (Fig. S7B). On crosslinking LDAO-solubilized VDAC1 with DSP, we obtained mainly dimers (Fig. S7E); these could be restored to monomers upon DTT treatment (Fig. S7E). DSP-crosslinked VDAC1 had scramblase activity (Fig. S7F,G), like VDAC^x, which was largely lost on DTT-treatment of the reconstituted sample (Fig. S7F,G). We conclude that VDAC dimerization is both necessary and sufficient to generate an efficient scramblase, and that the dimer is the likely minimal functional oligomeric state of the protein.

Molecular dynamics simulations reveal a phospholipid transit pathway at a specific dimer interface.

We used coarse-grained molecular dynamics to simulate different VDAC1 dimers and thereby gain insight into the scrambling mechanism. We chose two symmetric dimers, with interfaces mediated by strands β 1,2,18,19 (dimer-1) (Fig. 4A,B) as previously reported (Bayrhuber *et al.*, 2008; Geula *et al.*, 2012; Schredelseker *et al.*, 2014), and β 14-17 (dimer-3), a novel configuration in which the E73 residue is positioned distal to the interface in each protomer (Fig. S8A). We observed a high scrambling rate for dimer-1 (Fig. 4C,D) over 10 μ s of simulation time, with PC molecules moving along the edge of the interface while interacting with both protomers (Fig. 4E,F and Movie S1). In contrast, a VDAC1 monomer supported only slow scrambling, as noted experimentally (Fig. S6), with lipids moving along the originally proposed E73-centered transit path (Fig. S1). Unless restrained, dimer-1 re-oriented during the simulation to dimer-1*, with a symmetric interface mediated by strands β 2-4 (Fig. S8B), at which point the rate fell ~4-fold (Fig. 4C,D). Dimer-3 was largely ineffective (Fig. 4D), scrambling lipids at a rate only marginally higher than that seen for two individual VDAC1 monomers. In addition, we identified a VDAC1 mutant which, although highly multimeric after reconstitution as revealed by EGS crosslinking, scrambled lipids poorly (Fig. S9). These results indicate that a specific dimer interface promotes rapid scrambling.

The dimer-1 interface possesses numerous polar residues (Fig. 4A), but these are clearly insufficient to promote fast scrambling by themselves because the monomeric protein scrambles lipids slowly (Fig. 4C,D, Fig. S6). We therefore considered that the individual polar faces must synergize at the interface, possibly in concert with the membrane, to create a low energy trans-bilayer path for lipids. Indeed, we found that the bilayer was much thinner adjacent to the interface, reaching just over 2 nm (Fig. 4G,H), and exhibited a large degree of water penetration (Fig. 4I). These features could also be seen with dimer-1*, but to a much smaller extent, correlating with its weaker scrambling activity (Fig. S8C,D). In contrast, the membrane was only slightly perturbed in the vicinity of dimer-3 (Fig. S8C,D) and the monomeric protein (Fig. 4G,I). Evaluation of the interactions of the scrambling lipid along the dimer-1 interface (Fig. S10) demonstrated that a substantial portion of the decrease in the flip-flop barrier likely originates from the interaction between the lipid and the residues of VDAC1 along the scrambling pathway. This is readily seen by the increase in the energy barrier for lipid scrambling in simulations (Fig. S10A) in which the attractive dispersion interactions between the lipid and VDAC1 were turned off.

DISCUSSION

We identified the β -barrel proteins VDAC1 and VDAC2, members of a large superfamily of beta barrel proteins (Zeth, 2010; Zeth & Zachariae, 2018), as phospholipid scramblases. Our results indicate, that on specific dimerization, these proteins play a role in the efficient transport of anionic and zwitterionic phospholipids across the OMM (Fig. 1A), which is required for mitochondrial membrane biogenesis and mitochondrial function. As all phospholipid scramblases identified so far are α -helical membrane proteins, our results introduce the transmembrane β -barrel as a new scramblase-active protein fold. The current mechanistic paradigm for α -helical scramblases is that they provide a polar transbilayer pathway which allows phospholipids to traverse the membrane with their headgroup residing within the pathway while their acyl chains extend into the hydrophobic membrane core, as suggested by the credit-card model (Pomorski & Menon, 2006). This polar pathway can be formed by two or three adjacent membrane helices within a protein monomer as seen for GPCR and TMEM16 scramblases (Kalienkova *et al*, 2021; Khelashvili & Menon, 2022), between protomers in an oligomer as in bacteriorhodopsin and Atg9 (Maeda *et al*, 2020; Matoba *et al*, 2020; Verchere *et al*, 2017), or at the surface of the protein (Kalienkova *et al*, 2021; Maeda *et al*, 2020; Malvezzi *et al*, 2018). A polar pathway on the surface of VDAC1 and VDAC2, centered on the membrane facing glutamate, clearly lowers the barrier for lipid scrambling (Fig. S1) but this is not sufficient to support fast scrambling (Fig. S6). In contrast, the dimer-1 interface depresses the free energy barrier far more substantially, enabling fast lipid scrambling. The effectiveness of the dimer-1 interface in promoting fast scrambling derives from the synergistic effect of attractive interactions between lipid headgroups and protein residues (Fig. S10), and barrier lowering caused by membrane thinning and associated water defects (Bennett & Tieleman, 2014). Neither feature alone is sufficient to account for fast scrambling activity. Notably, because of the requirement for quaternary structure in generating a fast scrambling path, it is possible that scrambling at the OMM in cells may be regulated through control of dimerization by other proteins as well as by the lipid environment (Betaneli *et al*, 2012; Chadda *et al*, 2021; Shoshan-Barmatz *et al*, 2013).

Because elimination of both VDAC homologs in yeast slowed but did not abrogate phospholipid import into isolated mitochondria (Fig. S2E), we propose that there are additional scramblases operating redundantly with VDAC at the OMM. Possible candidates are the ubiquitous OMM translocation channel Tom40, a 19-stranded β -barrel which is evolutionarily related to VDAC and shares the same fold (Araiso *et al*, 2019; Tucker & Park, 2019; Wang *et al*, 2020; Zeth, 2010). In support of this idea, Tom40 dimers - like VDAC dimers - have been shown to influence the membrane through bending and destabilization (Araiso *et al*, 2019; Tucker & Park, 2019; Wang *et al*, 2020; Zeth, 2010). Other transmembrane β -barrels of the OMM like Sam50 or Mdm10 might show similar properties (Egea, 2021; Takeda *et al*, 2021). Recent discoveries have advanced the concept that lipid transfer proteins directly engage scramblases (Adlakha *et al*, 2022; Ghanbarpour *et al*, 2021). Thus, Vps13 and the ERMES complex that are involved in lipid flow between the ER and mitochondria physically interact with the yeast OMM proteins Mcp1 and Mdm10, respectively (Adlakha *et al*, 2022). Purified Mcp1 was recently shown to have scramblase activity and it remains to be seen whether this is also the case for Mdm10. In agreement with this concept, previous studies placed VDAC at ER-mitochondria contact sites and at hubs of lipid synthesis and distribution (Broeskamp *et al*, 2021; Miyata *et al*, 2018; Tamura *et al*, 2020). Thus, in yeast, the VDAC homolog Por1 is thought to interact with the

ERMES complex, the Ups-Mdm35 lipid transfer complex as well as Mdm31/32, the latter two being important for phospholipid transport across the IMS (Broeskamp *et al.*, 2021; Miyata *et al.*, 2018; Tamura *et al.*, 2020). In this scenario, VDAC constitutes a nexus between the phospholipid transport machineries on both side of the OMM, scrambling phospholipids to enable their entry(exit) into(from) these machineries.

MATERIALS AND METHODS

VDAC purification.

VDAC proteins were expressed and purified as described by Dadsena *et al* (2019). Chemically competent *E. coli* BL21(DE3)-omp9 cells [F^- , ompT hsdS_B ($r_B^- m_B^-$) gal dcm (DE3) Δ lamB ompF::Tn5 Δ ompA Δ ompC Δ ompN::Q] after Prilipov *et al* (1998) were transformed with pCold vectors encoding N-terminal hexahistidine tagged human VDAC1 and VDAC2 isoforms, the point mutant VDAC2 E84L, or the pentamutant VDAC1-5V (T60V, Y62V, E73V, T83V, S101V). All constructs were verified by Sanger sequencing. For VDAC purification, the transformed cells were grown at 37°C to OD₆₀₀ of 0.6 in 2xYT medium (16 g/l Tryptone, 5 g/l NaCl, 10 g/l Yeast extract) containing 100 µg/ml ampicillin. The cultures were cooled before adding 1 mM IPTG and incubating overnight at 15°C. Cells were harvested by centrifugation and the cell pellet was resuspended in TEN-buffer (50 mM Tris-HCl pH 8.0, 100 mM NaCl) containing a protease inhibitor cocktail (30ml of resuspension for 2L starting cell culture). The cells were disrupted using a probe sonicator and inclusion bodies were collected by ultracentrifugation at 25,000 x g for 30 min at 4°C. Residual cell membranes were removed by washing the pellet thrice with 15 mL TEN-buffer containing 2.5% v/v Triton X-100, using a Teflon pestle for resuspension followed by centrifugation at 25,000 x g 15min 4°C, after which the Triton X-100 was removed by further washing the inclusion bodies thrice in TEN-buffer. Washed inclusion bodies were resuspended in 4 mL TEN buffer for 2L starting cell culture and 2ml were used for immediate denatured by dropwise addition into denaturation buffer (25 mM Na⁺PO₄ pH 7.0, 100 mM NaCl, 6 M guanidine hydrochloride, 1 mM EDTA, 10 mM DTT) to a 10-times dilution. After overnight incubation at 4°C under constant stirring, the proteins were refolded by serial 10-times dilutions into first 25 mM Na⁺PO₄ pH 7.0, 100 mM NaCl, 1 mM EDTA, 2.22% LDAO followed by overnight incubation. This material was further 10-times diluted into 25 mM Na⁺PO₄ pH 7.0, 10 mM NaCl, 1 mM EDTA, 0.1% LDAO. After incubation at 4°C for 4h or overnight, the solution was filtered, applied onto a cation-exchange column (HiTrapTM SP HP 5ml (GE Healthcare)), and eluted with a salt gradient (10 mM to 1 M NaCl in the buffer 25 mM Na⁺PO₄ pH 7.0, 1 mM EDTA, 0.1% LDAO, 1 mM DTT). Protein-containing fractions were pooled, concentrated to 500 µl using Amicon Ultra-4 (Millipore) centrifugal filters, and loaded onto a Superdex 200 10/300 25ml size-exclusion column using SEC buffer (10 mM Tris-HCl pH 8.0, 100 mM NaCl, 0.05% LDAO). Peak fractions were pooled, quantified by absorbance (VDAC1 $\epsilon_{280} = 38,515 \text{ M}^{-1} \cdot \text{cm}^{-1}$, VDAC2 $\epsilon_{280} = 37,400 \text{ M}^{-1} \cdot \text{cm}^{-1}$) and BCA colorimetric assay, assessed for purity by Coomassie-stained SDS-PAGE, and analyzed by Circular Dichroism spectroscopy using Aviv 410 CD instrument. For storage at -80°C, 10% glycerol was added, and the protein was snap frozen. Typical protein concentration obtained was 1 mg/ml from 1L of starting culture.

Large unilamellar vesicles (LUVs).

POPC and POPG (1435 μ l and 160 μ l of 25 mg/ml stock solutions in chloroform), together with (1-myristoyl-2-C₆-NBD-PC) (195 μ l of a 1 mg/ml stock solution in chloroform) as indicated, were added to a round-bottom flask. The solvent was evaporated using a rotary evaporator and the flask was placed overnight in a desiccator attached to house vacuum. The dried lipid film was resuspended by swirling in 10 ml reconstitution buffer (10 mM MOPS/Tris pH 7.0, 100 mM KCl) or crosslinking buffer (10 mM MOPS/KOH pH 7.0, 100 mM KCl) as indicated (lipid concentration 5.25 mM), and incubated at 37°C for 20 min. The resuspended lipids were briefly sonicated in a bath sonicator, before being sequentially extruded through 0.4 μ m and 0.2 μ m membranes in a high-pressure lipid extruder (Northern Lipids). The size (150 nm) and polydispersity index (12.5) of the preparation was determined by dynamic light scattering (DLS) with the LitesizerTM 500 Instrument. The liposomes had a typical concentration of 3 mM.

VDAC reconstitution into LUVs.

VDAC was reconstituted into pre-formed LUVs by a modification of the method of Brunner & Schenck (2019). Briefly, 800 μ l LUVs were destabilized by adding 16 μ l 10% w/v Triton X-100 and incubating for 20 min with agitation. VDAC samples were supplemented to 1.05% (v/v) LDAO and agitated (600 rpm) on a thermomixer at room temperature or, in some instances at 37°C for VDAC1. The desired concentration of protein in a maximal volume of 100 μ l was added to the destabilized vesicles, and the volume made up to 1 ml with SEC buffer or crosslinking-buffer containing 0.05% LDAO as indicated, with additional LDAO to ensure equal detergent concentrations in all samples. Samples were incubated for 20 min with end-over-end mixing at room temperature (VDAC2) or in some instances agitated (600 rpm) at 37°C in the case of VDAC1. Washed Bio-Beads (140 mg) were added, and the samples were agitated (600 rpm) for 20 min at 37°C in the case of VDAC1, then transferred to the cold-room for overnight incubation at 4°C with end-over-end mixing. Reconstituted vesicles were separated from Bio-Beads and used immediately for further assays. Protein concentration was determined by SDS-PAGE (Coomassie or Fluorescence Protein gel stain (Lamda Biotech)) in comparison with standards. Lipid concentration was determined by colorimetric assay of lipid phosphorus after the protocol of Rouser *et al* (1970). The protein/phospholipid ratio (PPR) of the samples was calculated using experimentally determined values. PPR* (Ploier *et al.*, 2016; Verchere *et al.*, 2017) describing the protein per phospholipid ratio normalized against the fraction of vesicles that cannot be populated by proteins was calculated as $PPR^* = \frac{PPR}{(1-R)}$ with $R = 2 * F_{Dith}^{Min}$ as clarified in the following section.

Fluorescence assays for channel and scramblase activity.

NBD-PC-containing proteoliposomes were assayed for VDAC channel activity with the dithionite assay after the protocol of Menon *et al.* (2011) and scramblase activity with the BSA-back extraction assay after the protocol of Kubelt *et al.* (2002). Proteoliposomes were diluted 40-times into HEPES buffer (50 mM HEPES pH 7.4, 100 mM NaCl) in a fluorimeter cuvette, and fluorescence was monitored under constant stirring (900 rpm) at 20°C in a temperature-controlled Spectrofluorometer FluoroMax + instrument using λ_{ex} =470 nm, λ_{em} =530 nm, slit widths 2.5 nm. The sample was equilibrated for 5-10 min before proceeding with the assays. For the

dithionite assay, 40 μ l of 1 M sodium dithionite, freshly prepared in unbuffered 0.5 M Tris, was added. For the BSA-back extraction assay, 40 μ l 75 mg/ml fatty acid-free BSA (Calbiochem) in HEPES buffer was added. Collected fluorescence traces were normalized to the initial value F_{\max} . The fraction (f) of vesicles containing either channel or scramblase activity was calculated as $f = \frac{F^L - F^P}{F^L - F^{\min}}$, with the normalized fluorescence (F) at 350 sec after dithionite (f_{pore}) or BSA (f_{scr}) addition for protein-free (F^L) or protein-containing liposomes (F^P) and F^{\min} as the lowest fluorescence signal detectable if all vesicles possess activity while accounting for refractory liposomes. For the dithionite assay, F_{Dith}^{\min} was experimentally determined using the average fluorescence value of proteoliposomes reconstituted at a protein:phospholipid ratio 1:6,000, using 15 μ g/ml of protein to obtain a theoretical copy number of 30 proteins per vesicle. If the dithionite assay captures all liposomes containing VDAC proteins, the fluorescence signal describing F_{BSA}^{\min} for all proteoliposomes theoretically capable of scramblase activity can be calculated as:

$$F_{\text{BSA}}^{\min} = ((1 - F_{\text{Dith}}^{\min} * 2) * \phi) + (F_{\text{Dith}}^{\min} * \phi) + F_{\text{Dith}}^{\min}$$

where ϕ is the fluorescence of NBD-PC when it is complexed with BSA compared with its value in the membrane. The value of ϕ was experimentally determined to be 0.4 using the fluorescence signal after 350 sec for protein-free liposomes and comparing dithionite and BSA-back extraction traces with rearrangement of the above equation as follows:

$$\phi = \frac{F_{\text{BSA}}^L - F_{\text{Dit}}^L}{((1 - 2 * F_{\text{Dith}}^L) + F_{\text{Dith}}^L)} = 0.4$$

The fraction of VDAC proteoliposomes that possess scramblase activity (Q) was determined as $Q = \frac{f_{\text{scr}}}{f_{\text{pore}}}$. Additionally, fluorescence traces were analyzed using one-phase or two-phase exponential decay functions as determined using F-test with GraphPad Prism.

To determine oligomeric species that are reconstituted per vesicle, f_{pore} was plotted against PPR*, and analyzed according to the Poisson model (Goren *et al.*, 2014; Menon *et al.*, 2011; Ploier *et al.*, 2016) by fitting to a one-phase exponential function. The fit constant τ (in units of μ g protein/ μ mol lipids) corresponds to the reconstitution condition where each vesicle has a single functional unit on average. Thus, the protein copy (C) number per 1 μ mol lipids was determined as $C = \tau * \frac{1}{\text{MW}} * N_A$, with MW = 31 kDa, and the number of vesicles (V) in 1 μ mol

was determined as $V = \frac{N_A * 10^{-6}}{L}$ with lipids per vesicle (L) as $L = \frac{(4\pi * (\frac{d}{2})^2 + 4\pi * (\frac{d}{2} - h)^2)}{a}$ with diameter (d) as 150 nm, bilayer thickness (h) as 5 nm and phospholipid headgroup area (a) as 0.71 nm² (Mimms *et al.*, 1981). Proteins incorporated per vesicles are thus, $\frac{C}{V}$.

Scramblase assay using PI-PLC.

VDAC proteoliposomes were prepared as described above except that 20,000 cpm per ml of [³H]inositol-labeled phosphatidylinositol (PI) (American Radiolabeled Chemicals) was included during the reconstitution step. Lipids were dried under nitrogen stream and taken up in 50 μ l per

1 ml sample preparation in reconstitution buffer (10 mM MOPS/Tris pH 7.0, 100 mM KCl) containing 0.2% w/v Triton X-100. No NBD-PC was present in the samples. Scramblase activity was assayed using PI-specific phospholipase C (PI-PLC purchased from Sigma) as described by Wang *et al.* (2018). To 100 μ l aliquots of proteoliposomes 10 μ l HEPES buffer (50 mM HEPES pH 7.4, 100 mM NaCl) (or 10 μ l 10% w/v Triton X-100 for sample disruption) was added. Finally, they were treated with 1/40 volume of PI-PLC working solution (10-times dilution into HEPES buffer) for different periods of time at 25°C. The reactions were stopped by the addition of ice-cold trichloroacetic acid to a final concentration of 12% (w/v) and placement on ice. Cytochrome c (Sigma) was added to a final concentration of 150 μ g/ml for better visualization of the protein pellet. Samples were incubated on ice for 30 min before being microfuged to pellet precipitated material (including non-hydrolyzed [3 H]PI). The supernatant containing released [3 H]inositol-cyclic phosphate was taken for liquid scintillation counting. Control samples included proteoliposomes disrupted with Triton X-100, proteoliposomes not treated with PI-PLC and protein-free liposomes. Scintillation counts were offset-corrected using the value obtained from non-treated samples, and data were subsequently normalized to the maximum extent of hydrolysis determined from Triton X-100 disrupted samples. The percentage of hydrolyzed PI was graphed as a function of time and traces were fitted to a one-phase exponential association using GraphPad Prism software.

EGS crosslinking.

Ethylene glycol bis(succinimidyl succinate) (EGS; ThermoFisher) crosslinking was done according to the protocol of Zalk *et al.* (2005) and Kim *et al.* (2019) with modifications. VDAC was crosslinked in LDAO or after reconstitution into LUVs prepared in crosslinking buffer (10 mM MOPS/KOH pH 7.0, 100 mM KCl). Samples to be crosslinked in LDAO were applied to a desalting spin column for buffer exchange to crosslinking buffer containing 0.05% LDAO (additional LDAO was added as indicated). EGS was dissolved in DMSO, diluted 20-times in crosslinking buffer (with additional LDAO as indicated), before being diluted 10-times into VDAC samples (15 μ g/ml or 150 μ g/ml when used for further reconstitution) to achieve a final 16x mole excess over reactive sites (lysine residues + N-terminus), with final DMSO = 0.5%. Non-crosslinked controls were treated with an equivalent amount of DMSO. After agitation (1000 rpm) in a thermomixer at 20°C for 40 min, the reaction was stopped by adding 5xLaemmli buffer to 1x (60 mM Tris-HCl pH 6.8, 2% w/v SDS, 10% v/v glycerol, 5% v/v β -mercaptoethanol, 0.01% bromophenol blue) and heating to 95°C for 3 min. Samples were taken for immunoblotting using anti His-tag antibody (ThermoFisher). Crosslinking and control reactions intended for scramblase activity assays were stopped with 1 M Tris pH 7 to a final concentration of 10 mM, instead of Laemmli buffer, and used for reconstitution into LUVs whilst ensuring the same DMSO and EGS concentrations in control protein-free liposomes. The same protocol was followed for crosslinking with dithiobis(succinimidyl propionate) (DSP; ThermoFisher) and samples were taken up in 4xSDS-PAGE buffer to 1x (75 mM Tris-HCl pH 6.8, 2% w/v SDS, 10% glycerol, 0.05% bromophenol blue) instead of Laemmli buffer for immunoblotting. For cleavage of DSP, 50 mM DTT (from 1 M stock prepared in reconstitution buffer) was added and the samples were incubated for 30 min at 37°C while agitated (600 rpm). Control and non-cleaved samples were treated equally omitting DTT.

Assay of slow scrambling.

To assay slow scrambling, proteoliposomes were prepared without NBD-PC and were subsequently asymmetrically labeled by adding NBD-PC (from an EtOH stock solution) to a final concentration of 0.25 mol% lipid, and final EtOH concentration of 0.4 %. After 30 min incubation on ice, the samples were shifted to 20°C and 50 µl aliquots were taken for BSA-back extraction assay as described above every 2 h for up to 30 h. Normalized fluorescence signal 100 sec after BSA addition was used to determine the fraction of NBD-analogs at the outer leaflet setting time point 0 as 100% of analogs present at the outer leaflet. Traces were fitted to a one-phase exponential decay function using GraphPad Prism software.

Isolation of yeast mitochondria.

Crude mitochondria were prepared from yeast after the protocol of Daum *et al* (1981) with some modifications. Briefly, yeast cells were pre-grown in YPD (1% w/v yeast extract, 2% w/v peptone, 2% w/v dextrose) over a weekend, 10-times diluted into rich-lactate (RL) media (1% w/v yeast extract, 2% w/v peptone, 0.05% w/v dextrose, 2% v/v lactic acid, 3.4 mM CaCl₂ 2H₂O, 8.5 mM NaCl, 2.95 mM MgCl₂ 6H₂O, 7.35 mM KH₂PO₄, 18.7 mM NH₄Cl, pH 5.5 adjusted with KOH) containing 2 mM ethanolamine and grown for 4h at 30°C. 6 OD₆₀₀ were added to 500 ml RL containing 2 mM ethanolamine and further grown overnight to an OD₆₀₀ of 4 to 5. Cells were harvested, washed with 1 mM EDTA and taken up in T-buffer (0.1 M Tris-HCl, 10 mM DTT, pH 9.4) at 0.5 g cell pellet/ml. Cells were incubated for 10 min at 30°C, harvested and washed in 30°C warm S-buffer (1.2 M sorbitol, 20 mM KH₂PO₄, pH 7.4 with KOH) at 0.15 g cell pellet/ml. Cells were collected and taken up in S-buffer containing 1 mg Zymolyase per g yeast cell pellet at 0.3 g cell pellet/ml and incubated at 30°C for 1h. Spheroplasts were diluted in equal volume of ice-cold S-buffer and harvested at 4°C. Cells are resuspended in ice-cold D⁻-buffer (10 mM MES, 0.6 M sorbitol, pH 6 with NaOH) containing 0.2% w/v fatty acid free BSA and 1 mM PMSF at 0.75 g cell pellet/ml, and disrupted with 10 strokes of a dounce homogenizer. Mitochondria were collected by differential centrifugation, first, intact cells were removed by two slow spins at 1,400 x g 4°C for 5 min. Crude mitochondria were pelleted by centrifugation at 10,000 x g 12 min 4°C. Mitochondria were taken up in D⁻-buffer, snap frozen and stored at -80°C. The protein concentration of the preparations was determined using BCA protein assay in the presence of 0.8% SDS and Coomassie staining.

Transport-coupled PS decarboxylation by yeast mitochondria.

Conversion of NBD-PS to NBD-PE by crude mitochondria was assayed according to the protocol of Miyata *et al* (2016) with modifications. Crude mitochondria were diluted to 5 mg/ml in D⁻-buffer. The assay was started by 5-times dilution of the mitochondria into D⁺-buffer (10 mM MES, 0.6 M sorbitol, pH 6 with NaOH, 12.5 mM EDTA) containing 1.25 µM 16:0 – 6:0 NBD-PS (Avanti) and 0.3 µM C6-NBD-PC. Samples were incubated at 30°C and 100 µl aliquots were taken at the respective time points. To stop the reaction, 750 µl CHCl₃:MeOH v:v 1:2 was added. Subsequently, 250 µl CHCl₃ and 250 µl 0.2 M KCl were added, and the samples were centrifuged at 2,200 x g for 5 min at RT. The CHCl₃ phase was collected and dried under nitrogen. Samples were taken up in MeOH and spotted onto an activated thin layer chromatography (TLC) silica gel 60 plate (Merck). The plate was developed in CHCl₃:MeOH:Acetic Acid:H₂O (25:15:4:2,

by volume) and NBD fluorescence was visualized using the BioRad ChemiDoc System. Densitometry of the fluorescence (F) signal was used to determine PS to PE conversion according to the equation $\frac{F_{PE}}{F_{PE} + F_{PS}}$. Psd1 and Tom70 levels were determined by immunoblot to establish similar content of these OMM and IMM markers in the assay samples.

Coarse grained molecular dynamics simulations (CGMD).

CGMD - General settings and systems preparation.

All molecular dynamics simulations were performed using the simulation package Gromacs version 5.1.4 (Abraham *et al*, 2015) with plugin Plumed version 2.3 (Tribello *et al*, 2014). We employed i) coarse-grained Martini force-field version 3.0 (Souza *et al*, 2021) and ii) coarse-grained Martini force-field version 2.2 (de Jong *et al*, 2013; Marrink *et al*, 2007; Monticelli *et al*, 2008) with Elnedyn network (Periole *et al*, 2009). For Martini 2 simulations of VDAC dimers, the Lennard-Jones interactions between protein beads were scaled down by 10% to prevent unrealistically strong protein-protein interactions (Javanainen *et al*, 2017). We used a structure of human VDAC1 (hVDAC1) dimer solved by x-ray diffraction with a resolution of 2.7 Å and denoted as 6g6u in the RCSB Protein Data Bank (www.rcsb.org/structure/6g6u). Single monomer of VDAC1 was minimized in vacuum using atomistic Amber 99SB-ILDN force-field (Lindorff-Larsen *et al*, 2010) with steepest descent algorithm until the maximum force on any atom was lower than 100 kJ mol⁻¹ nm⁻¹.

For Martini 3 simulations, the minimized atomistic protein structure was coarse-grained using Martinize2 script (github.com/marrink-lab/vermouth-martinize) constructing an elastic network with a force constant of 1000 kJ mol⁻¹ nm⁻² to fix the tertiary structure of VDAC1. Dimeric structures were generated by placing the coarse-grained VDAC1 monomers at appropriate positions. No restraints maintaining the dimeric structure of VDAC1 were used, unless explicitly stated otherwise. Each VDAC1 structure was embedded in the center of an xy-positioned membrane using the Insane script (github.com/Tsjerk/Insane). For simulations with monomeric VDAC1, the membrane was composed of roughly 660 molecules of 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC), while for the simulations of dimeric interfaces, the membrane consisted of roughly 920 molecules of POPC. Systems had an approximate size of roughly 15 × 15 × 12 nm for simulations of monomeric VDAC1 and 18 × 18 × 11 nm for dimer simulations and were solvated with roughly 16,000 and 20,000 water beads, respectively. KCl ions were added to each system at a concentration of 100 mM with an excess of ions to neutralize the system. K⁺ ions were modelled as SQ5 particles with charge of +1 and mass of 39 a.u. Each constructed system was minimized using the steepest descent algorithm and a maximum force tolerance of 200 kJ mol⁻¹ nm⁻¹.

In Martini 2 simulations with Elnedyn, the VDAC1 monomer and dimer structures were coarse-grained and embedded into the membrane using the CHARMM-GUI web interface (Jo *et al*, 2008). The systems with monomeric and dimeric VDAC1 had similar sizes and the same molecular composition as the Martini 3 systems. In case of Martini 2, the solvated K⁺ ions were modelled as Qd type particles with charge +1 (the same as Na⁺ ions). The dimeric structures of VDAC1 were not fixed by any external potentials and all Martini 2 systems were minimized the same way as Martini 3 systems.

CGMD - Equilibration and production simulations.

For both Martini 3 and Martini 2 simulations, the equilibration was performed in five stages with increasing simulation time-step (dt) and length (t): I) dt = 2 fs, t = 0.5 ns; II) dt = 5 fs, t = 1.25 ns; III) dt = 10 fs, t = 1 ns; IV) dt = 20 fs, t = 30 ns; V) dt = 20 fs, t = 15 ns. In all, but the stage V, position restraints in the form of harmonic potential with a force constant of 1000 kJ mol⁻¹ nm⁻² were applied to all coordinates of protein backbone beads. For Martini 3 systems with VDAC1 dimers, water restraints were applied in stages I-III to keep water from entering the hydrophobic core of the membrane. For this purpose, inverted flat-bottom potential with a force constant (k) of 10 kJ mol⁻¹ nm⁻² and a reference distance (ref) of 2.2 nm from the membrane center of mass was applied to z-coordinates of all water beads.

Equilibration was performed in NPT ensemble with temperature being maintained at 300 K using stochastic velocity-rescaling thermostat (Bussi *et al*, 2007) with a coupling constant of 1 ps. Water with ions, membrane, and protein(s) were coupled to three separate thermal baths. Pressure was kept at 1 bar using Berendsen barostat (Berendsen *et al*, 1984) with a coupling constant of 12 ps. Simulation box size was scaled semi-isotropically with a compressibility of 3 × 10⁻⁴ bar⁻¹ in both xy-plane and z-direction.

To integrate the Newton equations of motion, we used leap-frog algorithm. Non-bonded interactions were cut-off at 1.1 nm. Van der Waals potential was shifted to zero at cut-off distance. Reaction field was used to treat electrostatic interactions with a relative dielectric constant of 15. For Martini 3 simulations, LINCS (Hess *et al*, 1997) parameters lincs-order and lincs-iter were set to 8 and 2, respectively, to avoid artificial temperature gradients (Thallmair *et al*, 2021). Following equilibration, monomeric and several dimeric VDAC1 structures were simulated in Martini 3 for 10 μs using the same simulation settings as in stage V of the equilibration, except the Berendsen barostat was replaced with Parrinello-Rahman barostat (Parrinello & Rahman, 1981). The same simulation settings were also used for other simulations described in the section Free energy calculations. Three independent replica simulations were performed for each of the systems with monomeric or dimeric VDAC1. Apart from unrestrained simulations, we simulated the fastest scrambling dimer, dimer-1, restrained to its initial structure using harmonic potential (k = 1000 kJ mol⁻¹ nm⁻²) in the x- and y-coordinates of backbone beads of the glutamate 73 and serine 193 of both VDAC1 protomers.

CGMD - Analysis.

For each of the three independent molecular dynamics simulations, we calculated the percentage of scrambled lipids in time, analyzing a simulation snapshot every 10 ns. Lipid was identified as “scrambled” if it was positioned in a different membrane leaflet than at the start of the simulation. Lipids were assigned to the leaflet by the position of their phosphate bead with respect to the membrane center. For further analysis, all three independent replicas were concatenated and centered on protein beads using its root-mean-square deviation. We constructed a water defect map describing membrane disruption around the protein structure. Water defect was defined as the average number of water beads located closer to the membrane center on z axis than 1 nm. The average number was calculated using xy-mesh with 0.1 nm bin every 100 ps. The same mesh was employed to calculate the average membrane

thickness using z-positions of lipid phosphates. The average membrane thickness was defined as the difference between the position of the upper- and lower-leaflet phosphates. The bins with less than 100 samples were excluded from the analysis. The code used for the analysis of lipid scrambling and membrane disruption is available from <https://github.com/Ladme/scramblyzer> and <https://github.com/Ladme/memidian>, respectively. The position of the beta-strands was approximated by the average position of backbone bead of the central residue in the xy-plane.

CGMD - Free energy calculations.

We calculated free energy of lipid flip-flop in several systems: a) in protein-free POPC membrane (Martini 3 & Martini 2), b) in system with VDAC1 monomer (Martini 3 & Martini 2), c) in system with VDAC1 dimer-1 (Martini 3), d) in system with VDAC1 monomer while turning off the attractive Lennard-Jones (LJ) interactions between the scrambling lipid and VDAC (Martini 3), and e) in system with VDAC1 dimer-1 while turning off the attractive Lennard-Jones interactions between the scrambling lipid and VDAC (Martini 3).

To enhance the sampling of the lipid flip-flop, we employed umbrella sampling method (Torrie & Valleau, 1977). For free energy calculations in the presence of VDAC1 monomer, Hamiltonian replica exchange was further applied (Fukunishi *et al*, 2002). Three one-dimensional collective variables (CVs) were used to capture the lipid flip-flop in different systems. For simulations without the presence of VDAC1, the oriented z-distance between a chosen lipid phosphate bead and local membrane center of mass was used as a CV. Local membrane center of mass was calculated from the positions of lipid beads localized inside a cylinder with a radius of 2.0 nm and its principal axis going through the selected phosphate bead. For simulations of systems with VDAC1 monomer, the CV was defined as the oriented z-distance between a chosen lipid phosphate bead and backbone bead of glutamate 73. In systems with VDAC1 dimer-1, the CV was defined as the oriented z-distance between a chosen lipid phosphate bead and the center of mass of the dimer-1 which was positioned close to the center of the membrane on the z-axis. During the pulling and umbrella sampling simulations, the dimer-1 was restrained in the same way as described in the section Equilibration and production simulations.

Initial configurations for umbrella sampling were generated by pulling the chosen lipid phosphate bead through the membrane for 1 μ s with a pulling rate of 4.2 nm μ s⁻¹ and initial reference distance of 2.1 (for systems without VDAC1) or \pm 2.0 (for systems with VDAC1) nm using a harmonic potential. In all systems, one pulling simulation was run for each direction of pulling (from the upper to the lower membrane leaflet and in the opposite direction). The exception was Martini 3 systems with VDAC1 monomer, where two pulling simulations in each direction were performed to generate a larger ensemble of initial configurations for umbrella sampling with Hamiltonian replica exchange. In monomeric VDAC1 systems, the pulled phosphate was restrained by an xy-plane flat-bottom potential ($k = 500$ kJ mol⁻¹ nm⁻², ref = 1.5 nm) to stay close the backbone bead of the glutamate 73 residue. In dimeric VDAC1 systems, the phosphate was restrained by an xy-plane flat-bottom potential ($k = 500$ kJ mol⁻¹ nm⁻², ref = 2.5 nm) to stay close to the dimeric interface.

For systems with VDAC1, two independent sets of umbrella sampling windows were prepared to observe any possible hysteresis of the flip-flop process. For Martini 2 simulations, the initial configurations for each set of umbrella sampling windows were obtained from the pulling simulation performed in the specific direction and the calculation was enhanced by

applying Hamiltonian replica exchange in these windows. In Martini 3 simulations with VDAC1 monomer, the efficiency of the Hamiltonian replica exchange was further enhanced by using configurations from two pulling simulations performed in the same direction. For systems without VDAC1, only one pulling direction was used.

We used between 15 (Martini 3 simulations of dimer-1) and 72 windows (Martini 2 simulations of monomeric VDAC) distributed along the range of the CV for each pulling direction. See Tables 1-5 for the complete list of umbrella sampling windows that were used for the calculations. After a short 30 ns equilibration, each window was sampled for a specific time which differed based on the system sampled and force-field used. The simulated times necessary to obtain converged free energy profiles were a) 1 μ s for protein-free membrane, b) 4 or 8 μ s for simulations with VDAC1 monomer in Martini 3 and Martini 2, respectively, c) 6 μ s for simulations with VDAC1 dimer, d) 2 μ s for simulations with VDAC1 monomer and attractive LJ interactions turned off, and e) 4 μ s for simulations with VDAC1 dimer and attractive LJ interactions turned off. In simulations in which Hamiltonian replica exchange was applied, the exchange was attempted every 10,000 integration steps (200 ps). Free energy profiles were obtained from the umbrella sampling windows using the weighted histogram analysis method (Kumar *et al*, 1992; Souaille & Roux, 2001) as implemented in the Gromacs tool g_wham (Hub *et al*, 2010).

ACKNOWLEDGEMENTS

We thank Steve Claypool (Johns Hopkins University) for antibodies, Non Miyata (Kyushu University) for yeast strains, and Toon de Kroon (Utrecht University) for the mitochondria isolation protocol. This work was supported by National Institutes of Health grant 1R21NS119779 (AKM), Boehringer Ingelheim Fonds Ph.D. Fellowship (HJ), Deutsche Forschungsgemeinschaft grants SFB944-P14 and HO 3539/2-1 (JH), Czech Science Foundation grant GA20-20152S (RV), European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 101001470) (RV), and National Institute of virology and bacteriology (Programme EXCELES, ID Project No. LX22NPO5103) - Funded by the European Union - Next Generation EU (RV). Computational resources were provided by the CESNET LM2015042 and the CERIT Scientific Cloud LM2015085 under the program Projects of Large Research, Development, and Innovations Infrastructures. Additional computational resources were obtained from IT4 Innovations National Supercomputing Center -- LM2015070 project supported by MEYS CR from the Large Infrastructures for Research, Experimental Development, and Innovations.

DISCLOSURE AND COMPETING INTERESTS.

The authors declare that they have no conflict of interest.

AUTHOR CONTRIBUTIONS

Conceptualization: HJ, JH, RV, AKM

Methodology: HJ, LB, RV, AKM

Investigation: HJ, LB

Visualization: HJ, LB, RV, AKM

Funding acquisition: HJ, JH, RV, AKM

Project administration: RV, AKM

Supervision: RV, AKM

Writing – original draft: HJ, LB, AKM

Writing – review & editing: HJ, LB, JH, RV, AKM

DATA AND MATERIALS AVAILABILITY

All data are available in the main text or the supplementary materials.

REFERENCES

- Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, Lindahl E (2015) GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* 1: 19-25
- Adlakha J, Hong Z, Li P, Reinisch KM (2022) Structural and biochemical insights into lipid transport by VPS13 proteins. *J Cell Biol* 221: e202202030
- Araiso Y, Tsutsumi A, Qiu J, Imai K, Shiota T, Song J, Lindau C, Wenz LS, Sakaue H, Yunoki K et al (2019) Structure of the mitochondrial import gate reveals distinct preprotein paths. *Nature* 575: 395-401
- Bayrhuber M, Meins T, Habeck M, Becker S, Giller K, Villinger S, Vonrhein C, Griesinger C, Zweckstetter M, Zeth K (2008) Structure of the human voltage-dependent anion channel. *Proc Natl Acad Sci U S A* 105: 15370-15375
- Bennett WF, Tieleman DP (2014) The importance of membrane defects-lessons from simulations. *Acc Chem Res* 47: 2244-2251
- Berendsen HJC, Postma JPM, Gunsteren WFv, DiNola A, Haak JR (1984) Molecular dynamics with coupling to an external bath. *The Journal of Chemical Physics* 81: 3684-3690
- Bergdoll LA, Lerch MT, Patrick JW, Belardo K, Altenbach C, Bisignano P, Laganowsky A, Grabe M, Hubbell WL, Abramson J (2018) Protonation state of glutamate 73 regulates the formation of a specific dimeric association of mVDAC1. *Proc Natl Acad Sci U S A* 115: E172-E179
- Bergdoll LG, M.; Abramson, J. (2017) An Assessment of How VDAC Structures Have Impacted Our Understanding of Their Function. In: *Molecular Basis for Mitochondrial Signaling*, Rostovtseva T.K. (ed.) pp. 141-160. Springer International Publishing AG
- Betaneli V, Petrov EP, Schwille P (2012) The role of lipids in VDAC oligomerization. *Biophys J* 102: 523-531
- Broeskamp F, Edrich ESM, Knittelfelder O, Neuhaus L, Meyer T, Heyden J, Habernig L, Kreppel F, Gourlay CW, Rockenfeller P (2021) Porin 1 Modulates Autophagy in Yeast. *Cells* 10
- Brunner JD, Schenck S (2019) Preparation of Proteoliposomes with Purified TMEM16 Protein for Accurate Measures of Lipid Scramblase Activity. *Methods Mol Biol* 1949: 181-199
- Bussi G, Donadio D, Parrinello M (2007) Canonical sampling through velocity rescaling. *J Chem Phys* 126: 014101
- Chadda R, Bernhardt N, Kelley EG, Teixeira SC, Griffith K, Gil-Ley A, Ozturk TN, Hughes LE, Forsythe A, Krishnamani V et al (2021) Membrane transporter dimerization driven by differential lipid solvation energetics of dissociated and associated states. *Elife* 10
- Chang QL, Gummadi SN, Menon AK (2004) Chemical modification identifies two populations of glycerophospholipid flippase in rat liver ER. *Biochemistry* 43: 10710-10718
- Cliff L, Chadda R, Robertson JL (2020) Occupancy distributions of membrane proteins in heterogeneous liposome populations. *Biochim Biophys Acta Biomembr* 1862: 183033
- Dadsena S, Bockelmann S, Mina JGM, Hassan DG, Korneev S, Razzera G, Jahn H, Niekamp P, Muller D, Schneider M et al (2019) Ceramides bind VDAC2 to trigger mitochondrial apoptosis. *Nat Commun* 10: 1832
- Daum G, Böhmi PC, Schatz G (1981) Import of Proteins into Mitochondria. *J Biol Chem* 257: 13028-13033

- de Jong DH, Singh G, Bennett WF, Arnarez C, Wassenaar TA, Schafer LV, Periole X, Tieleman DP, Marrink SJ (2013) Improved Parameters for the Martini Coarse-Grained Protein Force Field. *J Chem Theory Comput* 9: 687-697
- De Pinto V (2021) Renaissance of VDAC: New Insights on a Protein Family at the Interface between Mitochondria and Cytosol. *Biomolecules* 11
- Di Rosa MC, Guarino F, Conti Nibali S, Magri A, De Pinto V (2021) Voltage-Dependent Anion Selective Channel Isoforms in Yeast: Expression, Structure, and Functions. *Front Physiol* 12: 675708
- Dolis D, de Kroon AI, de Kruijff B (1996) Transmembrane movement of phosphatidylcholine in mitochondrial outer membrane vesicles. *J Biol Chem* 271: 11879-11883
- Egea PF (2021) Mechanisms of Non-Vesicular Exchange of Lipids at Membrane Contact Sites: Of Shuttles, Tunnels and, Funnels. *Front Cell Dev Biol* 9: 784367
- Fukunishi H, Watanabe O, Takada S (2002) On the Hamiltonian replica exchange method for efficient sampling of biomolecular systems: Application to protein structure prediction. *The Journal of Chemical Physics* 116: 9058-9067
- Geula S, Naveed H, Liang J, Shoshan-Barmatz V (2012) Structure-based analysis of VDAC1 protein: defining oligomer contact sites. *J Biol Chem* 287: 2179-2190
- Ghanbarpour A, Valverde DP, Melia TJ, Reinisch KM (2021) A model for a partnership of lipid transfer proteins and scramblases in membrane expansion and organelle biogenesis. *Proc Natl Acad Sci U S A* 118
- Goncalves RP, Buzhysnsky N, Scheuring S (2008) Mini review on the structure and supramolecular assembly of VDAC. *J Bioenerg Biomembr* 40: 133-138
- Goren MA, Morizumi T, Menon I, Joseph JS, Dittman JS, Cherezov V, Stevens RC, Ernst OP, Menon AK (2014) Constitutive phospholipid scramblase activity of a G protein-coupled receptor. *Nat Commun* 5: 5115
- Hess B, Bekker H, Berendsen HJC, Fraaije JGEM (1997) LINCS: A linear constraint solver for molecular simulations. *Journal of Computational Chemistry* 18: 1463-1472
- Hoogenboom BW, Suda K, Engel A, Fotiadis D (2007) The supramolecular assemblies of voltage-dependent anion channels in the native membrane. *J Mol Biol* 370: 246-255
- Horvath SE, Daum G (2013) Lipids of mitochondria. *Prog Lipid Res* 52: 590-614
- Hub JS, de Groot BL, van der Spoel D (2010) g_wham—A Free Weighted Histogram Analysis Implementation Including Robust Error and Autocorrelation Estimates. *Journal of Chemical Theory and Computation* 6: 3713-3720
- Janssen MJ, Koorengel MC, de Kruijff B, de Kroon AI (1999) Transbilayer movement of phosphatidylcholine in the mitochondrial outer membrane of *Saccharomyces cerevisiae* is rapid and bidirectional. *Biochim Biophys Acta* 1421: 64-76
- Jasinska R, Zborowski J, Somerharju P (1993) Intramitochondrial distribution and transport of phosphatidylserine and its decarboxylation product, phosphatidylethanolamine. Application of pyrene-labeled species. *Biochim Biophys Acta* 1152: 161-170
- Javanainen M, Martinez-Seara H, Vattulainen I (2017) Excessive aggregation of membrane proteins in the Martini model. *PLoS One* 12: e0187936
- Jo S, Kim T, Iyer VG, Im W (2008) CHARMM-GUI: a web-based graphical user interface for CHARMM. *J Comput Chem* 29: 1859-1865

- Kalienkova V, Clerico Mosina V, Paulino C (2021) The Groovy TMEM16 Family: Molecular Mechanisms of Lipid Scrambling and Ion Conduction. *J Mol Biol* 433: 166941
- Khelashvili G, Menon AK (2022) Phospholipid Scrambling by G Protein-Coupled Receptors. *Annu Rev Biophys* 51: 39-61
- Kim J, Gupta R, Blanco LP, Yang S, Shteinfer-Kuzmine A, Wang K, Zhu J, Yoon HE, Wang X, Kerkhofs M et al (2019) VDAC oligomers form mitochondrial pores to release mtDNA fragments and promote lupus-like disease. *Science* 366: 1531-1536
- Kubelt J, Menon AK, Muller P, Herrmann A (2002) Transbilayer movement of fluorescent phospholipid analogues in the cytoplasmic membrane of *Escherichia coli*. *Biochemistry* 41: 5605-5612
- Kumar S, Rosenberg JM, Bouzida D, Swendsen RH, Kollman PA (1992) THE weighted histogram analysis method for free-energy calculations on biomolecules. I. The method. *Journal of Computational Chemistry* 13: 1011-1021
- Lampl M, Leber A, Paltauf F, Daum G (1994) Import of phosphatidylinositol and phosphatidylcholine into mitochondria of the yeast, *Saccharomyces cerevisiae*. *FEBS Lett* 356: 1-4
- Lindorff-Larsen K, Piana S, Palmo K, Maragakis P, Klepeis JL, Dror RO, Shaw DE (2010) Improved side-chain torsion potentials for the Amber ff99SB protein force field. *Proteins* 78: 1950-1958
- Maeda S, Yamamoto H, Kinch LN, Garza CM, Takahashi S, Otomo C, Grishin NV, Forli S, Mizushima N, Otomo T (2020) Structure, lipid scrambling activity and role in autophagosome formation of ATG9A. *Nat Struct Mol Biol* 27: 1194-1201
- Malvezzi M, Andra KK, Pandey K, Lee BC, Falzone ME, Brown A, Iqbal R, Menon AK, Accardi A (2018) Out-of-the-groove transport of lipids by TMEM16 and GPCR scramblases. *Proc Natl Acad Sci U S A* 115: E7033-E7042
- Marrink SJ, Risselada HJ, Yefimov S, Tieleman DP, de Vries AH (2007) The MARTINI force field: coarse grained model for biomolecular simulations. *J Phys Chem B* 111: 7812-7824
- Matoba K, Kotani T, Tsutsumi A, Tsuji T, Mori T, Noshiro D, Sugita Y, Nomura N, Iwata S, Ohsumi Y et al (2020) Atg9 is a lipid scramblase that mediates autophagosomal membrane expansion. *Nat Struct Mol Biol* 27: 1185-1193
- Menon I, Huber T, Sanyal S, Banerjee S, Barre P, Canis S, Warren JD, Hwa J, Sakmar TP, Menon AK (2011) Opsin is a phospholipid flippase. *Curr Biol* 21: 149-153
- Messina A, Reina S, Guarino F, De Pinto V (2012) VDAC isoforms in mammals. *Biochim Biophys Acta* 1818: 1466-1476
- Mimms LT, Zampighi G, Nozaki Y, Tanford C, Reynolds JA (1981) Phospholipid vesicle formation and transmembrane protein incorporation using octyl glucoside. *Biochemistry* 20: 833-840
- Miyata N, Fujii S, Kuge O (2018) Porin proteins have critical functions in mitochondrial phospholipid metabolism in yeast. *J Biol Chem* 293: 17593-17605
- Miyata N, Watanabe Y, Tamura Y, Endo T, Kuge O (2016) Phosphatidylserine transport by Ups2-Mdm35 in respiration-active mitochondria. *J Cell Biol* 214: 77-88
- Monticelli L, Kandasamy SK, Periole X, Larson RG, Tieleman DP, Marrink SJ (2008) The MARTINI Coarse-Grained Force Field: Extension to Proteins. *J Chem Theory Comput* 4: 819-834
- Parrinello M, Rahman A (1981) Polymorphic transitions in single crystals: A new molecular dynamics method. *Journal of Applied Physics* 52: 7182-7190

- Periole X, Cavalli M, Marrink SJ, Ceruso MA (2009) Combining an Elastic Network With a Coarse-Grained Molecular Force Field: Structure, Dynamics, and Intermolecular Recognition. *J Chem Theory Comput* 5: 2531-2543
- Ploier B, Caro LN, Morizumi T, Pandey K, Pearing JN, Goren MA, Finnemann SC, Graumann J, Arshavsky VY, Dittman JS et al (2016) Dimerization deficiency of enigmatic retinitis pigmentosa-linked rhodopsin mutants. *Nat Commun* 7: 12832
- Pomorski T, Menon AK (2006) Lipid flippases and their biological functions. *Cell Mol Life Sci* 63: 2908-2921
- Pomorski TG, Menon AK (2016) Lipid somersaults: Uncovering the mechanisms of protein-mediated lipid flipping. *Prog Lipid Res* 64: 69-84
- Prilipov A, Phale PS, Van Gelder P, Rosenbusch JP, Koebnik R (1998) Coupling site-directed mutagenesis with high-level expression: large scale production of mutant porins from *E. coli*. *FEMS Microbiol Lett* 163: 65-72
- Raghavan A, Sheiko T, Graham BH, Craigen WJ (2012) Voltage-dependant anion channels: novel insights into isoform function through genetic models. *Biochim Biophys Acta* 1818: 1477-1485
- Reinisch KM, Prinz WA (2021) Mechanisms of nonvesicular lipid transport. *J Cell Biol* 220
- Rouser G, Fkeischer S, Yamamoto A (1970) Two dimensional thin layer chromatographic separation of polar lipids and determination of phospholipids by phosphorus analysis of spots. *Lipids* 5: 494-496
- Schlame M, Greenberg ML (2017) Biosynthesis, remodeling and turnover of mitochondrial cardiolipin. *Biochim Biophys Acta Mol Cell Biol Lipids* 1862: 3-7
- Schredelseker J, Paz A, Lopez CJ, Altenbach C, Leung CS, Drexler MK, Chen JN, Hubbell WL, Abramson J (2014) High resolution structure and double electron-electron resonance of the zebrafish voltage-dependent anion channel 2 reveal an oligomeric population. *J Biol Chem* 289: 12566-12577
- Shoshan-Barmatz V, Ben-Hail D, Admoni L, Krelin Y, Tripathi SS (2015) The mitochondrial voltage-dependent anion channel 1 in tumor cells. *Biochim Biophys Acta* 1848: 2547-2575
- Shoshan-Barmatz V, Mizrahi D, Keinan N (2013) Oligomerization of the mitochondrial protein VDAC1: from structure to function and cancer therapy. *Prog Mol Biol Transl Sci* 117: 303-334
- Souaille M, Roux Bt (2001) Extension to the weighted histogram analysis method: combining umbrella sampling with free energy calculations. *Computer Physics Communications* 135: 40-57
- Souza PCT, Alessandri R, Barnoud J, Thallmair S, Faustino I, Grunewald F, Patmanidis I, Abdizadeh H, Bruininks BMH, Wassenaar TA et al (2021) Martini 3: a general purpose force field for coarse-grained molecular dynamics. *Nat Methods* 18: 382-388
- Takeda H, Tsutsumi A, Nishizawa T, Lindau C, Busto JV, Wenz LS, Ellenrieder L, Imai K, Straub SP, Mossmann W et al (2021) Mitochondrial sorting and assembly machinery operates by beta-barrel switching. *Nature* 590: 163-169
- Tamura Y, Kawano S, Endo T (2020) Lipid homeostasis in mitochondria. *Biol Chem* 401: 821-833
- Tatsuta T, Langer T (2017) Intramitochondrial phospholipid trafficking. *Biochim Biophys Acta Mol Cell Biol Lipids* 1862: 81-89
- Thallmair S, Javanainen M, Fabian B, Martinez-Seara H, Marrink SJ (2021) Nonconverged Constraints Cause Artificial Temperature Gradients in Lipid Bilayer Simulations. *J Phys Chem B* 125: 9537-9546

- Torrie GM, Valleau JP (1977) Nonphysical sampling distributions in Monte Carlo free-energy estimation: Umbrella sampling. *Journal of Computational Physics* 23: 187-199
- Tribello GA, Bonomi M, Branduardi D, Camilloni C, Bussi G (2014) PLUMED 2: New feathers for an old bird. *Computer Physics Communications* 185: 604-613
- Tucker K, Park E (2019) Cryo-EM structure of the mitochondrial protein-import channel TOM complex at near-atomic resolution. *Nat Struct Mol Biol* 26: 1158-1166
- Verchere A, Ou WL, Ploier B, Morizumi T, Goren MA, Butikofer P, Ernst OP, Khelashvili G, Menon AK (2017) Light-independent phospholipid scramblase activity of bacteriorhodopsin from *Halobacterium salinarum*. *Sci Rep* 7: 9522
- Villinger S, Briones R, Giller K, Zachariae U, Lange A, de Groot BL, Griesinger C, Becker S, Zweckstetter M (2010) Functional dynamics in the voltage-dependent anion channel. *Proc Natl Acad Sci U S A* 107: 22546-22551
- Wang L, Iwasaki Y, Andra KK, Pandey K, Menon AK, Butikofer P (2018) Scrambling of natural and fluorescently tagged phosphatidylinositol by reconstituted G protein-coupled receptor and TMEM16 scramblases. *J Biol Chem* 293: 18318-18327
- Wang W, Chen X, Zhang L, Yi J, Ma Q, Yin J, Zhuo W, Gu J, Yang M (2020) Atomic structure of human TOM core complex. *Cell Discov* 6: 67
- Zalk R, Israelson A, Garty ES, Azoulay-Zohar H, Shoshan-Barmatz V (2005) Oligomeric states of the voltage-dependent anion channel and cytochrome c release from mitochondria. *Biochem J* 386: 73-83
- Zeth K (2010) Structure and evolution of mitochondrial outer membrane proteins of beta-barrel topology. *Biochim Biophys Acta* 1797: 1292-1299
- Zeth K, Zachariae U (2018) Ten Years of High Resolution Structural Research on the Voltage Dependent Anion Channel (VDAC)-Recent Developments and Future Directions. *Front Physiol* 9: 108

FIGURES (Figures 1-4, Supplementary Figures S1-S10, Movie S1)

Figures 1-4




Figure 1. VDAC2 is a phospholipid scramblase.

A. Schematic showing phospholipid transport from the ER to all bilayer leaflets of the mitochondrial double membrane. Non-vesicular mechanisms deliver ER-synthesized

phospholipids to the cytoplasmic face of the outer mitochondrial membrane (OMM). The lipids are scrambled across the OMM by a lipid transporter (scramblase) before moving through the intermembrane space (IMS) and across the inner mitochondrial membrane (IMM). **B.** Scramblase assay using [3 H]phosphatidylinositol ([3 H]PI). Protein-free-liposomes (L) or VDAC2-proteoliposomes (P) are reconstituted with [3 H]PI and probed with PI-specific phospholipase C (PI-PLC) which hydrolyzes [3 H]PI to diacylglycerol (DAG) and [3 H]inositol cyclic phosphate ([3 H]Inos-P). Only [3 H]PI molecules in the outer leaflet are hydrolyzed in protein-free-liposomes, whereas all [3 H]PI will be hydrolyzed in scramblase-containing vesicles where [3 H]PI from the inner leaflet is scrambled to the outer leaflet. **C.** Time-course of PI-PLC-mediated hydrolysis of [3 H]PI in protein-free-vesicles (liposome) versus vesicles reconstituted with VDAC2 at a theoretical protein/vesicle copy number of 30. Data points are shown as mean and 95% CI, $n = 4$, and fitted to a single exponential function. **D.** Channel assay using NBD-PC. Vesicles are reconstituted as in panel B, but with fluorescent NBD-PC instead of [3 H]PI. The membrane-impermeant reductant dithionite bleaches approximately 50% of the NBD-PC fluorescence in protein-free liposomes (L), corresponding to those molecules located in the outer leaflet. For vesicles containing a VDAC2 channel, all NBD-PC molecules are expected to be bleached as dithionite can enter the vesicles. **E.** Time-course of NBD-PC fluorescence (normalized to the starting value) in liposomes and VDAC2-vesicles on adding dithionite. **F.** Scramblase assay using NBD-PC. Vesicles are reconstituted as in panel D and treated with fatty acid-free BSA which extracts NBD-PC from the outer leaflet. In complex with BSA, NBD-PC fluorescence is partly quenched, ~60% lower than when it is in the membrane. For protein-free liposomes, fluorescence is expected to drop by ~30%, whereas for scramblase-containing vesicles fluorescence is expected to drop by ~60%. **G.** Time-course of NBD-PC fluorescence (normalized to the starting value) in liposomes and VDAC2-vesicles on adding BSA.




Figure 2. Native VDAC1 scrambles phospholipids poorly.

A, B. Representative traces corresponding to channel assays performed on vesicles reconstituted with different amounts of VDAC2 (panel A) or VDAC1 (panel B) (Shown are protein concentrations corresponding to theoretical protein/vesicle copy number of 30, 10 and 2). **C.** Protein-dependence plot showing functionalization of vesicles with channel activity, i.e., fraction of large unilamellar vesicles (LUVs) with at least one channel. PPR, protein/phospholipid ratio. The data were analyzed according to a Poisson statistics model for reconstitution of proteins into individual vesicles. A similar mono-exponential fit constant was obtained for both proteins (~ 2 mg/mmol). **D, E.** Representative traces corresponding to scramblase assays performed on vesicles reconstituted with different amounts of VDAC2 (panel D) or VDAC1 (panel E) as in panel A and B. **F.** Correlation between the fraction of LUVs showing channel activity (dithionite assay) and vesicles with scramblase activity (BSA-back extraction assay). Data points correspond to LUVs reconstituted with different amounts of protein. **G, H.** Crosslinking of VDAC proteins after reconstitution into vesicles. Reconstituted LUVs were treated with EGS to crosslink proteins in proximity. The samples were analyzed by SDS-PAGE immunoblotting using antibodies against the N-terminal His tag. Reconstituted VDAC2 shows a significant population of dimers/multimers (panel G) whereas reconstituted VDAC1 is predominantly monomeric (panel H, side panel shows a brighter signal to enable visualization of a faint dimer band)




Figure 3. Crosslinked VDAC1 dimers have phospholipid scramblase activity.

A. VDAC1 in LDAO was cross-linked using EGS, or mocked treated with DMSO, and visualized by SDS-PAGE immunoblotting using antibodies against the N-terminal His tag. B,C. Representative dithionite (B) and BSA-back extraction (C) traces with cross-linked (VDAC1^x) or mock treated (VDAC1) samples reconstituted into LUVs at a theoretical copy number of 30 proteins/vesicle. Normalized fluorescence traces are shown. D. Fraction of VDAC1-vesicles with scramblase activity (reconstituted at a theoretical protein/vesicle copy number of 30). Data points shown as individual points, as well as the mean ± 95% CI, n = 5, p < 0.0001. After EGS cross-linking, nearly all vesicles that show channel activity also have scramblase activity.




Figure 4: Coarse grained MD simulations of scrambling by a specific VDAC1 dimer.

A. Structure of VDAC1 dimer-1. The β -strands at the interface are colored as indicated, and the E73 residue on the β 4-strand is marked. Structures of dimer-1* and dimer-3 are shown in Fig. S8. **B.** Surface representation (white = hydrophobic, green = hydrophilic, blue = positive, red = negative) showing each protomer, rotated 90° to expose the dimer-1 interface, with the contact region marked with a shaded ellipse; polar residues that flank the interface are indicated according to the key under the structures. **C.** Percentage of lipids scrambled as a function of simulation time. The graphs show the scrambling activity of dimer-1, monomer (M) and protein-free membrane (L) (running average over 200 ns time intervals, shading=68% confidence interval, $n=3$). Unless constrained during the simulation, dimer-1 reorients to form dimer-1*. Three individual runs of unconstrained dimer-1 are shown (dimer-1 (relaxed)), the change in scrambling rate between 2-4 μ s coinciding with reorientation. **D.** Bar chart showing scrambling rate for dimer-1, -1*, -3 and monomer (M) (mean \pm SD, $n=3$). **E.** Snapshot of dimer-1 from the simulation showing phospholipids (multiple colors) transiting between bilayer leaflets along the

interface. Membrane phospholipids surrounding the dimer are in grey. **F.** Top view snapshot of dimer-1 showing representative phospholipids transiting across the bilayer on both sides of the interface. **G.** Bilayer thinning at the dimer-1 interface and near E73 in monomeric VDAC1 (thickness indicated by the color scale shown at right). Top views are shown, β -strands at the interface are indicated as colored dots (same color scheme as in panel A); the red dot indicates the β 4-strand where E73 is located (marked by arrowhead); the N-terminal helix is shown as a grey oblong within the VDAC1 pore. **H.** Snapshot side-view of VDAC dimer-1, with one protomer removed, demonstrating membrane thinning at the interface. Ochre surface indicates average positions of lipid phosphates from both membrane leaflets. The protein is shown in surface representation. **I.** Water penetration into the membrane (water defect, color scale shown at right) in the vicinity of the dimer-1 interface and monomeric VDAC1, as indicated.

SUPPLEMENTARY FIGURES S1-S10




Figure S1. MD simulations to assess the energetics of phospholipid translocation in the presence of human VDAC1.

A, B. Phospholipid translocation was analyzed by coarse-grained molecular dynamics simulations of a pure POPC bilayer and a POPC bilayer containing human VDAC1 (PDB ID 6G6U). A phospholipid was pulled from one leaflet of the bilayer to the other in the z-direction. In the case of VDAC1, the lipid phosphate was also restrained to be within a cylinder, centered on E73, with a radius of 1.5 nm. Five selected positions along the translocation pathway from the upper (IMS) to lower (cytoplasmic) side are shown colored from yellow to red (the lipid headgroup is represented by single bead, and acyl chains are represented as thin tubes). Membrane phospholipids are not shown explicitly - their phosphate groups are shown by grey beads, while the lipid tails are not displayed for clarity. VDAC1 is represented as a protein surface color-coded as follows: hydrophobic-white, polar-green, red-negatively charged, and blue-positively charged residues. **C.** Calculated free energy profiles of lipid translocation as a function of the distance between the transiting lipid phosphate and the membrane center or E73 are shown. The presence of VDAC1 leads to a significant decrease in the energy barrier for phospholipid translocation around E73 using coarse-grained Martini 3 (left) or Martini 2 (right) forcefields. The error is estimated $< 2 \text{ kJ.mol}^{-1}$ for the pure POPC bilayer and $< 5 \text{ kJ.mol}^{-1}$ for the system with VDAC based on the profile asymmetry.




Figure S2. Transport-coupled decarboxylation of NBD-PS in isolated yeast mitochondria.

A. Assay schematic. NBD-PS (indicated PS) is added to isolated yeast mitochondria. After insertion into the outer leaflet of the OMM, NBD-PS flips across the membrane where it encounters IMM-localized PS decarboxylase (Psd1) which converts it to NBD-PE (indicated PE). Psd1 can act *in trans*, hydrolyzing PS in the inner leaflet of the OMM as shown. It can also act *in cis* on NBD-PS molecules that are delivered to the IMS-side of the IMM. **B.** Mitochondria were isolated from indicated yeast strains lacking yeast VDAC Por1 and/or Por2, and Psd1, and the protein concentration of the preparations was adjusted to be the same in all cases. Immunoblots using antibodies against the OMM protein Tom70 and the IMM protein Psd1 confirm the similar protein content of the various preparations. **C.** Thin layer chromatogram, visualized with a ChemiDoc imager, of a decarboxylation assay time-course using mitochondria from wild type yeast. **D.** Time-course of NBD-PS decarboxylation at 30°C using mitochondria (1 mg/mL) from the indicated yeast strains, supplemented with 1.25 μM NBD-PS at time=0 min. The data points are shown as mean ± SD, fitted to a single exponential function (2 biological replicates, with at least 2 technical replicates per assay). No NBD-PE was detected in *psd1Δ* cells. **E.** Half-time of NBD-PS to NBD-PE conversion based on data shown in panel D (mean ± SD, $p \approx 0.0021$ (One-Way ANOVA followed up with multiple comparison test, corrected using Bonferroni)).




Figure S3. Purification of human VDAC1 and VDAC2.

A. Human VDAC proteins were produced heterologously in *E. coli*, purified as inclusion bodies, denatured with guanidine hydrochloride, solubilized using the detergent *N,N*-Dimethyldodecylamine *N*-oxide (LDAO) and purified by cation-exchange and size-exclusion chromatography (SEC). SEC profiles and associated Coomassie stained SDS-PAGE analyses are shown. **B.** Circular Dichroism spectroscopy of human VDAC proteins. Spectra were obtained with 66 μM VDAC samples (wild type (WT) VDAC1 and VDAC2, VDAC1-5V mutant, and VDAC2-E84L mutant) in a 0.2mm sandwich cell. All proteins showed high beta sheet content.




Figure S4. VDAC2 E84L is a scramblase.

VDAC2-E84L was purified as in Figure 3 (CD spectrum shown in Fig. 3B), reconstituted into unilamellar vesicles containing NBD-PC at a theoretical protein/vesicle copy number of 10, and assayed for channel and scramblase activity. Wild-type VDAC2 was analyzed in parallel. **A.** Channel assay, as described in Fig. 1D. **B.** Scramblase assay, as described in Fig. 1F.




Figure S5. Crosslinking of LDAO-solubilized VDAC proteins.

The oligomeric state of VDAC1 (left panel) and VDAC2 (right panel) in different concentrations of LDAO (0.05% - 1.05% w/v) was determined by cross-linking using the homobifunctional, amine-reactive crosslinker EGS at a 16-fold molar excess over lysine residues in the protein. EGS was delivered from a stock solution in DMSO (final DMSO concentration = 0.5%) and the reaction was carried out for 40 min at 20°C. Control samples (indicated (-)) were mocked treated with DMSO. The crosslinking reactions were analyzed by SDS-PAGE immunoblotting with anti-His tag antibodies. Oligomerization is highly dependent on the detergent concentration with mainly (VDAC1) or a greater proportion (VDAC2) of monomers present at high detergent concentrations.




Figure S6. VDAC1 monomers scramble phospholipids slowly.

A, B. Schematic of assay to detect slow scrambling. Asymmetric liposomes were prepared by adding 0.25 mol% NBD-PC lipids in EtOH (final EtOH concentration 0.4%) to protein-free liposomes (L) (panel A) or VDAC1-containing proteoliposomes (P) reconstituted at a theoretical protein/vesicle copy number of 10 (panel B). Samples were incubated for 30 min on ice after which they were incubated for approximately 30 h at 20°C. Aliquots were taken at different time points to perform a BSA-back extraction assay and determine the fraction of NBD-lipids present in the outer leaflet. If a slow scramblase is present, over the time course of 30 h, approximately 50% of NBD lipids should be shielded from BSA extraction at the inner leaflet. **C.** Slow scrambling by VDAC1. Two independent experiments with VDAC1-proteoliposomes are shown, fitted to a single exponential decay function. Parallel experiments with protein-free liposomes are also shown. The black dotted line indicates the expected fluorescence signal for the VDAC1-proteoliposomes, experimentally determined from protein reconstitutions with same conditions. The entire complement of NBD-PC molecules in the protein-free vesicles was accessible for BSA-extraction over the time-course of the experiment, indicating little or no scrambling. In contrast, for vesicles reconstituted with hVDAC1, nearly 50% of the NBD-lipids were shielded from BSA-extraction after 20 h, indicating a scrambling half-life ~4 h.




Figure S7. Scramblase activity of VDAC1 dimers prepared with a cleavable crosslinker.

A. Chemical structure of EGS. **B.** Chemical structure of DSP. **C.** Top view of dimer-1, indicating outward-facing lysine residues in protomers designated A and B. **D.** Distances between lysine pairs on VDAC1 protomers in dimer-1 calculated from a coarse-grained MD simulation that accounts for reorientation of the sidechains. Each of the indicated distances is the average xyz distance between the sidechain bead of lysines on protomer A and B. The shaded boxes indicate lysine pairs that are <1.4 nm apart and therefore likely to be crosslinked by EGS and DSP. **E.** VDAC1 in LDAO was cross-linked using DSP or mocked treated with DMSO (indicated (-)) and visualized by SDS-PAGE immunoblotting using antibodies against the N-terminal His tag. An aliquot of the DSP-cross-linked sample was treated with dithiothreitol (DTT) after reconstitution into liposomes and analyzed alongside. **F.** Cross-linked (VDAC^{DSP}) or mock treated (VDAC1) proteins were reconstituted into LUVs at a copy number of 30 proteins/vesicle and assayed for scramblase activity using BSA back extraction as in Fig. 1F. Normalized fluorescence traces are shown. Also shown is an assay in which a VDAC^{DSP}-reconstituted sample was treated with DTT to reverse the crosslink *in situ*. **G.** Fraction of VDAC1-vesicles with scramblase activity; data shown correspond to reconstituted DMSO-treated VDAC1, reconstituted EGS- or DSP-crosslinked VDAC1, and VDAC1^{DSP} treated with DTT after reconstitution at a theoretical protein/vesicle copy number of 30. Open circles in the DMSO and EGS data represent samples that were treated with DTT as a control.




Figure S8. Structures of other VDAC1 dimers and their effects on the membrane.

A, B. Structure of VDAC1 dimer-3 (A) and VDAC1 dimer-1* (B). The β -strands at the interfaces are colored as indicated, and the E73 residue on the β 4-strand is marked. **C.** Top views of water penetration into the membrane (water defect, color LUT shown at right) in the vicinity of the dimer interfaces. The β -strands at the interface are indicated as colored dots (same color scheme as in panels A, B); the E73 residue is shown as a red dot and marked by arrowheads; the N-terminal helix is shown as a grey oblong within the VDAC1 pore. **D.** Bilayer thinning at the dimer interfaces (thickness scale, color LUT shown at right, E73 indicated by arrowhead). Other details are as in panel C.




Figure S9. Analysis of the VDAC1-5V mutant.

A. VDAC1-5V is a dimer/oligomer in liposomes. VDAC1 WT and the 5V mutant were reconstituted into unilamellar vesicles at a theoretical protein/vesicle copy number of 30 and subjected to EGS crosslinking to determine their oligomeric state. Control samples were mocked treated with DMSO. The samples were analyzed by SDS-PAGE immunoblotting using antibodies against the N-terminal His tag, revealing the 5V mutant to be highly dimeric/oligomeric. **B.** Representative channel assay as described in Fig. 1D of reconstituted VDAC1-WT and VDAC1-5V at a theoretical protein/vesicle copy number of 10. **C.** Representative scramblase assay as described in Fig. 1F of reconstituted VDAC1 WT and VDAC1-5V at a theoretical protein/vesicle copy number of 10. **D.** Fraction of VDAC1-vesicles with scramblase activity reconstituted at a theoretical protein/vesicle copy number of 10. Despite its dimeric/multimeric nature after reconstitution as evidenced by EGS crosslinking, the 5V mutant is not an effective scramblase.




Figure S10. The effect of VDAC1 and membrane disruption on the energetics of lipid scrambling.

A. Free energy profiles of phospholipid translocation as a distance between the lipid phosphate and the membrane center in pure POPC membrane (blue line), in the presence of VDAC1 monomer (green and magenta lines in right panel), and in the presence of dimer-1 (azure and orange lines in left panel). Dashed lines correspond to simulations in which the attractive dispersion interactions (Lennard-Jones (LJ) interactions) between the lipid and VDAC1 were turned off, effectively removing the direct contribution of VDAC1 to the decrease in the flip-flop barrier.

B,C. Average enthalpic interactions (Lennard-Jones + electrostatic) between the scrambling lipid and VDAC (B) or between the scrambling lipid and membrane + water + ions (C) as a distance between the scrambling lipid phosphate and the membrane center. The enthalpic interactions were calculated from the simulated umbrella sampling windows. The color scheme is the same as in A.

MOVIE S1

Movie S1. Phospholipid scrambling at the dimer-1 interface.

Movie capturing spontaneous scrambling of a lipid along the interface of dimer-1 in a coarse-grained molecular dynamics simulation. The dimer is shown using molecular surface, the headgroups of most lipids are shown as gray beads and their tails as gray tubes, while a single lipid is highlighted in orange. Most lipids in front of the dimer are not depicted, except for several phosphate groups that are located close to the dimer. The movie captures roughly 140 ns of simulated time.

SUPPLEMENTARY TABLES

Table 1. Distribution of umbrella sampling windows along the collective variable with the biasing force constants used for Martini 3 and Martini 2 free energy calculations of lipid scrambling in a protein-free membrane.

Window No.	Reference position [nm]	Force constant [kJ mol ⁻¹ nm ⁻²]	Window No.	Reference position [nm]	Force constant [kJ mol ⁻¹ nm ⁻²]
1	2.00	3000	25	-0.05	5000
2	1.90	3000	26	-0.10	5000
3	1.80	3000	27	-0.15	5000
4	1.70	3000	28	-0.20	5000
5	1.60	3000	29	-0.25	5000
6	1.50	3000	30	-0.30	3000
7	1.40	3000	31	-0.40	3000
8	1.30	3000	32	-0.50	3000
9	1.20	3000	33	-0.60	3000
10	1.10	3000	34	-0.70	3000
11	1.00	3000	35	-0.80	3000
12	0.90	3000	36	-0.90	3000
13	0.80	3000	37	-1.00	3000
14	0.70	3000	38	-1.10	3000
15	0.60	3000	39	-1.20	3000
16	0.50	3000	40	-1.30	3000
17	0.40	3000	41	-1.40	3000
18	0.30	3000	42	-1.50	3000
19	0.25	5000	43	-1.60	3000
20	0.20	5000	44	-1.70	3000
21	0.15	5000	45	-1.80	3000
22	0.10	5000	46	-1.90	3000
23	0.05	5000	47	-2.00	3000
24	0.00	5000			

Table 2. Distribution of umbrella sampling windows along the collective variable with the biasing force constants used for Martini 3 free energy calculations of lipid scrambling along monomeric VDAC1.

Window No.	Reference position [nm]	Force constant [kJ mol ⁻¹ nm ⁻²]	Window No.	Reference position [nm]	Force constant [kJ mol ⁻¹ nm ⁻²]
1	1.90	1000	33	0.00	4000
2	1.80	1000	34	-0.04	4000
3	1.70	1000	35	-0.08	4000
4	1.60	1000	36	-0.12	4000
5	1.50	1000	37	-0.16	4000
6	1.40	1000	38	-0.20	4000
7	1.30	1000	39	-0.24	4000
8	1.20	1000	40	-0.28	4000
9	1.10	1000	41	-0.32	4000
10	1.00	1000	42	-0.36	4000
11	0.95	2000	43	-0.40	4000
12	0.90	2000	44	-0.40	4000
13	0.85	2000	45	-0.48	4000
14	0.80	2000	46	-0.52	4000
15	0.75	2000	47	-0.56	4000
16	0.70	2000	48	-0.60	2000
17	0.65	2000	49	-0.65	2000
18	0.60	2000	50	-0.70	2000
19	0.56	4000	51	-0.75	2000
20	0.52	4000	52	-0.80	2000
21	0.48	4000	53	-0.90	1000
22	0.44	4000	54	-1.00	1000
23	0.40	4000	55	-1.10	1000
24	0.36	4000	56	-1.20	1000
25	0.32	4000	57	-1.30	1000
26	0.28	4000	58	-1.40	1000
27	0.24	4000	59	-1.50	1000

28	0.20	4000	60	-1.60	1000
29	0.16	4000	61	-1.70	1000
30	0.12	4000	62	-1.80	1000
31	0.08	4000	63	-1.90	1000
32	0.04	4000	64	-2.00	1000

Table 3. Distribution of umbrella sampling windows along the collective variable with the biasing force constants used for Martini 2 free energy calculations of lipid scrambling along monomeric VDAC1.

Window No.	Reference position [nm]	Force constant [kJ mol ⁻¹ nm ⁻²]	Window No.	Reference position [nm]	Force constant [kJ mol ⁻¹ nm ⁻²]
1	1.90	3000	37	0.00	5000
2	1.80	3000	38	-0.04	5000
3	1.70	3000	39	-0.08	5000
4	1.60	3000	40	-0.12	5000
5	1.50	3000	41	-0.16	5000
6	1.40	3000	42	-0.20	5000
7	1.30	3000	43	-0.24	5000
8	1.20	3000	44	-0.28	5000
9	1.10	3000	45	-0.32	5000
10	1.00	3000	46	-0.36	5000
11	0.95	3000	47	-0.40	5000
12	0.90	3000	48	-0.44	5000
13	0.85	3000	49	-0.46	5000
14	0.80	5000	50	-0.48	7000
15	0.75	5000	51	-0.50	7000
16	0.73	7000	52	-0.52	7000
17	0.70	7000	53	-0.54	5000
18	0.68	7000	54	-0.58	5000
19	0.66	7000	55	-0.60	3000
20	0.64	7000	56	-0.65	3000
21	0.62	7000	57	-0.70	3000
22	0.60	5000	58	-0.75	3000
23	0.56	5000	59	-0.80	3000
24	0.52	5000	60	-0.90	3000
25	0.48	5000	61	-1.00	3000
26	0.44	5000	62	-1.10	3000
27	0.40	5000	63	-1.20	3000

28	0.36	5000	64	-1.30	3000
29	0.32	5000	65	-1.40	3000
30	0.28	5000	66	-1.50	3000
31	0.24	5000	67	-1.60	3000
32	0.20	5000	68	-1.70	3000
33	0.16	5000	69	-1.80	3000
34	0.12	5000	70	-1.90	3000
35	0.08	5000	71	-2.00	3000
36	0.04	5000	72	-2.10	3000

Table 4. Distribution of umbrella sampling windows along the collective variable with the biasing force constants used for Martini 3 free energy calculations of lipid scrambling along dimer-1.

Window No.	Reference position [nm]	Force constant [kJ mol ⁻¹ nm ⁻²]	Window No.	Reference position [nm]	Force constant [kJ mol ⁻¹ nm ⁻²]
1	1.80	50	9	-0.20	200
2	1.40	50	10	-0.40	200
3	1.00	50	11	-0.60	100
4	0.80	100	12	-0.80	100
5	0.60	100	13	-1.00	50
6	0.40	200	14	-1.40	50
7	0.20	200	15	-1.80	50
8	0.00	200			

Table 5. Distribution of umbrella sampling windows along the collective variable with the biasing force constants used for Martini 3 free energy calculations of lipid scrambling along VDAC1 monomer and dimer-1 with attractive LJ interactions turned off.

Window No.	Reference position [nm]	Force constant [kJ mol ⁻¹ nm ⁻²]	Window No.	Reference position [nm]	Force constant [kJ mol ⁻¹ nm ⁻²]
1	2.20	1000	33	0.04	4000
2	2.10	1000	34	0.00	4000
3	2.00	1000	35	-0.04	4000
4	1.90	1000	36	-0.08	4000
5	1.80	1000	37	-0.12	4000
6	1.70	1000	38	-0.16	4000
7	1.60	1000	39	-0.20	4000
8	1.50	1000	40	-0.24	4000
9	1.40	1000	41	-0.28	4000
10	1.30	1000	42	-0.32	4000
11	1.20	1000	43	-0.36	4000
12	1.10	1000	44	-0.40	4000
13	1.00	1000	45	-0.44	4000
14	0.90	1000	46	-0.48	4000
15	0.80	1000	47	-0.52	4000
16	0.75	2000	48	-0.56	4000
17	0.70	2000	49	-0.60	2000
18	0.65	2000	50	-0.65	2000
19	0.60	2000	51	-0.70	1000
20	0.56	4000	52	-0.80	1000
21	0.52	4000	53	-0.90	1000
22	0.48	4000	54	-1.00	1000
23	0.44	4000	55	-1.10	1000
24	0.40	4000	56	-1.20	1000
25	0.36	4000	57	-1.30	1000
26	0.32	4000	58	-1.40	1000
27	0.28	4000	59	-1.50	1000
28	0.24	4000	60	-1.60	1000
29	0.20	4000	61	-1.70	1000
30	0.16	4000	62	-1.80	1000
31	0.12	4000	63	-1.90	1000
32	0.08	4000	64	-2.00	1000

Table 6. Yeast strains used in this study.

Strain name	Genetic Background	Mating Type	Genotype	Source
TKY705 (Wild Type)	W303	a	ura3-52 trp1Δ2 leu2-3_112 his3-11 ade2-1 CAN1	Miyata et al. (2018)
TKY705 psd1Δ	W303	a	TKY705, psd1Δ::natMX4	This study
TKY705 por1Δ	W303	a	TKY705, por1Δ::natMX4	This study
TKY705 por2Δ	W303	a	TKY705, por2Δ::natMX4	This study
TKY705 por1Δpor2Δ	W303	a	TKY705, por1Δ::natNT2 por2Δ::hphNT1	Miyata et al. (2018)
TKY705 por1Δ tet::por2Δ	W303	a	TKY705, por1Δ::natNT2 tetO ₇ -POR2::kanMX4	Miyata et al. (2018)
TKY705 psd1Δ por1Δ tet::por2Δ	W303	a	TKY705, psd1Δ::hphMX4 por1Δ::natNT2 tetO ₇ -POR2::kanMX4	This study

Miyata N, Fujii S, Kuge O (2018) Porin proteins have critical functions in mitochondrial phospholipid metabolism in yeast. J Biol Chem 293: 17593-17605

Table 7. PCR templates and primers used in this study.

Locus	Template plasmid	Primer (5' to 3')
psd1Δ::natMX4	p4339	1: AAT CCT TCT TGG TCG TTA TTT TTT GAA GAA GAA GGA AAA GCA AAG CCA GCG ACA TGG AGG CCC AGA ATA C 2: GAT AAT ACT ATA TAC AGC AAA ATA AAT GCT AAC TTT ACA TAT GAT TGC TTC AGT ATA GCG ACC AGC ATT C
por1Δ::natMX4	p4339	1: ACG AAA CAG CCA AGC GTA CCC AAA GCA AAA ATC AAA CCA ACC TCT CAA CAG ACA TGG AGG CCC AGA ATA C 2: GGA AAC AAG AAC GAG CAC ATA TAT GGT ATA TAG TGA ACA TAT ATA TAT TAG ATA TAT ACG TCA GTA TAG CGA CCA GCA TTC
por2Δ::natMX4	p4339	1: TCT CGA GTG AAG GAA TTG CAT GAA GAG GAA AGT GTT AGA AAT TAC CTA CGG ACA TGG AGG CCC AGA ATA C 2: TCA AAA GTA CAA ATC CTT ATT TAG TAA AAT TTA TAA GAA AAT AAA ATC TCC AGT ATA GCG ACC AGC ATT CA
psd1Δ::hphMX4	pMJ696	1: AAT CCT TCT

TGG TCG TTA TTT
TTT GAA GAA GAA
GGA AAA GCA
AAG CCA GCG ACA
TGG AGG CCC AGA
ATA C
2: GAT AAT ACT
ATA TAC AGC AAA
ATA AAT GCT AAC
TTT ACA TAT GAT
TGC TTC AGT ATA
GCG ACC AGC ATT
C