

Palaeogenomic analysis of black rat (*Rattus rattus*) reveals multiple European introductions associated with human economic history

Authors

He Yu^{1,57}, Alexandra Jamieson^{2,55}, Ardern Hulme-Beaman^{3,4}, Chris J. Conroy⁵, Becky Knight⁶, Camilla Speller^{7,8}, Hiba Al-Jarah⁷, Heidi Eager⁹, Alexandra Trinks^{2,10}, Gamini Adikari¹¹, Henriette Baron¹², Beate Böhlendorf-Arslan¹³, Wijerathne Bohingamuwa¹⁴, Alison Crowther^{15,16}, Thomas Cucchi¹⁷, Kinie Esser¹⁸, Jeffrey Fleisher¹⁹, Louisa Gidney²⁰, Elena Gladilina²¹, Pavel Gol'din²², Steven M. Goodman²³, Sheila Hamilton-Dyer²⁴, Richard Helm²⁵, Chris Hillman²⁶, Nabil Kallala²⁷, Hanna Kivikero²⁸, Zsófia E. Kovács²⁹, Günther Karl Kunst²⁰, René Kysely³¹, Anna Linderholm³², Bouthéina Maraoui-Telmini³³, Arturo Morales-Muñiz³⁴, Mariana Nabais^{35,36}, Terry O'Connor⁷, Tarek Oueslati³⁷, Eréndira M. Quintana Morales³⁸, Kerstin Pasda³⁹, Jude Perera⁴⁰, Nimal Perera⁴⁰, Silvia Radbauer⁴¹, Joan Ramon⁴², Eve Rannamäe⁴³, Joan Sanmartí Grego⁴⁴, Edward Treasure⁴⁵, Silvia Valenzuela-Lamas⁴⁶, Inge van der Jagt⁴⁷, Wim Van Neer^{48,49}, Jean-Denis Vigne¹⁷, Thomas Walker⁵⁰, Stephanie Wynne-Jones⁶, Jørn Zeiler⁵¹, Keith Dobney^{3,52,53,54}, Nicole Boivin^{15,16}, Jeremy B. Searle⁹, Ben Krause-Kyora⁵⁵, Johannes Krause^{1,56,*}, Greger Larson^{2,*}, David Orton^{7,*}

¹Department of Archaeogenetics, Max Planck Institute for the Science of Human History, Jena 07745, Germany

²Palaeogenomics & Bio-Archaeology Research Network, Research Laboratory for Archaeology and History of Art, University of Oxford, Oxford OX1 3QY, United Kingdom

³Department of Archaeology, Classics and Egyptology, University of Liverpool, Liverpool L69 7WZ, United Kingdom

⁴Research Centre in Evolutionary Anthropology and Palaeoecology, Liverpool John Moores University, Liverpool L3 3AF, United Kingdom

⁵Museum of Vertebrate Zoology, University of California, Berkeley, Berkeley, CA 94720-3160, USA

⁶Department of Archaeology, University of York, York YO1 7EP, United Kingdom

⁷BioArCh, Department of Archaeology, University of York, York YO1 7EP, United Kingdom

⁸Department of Anthropology, University of British Columbia, Vancouver, BC, Canada

⁹Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY 14853, USA

¹⁰Charité – Universitätsmedizin Berlin, Institut für Pathologie, Charitéplatz 1, 10117 Berlin, Deutschland

¹¹Postgraduate Institute of Archaeology, 407, Baudhaloka Mawatha, Colombo -7, Sri Lanka.

¹²Römisch-Germanisches Zentralmuseum, Leibniz-Forschungsinstitut für Archäologie, Ernst - Ludwig-Platz, 2D – 55116, Mainz, Germany

- 36 ¹³Christian Archaeology and Byzantine Art History, Philipps University of Marburg, 35037 Marburg,
37 Germany
- 38 ¹⁴Department of History and Archaeology, University of Ruhuna, Matara - 81000, Sri Lanka.
- 39 ¹⁵Department of Archaeology, Max Planck Institute for the Science of Human History, Jena 07745,
40 Germany
- 41 ¹⁶School of Social Science, The University of Queensland, St Lucia, Queensland, Australia.
- 42 ¹⁷Archaeozoology, Archaeobotany, Societies, Practices, Environments (AASPE-UMR7209), CNRS,
43 National Museum of Natural History (MNHN), Paris, France
- 44 ¹⁸Archeoplan Eco, 2616 LZ, Delft, Netherlands
- 45 ¹⁹Department of Anthropology, Rice University, 6100 Main St, Houston, TX 77005
- 46 ²⁰Archaeological Services, University of Durham, United Kingdom
- 47 ²¹Ukrainian Scientific Centre of Ecology of the Sea, Odessa 65009, Ukraine
- 48 ²²Schmalhausen Institute of Zoology, National Academy of Sciences of Ukraine, Kiev 01030, Ukraine
- 49 ²³Negaunee Integrative Research Center, Field Museum of Natural History, Chicago, IL 60605, USA
- 50 ²⁴Department of Archaeology & Anthropology, Bournemouth University (Visiting Fellow), Poole,
51 BH12 5BB, United Kingdom
- 52 ²⁵Canterbury Archaeological Trust, 92a Broad Street, Canterbury, Kent CT1 2LU, UK
- 53 ²⁶6 Fell View Park, Gosforth, Seascale, Cumbria, CA20 1HY, UK
- 54 ²⁷Institut National du Patrimoine, Tunis 1008, Tunisia
- 55 ²⁸University of Helsinki, Department of Culture, P.O. Box 59, FI-00014 Helsinki, Finland
- 56 ²⁹Freelance archaeozoologist, Liliom u. 4. 1/1., Balatonfüred, 8230, Hungary
- 57 ³⁰VIAS Vienna Institute for Archaeological Science, University of Vienna, Althanstraße 14, 1090
58 Vienna, Austria
- 59 ³¹Department of Natural Sciences and Archaeometry, Institute of Archaeology of the Czech Academy
60 of Sciences, Prague, Letenská 4, 118 01, Prague, Czech Republic
- 61 ³²Department of Anthropology, Texas A&M University, College Station, TX 77843, USA
- 62 ³³Department of History, University of Tunis, Tunis, Tunisia
- 63 ³⁴Departamento de Biología, Universidad Autónoma de Madrid, 28049 Madrid, Spain
- 64 ³⁵Institute of Archaeology, University College London, London WC1H 0PY, UK
- 65 ³⁶Centro de Arqueologia da Universidade de Lisboa (UNIARQ), Faculdade de Letras, Universidade
66 de Lisboa, 1600-214 Lisboa, Portugal
- 67 ³⁷Centre National de la Recherche Scientifique, University of Lille, France
- 68 ³⁸Department of Anthropology, University of California, Santa Cruz, 1156 High St, Santa Cruz, CA
69 95064, USA
- 70 ³⁹Department of Philosophy, Institute of Prehistoric Archaeology, Friedrich-Alexander-University of
71 Erlangen-Nürnberg, 91054 Erlangen, Germany
- 72 ⁴⁰Department of Archaeology, Sir Marcus Fernando Mawatha, Colombo-07, Sri Lanka

- 73 ⁴¹Austrian Academy of Sciences, Austrian Archaeological Institute, Hollandstraße 11-13, 1020
74 Vienna, Austria
- 75 ⁴²Consell Insular d'Eivissa i Formentera
- 76 ⁴³Department of Archaeology, Institute of History and Archaeology, University of Tartu, 2 Jakobi St.,
77 51005 Tartu, Estonia
- 78 ⁴⁴Secció de Prehistòria i Arqueologia, University of Barcelona, Spain
- 79 ⁴⁵Department of Archaeology, Durham University, Durham, UK
- 80 ⁴⁶Archaeology of Social Dynamics, IMF-CSIC, Barcelona 08001, Spain
- 81 ⁴⁷Cultural Heritage Agency of the Netherlands, Smallepad 5, 3811 MG Amersfoort, The Netherlands
- 82 ⁴⁸Royal Belgian Institute of Natural Sciences, Vautierstraat 29, 1000 Brussels, Belgium
- 83 ⁴⁹Laboratory of Biodiversity and Evolutionary Genomics, Katholieke Universiteit Leuven, 3000 Leuven,
84 Belgium
- 85 ⁵⁰Department of Archaeology, University of Reading, Berkshire RG6 6AB, United Kingdom
- 86 ⁵¹ArchaeoBone, 9753 JN Haren, The Netherlands
- 87 ⁵²Department of Archaeology, University of Sydney, Sydney, Australia
- 88 ⁵³Department of Archaeology, University of Aberdeen, Aberdeen, UK
- 89 ⁵⁴Department of Archaeology, Simon Fraser University, Burnaby, Canada
- 90 ⁵⁵Institute of Clinical Molecular Biology, Kiel University, Kiel 24105, Germany
- 91 ⁵⁶Max Planck Institute for Evolutionary Anthropology, Leipzig 04103, Germany

92

93 ⁵⁷These authors contributed equally

94 *Correspondence: krause@eva.mpg.de, greger.larson@arch.ox.ac.uk, david.orton@york.ac.uk

96 Abstract

97 The distribution of the black rat (*Rattus rattus*) has been heavily influenced by its association with
98 humans. The dispersal history of this non-native commensal rodent across Europe, however, remains
99 poorly understood, and different introductions may have occurred during the Roman and medieval
100 periods. Here, in order to reconstruct the population history of European black rats, we generated a *de*
101 *novo* genome assembly of the black rat, 67 ancient black rat mitogenomes and 36 ancient nuclear
102 genomes from sites spanning the 1st-17th centuries CE in Europe and North Africa. Analyses of
103 mitochondrial DNA confirm that black rats were introduced into the Mediterranean and Europe from
104 Southwest Asia. Genomic analyses of the ancient rats reveal a population turnover in temperate
105 Europe between the 6th and 10th centuries CE, coincident with an archaeologically attested decline in
106 the black rat population. The near disappearance and re-emergence of black rats in Europe may have
107 been the result of the breakdown of the Roman Empire, the First Plague Pandemic, and/or post-
108 Roman climatic cooling.

109

Introduction

The black rat (*Rattus rattus*) is one of three rodent species, along with the house mouse (*Mus musculus*) and brown rat (*Rattus norvegicus*), to have become globally distributed thanks to a close commensal relationship with humans¹. Collectively, these taxa are highly significant to human societies both as pests responsible for billions of euros of damage to food stores annually², and as vectors and/or reservoirs that have contributed to the spread of numerous diseases, most infamously bubonic plague^{3,4}.

Despite the significance of this rodent, our knowledge of the black rat's evolutionary history and taxonomy remains limited. Previous genetic studies have described a *R. rattus* complex involving multiple recognized species with potential introgression among different lineages⁵⁻⁷. Mitochondrial DNA studies have helped to resolve the taxonomic controversies by linking a monophyletic mitochondrial lineage to specific South Asian (now globally distributed) *R. rattus* populations that possess a 2n=38 karyotype (previously referred to as lineage I)⁸⁻¹⁰. The Asian house rat (*Rattus tanezumi*), endemic to Southeast Asia, has been identified as the closest sister group of the black rat (previously designated as lineages II through IV). The divergence between the two species has been dated to ~0.4 Mya¹¹, and the two have been suggested to hybridise^{6,7,12}.

The ability of rats to colonise, and become dependent upon, anthropogenic niches¹³ makes them ideal bioproxies to track historical processes^{1,14,15}. Archaeological specimens of rats and mice have thus been used to track human migrations, trade, and settlement types in a wide range of contexts¹⁶⁻²². Previous archaeological and genetic evidence suggests that the pre-commensal distribution of the Eurasian black rat (based on the taxonomic definition proposed by mitochondrial DNA studies^{8,9} and hereafter referred to as black rat, see SI for discussion) was largely limited to South Asia^{10,23,24}. Black rat finds from cave sediments in the Levant spanning the late Pleistocene to early Holocene indicate a possible western distribution^{25,26}. These remains require direct dating to confirm their age, and there is a subsequent absence of rats from settlement sites in this region until at least the 2nd millennium BCE²⁶.

The earliest large concentrations of presumed commensal rat remains reported thus far derive from late third, or early second millennium BCE settlements in both the Indus Valley and Mesopotamia²⁶. Commensal black rats may also have reached the Levant and eastern Mediterranean region by the start of the first millennium BCE²⁶. Based on archaeological evidence from Corsica, the Balearics, Italy and Morocco²⁷⁻²⁹, black rats likely first appeared in the western Mediterranean basin towards the end of that same millennium.

The colonisation of Europe by the black rat has been linked to the historical development of urbanism and trade networks, and their arrival is important for understanding historical plague pandemics including the 6th C Justinianic Plague and the 14th C Black Death^{4,30–32}. The central role traditionally attributed to black rats and their fleas in the spread of the plague bacterium (*Yersinia pestis*) during these pandemics has been challenged on various grounds, however, including the historical distribution and abundance of rats, and this correlation continues to be debated^{33–37}.

Although surveys of zooarchaeological rat finds from archaeological sites across Europe suffer from considerable regional variation in coverage, the available data indicates successive episodes of dispersal north of the Mediterranean associated first with Roman expansion (first century BCE to second century CE), and then with the emergence of medieval economies from the 9th century CE, punctuated by a decline and possible range contraction³². Black rat remains are found throughout the Roman Empire in the 1st to 5th centuries CE, but rarely beyond its northern borders suggesting that these rats were dependent on a Roman economic system characterised by a network of dense settlements connected by bulk transport via efficient road, river, and maritime routes^{4,31}.

With the breakdown of the Roman Empire from the 5th century onwards, evidence for the existence of black rats becomes scarcer. They may have been extirpated entirely from the northern provinces including Britain^{32,38,39}, and the percentage of archaeological sites with black rat remains declined even in the Western Empire's Italian core⁴⁰. By contrast, black rats remained common in the Balkans and Anatolia until at least the 6th century CE, presumably reflecting continued stability in the Eastern Roman Empire^{41–44}. Since 5th–8th century zooarchaeological data is limited in many regions, the pattern of post-Roman absence may partly represent research bias⁴⁵, though early medieval black rats are rare even where other small mammals are reported³⁸.

Black rats reappear at northern European trading settlements during the 9th and 10th centuries CE, including sites well beyond their Roman range, including Hedeby in northern Germany and Birka in Sweden, as well as former Roman towns and high-status early medieval settlements such as York and Flixborough in England and Sulzbach in Bavaria^{46–50}. The subsequent expansion of urbanism and large-scale trade of bulk goods in medieval Europe appears to have favoured rats, just as in the Roman period. By the 13th century CE, black rats were present throughout most of Europe⁴ and they reached southern Finland by the late 14th century⁵¹. Rats remained widespread across Europe until at least the 18th century, before their population declined, most likely as a result of competition with the newly arrived brown rat, the now dominant rat species in temperate Europe^{52–54}.

It remains unclear whether the black rat was actually extirpated from post-Roman northern and western Europe; and whether medieval rat populations in temperate Europe derived from the remnant

population in southern Europe, or from another wave of rats that were introduced from beyond the Mediterranean (e.g. via Rus' river trade^{32,55}). These questions are relevant to several key debates in European economic and environmental history including: 1) the extent to which the end of the Western Roman Empire represented a crisis in urbanism and trade — particularly in bulk goods such as grain — as well as political collapse^{56–58}; 2) the role of easterly vs. westerly connections in the rise of northern European medieval urban networks^{59,60}; and 3) the model of the spread of the Justinianic Plague and the subsequent First Pandemic. This pandemic started in the eastern Mediterranean in 541 CE, spread quickly across Europe and England, and continued for approximately two centuries^{61–63}, a period that coincides with the gap in archaeological evidence for rats in northwest Europe. Given the limitations of both zooarchaeology and genetic studies of modern rat populations to address successive waves of contact after a species is established, ancient DNA may help to resolve these questions by directly revealing the presence or lack of genetic continuity through time.

We employed a nested three-stage approach to address these questions. First, we assembled a *de novo* reference genome of the black rat. This genome allowed us to investigate the long-term demographic history of *Rattus rattus* in relation to other rat species, and to lay the foundation for genome-wide analyses of ancient remains. Second, we explored the dispersal of black rats into the Mediterranean and Europe by analysing 70 new mitochondrial genomes from European and North African archaeological specimens spanning the Roman to early post-medieval periods (1st-17th century CE), alongside 132 mtDNA sequences generated from modern and museum black rat specimens from across western Eurasia, the Indian Ocean, and Africa. Lastly, we generated 39 nuclear genomes from our archaeological black rats and used these to explore the species' population history in Europe and the Mediterranean from the 1st to 17th centuries CE, focusing particularly on the hypothesis of dual dispersals in the Roman and medieval periods. We then interpreted the black rat's dispersal history within the context of major historical processes.

Results and Discussion

The demographic history of *Rattus rattus* and its closely related species

To facilitate the study of the demographic history of the black rat, both before and after the establishment of its commensal relationship with humans, we first generated a *de novo* genome assembly of *R. rattus* using a wild-caught black rat from California, USA. Combining shotgun, Chicago and Hi-C sequencing data with the Dovetail HiRise assembler pipeline⁶⁴, we obtained a genome assembly with a total length of 2.25 Gb and a scaffold N50 reaching 145.8 Mb (Supplementary Table 1, 2). The 22 scaffolds with over 10 Mb covering 98.9% of the entire assembly (Table 1), with each of the 18 autosomes of *R. rattus* corresponding to one large scaffold each and over 90% of the X chromosome represented by four scaffolds (Supplementary Figure 11, 12, Note 2). The average GC content is 42.1%, similar to the *R. norvegicus* reference genome *Rnor_6.0* (42.3%),

and 38.4% of the assembly was identified as repetitive elements (Supplementary Table 3). Benchmarking Universal Single-Copy Orthologs (BUSCO) analysis⁶⁵ also revealed a high completeness of this genome assembly, with 90.1% complete BUSCOs identified using eukaryotic dataset, comparable with *Rnor_6.0* (91.4%) (Supplementary Table 4). Because *R. tanezumi* and *R. rattus* are both present in California, we also assessed potential introgression from *R. tanezumi* into our *R. rattus* individual. A signature of introgression would limit the value of our *de novo* genome as a reference genome onto which reads derived from ancient black rats could be mapped. Our analyses suggested no significant introgression signature in the Californian black rat (Supplementary Note 5).

Table 1. Assembly statistics of the *de novo* *R. rattus* reference genome

| | |
|---------------------------------------|-------|
| Scaffold Number | 6805 |
| Scaffold N50 (Mb) | 145.8 |
| Largest Scaffold (Mb) | 260.8 |
| Assembly size (Gb) | 2.25 |
| Scaffold length >10 Mb (Gb) | 2.23 |
| GC content (%) | 42.1 |
| Repetitive region (%) | 38.4 |

To address the demographic history of black rat, we applied the Pairwise Sequentially Markovian Coalescent (PSMC)⁶⁶ analysis to estimate its population size dynamics alongside the brown rat and Asian house rat. When calibrated with a mutation rate of 2.96×10^{-9} per generation and generation time of 0.5 years⁶⁷, the analyses revealed different dynamic patterns of population size changes amongst these rat species (Figure 1A). The brown rat experienced a population decline beginning ~1 Mya, as described previously⁶⁷, while both the black rat and Asian house rat populations expanded until 300-400 thousand years ago (kya). The black rat population then experienced a bottleneck with an 8-fold drop in effective population size until 100 kya, and a re-expansion from 100 kya to 40 kya. The Asian house rat, however, did not experience a population decline until ~40 kya, when both black rat and Asian house rat populations experienced declines that have continued to the present.

To investigate the population sizes, split times, and migrations among these rat lineages, we applied Generalized Phylogenetic Coalescent Sampler (G-PhoCS)⁶⁸. The result revealed a similar population size dynamism, with the effective population size (N_e) of black rat/Asian house rat ancestral lineage estimated to be 1.25×10^6 , about tenfold the N_e of black rat/Asian house rat/brown rat lineages (Figure 1B, Supplementary Table 5, 6). The split time between brown rat and black rat/Asian house rat lineages was estimated to be 1.94 Mya (within the 95% Highest Posterior Density (HPD) range estimated using mitochondrial genomes in a previous study¹¹), while the split of Asian house rat and

black rat lineages took place ~120 kya. This recent split time relative to the coalescent time estimate based on mitochondrial genomes between these two lineages could be explained by the large ancestral population size of the black rat/Asian house rat lineage⁶⁸. Among these lineages, we only detected one instance of gene flow from the black rat/Asian house rat ancestral lineage into brown rat lineage, with an introgression proportion of 9.8%.

Taken together, we observed population expansions and bottlenecks in the black rat during the last million years, and a smaller N_e relative to the Asian house rat. This could be explained by the relatively limited geographic distribution of the black rat in southern Asia before the initiation of its commensal relationship with people, and the fact that the Asian house rat is endemic to a much greater area in southeastern Asia¹⁰. We did not detect any genomic introgression between the lineages leading to the black and Asian house rat, suggesting these two species were geographically isolated after their split from a common ancestor for a sufficiently long period to facilitate their reproductive incompatibility.




Figure 1. The demographic history of the black rat and its closely related species.

(A) Population dynamics of the black rat (*R. rattus*), Asian house rat (*R. tanezumi*) and brown rat (*R. norvegicus*) estimated by PSMC, with 100 bootstrap replicates.

(B) Demographic modelling of the divergence and migration among the black rat, Asian house rat and brown rat estimated by G-PhoCS. The values represent the average estimates of effective population sizes (in thousands), population divergence times (Mya) and the total migration rate through time. The 95% HPD range of all estimates are listed in Supplementary Table 6.

A global phylogeography of the black rat based on mitochondrial DNA

We collected 191 ancient black rat individuals from 33 archaeological sites across Europe, North and East Africa, and southern Asia dating from the 2nd millennium BCE to the 17th century CE (Supplementary Table 7), plus eight modern individuals from North Africa. After shotgun screening, we retrieved 70 mitochondrial genomes (with coverage spanning 3.5x-300.0x) from samples from 18 sites in Europe and the Mediterranean (Supplementary Table 8), and identified 40 haplotypes. The phylogenetic tree based on mitochondrial genomes revealed two clades: a major clade with 32 haplotypes, and a minor clade consisting of eight haplotypes and 23 ancient samples from the 6th-century site of Caričin Grad, Serbia (Supplementary Figure 3). The phylogenetic resolution within each major clade was relatively poor, though samples from the same or closely related sites occasionally formed sub-clades including the samples from modern-day Zembra (Tunisia) and medieval central Europe.

In order to establish the relationship between the ancient rats and modern black rats from across their range, we analysed the cytochrome b (CYTB) region from our ancient samples alongside 132 previously unpublished modern CYTB sequences from across the Indian Ocean basin (Supplementary Table 9), and sequences published in previous studies^{10,24,69,70}. The maximum likelihood tree of the CYTB region revealed that all the ancient rats from this study belong to the previously described black rat lineage I (Figure 2, Supplementary Figure 4, 5 and Table 10). None of the ancient rats from this study fell into *Rattus* lineages II-VI. Within lineage I, we recapitulated the unnamed substructure and assigned the terms A to E to the five major haplogroups¹⁰. In addition to these five, we confirmed a sixth lineage I haplogroup, F, consisting of modern samples from Sri Lanka and the Andaman Islands, which is basal relative to all other lineage I black rats and has previously been reported as the Sri Lankan unique sub-lineage, RrC LIb⁷¹ (Figure 2, Supplementary Figure 4b, Table 10).

Haplogroup A within lineage I (previously described as the European ship rat²⁴) was the most common among the analysed samples (179/354). Members of this haplogroup include ancient and modern black rats from Europe and regions of the world with a history of colonisation by, and/or trade with European powers. The only additional haplogroup found in Europe was Haplogroup C (previously described as the Arab ship rat²⁴) at Caričin Grad, Serbia, which included 24 archaeological individuals. Haplogroup C is found in modern rats from India, Egypt, East, South and West Africa and South America. None of the other haplogroups were present in Europe or the

Mediterranean region. Haplogroups B and E only included modern samples from India and countries bordering the Indian Ocean. Haplogroup D (previously described as the Madagascar and Indian Ocean islands group²⁴) included primarily samples from Madagascar and East Africa, and Haplogroup F consisted of samples from Sri Lanka and the Andaman Islands (Figure 2, Supplementary Table 10).




Figure 2. Sampling sites and mitochondrial phylogeographic patterns.

A - Map of sampling locations. The ancient sample SMI only has mitochondrial data, the rest have both nuclear and mitochondrial results. SMI (Villa Franca de Campo) MEP (Mertola) KLT (Kilton Castle) TRU (Tanner Row, York) GAU (Gatehampton Villa) VOB (Voorburg-Forum Hadriani) SNE (Deventer-Stadhuiskwartier) MDT (Monte di Tuda) SML (Santa Maria Lavezzi) ATU (Althiburos) Sulz (Castle Sulzbach) PRA (Prague Castle) PZA (Petronell-Carnuntum Zivilstadt) KAF (Kastelholm) BUD (Buda Castle-Teleki Palace) Car (Caričin Grad) Ass (Assos). B - The phylogeographic pattern of black rat revealed by CYTB mitochondrial haplogroups (see Supplementary Figure 4 for detailed phylogeny). We derived modern nuclear genomes from the reference genome rat from California and individuals trapped on the island of Zembra (ZMB) in the 1980s that were prepared for a zooarchaeological reference collection.

To investigate the introduction route of black rats into Europe, we analyzed mitochondrial cytochrome B sequences derived from globally distributed modern and ancient black rats. Previous studies indicated that the black rat originated in the Indian Peninsula^{10,24,69,72}. Leaving aside the putative late Pleistocene to early Holocene records from the eastern Levant, the earliest finds of presumed commensal rats derive from the Indus Valley and Mesopotamia in the 3rd/2nd millennium BCE, coincident with the emergence of urbanism and establishment of trade links between these regions^{26,73}, though a more westerly limit to the black rat's natal range cannot be excluded. The source for dispersal to the Mediterranean and ultimately Europe remains unclear. Suggestions include maritime trade from India and/or the Arabian peninsula into the Red Sea and subsequently through Egypt (perhaps via the canal built under Darius⁷⁴) in the mid/late first-millennium BCE^{4,32}, or more likely earlier overland communication routes between Mesopotamia and the Levant^{26,73}.

While a maritime route is clearly implicated in the black rat's dispersal to East Africa^{75,76}, our results tentatively favour an overland hypothesis for its dispersal from South Asia to the Mediterranean to Europe, since both ancient and modern black rats from Europe and the eastern Mediterranean share haplogroups with sampled populations from Iran and the Persian Gulf, but not with Indian Ocean samples from southern India to Madagascar (Figure 2). The results also suggest a secondary dispersal route via Egypt, given the appearance of Haplogroup C at the 6th century CE Byzantine site of Caričin Grad, Serbia and in modern samples from the Nile valley. While tentative, this may reflect Egypt's central roles both in direct Indo-Roman trade, following its annexation in 30 BCE, and in grain production for the Roman and early Byzantine Empires^{4,77}. To test these hypotheses, further investigations into ancient and modern black rat populations from the Levant, Mesopotamia, Egypt and the Indus Valley are necessary.

Ancient genomes reveal the relationships of European black rats over space and time

To explore the black rat's European population history in greater detail, we shotgun sequenced 36 ancient and 3 modern black rats from 17 sites to 0.2x-16x coverage for whole genome analysis, including 18 females and 21 males determined by the coverage on sex chromosomes (Supplementary Table 11). The deeper sequenced ancient samples spanned two broad time periods, including 15 from the Roman and Early Byzantine period (1st to 7th century CE), and 21 from medieval and post-medieval contexts (8th to 17th century CE) (Supplementary Table 12). Geographically, all the samples were divided into two groups: a "northern" group of 25 samples from temperate Europe, and a "southern" group of 11 samples from the Mediterranean and Portugal (Figure 2). After mapping and genotyping, we identified 7,869,069 bi-allelic transversion variants in the autosomal non-repetitive regions for downstream population genetic analysis.

The phylogenetic tree constructed from autosomal SNPs revealed complex relationships among ancient black rats from different regions and time periods (Figure 3A). Except for the late medieval (c. 14th century) to Ottoman (c. 17th century) site of Buda Castle, Hungary, samples from the same site are clustered together. All the samples from the northern group, together with one southern sample from the medieval period — from 8th-9th century Althiburos, Tunisia — formed a single clade, while all the other Byzantine to medieval samples from the southern group formed several separate clades consistent with their local geographic region. The rats from the southern group also possessed higher heterozygosity than those from the northern group, within both Roman/Byzantine and medieval/post-medieval periods (Supplementary Figure 7, Table 13). This could be explained by the longer history of rats in the Mediterranean which date to at least the first millennium BCE^{26,27}, and the founder effects of limited introductory waves of rats into the northern region.

Within the major northern cluster, samples were divided into two smaller clusters representing Roman / Byzantine and medieval / post-medieval periods respectively. The only exception was a medieval Tunisian sample that falls into the Roman cluster. Within each cluster, samples grouped together based upon their geographic location (central Europe, western/northern Europe, Serbia). These phylogenetic relationships suggest that the initial black rat population in temperate Europe was replaced by a genetically distinct population after the 6th century CE. The later population is first documented in early medieval (8th to early 10th century CE) Sulzbach, Germany. The Roman-like gene pool was still present during the 8th-9th century in North Africa, though due to the lack of more recent samples we cannot address whether or when the second wave arrived there. A similar pattern was also revealed by multidimensional scaling (MDS) based on isolation-by-state (IBS) distance among the samples (Supplementary Figure 8).

The phylogenetic tree based on Y-chromosome scpMSY regions (Supplementary Table 14) similarly demonstrated that the Roman rats formed a single cluster. However, unlike the autosomal phylogeny, all the post-Roman samples from both the northern and southern groups, including Byzantine Assos and Caričin grad, formed a separate cluster (Supplementary Figure 9), without well-supported substructures. Given the male-biased dispersal pattern commonly described in the black rat and other rodent species^{78,79}, this might indicate a male-specific replacement that took place in both the Mediterranean regions and temperate Europe.

A decline in the European black rat population during the 6th to 9th centuries has previously been suggested based on zooarchaeological evidence^{32,38,39}. This has been attributed to several causes including: (a) the demise of the Western Roman Empire's economic and urban system from the 5th century CE, including the cessation of large-scale grain shipments that may have helped to disperse and support rat populations⁴; (b) climatic cooling in the 'Late Antique Little Ice Age'⁸⁰; and/or (c)

the Justinianic Plague, which began in 541 CE and is likely to have infected rat populations previously naive to *Yersinia pestis*, regardless of their potential role in its spread among humans^{4,81,82}. Our finding of a post-6th-century turnover corroborates this apparent decline, though the density of our samples' spatiotemporal coverage is not sufficient for us to distinguish between the potential causes. To understand how the Justinianic Plague influenced the rat population, further studies should focus on archaeological black rats from contexts post-dating the mid-6th century in areas of the Byzantine Empire and wider Mediterranean where an urban settlement system persisted.

The medieval Tunisian (Althiburos) sample indicates a different population history of black rats in North Africa relative to temperate Europe. Black rats from a wider range of time periods resident in North Africa and the western Mediterranean would allow us to test whether there was continuity within the black rat populations from the Roman to early Islamic period (c.8th century). This is particularly pertinent to debates concerning the degree of continuity between the Roman Empire and the Early Islamic world, notably in urban settlements and trade networks⁸³.




Figure 3. Relationships of the ancient black rats over time and space.

(A) The ages of the rat samples included in whole genome analyses. The bars represent 95.4% confidence intervals surrounding the direct radiocarbon dates or stratigraphic dates (Supplementary Table 11,12). The colors correspond to the Roman/Byzantine (red) and medieval (blue) time periods.

The symbols represent the sampling sites listed in panel B, and the modern samples are represented by black symbols.

(B) The phylogenetic relationship among ancient and modern black rats reconstructed using a neighbor joining phylogeny. The pairwise genetic distances were calculated using autosomal variants. The support values based upon 100 bootstrap replicates are shown on the nodes. The branches are colored by sample ages as described in panel A, and the tip symbols correspond to the sampling site. * Though the medieval Tunisian (Althiburos) sample clusters geographically in the Southern group, it falls in with the Roman cluster of the northern group in the phylogeny.

To investigate the genetic interaction between different rat populations further, we applied a series of f_4 -statistics. Based on the result of the f_4 -statistics symmetry test, the ancient samples were divided into 18 groups (Supplementary Table 15). Sixteen of these correspond to samples from 16 different sites, while the three late/post-medieval samples from Buda Castle (Hungary) fell into two groups corresponding respectively to late medieval (14th-15th century) and Ottoman (17th century) periods. (Supplementary Table 12).

First, we investigated if any Roman population contributed to the Byzantine or medieval groups, with $f_4(\text{norvegicus}, \text{Byzantine/medieval}; \text{Roman1}, \text{Roman2})$. We found that of two Roman geographical groups (central European represented by Austria, and western Europe represented by Britain and the Netherlands) the western rats were significantly more closely related to all the Byzantine and medieval groups (Figure 4A, Supplementary Figure 10, Table 16). This result suggests that despite the population turnover that occurred in temperate Europe after the Roman period, Roman black rats from western Europe may have contributed to populations that colonized temperate Europe following the decline of the original population.

Next, we applied $f_4(\text{norvegicus}, \text{Roman}; \text{Byzantine/medieval1}, \text{Byzantine/medieval2})$ to test if there were any differences in the relative contribution of Roman rat ancestry into the Byzantine or medieval populations. In agreement with the phylogenetic and MDS analysis, most northern groups were significantly more closely related to the Roman rat populations compared to the Byzantine or more recent southern groups (SML, MDT, Ass). The lone exception to this pattern were two post-medieval samples from Buda Castle (BUD001/4), which were equally related to the Roman groups and the Assos (Ass) group that consists of two samples from Byzantine Turkey (Supplementary Table 17). Among the northern groups, the medieval rats from Åland (Finland), the UK and the Netherlands, as well as Byzantine rats from Serbia, were more closely related to the Roman rat populations than were medieval rats from central Europe (represented by populations in Germany, Czech Republic and Hungary). This suggests that the genetic contribution from putatively western European Roman rats, was also greater in the local western European medieval rats.

We also investigated the relationship between the Buda Castle (Hungary) samples from different time periods by contrasting them with the other medieval rats from temperate Europe (Figure 4A, Supplementary Figure 10, Table 18). As revealed by the phylogenetic tree, both the German and Czech rats shared more genetic affinity with the c.14th-15th century Buda Castle (BUD003) sample, than with the 17th century or later specimens (BUD001/4). Having said that, BUD001/4 still showed higher affinity to BUD003, when compared with all other populations. This evidence suggests a black rat population transition in this region between the 14th/15th century and the late 17th century, potentially related to the 16th-17th century Ottoman occupation of Buda (Hungary), while the local medieval ancestry was still present in the later population.

To corroborate the patterns of gene flow suggested by the f_4 -statistics, we used Treemix⁸⁴ to generate an admixture graph of all ancient rat populations, using the Asian house rat as an outgroup. The maximum-likelihood population tree without any admixture produced a similar topology to the neighbor-joining autosomal phylogeny (Supplementary Figure 11). The rats from the northern group and a medieval Tunisian rat formed a clade, to which all the other Mediterranean rats were an outgroup, without any significant clustering pattern among the lineages. When admixture events were allowed, the first two suggested gene flow edges were from the medieval central European population into the post-medieval Buda Castle population, estimated to 18.2% +/- 3.0%, and from the Roman western European population into the ancestral lineage of the medieval European populations in the northern group, estimated to be 8.1% +/- 0.8% (Figure 4D).




Figure 4. Gene flow among ancient rat populations.

(A) The f -statistics showing admixture between different ancient rat populations. The dots show all the combinations of f_4 -values as described above each cluster, and the error bars show $\pm 3SE$ of the estimates. The three clusters show the affinity between: (top, red) medieval rats (Med) and western European Roman rats (Roman_WEU); (middle, blue) Roman rats (Roman) and western European medieval rats (Med_WEU); and (bottom, green) post-medieval Buda Castle rats (BUD001/4) and the medieval Buda Castle rat (BUD003), respectively.

(B) Admixture graph with two migration events fitted, estimated by Treemix. The migration edges are displayed by arrow including the introgression fractions and standard errors. The color of each branch represents the time period of each group: Roman/Byzantine (red) and medieval / post-medieval (blue).

The results from both the f -statistics and Treemix analyses revealed a degree of Roman rat ancestry in the medieval populations. More specifically, medieval rats were more closely related to the Roman rat populations from the Netherlands and Britain (Figure 4, Supplementary Table 16). This signal suggests a reservoir of black rat population in western Europe that admixed with the re-introduced medieval population. The stronger affinity of medieval western European populations to Roman populations (Supplementary Table 17) also suggested that this relict population was more likely distributed in western and not central Europe. This result could indicate that rats from the northernmost Roman provinces were not extirpated, despite their absence in early medieval zooarchaeological assemblages. Alternatively, and in our view more likely, the inferred relict population may have been located in an unsampled region of France or southwest Europe. The observation that medieval rats from temperate Europe fall into the same cluster as Roman rats also suggests that the second (medieval) wave of introduction to temperate Europe probably originated from the same source population as the first (Roman) dispersal. Considering the zooarchaeological evidence that rat populations in southern Europe persisted after the collapse of the Western Roman Empire, notably in Italy⁴⁰, it is likely that southern Europe was the source of reintroduced rats in temperate Europe.

Given the presence of rats in 9th century northern *emporium* (proto-urban trading sites) around the North and Baltic Seas^{46,49,50}, a southern European origin would emphasise the importance of the Carolingian Empire (the Frankish polity which controlled much of western and central Europe as well as northern Italy in the 9th century CE) and routes such as the Rhône and Rhine corridors in reestablishing large-scale trade links between the Mediterranean and northern Europe⁸⁵. This connection remains tentative until samples from the early *emporium* themselves in mainland Italy and the Iberian Peninsula can be investigated. Samples from the early Islamic world derived from the Iberian Peninsula and North Africa would also help to clarify the population history of black rats.

Conclusion

This study explores the historic dispersal of commensal black rats using a *de novo* genome assembly for the black rat, ancient and modern mtDNA from across Europe, Africa, and the Indian Ocean, and ancient nuclear genomes from the Mediterranean and Europe. Our results suggest that the black rat was most likely introduced to the eastern Mediterranean by an overland route through Southwest Asia, though a maritime route via the Indian Ocean and Red Sea cannot be excluded. We identify two waves of rat introduction into temperate Europe. The first likely accompanied the Roman northward expansion during the first centuries BCE/CE and the second took place during the medieval period (starting in the 8th-10th centuries). The rats in this second wave were probably derived from the same ancestral population as the first, and subsequently admixed with a western or southern European relict population from the first wave.

Considered alongside the paucity of archaeological rat remains from the 6th-8th centuries CE (particularly in northern and western Europe), this population turnover suggests that the black rat population and range declined during the early medieval period. This may have been associated with the breakdown of the Roman Empire - from the 5th century CE in western Europe and the early 7th century CE in the Balkans - and with it, the network of well-connected settlements that had previously supported black rat populations. Grain shipments may have played a key role in the dispersal and maintenance of rat populations during the Roman period, and it is notable that weevils (*Sitophilus granarius*) and other grain pests show a similar pattern of Roman introduction, apparent post-Roman extirpation, and a medieval reintroduction in the northern provinces⁸⁶. Alternatively, or additionally, European rat populations may have been negatively impacted by the First Plague Pandemic and/or the climatic cooling of the Late Antique Little Ice Age, both of which began in the mid-6th century CE. To disentangle these scenarios, further zooarchaeological and genomic studies of ancient rats are required that span these centuries across a wider geographic range.

The medieval introduction of rats into Northern Europe is attested (at the latest) by their presence in Germany in the early 10th century, coincident with an increase in rat bone finds across the continent. Our results suggest a repopulation of temperate Europe from the south, perhaps linked with the development of trade routes in Carolingian western Europe, and probably not via early Russian riverine trade, as has been previously hypothesised³². Black rats appear to have been a continuous presence in Europe from this point until the post-medieval period spanning the 14th century Black Death and extending into the 17th century. This population may also have been supplemented by localised introductions, including one potentially associated with the Ottoman occupation of Buda from 1541 CE).

The near extirpation of black rats in modern Europe is likely linked to competition with the brown rat which arrived from Asia in the early 18th century^{52,53}. The genetic and demographic impact of this dispersal on black rats is an important area for future investigations since by the late 18th century,

naturalists in many European countries had already attributed a marked decline in *R. rattus* to competition from *R. norvegicus*^{87–90}. The black rat’s significantly reduced, but persistent presence, particularly in towns, suggests a degree of niche partitioning between the two species⁹¹.

Our results reveal the degree to which human-commensal species can undergo population dispersal and demographic fluctuations. In fact, because these dynamic evolutionary processes are tightly correlated with the characteristics of the human niche, commensal species can act as ideal proxies to interpret the history of human movement and cultural change.

Methods

Radiocarbon dating and calibration

Fourteen ancient rat bones were radiocarbon dated via accelerator mass spectrometry (AMS) on bone collagen at Mannheim (MAMS), University of Waikato (Wk), and Oxford University (OxA), and these were analysed alongside two previously published dates from Gatehampton⁹² (Supplementary Table 12). One additional sample (SNE002 / Wk-51521) failed due to insufficient collagen. All radiocarbon dates were calibrated in OxCal 4.4⁹³, using the IntCal20 calibration curve⁹⁴.

Given the omnivorous diet of black rats, stable isotope values were monitored for evidence of marine dietary contributions that might result in significant reservoir effects. Where possible, $\delta^{13}\text{C}$ (‰ vs. VPDB) and $\delta^{15}\text{N}$ values (‰ vs. AIR) were obtained by the respective dating laboratories using Isotope Ratio Mass Spectrometry (IRMS) and their standard protocols; otherwise $\delta^{13}\text{C}$ values were used as reported from the AMS. Nitrogen isotope values were available for 10 specimens and fell between 6.9‰–11.9‰, consistent with published data for commensal brown rats⁹⁵ and with an omnivorous diet. Carbon isotope values ranged from -21.9‰ to -17.4‰. In the absence of detailed local comparative isotope data from terrestrial and marine species, it was not possible confidently to estimate marine dietary contribution and hence, the magnitude of any required correction. Nonetheless, we performed indicative corrections for specimens whose $\delta^{13}\text{C}$ values suggested a possible non-negligible marine component in order to test for any possible impact on our interpretations. The cut-off for this was set as -18.5‰, based on published values for European terrestrial herbivores and the enrichment expected due to trophic level in an omnivore.

For six specimens with $\delta^{13}\text{C} > -18.5$ ‰, percentage marine contribution was estimated using “formula 1” from⁹⁶, with terrestrial and marine endpoints of -21‰ and -12‰ respectively and a trophic enrichment factor of 1‰. These values were used to recalibrate the dates using mixed IntCal20 and Marine20⁹⁷ curves, and the magnitude of potential offset was assessed. In no cases did the median calibrated date change by more than 140 years, and in no case would it have moved a specimen into a different chronological category or altered our interpretations. Given the uncertainty inherent in this

process, the uncorrected ranges are used in Figure 3, and details of the indicative corrections are given in Supplementary Table 12. In the majority of cases, the uncorrected date range coincided more closely with the stratigraphic dating.

***De novo* genome assembly**

The black rat genome was sequenced and assembled using DNA extracted from the liver of a male wild-caught individual from California, USA. Shotgun, Chicago and Dovetail Hi-C libraries were prepared and sequenced on Illumina HiSeq 4000 platform and the genome was assembled using Meraculous⁹⁸ and HiRise scaffolding pipeline⁶⁴. The detailed information of genome assembly is provided in the Supplementary Note 2.

The repetitive regions were identified using RepeatMasker 4.0.7⁹⁹ with Repbase 20170127 and the query species set as *rattus*, and TRF 4.09 (Tandem repeats finder)¹⁰⁰, with parameters set as “2 7 7 80 10 50 12”. The completeness of genome assembly was assessed by BUSCO 3.0.2⁶⁵, using the 303 orthologs in Eukaryota odb9 dataset. The new genome assembly was aligned to the brown rat reference genome *Rnor_6.0* using nucmer 4.0.0 in MUMmer tool package¹⁰¹, to investigate the synteny between *R. rattus* and *R. norvegicus* genomes, using both masked assemblies and anchor matches that are unique in both reference and query (Supplementary Note 2).

Mitochondrial Cytochrome B fragment sequencing

Overall, 292 tissue samples identified as *R. rattus* were included for analysis, including 263 museum specimens (sampled from the collections at: American Museum of Natural History, British National History Museum, Field Museum Chicago) and 29 modern specimens collected in the field representing a wide geographic area at the periphery and on islands within the Indian Ocean.

DNA extraction and sequencing of these modern and museum samples were conducted in the modern laboratory at the Archaeology Department of Durham University, following standard protocols (Supplementary Note 4). The *cytb* region was amplified in 10 overlapping fragments and a variety of primer combinations was used depending on the nature of the sample (Supplementary Table 19). The sequencing reaction was carried out by the DNA Sequencing Service at the School of Biological and Biomedical Sciences at Durham University. The sequencing chromatograms were edited manually, subsequently assembled, and a consensus sequence per individual exported using Geneious R6 version 6.0.6¹⁰². Standard anti-contamination guidelines were followed. We successfully amplified *cytb* sequences from 202 of 292 samples. Only those that possessed >90% gene coverage were included in the analysis, which left 132 sequences.

Ancient DNA extraction and processing

We sampled 191 ancient black rats and eight modern black rat individuals from 33 archaeological sites across Europe and three sites in North Africa (Supplementary Table 7, Note 1). Where multiple samples were taken from the same or related archaeological contexts, care was taken to ensure that these represented discrete individuals - either by sampling the same skeletal element and side, or on the basis of differing size and / or age.

Ancient DNA extraction was performed in dedicated ancient DNA facilities at the University of Oxford, the Max Planck Institute for the Science of Human History in Jena and the University of York. All of the ancient lab facilities followed standard ancient DNA laboratory practices to minimise contamination, including the use of blanks at each stage from extraction to amplification. All material analysed at Oxford underwent the following treatment. Due to the small size of black rat bones, the outer surface of the bones was not removed prior to extraction. Bones that weighed <50 mg were completely consumed during the extraction process. The bone or tooth was cut using a Dremel drill with a clean cutting wheel per sample (Dremel no 409) and pulverised in a Micro-dismembrator (Sartorius-Stedim Biotech). Material analysed at York was subjected to bleach treatment (6% sodium hypochlorite for 5 minutes, and then rinsed with ultrapure water 3 times) prior to powdering following the same procedure as Oxford.

Extractions performed in Jena followed a silica-based protocol¹⁰³ using 50mg of bone powder. Extractions performed at the University of Oxford were conducted using the Dabney protocol with a modification of the addition of a 30 minute pre-digestion stage¹⁰⁴. Extractions performed at the University of York were conducted using¹⁰⁵ as modified in¹⁰⁶.

For each sample processed in Jena, a double-stranded DNA sequencing library was prepared from 20 µL of extract, with partial uracil-DNA-glycosylase (UDG) treatment (hereafter denoted as ds_halfUDG) or without UDG treatment (ds_nonUDG), following a published protocol¹⁰⁷. Sample-specific index combinations were added to the sequencing libraries^{108,109}. The indexed libraries were shotgun sequenced on an Illumina HiSeq 4000 instrument for screening, with 75 single-end-run cycles for ds_halfUDG libraries and 75 double-end-run cycles for ds_nonUDG libraries. After screening, one ds_nonUDG library and seven ds_halfUDG libraries were deep sequenced in the University of Kiel, on an Illumina HiSeq 4000 platform with 75 double-end-run cycles using the manufacturer's protocol.

All extracts generated at the University of Oxford and the University of York were built into Illumina libraries using double stranded methods using the Blunt-End Single-Tube Illumina library building (BEST) protocol as described previously¹¹⁰ at the University of Oxford (ds_nonUDG). An additional barcode was added to the IS1_adapter.P5 adapter resulting in a double external indexed library. The

libraries were then amplified on an Applied Biosystems StepOnePlus Real-Time PCR system, to determine both the success of the library build and the number of optimum cycles to use for the indexing PCR reactions. These 164 libraries were pooled at equimolar concentrations ready for sequencing. The pool of libraries was sequenced on an Illumina HiSeq 4000 (paired-end 75 bp) at the Danish National High-Throughput Sequencing Centre for screening at Novogene, Sacramento.

Ten extracts from Oxford were built into single-stranded libraries at the Max Planck Institute for Evolutionary Anthropology in Leipzig, Germany. The libraries were built from 30µl of DNA extract in the absence of uracil DNA glycosylase (ss_nonUDG) followed by double indexing, using an automated version of the protocols described in ^{108,109} on a liquid handling system (Agilent Technologies Bravo NGS Workstation). From the initial screening run results 31 ds_nonUDG libraries from Oxford were included for deeper sequencing in Jena, together with the ten ss_nonUDG libraries, on an Illumina HiSeq 4000 platform with 75 single-end-run cycles (Supplementary Table 11).

Genotyping and dataset preparation

The shotgun sequencing reads from 39 ancient and modern black rats were cleaned and mapped to the *de novo* *R. rattus* genome assembly using EAGER pipeline 1.92.55 ¹¹¹. Within the pipeline, the adapters were removed by AdapterRemoval 2.2.0 ¹¹², reads were mapped with BWA 0.7.12 aln/samse algorithm ¹¹³, duplications were removed by DeDup 0.12.1 (<https://github.com/apeltzer/DeDup>) and damage patterns of each library were checked with mapDamage 2.0.6 ¹¹⁴. For the seven ds_halfUDG libraries, we masked 2bp from both ends of the reads using trimBam in bamUtil 1.0.13 ¹¹⁵ to remove the damaged sites.

The shotgun sequencing reads from four modern individuals, including the Californian black rat for *de novo* genome assembly, two published *R. norvegicus* individuals (Accession: ERS215789, ERS215791) ⁶⁷ and one published *R. tanezumii* individual (Accession: SRS1581480, HXM4) ¹¹⁶ were mapped to the genome assembly using the BWA 0.7.12 mem algorithm. After using a mapping quality filtering of 30 and removing reads with multiple hits, duplications were removed using DeDup. We then performed indel realignment for cleaned bam files of both ancient and modern individuals using RealignerTargetCreator and IndelRealigner in The Genome Analysis Toolkit (GATK) v3.5-0 ¹¹⁷.

For the demographic history analysis, we called diploid genotypes from three modern genomes using the highest coverage genome of each of the species: *R. rattus* (CP-5999), *R. norvegicus* (ERS215791) and *R. tanezumii* (HXM4). Each of the bam files was piled up using samtools mpileup, using reads with mapping quality and base quality over 30, and BAQ disabled. Bi-allelic SNPs were then

individually called using bcftools call -m mode and filtered for SNPs with phred-scaled quality score (QUAL) over 30, sequence depth between 0.5-2X mean coverage, and not within 5bp of an indel. After masking for repetitive regions, the consensus sequences of 18 largest autosomal scaffolds were generated, with heterozygous sites represented by IUPAC nucleotide code.

The sequencing reads of ancient and modern black rats after AdapterRemoval were also mapped to *R. rattus* reference mitochondrial sequence NC_012374.1 with BWA 0.7.12 aln/samse algorithm and realigned with CircularMapper¹¹¹. The reads of *R. norvegicus* and *R. tanezumi* individuals were mapped to mitochondrial references of *R. norvegicus* (NC_001665.2) and *R. tanezumi* (NC_011638.1), respectively. After removing duplication using DeDup, the consensus sequences were generated by Schmutzi with a quality threshold of 30¹¹⁸.

We called the pseudo-haploid genotypes in autosomal regions, from all modern and ancient individuals using ANGSD 0.931¹¹⁹, with parameter “-doHaploCall 1” to randomly sample one base. As the 18 longest autosomal scaffolds covered >99% of the autosomal assembly, we only called genotypes on the non-repetitive regions of these 18 scaffolds. We applied “-remove_bads 1 -uniqueOnly 1 -minMapQ 30 -minQ 30 -C 50 -baq 1” parameters to filter out reads that had multiple hits, with mapping quality or base quality less than 30, perform base alignment quality (BAQ) computation and adjust mapping quality based for excessive mismatches¹²⁰. To remove the deamination-induced damages in ancient DNA molecules, we only kept the transversion variants for downstream analysis. The genotypes on single-copied male-specific Y-chromosome regions (scpMSY) were called from all male individuals using ANGSD 0.931, with the same filters as autosomal genotyping, and -doHaploCall 2 to get the major call. The detailed information of scpMSY regions identification was provided in Supplementary Note 2.

To estimate the heterozygosity rates of ancient rat samples, the cleaned reads with base quality and mapping quality over 30 were piled up with mpileup in SAMtools 1.3¹²¹. We then called pseudo-diploid genotypes with pileupCaller 1.2.2 (<https://github.com/stschiff/sequenceTools>) under random diploid calling mode, which randomly sampled two reads at each site, on the transversion variants identified in ANGSD. The heterozygosity rates calculated from pseudo-diploid genotypes were half of the real heterozygosity rates of the samples on these variants.

Demographic history analysis

The population size dynamics was estimated using PSMC 0.6.5⁶⁶, with parameter “-N25 -t20 -r5 -p “4+25*2+4+6”” and 100 bootstrap replicates. The PSMC output was visualized with generation time of 0.5 years and mutation rate $\mu = 3 \times 10^{-9}$ site/generation, based on an estimate calculated in a previous study of *R. norvegicus*⁶⁷.

G-PhoCS⁶⁸ was applied to estimate the population sizes, population divergence times and migration rates among three rat species, using the three high-coverage diploid genomes. The analysis was performed on 38,078 loci of 1kb length, identified in non-repetitive, autosomal regions. A preliminary analysis with all possible migration events was first run for 250,000 generations, then two parallel runs for 500,000 generations with one migration event were carried out for parameter estimation. Finally, the estimated parameters were converted to effective population sizes (N_e), divergence times (T) and total migration rates (m_{total}) as described in⁶⁸: $\theta = 4 * N_e * \mu$, $\tau = T * \mu / g$ and $m_{total} = m * \tau$, with mutation rate $\mu = 2.96 * 10^{-9}$ site/generation and generation time (g) of 0.5 years. The detailed information for loci selection and analysis was provided in Supplementary Note 3.

Phylogenetic analysis

The ancient mitochondrial genomes were analyzed alongside seven modern reference genomes, including the modern Californian black rat from the reference genome assembly, two published *R. norvegicus* individuals⁶⁷, one published *R. tanezumi* individual¹¹⁶ and the published mitochondrial genome references of the three species (*R. rattus* NC_012374.1, *R. tanezumi* NC_011638.1, *R. norvegicus* NC_001665.2). The haplotypes were aligned using MUSCLE v3.8.1551¹²² with default parameters, and the best-fit model was selected based on Akaike Information Criterion (AIC) calculated by jmodeltest v2.1.10¹²³. Then Maximum Likelihood (ML) tree was built using RAxML v8.2.12¹²⁴, with GTR+I+G model and 100 bootstrap replicates.

The cytb region of the mitochondrial genome haplotypes were extracted using MEGA7, and combined with modern cytb haplotypes from previous publications^{10,24,69,70} and this study. We aligned the data using MAFFT v7.123b¹²⁵, then built a ML tree using RAxML v8.2.9¹²⁴, with GTR+I+G model and 1000 bootstrap replicates.

The autosomal phylogeny was reconstructed using neighbor-joining (NJ) method implemented in package Ape 5.3 in R 3.5.1. The distance matrix was calculated based on 3,393,710 autosomal transversion variants, after removing singletons, using the genetic distance described in⁶⁸. Bootstrapping was performed by resampling the variants from 100 kb non-overlapping windows, and the support on each node was calculated based on 100 bootstrap replicates. The phylogenetic tree based on Y-chromosome scpMSY regions was built with RaxML 8.2.12¹²⁴, using GTR substitution model, ML estimation of base frequencies and 100 rapid bootstrapping replicates.

Population genetics analysis

The IBS distance matrix among individuals was calculated using PLINK v1.90b¹²⁶ with parameter “--distance 1-ibs”. MDS analysis was performed using PLINK and ten dimensions were calculated on

both datasets including all studied samples and *R. rattus* samples only. The f_4 -statistics were calculated by *qpDstat* 755 in ADMIXTOOLS 5.1 package¹²⁷, with parameter “f4 mode: YES”, and the two *R. norvegicus* individuals were used as outgroup in all the analysis.

We also applied Treemix 1.13⁸⁴ to simultaneously infer the population structure and admixture events among black rat populations. The black rat samples were grouped based on the geographic location, time period and phylogenetic pattern identified in previous analysis (Supplementary Table 11). The allele frequency was calculated by PLINK and 1,145,713 sites covered in at least one sample from each group were included in the analysis. We built the admixture graph assuming 0 to 10 migration events, with parameters “-k 500 -global -se -noss -root tanezumi” to group 500 SNPs per block for covariance matrix estimation. We then performed a global rearrangement after adding all the populations, calculated standard errors of migration weights, disabled sample size correction and assigned *R. tanezumi* as the root of the topology.

Data availability

The *R. rattus* genome assembly is available in the NCBI under the accession number GCA_011800105.1. Aligned reads from the 39 newly reported ancient black rats are available at the ENA archive under the accession number (provided upon acceptance). The mitochondrial genome haplotypes are available under the accession number (provided upon acceptance).

Acknowledgements

We thank the wet laboratory teams at MPI-SHH, the PalaeoBARN at the University of Oxford and the University of York. We thank David K. James and Lucia Hui of the Alameda County Vector Control Services District for procuring the rat used for the *de novo* genome. We are grateful to Sarah Nagel at Max Planck Institute for the Evolutionary Anthropology for the single-stranded library preparation, and Dovetail Genomics for the *de novo* genome assembly service. We thank Maria Spyrou for her suggestions and comments. We acknowledge Ewan Chipping and Helena England (University of York), Carl Phillips, Veronica Lindholm (Ålands Museum), Christine McDonnell and Nienke van Doorn (York Archaeological Trust), Emile Mittendorf (Gemeente Deventer), Inge Riemersma (Archaeological depot, Provincie Zuid-Holland), the Turkish Ministry of Culture & Tourism, Jan Frolík and Iva Herichová (Institute of Archaeology of the Czech Academy of Sciences, Prague), Franz Humer and Eduard Pollhammer (Archaeological Park Carnuntum), Dorottya B. Nyékhelyi and László Daróczi-Szabó (Budapest History Museum), Institut National du Patrimoine (Tunisia), University of Barcelona, Spanish Ministry of Science and Innovation (Project HUM2006-03432/HIST), Spanish Ministry of Culture (program of archaeological excavations abroad 2009); Spanish Agency of International Cooperation for the Development (2009), Catalan Institute of Classical Archaeology (ICAC), Vujadin Ivanisević, Nemanja Marković and Ivan Bugarski

(Archaeological Institute Belgrade), the Field Museum Chicago, the British National History Museum and the American Museum of Natural History for providing materials and support. G.L. and A.J. were supported by the ERC (grant ERC-2013-StG-337574-UNDEAD) and A.J. was supported by the Natural Environment Research Council Doctoral Training Program. D.O. was supported by Wellcome (Small Grant in Humanities and Social Science 209817/Z) and the British Academy / Leverhulme Trust (Small Research Grant SG170938). E.R. was supported by Estonian Research Council grant No PRG29. R.K. was supported by the Czech Academy of Sciences institutional support (RVO:67985912). S.V.-L. was supported by the ERC (grant ERC-StG- 716298 ZooMWest). H.E. was funded by an ERC grant (206148) through the Sealinks Project. A.H.B was funded by the Leverhulme Trust (ECF-2017-315). The *de novo* genome assembly, population genomics study, and radiocarbon dating were funded by the Max Planck Society.

Author contributions

J.K., G.L. and D.O. designed the project; A.J., B.K.-K., B.K., C.S., H.A-J., H.E., A.T. generated data; H.Y., A.J. and D.O. analyzed data; G.A, N.B., H.B., B.B.-A., W.B., A.C., C.J.C., T.C., K.D., K.E., J.F., L.G., E.G., P.G., S.G., S.H.-D., C.H., R.H., N.K., H.K., Z.K., K.K., R.K., A.L., B.M.-T., A.M.-M., M.N., T.O'C., T.Ou, E.Q.M., K.P., J.P., N.P., S.R., J.R., E.R., J.S., J.S.G., E.T., S.V.-L., I.van der J., W.V.N., J.-D.V., T.W., S.W.-J. and J.Z. provided material and support; H.Y., D.O., A.J., A.H.-B., G.L. and J.K. wrote the paper with contributions from all other authors.

References

1. Boivin, N. L. *et al.* Ecological consequences of human niche construction: Examining long-term anthropogenic shaping of global species distributions. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 6388–6396 (2016).
2. Capizzi, D., Bertolino, S. & Mortelliti, A. Rating the rat: global patterns and research priorities in impacts and management of rodent pests. *Mamm. Rev.* **44**, 148–162 (2014).
3. Meerburg, B. G., Singleton, G. R. & Kijlstra, A. Rodent-borne diseases and their risks for public health. *Crit. Rev. Microbiol.* **35**, 221–270 (2009).
4. McCormick, M. Rats, Communications, and Plague: Toward an Ecological History. *J. Interdiscip. Hist.* **34**, 1–25 (2003).
5. Aplin, K. P., Brown, P., Jacob, J., Krebs, C. J. & Singleton, G. R. *Field methods for rodent studies in Asia and the Indo-Pacific.* (2003).

6. Lack, J. B. *et al.* Invasion facilitates hybridization with introgression in the *Rattus rattus* species complex. *Mol. Ecol.* **21**, 3545–3561 (2012).
7. Conroy, C. J. *et al.* Cryptic genetic diversity in *Rattus* of the San Francisco Bay region, California. *Biol. Invasions* **15**, 741–758 (2013).
8. Robins, J. H., Hingston, M., Matisoo-Smith, E. & Ross, H. A. Identifying *Rattus* species using mitochondrial DNA. *Mol. Ecol. Notes* **7**, 717–729 (2007).
9. Pagès, M. *et al.* Revisiting the taxonomy of the Rattini tribe: a phylogeny-based delimitation of species boundaries. *BMC Evol. Biol.* **10**, 184 (2010).
10. Aplin, K. P. *et al.* Multiple geographic origins of commensalism and complex dispersal history of Black Rats. *PLoS ONE* **6**, e26357 (2011).
11. Robins, J. H. *et al.* Dating of divergences within the *Rattus* genus phylogeny using whole mitochondrial genomes. *Mol. Phylogenet. Evol.* **49**, 460–466 (2008).
12. Yosida, T. H., Kato, H., Tsuchiya, K. & Moriwaki, K. Karyotypes and serum transferrin patterns of hybrids between Asian and Oceanian black rats, *Rattus rattus*. *Chromosoma* **34**, 40–50 (1971).
13. Hulme-Beaman, A., Dobney, K., Cucchi, T. & Searle, J. B. An Ecological and Evolutionary Framework for Commensalism in Anthropogenic Environments. *Trends Ecol. Evol.* **31**, 633–645 (2016).
14. Jones, E. P., Eager, H. M., Gabriel, S. I., Jóhannesdóttir, F. & Searle, J. B. Genetic tracking of mice and other bioproxies to infer human history. *Trends Genet.* **29**, 298–308 (2013).
15. Puckett, E. E., Orton, D. & Munshi-South, J. Commensal Rats and Humans: Integrating Rodent Phylogeography and Zooarchaeology to Highlight Connections between Human Societies. *Bioessays* e1900160 (2020).
16. Jones, E. P. *et al.* Fellow travellers: a concordance of colonization patterns between mice and men in the North Atlantic region. *BMC Evol. Biol.* **12**, 35 (2012).
17. Cucchi, T. Uluburun shipwreck stowaway house mouse: molar shape analysis and indirect clues about the vessel's last journey. *J. Archaeol. Sci.* **35**, 2953–2959 (2008).
18. Cucchi, T. *et al.* On the trail of Neolithic mice and men towards Transcaucasia: zooarchaeological clues from Nakhchivan (Azerbaijan). *Biol. J. Linn. Soc. Lond.* **108**, 917–928

- 869 (2013).
- 870 19. Cucchi, T. *et al.* Tracking the Near Eastern origins and European dispersal of the western house
871 mouse. *Sci. Rep.* **10**, 8276 (2020).
- 872 20. Matisoo-Smith, E. & Robins, J. H. Origins and dispersals of Pacific peoples: evidence from
873 mtDNA phylogenies of the Pacific rat. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 9167–9172 (2004).
- 874 21. Matisoo-Smith, E. & Robins, J. Mitochondrial DNA evidence for the spread of Pacific rats
875 through Oceania. *Biol. Invasions* **11**, 1521–1527 (2009).
- 876 22. West, K. *et al.* The Pacific Rat Race to Easter Island: Tracking the Prehistoric Dispersal of
877 *Rattus exulans* Using Ancient Mitochondrial Genomes. *Frontiers in Ecology and Evolution* **5**, 52
878 (2017).
- 879 23. Niethammer, V. J. Zur Taxonomie und Ausbreitungsgeschichte der Hausratte (*Rattus rattus*).
880 *Zool. Anz. Jena* **194**, 405–415 (1975).
- 881 24. Baig, M., Khan, S., Eager, H., Atkulwar, A. & Searle, J. B. Phylogeography of the black rat
882 *Rattus rattus* in India and the implications for its dispersal history in Eurasia. *Biol. Invasions* **21**,
883 417–433 (2019).
- 884 25. Tchernov, E. Commensal animals and human sedentism in the Middle East. *Animals and*
885 *archaeology* **3**, 91–115 (1984).
- 886 26. Ervynck, A. Sedentism or urbanism? On the origin of the commensal black rat (*Rattus rattus*). in
887 *Bones and the Man: studies in honour of Don Brothwell* (eds. Dobney, K. & O'Connor, T. P.)
888 95–109 (Oxbow, 2002).
- 889 27. Ruffino, L. & Vidal, E. Early colonization of Mediterranean islands by *Rattus rattus*: a review of
890 zooarcheological data. *Biol. Invasions* **12**, 2389–2394 (2010).
- 891 28. Vigne, J.-D. & Valladas, H. È. Small Mammal Fossil Assemblages as Indicators of
892 Environmental Change in Northern Corsica during the Last 2500 Years. *J. Archaeol. Sci.* **23**,
893 199–215 (1996).
- 894 29. Oueslati, T. *et al.* ‘1st century BCE occurrence of chicken, house mouse and black rat in
895 Morocco: Socio-economic changes around the reign of Juba II on the site of Rirha’. *Journal of*
896 *Archaeological Science: Reports* **29**, 102162 (2020).

- 897 30. Audoin-Rouzeau, F. & Vigne, J.-D. La colonisation de l'Europe par le rat noir (*Rattus rattus*).
898 *Revue de Paléobiologie* **13**, 125–145 (1994).
- 899 31. Audoin-Rouzeau, F. & Vigne, et J.-D. Le rat noir (*Rattus rattus*) en Europe Antique et
900 médiévale: les voies du commerce et l'expansion de la peste. *Anthropozoologica* **25/26**, 399–404
901 (1997).
- 902 32. Armitage, P. L. Unwelcome companions: ancient rats reviewed. *Antiquity* **68**, 231–240 (1994).
- 903 33. Hufthammer, A. K. & Walløe, L. Rats cannot have been intermediate hosts for *Yersinia pestis*
904 during medieval plague epidemics in Northern Europe. *J. Archaeol. Sci.* **40**, 1752–1759 (2013).
- 905 34. Dean, K. R. *et al.* Human ectoparasites and the spread of plague in Europe during the Second
906 Pandemic. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 1304–1309 (2018).
- 907 35. Sloane, B. *The Black Death in London*. (The History Press, 2011).
- 908 36. Hardy, A. The Under-Appreciated Rodent: Harbingers of Plague From the Middle Ages to the
909 Twenty-First Century. *J. Interdiscip. Hist.* **50**, 171–185 (2019).
- 910 37. White, L. A. & Mordechai, L. Modeling the Justinianic Plague: Comparing hypothesized
911 transmission routes. *PLoS One* **15**, e0231256 (2020).
- 912 38. O'Connor, T. P. On the lack of bones of the ship rat *Rattus rattus* from Dark Age York. *J. Zool.*
913 **224**, 318–320 (1991).
- 914 39. Rielly, K. The black rat. in *Extinctions and invasions: a social history of British fauna* (eds.
915 O'Connor, T. & Sykes, N.) 134–145 (Oxbow, 2010).
- 916 40. Salvadori, F. The transition from late antiquity to early Middle Ages in Italy. A
917 zooarchaeological perspective. *Quat. Int.* (2018) doi:10.1016/j.quaint.2018.06.040.
- 918 41. De Cupere, B. *et al.* Eagle owl (*Bubo bubo*) pellets from Roman Sagalassos (SW Turkey):
919 distinguishing the prey remains from nest and roost sites. *Int. J. Osteoarchaeol.* **19**, 1–22 (2009).
- 920 42. Parfitt, S. A. The small mammals. in *The transition to late antiquity on the Danube and beyond*
921 (ed. Poulter, A. G.) 198–318 (Oxford University Press, 2007).
- 922 43. Davis, E. M. Paleoecological studies at Stobi. in *Studies in the antiquities of Stobi* (eds.
923 Aleksova, B., Wiseman, J. & Mano-Zisi, D.) vol. 3 (Princeton University Press, 1981).
- 924 44. Baron, H., Reuter, A. E. & Marković, N. Rethinking ruralization in terms of resilience:

- 925 Subsistence strategies in sixth-century Caričin Grad in the light of plant and animal bone finds.
- 926 *Quat. Int.* **499**, 112–128 (2019).
- 927 45. Benedictow, O. J. *What Disease was Plague?: On the Controversy Over the Microbiological*
- 928 *Identity of Plague Epidemics of the Past.* (Brill, 2010).
- 929 46. O'Connor, T. P. *Bones from Anglo-Scandinavian levels at 16-22 Coppergate.* (Council for
- 930 British Archaeology, 1989).
- 931 47. Pasda, K. *Tierknochen als Spiegel sozialer Verhältnisse im 8.-15. Jh. in Bayern.*
- 932 (Praehistoricaverlag, 2004).
- 933 48. Dobney, K., Jacques, D., Barrett, J. & Johnstone, C. *Farmers, Monks and Aristocrats: the*
- 934 *environmental archaeology of Anglo-Saxon Flixborough.* (Oxbow, 2007).
- 935 49. Reichstein, H. *Die wildlebenden Säugetiere von Haithabu: Ausgrabungen 1966-1969 und 1979-*
- 936 *1980.* (Karl Wachholtz, 1991).
- 937 50. Wigh, B. *Animal husbandry in the Viking Age town of Birka and its hinterland: excavations in*
- 938 *the black earth 1990-95.* (Riksantikvarieämbetet, 2001).
- 939 51. Tourunen, A. *Animals in an Urban Context - A Zooarchaeological study of the Medieval and*
- 940 *Post-Medieval town of Turku.* (University of Turku, 2008).
- 941 52. Eryvnyck, A. *Archeozoölogisch onderzoek van de zwarte rat (Rattus rattus) en de bruine rat*
- 942 *(Rattus norvegicus).* (Amsterdam, 1989).
- 943 53. Barrett-Hamilton, G. & Hinton, M. *A history of British mammals.* (Gurney and Jackson, 1910).
- 944 54. Mitchell-Jones, A. J. *The atlas of European mammals.* (Academic, 1999).
- 945 55. Savinetsky, A. B. & Krylovich, O. A. On the history of the spread of the black rat (*Rattus rattus*
- 946 *L.*, 1758) in northwestern Russia. *Biol. Bull. Russ. Acad. Sci.* **38**, 203–207 (2011).
- 947 56. Ward-Perkins, B. *The Fall of Rome and the End of Civilization.* (Oxford University Press, 2005).
- 948 57. Wickham, C. *Framing the Early Middle Ages: Europe and the Mediterranean, 400-800.* (Oxford
- 949 University Press, 2005).
- 950 58. Horden, P. & Purcell, N. *The Corrupting Sea: A Study of Mediterranean History.* (Wiley-
- 951 Blackwell, 2000).
- 952 59. Ambrosiani, B. Osten und Westen im Ostseehandel zur Wikingerzeit. in *Haithabu und die frühe*

953 *Stadtentwicklung im nördlichen Europa*. (eds. Brandt, K., Müller-Wille, M. & Radtke, C.) 339–
954 348 (Wachholtz, 2002).

955 60. Hodges, R. *Dark Age Economics: A New Audit*. (Bloomsbury, 2012).

956 61. Keller, M. *et al.* Ancient *Yersinia pestis* genomes from across Western Europe reveal early
957 diversification during the First Pandemic (541–750). *Proceedings of the National Academy of*
958 *Sciences* vol. 116 12363–12372 (2019).

959 62. Wagner, D. M. *et al.* *Yersinia pestis* and the Plague of Justinian 541–543 AD: a genomic
960 analysis. *The Lancet Infectious Diseases* vol. 14 319–326 (2014).

961 63. Feldman, M. *et al.* A High-Coverage *Yersinia pestis* Genome from a Sixth-Century Justinianic
962 Plague Victim. *Mol. Biol. Evol.* **33**, 2911–2923 (2016).

963 64. Putnam, N. H. *et al.* Chromosome-scale shotgun assembly using an in vitro method for long-
964 range linkage. *Genome Research* vol. 26 342–350 (2016).

965 65. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M. BUSCO:
966 assessing genome assembly and annotation completeness with single-copy orthologs.
967 *Bioinformatics* vol. 31 3210–3212 (2015).

968 66. Li, H. & Durbin, R. Inference of human population history from individual whole-genome
969 sequences. *Nature* **475**, 493–496 (2011).

970 67. Deinum, E. E. *et al.* Recent Evolution in *Rattus norvegicus* Is Shaped by Declining Effective
971 Population Size. *Mol. Biol. Evol.* **32**, 2547–2558 (2015).

972 68. Gronau, I., Hubisz, M. J., Gulko, B., Danko, C. G. & Siepel, A. Bayesian inference of ancient
973 human demography from individual genome sequences. *Nat. Genet.* **43**, 1031–1034 (2011).

974 69. Colangelo, P. *et al.* Mitochondrial phylogeography of the black rat supports a single invasion of
975 the western Mediterranean basin. *Biological Invasions* vol. 17 1859–1868 (2015).

976 70. Etougbéché, J. *et al.* Genetic diversity and origins of invasive black rats (*Rattus rattus*) in Benin,
977 West Africa. *fozo. I* **69**, 20014.1 (2020).

978 71. Hemamali, P. P. C. & Boyagoda, S. H. Historic black rat invasions into Sri Lanka lead to
979 hybridization forming two sub-lineages in the *Rattus rattus* species complex. *Ceylon Journal of*
980 *Science* vol. 49 433 (2020).

- 981 72. Carleton, G. G., Musserand, M. D. & Musser, G. G. Superfamily Muroidea. in *Mammal species*
982 *of the world: a taxonomic and geographic reference (3rd ed)*. (eds. Wilson, D. E. & Reeder, D.
983 M.) 894–1531 (Johns Hopkins University Press, 2005).
- 984 73. Boivin, N. Proto-globalisation and biotic exchange in the old world. in *Human dispersal and*
985 *species movement from prehistory to the present* (eds. Boivin, N., Crassard, R. & Petraglia, M.)
986 349–408 (Cambridge University Press, 2017).
- 987 74. Cooper, J. P. Egypt’s Nile-Red Sea canals: chronology, location, seasonality and function. in *Red*
988 *Sea IV: Connected Hinterlands* (eds. Blue, L., Cooper, J., Thomas, R. & Whitewright, J.) 195–
989 210 (Archaeopress, 2009).
- 990 75. Prendergast, M. E. *et al.* Reconstructing Asian faunal introductions to eastern Africa from multi-
991 proxy biomolecular and archaeological datasets. *PLoS One* **12**, e0182565 (2017).
- 992 76. Tollenaere, C. *et al.* Phylogeography of the introduced species *Rattus rattus* in the western Indian
993 Ocean, with special emphasis on the colonization history of Madagascar. *J. Biogeogr.* **37**, 398–
994 410 (2010).
- 995 77. Cobb, M. The Chronology of Roman Trade in the Indian Ocean from Augustus to Early Third
996 Century ce. *J. Econ. Soc. Hist. Orient* **58**, 362–418 (2015).
- 997 78. Ewer, R. F. The biology and behaviour of a free-living population of black rats (*Rattus rattus*).
998 *Animal Behaviour Monographs* **4**, 127–174 (1971).
- 999 79. Pocock, M. J. O., Hauffe, H. C. & Searle, J. B. Dispersal in house mice. *Biol. J. Linn. Soc. Lond.*
1000 **84**, 565–583 (2005).
- 1001 80. Büntgen, U. *et al.* Cooling and societal change during the Late Antique Little Ice Age from 536
1002 to around 660 AD. *Nat. Geosci.* **9**, 231–236 (2016).
- 1003 81. Perry, R. D. & Fetherston, J. D. *Yersinia pestis*--etiologic agent of plague. *Clin. Microbiol. Rev.*
1004 **10**, 35–66 (1997).
- 1005 82. Spyrou, M. A., Bos, K. I., Herbig, A. & Krause, J. Ancient pathogen genomics as an emerging
1006 tool for infectious disease research. *Nat. Rev. Genet.* **20**, 323–340 (2019).
- 1007 83. Fenwick, C. From Africa to Ifrīqiya: Settlement and Society in Early Medieval North Africa
1008 (650–800). *Al-Masāq* **25**, 9–33 (2013).

- 1009 84. Pickrell, J. K. & Pritchard, J. K. Inference of population splits and mixtures from genome-wide
1010 allele frequency data. *PLoS Genet.* **8**, e1002967 (2012).
- 1011 85. McCormick, M. Where do trading towns come from? Early medieval Venice and the northern
1012 emporia. *Post-Roman towns, trade and settlement in Europe and Byzantium* **1**, 41–68 (2007).
- 1013 86. Smith, D. & Kenward, H. Roman Grain Pests in Britain: Implications for Grain Supply and
1014 Agricultural Production. *Britannia* **42**, 243–262 (2011).
- 1015 87. Buffon, G.-L. L. C. de. *Histoire Naturelle, Générale et Particulière, avec la Description du*
1016 *Cabinet du Roi*. vol. 8 (L’Imprimerie Royale, 1760).
- 1017 88. Smith, R. *The Universal Directory for Taking Alive and Destroying Rats, and All Other Kinds of*
1018 *Four-footed and Winged Vermin, In a Method Hitherto Unattempted: Calculated for the Use of*
1019 *the Gentleman, the Farmer, and the Warrener*. (J. Walker, 1768).
- 1020 89. Pennant, T. *British Zoology*. (William Eyres, 1776).
- 1021 90. Ruttly, J. *An essay towards a natural history of the County of Dublin*. vol. 1 (W. Sleater, 1772).
- 1022 91. O’Connor, T. Commensal Species. in *The Oxford Handbook of Animal Studies* (ed. Kalof, L.)
1023 (Oxford University Press, 2017).
- 1024 92. Walker, T., Sharpe, J. R. & Williams, H. Barn Owls and Black Rats from a Rural Roman Villa at
1025 Gatehampton, South Oxfordshire. *Environ. Archaeol.* 1–10 (2019).
- 1026 93. Bronk Ramsey, C. *OxCal*. (2020).
- 1027 94. Reimer, P. J., Austin, W. E. N., Bard, E. & Bayliss, A. The IntCal20 northern hemisphere
1028 radiocarbon age calibration curve (0–55 cal kBP). (2020).
- 1029 95. Guiry, E. & Buckley, M. Urban rats have less variable, higher protein diets. *Proc. R. Soc. B* **285**,
1030 20181441 (2018).
- 1031 96. Dewar, G. & Pfeiffer, S. Approaches to Estimating Marine Protein in Human Collagen for
1032 Radiocarbon Date Calibration. *Radiocarbon* **52**, 1611–1625 (2010).
- 1033 97. Heaton, T. J. *et al.* Marine20—The Marine Radiocarbon Age Calibration Curve (0–55,000 cal
1034 BP). *Radiocarbon* **62**, 779–820 (2020).
- 1035 98. Chapman, J. A. *et al.* Meraculous: de novo genome assembly with short paired-end reads. *PLoS*
1036 *One* **6**, e23501 (2011).

99. Smit, AFA, Hubley, R & Green, P. RepeatMasker Open-4.0. <http://www.repeatmasker.org> (2013-2015).
100. Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Research* vol. 27 573–580 (1999).
101. Kurtz, S. *et al.* Versatile and open software for comparing large genomes. *Genome Biol.* **5**, R12 (2004).
102. Drummond, A. J. *et al.* Geneious v6. 1.6. (2011).
103. Dabney, J. *et al.* Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proceedings of the National Academy of Sciences* vol. 110 15758–15763 (2013).
104. Damgaard, P. B. *et al.* Improving access to endogenous DNA in ancient bones and teeth. *Sci. Rep.* **5**, 11184 (2015).
105. Yang, D. Y., Eng, B., Wayne, J. S., Dудар, J. C. & Saunders, S. R. Technical note: improved DNA extraction from ancient bones using silica-based spin columns. *Am. J. Phys. Anthropol.* **105**, 539–543 (1998).
106. Speller, C. F. *et al.* Ancient mitochondrial DNA analysis reveals complexity of indigenous North American turkey domestication. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 2807–2812 (2010).
107. Rohland, N., Harney, E., Mallick, S., Nordenfelt, S. & Reich, D. Partial uracil–DNA–glycosylase treatment for screening of ancient DNA. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **370**, 20130624 (2015).
108. Gansauge, M.-T., Aximu-Petri, A., Nagel, S. & Meyer, M. Manual and automated preparation of single-stranded DNA libraries for the sequencing of DNA from ancient biological remains and other sources of highly degraded DNA. *Nat. Protoc.* (2020) doi:10.1038/s41596-020-0338-0.
109. Kircher, M., Sawyer, S. & Meyer, M. Double indexing overcomes inaccuracies in multiplex sequencing on the Illumina platform. *Nucleic Acids Res.* **40**, e3 (2012).
110. Carøe, C. *et al.* Single tube library preparation for degraded DNA. *Methods Ecol. Evol.* **9**, 410–419 (2018).
111. Peltzer, A. *et al.* EAGER: efficient ancient genome reconstruction. *Genome Biol.* **17**, 60 (2016).

112. Schubert, M., Lindgreen, S. & Orlando, L. AdapterRemoval v2: rapid adapter trimming, identification, and read merging. *BMC Res. Notes* **9**, 88 (2016).
113. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).
114. Jónsson, H., Ginolhac, A., Schubert, M., Johnson, P. L. F. & Orlando, L. mapDamage2.0: fast approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics* **29**, 1682–1684 (2013).
115. Jun, G., Wing, M. K., Abecasis, G. R. & Kang, H. M. An efficient and scalable analysis framework for variant extraction and refinement from population-scale DNA sequence data. *Genome Res.* **25**, 918–925 (2015).
116. Teng, H. *et al.* Population Genomics Reveals Speciation and Introgression between Brown Norway Rats and Their Sibling Species. *Mol. Biol. Evol.* **34**, 2214–2228 (2017).
117. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).
118. Renaud, G., Slon, V., Duggan, A. T. & Kelso, J. Schmutzi: estimation of contamination and endogenous mitochondrial consensus calling for ancient DNA. *Genome Biol.* **16**, 224 (2015).
119. Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of Next Generation Sequencing Data. *BMC Bioinformatics* **15**, 356 (2014).
120. Li, H. Improving SNP discovery by base alignment quality. *Bioinformatics* **27**, 1157–1158 (2011).
121. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).
122. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).
123. Darriba, D., Taboada, G. L., Doallo, R. & Posada, D. jModelTest 2: more models, new heuristics and parallel computing. *Nat. Methods* **9**, 772 (2012).
124. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

- 1093 125. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
1094 improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).
- 1095 126. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage
1096 analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).
- 1097 127. Patterson, N. *et al.* Ancient Admixture in Human History. *Genetics* vol. 192 1065–1093 (2012).
- 1098