

1 Renegade Bacterial Genetic Sequences in a Stealth Adapted Virus: Biological and Diagnostic Implications  
2

3 W. John Martin, MD, PhD

4 Institute of Progressive Medicine  
5 South Pasadena CA 91030

6 [wjohnmartin@ccid.org](mailto:wjohnmartin@ccid.org)

7  
8  
9  
10 Running Title: Renegade Bacterial Sequence in Stealth Adapted Virus  
11  
12

13 Author Mailing Address: 1634 Spruce Street  
14 South Pasadena CA 91030  
15

16 E-Mail: [wjohnmartin@ccid.org](mailto:wjohnmartin@ccid.org)

17  
18 Phone: 01-626-616-2868  
19

20 Author ORCID Number: 0000-0002-9947-4374  
21

22 Acknowledgment: The work was supported by MI Hope Inc., a non-profit public charity.

23

24 Conflicts of Interests: None

25

26

27 Word Count w/o References 5,556

28

29

30 Abstract

31 There are major differences between viruses, bacteria, and eukaryotic cells in the structuring of their genomes, modes of  
32 replication, and capacity to horizontally transfer genetic sequences. DNA sequencing of a cloned African green monkey simian  
33 cytomegalovirus (SCMV) indicate the inclusion of certain bacterial and cellular genetic sequences. The virus was cultured from  
34 a chronic fatigue syndrome (CFS) patient. It is stealth adapted with the loss or mutation of the genes coding for the relatively  
35 few viral components normally targeted by the cellular immune system. This article identifies the likely origins of many of the  
36 bacterial-derived genetic sequences present in this virus. There are multiple clones with close but non-identical sequence  
37 alignments with different genomic regions of the *Ochrobactrum quorumnocens* A44 species of bacteria. Another set of clones  
38 matched most closely to diverse genomic regions of *Mycoplasma fermentans* bacteria. The sequences of several other clones  
39 could only be approximately aligned to those of different types of bacteria. The sequence of clone 3B513 is consistent with  
40 genetic contributions from the genomes of several types of bacteria. The term viteria refers to viruses with bacteria-derived  
41 genetic sequences. Stealth adapted viruses are the likely primary cause of CFS and autism, and possibly act as cofactors in many  
42 illnesses including AIDS. The additional incorporation of bacterial sequences into these viruses can potentially lead to the  
43 mistaken diagnoses of a bacterial chronic illness rather than a viral infection. Stealth adapted virus testing should be performed  
44 in illnesses attributed to *Mycoplasma*, *Borrelia*, and *Streptococcal* infections.

45  
46 Introduction

47 Molecular analysis of cloned DNA derived from viral cultures from a patient with the chronic fatigue syndrome (CFS)  
48 showed that the cultured virus had originated from an African green monkey simian cytomegalovirus (SCMV) [1-4 ]. Yet,  
49 genetic sequences corresponding to major regions of the SCMV genome were not detected within any of the sequenced DNA  
50 clones [4-5]. Moreover, there was an uneven distribution of the clones with regards to the remaining identified regions of  
51 SCMV, with genetic variability within clones that match to the same region of the SCMV genome. These finding are consistent  
52 with an immune escape mechanism, referred to as stealth adaptation, occurring from the deletion or mutation of the genes  
53 coding the relatively few virus components that are normally targeted by the cellular immune system [1, 6]. In addition to the  
54 SCMV-derived genetic sequences, there are certain clones with genetic sequences that have come from portions of cellular and  
55 bacterial genomes [7-13]. These incorporated cellular and bacterial renegade sequences may be required for the virus to regain  
56 its infectivity and may also contribute to the virus-mediated cytopathic effect (CPE).

58        The increasing availability of DNA sequence data in GenBank has allowed further elucidation of the origins of bacterial-  
59        derived genetic sequences present in this virus. The findings are consistent with genetic contributions to the virus genome not  
60        only from intracellular growing bacteria, such as *Mycoplasma fermentans* [14], but also from soil-based bacteria including  
61        members of the *Brucella – Ochrobactrum* Family within the *Hyphomicrobiales* Order [15]. The findings have major biological  
62        implications and are relevant to the potential of misdiagnosing stealth adapted virus infections as being caused by bacteria with  
63        patients being inappropriately treated with antibiotics.

64  
65        Materials and Methods

66  
67        Patient and Virus Culture: The virus was repeatedly cultured from a woman who was hospitalized in 1990 with a provisional  
68        diagnosis of either encephalitis or meningitis developing a week after she had a sore throat [1]. A cerebrospinal fluid (CSF)  
69        sample obtained during her hospital admission showed no cells and normal protein level. She failed to regain her prior level of  
70        health and vigor but felt cognitively impaired, fatigued, and lacked the capacity for restorative sleep. She had earlier rented an  
71        apartment to a HIV positive individual with similar symptoms and considered the possibility of a non-sexually transmitted  
72        illness. The woman provided a blood sample as part of a study using the polymerase chain reaction (PCR) to look for viruses in  
73        patients with CFS. Based on a clearly positive PCR result, an additional blood sample was obtained for virus culturing. A  
74        cytopathic effect (CPE) was observed in primary human foreskin fibroblasts (MRHF). It comprised the formation of foamy  
75        vacuolated cells with marked syncytia. Subsequently obtained blood samples confirmed the presence of an infectious cytopathic  
76        agent that could be passaged in cultured cells. Abundant viral particles were seen by electron microscopy. The cultured cells and  
77        cell free supernatants yielded strongly positive PCR assays with well-defined PCR products shown in agarose gel  
78        electrophoresis. Radiolabeled PCR products hybridized with material that was pelleted by ultracentrifugation of 0.45  $\mu$  filtered  
79        culture supernatant. As will be shown, DNA extracted from this material migrated in agarose gel electrophoresis as a well-  
80        defined band with an estimated size of approximately 20 kilobase (kb).

81  
82        Cloning of Virus DNA: In the first set of cloning experiments, the DNA obtained from the filtered and ultracentrifuged culture  
83        supernatant was digested overnight with 10U/ $\mu$ l of EcoRI enzyme and cloned into pBluescript plasmids [1]. The plasmids were  
84        propagated in XL-1 bacteria and those with inserts were either partially or fully sequenced. This series of over 180 clones was  
85        labeled as the 3B series. In a later set of cloning experiments, the nucleic acids extracted from the filtered and ultracentrifuged

86 culture supernatant were further purified by agarose gel electrophoresis. The approximately 20 kb band was excised, and the  
87 DNA was reextracted, and then digested overnight with 2U/ $\mu$ l SacI restriction enzyme [5]. The digested DNA was cloned into  
88 pBluescript plasmids and propagated in XL-1 bacteria. The resulting series of over 120 clones was labeled as the C16 series.  
89 The individual clones of stealth virus-1 that are referred to in this article, together with their NCBI Accession number and  
90 nucleotide length are listed in Table 1. Separate Accession numbers corresponding to the separate sequences derived from T3  
91 and T7 polymerase primer sites of the pBluescript plasmids are provided for those clones that were not completely sequenced.

92 Sequencing of Clones: Preliminary sequencing of the clones was performed from both the T3 and the T7 primer sites. The initial  
93 sequencing was performed by either Bio Serve Biotechnologies, Laurus MD or by the City of Hope Cancer Center Molecular  
94 Core Facility, Duarte CA. Sequenase was used at the first location and generally yielded unambiguous 100-200 nucleotide  
95 sequences. Thermal cycling was used at the second facility and would commonly yield over 500 nucleotides, but with increasing  
96 numbers of undefined nucleotides indicated as ‘N’; in the longer stretches. Only the T3 and T7 sequences are available for many  
97 of the entire 3B and C16 series of clones. Additional double stranded complete sequencing was, however, obtained for most of  
98 the clones discussed in this article. The extended sequencing was provided by Lark Technology, Houston, Texas and by U.S.  
99 Biochemical Corp. Cleveland, Ohio. All DNA sequence data on the clones have been submitted to GenBank under the heading  
00 of stealth virus-1.

01 DNA Sequence Analysis: The sequences of the clones were analyzed using the online BLASTN and BLASTX programs  
02 provided by the National Center for Biotechnology Information (NCBI) [16]. The BLASTN sequence matching and alignment  
03 program compares the input (“query”) DNA sequence with the nt/nr non-redundant DNA “subject” sequences currently  
04 available in the GenBank repository. The BLASTX program compares the sequences of each of the six potentially derived  
05 amino acid sequences coded by the query input sequence with the known protein subject sequences on GenBank. When an  
06 amino acid match is identified, the coding nucleotide sequence of the known protein can be used in a pairwise BLASTN  
07 comparison with the query sequence. The designated name, NCBI accession number, and the range of nucleotide numbers in the  
08 identified GenBank subject sequences are displayed in the BLASTN results. If the input sequence is already in GenBank, this  
09 sequence will be the first selected sequence, followed by the sequences with increasing nucleotide disparities till the set limit of  
10 identified matching sequences is met. The BLASTN program further tests if the placing of one or more gaps in the input and  
11 selected sequences, or if separating the sequences into two or more segments, will improve the overall alignment. The “bit

12 score" is a measure of the non-random chance alignment of two sequences. The number of inserted gaps to obtain optimal  
13 alignment is also shown in the results, along with the numbers and ratio of identical nucleotides. The statistical probability  
14 against a random alignment is more clearly reflected in the Expect Value. This is expressed as the negative exponential value to  
15 the log base "e." The smaller the "e" value, i.e., the higher the negative log value, then the less likelihood of purely random  
16 matching. When the e value is less than e-180, it is recorded as 0.0. This reflects that the sequences are very similar over  
17 sufficiently long regions. There can be multiple GenBank sequences that show close matching to only a portion of the query  
18 sequence. This can potentially reduce the sensitivity of showing less well aligned sequences. To avoid this, the BLASTN  
19 analysis was typically repeated after excluding the highly matched regions.

20  
21 **RESULTS**

22  
23  
24 A previously published Figure is included in this article as Figure 1. It shows an ethidium bromide-stained agarose gel  
25 with HindIII and BetE-II enzyme digested Lambda DNA phage size markers in the left and right outside lanes, respectively. The  
26 added arrow indicates the clearly seen band of a portion of the DNA extracted from the material that was pelleted by  
27 ultracentrifugation of filtered cultured supernatant. In comparison with the DNA size markers, the band has DNA of  
28 approximately 20 kb. The lane directly beneath the arrow shows several smaller DNA bands, which resulted from EcoRI  
29 digestion of another portion of the pelleted and extracted DNA. The lower stained material in the lanes of the supernatant  
30 extracted material is RNA. The remaining lane is an EcoR1 digest of nucleic acids extracted from lysed infected cells of the  
31 culture from which the supernatant was obtained.

32  
33  
34 **Clones With Sequences Matching to *Brucella* - *Ochrobactrum* Bacteria**

35 BLASTN analyses of the DNA sequences in eight of the clones show that the highest levels of non-self-matching to DNA  
36 sequences are to distinct regions of Chromosome 1 of *Ochrobactrum quorumnocens* strain A44 [17; NCBI accession number  
37 [CP022604.1](https://www.ncbi.nlm.nih.gov/nuccore/CP022604.1)]. The clones are listed in Table 2 in the order of the regions of the matching nucleotides along chromosome 1 of the  
38 bacterium. These regions code for a variety of functional proteins. Each of the eight clones had been fully sequenced. The  
39 overall ratio of identical nucleotides (nt) in the eight clones to the matched regions in chromosome 1 of *O. quorumnocens* strain

40 A44 is 99.7% (31,977/32,064 nt). Except for the 11 gaps for clone C16116, there is on average only 1 gap per clone for the other  
41 7 clones.

42 There are three other clones that were only partially sequenced from the T3 and T7 promoter sites on the pBluescript  
43 plasmid. For two of these clones (C16118 and 3B315) both ends of the clones matched to regions within Chromosome 2 of *O.*  
44 *quorumnocens* strain A44 (NCBI accession number CP022603.1). The T3-derived sequence for plasmid C16282 also matched  
45 to a region of chromosome 2 of *O. quorumnocens* strain A44. The T7-derived sequence, however, matched to a genomic region  
46 of SCMV. The matching to SCMV strain 2715 (NCBI accession number [FJ483968.2](#)) extended from nucleotide 84420 and  
47 84692 with an Expect Value of 1e-63, and 192/248 identical nucleotides after excluding unmatchable “N” nucleotides in the  
48 clone. A small fully sequenced clone (3B540) also matched preferable to a region in chromosome 2 of *O. quorumnocens* strain  
49 A44. There were 205/226 identical nucleotides, although the optimal matching required 15 gaps.

50 There is an additional fully sequenced clone (3B43) with 3620 nucleotides. BLASTN identified the statistically higher  
51 overall match to the ribosomal genetic sequences in *Brucella BTU1*, *Brucella pituitosa* strain AA2, and *Brucella*  
52 *pseudogrignonensis* strains. Note that the term *Brucella* in referring to these later two strains is a synomon for *Ochrobactrum*.  
53 There was less matching to *O. quorumnocens* strain A44 genome. The lower matching occurred because the BLASTN program  
54 matched the overall sequence of clone 3B43 to the *Brucella* species, but to two separated ribosomal sequences in the *O.*  
55 *quorumnocens* strain A44 genome. A 103 nucleotide long stretch of the 3B43 clone from nucleotide 2301 to nucleotide 2404  
56 was present at a 99-100% identity in the earlier mentioned *Brucella* strains, but not in the *O. quorumnocens* strain A44. The  
57 missing sequence corresponds to part of the ribosomal gene complex that intervenes between the 16S and the 28S coding  
58 regions. Intervening nucleotides are present in some but not of the bacteria within the different bacterial families [18].  
59 Sequences in the two regions of clone 3B43 that are apart from the intervening region match most closely to those in  
60 Chromosome 2 of *O. quorumnocens* strain A44. These sequences were, therefore, chosen to be included in of Table 2.

61 The initial BLASTN results with clone 3B562 did not include any *Ochrobactrum* species. Rather, it identified a  
62 sequence in *Serratia marcescens* bacteria with 88% nucleotide identity. Yet, the highest matching to its BLASTX translated  
63 amino acid sequence was to an *O. quorumnocens* coded protein. The 310-nucleotide sequence coding the matching region of the  
64 *O. quorumnocens* protein is identical to the sequence of clone 3B562. For convenience, this entity has been added to Table 2,  
65 even though it could not be specifically identified in the published sequences of either chromosome 1 or 2 of *O. quorumnocens*.

66 There are sequences in two clones, 3B41 and 3B47, that do not closely align with sequences in either chromosome 1 or 2  
67 of *O. quorumnocens* strain A44 genome but do align better to sequences in chromosome 2 of *Brucella pseudogrignonensis* [19].

68 Even though the Expect Value was still 0.0, the total identical matching of the two clones was only 77.1% (3779/4901) with a  
69 total of 82 gaps (Table 3).

70

71

72 **Clones Matching to Mycoplasma Bacteria**

73 There are ten 3B series clones with sequences that best match to separate sequences within both *Mycoplasma fermentans*  
74 and *Mycoplasma conjunctivae* strain NCTC10147. The complete genome of this later species of mycoplasma has yet to be fully  
75 assembled. The matching stealth virus clones are listed in Table 4 in the order of their sequence alignments with increasing  
76 numbers of the matching *M. fermentans* nucleotides. For one of the partially sequenced clones (3B680), while its T7 sequence  
77 matched closely to *M. fermentans* and *M. conjunctivae*, 79 of the first 85 of 100 nucleotides at its T3 end matched to a sequence  
78 in SCMV. The matching to SCMV strain 2715 (NCBI accession number FJ483968.2) extended from nucleotide 202627 to  
79 202721 of the SCMV genome with an Expect Value of only 2e-25 due in part to its small size. The 85 nucleotides in the T3  
80 sequence of clone 3B680 also matched to 85 nucleotides within two clones 3B418 and 3B625 that matched at both of their ends  
81 with sequences of SCMV. The overall level of matching of the sequences of the ten clones with the sequences of both *M.*  
82 *fermentans* and *M. conjunctivae* is 97.5% (13,931/14,287 nucleotides)

83

84 **Clones Preferably Matching to Other Bacteria**

85 The following are the best currently available alignments of the sequences in some of the remaining clones. For example,  
86 there are three clones (C16122, C16127, and C16141) in which with DNA sequences best match to different species of  
87 *microbacterium* bacteria. These are small, gram-positive bacteria with a high guanosine-cytosine content that are classified  
88 within the *Actinomycetia* class of bacteria [20]. The matching of the T3 and T7 sequences from clone C16122 was to different  
89 species of this bacteria. The T3 sequence matching clone C16122 T7 was reasonably high with an Expect Value of 2e-102.  
90 However, it was only the region between nucleotides 162 to 505 of the 812 long nucleotide sequence of the clone that matched  
91 to a *microbacterium* sequence. The regions of C16122 T7 from nucleotide 1 to 161 and 505-812 showed no matching with any  
92 of the accessible GenBank sequences, even when using the BLASTX program. The e-120 and e-97 Expect Values, respectively,  
93 of the sequence alignments for clones C16127 and C16141 to *microbacterium* species of bacteria are reasonable in view of the  
94 relatively short lengths of the sequences. The data on the three clones are recorded in Table 4.

95 The T3 and T7 sequences of clone (C16125) show preferred matching to a 157689 base pair (bp) plasmid isolated from  
96 the MDW-2 unclassified species of Aminobacter bacteria (Table 5). Aminobacter bacteria are  $\alpha$ -proteobacteria in the same  
97 *Hyphomicrobiales* Order as the *Brucella* – *Ochrobactrum* bacteria but belong to a different family. One clone (C16135) has a  
98 T3 sequence that best matches to a sequence within a strain of *Cellulosimicrobium cellulans* bacteria, which is also within the  
99 *Actinomycetia* class of bacteria [20]. The T7 sequence of clone C16135, however, matches to an intron sequence within the  
00 Karzin coding human cellular gene (NCBI Accession no. NG\_029844.2) with 205/396 nucleotide identity after discounting the  
01 non-assigned “N” nucleotides.

02 Clone 3B513 (NCBI Accession no. U27894.2) is a fully sequenced clone of 8,106 nucleotides. Routine BLASTN  
03 matching identified only eight small, regions that separately matched to known bacterial sequences. Collectively, these regions  
04 comprised only 3,361 of the 8,106 nucleotides. Only one of the matches had an Expect Value of 0.0. Far more extensive  
05 sequence matching was obtained using the BLASTX and pairwise matching of the sequence of 3B513 with the nucleotides  
06 coding the best matching amino acid sequence. The results of this analysis are summarized in Table 6. There was no uniform  
07 genus of the bacteria with matching sequences. Rather the identified bacteria were either within the *Rhizobiaceae* or  
08 *Phyllobacteriaceae* family, within the *Hyphomicrobiales* Order. The sequences of the *Shinella oryzae* and *Rhizobium flavin*  
09 strains remain separated along the 3B513 sequence. By contrast, there are several hundred overlapping nucleotides shared by the  
10 end regions of the *Aminobacter niigataensi* strain and *Ochrobactrum POC9* strain sequences. Similarly, there are shared  
11 nucleotide sequences between the other end of *Ochrobactrum POC9* strain sequences with the end region of the *Mesorhizobium*  
12 *denitrificans*-related sequence.

13 BLASTN analysis of clone 3B513 showed matching with a portion of the 3B525 T3 sequence extending from nucleotide  
14 76 to 593. The matching region was within the section of the 3B513 clone that is related to a sequence in *M. denitrificans*.  
15 Nucleotides 15 to 82 on clone 3B525 T3 matched to SCMV (NCBI Accession [FJ483968.2](#) with an Expect Value of 2e-09 and  
16 52/64 identical nucleotides after omitting 4 non-assigned “N” nucleotides. The matching SCMV nucleotides were from 72,686  
17 to 72,619. The sequence of clone 3B525 T7 also matches to SCMV nucleotides 66002 to 66651 with an Expect score of 0.0.  
18 Over the first 500 nucleotides, there are 473 identical nucleotides between the SCMV and 3B525 T7 sequences. Moreover, the  
19 3B527 T7 sequences closely matches the sequences clones 3B526T7, 3B550 T7, 3B544 T7, 3B320 T3, 3B642 T7, 3B314 T3,  
20 3B663 T3 all of which have SCMV-related sequences in both their T3 and T7 sequences (data not shown).  
21  
22

## 23 Discussion

24 The presence of bacterial sequences in the cultures of a stealth adapted virus cannot be explained by bacterial  
25 contamination of the cultures. This possibility is excluded by the following observations: i) Repeated blood cultures from the  
26 patient over a 4-year period gave very comparable results in terms of the observed formation of foamy vacuolated cells with  
27 prominent syncytia. ii) There were no indications in any of the cultures of bacteria being present in the frequent viewing of  
28 living cells by phase contrast microscopy; the examination of hematoxylin and eosin-stained cells by regular microscopy; or by  
29 electron microscopy. iii) Some of the cultures were maintained in both serum-free and antibiotic free medium for extended time  
30 periods. iv) The identified sequences are not from the bacteria that are typically involved in bacterial contamination. v) The  
31 agarose gel of the nucleic acids extracted from the pelleted material obtained from filtered and ultracentrifuged tissue culture  
32 medium show only minimal DNA that is larger than the approximately 20 kb band and, which would have been expected if  
33 there were bacterial chromosomal DNA. vi) The C16 series of clones were derived solely from the DNA that banded in the  
34 agarose gel at a size of approximately 20 kb.

35 The relocation or transposition of genetic sequences from their bacterial origin to a virus can be viewed as a passive  
36 hijacking of the sequences by the virus or as the loss of the bacteria's capacity to restrain some of their own genetic sequences  
37 from deserting and moving elsewhere. To help in understanding this process, the relocating sequences are referred to as being  
38 "renegade sequences" [9]. The term "veteria" was also introduced to describe viruses with incorporated bacteria-derived genetic  
39 sequences [8].

40 Even though the double stranded DNA pelleted material migrated in agarose gel with an approximate size of 20 kb, the  
41 sum of the previously reported SCMV-derived nucleotide sequences (~100,000), cellular-derived sequences (~ 25,000) and the  
42 presently reported bacterial related sequences (~ 50,000) indicates that the entire genome comprises genetically different  
43 segments with a diversity of viral, cellular, and bacterial-derived sequences. These segments are not necessary all packaged into  
44 each virus particle, which did appear to be morphologically heterogeneous on electron microscopy [1]. It is of interest that the  
45 agarose gel of the nucleic acids extracted from the pelleted material showed a substantial amount of RNA (Figure 1). A 20 kb  
46 size of the double-stranded DNA is consistent with RNA being involved in the replication process. Thus, 20 kb is close to the  
47 upper size limit of replicating RNA without the inclusion of an intrinsic genomic proof-reading mechanism [21]. Involvement of  
48 RNA may also explain the genetic instability previously reported in the SCMV and cellular related DNA genetic sequences [3].  
49 It is also consistent with a reverse transcriptase step being required to obtain a positive PCR in a stealth adapted virus culture  
50 from a different CFS patient [22].

51 The mechanism of the apparent incorporation of bacteria-derived sequences is probably the same as that occurring with  
52 the cellular-derived sequences in this and in other stealth adapted viruses [ 9,10]. It is perceived as single stranded RNA cross-  
53 linking residual DNA or RNA segments that are remaining from the originating SCMV virus after it has undergone  
54 fragmentation as part of the stealth adaptation process. Further fragmentation of the virus could lead to additionally incorporated  
55 cross-linked genetic sequences. The finding of bacterial and SCMV related sequences at the opposite terminal regions of clones  
56 C16282, 3B680, and 3B525 is consistent with this hypothesis. This possibility is further supported by the limited SCMV  
57 matching of nucleotides 15 to 82 on clone 3B525. This region slightly overlaps with the bacterial matching sequences from  
58 nucleotides 76 to 593 in the same clone. Also consistent with a cross-linking process is the presence of a bacterial sequence and  
59 of a cellular sequence at the T3 and T7 readouts, respectively, of clone C16135.

60 Another example of overlapping sequences is seen within clone 3B513. The two end regions of the *Ochrobactrum*-  
61 matching sequences in clone 3B513 overlap with the end regions of *Aminobacter* and of *Mesorhizobium* bacterial sequences,  
62 respectively. Moreover, as mentioned above, a region within the *Mesorhizobium*-related sequence matchers to a region within  
63 clone 3B525 T3. In addition to sequences linking to each other, homologous recombination can lead to the substitution of  
64 sequences with matching end regions (9). It is still possible, however, that the predominant viral, bacterial, and cellular  
65 sequences are mainly located on discrete segments of the stealth adapted virus. This question can be resolved by further  
66 sequencing of the virus, which is archived at the American Type Culture Collection (ATCC).

67 The potential functions of the incorporated bacterial genetic sequences will be discussed in more detail in a subsequent  
68 article. At least some of the coded proteins are presumably contributing to the replication and transmission of the virus. One  
69 such role could be providing alternative or additive capsid-like proteins [13]. Some of the proteins are enzymes and have  
70 potentially useful metabolic functions. There is a self-healing process that occurs during the culturing of the stealth adapted  
71 viruses [23]. It is associated with the production of self-assembling aliphatic and aromatic chemical compounds. The assembled  
72 materials are referred to as alternative cellular energy (ACE) pigments [23] These materials can attract an external force  
73 provisionally called KELEA (Kinetic Energy Limiting Electrostatic Attraction), which drives the ACE pathway [24]. Enhancing  
74 the ACE pathway can lead to the suppression of the virus induced CPE [24] It could be, therefore, that some of the bacteria  
75 gene-coded proteins are contributing to the formation of ACE pigments and, thereby, prolonging the infection process.  
76 Similarly, intracellular protein aggregates can trigger the unfolded protein response [25] that can also delay virus-induced cell  
77 death.

78 An intracellular location for interactions between bacterial and viral sequences is easier to envision for those bacteria that  
79 can replicate intracellularly. This applies to infections with *mycoplasma*, *brucella*, and *microbacterium* bacteria [14, 26].  
80 Intracellular growth in mammalian cells is not, however, a known characteristic of *Ochrobactrum* bacteria [26-28]. These  
81 bacteria are mainly viewed as being present in soils and comprising part of the complex rhizosphere surrounding and commonly  
82 penetrating plant cells. Indeed, considerable symbiosis occurs between plant cells and a wide array of endophytic bacteria [29].  
83 Another option is for stealth adapted viruses to directly enter soil-based bacteria, which can be present in consumed uncooked  
84 foods [30]. Some of these ingested bacteria may continue to reside in the gut microbiota. Interestingly, atypical bacteria were  
85 cultured from the feces of the CFS patient from which stealth virus-1 was cultured. Moreover, transmissible cytopathic activity  
86 was subsequently retrieved from the atypical bacterial colonies (unpublished). There are major epidemiological ramifications if  
87 bacteria can be involved in the transmission of stealth adapted viruses.

88 A related characteristic of stealth virus-1 is its wide host range, including being infectious for insect cells [2]. This could  
89 occur through the deletion of genes responsible for the typical species restricted growth of most animal cytomegaloviruses.  
90 Insects also commonly harbor endophytic bacteria [31].

91 *O. quorumnocens* strain A44 was originally isolated in Holland from the rhizosphere of field potatoes [17]. It has a  
92 defining function of metabolically inhibiting *N*-acyl homoserine lactones, a chemical used by certain gram negative bacteria to  
93 establish a more pathogenic quorum that can lead to “soft rot” [17]. Because *O. quorumnocens* strain A44 lacks the 16S-28S  
94 intervening sequence present in clone 3B43, it cannot be regarded as the unequivocal source of the *O. quorumnocens* related  
95 sequences. Moreover, the *B. pseudogrignonensis* related sequences in two of the clones (3B41 and 3B47) are not present in *O.*  
96 *quorumnocens*; nor are the *O. quorumnocens*-related sequences in clone 3B513 directly identifiable with strain A44..

97 Intact *B. pseudogrignonensis* bacteria are noteworthy because of their ability to induce tumors in mushrooms [32]. This is  
98 mentioned to underscore the uncertainty as to the potential biological consequences of the virus mediated transfer into humans  
99 of infectious bacteria-derived genetic sequences. Moreover, the exact origins and biological functions of the incorporated  
00 bacterial sequences may remain in doubt because of their genetic instability with potential ongoing divergence from the original  
01 bacterial sequence. As seen with incorporated cellular sequences, there is also the potential for the substitution between different  
02 bacterial sequences [10].

03 As noted above, it is easier to envision the assimilation of *mycoplasma* sequences into stealth adapted viruses than from  
04 soil-based bacteria. It is particularly noteworthy that the closest alignment is with *M. fermentans*, strain M64. (There was  
05 essentially identical nucleotide matching to *M. conjunctivae*, a sheep eye pathogen [33].) The *M. fermentans* species of

06 mycoplasma gained special interest during the early history of the AIDS epidemic. Although HIV had been isolated from  
07 patients, concerns were expressed that a co-infecting pathogen was also required for the development of severe illness. Dr.  
08 Shyh-Ching Lo identified anti-mycoplasma antibodies and mycoplasma DNA sequences in several AIDS patients with more  
09 fulminant disease. He was eventually able to isolate a culturable mycoplasma, which is called *M. incognitus*. It was later  
10 confirmed as *M. fermentans incognitus* [34-38].

11 Most of the HIV clinical studies, however, relied on serological and molecular testing rather than on isolating culturable  
12 bacteria [34-38]. Indeed, the isolation of an intact *mycoplasma* bacteria could be more of a coincidental and somewhat  
13 misleading distraction from finding a stealth adapted virus with incorporated *M. fermentans* genetic sequences.

14 The clinical association between positive *M. fermentans* assays and severity of HIV illness is relevant to the proposed role  
15 that the testing of an experimental polio vaccine in chimpanzees had in the formation of HIV [39]. The polio vaccine was grown  
16 in Rhesus monkey kidney cell cultures. It was reportedly contaminated with a cytopathic virus that was difficult to culture [40].  
17 It also has detectable DNA of rhesus monkey cytomegalovirus [41]. These two findings are consistent with the vaccine being  
18 contaminated with a stealth adapted virus able to promote the growth and transformation of simian to human immunodeficiency  
19 virus.

20 Several other illnesses, including CFS, Fibromyalgia, Gulf War Syndrome, and chronic arthritis were previously  
21 attributed to ongoing infection with *M. fermentans* bacteria [42-46]. This conclusion was based mainly on positive serology and  
22 molecular detection methods. Clinical trials with long term antibiotic therapy failed to achieve clinical remission and the  
23 presumed bacterial infections were largely dismissed as being coincidental to the real cause. Based on the data presented in this  
24 article, the findings are consistent with a viteria infection, in which the underlying stealth adapted virus has incorporated  
25 mycoplasma-derived genetic sequences. An interesting admission by the owner of a commercial mycoplasma molecular testing  
26 facility for both CFS and Gulf War Syndrome patients was that the patient's PCR amplified products commonly showed minor  
27 genetic differences. Although he reported the results as being positive for *M. fermentans*, he inwardly thought the patients were  
28 infected with a multiplicity of mycoplasma strains, none of which had the exact sequence of *M. fermentans*.

29 There are also published studies suggesting the potential involvement of *Brucella* bacteria in CFS, fibromyalgia, and the  
30 Gulf War Syndrome [47]. Again, this conclusion was primarily based on positive serological and molecular detection methods.  
31 This was also the suggestion of an inadvertent consequence of efforts to use *Brucella* bacteria to create a Germ Warfare agent

32 [48]. The data provided in this article are more consistent with infection with a viteria containing *brucella* bacteria-derived  
33 genetic sequences.

34 CFS is an imprecisely defined illness [49]. It can range in severity from a rather mild illness to the patients having severe  
35 cognitive impairments. CFS has many clinical features in common with another somewhat controversial illness termed chronic  
36 Lyme disease [50-55]. This supposedly tick-borne illness is attributed to chronic infection with *Borrelia burgdorferi* bacteria.  
37 Comparable to the methods for attributing CFS to mycoplasma infection, the evidence for *Borrelia* infection is largely  
38 serological and/or molecular. Many of the self-designated “Lyme Literate” physicians contend that the disease is empirically  
39 diagnosable even with negative *Borrelia* testing results. Serological and molecular studies have further suggested that many  
40 chronic Lyme disease patients are commonly coinfected with other types of bacteria, including strains of *Bartonella*, *Ehrlichia*,  
41 *Anaplasma*, and *Rickettsia* Bacteria [56]. Infection with *Babesia*, a unicellular parasite is also thought to be common. Blood  
42 samples from many patients diagnosed with chronic Lyme disease were positive when personally tested for stealth adapted virus  
43 infection (unpublished data). It will be of interest to sequence stealth adapted viruses cultured from patients diagnosed as having  
44 chronic Lyme disease. *Borrelia* infections are increasingly being linked with other illnesses. These include Morgellons disease  
45 [57], in which the patients have skin lesions from which electrostatic particles can be obtained. Other diseases attributed to  
46 *Borrelia* include acute psychosis, carditis, autoimmunity, and Guillain-Barre Syndrome [58].

47 Similar considerations apply to other illnesses that are likely to be due to infection with stealth adapted viruses but are  
48 publicly being mistakenly as a bacterial infection. These illnesses include PANDAS (Pediatric Autoimmune Neuropsychiatric  
49 Disorders Associated with Streptococcal Infections) [59-61]. Blood samples from several children with this diagnosis were  
50 tested for stealth adapted viruses with positive results.

51 Other bacterial infections have presumptively been associated with a range of additional neurological, psychiatric,  
52 dermatological, and allergic disorders [62 -66]. Consideration of viteria, i.e., stealth adapted viruses with incorporated bacteria-  
53 derived genetic sequences, may help reorient thinking about these presumed associations.

54 Regardless of the presence of bacteria-derived genetic sequences, major consideration should be given to the  
55 demonstrated wide range of neurological, psychiatric, and other illnesses caused by stealth adapted viruses. Positive viral  
56 cultures were regularly obtained from blood samples of autistic children [67], and from children with severe learning and  
57 behavioral disorders [68]. A virus inducing a very similar CPE as did stealth virus-1 was isolated from the cerebrospinal fluid

58 (CSF) of a comatose patient with a four-year history of a bipolar psychosis [69]. Multiple family members have received  
59 different diagnoses and yet have experienced similar core symptoms. The diagnoses in one family were dementia in both  
60 grandparents, CFS in their daughter and amyotrophic lateral sclerosis (ALS) in her husband, and major learning disorders in  
61 three younger children [70]. A controlled study provided positive culture results in all 10 tested patients with multiple myeloma  
62 [71]. Animal illnesses can also be caused by stealth adapted viruses [72]. On a more positive note, evidence of recovery from  
63 stealth adapted virus infections are highlighting the role of the ACE pathway as a non-immunological anti-viral defense  
64 mechanism [23, 73].

65  
66 **Abbreviations:** ACE – Alternative Cellular Energy, CFS – Chronic Fatigue Syndrome, CPE – cytopathic effect, CSF –  
67 cerebrospinal fluid, KELEA – Kinetic Energy Limiting Electrostatic Attraction, PCR – Polymerase Chain Reaction, SCMV –  
68 African green monkey simian cytomegalovirus, AIDS – Acquired Immunodeficiency Disease, N - non-assigned nucleotide, nt  
69 – nucleotide, bp – base pair, kb – kilobase  
70

71 **Key Words:** Viteria, Renegade Genetic Sequences, Chronic Fatigue Syndrome, AIDS, ACE Pathway, KELEA, amyotrophic  
72 lateral sclerosis, Gulf War Syndrome, *Mycoplasma fermentans*, *Brucella*, *Ochrobactrum*, *Microbacterium*, Autism, Morgellons  
73 disease, PANDAS, Chronic Lyme disease, *Borrelia burgdorferi*  
74  
75  
76

77

## References:

1. Martin WJ, Zeng LC, Ahmed K, et al. Cytomegalovirus-related sequence in an atypical cytopathic virus repeatedly isolated from a patient with chronic fatigue syndrome. *Am J Pathol* 1994;145(2):440–451.
2. Martin WJ, Ahmed KN, Zeng LC, et al. African green monkey origin of the atypical cytopathic “stealth virus” isolated from a patient with chronic fatigue syndrome. *Clin Diagn Virol* 1995;4(1):93–103.
3. Martin WJ. Stealth adaptation of an African green monkey simian cytomegalovirus. *Exp Mol Pathol* 1999;66(1):3–7.
4. Martin WJ. Stealth adaptation of viruses: Review and updated molecular analysis on a stealth adapted African green monkey simian cytomegalovirus (SCMV). *J Human Virology & Retrovirology* 2014;1(4):00020.
5. Martin WJ. Genetic instability and fragmentation of a stealth viral genome. *Pathobiology* 1996;64(1):9–17.
6. Martin WJ. Stealth viruses as neuropathogens. *CAP Today* 1994;8(10): 67–70.
7. Martin WJ. Cellular sequences in stealth viruses. *Pathobiology* 1998;66:53–58.
8. Martin WJ. Bacteria-related sequences in a simian cytomegalovirus– derived stealth virus culture. *Exp Mol Pathol* 1999;66(1):8–14.
9. Martin WJ. Renegade cellular and bacterial genetic sequences in monkey-derived stealth adapted viruses. *J Hum Virol Retrovirology* 2019;7(2):26–40.
10. Martin WJ. Virus transmission to humans of genetically unstable rhesus monkey cellular sequences: A possible forerunner of complex human illnesses. *J Hum Virol Retrovirology* 2020;8(3):74–82.
11. Martin WJ. Infectious, rhesus monkey-derived cellular DNA sequences in certain stealth adapted viruses. *FASEB J* 2020;34(S1): 1-1
12. Martin WJ. Viruses disguised as self and/or as bacteria. *Microbiol Infect Dis* 2020; 4(1):1-5.
13. Martin WJ. Renegade cellular genetic sequences in stealth adapted viruses: introducing a new era of virology. *J Hum Virol Retrovirology*. 2022;9(1):24–29.
14. Razin S. The mycoplasmas. *Microbiol Rev* 1978;42(2):414-70.
15. Volpiano CG, Sant’Anna FH, Ambrosini A, de São José JFB, Beneduzi A, Whitman WB, de Souza EM, Lisboa BB, Vargas LK and Passaglia LMP. Genomic metrics applied to Rhizobiales (Hyphomicrobiales): Species reclassification, identification of unauthentic genomes and false type strains. *Front. Microbiol* 2021;12:614957.

03 16. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z et al. Gapped BLAST and PSI-BLAST: a new generation of  
04 protein database search programs. *Nucleic Acids Res* 1997; 25(17):3389-402.

05 17. Krzyżanowska DM, Maciąg T, Ossowicki A, Rajewska M, Kaczyński Z, Czerwicka M, Rąbalski Ł, Czaplewska P, Jafra  
06 S. *Ochrobactrum quorumnocens* sp. nov., a quorum quenching bacterium from the potato rhizosphere, and comparative  
07 genome analysis with related type strains. *PLoS One* 2019;14(1):e0210874.

08 18. Bautista-Zapanta JN, Arafat HH, Tanaka K, Sawada H, Suzuki K. Variation of 16S-23S internally transcribed spacer  
09 sequence and intervening sequence in rDNA among the three major *Agrobacterium* species. *Microbiological Research*  
10 2009;164(6):604-612.

11 19. Li SY, Huang YE, Chen JY, Lai CH, Mao YC, Huang YT, Liu PY. Genomics of *Ochrobactrum*  
12 *pseudogrignonense* (newly named *Brucella pseudogrignonensis*) reveals a new *bla<sub>OXA</sub>* subgroup. *Microb Genom*  
13 2021;7(8):000626.

14 20. Gtari M, Ghodhbane-Gtari F, Nouiou I, Beauchemin N, Tisa LS. Phylogenetic perspectives of nitrogen-fixing  
15 actinobacteria. *Arch Microbiol* 2012;194(1):3-11.

16 21. Saberi A, Gulyaeva AA, Brubacher JL, Newmark PA, Gorbalya AE. A planarian nidovirus expands the limits of RNA  
17 genome size. *PLoS Pathog* 2018;14(11):e1007314.

18 22. Martin WJ. Detection of RNA sequences in cultures of a stealth virus isolated from the cerebrospinal fluid of a health care  
19 worker with chronic fatigue syndrome. Case report. *Pathobiology* 1997;65:57-60.

20 23. Martin WJ. Stealth virus culture pigments: A potential source of cellular energy. *Exp Mol Path* 2003;74:210-223.

21 24. Martin WJ. Enhancing the alternative cellular energy (ACE) pathway with KELEA activated water as therapy for  
22 infectious diseases. *Infect Disord Drug Targets* 2021;21(3):314-319.

23 25. Prasad V, Greber UF. The endoplasmic reticulum unfolded protein response - homeostasis, cell death and evolution in  
24 virus infections. *FEMS Microbiol Rev* 2021;45(5):fuab016.

25 26. Moreno E, Blasco JM, Letesson JJ, Gorvel JP, Moriyón I. Pathogenicity and its implications in taxonomy:  
26 The *Brucella* and *Ochrobactrum* Case *Pathogens* 2022;11(3):377.

27 27. Alonso CA, Kwabugge YA, Anyanwu MU, Torres C, Chah KF. Diversity of *Ochrobactrum* species in food animals,  
28 antibiotic resistance phenotypes and polymorphisms in the *bla<sub>OCH</sub>* gene, *FEMS Microbiology Letters* 2017;364(17):2017,

29 28. Barquero-Calvo E, Conde-Alvarez R, Chacón-Díaz C, Quesada-Lobo L, Martirosyan A, Guzmán-Verri C, et al. The  
30 differential interaction of *Brucella* and *Ochrobactrum* with innate immunity reveals traits related to the evolution of  
31 stealthy pathogens. *PLoS ONE* 2009;4(6):e5893.

32 29. Liu, H., Carvalhais, L. C., Crawford, M., Singh, E., Dennis, P. G., Pieterse, C. M., et al. Inner plant values: diversity,  
33 colonization and benefits from endophytic bacteria. *Front. Microbiol* 2017;8:2552.

34 30. Jaykus LA, Wang HH, Schlesinger LS. *Food–Borne Microbes: Shaping the Host Ecosystem*. Washington DC. ASM  
35 Press; 2009.

36 31. Galambos, N.; Compant, S.; Wackers, F.; Sessitsch, A.; Anfora, G.; Mazzoni, V.; Pertot, I.; Perazzolli, M. Beneficial  
37 insects deliver plant growth-promoting bacterial endophytes between tomato plants. *Microorganisms* 2021;9:1294.

38 32. Wu Z, Peng W, He X, Wang B, Gan B, Zhang X. Mushroom tumor: a new disease on *Flammulina velutipes* caused by  
39 *Ochrobactrum pseudogrignonense*. *FEMS Microbiol Lett* 2016;363(2):fnv226.

40 33. Fernández-Aguilar X, Cabezón O, Marco I, et al. *Mycoplasma conjunctivae* in domestic small ruminants from high  
41 mountain habitats in Northern Spain. *BMC Vet Res* 2013;9:253.

42 34. Lo S-C, Shih JW-K, Newton PB III, Wong DM, Hayes MM, Benish JR, Wear DJ, Wang RY-H. Virus-like infectious  
43 agent (VLIA) is a novel pathogenic mycoplasma: *Mycoplasma incognitus*. *Amer J Trop Med Hyg* 1989;41:586-600.

44 35. Lo S-C, Dawson MS, Newton PB III, Sonoda MA, Shih JW-K, Engler WF, Wang RY-H, Wear DJ. Association of the  
45 virus-like infectious agent originally reported in patients with AIDS with acute fatal disease in previously healthy non-  
46 AIDS patients. *Amer J Trop Med Hyg* 1989;41:364-376.

47 36. Saillard C, Carle P, Bove JM, Bebear C, Lo S-C, Shih JW-K, Wang RY-H, Rose DL, Tully JG. Genetic and serologic  
48 relatedness between *Mycoplasma fermentans* strains and a mycoplasma recently identified in tissues of AIDS and non-  
49 AIDS patients. *Res Virol* 1990;141:385-395.

50 37. Montagnier L, Blanchard A. Mycoplasmas as cofactors in infection due to the immunodeficiency virus. *Clinical Infectious  
51 Disease* 1993;17:S309-S315.

52 38. Liu Y, Dong Y, Wen Y, Peng S, Liao J, Liu Y. Association of *Mycoplasma fermentans* and the risk of HIV-1 infection: A  
53 meta-analysis. *Medicine (Baltimore)* 2020;99(2):e18499

54 39. Martin WJ. Chimpanzees inoculated with cytomegalovirus contaminated polio vaccines may explain origin of HIV-1. *J  
55 Hum Virol Retrovirol* 2015;2(2):00035.

56 40. Sabin AB. Present position on immunization against poliomyelitis with live virus vaccine. *Brit Med J* 1959;1(5123): 663-  
57 680.

58 41. Baylis SA, Shah N, Jenkins A, Berry NJ, Minor PD. Simian cytomegalovirus and contamination of oral poliovirus  
59 vaccines. *Biologicals* 2002; 31(1):63-73.

60 42. Nicolson GL, Nasralla MY, Haier J, Erwin R, Nicolson NL, Ngwenya R. Mycoplasmal Infections in chronic illnesses:  
61 Fibromyalgia and chronic fatigue syndromes, Gulf War Illness, HIV-AIDS and rheumatoid arthritis. *Medical Sentinel*  
62 1999;4:172-176

63 43. Nicolson NL, Nicolson G, Nasralla M. Mycoplasmal infections and Fibromyalgia/Chronic Fatigue Illness (Gulf War  
64 Illness) associated with deployment to operation desert storm. *Int Journal of Medicine* 1998;1:80-92.

65 44. Vojdani, et al., Detection of *Mycoplasma* genus and *Mycoplasma fermentans* by PCR in patients with Chronic Fatigue  
66 Syndrome, *FEMS Immunol Med Microbiol* 1998;22:355-365.

67 45. Nasralla M, Haier J, Nicolson, G. L. Multiple mycoplasmal infections detected in blood of patients with chronic fatigue  
68 syndrome and/or fibromyalgia syndrome". *Euro J of Microbiol Infect Dis* 1999;18(12):859-865.

69 46. Nasralla, M., Haier, J. and Nicolson, G.L. (1999) Multiple mycoplasmal infections detected in blood of patients with  
70 Chronic Fatigue Syndrome and/or Fibromyalgia. *Euro J of Microbiol Infect Dis* 1999;18:859-865.

71 47. Nicolson G, Gan R, Haier J Evidence for *Brucella* spp. And *Mycoplasma* spp. co-Infections in blood of fatigue syndrome  
72 Patients. *J Chronic Fatigue Syndrome* 2005;12(2):5-17.

73 48. Scott, Donald W. & Scott, William L. C. *The Brucellosis Triangle*. The Chelmsford Publishers, Sudbury, Ontario, Canada  
74 1998

75 49. Martin WJ. Chronic fatigue syndrome (CFS): An imprecisely defined infectious disease caused by stealth adapted viruses.  
76 *J Infect Dis Epidemiol* 2020;6:106.

77 50. Cabello FC, Embers ME, Newman SA, Godfrey HP. *Borrelia burgdorferi* antimicrobial-tolerant persistence in Lyme  
78 disease and posttreatment Lyme disease syndromes. *mBio* 2022;13(3):e0344021.

79 51. Shor S, Green C, Szantyr B, et al. Chronic Lyme disease: An evidence-based definition by the ILADS Working  
80 Group. *Antibiotics (Basel)*. 2019;8(4):269.

81 52. Lantos PM. Chronic Lyme disease. *Infect Dis Clin North Am* 2015;29(2):325-340.

82 53. Fallon, B.A.; Kochevar, J.M.; Gaito, A.; Nields, J.A. The underdiagnosis of neuropsychiatric Lyme disease in children  
83 and adults. *Psychiatr. Clin N Am* 1998;21:693-703.

84 54.Ruiz VH, Edjolo A, Roubaud-Baudron C, et al. Association of seropositivity to *Borrelia burgdorferi* with the risk of  
85 neuropsychiatric disorders and functional decline in older adults: The Aging Multidisciplinary Investigation Study. *JAMA*  
86 *Neurol* 2020;77(2):210-214.

87 55.Diuk-Wasser MA, Vannier E, Krause PJ. Coinfection by *Ixodes* tick-borne pathogens: Ecological, epidemiological, and  
88 clinical consequences. *Trends Parasitol* 2016;32(1):30-42.

89 56.Nelder MP, Russell CB, Sheehan NJ, et al. Human pathogens associated with the blacklegged tick *Ixodes scapularis*: a  
90 systematic review. *Parasit Vectors* 2016;9:265.

91 57.Middelveen MJ, Fesler MC, Stricker RB. History of Morgellons disease: from delusion to definition. *Clin Cosmet Investig*  
92 *Dermatol* 2018;11:71-90.

93 58.Owens J, Filatov A, Husain-Wilson S. Guillain-Barre syndrome, neuroborreliosis, or both. *Cureus* 2020;12(4):e7823.

94 59.Swedo, S.E.; Leckman, J.F.; Rose, N.R. From research subgroup to clinical syndrome: Modifying the PANDAS criteria to  
95 describe PANS (Pediatric Acute-onset Neuropsychiatric Syndrome). *Pediatr Ther* 2012;2: 113.

96 60.Swedo, S.E.; Leonard, H.L.; Garvey, M.; Mittleman, B.; Allen, A.J.; Perlmuter, S.; Lougee, L.; Dow, S.; Zamkoff, J.;  
97 Dubbert, B.K. Pediatric autoimmune neuropsychiatric disorders associated with streptococcal infections: Clinical  
98 description of the first 50 cases. *Am J Psychiatry* 1998;155:264-271.

99 61.Chang, K.; Frankovich, J.; Cooperstock, M.; Cunningham, M.W.; Latimer, M.E.; Murphy, T.K.; Pasternack, M.;  
00 Thienemann, M.; Williams, K.; Walter, J.; et al. Clinical evaluation of youth with Pediatric Acute-Onset Neuropsychiatric  
01 Syndrome (PANS): Recommendations from the 2013 PANS Consensus Conference. *J Child Adolesc Psychopharmacol*  
02 2015;25:3-13.

03 62.Beydoun MA, Beydoun HA, Hossain S, El-Hajj ZW, Weiss J, Zonderman AB. Clinical and bacterial markers of  
04 periodontitis and their association with incident all-cause and Alzheimer's disease dementia in a large national survey. *J*  
05 *Alzheimers Dis* 2020;75(1):157-172.

06 63.Mattson MP. Infectious agents and age-related neurodegenerative disorders. *Ageing Research Reviews* 2004;3:105-120.

07 64.Nicolson GL, Nasralla M, Haier J, Promfret J. High Frequency of systemic mycoplasmal infections in Gulf War veterans  
08 and civilians with Amyotrophic Lateral Sclerosis (ALS). *J Clin Neurosciences* 2002;9: 525-529.

09 65.Keikha M, Karbalaei M. Potential association between bacterial infections and ischemic stroke based on fifty case-control  
10 studies: A systematic review and meta-analysis. *New Microbes New Infect*. 2022;47:100980.

11 66.Johnston, S. L. et al. The effect of telithromycin in acute exacerbations of asthma. *N Engl J Med* 2006;354:1589-1600.

12 67.Martin WJ. Stealth virus isolated from an autistic child. *J Aut Dev Dis* 1995;25:223-224.

13 68.Martin WJ, Anderson D. Stealth virus epidemic in the Mohave Valley: Severe vacuolating encephalopathy in a child  
14 presenting with a behavioral disorder. *Exp Mol Pathol* 1999;66:19-30.

15 69.Martin WJ. Simian cytomegalovirus-related stealth virus isolated from the cerebrospinal fluid of a patient with bipolar  
16 psychosis and acute encephalopathy. *Pathobiology* 1996;64:64-66.

17 70.Martin WJ. Complex multi-system illnesses occurring within a family: Presumptive evidence for an infectious disease  
18 process. *J Neurol Stroke* 2020;10(1):34–36.

19 71.Durie BG, Collins RA, Martin WJ. Positive stealth virus cultures in myeloma patients: A possible explanation for  
20 neuropsychiatric co-morbidity. *Blood* (suppl 1) (2000) Abstr 1553

21 72.Martin WJ, Glass RT. Acute encephalopathy induced in cats with a stealth virus isolated from a patient with chronic  
22 fatigue syndrome. *Pathobiology* 1995;63:115-118.

23 73.Martin WJ. The ACE pathway in comparison to the immune system in the defense against infectious diseases. *J Hum  
24 Virol & Retrovirology* 2016;3(5): 00124.

25

26

27

28

29

30

31

32

33

34

35

36

37

38

CLONE	NCBI ACCESSION NUMBER	LENGTH (nt)
3B23	U27612.2	8916
3B35	U27474.2	2142
3B37 T3 T7	AF065738.1 AF065739.1	676 678
3B41	AF191072.1	2,869
3B43	AF191073.1	3620
3B47	AF191074.1	2024
3B313	U27616.3	7985
3B315 T3 T7	AF065745.1 AF065744.1	808 810
3B512	AF191075	2345
3B513	U27894.2	8106
3B520	AF191076.1	2797
3B528	AF191077.2	2043
3B534 T3 T7	U27638.2 U27900.1	612 808
3B540	AF067304.1	211
3B545	AF067311.1	212
3B562	AF067332.1	310
3B614	U27645.2	5062
3B622 T3 T7	AF067364.1 AF067365.1	818 834
3B627	U27648.1	328
3B632	AF191079.1	1396
3B680 T3 T7	U27773 U27929	100 120

C1616	AF065660.2	4626
C16116 T3	AF065678.1	846
T7	AF065679.1	774
C16118 T3	AF065682.1	903
T7	AF065683.1	787
C16122 T3	U27777.1	840
C61122 T7	U27931.1	812
C16125 T3	AF065692.1	929
T7	AF065693.1	806
C16127	AF065696.1	257
C16134	AF065710.2	4142
C16135 T3	AF065712.1	504
T7	AF065713.1	427
C16141	AF065720.1	213
C16282 T3	U27864.1	250
T7	U27971.1	589

39  
40  
41  
42  
43 **Table 2**  
44 **Clones With Sequences That Best Match to**  
45 *Ochrobactrum quorumnocens* Strain A44  
46

Clone	Ch.	Matching Nucleotides Range	Expect Value	Identity Ratio	Gaps
C16134	1	659449 to 663591	0.0	4127/4143	1
C1616	1	1127153 to 1131778	0.0	4598/4626	1
3B534	1	1208039 to 1208652	0.0	611/615	1
3B629	1	1287540 to 1287704	5e-75	164/165	0
3B313	1	1424566 to 1432547	0.0	7973/7985	3
C16116	1	2327809 to 2328349	0.0	533/552	11

3B614	1	2387534 to 2392595	0.0	5060/5062	0
3B23	1	2395441 to 2404355	0.0	8911/8916	1
C16118 T3	2	24369 to 25146	0.0	720/804	27
T7	2	25368 to 26021	0.0	628/663	10
3B540	2	213959 to 214184	1e-83	205/226	15
C16282 T3*	2	241782 to 242028	6e-83	220/250	3
3B315 T3	2	455544 to 456126	0.0	510/585	2
T7	2	459739 to 460401	0.0	622/678	15
3B43	2	1794830 to 1796043	0.0	1212/1214	0
	2	1796047 to 1798348	0.0	2290/2303	3
3B562	-	111334 to 111643	92-162	310/310	0

47 O. *quorumnocens* Strain A44 NCBI ID VYXQ01000012.1  
 48 Chromosome 1 NCBI ID CP022604.1  
 49 Chromosome 2 NCBI ID: CP022603.1  
 50 \* The T7 sequence of clone C16282 matched to SCMV  
 51

52  
53  
54  
55  
56  
**Table 3**  
**Clones With Sequences That Best Match to**  
***Brucella Pseudogrignonensis* Strain K8**

Clone	Matching Nucleotides		Expect	Identity	Gaps
	Ch.	Range	Value	Ratio	
3B41	2	856636 to 859505	0.0	2330/2886	39
3B47	2	520404 to 522396	0.0	1449/2015	43

57 Chromosome 2 NCBI ID CP022603.1  
58  
59

60  
61  
62 **Table 4**  
63 **Clones With Sequences That Best Match**  
64 **to *Mycoplasma fermentans* M64 Strain**  
65

Clone	Matching Nucleotides	Expect Value	Identity Ratio	Gaps
3B680 T7*	1106962 to 1107081	1e-49	118/120	0
3B627	152708 to 153034	4e-163	327/328	0
3B632*	174189 to 175572	0.0	1380/1384	1
3B520	473083 to 475878	0.0	2771/2797	2
3B528*	651388 to 653431	0.0	2020/2044	1
3B512 *	668449 to 670792	0.0	2342/2345	1
3B35	715735 to 717874	0.0	2136/2142	2
3B37 T7 T3 *	763908 to 764569 765106 to 765768	0.0 0.0	656/672 654/671	10 8
3B545*	801182 to 801394	0.0	212/213	1
3B622T3* T7*	962497 to 963182 963536 to 964273	0.0 0.0	669/697 718/774	11 36

66 \*3B680 T3 also has 79 of 85 nucleotides that match to a sequence in SCMV

67  
68  
69  
70  
**Table 5**  
**Clones With Sequences That Best Match to**  
**Different Species of *Microbacterium***

Clone	Matching Nucleotides		Expect Value	Identity Ratio	Gaps
	Species	Range			
C16122 T3	A	630149 to 630770	2e-102	461/624	2
C16122 T7	B	1690217 to 1690563	6e-26	241/348	5
C16127	C	1364049 to 1364299	2e-120	249/251	0
C16141	D	657325 to 657522	7e-93	197/198	0

71 NCBI ID of best matching *Microbacterium* sequence and name of species  
72 A CP063814.1 *Microbacterium* A18JL200 chromosome  
73 B CP043732.1 *Microbacterium esteraromaticum* strain B24  
74 C LR880474.1 *Microbacterium* Nx66 genome assembly  
75 D CP080491.1 *Microbacterium* Se5.02b chromosome  
76  
77

**Table 6**  
**Clone With Sequences That Best Match to**  
**A Plasmid Sequence in *Aminobacter***

Clone	Matching Nucleotides Range	Expect Value	Identity Ratio	Gaps
C16125 T7	19341 to 19987	0.0	565/659	12
C16125 T3	21377 to 22070	0.0	578/698	5

NCBI ID: [CP060199.1](https://www.ncbi.nlm.nih.gov/nuccore/CP060199.1) *Aminobacter* sp. MDW-2 plasmid pMDW2

96  
97  
**Table 7**  
98  
99

00  
01  
**Clone With Sequence at One End That Best Match to**  
02  
***Cellulosmicrobium* strain ORNL-0100 chromosome**  
03

Clone	Matching Nucleotides Range	Expect Value	Identity Ratio	Gaps
C16135 T3	1323875 to 1324367	0.0	477/493	0

01  
02  
03  
NCBI ID: CP072387.1

The T7 sequence of clone C16135 matches to a sequence in the human Karzin gene

04  
05  
06  
07  
08  
09  
**Table 8**  
**Nucleotide Matching to the Sequence of Clone 3B513**

Best Matching Microbe*	NCBI Accession No.	Length (nt))	Matching Target Sequence(s)	Matching 3B513 Nucleotides	Expect Value	Identity Ratio	Gaps
<i>Shinella oryzae</i> strain Z-25 plasmid pZ25	NZ_CP081988.1	1888955	81735 to 82117 80500 to 80963 79997 to 80149 79570 to 79715	1 to 383 1149 to 1613 1970 to 2122 2404 to 2549	8e-53 2e-35 3e-13 3e-08	278/386 324/476 113/156 103/148	6 23 6 4
<i>Rhizobium flavum</i> strain YW14 genome assembly	LR723673.1	150934	28209 to 28633	2663 to 3088	0.0	417/426	1
<i>Aminobacter niigataensis</i> strain DSM 7050 Ga0373207_13	NZ_JACHOT010000013	40804	13754 to 14755 15089 to 15368 15715 to 16298	3218 to 4219 4556 to 4835 5182 to 5762	6e-143 6e-10 3e-90	716/1006 183/281 425/585	8 2 5
<i>Ochrobactrum</i> sp. POC9 _contig00056,	QGST01000056.1	5479	284 to 2598	5202 to 7509	0.0	2298/2315	7
<i>Mesorhizobium</i> denitrificans strain LA-28	NZ_QURN01000017.1	86039	77672 to 78643	7143 - 8106	0.0	962/972	8
Stealth Virus 3B525 T3	U27635.1	853	76 to 593	7520 to 8027	0.0	415/434*	11

\*Although the pairwise matching with Stealth virus-1 clone extended from nucleotides 76 to 593 with an Expect Value of 0.0, the identity ratio calculation was made from nucleotide 160 to 593 to minimize the extent of mismatching with undefined N nucleotides




Figure 1.

Legend: Photo of an ethidium bromide-stained 8-laned agarose gel electrophoresis. The arrow points to lane 4 and shows the migration of a portion of the DNA that was extracted from the filtered and ultracentrifuged supernatant of stealth virus-1 infected MRHF cells. Lane 3 directly beneath the arrow shows the migration of another portion of the extracted DNA that was digested using EcoRI enzyme prior to electrophoresis. Lane 1 is EcoRI digested DNA obtained from the lysate of the infected MRHF cells. Lanes 1 and 3 are HindIII and Bst-II lambda phage DNA markers, the largest of which are 23,130 and 8,454 nucleotide base pairs, respectively. The lower staining material in lanes 3, 4, and 6 is RNA. Reproduced from reference (1) with permission.

