

UC Irvine

UC Irvine Previously Published Works

Title

From FOS fusions to somatic mutations in the MAPK pathway, heterogeneous genetic abnormalities cause distinct pathophysiology among subsets of epithelioid haemangiomas

Permalink

<https://escholarship.org/uc/item/73t40265>

Journal

British Journal of Dermatology, 186(3)

ISSN

0007-0963

Authors

Tan, W
Nelson, JS

Publication Date

2022-03-01

DOI

10.1111/bjd.20952

Peer reviewed

Conflicts of interest: the authors declare they have no conflicts of interest.

References

- 1 Beck DB, Ferrada MA, Sikora KA et al. Somatic mutations in UBA1 and severe adult-onset autoinflammatory disease. *N Engl J Med* 2020; **383**:2628–38.
- 2 van der Made CI, Potjewijd J, Hoogstins A et al. Adult-onset autoinflammation caused by mutations in UBA1: a Dutch case series of patients with VEXAS. *J Allergy Clin Immunol* 2022; **149**(1):432–9.
- 3 Ferrada MA, Sikora KA, Luo Y et al. Somatic mutations in UBA1 define a distinct subset of relapsing polychondritis patients with VEXAS. *Arthritis Rheumatol* 2021; **73**:1886–95.
- 4 Georgin-Lavialle S, Terrier B, Guedon AF et al. Further characterization of clinical and laboratory features in VEXAS syndrome: large-scale analysis of a multicentre case series of 116 French patients. *Br J Dermatol* 2022; **186**:564–74.

From *FOS* fusions to somatic mutations in the MAPK pathway, heterogeneous genetic abnormalities cause distinct pathophysiology among subsets of epithelioid haemangiomas

DOI: 10.1111/bjd.20952

Linked Article: Maurus et al. *Br J Dermatol* 2022; **186**:553–563.

Epithelioid haemangioma (EH) is a rare benign vascular tumour comprised of prominent epithelioid endothelial cells, occurring at diverse anatomical locations including soft tissues, bone and skin.^{1,2} EH can be categorized into three subsets: typical, cellular, and angiolymphoid hyperplasia with eosinophilia (ALHE), with a wide range of vasoproliferative spectra and inflammatory infiltrates. The emerging evidence of genetic abnormalities in EH includes *FOS* rearrangements and somatic mutations in the mitogen-activated protein kinase (MAPK) pathway. This has greatly enhanced our understanding of the aetiology of this disease and facilitated molecular differential diagnosis from malignant epithelioid vascular tumours such as epithelioid haemangioendothelioma and angiosarcoma.

There are several types of *FOS* gene fusions reported in EH, including *ZFP36/FOSB*, *WWTR1/FOSB*, *FOS/LMNA* and *FOS/VIM*, which lead to consecutive overexpression of their transcripts.^{3–5} These *FOS* rearrangements are commonly found in soft tissue and intraosseous lesions, but are very infrequent in cutaneous EH.^{3,6} In addition, a fusion of *GATA6/FOXO1* was reported in a subset of EH lacking *FOS* fusions.⁷

In this issue of the *BJD*, Maurus et al. report various somatic mutations in *MAP2K1* and *KRAS* in cutaneous EH, which resulted in an enhanced MAPK pathway.⁸ These somatic mutations may account for distinct pathological mechanisms underlying the development of cutaneous EH where *FOS* gene fusions are absent. The authors identified several somatic

mutations in the *MAP2K1* gene (including c.383G>A, p.Gly128Asp and c.171G>T or C, p.Lys57Asn) and in the *KRAS* gene (c.182A>G, p.Gln61Arg).⁸ These mutations occur in 50% of patients across all three EH subsets: typical, cellular and ALHE. The mutation allele frequency ranged from 1.5% to 6.2%.⁸ As a pathological readout of these mutations, a strong activation of the MAPK pathway and subsequently morphological changes in endothelial cells was observed.⁸

Low-frequency somatic mutations in many MAPK pathway-associated genes (*KRAS*, *NRAS*, *BRAF* and *MAP2K1*) have been reported in various vascular anomalies including arteriovenous malformations.⁹ This cumulative evidence combined with the report from Maurus et al. suggests a mechanistic role of mosaic mutations for targeting the MAPK pathway in the pathogenesis of certain cutaneous vascular disorders including EH. However, there are several critical questions yet to be answered prior to affirmation of their detrimental roles. Firstly, are these low-frequency somatic mutations primary causes, role players or merely pathological consequences of the EH? Secondly, which cell types in lesions harbour those mutations? Maurus et al. suggest that epithelioid endothelial cells carry the mutations but lymphocytes do not.⁸ As a result of the low frequencies of these mutations, it is reasonably posited that only a very few epithelioid endothelial cells carry the mutations. What are the molecular phenotypes of those mutation-harbouring cells and what are their detrimental roles in the development and progression of the cutaneous EH? Thirdly, if somatic mutations in the MAPK pathway act as a 'second hit' for cutaneous EH development, the 'primary hit' has yet to be elucidated.

Nevertheless, the identification of somatic mutations in *MAP2K1* and *KRAS* has provided a new insight into the neoplastic nature of cutaneous EH, suggesting various genetic abnormalities underlying distinct molecular pathophysiology among subsets of EH.

W. Tan ¹ and J.S. Nelson²

¹Department of Cell Biology and Anatomy, University of South Carolina School of Medicine, Columbia, SC 29209, USA and ²Department of Surgery, Beckman Laser Institute and Medical Clinic, University of California Irvine, Irvine, CA 92612, USA
Email: wembin.tan@uscmed.sc.edu

Funding sources: this work was supported by a grant from the National Institutes of Health (AR073172 to W.T.).

Conflicts of interest: the authors declare they have no conflicts of interest.

References

- 1 Wells GC, Whimster IW. Subcutaneous angiolymphoid hyperplasia with eosinophilia. *Br J Dermatol* 1969; **81**:1–14.
- 2 Fetsch JF, Weiss SW. Observations concerning the pathogenesis of epithelioid hemangioma (angiolymphoid hyperplasia). *Mod Pathol* 1991; **4**:449–55.

- 3 Huang SC, Zhang L, Sung YS et al. Frequent FOS gene rearrangements in epithelioid hemangioma: a molecular study of 58 cases with morphologic reappraisal. *Am J Surg Pathol* 2015; **39**:1313–21.
- 4 Antonescu CR, Chen HW, Sung YS et al. ZFP36-FOSB fusion defines a subset of epithelioid hemangioma with atypical features. *Genes Chromosomes Cancer* 2014; **53**:951–9.
- 5 Tsuda Y, Suurmeijer AJH, Sung YS et al. Epithelioid hemangioma of bone harboring FOS and FOSB gene rearrangements: a clinicopathologic and molecular study. *Genes Chromosomes Cancer* 2021; **60**:17–25.
- 6 Luzar B, Jeremia E, Antonescu CR et al. Cutaneous intravascular epithelioid hemangioma. A clinicopathological and molecular study of 21 cases. *Mod Pathol* 2020; **33**:1527–36.
- 7 Antonescu CR, Huang SC, Sung YS et al. Novel GATA6-FOXO1 fusions in a subset of epithelioid hemangioma. *Mod Pathol* 2021; **34**:934–41.
- 8 Maurus K, Kosnoppel C, Kneitz H et al. Cutaneous epithelioid hemangiomas show somatic mutations in the mitogen-activated protein kinase pathway. *Br J Dermatol* 2022; **186**:553–63.
- 9 Al-Olabi L, Polubothu S, Dowsett K et al. Mosaic RAS/MAPK variants cause sporadic vascular malformations which respond to targeted therapy. *J Clin Invest* 2018; **128**:5185.

Safety in numbers: risankizumab for moderate-to-severe psoriasis

DOI: 10.1111/bjd.20947

Linked Article: Gordon et al. *Br J Dermatol* 2022; **186**:466–475.

Biologic therapies are effective at improving the quality of life and clinical outcomes of patients with moderate-to-severe psoriasis.¹ Risankizumab – a humanized monoclonal antibody that inhibits interleukin (IL)-23, a key cytokine in the pathogenesis of psoriasis – is effective for treatment of psoriasis and Crohn disease.^{2–6}

In this issue of the *BJD*, Gordon et al. report on the safety of risankizumab by analysing data from 17 completed or ongoing trials in patients with plaque psoriasis.⁷ Adverse events were reported as exposure-adjusted event rates (EAERs) per 100 patient-years (PY). The dataset was divided into adverse events that occurred in the short term (defined as <16 weeks; 1306 patients) and adverse events that occurred in the long term (3072 patients). Consistently with previous clinical trials, the most common adverse events were nasopharyngitis and viral upper respiratory tract infections (URTIs).^{3–6} Most of the EAERs recorded were mild or moderate, with serious drug-related EAERs accounting for only 1.5 and 1.0 events per 100 PY in the short- and long-term datasets, respectively, in line with benchmark reference ranges.

Importantly, this study did not identify any new safety signals for risankizumab. Inhibition of IL-23 did not predispose patients to bacterial or fungal pathogens, nor did it increase the risk of malignancy. In head-to-head clinical trials with other biologics such as ustekinumab (anti-IL-12/23),

adalimumab (anti-tumour necrosis factor) and secukinumab (anti-IL-17A), risankizumab was more effective in the treatment of moderate-to-severe plaque psoriasis and had a similar safety profile, with the most common adverse event being viral URTIs.^{3–6} Based on the evidence we have so far, risankizumab appears to be both effective and safe.

While these findings are encouraging, there are several limitations to consider. Although the authors include comprehensive safety data from 17 clinical trials, the generalizability of the results is limited to the populations included in the original trials. For example, a reviewer noted that although the authors report very few cases of infections such as tuberculosis in patients treated with risankizumab, the majority of trials were conducted in countries with low rates of tuberculosis, making it unclear how these safety data might differ in countries with higher prevalence of tuberculosis.

As the authors acknowledge, patient attrition over time may have selected for healthier patients at later timepoints, potentially biasing the long-term safety profile of risankizumab. Safety data were available for over 5 years, making this the most comprehensive study on the safety of risankizumab to date; however, many of the safety concerns, including cancer, may take decades to develop, making it paramount to continue efforts to collect safety data in the long term. Data from large registries, such as BADBIR (British Association of Dermatologists Biologics and Immunomodulators Register), provide the power and duration of follow-up necessary to identify any rare or delayed adverse effects in a real-world setting.⁸

In summary, this study provides encouraging data for the safety profile of risankizumab for the management of moderate-to-severe plaque psoriasis. Continued analyses in real-world settings and ongoing clinical trials are necessary to further characterize the long-term side-effect profile of risankizumab in diverse settings, and to highlight the potential barriers to access to biologics and implications in the context of growing healthcare costs.⁹

Acknowledgments: We thank Dr Zenas Yiu, *BJD* Associate Editor and NIHR Clinical Lecturer at the University of Manchester, for his contribution to this commentary.

S. Mirali ¹, K. Fabusiwa² and E. Linos ³

¹University of Toronto Faculty of Medicine, Toronto, ON, Canada; ²Guy's and St Thomas' Hospital, London, UK; and ³Stanford University Department of Dermatology, Palo Alto, CA, USA
Email: linos@stanford.edu

Conflicts of interest: the authors declare they have no conflicts of interest.

References

- 1 Iskandar IYK, Ashcroft DM, Warren RB et al. Comparative effectiveness of biological therapies on improvements in quality of life in patients with psoriasis. *Br J Dermatol* 2017; **177**:1410–21.