

Ancient human genomes suggest three ancestral populations for present-day Europeans

Iosif Lazaridis^{1,2}, Nick Patterson², Alissa Mittnik³, Gabriel Renaud⁴, Swapan Mallick^{1,2}, Karola Kirsanow⁵, Peter H. Sudmant⁶, Joshua G. Schraiber⁷, Sergi Castellano⁴, Mark Lipson⁸, Bonnie Berger^{2,8}, Christos Economou⁹, Ruth Bollongino⁵, Qiaomei Fu^{1,4,10}, Kirsten I. Bos³, Susanne Nordenfelt^{1,2}, Heng Li^{1,2}, Cesare de Filippo⁴, Kay Prüfer⁴, Susanna Sawyer⁴, Cosimo Posth³, Wolfgang Haak¹¹, Fredrik Hallgren¹², Elin Fornander¹², Nadin Rohland^{1,2}, Dominique Delsate^{13,14}, Michael Francken¹⁵, Jean-Michel Guinet¹³, Joachim Wahl¹⁶, George Ayodo¹⁷, Hamza A. Babiker^{18,19}, Graciela Bailliet²⁰, Elena Balanovska²¹, Oleg Balanovsky^{21,22}, Ramiro Barrantes²³, Gabriel Bedoya²⁴, Haim Ben-Ami²⁵, Judit Bene²⁶, Fouad Berrada²⁷, Claudio M. Bravi²⁰, Francesca Brisighelli²⁸, George Busby^{29,30}, Francesco Cali³¹, Mikhail Churnosov³², David E. C. Cole³³, Daniel Corach³⁴, Larissa Damba³⁵, George van Driem³⁶, Stanislav Dryomov³⁷, Jean-Michel Dugoujon³⁸, Sardana A. Fedorova³⁹, Irene Gallego Romero⁴⁰, Marina Gubina³⁵, Michael Hammer⁴¹, Brenna Henn⁴², Tor Hervig⁴³, Ugur Hodoglugil⁴⁴, Aashish R. Jha⁴⁰, Sena Karachanak-Yankova⁴⁵, Rita Khusainova^{46,47}, Elza Khusnutdinova^{46,47}, Rick Kittles⁴⁸, Toomas Kivisild⁴⁹, William Klitz⁵⁰, Vaidutis Kučinskas⁵¹, Alena Kushniarevich⁵², Leila Laredj⁵³, Sergey Litvinov^{46,47,52}, Theologos Loukidis⁵⁴, Robert W. Mahley⁵⁵, Béla Melegh²⁶, Ene Metspalu⁵⁶, Julio Molina⁵⁷, Joanna Mountain⁵⁸, Klemetti Näkkäläjärvi⁵⁹, Desislava Nesheva⁴⁵, Thomas Nyambo⁶⁰, Ludmila Osipova³⁵, Jüri Parik⁵⁶, Fedor Platonov⁶¹, Olga Posukh³⁵, Valentino Romano⁶², Francisco Rothhammer^{63,64,65}, Igor Rudan⁶⁶, Ruslan Ruizbakiev⁶⁷, Hovhannes Sahakyan^{52,68}, Antti Sajantila^{69,70}, Antonio Salas⁷¹, Elena B. Starikovskaya³⁷, Ayele Tarekegn⁷², Draga Toncheva⁴⁵, Shahlo Turdikulova⁷³, Ingrida Uktveryte⁵¹, Olga Utevska⁷⁴, René Vasquez⁷⁵, Mercedes Villena⁷⁵, Mikhail Voevoda^{35,76}, Cheryl Winkler⁷⁷, Levon Yepiskoposyan⁶⁸, Pierre Zalloua^{78,79}, Tatijana Zemunik⁸⁰, Alan Cooper¹¹, Cristian Capelli²⁹, Mark G. Thomas⁸¹, Andres Ruiz-Linares⁸¹, Sarah A. Tishkoff⁸², Lalji Singh^{83,84}, Kumarasamy Thangaraj⁸³, Richard Villems^{52,56,85}, David Comas⁸⁶, Rem Sukernik³⁷, Mait Metspalu⁵², Matthias Meyer⁴, Evan E. Eichler^{6,87}, Joachim Burger⁵, Montgomery Slatkin⁷, Svante Pääbo⁴, Janet Kelso⁴, David Reich^{1,2,88,†} and Johannes Krause^{3,89,†}

† Co-senior authors

¹ Department of Genetics, Harvard Medical School, Boston, MA, 02115, USA

² Broad Institute of Harvard and MIT, Cambridge, MA, 02142, USA

³ Institute for Archaeological Sciences, University of Tübingen, Tübingen, 72074, Germany

- ⁴ Max Planck Institute for Evolutionary Anthropology, Leipzig, 04103, Germany
- ⁵ Johannes Gutenberg University Mainz, Institute of Anthropology, Mainz, D-55128, Germany
- ⁶ Department of Genome Sciences, University of Washington, Seattle, WA, 98195, USA
- ⁷ Department of Integrative Biology, University of California, Berkeley, CA, 94720-3140, USA
- ⁸ Department of Mathematics and Computer Science and Artificial Intelligence Laboratory, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
- ⁹ Archaeological Research Laboratory, Stockholm University, 114 18, Sweden
- ¹⁰ Key Laboratory of Vertebrate Evolution and Human Origins of Chinese Academy of Sciences, IVPP, CAS, Beijing, 100049, China
- ¹¹ Australian Centre for Ancient DNA, School of Earth and Environmental Sciences, University of Adelaide, Adelaide, South Australia, SA 5005, Australia
- ¹² The Cultural Heritage Foundation, Västerås, 722 12, Sweden
- ¹³ National Museum of Natural History, L-2160, Luxembourg
- ¹⁴ National Center of Archaeological Research, National Museum of History and Art, L-2345, Luxembourg
- ¹⁵ Department of Paleanthropology, Senckenberg Center for Human Evolution and Paleoenvironment, University of Tübingen, Tübingen, D-72070, Germany
- ¹⁶ State Office for Cultural Heritage Management Baden-Württemberg, Osteology, Konstanz, D-78467, Germany
- ¹⁷ Center for Global Health and Child Development, Kisumu, 40100, Kenya
- ¹⁸ Institutes of Evolution, Immunology and Infection Research, School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3JT, UK
- ¹⁹ Biochemistry Department, Faculty of Medicine, Sultan Qaboos University, Alkhod, Muscat, 123, Oman
- ²⁰ Laboratorio de Genética Molecular Poblacional, Instituto Multidisciplinario de Biología Celular (IMBICE), CCT-CONICET & CICPBA, La Plata, B1906APO, Argentina
- ²¹ Research Centre for Medical Genetics, Moscow, 115478, Russia
- ²² Vavilov Institute for General Genetics, Moscow, 119991, Russia
- ²³ Escuela de Biología, Universidad de Costa Rica, San José, 2060, Costa Rica
- ²⁴ Institute of Biology, Research group GENMOL, Universidad de Antioquia, Medellín, Colombia
- ²⁵ Rambam Health Care Campus, Haifa, 31096, Israel
- ²⁶ Department of Medical Genetics and Szentagothai Research Center, University of Pécs, Pécs, H-7624 Hungary
- ²⁷ Al Akhawayn University in Ifrane (AUI), School of Science and Engineering, Ifrane, 53000, Morocco
- ²⁸ Forensic Genetics Laboratory, Institute of Legal Medicine, Università Cattolica del Sacro Cuore, Rome, 00168, Italy
- ²⁹ Department of Zoology, University of Oxford, Oxford, OX1 3PS, UK
- ³⁰ Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, OX3 7BN, UK
- ³¹ Laboratorio di Genetica Molecolare, IRCCS Associazione Oasi Maria SS, Troina, 94018, Italy
- ³² Belgorod State University, Belgorod, 308015, Russia
- ³³ Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Ontario, M5G 1L5, Canada
- ³⁴ Servicio de Huellas Digitales Genéticas, School of Pharmacy and Biochemistry, Universidad de Buenos Aires, 1113 CABA, Argentina
- ³⁵ Institute of Cytology and Genetics, Siberian Branch of Russian Academy of Sciences, Novosibirsk, 630090, Russia
- ³⁶ Institute of Linguistics, University of Bern, Bern, CH-3012, Switzerland
- ³⁷ Laboratory of Human Molecular Genetics, Institute of Molecular and Cellular Biology, Russian Academy of Science, Siberian Branch, Novosibirsk, 630090, Russia
- ³⁸ Anthropologie Moléculaire et Imagerie de Synthèse, CNRS UMR 5288, Université Paul Sabatier Toulouse III, Toulouse, 31000, France
- ³⁹ Yakut Research Center of Complex Medical Problems and North-Eastern Federal University, Yakutsk, 677010, Russia
- ⁴⁰ Department of Human Genetics, University of Chicago, Chicago, IL, 60637, USA
- ⁴¹ ARL Division of Biotechnology, University of Arizona, Tucson, AZ, 85721, USA
- ⁴² Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY, 11794, USA
- ⁴³ Department of Clinical Science, University of Bergen, Bergen, 5021, Norway
- ⁴⁴ NextBio, part of Illumina, Santa Clara, CA, USA 95050
- ⁴⁵ Dept. of Medical Genetics, National Human Genome Center, Medical University Sofia, Sofia, 1431, Bulgaria

- 46 Institute of Biochemistry and Genetics, Ufa Research Centre, Russian Academy of Sciences, Ufa, 450054, Russia
- 47 Department of Genetics and Fundamental Medicine, Bashkir State University, Ufa, 450074, Russia
- 48 College of Medicine, University of Illinois at Chicago, Chicago, IL, 60607, USA
- 49 Division of Biological Anthropology, University of Cambridge, Cambridge, United Kingdom CB2 1QH
- 50 School of Public Health, University of California, Berkeley, CA, 94720, USA
- 51 Department of Human and Medical Genetics, Vilnius University, Vilnius, LT-08661, Lithuania
- 52 Estonian Biocentre, Evolutionary Biology group, Tartu, 51010, Estonia
- 53 Translational Medicine and Neurogenetics, Institut de Génétique et de Biologie Moléculaire et Cellulaire, Illkirch, 67404, France
- 54 Currently employed by AMGEN; 33 Kazantzaki Str, Ilioupolis 16342, Athens, Greece
- 55 Gladstone Institutes, San Francisco, CA, 94158, USA
- 56 Department of Evolutionary Biology, University of Tartu, Tartu, 51010, Estonia
- 57 Centro de Investigaciones Biomédicas de Guatemala, Ciudad de Guatemala, Guatemala
- 58 Research Department, 23andMe, Inc. Mountain View, CA, 94043, USA
- 59 Cultural Anthropology Program, University of Oulu, Oulu, 90014, Finland
- 60 Department of Biochemistry, Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania
- 61 Research Institute of Health, North-Eastern Federal University, Yakutsk, 677000, Russia
- 62 Dipartimento di Fisica e Chimica, Università di Palermo, Palermo, 90128, Italy
- 63 Instituto de Alta Investigación, Universidad de Tarapacá, Arica, Chile
- 64 Programa de Genética Humana ICBM Facultad de Medicina Universidad de Chile, Santiago, Chile
- 65 Centro de Investigaciones del Hombre en el Desierto, Arica, Chile
- 66 Centre for Population Health Sciences, The University of Edinburgh Medical School, Edinburgh, Scotland, EH8 9AG, UK
- 67 Deceased: formerly of the Institute of Immunology, Academy of Science, Tashkent, 70000, Uzbekistan
- 68 Laboratory of Ethnogenomics, Institute of Molecular Biology, National Academy of Sciences of Armenia, Yerevan, 0014, Armenia
- 69 Department of Forensic Medicine, Hjelt Institute, University of Helsinki, Helsinki, 00014, Finland
- 70 Institute of Applied Genetics, Department of Molecular and Medical Genetics, University of North Texas Health Science Center, Fort Worth, Texas 76107, USA
- 71 Unidade de Xenética, Departamento de Anatomía Patolóxica e Ciencias Forenses, and Instituto de Ciencias Forenses, Grupo de Medicina Xenómica (GMX), Facultade de Medicina, Universidade de Santiago de Compostela, Galicia, 15872, Spain
- 72 Research Fellow, Henry Stewart Group, Russell House, London WC1A 2HN, UK
- 73 Institute of Bioorganic Chemistry Academy of Sciences Republic of Uzbekistan, Tashkent, 100125, Uzbekistan
- 74 Department of Genetics and Cytology, V.N. Karazin Kharkiv National University, Kharkiv, 61077, Ukraine
- 75 Instituto Boliviano de Biología de la Altura, Universidad Autónoma Tomás Frías, Potosí, Bolivia
- 76 Inst. of Internal Medicine, Siberian Branch of Russian Acad. of Medical Sciences, Novosibirsk, 630089, Russia
- 77 Molecular Genetics Epidemiology Section, Frederick National Lab for Cancer Research, NCI, NIH, Frederick, MD 21702, USA
- 78 Lebanese American University, School of Medicine, Beirut, 13-5053, Lebanon
- 79 Harvard School of Public Health, Boston, 02115, USA
- 80 Department of Medical Biology, University of Split, School of Medicine, Split, 21000, Croatia
- 81 Department of Genetics, Evolution and Environment, University College London, WC1E 6BT, UK
- 82 Department of Biology and Genetics, University of Pennsylvania, Philadelphia, Pennsylvania, 19104, USA
- 83 CSIR-Centre for Cellular and Molecular Biology, Hyderabad, 500 007, India
- 84 Present address: Banaras Hindu University, Varanasi, 221 005, India
- 85 Estonian Academy of Sciences, Tallinn, 10130, Estonia
- 86 Institut de Biologia Evolutiva (CSIC-UPF), Departament de Ciències Experimentals i de la Salut, Universitat Pompeu Fabra, Barcelona, 08003, Spain
- 87 Howard Hughes Medical Institute, University of Washington, Seattle, WA, USA 98195
- 88 Howard Hughes Medical Institute, Harvard Medical School, Boston, MA, 02115, USA
- 89 Senckenberg Centre for Human Evolution and Palaeoenvironment, University of Tübingen, 72070, Germany

We sequenced genomes from a ~7,000 year old early farmer from Stuttgart in Germany, an ~8,000 year old hunter-gatherer from Luxembourg, and seven ~8,000 year old hunter-gatherers from southern Sweden. We analyzed these data together with other ancient genomes and 2,345 contemporary humans to show that the great majority of present-day Europeans derive from at least three highly differentiated populations: West European Hunter-Gatherers (WHG), who contributed ancestry to all Europeans but not to Near Easterners; Ancient North Eurasians (ANE), who were most closely related to Upper Paleolithic Siberians and contributed to both Europeans and Near Easterners; and Early European Farmers (EEF), who were mainly of Near Eastern origin but also harbored WHG-related ancestry. We model these populations' deep relationships and show that EEF had ~44% ancestry from a "Basal Eurasian" lineage that split prior to the diversification of all other non-African lineages.

Ancient DNA studies have demonstrated that migration played a major role in the introduction of agriculture to Europe, as early farmers were genetically distinct from hunter-gatherers^{1,2} and closer to present-day Near Easterners^{2,3}. Modelling the ancestry of present-day Europeans as a simple mixture of two ancestral populations², however, does not take into account their genetic affinity to an Ancient North Eurasian (ANE) population^{4,5} who also contributed genetically to Native Americans⁶. To better understand the deep ancestry of present-day Europeans, we sequenced nine ancient genomes that span the transition from hunting and gathering to agriculture in Europe (Fig. 1A; Extended Data Fig. 1): "Stuttgart" (19-fold coverage), a ~7,000 year old skeleton found in Germany in the context of artifacts from the first widespread Neolithic farming culture of central Europe, the *Linearbandkeramik*; "Loschbour" (22-fold coverage), an ~8,000 year old skeleton from the Loschbour rock shelter in Heffingen, Luxembourg, discovered in the context of Mesolithic hunter-gatherer artifacts (SI1; SI2); and seven samples (0.01-2.4-fold coverage) from an ~8,000 year old Mesolithic hunter-gatherer burial in Motala, Sweden.

A central challenge is to show that DNA sequences retrieved from ancient samples are authentic and not due to present-day human contamination. The rate of C→T and G→A mismatches to the human genome at the ends of the molecules in libraries from each of the ancient samples exceeds 20%, a signature that suggests the DNA is largely ancient^{7,8} (SI3). We inferred mitochondrial

DNA (mtDNA) consensus sequences, and based on the number of sites that differed, estimated contamination rates of 0.3% for Loschbour, 0.4% for Stuttgart, and 0.01%-5% for the Motala individuals (SI3). We inferred similar levels of contamination for the nuclear DNA of Loschbour (0.4%) and Stuttgart (0.3%) using a maximum-likelihood-based test (SI3). The effective contamination rate for the high coverage samples is likely to be far lower, as consensus diploid genotype calling (SI2) tends to reduce the effects of a small fraction of contaminating reads.

Stuttgart belongs to mtDNA haplogroup T2, typical of Neolithic Europeans⁹, while Loschbour and all Motala individuals belong to haplogroups U5 and U2, typical of pre-agricultural Europeans^{1,7} (SI4). Based on the ratio of reads aligning to chromosomes X and Y, Stuttgart is female, while Loschbour and five of seven Motala individuals are male¹⁰ (SI5). Loschbour and the four Motala males whose haplogroups we could determine all belong to Y-chromosome haplogroup I, suggesting that this was a predominant haplogroup in pre-agricultural northern Europeans analogous to mtDNA haplogroup U¹¹ (SI5).

We carried out most of our sequencing on libraries prepared in the presence of uracil DNA glycosylase (UDG), which reduces C→T and G→A errors due to ancient DNA damage (SI3). We first confirm that the ancient samples had statistically indistinguishable levels of Neandertal ancestry to each other (~2%) and to present-day Eurasians (SI6), and so we do not consider this further in our analyses of population relationships. We report analyses that leverage the type of information that can only be obtained from deep coverage genomes, mostly focusing on Loschbour and Stuttgart, and for some analyses also including Motala12 (2.4×) and La Braña from Mesolithic Iberia (3.4×)¹². Heterozygosity, the number of differences per nucleotide between an individual's two chromosomes, is 0.00074 for Stuttgart, at the high end of present-day Europeans, and 0.00048 for Loschbour, lower than in any present-day humans (SI2). Through comparison of Loschbour's two chromosomes we find that this low diversity is not due to recent inbreeding but instead due to a population bottleneck in this individual's more distant ancestors (Extended Data Fig. 2). Regarding alleles that affect phenotype, we find that the *AMY1* gene coding for salivary amylase had 5, 6, 13, and 16 copies in La Braña¹², Motala12, Loschbour and Stuttgart respectively; these numbers are within the range of present-day Europeans (SI7), suggesting that high copy counts of *AMY1* are not entirely due to selection since the switch to

agriculture¹³. The genotypes at SNPs associated with lactase persistence indicate that Stuttgart, Loschbour, and Motala12 were unable to digest milk as adults. Both Loschbour and Stuttgart likely had dark hair (>99% probability); Loschbour, like La Braña and Motala12, likely had blue or intermediate-colored eyes (>75% probability), while Stuttgart most likely had brown eyes (>99% probability) (SI8). Neither Loschbour nor La Braña carries the skin-lightening allele in *SLC24A5* that is homozygous in Stuttgart and nearly fixed in Europeans today, indicating that they probably had darker skin¹². However, Motala12 carries at least one copy of the derived allele, indicating that this locus was already polymorphic in Europeans prior to the advent of agriculture.

To place the ancient European genomes in the context of present-day human genetic variation, we assembled a dataset of 2,345 present-day humans from 203 populations genotyped at 594,924 autosomal single nucleotide polymorphisms (SNPs)⁵ (SI9) (Extended Data Table 1). We used ADMIXTURE¹⁴ to identify 59 “West Eurasian” populations (777 individuals) that cluster with Europe and the Near East (SI9 and Extended Data Fig. 3). Principal component analysis (PCA)¹⁵ (SI10) (Fig. 1B) reveals a discontinuity between the Near East and Europe, with each showing north-south clines bridged only by a few populations of mainly Mediterranean origin. Our PCA differs from previous studies that showed a correlation with the map of Europe^{16,17}, which we determined is due to our study having relatively fewer central and northwestern Europeans, and more Near Easterners and eastern Europeans (SI10). We projected¹⁸ the newly sequenced and previously published^{2,6,12,19} ancient genomes onto the first two PCs inferred from present-day samples (Fig. 1B). MA1 and AG2, both Upper Paleolithic hunter-gatherers from Lake Baikal⁶ in Siberia, project at the northern end of the PCA, suggesting an “Ancient North Eurasian” meta-population (ANE). European hunter-gatherers from Spain, Luxembourg, and Sweden fall outside the genetic variation of West Eurasians in the direction of European differentiation from the Near East, with a “West European Hunter-Gatherer” (WHG) cluster including Loschbour and La Braña¹², and a “Scandinavian Hunter-Gatherer” (SHG) cluster including the Motala individuals and ~5,000 year old hunter-gatherers from the Swedish Pitted Ware Culture². An “Early European Farmer” (EEF) cluster includes Stuttgart, the ~5,300 year old Tyrolean Iceman¹⁹ and a ~5,000 year old southern Swedish farmer², and is near present-day Sardinians^{2,19}.

PCA gradients of genetic variation may arise under very different histories²⁰. To test if they reflect population mixture events or are entirely due to genetic drift within West Eurasia, we computed an f_4 -statistic¹⁸ that tests whether the ancient MA1 from Siberia shares more alleles with a *Test* West Eurasian population or with Stuttgart. We find that $f_4(\text{Test}, \text{Stuttgart}; \text{MA1}, \text{Chimp})$ is positive for many West Eurasians, which must be due to variable degrees of admixture with ancient populations related to MA1 (Extended Data Fig. 4). We also find that $f_4(\text{Test}, \text{Stuttgart}; \text{Loschbour}, \text{Chimp})$ is nearly always positive in Europeans and always negative in Near Easterners, indicating that Europeans have more ancestry from ancient populations related to Loschbour than do Near Easterners (Extended Data Fig. 4). To investigate systematically the history of population mixture in West Eurasia, we computed all possible statistics of the form $f_3(X; \text{Ref}_1, \text{Ref}_2)$ (SI11). An f_3 -statistic is expected to be positive if no admixture has taken place, but if X is admixed between populations related to Ref_1 and Ref_2 , it can be negative⁵. We tested all possible pairs of $\text{Ref}_1, \text{Ref}_2$ chosen from the list of 192 present-day populations with at least four individuals, and five ancient genomes (SI11). The lowest f_3 -statistics for Europeans are usually negative (93% are >4 standard errors below zero using a standard error from a block jackknife^{5,21}). The most negative statistic (Table 1) always involves at least one ancient individual as a reference, and for Europeans it is nearly always significantly lower than the most negative statistic obtained using only present-day populations as references (SI11). MA1 is a better surrogate (Extended Data Fig. 5) for Ancient North Eurasian ancestry than the Native American Karitiana who were first used to represent this component of ancestry in Europe^{4,5}. Motala12 never appears as one of the references, suggesting that SHG may not be a source for Europeans. Instead, present-day European populations usually have their lowest f_3 with either the (EEF, ANE) or (WHG, Near East) pair (SI11, Extended Data Table 1). For Near Easterners, the lowest f_3 -statistic always takes as references Stuttgart and a population from Africa, the Americas, South Asia, or MA1 (Table 1), reflecting the fact that both Stuttgart and present-day Near Easterners harbor ancestry from ancient Near Easterners. Extended Data Fig. 6 plots statistics of the form $f_4(\text{West Eurasian } X, \text{Chimp}; \text{Ancient}_1, \text{Ancient}_2)$ onto a map, showing strong gradients in the relatedness to Stuttgart (EEF), Loschbour (WHG) and MA1 (ANE).

We determined formally that a minimum of three source ancestral populations are needed to explain the data for many European populations taken together by studying the correlation

patterns of all possible statistics of the form $f_4(\text{Test}_{\text{base}}, \text{Test}_i; O_{\text{base}}, O_j)$ (SI12). Here $\text{Test}_{\text{base}}$ is a reference European population and Test_i the set of all other European Test populations; O_{base} is a reference outgroup population, and O_i the set of other outgroups (ancient DNA samples, Onge, Karitiana, and Mbuti). The rank of the (i, j) matrix reflects the minimum number of source populations that contributed to the Test populations^{22,23}. For a pool of 23 Test populations comprising most present-day Europeans, this analysis rejects descent from just two sources ($P < 10^{-12}$ by a Hotelling T-test²³). However, three source populations are consistent with the data after excluding the Spanish who have evidence for African admixture²⁴⁻²⁶ ($P = 0.019$, not significant after multiple-hypothesis correction). Our finding of at least three source populations is also qualitatively consistent with the results from ADMIXTURE (SI9), PCA (Fig. 1B, SI10) and f -statistics (Extended Data Table 1, Extended Data Fig. 6, SI11, SI12). We caution that the finding of three sources could be consistent with a larger number of mixture events, as the method cannot distinguish between one or more mixture events if they are from the same set of sources. Our analysis also does not assume that the inferred source populations were themselves unadmixed; indeed, the positive $f_4(\text{Stuttgart}, X; \text{Loschbour}, \text{Chimp})$ statistics obtained when X is a Near Eastern population (Extended Data Table 1) implies that EEF had some WHG-related ancestry, which we show in SI13 was at least 0% and less than 45%.

Motivated by the evidence of at least three source populations for present-day Europeans, we set out to develop a model consistent with our data. To constrain our search space for modeling, we first studied f_4 -statistics comparing the ancient individuals from Europe and Siberia and diverse eastern non-African groups (Oceanians, East Asians, Siberians, Native Americans, and Onge from the Andaman Islands²⁷) (SI14). We find that: (1) Loschbour (WHG) and Stuttgart (EEF) share more alleles with each other than either does with MA1 (ANE), as might be expected by geography, but MA1 shares more alleles with Loschbour than with Stuttgart, indicating a link between Eurasian hunter-gatherers to the exclusion of European farmers; (2) Eastern non-Africans share more alleles with Eurasian hunter-gatherers (MA1, Loschbour, La Braña, and Motala12) than with Stuttgart; (3) Every eastern non-African population except for Native Americans and Siberians is equally closely related to diverse Eurasian hunter-gatherers, but Native Americans and Siberians share more alleles with MA1 than with European hunter-gatherers; and (4) Eurasian hunter-gatherers and Stuttgart both share more alleles with Native

Americans than with other eastern non-Africans. We use the ADMIXTUREGRAPH¹⁸ software to search for a model of population relationships (a tree structure augmented by admixture events) that is consistent with these observations. We explored models with 0, 1, or 2 admixture events in the ancestry of the three ancient source populations and eastern non-Africans, and identified a single model with two admixture events that fit the data. The successful model (Fig. 2A) includes the previously reported gene flow into Native Americans from an MA1-like population⁶, as well as the novel inference that Stuttgart is partially ($44 \pm 10\%$) derived from a “Basal Eurasian” lineage that split prior to the separation of eastern non-Africans from the common ancestor of WHG and ANE. If this model is accurate, the ANE/WHG split must have occurred $>24,000$ years ago since this is the age⁶ of MA1 and this individual is on the ANE lineage. The WHG must then have split from eastern non-Africans $>40,000$ years ago, as this is the age of the Chinese Tianyuan sample which clusters with eastern non-Africans to the exclusion of Europeans²⁸. The Basal Eurasian split would then have to be even older. A Basal Eurasian lineage in the Near East is plausible given the presence of anatomically modern humans in the Levant²⁹ ~ 100 thousand years ago and African-related tools likely made by modern humans in Arabia^{30,31}. Alternatively, evidence for gene flow between the Near East and Africa³², and African morphology in pre-farming Natufians³³ from Israel, may also be consistent with the population representing a later movement of humans out of Africa and into the Near East.

We tested the robustness of the ADMIXTUREGRAPH model in various ways. First, we verified that Stuttgart and the Iceman (EEF), and Loschbour and LaBraña (WHG) can be formally fit as clades (SI14). We also used the unsupervised MixMapper⁴ (SI15) and TreeMix³⁴ software (SI16) to fit graph models; both found all the same admixture events. The statistics supporting our key inferences about history also provide consistent results when restricted to transversions polymorphisms not affected by ancient DNA damage, and when repeated with whole-genome sequencing data that is not affected by SNP ascertainment bias³⁵ (Extended Data Table 2).

We next fit present-day European populations into our working model. We found that few European populations could be fit as 2-way mixtures, but nearly all were compatible with being 3-way mixtures of ANE/EEF/WHG (SI14). Mixture proportions (Fig. 2B; Extended Data Table 3) inferred via our model are consistent with those from an independent method that relates

European populations to diverse outgroups using f_4 -statistics while making much weaker modeling assumptions (only assuming that MA1 is an unmixed descendent of ANE, Loschbour of WHG, and Stuttgart of EEF; SI17). These analyses allow us to infer that EEF ancestry in Europe today ranges from ~30% in the Baltic region to ~90% in the Mediterranean, a gradient that is also consistent with patterns of identity-by-descent (IBD) sharing³⁶ (SI18) and chromosome painting³⁷ (SI19) in which Loschbour shares more segments with northern Europeans and Stuttgart with southern Europeans. Our estimates suggest that Southern Europeans inherited their European hunter-gatherer ancestry mostly via EEF ancestors (Extended Data Fig. 6), while Northern Europeans acquired up to 50% additional WHG ancestry. Europeans have a larger proportion of WHG than ANE ancestry ($WHG/(WHG+ANE) = 0.6-0.8$) with the ANE ancestry never being larger than ~20%. (By contrast, in the Near East there is no detectible WHG ancestry, but substantial ANE ancestry, up to ~29% in the North Caucasus) (SI14). While ANE ancestry was not as pervasive in Europe during the agricultural transition as it is today (we do not detect it in either Loschbour or Stuttgart), it was already present, since MA1 shares more alleles with Motala12 (SHG) than with Loschbour, and Motala12 fits as a mixture of 81% WHG and 19% ANE (SI14).

Two sets of European populations are poor fits. Sicilians, Maltese, and Ashkenazi Jews have EEF estimates beyond the 0-100% interval (SI17) and cannot be jointly fit with other Europeans (SI14). These populations may have more Near Eastern ancestry than can be explained via EEF admixture (SI14), consistent with their falling in the gap between European and Near Eastern populations in Fig. 1B. Finns, Mordovians and Russians from northeastern Europe also do not fit (SI14; Extended Data Table 3). To better understand this, we plotted $f_4(X, \text{Bedouin2}; \text{Han}, \text{Mbuti})$ against $f_4(X, \text{Bedouin2}; \text{MA1}, \text{Mbuti})$. These statistics measure the degree of a European population's allele sharing with Han Chinese or MA1 (Extended Data Fig. 7). Europeans fall on a line of slope >1 in the plot of these two statistics. However, northeastern Europeans including Chuvash and Saami (which we add in to the analysis) fall away from this line in the direction of East Asians. This is consistent with East Asian (most likely Siberian) gene flow into northeastern Europeans, some of which may be more recent³⁸ than the original ANE admixture (SI14).

Three questions seem particularly important to address in follow-up work. Where did the EEF obtain their WHG ancestry? Southeastern Europe is a candidate as it lies along the path from Anatolia into central Europe³⁹. When and where the ancestors of present-day Europeans first acquire their ANE ancestry? Based on discontinuity in mtDNA haplogroup frequencies, this may have occurred ~5,500-4,000 years ago⁴⁰ in Central Europe. When and where did Basal Eurasians mix into the ancestors of the EEF? An important aim for future work should be to collect DNA from additional ancient samples to illuminate these transformations.

Methods Summary

We extracted DNA from nine sets of ancient human remains and converted the extracts into Illumina sequencing libraries in dedicated clean rooms. We assessed whether sequences for these libraries were consistent with genuine ancient DNA by searching for characteristic deaminations at the ends of molecules^{7,8}. We also tested for contamination by searching for evidence of mixture of DNA from multiple individuals. For large-scale shotgun sequencing we used libraries that we made in the presence of the enzymes Uracil-DNA-glycosylase and endonuclease VIII, which reduce the rate of ancient DNA-induced errors. After removal of duplicated molecules, we called consensus genotypes for the high coverage samples using the Genome Analysis Toolkit⁴¹. We merged the data with published ancient genomes, as well as with 2,345 present-day humans from 203 populations genotyped at 594,924 autosomal single nucleotide polymorphisms. We visualized population structure using Principal Component Analysis¹⁵ and ADMIXTURE¹⁴. To make inferences about population history, we used methods that can analyze allele frequency correlation statistics to detect population mixture⁵; that can estimate mixture proportions in the absence of accurate ancestral populations; that can infer the minimum number of source populations for a collection of tests population²³; and that can assess formally the fit of genetic data to models of population history⁵.

1. Bramanti, B. *et al.* Genetic discontinuity between local hunter-gatherers and Central Europe's first farmers. *Science* **326**, 137-140, (2009).
2. Skoglund, P. *et al.* Origins and genetic legacy of Neolithic farmers and hunter-gatherers in Europe. *Science* **336**, 466-469, (2012).
3. Haak, W. *et al.* Ancient DNA from European early Neolithic farmers reveals their Near Eastern affinities. *PLoS Biol.* **8**, e1000536, (2010).
4. Lipson, M. *et al.* Efficient moment-based inference of admixture parameters and sources of gene flow. *Mol. Biol. Evol.* **30**, 1788-1802, (2013).

5. Patterson, N. *et al.* Ancient admixture in human history. *Genetics* **192**, 1065-1093, (2012).
6. Raghavan, M. *et al.* Upper Palaeolithic Siberian genome reveals dual ancestry of Native Americans. *Nature* **505**, 87-91, (2014).
7. Krause, J. *et al.* A complete mtDNA genome of an early modern human from Kostenki, Russia. *Curr. Biol.* **20**, 231-236, (2010).
8. Sawyer, S., Krause, J., Guschanski, K., Savolainen, V. & Pääbo, S. Temporal patterns of nucleotide misincorporations and DNA fragmentation in ancient DNA. *PLoS ONE* **7**, e34131, (2012).
9. Haak, W. *et al.* Ancient DNA from the first European farmers in 7500-Year-old Neolithic sites. *Science* **310**, 1016-1018, (2005).
10. Skoglund, P., Storå, J., Götherström, A. & Jakobsson, M. Accurate sex identification of ancient human remains using DNA shotgun sequencing. *J. Archaeol. Sci.* **40**, 4477-4482, (2013).
11. Soares, P. *et al.* The Archaeogenetics of Europe. *Curr. Biol.* **20**, R174-R183, (2010).
12. Olalde, I. *et al.* Derived immune and ancestral pigmentation alleles in a 7,000-year-old Mesolithic European. *Nature* **507**, 225-228, (2014).
13. Perry, G. H. *et al.* Diet and the evolution of human amylase gene copy number variation. *Nat. Genet.* **39**, 1256-1260, (2007).
14. Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* **19**, 1655-1664, (2009).
15. Patterson, N., Price, A. L. & Reich, D. Population structure and eigenanalysis. *PLoS Genet.* **2**, e190, (2006).
16. Lao, O. *et al.* Correlation between genetic and geographic structure in Europe. *Curr. Biol.* **18**, 1241-1248, (2008).
17. Novembre, J. *et al.* Genes mirror geography within Europe. *Nature* **456**, 98-101, (2008).
18. Reich, D., Thangaraj, K., Patterson, N., Price, A. L. & Singh, L. Reconstructing Indian population history. *Nature* **461**, 489-494, (2009).
19. Keller, A. *et al.* New insights into the Tyrolean Iceman's origin and phenotype as inferred by whole-genome sequencing. *Nat. Commun.* **3**, 698, (2012).
20. Novembre, J. & Stephens, M. Interpreting principal component analyses of spatial population genetic variation. *Nat. Genet.* **40**, 646-649, (2008).
21. Busing, F. T. A., Meijer, E. & Leeden, R. Delete-m Jackknife for Unequal m. *Statistics and Computing* **9**, 3-8, (1999).
22. Moorjani, P. *et al.* Genetic evidence for recent population mixture in India. *Am. J. Hum. Genet.* **93**, 422-438, (2013).
23. Reich, D. *et al.* Reconstructing Native American population history. *Nature* **488**, 370-374, (2012).
24. Botigué, L. R. *et al.* Gene flow from North Africa contributes to differential human genetic diversity in southern Europe. *Proc. Natl. Acad. Sci. USA* **110**, 11791-11796, (2013).
25. Cerezo, M. *et al.* Reconstructing ancient mitochondrial DNA links between Africa and Europe. *Genome Res.* **22**, 821-826, (2012).
26. Moorjani, P. *et al.* The history of African gene flow into southern Europeans, Levantines, and Jews. *PLoS Genet.* **7**, e1001373, (2011).
27. Thangaraj, K. *et al.* Reconstructing the origin of Andaman Islanders. *Science* **308**, 996-996, (2005).
28. Fu, Q. *et al.* DNA analysis of an early modern human from Tianyuan Cave, China. *Proc. Natl. Acad. Sci. USA* **110**, 2223-2227, (2013).
29. Bar-Yosef, O. *The chronology of the Middle Paleolithic of the Levant*. 39-56 (New York: Plenum Press, 1998).
30. Armitage, S. J. *et al.* The southern route "Out of Africa": evidence for an early expansion of modern humans into Arabia. *Science* **331**, 453-456, (2011).

31. Rose, J. I. *et al.* The Nubian Complex of Dhofar, Oman: an African middle stone age industry in Southern Arabia. *PLoS ONE* **6**, e28239, (2011).
32. Haber, M. *et al.* Genome-Wide Diversity in the Levant Reveals Recent Structuring by Culture. *PLoS Genet* **9**, e1003316, (2013).
33. Brace, C. L. *et al.* The questionable contribution of the Neolithic and the Bronze Age to European craniofacial form. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 242-247, (2006).
34. Pickrell, J. K. & Pritchard, J. K. Inference of population splits and mixtures from genome-wide Allele frequency data. *PLoS Genet.* **8**, e1002967, (2012).
35. Prufer, K. *et al.* The complete genome sequence of a Neanderthal from the Altai Mountains. *Nature* **505**, 43-49, (2014).
36. Ralph, P. & Coop, G. The geography of recent genetic ancestry across Europe. *PLoS Biol.* **11**, e1001555, (2013).
37. Lawson, D. J., Hellenthal, G., Myers, S. & Falush, D. Inference of Population Structure using Dense Haplotype Data. *PLoS Genet.* **8**, e1002453, (2012).
38. Hellenthal, G. *et al.* A genetic atlas of human admixture history. *Science* **343**, 747-751, (2014).
39. Bellwood, P. *First Farmers: The Origins of Agricultural Societies*. (Wiley-Blackwell, 2004).
40. Brandt, G. *et al.* Ancient DNA reveals key stages in the formation of central European mitochondrial genetic diversity. *Science* **342**, 257-261, (2013).

Supplementary Information is linked to the online version of the paper. The fully public version of the Human Origins dataset can be found at http://genetics.med.harvard.edu/reichlab/Reich_Lab/Datasets.html. The full version of the dataset (including additional samples) is available to researchers who send a signed letter to DR indicating that they will abide by specified usage conditions.

Acknowledgments We are grateful to Cynthia Beall, Neil Bradman, Amha Gebremedhin, Damian Labuda, Maria Nelis and Anna Di Rienzo for sharing DNA samples; to Detlef Weigel, Christa Lanz, Verena Schünemann, Peter Bauer and Olaf Riess for support and access to DNA sequencing facilities; to Philip Johnson for advice on contamination estimation; and to Pontus Skoglund for sharing the graphics software that we used to generate Extended Data Fig. 6. We thank Kenneth Nordtvedt for alerting us about the existence of newly discovered Y-chromosome SNPs. The collections and methods for the Population Reference Sample (POPRES) discussed in SI18 are described in ref.⁴², and the dataset used for our analyses was obtained from dbGaP at http://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000145.v4.p2 through dbGaP accession number phs000145.v1.p2. We thank all the volunteers who donated DNA; the staff of the Unità Operativa Complessa di Medicina Trasfusionale, Azienda Ospedaliera Umberto I, Siracusa, Italy for assistance in sample collection; and The National Laboratory for

the Genetics of Israeli Populations for facilitating access to DNA. We thank colleagues at the Applied Genomics at the Children's Hospital of Philadelphia, especially Hakon Hakonarson, Cecilia Kim, Kelly Thomas, and Cuiping Hou, for genotyping samples on the Human Origins array. JK is grateful for support from DFG grant # KR 4015/1-1, the Carl-Zeiss Foundation and the Baden Württemberg Foundation. SP acknowledges support from the Presidential Innovation Fund of the Max Planck Society. JGS acknowledges use of the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by NSF grant number OCI-1053575. EB and OB were supported by RFBR grants 13-06-00670, 13-04-01711, 13-04-90420 and by the Molecular and Cell Biology Program of the Presidium, Russian Academy of Sciences. BM was supported by grants OTKA 73430 and 103983. ASaj was supported by a Finnish Professorpool (Paulo Foundation) Grant. The Lithuanian sampling was supported by the LITGEN project (VP1-3.1-ŠMM-07-K-01-013), funded by the European Social Fund under the Global Grant Measure. AS was supported by Spanish grants SAF2008-02971 and EM 2012/045. OU was supported by Ukrainian SFFS grant F53.4/071. SAT was supported by NIH Pioneer Award 8DP1ES022577-04 and NSF HOMINID award BCS-0827436. KT was supported by an Indian CSIR Network Project (GENESIS: BSC0121). LS was supported by an Indian CSIR Bhatnagar Fellowship. RV, MM, JP and EM were supported by the European Union Regional Development Fund through the Centre of Excellence in Genomics to the Estonian Biocentre and University of Tartu and by a Estonian Basic Research grant SF0270177As08. MM was additionally supported by Estonian Science Foundation grant #8973. JGS and MS were supported by NIH grant GM40282. PHS and EEE were supported by NIH grants HG004120 and HG002385. DR and NP were supported by NSF HOMINID award BCS-1032255 and NIH grant GM100233. DR and EEE are Howard Hughes Medical Institute investigators.

Author contributions

BB, EEE, JBu, MS, SP, JKe, DR and JKr supervised the study. IL, NP, AM, GR, SM, KK, PHS, JGS, SC, ML, QF, HL, CdF, KP, WH, MMey and DR analyzed genetic data. FH, EF, DD, MF, J-MG, JW, AC and JKr obtained human remains. AM, CE, RBo, KB, SS, CP, NR and JKr processed ancient DNA. IL, NP, SN, NR, GA, HAB, GBa, EB, OB, RBa, GBe, HB-A, JBe, FBe, CMB, FBr, GBJB, FC, MC, DECC, DCor, LD, GvD, SD, J-MD, SAF, IGR, MG, MH, BH, TH, UH, ARJ, SK-Y, RKh, EK, RKi, TK, WK, VK, AK, LL, SL, TL, RWM, BM, EM, JMol, JMou,

KN, DN, TN, LO, JP, FP, OLP, VR, FR, IR, RR, HS, ASaj, ASal, EBS, ATar, DT, ST, IU, OU, RVa, MVi, MVo, CW, LY, PZ, TZ, CC, MGT, AR-L, SAT, LS, KT, RVi, DCom, RS, MMet, SP and DR assembled the genotyping dataset. IL, NP, DR and JKr wrote the manuscript with help from all co-authors.

Author information

The aligned sequences are available through the Sequence Read Archive (SRA) under accession numbers that will be made available upon publication. The authors declare competing financial interests: UH is an employee of Illumina, TL is an employee of AMGEN, and JM is an employee of 23andMe. Correspondence and requests for materials should be addressed to David Reich (reich@genetics.med.harvard.edu) or Johannes Krause (johannes.krause@uni-tuebingen.de).

Table 1: Lowest f_3 -statistics for each West Eurasian population

<i>Ref₁</i>	<i>Ref₂</i>	Target for which these two references give the lowest $f_3(X; Ref_1, Ref_2)$
WHG	EEF	Sardinian ^{***}
WHG	Near East	Basque, Belarusian, Czech, English, Estonian, Finnish, French_South, Icelandic, Lithuanian, Mordovian, Norwegian, Orcadian, Scottish, Spanish, Spanish_North, Ukrainian
EEF	ANE	Abkhasian ^{***} , Albanian, Ashkenazi_Jew ^{****} , Bergamo, Bulgarian, Chechen ^{****} , Croatian, Cypriot ^{***} , Druze ^{**} , French, Greek, Hungarian, Lezgin, MA1, Maltese, Sicilian, Turkish_Jew, Tuscan
EEF	Native American	Adygei, Balkar, Iranian, Kumyk, North_Ossetian, Turkish
EEF	African	BedouinA, BedouinB†, Jordanian, Lebanese, Libyan_Jew, Moroccan_Jew, Palestinian, Saudi ^{****} , Syrian, Tunisian_Jew ^{***} , Yemenite_Jew ^{***}
EEF	South Asian	Armenian, Georgian ^{****} , Georgian_Jew [*] , Iranian_Jew ^{***} , Iraqi_Jew ^{***}

Note: WHG = Loschbour or LaBraña; EEF=Stuttgart; ANE=MA1; Native American=Piapoco; African=Esan, Gambian, or Kgalagadi; South Asian=GujaratiC or Vishwabrahmin. Statistics are negative with $Z < -4$ unless otherwise noted: † (positive) or *, **, ***, ****, to indicate Z less than 0, -1, -2, and -3 respectively. The complete list of statistics can be found in Extended Data Table 1.

Figure Legends

Figure 1: Map of West Eurasian populations and Principal Component Analysis. (a)

Geographical locations of ancient and present-day samples, with color coding matching the PCA. We show all sampling locations for each population, which results in multiple points for some populations (e.g., Spain). (b) PCA on all present-day West Eurasians, with the ancient and selected eastern non-Africans projected. European hunter-gatherers fall beyond present-day Europeans in the direction of European differentiation from the Near East. Stuttgart clusters with other Neolithic Europeans and present-day Sardinians. MA1 falls outside the variation of present-day West Eurasians in the direction of southern-northern differentiation along dimension 2 and between the European and Near Eastern clines along dimension 1.

Figure 2: Modeling of West Eurasian population history. (a) A three-way mixture model that is a statistical fit to the data for many European populations, ancient DNA samples, and non-European populations. Present-day samples are colored in blue, ancient samples in red, and reconstructed ancestral populations in green. Solid lines represent descent without mixture, and dashed lines represent admixture events. For the two mixture events relating the highly divergent ancestral populations, we print estimates for the mixture proportions as well as one standard error. (b) We plot the proportions of ancestry from each of three inferred ancestral populations (EEF, ANE and WHG) as inferred from the model-based analysis.

Methods

Archeological context, sampling and DNA extraction

The Loschbour sample stems from a male skeleton excavated in 1935 at the Loschbour rock shelter in Heffingen, Luxembourg. The skeleton was AMS radiocarbon dated to $7,205 \pm 50$ years before present (OxA-7738; 6,220-5,990 cal BC)⁴³. At the Palaeogenetics Laboratory in Mainz, material for DNA extraction was sampled from a molar (M48) after irradiation with UV-light, surface removal, and pulverization in a mixer mill. DNA extraction took place in the palaeogenetics facilities in the Institute for Archaeological Sciences at the University of Tübingen. Three extracts were made in total, one from 80 mg of powder using an established silica based protocol⁴⁴ and two additional extracts from 90 mg of powder each with a protocol optimized for the recovery of short DNA molecules⁴⁵.

The Stuttgart sample was taken from a female skeleton excavated in 1982 at the site Viesenhäuser Hof, Stuttgart-Mühlhausen, Germany. It was attributed to the Linearbandkeramik (5,500-4,800 BC) through associated pottery artifacts and the chronology was corroborated by radiocarbon dating of the stratigraphy⁴⁶. Both sampling and DNA extraction took place in the Institute for Archaeological Sciences at the University of Tübingen. The M47 molar was removed and material from the inner part was sampled with a sterile dentistry drill. An extract was made using 40 mg of bone powder⁴⁵.

The Motala individuals were recovered from the site of Kanaljorden in the town of Motala, Östergötland, Sweden, excavated between 2009 and 2013. The human remains at this site are represented by several adult skulls and one infant skeleton. All individuals are part of a ritual deposition at the bottom of a small lake. Direct radiocarbon dates on the remains range between $7,013 \pm 76$ and $6,701 \pm 64$ BP (6,361-5,516 cal BC), corresponding to the late Middle Mesolithic of Scandinavia. Samples were taken from the teeth of the nine best preserved skulls, as well as a femur and tibia. Bone powder was removed from the inner parts of the teeth or bones with a sterile dentistry drill. DNA from 100 mg of bone powder was extracted⁴⁷ in the ancient DNA laboratory of the Archaeological Research Laboratory, Stockholm.

Library preparation

Illumina sequencing libraries were prepared using either double- or single-stranded library preparation protocols^{48,49} (SI1). For high-coverage shotgun sequencing libraries, a DNA repair step with Uracil-DNA-glycosylase (UDG) and endonuclease VIII (endo VIII) treatment was included in order to remove uracil residues⁵⁰. Size fractionation on a PAGE gel was also performed in order to remove longer DNA molecules that are more likely to be contaminants⁴⁹. Positive and blank controls were carried along during every step of library preparation.

Shotgun sequencing and read processing

All non-UDG-treated libraries were sequenced either on an Illumina Genome Analyzer IIX with $2 \times 76 + 7$ cycles for the Loschbour and Motala libraries, or on an Illumina MiSeq with $2 \times 150 + 8 + 8$ cycles for the Stuttgart library. We followed the manufacturer's protocol for multiplex sequencing. Raw overlapping forward and reverse reads were merged and filtered for quality⁵¹ and mapped to the human reference genome (hg19/GRCh37/1000Genomes) using the Burrows-Wheeler Aligner (BWA)⁵² (SI2). For deeper sequencing, UDG-treated libraries of Loschbour were sequenced on 3 Illumina HiSeq 2000 lanes with 50-bp single-end reads, 8 Illumina HiSeq 2000 lanes of 100-bp paired-end reads and 8 Illumina HiSeq 2500 lanes of 101-bp paired-end reads. The UDG-treated library for Stuttgart was sequenced on 8 HiSeq 2000 lanes and 101-bp paired-end reads. The UDG-treated libraries for Motala were sequenced on 8 HiSeq 2000 lanes of 100-bp paired-end reads, with 4 lanes each for two pools (one of 3 individuals and one of 4 individuals). We also sequenced an additional 8 HiSeq 2000 lanes for Motala12, the Motala sample with the highest percentage of endogenous human DNA.

Enrichment of mitochondrial DNA and sequencing

Non-UDG-treated libraries of Loschbour and all Motala samples were enriched for human mitochondrial DNA using a bead-based capture approach with present-day human DNA as bait⁵³ to test for DNA preservation and mtDNA contamination. UDG-treatment was omitted in order to allow characterization of damage patterns typical for ancient DNA⁸. The captured libraries were sequenced on an Illumina Genome Analyzer IIX platform with $2 \times 76 + 7$ cycles and the resulting reads were merged and quality filtered⁵¹. The sequences were mapped to the Reconstructed Sapiens Reference Sequence, RSRS⁵⁴, using a custom iterative mapping assembler, MIA⁵⁵ (SI4).

Contamination estimates

We assessed if the sequences had the characteristics of authentic ancient DNA using four approaches. First we searched for evidence of contamination by determining whether the sequences mapping to the mitochondrial genome were consistent with deriving from more than one individual^{55,56}. Second, for the high-coverage Loschbour and Stuttgart genomes, we used a maximum-likelihood-based estimate of autosomal contamination that uses variation at sites that are fixed in the 1000 Genomes data to estimate error, heterozygosity and contamination⁵⁷ simultaneously. Third, we estimated contamination based on the rate of polymorphic sites on the X chromosome of the male Loschbour individual⁵⁸ (SI3). Fourth, we analyzed non-UDG treated reads mapping to the RSRS to search for aDNA-typical damage patterns resulting in C→T changes at the 5'-end of the molecule⁸ (SI3).

Phylogenetic analysis of the mitochondrial genomes

All nine complete mitochondrial genomes that fulfilled the criteria of authenticity were assigned to haplogroups using Haplofind⁵⁹. A Maximum Parsimony tree including present day humans and previously published ancient mtDNA sequences was generated with MEGA⁶⁰. The effect of branch shortening due to a lower number of substitutions in ancient lineages was studied by calculating the nucleotide edit distance to the root for all haplogroup R sequences (SI4).

Sex Determination and Y-chromosome Analysis

We assessed the sex of all sequenced individuals by using the ratio of (chrY) to (chrY+chrX) aligned reads¹⁰. We downloaded a list of Y-chromosome SNPs curated by the International Society of Genetic Genealogy (ISOGG, <http://www.isogg.org>) v. 9.22 (accessed Feb. 18, 2014) and determined the state of the ancient individuals at positions where a single allele was observed and MAPQ≥30. We excluded C/G or A/T SNPs due to uncertainty about the polarity of the mutation in the database. The ancient individuals were assigned haplogroups based on their derived state (SI5). We also used BEAST v1.7.51⁶¹ to assess the phylogenetic position of Loschbour using 623 males from around the world with 2,799 variant sites across 500kb of non-recombining Y-chromosome sequence⁶² (SI5).

Estimation of Neandertal admixture

We estimate Neandertal admixture in ancient individuals with the f_4 -ratio or S -statistic^{5,63,64} $\hat{\alpha} = f_4(\text{Altai}, \text{Denisova}; \text{Test}, \text{Yoruba}) / f_4(\text{Altai}, \text{Denisova}; \text{Vindija}, \text{Yoruba})$ which uses whole genome data from Altai, a high coverage (52×) Neanderthal genome sequence³⁵, Denisova, a high coverage sequence⁴⁹ from another archaic human population (31×), and Vindija, a low coverage (1.3×) Neanderthal genome from a mixture of three Neanderthal individuals from Vindija Cave in Croatia⁶³.

Inference of demographic history and inbreeding

We used the Pairwise Sequentially Markovian Coalescent (PSMC)⁶⁵ to infer the size of the ancestral population of Stuttgart and Loschbour. This analysis requires high quality diploid genotype calls and cannot be performed in the low-coverage Motala samples. To determine whether the low effective population size inferred for Loschbour is due to recent inbreeding, we plotted the time-to-most-recent common ancestor (TMRCA) along each of chr1-22 to detect runs of low TMRCA.

Analysis of segmental duplications and copy number variants

We built read-depth based copy number maps for the Loschbour, Stuttgart and Motala12 genomes in addition to the Denisova and Altai Neanderthal genome and 25 deeply sequenced modern genomes³⁵ (SI7). We built these maps by aligning reads, subdivided into their non-overlapping 36-bp constituents, against the reference genome using the mrsFAST aligner⁶⁶, and renormalizing read-depth for local GC content. We estimated copy numbers in windows of 500 unmasked base pairs slid at 100 bp intervals across the genome. We called copy number variants using a scale space filter algorithm. We genotyped variants of interest and compared the genotypes to those from individuals sequenced as part of the 1000 Genomes Project⁶⁷.

Phenotypic inference

We inferred likely phenotypes (SI8) by analyzing DNA polymorphism data in the VCF format⁶⁸ using VCFtools (<http://vcftools.sourceforge.net/>). For the Loschbour and Stuttgart individuals, we included data from sites not flagged as LowQuality, with genotype quality (GQ) of ≥ 30 , and SNP quality (QUAL) of ≥ 50 . For Motala12, which is of lower coverage, we included sites

having at least $2\times$ coverage and passed visual inspection of the local alignment using samtools tview (<http://samtools.sourceforge.net>)⁶⁹

Human Origins dataset curation

The Human Origins array consists of 14 panels of SNPs for which the ascertainment is well known^{5,70}. All population genetics analysis were carried out on a set of 594,924 autosomal SNPs, after restricting to sites that had >90% completeness across 7 different batches of sequencing, and that had >97.5% concordance with at least one of two subsets of samples for which whole genome sequencing data was also available. The total dataset consists of 2,722 individuals, which we filtered to 2,345 individuals (203 populations) after removing outlier individuals or relatives based on visual inspection of PCA plots^{15,71} or model-based clustering analysis¹⁴. Whole genome amplified (WGA) individuals were not used in analysis, except for a Saami individual who we forced in because of the special interest of this population for Northeastern European population history (Extended Data Fig. 7).

ADMIXTURE analysis

We merged all Human Origins genotype data with whole genome sequencing data from Loschbour, Stuttgart, MA1, Motala12, Motala_merge, and LaBrana. We then thinned the resulting dataset to remove SNPs in linkage-disequilibrium with PLINK 1.07⁷², using a window size of 200 SNPs advanced by 25 SNPs and an r^2 threshold of 0.4. We ran ADMIXTURE 1.23^{14,73} for 100 replicates with different starting random seeds, default 5-fold cross-validation, and varying the number of ancestral populations K between 2 and 20. We assessed clustering quality using CLUMPP⁷⁴. We used the ADMIXTURE results to identify a set of 59 “West Eurasian” (European/Near Eastern) populations based on values of a “West Eurasian” ancestral population at $K=3$ (SI9). We also identified 15 populations for use as “non-West Eurasian outgroups” based on their having at least 10 individuals and no evidence of European or Near Eastern admixture at $K=11$, the lowest K for which Near Eastern/European-maximized ancestral populations appeared consistently across all 100 replicates.

Principal Components Analysis

We used *smartpca*¹⁵ (version: 10210) from EIGENSOFT^{71,75} 5.0.1 to carry out Principal Components Analysis (PCA) (SI10). We performed PCA on a subset on individuals and then projected others using the *lsqproject: YES* option that gives an unbiased inference of the position of samples even in the presence of missing data (especially important for ancient DNA).

***f*₃-statistics**

We use the *f*₃-statistic⁵ $f_3(\text{Test}; \text{Ref}_1, \text{Ref}_2) = \frac{1}{N} \sum_{i=1}^N (t_i - r_{1,i}) (t_i - r_{2,i})$, where t_i , $r_{1,i}$ and $r_{2,i}$ are the allele frequencies for the i^{th} SNP in populations *Test*, *Ref*₁, *Ref*₂, respectively, to determine if there is evidence that the *Test* population is derived from admixture of populations related to *Ref*₁ and *Ref*₂ (SI11). A significantly negative statistic provides unambiguous evidence of mixture in the *Test* population⁵. We allow *Ref*₁ and *Ref*₂ to be any Human Origins population with 4 or more individuals, or Loschbour, Stuttgart, MA1, Motala12, LaBrana. We assess significance of the *f*₃-statistics using a block jackknife²¹ and a block size of 5cM. We report significance as the number of standard errors by which the statistic differs from zero (Z-score). We also perform an analysis in which we constrain the reference populations to be (i) EEF (Stuttgart) and WHG (Loschbour or LaBrana), (ii) EEF and a Near Eastern population, (iii) EEF and ANE (MA1), or (iv) any two present-day populations, and compute a *Z*_{diff} score between the lowest *f*₃-statistic observed in the dataset, and the *f*₃-statistic observed for the specified pair.

***f*₄-statistics**

We analyze *f*₄-statistics⁵ of the form $f_4(A, B; C, D) = \frac{1}{N} \sum_{i=1}^N (a_i - b_i) (c_i - d_i)$ to assess if populations A, B are consistent with forming a clade in an unrooted tree with respect to C, D. If they form a clade, the allele frequency differences between the two pairs should be uncorrelated and the statistic has an expected value of 0. We set the outgroup *D* to be a sub-Saharan African population or Chimpanzee. We systematically tried all possible combinations of the ancient samples or 15 “non-West Eurasian outgroups” identified by ADMIXTURE analysis as A, B, C to determine their genetic affinities (SI14). Setting A as a present-day test population and B as either Stuttgart or BedouinB, we documented relatedness to C=(Loschbour or MA1) or C=(MA1 and Karitiana) or C=(MA1 or Han) (Extended Data Figs. 4, 5, 7). Setting C as a test population and (A, B) a pair from (Loschbour, Stuttgart, MA1) we documented differential relatedness to ancient populations (Extended Data Fig. 6). We computed *D*-statistics⁶³ using

transversion polymorphisms in whole genome sequence data³⁵ to confirm robustness to ascertainment and ancient DNA damage (Extended Data Table 2).

Minimum number of source populations for Europeans

We used *qpWave*^{22,23} to study the minimum number of source populations for a designated set of Europeans (SI12). We use f_4 -statistics of the form $X(l, r) = f_4(l_0, l; r_0, r)$ where l_0, r_0 are arbitrarily chosen “base” populations, and l, r are other populations from two sets L and R respectively. If $X(l, r)$ has rank r and there were n waves of immigration into R with no back-migration from R to L , then $r+1 \leq n$. We set L to include *Stuttgart*, *Loschbour*, *MA1*, *Onge*, *Karitiana*, *Mbuti* and R to include 23 modern European populations who fit the model of SI14 and had admixture proportions within the interval [0,1] for the method with minimal modeling assumptions (SI17).

Admixture proportions for Stuttgart in the absence of a Near Eastern ancient genome

We used *Loschbour* and *BedouinB* as surrogates for “Unknown hunter-gatherer” and Near Eastern (NE) farmer populations that contributed to *Stuttgart* (SI13). Ancient Near Eastern ancestry in *Stuttgart* is estimated by the f_4 -ratio^{5,18} $f_4(\text{Outgroup}, X; \text{Loschbour}, \text{Stuttgart}) / f_4(\text{Outgroup}, X; \text{Loschbour}, \text{NE})$. A complication is that *BedouinB* is a mixture of NE and African ancestry. We therefore subtracted²³ the effects of African ancestry using estimates of the *BedouinB* African admixture proportion from ADMIXTURE (SI9) or ALDER⁷⁶.

Admixture graph modeling

We used ADMIXTUREGRAPH⁵ (version 3110) to model population relationships between *Loschbour*, *Stuttgart*, *Onge*, and *Karitiana* using *Mbuti* as an African outgroup. We assessed model fit using a block jackknife of differences between estimated and fitted f -statistics for the set of included populations (we expressed the fit as a Z score). We determined that a model failed if $|Z| > 3$ for at least one f -statistic. A basic tree model failed and we manually amended the model to test all possible models with a single admixture event, which also failed. Further manual amendment to include 2 admixture events resulted in 8 successful models, only one of which could be amended to also fit *MA1* as an additional constraint. We successfully fit both the *Iceman* and *LaBrana* into this model as simple clades and *Motala12* as a 2-way mixture. We also fit present-day West Eurasians as clades, 2-way mixtures, or 3-way mixtures in this basic

model, achieving a successful fit for a larger number of European populations ($n=26$) as 3-way mixtures. We estimated the individual admixture proportions from the fitted model parameters. To test if fitted parameters for different populations are consistent with each other, we jointly fit all pairs of populations A and B by modifying ADMIXTUREGRAPH to add a large constant (10,000) to the variance term $f_3(A_0, A, B)$. By doing this, we can safely ignore recent gene flow within Europe that affects statistics that include both A and B .

Ancestry estimates from f_4 -ratios

We estimate EEF ancestry using the f_4 -ratio^{5,18} $f_4(\text{Mbuti}, \text{Onge}; \text{Loschbour}, \text{European}) / f_4(\text{Mbuti}, \text{Onge}; \text{Loschbour}, \text{Stuttgart})$, which produces consistent results with ADMIXTUREGRAPH (SI14). We use $f_4(\text{Stuttgart}, \text{Loschbour}; \text{Onge MA1}) / f_4(\text{Mbuti}, \text{MA1}; \text{Onge}, \text{Loschbour})$ to estimate Basal Eurasian admixture into Stuttgart. We use $f_4(\text{Stuttgart}, \text{Loschbour}; \text{Onge Karitiana}) / f_4(\text{Stuttgart}, \text{Loschbour}; \text{Onge MA1})$ to estimate ANE mixture in Karitiana (Fig. 2B). We use $f_4(\text{Test}, \text{Stuttgart}; \text{Karitiana}, \text{Onge}) / f_4(\text{MA1}, \text{Stuttgart}; \text{Karitiana}, \text{Onge})$ to lower bound ANE mixture into North Caucasian populations.

MixMapper analysis

We carried out *MixMapper* 2.0⁴ analysis, a semi-supervised admixture graph fitting technique. First, we infer a scaffold tree of populations without strong evidence of mixture relative to each other (Mbuti, Onge, Loschbour and MA1). We do not include European populations in the scaffold as all had significantly negative f_3 -statistics indicating admixture. We then ran *MixMapper* to infer the relatedness of the other ancient and present-day samples, fitting them onto the scaffold as 2- or 3-way mixtures. The uncertainty in all parameter estimates is measured by block bootstrap resampling of the SNP set (100 replicates with 50 blocks).

TreeMix analysis

We applied *TreeMix*³⁴ to Loschbour, Stuttgart, Motala12, and MA1⁶, LaBrana¹² and the Iceman¹⁹, along with the present-day samples of Karitiana, Onge and Mbuti. We restricted the analysis to 265,521 Human Origins array sites after excluding any SNPs where there were no-calls in any of the studied individuals. The tree was rooted with Mbuti and standard errors were estimated using blocks of 500 SNPs. We repeated the analysis on whole-genome sequence data,

rooting with Chimp and replacing Onge with Dai since we did not have Onge whole genome sequence data³⁵. We varied the number of migration events (m) between 0 and 5.

Inferring admixture proportions with minimal modeling assumptions

We devised a method to infer ancestry proportions from three ancestral populations (EEF, WHG, and ANE) without strong phylogenetic assumptions (SI17). We rely on 15 “non-West Eurasian” outgroups and study $f_4(\text{European}, \text{Stuttgart}; O_1, O_2)$ which equals $\alpha\beta f_4(\text{Loschbour}, \text{Stuttgart}; O_1, O_2) + \alpha(1-\beta) f_4(\text{MA1}, \text{Stuttgart}; O_1, O_2)$ if *European* has $1-\alpha$ ancestry from EEF and β , $1-\beta$ ancestry from WHG and ANE respectively. This defines a system of $\binom{15}{2} = 105$ equations with unknowns $\alpha\beta$, $\alpha(1-\beta)$, which we solve with least squares implemented in the function *lsfit* in *R* to obtain estimates of α and β . We repeated this computation 22 times dropping one chromosome at a time²⁶ to obtain block jackknife²¹ estimates of the ancestry proportions and standard errors, with block size equal to the number of SNPs per chromosome. We assessed consistency of the inferred admixture proportions with those derived from the ADMIXTUREGRAPH model based on the number of standard errors between the two (Extended Data Table 1).

Haplotype-based analyses

We used RefinedIBD from BEAGLE 4⁷⁷ with the settings *ibdtrim*=20 and *ibdwindow*=25 to study IBD sharing between Loschbour and Stuttgart and populations from the POPRES dataset⁴². We kept all IBD tracts spanning at least 0.5 centimorgans (cM) and with a LOD score >3 (SI18). We also used ChromoPainter³⁷ to study haplotype sharing between Loschbour and Stuttgart and present-day West Eurasian populations (SI19). We identified 495,357 SNPs that were complete in all individuals and phased the data using Beagle 4⁷⁷ with parameters *phase-its*=50 and *impute-its*=10. We did not keep sites with missing data to avoid imputing modern alleles into the ancient individuals. We combined ChromoPainter output for chromosomes 1-22 using ChromoCombine³⁷. We carried out a PCA of the co-ancestry matrix using fineSTRUCTURE³⁷.

41. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* **20**, 1297-1303, (2010).
42. Nelson, M. R. *et al.* The Population Reference Sample, POPRES: a resource for population, disease, and pharmacological genetics research. *Am. J. Hum. Genet.* **83**, 347-358, (2008).

43. Delsate, D., Guinet, J.-M. & Saverwyns, S. De l'ocre sur le crâne mésolithique (haplogroupe U5a) de Reuland-Loschbour (Grand-Duché de Luxembourg) ? *Bull. Soc. Préhist. Luxembourgeoise* **31**, 7-30, (2009).
44. Rohland, N. & Hofreiter, M. Ancient DNA extraction from bones and teeth. *Nat. Protocols* **2**, 1756-1762, (2007).
45. Dabney, J. *et al.* Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proceedings of the National Academy of Sciences* **110**, 15758–15763, (2013).
46. Stäuble, H. S. f. V.-u. F. d. U. F. *Häuser und absolute Datierung der Ältesten Bandkeramik.* (Habelt, 2005).
47. Yang, D. Y., Eng, B., Waye, J. S., Dudar, J. C. & Saunders, S. R. Improved DNA extraction from ancient bones using silica-based spin columns. *Am. J. Phys. Anthropol.* **105**, 539-543, (1998).
48. Meyer, M. & Kircher, M. Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harb. Protoc.* **2010**, pdb prot5448, (2010).
49. Meyer, M. *et al.* A High-Coverage Genome Sequence from an Archaic Denisovan Individual. *Science* **338**, 222-226, (2012).
50. Briggs, A. W. *et al.* Removal of deaminated cytosines and detection of in vivo methylation in ancient DNA. *Nucleic Acids Res.* **38**, e87-e87, (2010).
51. Kircher, M. in *Methods Mol. Biol. Vol. 840 Methods in Molecular Biology* 197-228 (2012).
52. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**, 1754-1760, (2009).
53. Maricic, T., Whitten, M. & Pääbo, S. Multiplexed DNA Sequence Capture of Mitochondrial Genomes Using PCR Products. *PLoS ONE* **5**, e14004, (2010).
54. Behar, Doron M. *et al.* A Copernican Reassessment of the Human Mitochondrial DNA Tree from its Root. *Am. J. Hum. Genet.* **90**, 675-684, (2012).
55. Green, R. E. *et al.* A Complete Neandertal Mitochondrial Genome Sequence Determined by High-Throughput Sequencing. *Cell* **134**, 416-426, (2008).
56. Fu, Q. *et al.* A Revised Timescale for Human Evolution Based on Ancient Mitochondrial Genomes. *Curr. Biol.* **23**, 553-559, (2013).
57. Fu, Q. & al., e. (*in preparation*), (2014).
58. Rasmussen, M. *et al.* An Aboriginal Australian Genome Reveals Separate Human Dispersals into Asia. *Science* **334**, 94-98, (2011).
59. Vianello, D. *et al.* HAPLOFIND: a new method for high-throughput mtDNA haplogroup assignment. *Hum. Mutat.* **34**, 1189-1194, (2013).
60. Tamura, K. *et al.* MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Mol. Biol. Evol.* **28**, 2731-2739, (2011).
61. Drummond, A. & Rambaut, A. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol. Biol.* **7**, 214, (2007).
62. Lippold, S. *et al.* Human paternal and maternal demographic histories: insights from high-resolution Y chromosome and mtDNA sequences. *bioRxiv*, doi: 10.1101/001792, (2014).
63. Green, R. E. *et al.* A Draft Sequence of the Neandertal Genome. *Science* **328**, 710-722, (2010).
64. Reich, D. *et al.* Genetic history of an archaic hominin group from Denisova Cave in Siberia. *Nature* **468**, 1053-1060, (2010).
65. Li, H. & Durbin, R. Inference of human population history from individual whole-genome sequences. *Nature* **475**, 493-496, (2011).
66. Hach, F. *et al.* mrsFAST: a cache-oblivious algorithm for short-read mapping. *Nat. Meth.* **7**, 576-577, (2010).

67. An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**, 56-65, (2012).
68. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-2158, (2011).
69. Li, H. The sequence alignment/map (SAM) format and SAMtools. *Bioinformatics*. **25**, 2078-2079, (2009).
70. Keinan, A., Mullikin, J. C., Patterson, N. & Reich, D. Measurement of the human allele frequency spectrum demonstrates greater genetic drift in East Asians than in Europeans. *Nat Genet* **39**, 1251-1255, (2007).
71. Price, A. L. *et al.* Principal components analysis corrects for stratification in genome-wide association studies. *Nat. Genet.* **38**, 904-909, (2006).
72. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559-575, (2007).
73. Alexander, D. & Lange, K. Enhancements to the ADMIXTURE algorithm for individual ancestry estimation. *BMC Bioinformatics* **12**, 246, (2011).
74. Jakobsson, M. & Rosenberg, N. A. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* **23**, 1801-1806, (2007).
75. Price, A. L., Zaitlen, N. A., Reich, D. & Patterson, N. New approaches to population stratification in genome-wide association studies. *Nat. Rev. Genet.* **11**, 459-463, (2010).
76. Loh, P.-R. *et al.* Inferring Admixture Histories of Human Populations Using Linkage Disequilibrium. *Genetics* **193**, 1233-1254, (2013).
77. Browning, B. L. & Browning, S. R. Improving the Accuracy and Efficiency of Identity-by-Descent Detection in Population Data. *Genetics* **194**, 459-471, (2013).

Figure 1





Figure 2

A



B

