

Opposing associations of Internet Use Disorder symptom domains with structural and functional organization of the striatum: a dimensional neuroimaging approach

Fangwen Yu¹, Jialin Li¹, Lei Xu^{1,2}, Xiaoxiao Zheng^{1,3}, Meina Fu¹, Keshuang Li¹, Shuxia Yao¹, Keith M Kendrick¹, Christian Montag^{1,4}, Benjamin Becker^{1*}

¹The Clinical Hospital of Chengdu Brain Science Institute, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu, China

²Institute of Brain and Psychological Sciences, Sichuan Normal University, Chengdu, China

³Brain Cognition and Brain Disease Institute (BCBDI), Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China

⁴Department of Molecular Psychology, Institute of Psychology and Education, Ulm University, Ulm, Germany

Funding sources The study was supported by the National Key Research and Development Program of China (grant number: 2018YFA0701400).

Author's contribution BB and CM were responsible for study concept and design, JL, LX, XZ, MF, KL, SY, KK contributed to the data acquisition, YF performed the statistical analyses. BB and YF interpreted the results and drafted the manuscript. KK, SY and CM provided critical revision of the manuscript for important intellectual content. All authors reviewed the content and approved the final version for publication.

Conflict of interest The authors declare no conflict of interest

Ethics The study was approached by the Ethics committee of the University of Electronic Science and Technology of China, Chengdu, China

Correspondence to Benjamin Becker, University of Electronic Science and Technology of China, Xiyuan Avenue 2006, 611731 Chengdu, China
Mail: ben_becker@gmx.de

Abstract

Background Accumulating evidence suggests brain structural and functional alterations in Internet Use Disorder (IUD). However, conclusions are strongly limited due to the retrospective case-control design of the studies, small samples, and the focus on general rather than symptom-specific approaches.

Methods We here employed a dimensional multi-methodical MRI-neuroimaging design in a final sample of $n = 203$ subjects to examine associations between levels of IUD and its symptom-dimensions (loss of control/time management, craving/social problems) with brain structure, resting state and task-based (pain empathy, affective go/no-go) brain function.

Results Although the present sample covered the entire range of IUD, including normal, problematic as well as pathological levels, general IUD symptom load was not associated with brain structural or functional alterations. However, the symptom-dimensions exhibited opposing associations with the intrinsic and structural organization of the brain, such that loss of control/time management exhibited negative associations with intrinsic striatal networks and hippocampal volume, while craving/social problems exhibited a positive association with intrinsic striatal networks and caudate volume.

Conclusions Our findings provided the first evidence for IUD symptom-domain specific associations with progressive alterations in the intrinsic structural and functional organization of the brain, particularly of striatal systems involved in reward, habitual and cognitive control processes.

Key words Internet use disorder, Internet gaming disorder, neuroimaging, striatum, hippocampus, brain

Introduction

Internet use disorders (IUD) have become a growing public health concern. IUD is characterized by excessive Internet usage, loss of control over usage and detrimental consequences for occupational and social functioning. IUD represents an umbrella term encompassing Internet-associated use disorders, i.e. Internet gaming disorder (IGD) or Social Networks Use Disorder (Montag et al., 2021a; b; Montag, Schivinski & Pontes, 2021; Musetti et al., 2016; Spada, 2014). Estimated prevalence rates of IUD or its subforms such as IGD range from 1% to 10%, with particularly high rates among young individuals and in Asia with up to 25% meeting the criteria for some forms of IUD (Pan, Chiu, & Lin, 2020; for Asia see Montag & Becker; 2020). Debates continue on whether IUD represents a mental disorder (Musetti et al., 2016; Starcevic & Khazaal, 2017), yet (I)GD has been included in the current diagnostic classification systems (DSM-V, ICD-11) and conceptual perspectives propose symptomatic and neurobiological similarities with behavioral and substance use addictions (Montag & Reuter, 2017; Musetti et al., 2016; Young, 1998b, 2011).

Several psychometric instruments have been developed to assess specific forms of IUD, including IGD or the use of specific platforms (Montag et al., 2018; Pontes et al., 2021), while the extensively validated Internet Addiction Diagnostic Questionnaire and the Internet addiction test (IAT) measure level of IUD in general (Young, 1998a, 1999). The symptoms assessed by these scales strongly resemble symptoms of substance addiction, such that the items of the IAT assess preoccupation with the Internet, withdrawal, loss of regulatory control and social problems (Musetti et al., 2016; Young, 1998b). While different factorial structures of the IAT ranging from one to six factors have been proposed based on exploratory factor analyses (Chang & Law, 2008; Khazaal et al., 2008; Korkeila, Kaarlas, Jääskeläinen, Vahlberg, & Taiminen, 2010; Widjianto, Griffiths, & Brunsden, 2011), confirmatory factor analyses consistently revealed a two-factorial model of the IAT across Western and Chinese samples (Pawlikowski, Altstötter-Gleich, & Brand, 2013; Stodt et al., 2018). The factors describe the key facets of 'loss of control/time management', which strongly predicts obsessive-compulsive and impulsivity symptoms and 'craving/social problems', which predicts emotional and interpersonal dysregulations. The two factorial solution has been further confirmed by good convergent, divergent and incremental validity of the two-factor structure of the short version of the IAT (s-IAT) across independent samples (Pawlikowski et al., 2013;

Stodt et al., 2018).

In line with the similarities on the symptomatic level, accumulating evidence suggests that functional and structural brain changes associated with problematic internet use partly resemble alterations observed in substance addiction (Klugah-Brown et al., 2020; Klugah-Brown et al., 2021; Tolomeo & Yu, 2022). Similar to substance use disorders, IGD showed reduced gray matter volumes in the striatal reward system, as well as insular, frontal and temporal systems involved in executive control, decision-making and social processes (He, Turel, & Bechara, 2017; Montag et al., 2018; Zhou et al., 2011; Zhou et al., 2019a). Functional MRI studies reported cue and craving associated hyperreactivity in striatal, cingulate and lateral prefrontal cortex (DLPFC) regions (Han, Hwang, & Renshaw, 2011; Ko et al., 2009; Ko et al., 2013a; Ko et al., 2013b; Lorenz et al., 2013) while cognitive and emotional control impairments were associated with abnormal engagement of fronto-parietal, insular and temporal regions (Chen et al., 2015; Ding et al., 2014; Dong, Devito, Du, & Cui, 2012; Dong, Zhou, & Zhao, 2010; Ko et al., 2014; Liu et al., 2014; Shin, Kim, Kim, & Kim, 2021; Turel, He, Xue, Xiao, & Bechara, 2014; Zhang et al., 2020). Alterations in the intrinsic functional organization of the brain have been extensively examined in individuals with problematic Internet use by means of resting-state fMRI with recent large scale and meta-analysis studies reporting altered functional connectivity in striato-insular-frontal circuits (Dong et al., 2021a; Yan, Li, Yu, & Zhao, 2021).

However, the findings for IUD (particularly for IGD) have been inconsistent, such that prospective longitudinal studies could not confirm emotional or empathic alterations previously observed in cross-section designs (Anderson et al., 2008; Anderson et al., 2010; Gao et al., 2017; Jiao, Wang, Peng, & Cui, 2017; Kühn et al., 2018). These inconsistencies may partly be related to the retrospective categorical case-control design of the studies that commonly compared relatively small sample sizes of individuals with excessive internet engagement or established IUD with healthy controls. The assumption of the case-control approach is that categorical diagnosis, e.g. fulfilling at least five of the DSM-5 IGD criteria or an IAT score >50 will define mechanistically meaningful study samples, whereas in fact the categorical IUD groups neither exhibit the same symptom constellation nor cover the whole symptom spectrum of IUD (Etkin, 2019). Several of the early studies employ sample sizes between 20 to 30, or often less than 20 per group (Cheng & Liu, 2020; Dong et al., 2010; Lorenz et

al., 2013), which permits statistical detection power to be achieved but at the cost of sacrificing clinical significance.

Recent suggestions to improve psychiatric neuroimaging advocate prospective dimensional approaches that examine associations between varying degrees in symptom load and progressive neural alterations in larger samples (Etkin, 2019). An increasing number of studies successfully employed this approach to determine symptom-dimension specific brain alterations e.g. with respect to depression or alexithymia (Hägele et al., 2016; Li et al., 2019; Luo et al., 2018) and initial studies have demonstrated the feasibility of this approach to determine IUD-associated brain structural and intrinsic functional changes in larger samples spanning the entire symptom load range (Dong et al., 2021a; Montag et al., 2018; Zhou et al., 2021). Against this background we employed a dimensional multi-modal neuroimaging approach in a comparably large sample (> 240 subjects) to explore associations between the level of problematic Internet use (as assessed by the IAT) on brain structure as well as intrinsic and task-associated brain function. We focused on task domains for which IUD alterations are a matter of debate, in particular cognitive control and implicit emotion regulation (affective go/no-go) and empathy or social communicative processing (pain empathy task including physical and social communicative pain). To further disentangle the impact of different symptom constellations which cannot be examined with the categorical approach the symptom dimensions “loss of control/time” and “craving/social problem” (Pawlikowski et al., 2013) were separately examined (Stodt et al., 2018). In line with previous studies, we expected that higher levels of problematic Internet use would be associated with volume and intrinsic striatal connectivity changes as well as deficient frontal and insular engagement during cognitive control and empathic processing, respectively. With respect to the sub-facets we expected distinguishable symptom-specific associations.

Methods

Participants

To implement dimensional neuroimaging in a large sample we capitalized on the Chengdu Gene Brain Behavior Project which aims at determining psychopathological, genetic and neural alterations in a large cohort of healthy individuals (Liu et al., 2021; Montag et al., 2017b). N = 250 subjects from this project underwent multimodal MRI imaging, including a pain empathy paradigm (Li et al., 2019; Xu et al., 2020; Zhou et al., 2020a), an affective Go/No-go paradigm (Zhuang et al., 2021) as well as resting state and brain structural assessments (e.g. Liu et al., 2021). Levels of IUD were assessed at the time of the MRI-data acquisition by means of the Internet Addiction Test (IAT, Pawlikowski et al., 2013; Stodt et al., 2018; Young, 1998b) which assess a total score as well as the sub-facets describing loss of control/time management and craving/social problems in the context of problematic Internet use.

Following data quality assessments N=203 subjects were included in the final multimodal data analyses (104 males; mean age \pm SD=21.65 \pm 2.37 years). Participants provided written informed consent, the study had full ethical approval and was in accordance with the latest revision of the declaration of Helsinki.

Experimental Protocols

Internet addiction and the sub-facets were assessed using the short version of Young's Internet Addiction Test (s-IAT, Pawlikowski et al., 2013). The s-IAT includes 12-items which assess the frequency of negative experiences and consequences due to excessive online activities in everyday life on a 5-point Likert scale ranging from 1 (never) to 5 (very often) (Young, 1998). The total score can correspondingly range from 12 to 60, with a total score >30 indicating problematic and >37 indicating pathological Internet use (Pawlikowski et al., 2013; Stodt et al., 2018). The s-IAT consists of the two subscales assessing loss of control/time management and craving/social problems each consisting of six items (items see table S1). Subjects underwent MRI assessments, including acquisition of T1-weighted brain structural data, resting state fMRI data as well as a validated fMRI pain empathy task and an affective Go/No-go task (details see supplemental material and figure 1).

MRI Data Processing, Dimensional Approach and Thresholding

MRI data were acquired and processed using extensively validated standard procedures and processing pipelines (see supplements). We employed a dimensional neuroimaging approach modelling associations between the severity of Internet addiction (aka Internet Use Disorders) and its sub-facets (as assessed by the s-IAT) and individual variations in brain function and structure using voxel-level multiple regression analyses in SPM12. To this end the s-IAT total and subscale scores served as predictor while the voxel-wise brain structural, task activation and intrinsic connectivity maps served as dependent variable. Given that the s-IAT total scores consisted of the sum of loss of control/time management subscale scores and craving/social problems subscales separate multiple regression analyses for the total scores and two subscales, both including age and sex as covariates of no interest (in the VBM analysis also TIV) were conducted. Analyses were conducted on the whole-brain level with FDR $p < 0.05$ for multiple comparison correction. Details on the paradigms, preprocessing and first level models are provided in the supplements. The voxel-wise whole-brain maps were modelled as follows.

Task fMRI pain empathy

The paradigm included visual stimuli depicting ‘physical pain’, ‘affective pain’ and corresponding non-painful control stimuli (‘physical control’ and ‘affective control’). The first level matrix included separate regressors for the four experimental conditions and six head-motion parameters. In the absence of empathy type-specific hypotheses we focused on the main pain empathy contrast $[(\text{physical pain} + \text{affective pain}) > (\text{physical control} + \text{affective control})]$, with further exploratory analyses examining associations with empathy-type specific activity ($[\text{physical pain} > \text{physical control}]$; $[\text{affective pain} > \text{affective control}]$).

Task fMRI affective Go/No-go

The lexical Go/No-go paradigm required go and no-go responses in neutral, negative and positive contexts leading to six experimental conditions on the first level (neu Go, neu No Go, neg Go, neg No Go, pos Go, pos No Go) and additional six head-motion parameters. We focused on the general inhibitory control contrast $[(\text{neu No Go} + \text{neg No Go} + \text{pos No Go}) > (\text{neu Go} + \text{neg Go} + \text{pos Go})]$, with further exploratory analyses exploring associations with emotional context-specific inhibitory activity ($[\text{neu No Go} >$

neu Go], [neg No Go > neg Go], ([pos No Go > pos Go]). During the paradigm accuracy and reaction times were collected and served as behavioral outcomes.

Resting state functional connectivity

Based on the different roles of the ventral and dorsal striatum in (internet) addiction, (Dong et al., 2021a; Dong et al., 2021b; Zhou et al., 2018; Zhou et al., 2020b; Zhou et al., 2019b) bilateral ventral striatum and dorsal striatum masks served as seed regions (regions of interest, ROIs). ROIs were derived from the brainnetome atlas (Fan et al., 2016) (table S4) functional connectivity between the ROIs and voxels on the whole brain level were computed using DPABI (Yan, Wang, Zuo, & Zang, 2016; <https://www.rfmri.org/dpabi>).

Brain structure gray matter volume

Voxel-based morphometry (VBM) analysis was conducted using the CAT toolbox (version 12.7, <http://www.neuro.uni-jena.de/cat/>) including quality assessments, smoothing and statistical analysis. T1 images were normalized to a template space and segmented into gray matter (GM), white matter (WM), and cerebrospinal fluid (CSF). The preprocessing parameters used the default recommendations (“Segment Data”). After the preprocessing, we conducted a quality check and deleted data with mean correlation of sample homogeneity below two standard deviations. After segmentation, images were smoothed with 8mm FWHM and resulting voxel-wise gray matter masks were subjected to the multiple regression models.

Results

Participants and levels of IUD

Mean total score of the s-IAT was 32.49 (SD=7.62) in the present sample (N=203). 74 subjects scored below 30 indicating normal Internet use, n=81 subjects reported scores between 30 and 37 suggesting problematic Internet use, while 48 subjects reported total scores >37 suggesting pathological Internet use (Pawlikowski et al., 2013; Stodt et al., 2018). With respect to sex differences, the total levels, $t=1.06$, $df=201$, $p=0.289$, Cohen's $d=0.149$, and levels of craving/social problem were comparable $t=0.757$, $df=201$, $p=0.450$, Cohen's $d=0.106$ while for the loss of control/time management females reported higher levels than males ($t=2.682$, $df=201$, $p=0.008$, Cohen's $d=0.377$).

Sex was consequently included in all analyses as covariate. Details see table 1. The data displayed no strong violations of normal distribution regarding the s-IAT and its subscales. Moreover, the scales showed good psychometric properties (details see table S2, supplements, supplementary figure S1)

Associations between levels of IUD and affective Go/No-Go performance

Partial correlation analyses with accuracy and reaction time of go and no-go trials for the three conditions (negative, positive, neutral) and age and gender as covariates revealed that the s-IAT total scores and two subscale scores had significant negative correlations with the accuracy in no-go trials (Negative no-go: $r_{\text{total}}=-0.215, p=0.00215$, $r_{\text{control}}=-0.186, p=0.00823$, $r_{\text{craving}}=-0.211, p=0.00266$; Positive no-go: $r_{\text{total}}=-0.233, p=0.00087$, $r_{\text{control}}=-0.189, p=0.00717$, $r_{\text{craving}}=-0.241, p=0.00058$; Neutral no-go: $r_{\text{total}}=-0.203, p=0.00391$, $r_{\text{control}}=-0.162, p=0.02137$, $r_{\text{craving}}=-0.211, p=0.00258$, Bonferroni corrected $p=0.0083$, six tests). No significant correlation with accuracy and reaction times of go trials were observed after Bonferroni correction. Details see table S3.

Associations between levels of IUD and pain empathy and cognitive/emotional control associated brain activity

The pain empathy paradigm engaged per se the typical bilateral pain empathy networks encompassing e.g. the bilateral inferior frontal gyrus (IFG) and insula and medial frontal cortex (FDR $p<0.05$, cluster size>297, see table S5 and figure S2A). The go/no-go paradigm engaged the key frontoparietal inhibitory control networks (contrast [all no/go > all go], FDR $p<0.05$) (figure S2B). For details on other contrasts please see also Zhou et al., 2020(a); Zhuang et al., 2021. Multiple regression analysis examining associations with IUD total and subscale scores did not reveal significant associations with pain empathy and general inhibitory control associated brain activity (FDR $p<0.05$). Further exploratory analyses did not reveal significant associations with empathy-type or emotional specific inhibitory activity.

Associations between levels of IUD and intrinsic striatal networks

While no associations were found for the total IAT score, the subfacets showed associations with striatal intrinsic connectivity networks. Levels of loss of control/time management were significantly negatively associated with the intrinsic functional

connectivity of ventral striatum with the right calcarine, right middle occipital lobe, left superior occipital lobe and right fusiform gyrus (see table 2 and figure 2A, FDR $p<0.05$, cluster size ≥ 49 voxels), while they were significantly negatively associated with dorsal striatum intrinsic connectivity with the right precuneus, right posterior cerebellum lobe and anterior cingulate cortex (see table 2 and figure 2A, FDR $p<0.05$, cluster size ≥ 36 voxels). Levels of craving/social problems were significantly positively associated with functional connectivity of the ventral striatum with the right fusiform gyrus, left superior occipital lobe, right middle occipital lobe, left inferior orbital frontal cortex, right superior occipital lobe, right superior temporal lobe and left insula (see table 3 and figure 2B, FDR $p<0.05$, cluster size ≥ 22 voxels). With respect to the dorsal striatum connectivity levels of craving/social problems were positively associated with coupling with the left occipital lobe, left inferior frontal gyrus and right inferior frontal gyrus (see table 3 and figure 2B, FDR $p<0.05$, cluster size ≥ 27 voxels).

Associations between levels of IUD and brain structure

No associations with s-IAT total score were observed but loss of control/time management exhibited a significant positive association with the GMV of the left hippocampus (MNI: -15/-18/-15, FDR $p<0.05$, cluster size=530 voxels, see figure 3A) while levels of craving/social problems showed a significant positive association with the GMV of the striatum (caudate subregion; MNI: 5/8/5, FDR $p<0.05$, cluster size=985 voxels, see figure 3B.). Mapping the caudate cluster using the brainnetome atlas revealed that the cluster encompassed ventral and dorsal striatal subregions (see table S6 and figure 3B).

Discussion

We employed a dimensional neuroimaging approach in a comparably large sample and across different imaging modalities to determine early brain markers for IUD. Although a sample of young students was enrolled our sample spanned the entire range of IUD, with 81 or 43 from 203 individuals reporting problematic ($n = 81$) or pathological levels of internet use ($n = 43$), respectively (according to cut off scores provided in Pawlikowski et al., 2013; Stodt et al., 2018). Levels of IUD had negative associations with response inhibition performance in an affective go/no-go task in the absence of significant associations with brain activity during cognitive/emotional control or pain empathic processing. Although general severity of IUD (s-IAT total scores) did not associate with brain structural or functional organization, the sub-facets exhibited significant and opposing associations, such that loss of control/time management exhibited negative associations with intrinsic striatal networks and hippocampal volume, while craving/social problems demonstrated positive associations with striatal networks and caudate volume.

The dimensional approach allowed us for the first time to disentangle symptom-domain specific alterations, thus providing a neurobiological basis of previously reported differential associations with pathological domains (Etkin 2019; Pawlikowski et al., 2013) and suggesting that the behavioral dysregulations are mediated by separable neurobiological alterations. Higher levels of loss of control/time management symptom load were associated with decreasing connectivity of the ventral striatum with the occipital and fusiform regions and of the dorsal striatum with the precuneus and the anterior cingulate cortex. In contrast, higher levels of craving/social problems were associated with enhanced connectivity between the ventral striatum and the occipital, orbitofrontal and insular regions. Negative associations between loss of control and striatum connectivity might reflect dysregulations in inhibitory control and visual salience processing including the superior and middle occipital regions, and the action control system, including the cerebellum and ACC (Dong et al., 2021b; Meng, Deng, Wang, Guo, & Li, 2015). Positive associations between craving/social problems and the ventral striatal system encompassed occipital regions involved in visual salience processing and orbitofrontal regions involved in reward processing which have been associated with cue-reactivity and craving (Hanlon, Dowdle, Naselaris, Canterberry, & Cortese, 2014; Zhou et al., 2019b). The identified circuits moreover overlap with

circuits reported to be involved in cognitive control deficits, reward dysregulations and cue-reactivity in both IUD (Ma et al., 2019; Starcke, Antons, Trotzke, & Brand, 2018; Yu et al., 2021; Yuan et al., 2017) and substance-use disorders (Montag & Reuter, 2017; Verdejo-Garcia, Garcia-Fernandez, & Dom, 2019; Volkow & Boyle, 2018). In substance addiction progressive adaptations in these circuits have been explained in terms of ‘incentive sensitization’ reflecting that the incentive motivational effects of drug and drug-associated stimuli leads to progressive dysfunctions in salience and reward processing and impaired executive control (Robinson & Berridge, 1993, 2008) or in terms of dysregulated ‘habit formation’, a process during which initially reinforced goal-directed actions become progressively habitual and compulsive (Ersche et al., 2016; Robbins & Clark, 2015). The excessive engagement in Internet usage may additionally be promoted by negative reinforcement learning such that these activities can reduce stress induced by societal and social pressure and attenuate anxiety and depression (King & Delfabbro, 2014, 2018).

The different neurobiological underpinnings of the two symptom scales were additionally mirrored on the brain structural level such that craving/social problems positively associated with the GMV of the striatum, including both ventral and dorsal parts, while higher levels of loss of control/time management problems were associated with larger GMV of the hippocampus. Previous studies reported increased GMV of the dorsal and ventral striatum in IUD and the volumes correlated with cognitive control performance and IAT scores (Cai et al., 2016). Although previous dimensional studies reported negative associations between problematic internet usage and striatal volume (Montag et al., 2017a; Montag et al., 2018; Zhou et al., 2020), morphological alterations in the mesocorticolimbic reward systems – also positive associations - and their structural connectivity have been extensively documented in IUD, in particular IGD (He, Turel & Bechara 2017; Hong et al., 2015; Wang et al., 2019; Zhou et al., 2019a; Zhou et al., 2011). In the context of the previous literature the current findings indicate that alterations in these pathways may specifically promote craving/social problems in IUD. In contrast, higher levels of control/time management were associated with larger hippocampal volumes. Previous studies have reported decreased as well as increased hippocampal volumes in IGD (Lin, Dong, Wang, & Du, 2015; Yoon et al., 2017) and this region has been suggested to play a role in the interaction between reward processing, behavioral reinforcement and memory retrieval (Everitt & Robbins, 2005).

This role in the development of IUD has been confirmed by a prospective longitudinal study employing gaming training (“Super Mario 64 DS”) at least 30 minutes per day over two months (Gleich, Lorenz, Gallinat, & Kuhn, 2017). Before and after training phase, both groups were shown gaming videos, which recorded during training, with three consequences (reward, punishment and neural) in the fMRI scanner. The results showed that gaming increased brain activity in the left hippocampus after training that reward processing during gaming may lead to a stronger memory formation and in turn promote the development of pathological craving and problems in control/time management (Cox & Witten, 2019; Dolan & Dayan, 2013).

In contrast to our expectations, we did not find associations between levels of IUD and neural activity during affective go/no-go or pain empathy. Previous studies reported inhibitory and cognitive control deficits in the context of deficient frontal engagement in IUD, in particular IGD (e.g. Kuss, Pontes, & Griffiths, 2018). These findings were primarily based on samples with manifest IUD, while our findings were obtained in subjects with comparably lower symptom load suggesting that frontal deficits during inhibitory control may either represent a predisposition for rather than a consequence of excessive Internet use or alternatively may only manifest at later stages of the disorder. In line with accumulating evidence from prospective longitudinal intervention studies reporting no effects of gaming on emotional or pain empathic processes (Gao et al., 2017; Kühn et al., 2018; Yu et al., 2021), we did not observe alterations in the pain empathic and social processing domain. This may reflect that specifically the processing of Internet- or game-related stimuli is biased in IUD. In line with this explanation previous studies observed altered processing of internet related stimuli but not of general emotional or monetary reward related processing (Starcke et al., 2018; Yao et al., 2020; Yu et al., 2021).

The present findings underscore the importance of considering symptom-domains separately rather than the total symptom load of IUD related scales. While the two-factorial structure of the IAT (Pawlakowski et al., 2013) and separate diagnostic symptom domains have been emphasized, the vast majority of IUD or IGD studies determined samples based on the total symptom score. The total score may have a high clinical utility to identify individuals in need for intervention (Pontes & Griffiths, 2015; Pontes, Kiraly, Demetrovics, & Griffiths, 2014), however, the present results underscore that the identification of the precise neurocognitive and brain-based

mechanisms will require a symptom-domain specific approach because distinct alterations may underlie specific behavioral dysregulations in IUD.

Findings of the present study need to be considered within the context of limitations. Firstly, IUD was associated with inhibitory control performance, but a comprehensive neurocognitive characterization in association with the IAT was not concluded. Secondly, we cannot exclude that the s-IAT sub-facets would exhibit associations with neural activity during other cognitive and emotional processes. The symptom-subdomain specific alterations need to be established across different IUD subforms such as IUD.

Conclusions

Our findings provided the first evidence for symptom-domain specific associations with progressive alterations in the intrinsic structural and functional organization of the brain, in particular the striatal systems involved in reward, habitual and cognitive control processes. Our findings suggest that different symptom domains in IUD are neurally underpinned by separable alterations.

Acknowledgements and funding

This work was supported by the National Key Research and Development Program of China (grant number: 2018YFA0701400). The authors declare no competing interests.

References

Anderson, C. A., Sakamoto, A., Gentile, D. A., Ihori, N., Shibuya, A., Yukawa, S., . . . Kobayashi, K. (2008). Longitudinal effects of violent video games on aggression in Japan and the United States. *Pediatrics*, 122(5), e1067-1072. doi:10.1542/peds.2008-1425.

Anderson, C. A., Shibuya, A., Ihori, N., Swing, E. L., Bushman, B. J., Sakamoto, A., . . . Saleem, M. (2010). Violent video game effects on aggression, empathy, and prosocial behavior in eastern and western countries: a meta-analytic review. *Psychol Bull*, 136(2), 151-173. doi:10.1037/a0018251.

Cai, C., Yuan, K., Yin, J., Feng, D., Bi, Y., Li, Y., . . . Tian, J. (2016). Striatum morphometry is associated with cognitive control deficits and symptom severity in internet gaming disorder. *Brain imaging and behavior*, 10(1), 12-20. doi:10.1007/s11682-015-9358-8.

Chang, M. K., & Law, S. P. M. (2008). Factor structure for Young's Internet Addiction Test: A confirmatory study. *Computers in human behavior*, 24(6), 2597-2619. doi:10.1016/j.chb.2008.03.001.

Chen, C. Y., Huang, M. F., Yen, J. Y., Chen, C. S., Liu, G. C., Yen, C. F., & Ko, C. H. (2015). Brain correlates of response inhibition in Internet gaming disorder. *Psychiatry and clinical neurosciences*, 69(4), 201-209. doi:10.1111/pcn.12224.

Cheng, H., & Liu, J. (2020). Alterations in Amygdala Connectivity in Internet Addiction Disorder. *Sci Rep*, 10(1), 2370. doi:10.1038/s41598-020-59195-w.

Cox, J., & Witten, I. B. (2019). Striatal circuits for reward learning and decision-making. *Nature Reviews Neuroscience*, 20(8), 482-494. doi:10.1038/s41583-019-0189-2.

Ding, W. N., Sun, J. H., Sun, Y. W., Chen, X., Zhou, Y., Zhuang, Z. G., . . . Du, Y. S. (2014). Trait impulsivity and impaired prefrontal impulse inhibition function in adolescents with internet gaming addiction revealed by a Go/No-Go fMRI study. *Behav Brain Funct*, 10(1), 20. doi:10.1186/1744-9081-10-20.

Dolan, R. J., & Dayan, P. (2013). Goals and habits in the brain. *Neuron*, 80(2), 312-325. doi:10.1016/j.neuron.2013.09.007

Dong, G., Devito, E. E., Du, X., & Cui, Z. (2012). Impaired inhibitory control in 'internet addiction disorder': a functional magnetic resonance imaging study. *Psychiatry Res*, 203(2-3), 153-158. doi:10.1016/j.psychresns.2012.02.001.

Dong, G., Zhou, H., & Zhao, X. (2010). Impulse inhibition in people with Internet addiction disorder: electrophysiological evidence from a Go/NoGo study. *Neurosci Lett*, 485(2), 138-142. doi:10.1016/j.neulet.2010.09.002.

Dong, G. H., Dong, H., Wang, M., Zhang, J., Zhou, W., Du, X., & Potenza, M. N. (2021a). Dorsal and ventral striatal functional connectivity shifts play a potential role in internet gaming disorder. *Commun Biol*, 4(1), 866. doi:10.1038/s42003-021-02395-5.

Dong, H., Wang, M., Zhang, J., Hu, Y., Potenza, M. N., & Dong, G. H. (2021b). Reduced frontostriatal functional connectivity and associations with severity of Internet gaming disorder. *Addiction Biology*, 26(4), e12985. doi: 10.1111/adb.12985.

Ersche, K. D., Gillan, C. M., Jones, P. S., Williams, G. B., Ward, L. H., Luijten, M., . . . Robbins, T. W. (2016). Carrots and sticks fail to change behavior in cocaine addiction. *Science*, 352(6292), 1468-1471. doi:10.1126/science.aaf3700.

Etkin, A. (2019). A Reckoning and Research Agenda for Neuroimaging in Psychiatry. *Am J Psychiatry*, 176(7), 507-511. doi:10.1176/appi.ajp.2019.19050521.

Everitt, B. J., & Robbins, T. W. (2005). Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nature neuroscience*, 8(11), 1481-1489. doi:10.1038/nn1579.

Fan, L., Li, H., Zhuo, J., Zhang, Y., Wang, J., Chen, L., . . . Laird, A. R. (2016). The human brainnetome atlas: a new brain atlas based on connectional architecture. *Cerebral cortex*, 26(8), 3508-3526. doi:10.1093/cercor/bhw157.

Gao, X., Pan, W., Li, C., Weng, L., Yao, M., & Chen, A. (2017). Long-time exposure to violent video games does not show desensitization on empathy for pain: an fMRI study. *Frontiers in psychology*, 8, 650. doi:10.3389/fpsyg.2017.00650.

Gleich, T., Lorenz, R. C., Gallinat, J., & Kuhn, S. (2017). Functional changes in the reward circuit in response to gaming-related cues after training with a commercial video game. *Neuroimage*, 152, 467-475. doi:10.1016/j.neuroimage.2017.03.032.

Hägele, C., Friedel, E., Schlagenhauf, F., Sterzer, P., Beck, A., Bermpohl, F., . . . Ströhle, A. (2016). Affective responses across psychiatric disorders—A dimensional approach. *Neuroscience letters*, 623, 71-78. doi:10.1016/j.neulet.2016.04.037.

Han, D. H., Hwang, J. W., & Renshaw, P. F. (2011). Bupropion sustained release treatment decreases craving for video games and cue-induced brain activity in patients with Internet video game addiction. doi:10.1037/2160-4134.1.S.108.

Hanlon, C. A., Dowdle, L. T., Naselaris, T., Canterberry, M., & Cortese, B. M. (2014). Visual cortex activation to drug cues: a meta-analysis of functional neuroimaging papers in addiction and substance abuse literature. *Drug and alcohol dependence*, 143, 206-212. doi:10.1016/j.drugalcdep.2014.07.028.

He, Q., Turel, O., & Bechara, A. (2017). Brain anatomy alterations associated with Social Networking Site (SNS) addiction. *Sci Rep*, 7(1), 45064. doi:10.1038/srep45064.

Hong, S. B., Harrison, B. J., Dandash, O., Choi, E. J., Kim, S. C., Kim, H. H., . . . Yi, S. H. (2015). A selective involvement of putamen functional connectivity in youth with internet gaming disorder. *Brain Res*, 1602, 85-95. doi:10.1016/j.brainres.2014.12.042.

Jiao, C., Wang, T., Peng, X., & Cui, F. (2017). Impaired empathy processing in individuals with internet addiction disorder: An event-related potential study. *Frontiers in Human Neuroscience*, 11, 498. doi:10.3389/fnhum.2017.00498.

Khazaal, Y., Billieux, J., Thorens, G., Khan, R., Louati, Y., Scarlatti, E., . . . Zullino, D. (2008). French validation of the internet addiction test. *Cyberpsychology & behavior*, 11(6), 703-706. doi:10.1089/cpb.2007.0249.

King, D. L., & Delfabbro, P. H. (2014). The cognitive psychology of Internet gaming disorder. *Clinical psychology review*, 34(4), 298-308. doi:10.1016/j.cpr.2014.03.006.

King, D. L., & Delfabbro, P. H. (2018). The concept of “harm” in Internet gaming disorder. *Journal of behavioral addictions*, 7(3), 562-564. doi:10.1556/2006.7.2018.24.

Klugah-Brown, B., Di, X., Zweerings, J., Mathiak, K., Becker, B., & Biswal, B. (2020). Common and separable neural alterations in substance use disorders: A coordinate-based meta-analyses of functional neuroimaging studies in humans. *Human brain mapping*, 41(16), 4459-4477. doi:10.1002/hbm.25085.

KlugahBrown, B., Jiang, C., Agoalikum, E., Zhou, X., Zou, L., Yu, Q., . . . Biswal, B. (2021). Common abnormality of gray matter integrity in substance use disorder and obsessive-compulsive disorder: A comparative voxel-based meta-analysis.

Human brain mapping, 42(12), 3871-3886. doi:10.1002/hbm.25471.

Ko, C. H., Hsieh, T. J., Chen, C. Y., Yen, C. F., Chen, C. S., Yen, J. Y., . . . Liu, G. C. (2014). Altered brain activation during response inhibition and error processing in subjects with Internet gaming disorder: a functional magnetic imaging study. *Eur Arch Psychiatry Clin Neurosci*, 264(8), 661-672. doi:10.1007/s00406-013-0483-3.

Ko, C. H., Liu, G. C., Yen, J. Y., Chen, C. Y., Yen, C. F., & Chen, C. S. (2013a). Brain correlates of craving for online gaming under cue exposure in subjects with Internet gaming addiction and in remitted subjects. *Addict Biol*, 18(3), 559-569. doi:10.1111/j.1369-1600.2011.00405.x.

Ko, C. H., Liu, G. C., Yen, J. Y., Yen, C. F., Chen, C. S., & Lin, W. C. (2013b). The brain activations for both cue-induced gaming urge and smoking craving among subjects comorbid with Internet gaming addiction and nicotine dependence. *Journal of psychiatric research*, 47(4), 486-493. doi:10.1016/j.jpsychires.2012.11.008.

Ko, C. H., Yen, J. Y., Yen, C. F., Chen, C. S., & Chen, C. C. (2012). The association between Internet addiction and psychiatric disorder: a review of the literature. *Eur Psychiatry*, 27(1), 1-8. doi:10.1016/j.eurpsy.2010.04.011.

Korkeila, J., Kaarlas, S., Jääskeläinen, M., Vahlberg, T., & Taiminen, T. (2010). Attached to the web—harmful use of the Internet and its correlates. *European Psychiatry*, 25(4), 236-241. doi:10.1016/j.eurpsy.2009.02.008.

Kühn, S., Kugler, D., Schmalen, K., Weichenberger, M., Witt, C., & Gallinat, J. (2018). The myth of blunted gamers: No evidence for desensitization in empathy for pain after a violent video game intervention in a longitudinal fMRI study on non-gamers. *Neurosignals*, 26(1), 22-30. doi:10.1159/000487217.

Kuss, D. J., Pontes, H. M., & Griffiths, M. D. (2018). Neurobiological Correlates in Internet Gaming Disorder: A Systematic Literature Review. *Front Psychiatry*, 9, 166. doi:10.3389/fpsyg.2018.00166.

Li, J., Xu, L., Zheng, X., Fu, M., Zhou, F., Xu, X., . . . Becker, B. (2019). Common and Dissociable Contributions of Alexithymia and Autism to Domain-Specific Interoceptive Dysregulations: A Dimensional Neuroimaging Approach. *Psychother Psychosom*, 88(3), 187-189. doi:10.1159/000495122.

Lin, X., Dong, G., Wang, Q., & Du, X. (2015). Abnormal gray matter and white

matter volume in 'Internet gaming addicts'. *Addictive Behaviors*, 40, 137-143. doi:10.1016/j.addbeh.2014.09.010.

Liu, C., Xu, L., Li, J., Zhou, F., Yang, X., Zheng, X., . . . Becker, B. (2021). Serotonin and early life stress interact to shape brain architecture and anxious avoidant behavior - a TPH2 imaging genetics approach. *Psychol Med*, 51(14), 2476-2484. doi:10.1017/S0033291720002809.

Liu, G. C., Yen, J. Y., Chen, C. Y., Yen, C. F., Chen, C. S., Lin, W. C., & Ko, C. H. (2014). Brain activation for response inhibition under gaming cue distraction in internet gaming disorder. *Kaohsiung J Med Sci*, 30(1), 43-51. doi:10.1016/j.kjms.2013.08.005.

Lorenz, R. C., Kruger, J. K., Neumann, B., Schott, B. H., Kaufmann, C., Heinz, A., & Wustenberg, T. (2013). Cue reactivity and its inhibition in pathological computer game players. *Addict Biol*, 18(1), 134-146. doi:10.1111/j.1369-1600.2012.00491.x.

Luo, L., Becker, B., Zheng, X., Zhao, Z., Xu, X., Zhou, F., . . . Kendrick, K. M. (2018). A dimensional approach to determine common and specific neurofunctional markers for depression and social anxiety during emotional face processing. *Hum Brain Mapp*, 39(2), 758-771. doi:10.1002/hbm.23880.

Ma, S.-S., Worhunsky, P. D., Xu, J.-s., Yip, S. W., Zhou, N., Zhang, J.-T., . . . Yao, Y.-W. (2019). Alterations in functional networks during cue-reactivity in Internet gaming disorder. *Journal of behavioral addictions*, 8(2), 277-287. doi:10.1556/2006.8.2019.25.

Meng, Y., Deng, W., Wang, H., Guo, W., & Li, T. (2015). The prefrontal dysfunction in individuals with Internet gaming disorder: a meta - analysis of functional magnetic resonance imaging studies. *Addiction Biology*, 20(4), 799-808. doi:10.1111/adb.12154.

Montag, C., Markowetz, A., Blaszkiewicz, K., Andone, I., Lachmann, B., Sariyska, R., . . . Reuter, M. (2017a). Facebook usage on smartphones and gray matter volume of the nucleus accumbens. *Behavioural brain research*, 329, 221-228.

Montag, C., & Reuter, M. (2017). *Internet addiction*: Springer. doi:10.1016/j.bbr.2017.04.035.

Montag, C., Sindermann, C., Melchers, M., Jung, S., Luo, R. X., Becker, B., . . . Kendrick, K. M. (2017b). A functional polymorphism of the OXTR gene is

associated with autistic traits in Caucasian and Asian populations. *American Journal of Medical Genetics Part B-Neuropsychiatric Genetics*, 174(8), 808-816. doi:10.1002/ajmg.b.32596.

Montag, C., Zhao, Z., Sindermann, C., Xu, L., Fu, M., Li, J., . . . Becker, B. (2018). Internet Communication Disorder and the structure of the human brain: initial insights on WeChat addiction. *Sci Rep*, 8(1), 2155. doi:10.1038/s41598-018-19904-y.

Montag, C., & Becker, B. (2020). Internet and smartphone use disorder in Asia [Editorial]. *Addictive Behaviors*, 107, Article 106380. doi:10.1016/j.addbeh.2020.106380.

Montag, C., Schivinski, B., & Pontes, H. M. (2021a). Is the proposed distinction of gaming disorder into a predominantly online vs. offline form meaningful? Empirical evidence from a large German speaking gamer sample. *Addictive behaviors reports*, 14, 100391. doi:10.1016/j.abrep.2021.100391.

Montag, C., Sindermann, C., Rozgonjuk, D., Yang, S., Elhai, J. D., & Yang, H. (2021). Investigating links between fear of COVID-19, neuroticism, social networks use disorder, and smartphone use disorder tendencies. *Frontiers in psychology*, 12. doi:10.3389/fpsyg.2021.682837.

Montag, C., Wegmann, E., Sariyska, R., Demetrovics, Z., & Brand, M. (2021b). How to overcome taxonomical problems in the study of Internet use disorders and what to do with “smartphone addiction”? *Journal of behavioral addictions*, 9(4), 908-914. doi:10.1556/2006.8.2019.59

Musetti, A., Cattivelli, R., Giacobbi, M., Zuglian, P., Ceccarini, M., Capelli, F., . . . Castelnuovo, G. (2016). Challenges in Internet Addiction Disorder: Is a Diagnosis Feasible or Not? *Front Psychol*, 7, 842. doi:10.3389/fpsyg.2016.00842.

Pan, Y.-C., Chiu, Y.-C., & Lin, Y.-H. (2020). Systematic review and meta-analysis of epidemiology of internet addiction. *Neuroscience & Biobehavioral Reviews*, 118, 612-622. doi:10.1016/j.neubiorev.2020.08.013.

Pawlikowski, M., Altstötter-Gleich, C., & Brand, M. (2013). Validation and psychometric properties of a short version of Young's Internet Addiction Test. *Computers in human behavior*, 29(3), 1212-1223. doi:10.1016/j.chb.2012.10.014.

Pontes, H. M., & Griffiths, M. D. (2015). Measuring DSM-5 Internet gaming disorder: Development and validation of a short psychometric scale. *Computers in human behavior*, 45, 137-143. doi:10.1016/j.chb.2014.12.006.

Pontes, H. M., Kiraly, O., Demetrovics, Z., & Griffiths, M. D. (2014). The conceptualisation and measurement of DSM-5 Internet Gaming Disorder: The development of the IGD-20 Test. *PLoS one*, 9(10), e110137. doi:10.1371/journal.pone.0110137.

Pontes, H. M., Schivinski, B., Sindermann, C., Li, M., Becker, B., Zhou, M., & Montag, C. (2021). Measurement and conceptualization of Gaming Disorder according to the World Health Organization framework: The development of the Gaming Disorder Test. *International Journal of Mental Health and Addiction*, 19(2), 508-528. doi:10.1007/s11469-019-00088-z.

Robbins, T. W., & Clark, L. (2015). Behavioral addictions. *Current opinion in neurobiology*, 30, 66-72. doi:10.1016/j.conb.2014.09.005.

Robinson, T. E., & Berridge, K. C. (1993). The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain Res Brain Res Rev*, 18(3), 247-291. doi:10.1016/0165-0173(93)90013-p.

Robinson, T. E., & Berridge, K. C. (2008). The incentive sensitization theory of addiction: some current issues. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 363(1507), 3137-3146. doi:10.1098/rstb.2008.0093.

Shin, Y. B., Kim, H., Kim, S. J., & Kim, J. J. (2021). A neural mechanism of the relationship between impulsivity and emotion dysregulation in patients with Internet gaming disorder. *Addiction Biology*, 26(3), e12916. doi:10.1111/adb.12916.

Spada, M. M. (2014). An overview of problematic internet use. *Addict Behav*, 39(1), 3-6. doi:10.1016/j.addbeh.2013.09.007.

Starcevic, V., & Khazaal, Y. (2017). Relationships between behavioural addictions and psychiatric disorders: What is known and what is yet to be learned? *Frontiers in psychiatry*, 8, 53. doi:10.3389/fpsyg.2017.00053.

Starcke, K., Antons, S., Trotzke, P., & Brand, M. (2018). Cue-reactivity in behavioral addictions: A meta-analysis and methodological considerations. *J Behav Addict*, 7(2), 227-238. doi:10.1556/2006.7.2018.39.

Stodt, B., Brand, M., Sindermann, C., Wegmann, E., Li, M., Zhou, M., . . . Montag, C.

(2018). Investigating the Effect of Personality, Internet Literacy, and Use Expectancies in Internet-Use Disorder: A Comparative Study between China and Germany. *International journal of environmental research and public health*, 15(4), 579. doi:10.3390/ijerph15040579.

Tolomeo, S., & Yu, R. (2022). Brain network dysfunctions in addiction: a meta-analysis of resting-state functional connectivity. *Translational Psychiatry*, 12(1), 1-11. doi:10.1038/s41398-022-01792-6.

Turel, O., He, Q., Xue, G., Xiao, L., & Bechara, A. (2014). Examination of neural systems sub-serving facebook "addiction". *Psychol Rep*, 115(3), 675-695. doi:10.2466/18.PR0.115c31z8.

Verdejo-Garcia, A., Garcia-Fernandez, G., & Dom, G. (2019). Cognition and addiction. *Dialogues in clinical neuroscience*, 21(3), 281. doi:10.31887/DCNS.2019.21.3/gdom.

Volkow, N. D., & Boyle, M. (2018). Neuroscience of addiction: relevance to prevention and treatment. *American Journal of Psychiatry*, 175(8), 729-740. doi:10.1176/appi.ajp.2018.17101174.

Wang, R., Li, M., Zhao, M., Yu, D., Hu, Y., Wiers, C. E., . . . Yuan, K. (2019). Internet gaming disorder: deficits in functional and structural connectivity in the ventral tegmental area-Accumbens pathway. *Brain imaging and behavior*, 13(4), 1172-1181. doi:10.1007/s11682-018-9929-6.

Widyanto, L., Griffiths, M. D., & Brunsden, V. (2011). A psychometric comparison of the Internet Addiction Test, the Internet-Related Problem Scale, and self-diagnosis. *Cyberpsychology, Behavior, and Social Networking*, 14(3), 141-149. doi:10.1089/cyber.2010.0151.

Xu, L., Bolt, T., Nomi, J. S., Li, J., Zheng, X., Fu, M., . . . Uddin, L. Q. (2020). Inter-subject phase synchronization differentiates neural networks underlying physical pain empathy. *Social cognitive and affective neuroscience*, 15(2), 225-233. doi:10.1093/scan/nsaa025.

Yan, C.G., Wang, X.D., Zuo, X.N., & Zang, Y.F., (2016). DPABI: data processing & analysis for (resting-state) brain imaging. *Neuroinformatics*, 14(3), 339-351. doi: 10.1007/s12021-016-9299-4.

Yan, H., Li, Q., Yu, K., & Zhao, G. (2021). Large-scale network dysfunction in youths with Internet gaming disorder: a meta-analysis of resting-state functional

connectivity studies. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 109, 110242. doi:10.1016/j.pnpbp.2021.110242.

Yao, Y.W., Liu, L., Worhunsky, P. D., Lichenstein, S., Ma, S.S., Zhu, L., . . . Yip, S. W. (2020). Is monetary reward processing altered in drug-naïve youth with a behavioral addiction? Findings from internet gaming disorder. *NeuroImage: Clinical*, 26, 102202. doi:10.1016/j.nicl.2020.102202.

Yoon, E. J., Choi, J.S., Kim, H., Sohn, B. K., Jung, H. Y., Lee, J.Y., . . . Kim, Y. K. (2017). Altered hippocampal volume and functional connectivity in males with Internet gaming disorder comparing to those with alcohol use disorder. *Scientific reports*, 7(1), 1-12. doi:10.1038/s41598-017-06057-7.

Young, K. S. (1998a). *Caught in the net: How to recognize the signs of internet addiction--and a winning strategy for recovery*: John Wiley & Sons.

Young, K. S. (1998b). Internet addiction: The emergence of a new clinical disorder. *Cyberpsychology & behavior*, 1(3), 237-244. doi:10.1089/cpb.1998.1.237.

Young, K. S. (1999). The evaluation and treatment of Internet addiction. *Innovations in clinical practice: A source book*, 17, 19-31.

Young, K. S. (2011). Clinical assessment of Internet-addicted clients. *Internet addiction: A handbook and guide to evaluation and treatment*, 19-34.

Yu, F., Sariyska, R., Lachmann, B., Wang, Q., Reuter, M., Weber, B., . . . Becker, B. (2021). Convergent cross-sectional and longitudinal evidence for gaming-cue specific posterior parietal dysregulations in early stages of internet gaming disorder. *Addict Biol*, 26(3), e12933. doi:10.1111/adb.12933.

Yuan, K., Yu, D., Cai, C., Feng, D., Li, Y., Bi, Y., . . . Tian, J. (2017). Frontostriatal circuits, resting state functional connectivity and cognitive control in internet gaming disorder. *Addict Biol*, 22(3), 813-822. doi:10.1111/adb.12348.

Zhang, J., Dong, H., Zhao, Z., Chen, S., Jiang, Q., Du, X., & Dong, G. H. (2020). Altered neural processing of negative stimuli in people with internet gaming disorder: fMRI evidence from the comparison with recreational game users. *J Affect Disord*, 264, 324-332. doi:10.1016/j.jad.2020.01.008.

Zhou, F., Li, J., Zhao, W., Xu, L., Zheng, X., Fu, M., . . . Becker, B. (2020a). Empathic pain evoked by sensory and emotional-communicative cues share common and process-specific neural representations. *eLife*, 9, e56929. doi:10.7554/eLife.56929.

Zhou, F., Montag, C., Sariyska, R., Lachmann, B., Reuter, M., Weber, B., . . . Becker, B. (2019a). Orbitofrontal gray matter deficits as marker of Internet gaming disorder: converging evidence from a cross-sectional and prospective longitudinal design. *Addiction Biology*, 24(1), 100-109. doi:10.1111/adb.12570.

Zhou, F., Zimmermann, K., Xin, F., Scheele, D., Dau, W., Banger, M., . . . Becker, B. (2018). Shifted balance of dorsal versus ventral striatal communication with frontal reward and regulatory regions in cannabis-dependent males. *Human brain mapping*, 39(12), 5062-5073. doi:10.1002/hbm.24345.

Zhou, X., Wu, R., Liu, C., Kou, J., Chen, Y., Pontes, H. M., . . . Montag, C. (2020b). Higher levels of (Internet) Gaming Disorder symptoms according to the WHO and APA frameworks associate with lower striatal volume. *Journal of behavioral addictions*, 9(3), 598-605. doi:10.1556/2006.2020.00066.

Zhou, X., Wu, R., Zeng, Y., Qi, Z., Ferraro, S., Yao, S., . . . Becker, B. (2021). Location, location, location—choice of Voxel-Based Morphometry processing pipeline drives variability in the location of neuroanatomical brain markers. *bioRxiv*. doi:10.1101/2021.03.09.434531.

Zhou, X., Zimmermann, K., Xin, F., Zhao, W., Derckx, R. T., Sassmannshausen, A., . . . Kendrick, K. M. (2019b). Cue reactivity in the ventral striatum characterizes heavy cannabis use, whereas reactivity in the dorsal striatum mediates dependent use. *Biological psychiatry: cognitive neuroscience and neuroimaging*, 4(8), 751-762. doi:10.1016/j.bpsc.2019.04.006.

Zhou, Y., Lin, F. C., Du, Y. S., Qin, L. D., Zhao, Z. M., Xu, J. R., & Lei, H. (2011). Gray matter abnormalities in Internet addiction: a voxel-based morphometry study. *Eur J Radiol*, 79(1), 92-95. doi:10.1016/j.ejrad.2009.10.025.

Zhuang, Q., Xu, L., Zhou, F., Yao, S., Zheng, X., Zhou, X., . . . Li, K. (2021). Segregating domain-general from emotional context-specific inhibitory control systems—ventral striatum and orbitofrontal cortex serve as emotion-cognition integration hubs. *Neuroimage*, 238, 118269. doi:10.1016/j.neuroimage.2021.118269.

Figures and tables

Figure 1 Experimental procedure (note: for the preprint version the faces have been overlaid with a blue circle)






Figure 2 Associations of two subscale scores and striatum based resting state functional connectivity



We firstly calculated the resting functional connectivity of bilateral ventral and dorsal striatum with voxels on the whole brain level, and conducted multiple regression analysis of connectivity, including loss of control subscale scores and craving/social problem subscale scores as covariates of interest, including age and gender as covariates of no interest. (A) The loss of control/time management subscale showed significant negative associations with resting state functional connectivity of the ventral striatum with right calcarine and right middle occipital lobe, left superior occipital lobe and right fusiform gyrus (FDR < 0.05, cluster size ≥ 49 voxels) (depicted in cyan). Loss of control/time management was significantly negative related to the resting state functional connectivity of dorsal striatum with right precuneus, right posterior cerebellum lobe and anterior cingulate cortex (FDR < 0.05, cluster size ≥ 36 voxels) (depicted in green).

B: The craving/social problems subscale showed a significant positive association with resting state functional connectivity of the ventral striatum with right fusiform, left superior occipital lobe, right middle occipital lobe, left inferior orbital frontal cortex, right superior occipital lobe, right superior temporal lobe and left insula (FDR < 0.05, cluster size ≥ 22 voxels) (depicted in yellow). The craving/social problems subscale showed a significant positive relation with functional connectivity strengths of the dorsal striatum with the left occipital lobe, left inferior frontal gyrus and right inferior frontal gyrus (FDR < 0.05, cluster size ≥ 27 voxels) (depicted in magenta).

Figure 3 Associations of two subscales with gray matter volume on the whole brain level.



A: Loss of control/time management was significantly positive associated with gray matter volume (GMV) of the left hippocampus (MNI coordinate: -15, -18, -15, FDR<0.05, cluster size = 530 voxels, see red region).

B: Craving/social problems showed a significant positive association with GMV of Caudate (MNI coordinate: 5, 8, 5, FDR $p < 0.05$, cluster size = 985 voxels, see red region). Purple represents the overlap with the caudate and bilateral dorsal striatum (green region); yellow represents regions representing the overlap between caudate and bilateral ventral striatum (blue region). Details see also table S7.

Note: L: left; R: right; A: anterior.

Table 1 General and symptom-domain levels of Internet Use disorders as measured by the s-IAT and its subscales

Questionnaire	Gender	Mean (SD)	<i>t</i>	<i>p</i>	Cohen's <i>d</i>
Total scores	female	33.07(8.01)	1.06	0.289	0.149
	male	31.93(7.23)			
Loss of control subscale scores	female	19.12(4.62)	2.68	0.008	0.377
	male	17.55(3.71)			
Craving/social problem subscale scores	female	13.95(4.03)	0.44	0.450	0.106
	male	14.38(4.15)			

Table 2 Associations between loss of control/time management and the striatum resting state networks

Scale	ROI	Direction	Connectivity region	MNI coordinate	Cluster size
Loss of control subscale scores	Ventral striatum	Negative	Right Calcarine	18, -78, 15	296
			Right middle occipital lobe	39, -84, 0	194
			Left superior occipital lobe	-12, -102, 15	62
	Dorsal striatum	Negative	Right fusiform gyrus	36, -63, -9	49
			Right precuneus	15, -60, 24	46
			Right cerebellum posterior lobe	9, -51, 45	42
			Anterior cingulate cortex	3, 45, 12	36

Note: FDR $p < 0.05$ for multiple comparison.

Table 3 Associations between craving / social problems and striatal networks

Scale	ROI	Direction	Connectivity region	MNI	Cluster
				coordinate	size
Craving/social problems	Ventral striatum	Positive	Right fusiform	30, -69, -15	458
			Left superior occipital lobe	-18, -93, 33	41
			Right middle occipital lobe	39, -84, 0	39
	Dorsal striatum	Positive	Left inferior orbital frontal cortex	-30, 21, -18	36
			Right superior occipital lobe	18, -84, 27	32
			Right superior temporal lobe	57, -48, 15	29
subscale scores			Left insula	-39, 9, -9	22
			Left occipital lobe	-9, -75, 15	56
			Left inferior frontal gyrus	-27, 21, -15	39
			Right inferior frontal gyrus	30, 24, -15	27

Note: FDR $p < 0.05$ for multiple comparison.