

1 Covalent Disruptor of YAP-TEAD Association Suppresses Defective Hippo 2 Signaling

3 **Authors:** Mengyang Fan^{1,2,4,10}, Wenchao Lu^{1,2,3,10}, Jianwei Che^{1,2}, Nicholas Kwiatkowski^{1,2},
4 Yang Gao^{1,2}, Hyuk-Soo Seo^{1,2}, Scott B. Ficarro^{1,5}, Prafulla C. Gokhale⁶, Yao Liu^{1,2}, Ezekiel
5 A. Geffken¹, Jimit Lakhani¹, Kijun Song¹, Miljan Kuljanin^{7,8}, Wenzhi Ji^{1,2,3}, Jie Jiang^{1,2},
6 Zhixiang He^{1,2}, Jason Tse³, Andrew S. Boghossian⁹, Matthew G. Rees⁹, Melissa M. Ronan⁹,
7 Jennifer A. Roth⁹, Joseph D. Mancias⁸, Jarrod A. Marto^{1,5}, Sirano Dhe-Paganon^{1,2}, Tinghu
8 Zhang^{1,2,3*}, Nathanael S. Gray^{1,2,3*}.

9 ¹Department of Cancer Biology, Dana-Farber Cancer Institute, Harvard Medical School,
10 Boston, MA 02215, USA

11 ²Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical
12 School, Boston, MA 02215, USA

13 ³Department of Chemical and Systems Biology, ChEM-H, Stanford Cancer Institute, School
14 of Medicine, Stanford University, Stanford, CA 94305, USA

15 ⁴Hangzhou Institute of Medicine (HIM), Chinese Academy of Sciences, Hangzhou, Zhejiang
16 310022, China

17 ⁵Blais Proteomics Center, Dana-Farber Cancer Institute, Harvard Medical School, Boston,
18 MA 02215, USA

19 ⁶Experimental Therapeutics Core, Dana-Farber Cancer Institute, Boston, MA 02215, USA

20 ⁷Department of Cell Biology, Harvard Medical School, Boston, MA 02215, USA

21 ⁸Division of Radiation and Genome Stability, Department of Radiation Oncology, Dana-
22 Farber Cancer Institute, Harvard Medical School, Boston, MA 02215, USA

23 ⁹Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA

24 ¹⁰These authors contributed equally to this work

25 Correspondence should be addressed to T.H.Z. (ztinghu8@stanford.edu) and N.S.G.
26 (nsgray01@stanford.edu).

27

28 **Keywords:** covalent fragment, transcription factor, YAP, TEAD, palmitoylation,

29 mesothelioma

30 Abstract

31 The transcription factor TEAD, together with its coactivator YAP/TAZ, is a key transcriptional
 32 modulator of the Hippo pathway. Activation of TEAD transcription by YAP has been implicated in a
 33 number of malignancies, and this complex represents a promising target for drug discovery. However,
 34 both YAP and its extensive binding interfaces to TEAD have been difficult to address using small
 35 molecules, mainly due to a lack of druggable pockets. TEAD is post-translationally modified by
 36 palmitoylation that targets a conserved cysteine at a central pocket, which provides an opportunity to
 37 develop cysteine-directed covalent small molecules for TEAD inhibition. Here, we employed covalent
 38 fragment screening approach followed by structure-based design to develop an irreversible TEAD
 39 inhibitor MYF-03-69. Using a range of *in vitro* and cell-based assays we demonstrated that through a
 40 covalent binding with TEAD palmitate pocket, MYF-03-69 disrupts YAP-TEAD association,
 41 suppresses TEAD transcriptional activity and inhibits cell growth of Hippo signaling defective
 42 malignant pleural mesothelioma (MPM). Further, a cell viability screening with a panel of 903 cancer
 43 cell lines indicated a high correlation between TEAD-YAP dependency and the sensitivity to MYF-
 44 03-69. Transcription profiling identified the upregulation of proapoptotic *BMF* gene in cancer cells
 45 that are sensitive to TEAD inhibition. Further optimization of MYF-03-69 led to an *in vivo*
 46 compatible compound MYF-03-176, which shows strong antitumor efficacy in MPM mouse
 47 xenograft model via oral administration. Taken together, we disclosed a story of the development of
 48 covalent TEAD inhibitors and its high therapeutic potential for clinic treatment for the cancers that are
 49 driven by TEAD-YAP alteration.

50 Introduction

51 The Hippo pathway is a highly conserved signaling pathway that regulates embryonic development,
 52 organ size, cell proliferation, tissue regeneration, homeostasis and is responsible for the development
 53 and progression of many malignancies¹. Critical components of Hippo signaling pathway are TEAD
 54 transcription factors, that are present as a family of four highly homologous members (TEAD1-4) in
 55 mammals. Importantly, TEADs alone show minimal transcriptional activity and require binding with
 56 a coactivator YAP/TAZ to initiate efficient gene expression². YAP has been well characterized as an

57 oncoprotein³ and its tumorigenesis role is mostly associated with TEAD mediated YAP-dependent
 58 gene expression⁴. Tissue-specific deletion of the upstream regulators or overexpression of YAP itself
 59 results in hyperplasia, organ overgrowth and tumorigenesis⁵. Tumor suppressor Merlin, encoded by
 60 *NF2* gene, and kinase LATS1/2 are known upstream regulators that cooperatively promote
 61 phosphorylation of YAP and hence induce its retention and degradation in the cytoplasm⁶. However,
 62 loss of function mutations in *NF2* or *LATS1/2*, which occurs in >70% mesothelioma, promote YAP
 63 nuclear entry and binding with TEAD to drive oncogenesis⁷. In addition to mesothelioma, high level
 64 of nuclear YAP has been associated with poor prognosis in non-small cell lung cancer (NSCLC),
 65 pancreatic cancer, and colorectal cancer (CRC)^{3a}. Moreover, there is evidence to suggest that
 66 activation of the YAP-TEAD transcriptional program can be involved with drug resistance⁸. For
 67 example, *ALK*-rearranged lung cancer cells were shown to survive the treatment of ALK inhibitor
 68 alectinib through YAP1 activation⁹. In EGFR-mutant lung cancers, YAP-TEAD was observed to
 69 promote cell survival and induce a dormant state upon EGFR signaling blockade¹⁰. Taken together,
 70 targeting YAP-TEAD has emerged as a promising therapeutic strategy for cancer treatment.

71 Over the decades, efforts to directly target YAP/TAZ-TEAD have focused on either designing
 72 peptide-mimicking agents that bind in the interface between YAP and TEAD and disrupt this
 73 interaction or using phenotypic screening to identify small molecules that inhibit YAP-dependent
 74 signaling¹¹. For example, a cyclic YAP-like peptide has been developed as a potent disruptor of YAP-
 75 TEAD complex in cell lysates but failed to exert effects in cells due to poor cellular permeability¹².
 76 Recently, TEAD stability and the association with YAP protein was shown to be regulated by S-
 77 palmitoylation, a post-translational modification on a conserved cysteine located within the palmitate
 78 binding pocket (PBP) on YAP binding domain (YBD) of TEAD¹³. This finding led to the discovery of
 79 several small molecules: flufenamic acid (FA)¹⁴, MGH-CP1¹⁵, compound 2¹⁶ and VT101~107¹⁷, that
 80 reversibly bind TEAD PBP (**Figure S1**). Additionally, inspired by thioester formation of the
 81 conserved cysteine with palmitate, covalent TEAD inhibitors TED-347¹⁸, DC-TEADin02¹⁹ and K-
 82 975²⁰ (**Figure S1**) have also been designed. TED-347 and DC-TEADin02 covalently bind with TEAD
 83 using a highly electrophilic chloromethyl ketone and vinyl sulfonamide warhead. In comparison, a

less electrophilic arylamide warhead is used in the design of K-975, which shows a potent cellular and *in vivo* activity in mouse tumor models albeit employing a high dose. Although these compounds have adequately demonstrated that the palmitoylated cysteine is targetable, none of them are optimized compounds and the structure-activity study for each of these scaffolds has not been disclosed. We have previously reported MYF-01-37 which possesses a less electrophilic warhead compared to K-975 as a covalent TEAD inhibitor and demonstrated its utility to blunt the transcriptional adaptation in mutant EGFR-dependent NSCLC cells¹⁰. However, MYF-01-37 is a sub-optimal chemical probe which requires micromolar concentrations in cellular assays and displays poor pharmacokinetic properties.

Development of covalent chemical probes and drugs has experienced a revival in recent years resulting in a deluge of new modality targeting a range of cancer-relevant targets, such as KRAS^{G12C}, EGFR, BTK, JAK3 and others²¹. The conventional approach to develop a covalent inhibitor is using a structure-guided design to equip a pre-existing reversible binder with an electrophile to target a nucleophilic residue on the target protein, most typically a cysteine. Alternatively, a recently developed approach involves screening a library of low molecular weight electrophilic fragments followed by medicinal chemical optimization²². Here, we used such screening approach to identify the lead covalent fragment MYF-01-37. Subsequent structure-based design to engage a side pocket resulted in development of a more potent covalent TEAD inhibitor MYF-03-69, which inhibits the palmitoylation process of all four TEAD paralogs *in vitro*, disassociates the endogenous YAP-TEAD complex, downregulates YAP-dependent target gene expression, and preferentially impairs the proliferation of mesothelioma cells that exhibit defects in Hippo signaling. A comprehensive screen against 903 cancer cell lines²³ identified more YAP-TEAD dependent cells as sensitive to MYF-03-69 in addition to mesothelioma. Transcription profiling suggests that the upregulation of Bcl-2 modifying factor (*BMF*) gene correlated with antiproliferative response to TEAD inhibition. Further, *in vivo* pharmacological effect was achieved with orally bioavailable compound MYF-03-176, indicating a promising lead for therapy of cancers driven by Hippo signaling dysfunction.

Results

Covalent fragment screening identified MYF-01-37 as a covalent TEAD binder

To screen for a covalent lead compound for TEAD, we first assembled a biased covalent fragment library that was developed based on the flufenamic acid template and a cysteine-reactive acrylamide warhead²⁴ (**Figure 1a, Figure S2a**). We then conducted a medium-throughput screen of this library using mass spectrometry (MS) to detect protein-fragment adduct formation with recombinant, purified TEAD2 YBD protein. MYF-01-37 was identified as the fragment capable of efficient labeling while the majority of the fragment library failed to label the protein (**Figure S2b**). Next, the protein labeled with MYF-01-37 was proteolytically digested and the resultant peptides were analyzed using tandem mass spectrometry (MS/MS) which confirmed labeling of C380, which is the cysteine previously reported to be S-palmitoylated in TEAD2^{13a} (**Figure 1a**).

Structure-based optimization yields MYF-03-69, an irreversible inhibitor of TEAD palmitoylation

Next, we employed a structure-based fragment growing strategy to elaborate MYF-01-37 to develop a more potent and selective inhibitor (**Figure 1a**). Given that the site of covalent modification is located at the opening of the palmitate binding pocket, we analyzed the TEAD palmitate pockets from multiple crystal structures available in the Protein Data Bank (PDB)¹³⁻¹⁴. All the structures aligned well, indicating a conserved and relatively inflexible pocket. As shown in the **Figure 1b**, docking of MYF-01-37 and TEAD2 revealed that MYF-01-37 binds in a tunnel formed by side chains of hydrophobic residues F233, F251, V252, V329, V347, M379, L383, L387, F406 and F428. We also identified a side pocket lined with hydrophilic side chains contributed by S331, S345, S377 and Q410. We hypothesized that we could improve potency and selectivity for our covalent ligands by introducing polar interactions and shape complementarity to this side pocket. Therefore, we predicted that the optimized binding pocket is Y-shaped with the targeted cysteine at one end, and hydrophobic and hydrophilic pockets at the other two ends. We envisioned that occupying the hydrophilic side pocket will also provide chance to incorporate polar moiety, leading to reduced global hydrophobicity and more druglike molecules. With this rationale, a variety of hydrophilic groups were introduced to

the pyrrolidine ring, leading to synthesis of a focused library of Y-shaped molecules, with representatives shown in **Figure 1c**. We employed an *in vitro* palmitoylation assay that uses an alkyne-tagged palmitoyl CoA as the lipidation substrate and clickable biotin (**Figure 1d**), to screen this series of compounds for their ability to inhibit palmitoylation (**Figure 1e**, **Figure S3**). We observed that the *para* substituted trifluoromethyl benzyl is superior to *meta* substituted trifluoromethyl benzyl (MYF-02-111 vs. MYF-03-42). We then replaced pyrazole by a substituted triazole to explore the hydrophilic pocket (MYF-03-135, MYF-03-162, MYF-03-138, MYF-03-146, MYF-03-69), and observed a sensitivity to various substitutions at C4 position of the triazole ring. For example, tetrahydropyran and pyridine enhanced the activities, whereas piperidine and pyridone were much less favored. In addition, we noticed that extending the hydrophobic tail of MYF-01-37 allowed more complete occupancy of the hydrophobic channel. Further modifications of the groups occupying the hydrophobic tunnel did not yield further improvement on the potency compared to trifluoromethyl benzyl (MYF-03-137, MYF-03-139), implicating restrict requirement for a hydrophobic moiety. Amongst this series of compounds, we identified MYF-03-69 (**Figure 1c**) as the most potent compound, and chose it for further characterization.

MYF-03-69 occupies the Y-shaped pocket and forms covalent bond with the conserved cysteine of TEAD

Given that labeling efficiency of the parental fragment MYF-01-37 was moderate, we examined whether the elaborated, Y-shaped molecule labeled TEAD more efficiently. Recombinant TEAD2 YBD was incubated with MYF-03-69, followed by mass spectrometry. Incubation with 10-fold molar excess of MYF-03-69 at room temperature for 1 hour led to 100% labeling (**Figure 2a**) and follow-up trypsin digestion and MS/MS analysis confirmed C380 as the modified residue (**Figure 2b**). This covalent bond formation and the exact binding mode was further validated by obtaining a high-resolution co-crystal structure of TEAD1 YBD in complex with MYF-03-69 (1.68 Å) (**Figure 2c**). Unambiguous electron density of a covalent bond was observed between TEAD1 C359 (analogous to C380 in TEAD2) and the acrylamide warhead of MYF-03-69 with the remaining part of the molecule

adopting a Y-shape and establishing specific interactions with the hydrophobic pocket and hydrophilic patch as predicted (**Figure 2c**). MYF-03-69 was completely buried in the lipid pocket with the *para*-trifluoromethyl benzyl group extended along a lipid trajectory, forming extensive hydrophobic interactions with side chains of A223, F239, V240, A292, I366, and F407. The 3-pyridinyl group occupied the side pocket, forming a favorable water-bridged hydrogen bond network with adjacent S310 and Y312 via pyridinyl nitrogen (**Figure 2d**). Overall, high resolution structural analysis confirms that MYF-03-69 forms covalent attachment with the TEAD cysteine that is the site of S-palmitoylation, and that the elaborated molecule forms specific interactions via both hydrophobic and hydrophilic regions surrounding the targeted cysteine.

To further characterize biochemical activity of MYF-03-69, we performed a dose titration for all TEAD paralogs (**Figure 2e, Figure S4**). Preincubation of all recombinant TEAD1-4 YBDs with MYF-03-69 potently inhibited palmitoylation with similar IC₅₀ values at submicromolar concentrations, indicating that MYF-03-69 is a pan-TEAD inhibitor. This is consistent with the high sequence conservation of residues in the palmitate pockets of TEAD1-4. In contrast, MYF-03-69-NC (**Figure 1c**), the non-covalent counterpart of MYF-03-69 incapable of forming covalent bond, completely lost activity across all TEADs, which demonstrated the essentiality of covalent bond formation.

MYF-03-69 inhibits TEAD palmitoylation and disrupts endogenous YAP-TEAD association in cells

We next investigated the ability of MYF-03-69 to modulate TEAD palmitoylation inside cells with ω -alkynyl palmitic acid (Alkyne-C16) as probe. Alkyne-C16 contains an alkyne group at C15 and has been shown to be metabolically incorporated into cellular proteins at native palmitoylation sites. The previously reported pulse-chase style experiment¹⁶ was adopted and modified to monitor TEAD palmitoylation in HEK293T cells transfected with Myc-TEAD4. After the cells were incubated with Alkyne-C16 for 24 hours, the labeled Myc-TEAD4 was conjugated with biotin-azide through a Cu(I)-catalyzed click reaction. With western-blotting, we found that the alkyne-palmitate labeling of Myc-

TEAD4 decreased after cotreatment with varying concentrations of MYF-03-69 for 24 hours compared to the DMSO treatment, while TEAD protein levels were not affected (**Figure 3a**). This result suggested that MYF-03-69 competed off palmitic acid during TEAD lipidation in cells. Consistently with biochemical result, the negative control compound MYF-03-69-NC did not exhibit any effect on TEAD (**Figure 3a**).

Palmitoylation of TEADs was recently proposed to be required for their binding to YAP and TAZ^{13b}. Although TEAD inhibitors with different chemotypes have been recently developed, it remains controversial whether pharmacological targeting of the TEAD palmitate pocket could disrupt YAP/TAZ association²⁵. In order to investigate whether MYF-03-69 impairs YAP-TEAD interaction, we conducted an endogenous co-immunoprecipitation (co-IP) experiment to monitor YAP-TEAD interactions in the presence of the compound. Notably, 24-hour treatment of NCI-H226 cells with MYF-03-69 significantly decreased endogenous YAP co-immunoprecipitation with a pan-TEAD antibody, whereas this treatment had minimal effect on TEAD protein level (**Figure 3b**). In parallel, we also evaluated MYF-03-69-NC and observed no effect on YAP-TEAD association in cells. Taken together, these results suggest that MYF-03-69-mediated perturbation of TEAD palmitoylation disrupts YAP and TEAD protein interactions.

To demonstrate MYF-03-69 as a selective covalent TEAD binder, we interrogated the proteome-wide reactivity profile of MYF-03-69 on cysteines using a well-established streamlined cysteine activity-based protein profiling (SLC-ABPP) approach (**Figure S5a**)²⁶. We employed the cysteine reactive desthiobiotin iodoacetamide (DBIA) probe²⁶ which was reported to map more than 8,000 cysteines and performed a competition study in NCI-H226 cells pretreated with 0.5, 2, 10 or 25 μ M of MYF-03-69 for 3 hours in triplicate. The cysteines that were conjugated >50% (competition ratio CR>2) compared to DMSO control were analyzed and assigned to the protein targets (**Figure S5b**). In the DMSO control group, although DBIA mapped 12,498 cysteines in total, the TEAD PBP cysteines were not detected. Among 12,498 mapped cysteines, only 7 cysteines were significantly labeled (i.e. exhibited >50% conjugation or CR>2) by 25 μ M of MYF-03-69, and all of these sites exhibited dose-

dependent engagement (**Supplementary Dataset 1**). The missing TEAD cysteines might be due to low TEAD1-4 protein abundance and/or inability of the PBP cysteines to be labeled by DBIA. DBIA probe can only be applied in cell lysate context, which might also result in the missing labeling on TEAD. To our knowledge, all the proteins with meaningful labeling efficiency (CR>2) cysteine sites are not known to be involved in the YAP/TAZ-TEAD signaling pathway. Taken together, our proteomic analysis suggests that MYF-03-69 exhibited quite low reactivity towards the cysteine proteome, although TEAD PBP cysteines were not detected by the DBIA probe.

MYF-03-69 inhibits TEAD transcription and downregulates YAP target genes expression in mesothelioma cells

The majority of mesothelioma patients (50-75%) harbor genetic alterations in Hippo pathway regulatory components, including *NF2* loss of function mutation/deletion and *LATS1-PSEN1* fusion/*LATS2* deletion, which lead to YAP activation and Hippo pathway gene expression⁷. Thus, in order to monitor YAP-TEAD transcriptional activity, Hippo/YAP reporter cells were generated in NCI-H226, a *NF2*-deficient MPM cell line. After 72-hour treatment of MYF-03-69, YAP-TEAD transcriptional activity of reporter cells significantly decreased in a dose-dependent manner with an IC₅₀ value of 56 nM, while MYF-03-69-NC had minimal effect (**Figure 4a**). The transcriptional inhibition also led to significant downregulation of canonical YAP target genes including *CTGF*, *CYR61*, *IGFBP3*, *KRT34* and *NPPB* and upregulation of proapoptotic gene *BMF* in NCI-H226 cells and MSTO-211H, a *LATS1-PSEN1* fusion MPM cell line (**Figure 4b-c**), while it showed much milder effects on transcription in normal mesothelium cells MeT-5A (**Figure S6**). Protein level of *CYR61* and *AXL* gene products also decreased (**Figure 4d, Figure S7**). Encouraged by these results, we set out to study whole transcriptome perturbation by MYF-03-69. RNA sequencing was performed in NCI-H226 cells that were treated with 0.1 μM, 0.5 μM, and 2 μM of MYF-03-69. There were 339 genes that exhibited significant differential expression at 2 μM treatment, and the majority of them changed in a dose dependent manner (**Figure 4e** and **Supplementary Dataset 2**). The genes that were differentially expressed with statistical significance (Fold change ≥ 1.5 and adjusted p value ≤ 0.05)

at 2 μ M treatment condition, are colored in red and partially labeled in the volcano plot (**Figure 4f**). Compared to high concentration, the number of genes with significantly altered expression level dropped off quickly at lower concentrations (**Figure S8a-b**). For example, at the concentration of 0.1 μ M, only *CTGF* showed statistically significant change (**Figure S8b**). While multiple YAP target genes such as *CTGF*, *ADM*, *ANKRD1* were significantly downregulated², *DDIT4* was observed to be upregulated²⁷. To investigate whether the 339 differentially expressed genes were concentrated in particular biological pathways, KEGG pathway enrichment calculation²⁸ were carried out, which demonstrated Hippo signaling pathway was among the top 5 enriched processes (**Figure 4g**). Beyond the Hippo pathway, gap junction²⁹ and WNT signaling pathway³⁰ were also enriched, consistent with pleiotropic functions and cross-talk of YAP-TEAD pathway in diverse biological processes. Overall, we documented that MYF-03-69 affects transcription in a manner consistent with its activity as a disruptor of YAP-TEAD interactions.

MYF-03-69 selectively inhibits mesothelioma cancer cells with defective Hippo signaling

Next, we investigated anti-cancer activity of MYF-03-69 on Hippo signaling defective mesothelioma cells. As shown in **Figure 5a** and **Figure S9a**, 5-day cell growth assays demonstrated that MYF-03-69 potently retarded the cell growth of NCI-H226 and MSTO-211H, while it showed no antiproliferation activity against MeT-5A and NCI-H2452 cells, which are non-cancerous mesothelium cells and mesothelioma cells with intact Hippo signaling, respectively. These antiproliferative effects in NCI-H226 and MSTO-211H cells were observed under 3D spheroid suspensions culture condition as well (**Figure 5b**). Under the same conditions, the negative control compound MYF-03-69-NC was not antiproliferative (**Figure 5b, Figure S9b**). Further, cell cycle analysis demonstrated that 48-hour treatment with MYF-03-69 on NCI-H226 and MSTO-211H cells caused cell cycle arrest at the G1 phase, which is in accordance with previous findings obtained from genetic knockdown of YAP³¹, while negative control compound had no effect (**Figure 5c-d**). Collectively, these results show that inhibition of TEAD palmitoylation by MYF-03-69 effectively and precisely affects YAP-TEAD function.

MYF-03-69 inhibits YAP or TEAD-dependent cancer cells beyond mesothelioma

To further investigate whether inhibition by MYF-03-69 was selectively lethal to YAP/TEAD-dependent cancers, 903 barcoded cancer cell lines were screened using the PRISM assay²³. As shown in **Figure 5e**, a small portion of cell lineages exhibited vulnerability (**Supplementary Dataset 3**). Correlation analysis reveals that the dependency scores of *TEAD1* and *YAP1* according to genomic knockout dataset (DepMap) provided the highest correlation with the sensitivity profile (**Figure S9c**, **Supplementary Dataset 4**). This is followed by *TP53BP2*, a gene that is also involved in Hippo pathway as activator of TAZ³². For example, when we used a threshold of $AUC \leq 0.8$ for the sensitivity to MYF-03-69, 33 cell lines were selected and the majority of which are either YAP or TEADs dependent cells, as suggested by CERES scores (**Figure 5e**). These include YAP/TEAD co-dependent cells (red dots), YAP dependent cells (blue dots), and TEAD dependent cells (yellow dots). Next, to verify the antiproliferative activity, we chose three sensitive cell lines (94T778, SKHEP-1 and HuCCT1) and three insensitive cell lines (93T449, MIA PaCa-2 and MM.1S) indicated by PRISM screening, as well as two additional cell lines (PC9, HEK293T) that were not included in PRISM screening panel to test in 5-day antiproliferation assay. As shown in **Figure 5f**, liposarcoma cell 94T778, hepatic adenocarcinoma cell SKHEP-1 and cholangiocarcinoma cell HuCCT1 were inhibited with nanomolar IC_{50} . In contrast, liposarcoma cell 93T449, pancreatic ductal adenocarcinoma cell MIA PaCa-2 and myeloma cell MM.1S were barely inhibited (**Figure 5g**). Since proapoptotic gene *BMF* was known to be released from repression upon TEAD inhibition¹⁰, we examined *BMF* mRNA levels after 6-hour treatment with 0.5 and 2 μ M MYF-03-69 on both sensitive and insensitive cells. As expected, *BMF* levels increased in three sensitive cells, but remained unchanged in three insensitive cells (**Figure 5h**). We also noted the upregulation of *BMF* in PC9 and HEK293T cells which are resistant to TEAD inhibition (**Figure 5g-h**). A heterogeneity of signaling pathway dependency, especially EGFR signaling and YAP1 activation in PC9 cells might account for the insensitivity to TEAD inhibition alone. Taken together, we demonstrate that TEAD inhibition could be an exploitable vulnerability across multiple malignant tumor models besides mesothelioma

and that upregulation of *BMF* gene is a common phenomenon in those cancer cells with sensitive antiproliferative response to TEAD inhibitor.

MYF-03-176 exhibits significant antitumor efficacy in NCI-H226 xenograft mouse model through oral administration

In order to demonstrate therapeutic potential of covalent TEAD palmitoylation disruptor, extensive medicinal chemistry efforts were undertaken on this Y-shaped scaffold, leading to a more potent and more importantly, orally bioavailable MYF-03-69 analog MYF-03-176 (**Figure 6a**). Before we administrated the mice with MYF-03-176, we conducted a head-to-head comparison on the activity with MYF-03-69 as well as other reported TEAD inhibitors including K-975, compound 2 and VT103. We used a stably transfected TEAD luciferase reporter system in NCI-H226 cells to profile transcriptional effect of TEAD inhibitors. Consistently with the results of mCherry reporter system, MYF-03-69 inhibited the transcription of reporter gene at IC₅₀ of 45 nM (**Figure S10a**) and interestingly MYF-03-176 was 3-fold more potent with IC₅₀ of 11 nM (**Figure 6b**). Further, similar to MYF-03-69, MYF-03-176 led to a significant downregulation of YAP target genes *CTGF*, *CYR61*, *ANKRD1* and an upregulation of *BMF* (**Figure S10b**). As comparison, MYF-03-176 exhibited comparable and even better activity than K-975 and compound 2, while was less potent than VT103 as shown in **Figure 6b~c**. The antiproliferative effect of MYF-03-176 in NCI-H226 cells was similar to VT-103 and K-975 but even stronger than MYF-03-69 and compound 2. With the support of all these data from MYF-03-176 and decent pharmacokinetics properties including low clearance and high oral bioavailability (**Figure S10c-d**), we then administrated MYF-03-176 to NCI-H226 cell line derived xenograft (CDX) mouse model. The tumor-bearing mice were randomized and orally administrated MYF-03-176 twice daily for 28 days. Significant antitumor activity with tumor regressions was observed at both 30 mg/kg (average tumor regression of 54%) and 75 mg/kg (average tumor regression of 68%). The anti-tumor activity at the two doses was comparable ($P = 0.23$, 2-way ANOVA). The 30 mg/kg dose was well tolerated with an average body weight gain comparable to vehicle control (data not shown), while at the 75 mg/kg dose, an average weight loss of 5% was

observed. However, at this 75 mg/kg dose, 3 of 8 animals demonstrated 12-14% body weight loss. The weight loss recovered once drug administration was stopped (**Figure 6f**). Taking all these together, MYF-03-176 potently inhibited Hippo signaling defective MPM cells and exhibited strong antitumor effect in the human mesothelioma NCI-H226 CDX model *in vivo*, representing a promising leading compound for drug discovery.

Discussion

As widely recognized oncogenic proteins, YAP/TAZ have emerged as potentially attractive targets for anti-cancer drug development. However, the unstructured nature of YAP/TAZ proteins renders them difficult to target by conventional occupancy-based small molecules. Therefore, the majority of compounds that are known to inhibit YAP activity are targeted at upstream stimulators of YAP/TAZ. In this context, TEADs, key components of Hippo signaling pathway that depend on YAP/TAZ binding for activation of transcriptional activity, have attracted attention. Here, the presence of a well-defined palmitate binding pocket (PBP) on TEADs suggested opportunity for small molecule inhibitor development. However, recent studies have not fully resolved the question of whether occupying TEAD PBP disrupts or stabilizes YAP-TEAD interaction. Chan et al. suggested that palmitoylation of TEAD stabilizes YAP-TEAD interaction^{13b}. Holden et al. reported the reversible PBP binder compound 2 had minimal to no disruptive effect on YAP-TEAD interaction biochemically but transformed TEAD into a dominant-negative transcriptional repressor¹⁶. VT series compounds reported by Tang et al. dissociated endogenous YAP-TEAD according to co-IP experiment¹⁷, which is consistent with our result of MYF-03-69. Other data from MGH-CP1¹⁵ and covalent binder K-975²⁰ pointed to the same conclusion while the experiments were performed with exogenous YAP or TEAD. K-975 and TED-347 were reported to disrupt protein-protein interaction between TEAD and YAP peptide in biochemical assay. However, their co-crystal structures indicated no conformation change compared with palmitoylated TEAD. Similarly, superimposition of co-crystal structures of TEADs with MGH-CP1, compound 2, VT105, MYF-03-69 and their corresponding palmitoylated TEADs did not reveal any obvious conformation change or side chain move that might affect YAP binding.


Taken all these together, the transcriptional inhibitory effect of these TEAD PBP binders might result from disruption of certain dynamic process of TEAD lifecycle. Recently, several reports elaborated that YAP/TAZ and TEAD undergo lipid-lipid phase separation (LLPS) during transcription process³³. One hypothesis is the alkyl chain of palmitoyl might escape from TEAD PBP and expose to outside when forming such functional transcription compartment. The hydrophobic nature of the lipid chain may play a role in organizing the disordered and hydrophobic region of YAP. Thus, replacement of palmitate by these PBP binders leads to incapability of TEAD to organize into transcription machinery, though the direct binding to YAP peptide may not being affected *in vitro*. The underlying mechanism is remained to be uncovered in the future.

MYF-03-69 is a covalent compound that we designed to target the conserved cysteine on TEAD, the site of palmitoylation. The starting point for MYF-03-69 was a fragment hit that we identified through screening of a biased covalent fragment library and further optimized using structure-based strategy. Unlike screening reversible fragments which often require significant follow-up efforts to determine binding site, the MS-guided covalent tethering method gives unambiguous information of labeling site in a cost and time efficient way. Unlike previously reported compounds that engage only the hydrophobic PBP, our optimized inhibitor was developed to exploit interactions with not only the PBP but a hydrophilic binding pocket that we have identified during this study. Thus, our optimized inhibitor MYF-03-69 is a Y-shaped molecule which efficiently occupies two pockets, as well as covalently binds the conserved cysteine, which contributed significantly to both potency and specificity. Therefore, we speculated that MYF-03-69 may be employed as a useful tool to interrogate the process of how PBP binder affect TEAD homeostasis. We also propose that further optimization of the basic Y-shaped molecule reported here may yield paralog-selective TEAD inhibitors.

Aberrant regulation of Hippo-YAP/TAZ-TEAD signaling axis has been recognized as driver for genesis and development of multiple cancers, especially in mesothelioma. Besides the Hippo pathway, diverse signaling pathways such as Wnt, TGF- β , EGFR and Hedgehog pathways can also potentiate the activation of YAP/TAZ. Given such essentiality, it is important to ascertain what portion of these YAP/TAZ activated cancers are vulnerable to TEAD inhibition. The pan-TEAD selective nature of

MYF-03-69 allowed us to examine this question at the TEAD family level using a panel of more than 900 cancer cell lines. Our results demonstrated that MYF-03-69 exhibited selective antiproliferative effect on YAP or TEAD dependent cancer cells from different lineages including mesothelioma, liver cancer, liposarcoma, and lung cancer. Strong antitumor efficacy in human mesothelioma NCI-H226 CDX model has been achieved with an orally bioavailable compound MYF-03-176 in the same scaffold, which is ready to be tested in various cancer models. Moreover, we noted that increased expression of a proapoptotic *BMF* gene correlates with response to TEAD inhibition in some cancer cell lines.

Collectively, this study provides evidence that MYF-03-69 represents a potent, covalent, cysteine-targeted pan-TEAD inhibitor that disrupts YAP-TEAD association and affects transcription in covalent binding-dependent manner. Given its “mild” covalent warhead and low reactivity across proteome, we nominate MYF-03-69 as a viable lead for drug discovery for not only mesothelioma, but other YAP or TEAD dependent cancers such as liver cancer, liposarcoma, and lung cancer.






Figure 1 | Covalent fragments screening and structure-guided design to identify Y-shaped compound MYF-03-69. (a) Illustration of covalent fragment library screening and optimization for TEAD inhibitor. (b) Surface of TEAD2 palmitate pocket depicted in mesh with MYF-01-37 (in green) modeled in. Residues forming the pocket were labeled. The color of mesh indicates hydrophobicity of the pocket surface. (c) Chemical structures of representative Y-shaped compounds. (d) The schematic diagram of *in vitro* palmitoylation assay. (e) Anti-palmitoylation activity of MYF-01-37 and Y-shaped compounds after TEAD2 protein was pre-incubated with 2 μ M compound at 37°C for 2 hours.




Figure 2 | MYF-03-69 binds in TEAD palmitate pocket covalently through the conserved cysteine. (a) The mass labeling of intact TEAD2 protein by MYF-03-69. (b) Trypsin digestion and tandem mass spectrum (MS/MS) localize labeling site of cysteine 380. (c) Co-crystal structure of MYF-03-69 with TEAD1 indicates covalent bond formation with Cys359. The compound adopts Y-shape and binds in both lipid tunnel and hydrophilic side pocket. (d) Interactions between MYF-03-69 and TEAD1 palmitate pocket. (e) A dose titration of MYF-03-69 and MYF-03-69-NC in anti-palmitoylation assay on TEAD1-4. Recombinant YBD protein of TEADs were preincubated with compounds at 37°C for 2 hours. Data are representative of n = 3 independent experiments.




Figure 3 | MYF-03-69 inhibits palmitoylation of TEAD protein and disrupts its association with YAP in cells. (a) Palmitoylation of Myc-TEAD4 in HEK293T cells after treatment with MYF-03-69 and MYF-03-69-NC indicated by an alkyne-palmitic-acid probe and click chemistry. Cells were treated for 24 hours. (b) Co-immunoprecipitation (Co-IP) of endogenous YAP and TEAD in NCI-H226 cells after treatment with MYF-03-69 and MYF-03-69-NC at indicated doses. Cells were treated for 24 hours.





Figure 4 | MYF-03-69 inhibits YAP-TEAD transcription. (a) MYF-03-69, but not MYF-03-69-NC, inhibits YAP-TEAD transcriptional activity in NCI-H226 mCherry reporter cells. Cells were treated for 72 hours. (b-c) MYF-03-69, but not MYF-03-69-NC, downregulates YAP target genes and upregulates a pro-apoptotic gene *BMF*. Cells were treated for 24 hours. Data were presented as mean $\pm$ SD of $n = 3$ biological independent samples. (d) MYF-03-69, but not MYF-03-69-NC, downregulates CYR61 protein level in NCI-H226 and MSTO-211H cells. (e) Heatmap for gene

424 expression change with MYF-03-69 treatment at indicated concentrations. (f) Differential gene
425 expression from RNA sequencing of NCI-H226 cells treated with 2 μ M MYF-03-69. The
426 differentially expressed genes with $FC \geq 1.5$ and $p \leq 0.05$ were colored in red and labeled. Not all
427 differentially expressed genes were labeled. (g) Pathway enrichment analysis of differentially
428 expressed genes from 2 μ M compound treatment samples.

429



430




Figure 5 | MYF-03-69 selectively inhibits proliferation of cancer cells with YAP or TEAD dependency regardless of lineage. (a) Antiproliferation IC_{50} curve of MYF-03-69 on three mesothelioma cell lines (NCI-H226, MSTO-211H and NCI-H2452) and a normal noncancerous mesothelium cell line (Met-5A). (b) Antiproliferation IC_{50} curve of MYF-03-69 and MYF-03-69-NC in 3D cell culture. (c-d) Cell cycle arrest induced by MYF-03-69, but not MYF-03-69-NC. Cells were treated for 48 hours at indicated doses. (e) PRISM profiling across a broad panel of cell lineages. 903 cancer cells were treated with MYF-03-69 for 5 days. The viability values were measured at 8-point dose manner (3-fold dilution from 10 μ M) and fitted a dose-response curve for each cell line. Area under the curve (AUC) was calculated as a measurement of compound effect on cell viability. CERES score of YAP1 or TEADs from CRISPR (Avana) Public 21Q1 dataset (DepMap) were used to estimate gene-dependency. Cell lines without CERES Score of YAP1 were excluded from the figure. The CERES Score of most dependent TEAD isoform was used to represent TEAD dependency. Cell lines with a dependency score less than -0.6 were defined as the dependent cell lines. (f) Antiproliferation curves of cell lines that are sensitive to MYF-03-69 treatment besides mesothelioma.

(g) Antiproliferation curves of cell lines that are insensitive to MYF-03-69 treatment. (h) *BMF* expression level after treatment with 0.5 or 2 μ M MYF-03-69 over 6 hours in different cells.




Figure 6 | MYF-03-176 is a potent and orally bioavailable YAP-TEAD transcription inhibitor and suppresses tumor growth in mesothelioma xenograft mouse model. (a) Chemical structure of MYF-03-176. (b-c) Inhibitory effect of MYF-03-176 and other TEAD PBP binders on YAP-TEAD transcription in NCI-H226 luciferase reporter cells. Cells were treated for 72 hours. Data were presented as mean $\pm$ SD of n = 3 biological independent samples. (d-e) Antiproliferation effect of MYF-03-176 and other TEAD PBP binders in NCI-H226 cells. Cells were treated for 5 days. Data were presented as mean $\pm$ SD of n = 3 biological independent samples. (f) *In vivo* efficacy of MYF-03-176 in NCI-H226 CDX mouse model (n = 8~9 per group).

Acknowledgments: The authors would like to thank the following for valuable help with this study:

Dr. Milka. Kostic for her advice and editing for this manuscript; Jim Sun at the NMR facility of the Dana-Farber Cancer Institute for his assistance on ¹H NMR data collection; Zachary Herbert from the Molecular Biology Core Facility at the Dana-Farber Cancer Institute for the sequencing services; Kara Soroko and Jessica Sarro from Dana-Farber Cancer Institute for the animal study services. J.D.M. acknowledges this work was supported by the Hale Family Center for Pancreatic Cancer Research.

Author contributions: N.S.G. and T.Z. conceived the project. M.F. performed the compound synthesis and structure determination with help from Y.L., W.J. and Z.H.. W.L. executed biological experimental research with help from Y.G., N.K., J.J. and J.T.. J.C. executed computational modeling and RNA sequencing analysis. S.B.F. and J.A.M. executed and analyzed protein mass spectrum. H.S., E.A.G., J.L., K.S. and S.D.P. executed protein expression, purification, crystallization and structural determination. P.C.G. managed the animal study and analyzed the results. M.K. and J.D.M. performed the SLC-ABPP profiling and analysis. A.S.B., M.G.R., M.M.R. and J.A.R. executed PRISM screening and analysis. M.F., W.L., T.Z., J.C. and N.S.G. co-wrote the paper. All authors edited the manuscript.

Declaration of interests: N.S.G. is a founder, science advisory board (SAB) member and equity holder in Syros, Jengu, C4, B2S, Allorion, Inception, GSK, Larkspur (board member) and Soltego (board member). The Gray lab receives or has received research funding from Novartis, Takeda, Astellas, Taiho, Janssen, Kinogen, Voronoi, Interline, Springworks and Sanofi. T.Z. is a consultant and equity holder of EoCys. J.C. is a consultant to Soltego, Jengu, Allorion, EoCys, and equity holder for Soltego, Allorion, EoCys, and M3 bioinformatics & technology Inc. The Gray lab has sponsored research agreement for TEAD inhibitor project with Epiphanes. M.F., W.L., Y.L., J.C., Y.G., N.K., T.Z. and N.S.G. are inventors on TEAD inhibitor patents.

Reference

- 485 1. (a) Ota, M.; Sasaki, H., Mammalian Tead proteins regulate cell proliferation and contact
486 inhibition as transcriptional mediators of Hippo signaling. *Development* **2008**, *135* (24), 4059-4069;
- 487 (b) Pan, D., The Hippo Signaling Pathway in Development and Cancer. *Developmental Cell* **2010**, *19*
488 (4), 491-505; (c) Harvey, K. F.; Zhang, X.; Thomas, D. M., The Hippo pathway and human cancer.
489 *Nature Reviews Cancer* **2013**, *13* (4), 246-257; (d) Yu, F.-X.; Zhao, B.; Guan, K.-L., Hippo Pathway
490 in Organ Size Control, Tissue Homeostasis, and Cancer. *Cell* **2015**, *163* (4), 811-828.
- 491 2. Zhao, B.; Ye, X.; Yu, J.; Li, L.; Li, W.; Li, S.; Yu, J.; Lin, J. D.; Wang, C.-Y.; Chinnaiyan, A.
492 M.; Lai, Z.-C.; Guan, K.-L., TEAD mediates YAP-dependent gene induction and growth control.
493 *Genes & Development* **2008**, *22* (14), 1962-1971.
- 494 3. (a) Zanconato, F.; Cordenonsi, M.; Piccolo, S., YAP/TAZ at the Roots of Cancer. *Cancer*
495 *Cell* **2016**, *29* (6), 783-803; (b) Zanconato, F.; Cordenonsi, M.; Piccolo, S., YAP and TAZ: a
496 signalling hub of the tumour microenvironment. *Nature Reviews Cancer* **2019**, *19* (8), 454-464; (c)
497 Moroishi, T.; Hansen, C. G.; Guan, K.-L., The emerging roles of YAP and TAZ in cancer. *Nature*
498 *Reviews Cancer* **2015**, *15* (2), 73-79.
- 499 4. (a) Stein, C.; Bardet, A. F.; Roma, G.; Bergling, S.; Clay, I.; Ruchti, A.; Agarinis, C.;
500 Schmelzle, T.; Bouwmeester, T.; Schübeler, D.; Bauer, A., YAP1 Exerts Its Transcriptional Control
501 via TEAD-Mediated Activation of Enhancers. *PLOS Genetics* **2015**, *11* (8), e1005465; (b) Zanconato,
502 F.; Forcato, M.; Battilana, G.; Azzolin, L.; Quaranta, E.; Bodega, B.; Rosato, A.; Bicciato, S.;
503 Cordenonsi, M.; Piccolo, S., Genome-wide association between YAP/TAZ/TEAD and AP-1 at
504 enhancers drives oncogenic growth. *Nature Cell Biology* **2015**, *17* (9), 1218-1227; (c) Galli,
505 Giorgio G.; Carrara, M.; Yuan, W.-C.; Valdes-Quezada, C.; Gurung, B.; Pepe-Mooney, B.; Zhang, T.;
506 Geeven, G.; Gray, Nathanael S.; de Laat, W.; Calogero, Raffaele A.; Camargo, Fernando D., YAP
507 Drives Growth by Controlling Transcriptional Pause Release from Dynamic Enhancers. *Molecular*
508 *Cell* **2015**, *60* (2), 328-337.
- 509 5. (a) Zhou, D.; Conrad, C.; Xia, F.; Park, J.-S.; Payer, B.; Yin, Y.; Lauwers, G. Y.; Thasler, W.;
510 Lee, J. T.; Avruch, J.; Bardeesy, N., Mst1 and Mst2 Maintain Hepatocyte Quiescence and Suppress
511 Hepatocellular Carcinoma Development through Inactivation of the Yap1 Oncogene. *Cancer Cell*
512 **2009**, *16* (5), 425-438; (b) Camargo, F. D.; Gokhale, S.; Johnnidis, J. B.; Fu, D.; Bell, G. W.; Jaenisch,

513 R.; Brummelkamp, T. R., YAP1 Increases Organ Size and Expands Undifferentiated Progenitor Cells.
514 *Current Biology* **2007**, 17 (23), 2054-2060; (c) Liu-Chittenden, Y.; Huang, B.; Shim, J. S.; Chen, Q.;
515 Lee, S.-J.; Anders, R. A.; Liu, J. O.; Pan, D., Genetic and pharmacological disruption of the TEAD–
516 YAP complex suppresses the oncogenic activity of YAP. *Genes & Development* **2012**, 26 (12), 1300–
517 1305.

518 6. (a) Petrilli, A. M.; Fernández-Valle, C., Role of Merlin/NF2 inactivation in tumor biology.
519 *Oncogene* **2016**, 35 (5), 537-548; (b) Zhang, N.; Bai, H.; David, K. K.; Dong, J.; Zheng, Y.; Cai, J.;
520 Giovannini, M.; Liu, P.; Anders, R. A.; Pan, D., The Merlin/NF2 Tumor Suppressor Functions
521 through the YAP Oncoprotein to Regulate Tissue Homeostasis in Mammals. *Developmental Cell*
522 **2010**, 19 (1), 27-38; (c) Plouffe, S. W.; Meng, Z.; Lin, K. C.; Lin, B.; Hong, A. W.; Chun, J. V.; Guan,
523 K.-L., Characterization of Hippo Pathway Components by Gene Inactivation. *Molecular Cell* **2016**,
524 64 (5), 993-1008.

525 7. (a) Bueno, R.; Stawiski, E. W.; Goldstein, L. D.; Durinck, S.; De Rienzo, A.; Modrusan, Z.;
526 Gnad, F.; Nguyen, T. T.; Jaiswal, B. S.; Chirieac, L. R.; Sciaranghella, D.; Dao, N.; Gustafson, C. E.;
527 Munir, K. J.; Hackney, J. A.; Chaudhuri, A.; Gupta, R.; Guillory, J.; Toy, K.; Ha, C.; Chen, Y.-J.;
528 Stinson, J.; Chaudhuri, S.; Zhang, N.; Wu, T. D.; Sugarbaker, D. J.; de Sauvage, F. J.; Richards, W.
529 G.; Seshagiri, S., Comprehensive genomic analysis of malignant pleural mesothelioma identifies
530 recurrent mutations, gene fusions and splicing alterations. *Nature Genetics* **2016**, 48 (4), 407-416; (b)
531 Miyanaga, A.; Masuda, M.; Tsuta, K.; Kawasaki, K.; Nakamura, Y.; Sakuma, T.; Asamura, H.;
532 Gemma, A.; Yamada, T., Hippo Pathway Gene Mutations in Malignant Mesothelioma: Revealed by
533 RNA and Targeted Exon Sequencing. *Journal of Thoracic Oncology* **2015**, 10 (5), 844-851.

534 8. Nguyen, C. D. K.; Yi, C., YAP/TAZ Signaling and Resistance to Cancer Therapy. *Trends in*
535 *Cancer* **2019**, 5 (5), 283-296.

536 9. Tsuji, T.; Ozasa, H.; Aoki, W.; Aburaya, S.; Yamamoto Funazo, T.; Furugaki, K.; Yoshimura,
537 Y.; Yamazoe, M.; Ajimizu, H.; Yasuda, Y.; Nomizo, T.; Yoshida, H.; Sakamori, Y.; Wake, H.; Ueda,
538 M.; Kim, Y. H.; Hirai, T., YAP1 mediates survival of ALK-rearranged lung cancer cells treated with
539 alectinib via pro-apoptotic protein regulation. *Nature Communications* **2020**, 11 (1), 74.

- 540 10. Kurppa, K. J.; Liu, Y.; To, C.; Zhang, T.; Fan, M.; Vajdi, A.; Knelson, E. H.; Xie, Y.; Lim, K.;
541 Cejas, P.; Portell, A.; Lizotte, P. H.; Ficarro, S. B.; Li, S.; Chen, T.; Haikala, H. M.; Wang, H.;
542 Bahcall, M.; Gao, Y.; Shalhout, S.; Boettcher, S.; Shin, B. H.; Thai, T.; Wilkens, M. K.; Tillgren, M.
543 L.; Mushajiang, M.; Xu, M.; Choi, J.; Bertram, A. A.; Ebert, B. L.; Beroukhi, R.; Bandopadhyay,
544 P.; Awad, M. M.; Gokhale, P. C.; Kirschmeier, P. T.; Marto, J. A.; Camargo, F. D.; Haq, R.; Paweletz,
545 C. P.; Wong, K.-K.; Barbie, D. A.; Long, H. W.; Gray, N. S.; Jänne, P. A., Treatment-Induced Tumor
546 Dormancy through YAP-Mediated Transcriptional Reprogramming of the Apoptotic Pathway.
547 *Cancer Cell* **2020**, 37 (1), 104-122.e12.
- 548 11. (a) Santucci, M.; Vignudelli, T.; Ferrari, S.; Mor, M.; Scalvini, L.; Bolognesi, M. L.; Uliassi,
549 E.; Costi, M. P., The Hippo Pathway and YAP/TAZ–TEAD Protein–Protein Interaction as Targets for
550 Regenerative Medicine and Cancer Treatment. *Journal of Medicinal Chemistry* **2015**, 58 (12), 4857-
551 4873; (b) Gibault, F.; Sturbaut, M.; Bailly, F.; Melnyk, P.; Cotellet, P., Targeting Transcriptional
552 Enhanced Associate Domains (TEADs). *Journal of Medicinal Chemistry* **2018**, 61 (12), 5057-5072; (c)
553 Dey, A.; Varelas, X.; Guan, K.-L., Targeting the Hippo pathway in cancer, fibrosis, wound healing
554 and regenerative medicine. *Nature Reviews Drug Discovery* **2020**, 19 (7), 480-494.
- 555 12. Zhang, Z.; Lin, Z.; Zhou, Z.; Shen, H. C.; Yan, S. F.; Mayweg, A. V.; Xu, Z.; Qin, N.; Wong,
556 J. C.; Zhang, Z.; Rong, Y.; Fry, D. C.; Hu, T., Structure-Based Design and Synthesis of Potent Cyclic
557 Peptides Inhibiting the YAP–TEAD Protein–Protein Interaction. *ACS Medicinal Chemistry Letters*
558 **2014**, 5 (9), 993-998.
- 559 13. (a) Noland, Cameron L.; Gierke, S.; Schnier, Paul D.; Murray, J.; Sandoval, Wendy N.;
560 Sagolla, M.; Dey, A.; Hannoush, Rami N.; Fairbrother, Wayne J.; Cunningham, Christian N.,
561 Palmitoylation of TEAD Transcription Factors Is Required for Their Stability and Function in Hippo
562 Pathway Signaling. *Structure* **2016**, 24 (1), 179-186; (b) Chan, P.; Han, X.; Zheng, B.; DeRan, M.; Yu,
563 J.; Jarugumilli, G. K.; Deng, H.; Pan, D.; Luo, X.; Wu, X., Autopalmitoylation of TEAD proteins
564 regulates transcriptional output of the Hippo pathway. *Nature Chemical Biology* **2016**, 12 (4), 282-
565 289.
- 566 14. Pobbati, Ajaybabu V.; Han, X.; Hung, Alvin W.; Weiguang, S.; Huda, N.; Chen, G.-Y.; Kang,
567 C.; Chia, Cheng San B.; Luo, X.; Hong, W.; Poulsen, A., Targeting the Central Pocket in Human

568 Transcription Factor TEAD as a Potential Cancer Therapeutic Strategy. *Structure* **2015**, 23 (11),
569 2076-2086.

570 15. Li, Q.; Sun, Y.; Jarugumilli, G. K.; Liu, S.; Dang, K.; Cotton, J. L.; Xiol, J.; Chan, P. Y.;
571 DeRan, M.; Ma, L.; Li, R.; Zhu, L. J.; Li, J. H.; Leiter, A. B.; Ip, Y. T.; Camargo, F. D.; Luo, X.;
572 Johnson, R. L.; Wu, X.; Mao, J., Lats1/2 Sustain Intestinal Stem Cells and Wnt Activation through
573 TEAD-Dependent and Independent Transcription. *Cell Stem Cell* **2020**, 26 (5), 675-692.e8.

574 16. Holden, J. K.; Crawford, J. J.; Noland, C. L.; Schmidt, S.; Zbieg, J. R.; Lacap, J. A.; Zang, R.;
575 Miller, G. M.; Zhang, Y.; Beroza, P.; Reja, R.; Lee, W.; Tom, J. Y. K.; Fong, R.; Steffek, M.; Clausen,
576 S.; Hagenbeek, T. J.; Hu, T.; Zhou, Z.; Shen, H. C.; Cunningham, C. N., Small Molecule
577 Dysregulation of TEAD Lipidation Induces a Dominant-Negative Inhibition of Hippo Pathway
578 Signaling. *Cell Reports* **2020**, 31 (12), 107809.

579 17. Tang, T. T.; Konradi, A. W.; Feng, Y.; Peng, X.; Ma, M.; Li, J.; Yu, F.-X.; Guan, K.-L.; Post,
580 L., Small Molecule Inhibitors of TEAD Auto-palmitoylation Selectively Inhibit Proliferation and
581 Tumor Growth of NF2 -deficient Mesothelioma. *Molecular Cancer Therapeutics* **2021**, 20
582 (6), 986-998.

583 18. Bum-Erdene, K.; Zhou, D.; Gonzalez-Gutierrez, G.; Ghosayel, M. K.; Si, Y.; Xu, D.;
584 Shannon, H. E.; Bailey, B. J.; Corson, T. W.; Pollok, K. E.; Wells, C. D.; Meroueh, S. O., Small-
585 Molecule Covalent Modification of Conserved Cysteine Leads to Allosteric Inhibition of the
586 TEAD-Yap Protein-Protein Interaction. *Cell Chemical Biology* **2019**, 26 (3), 378-389.e13.

587 19. Lu, W.; Wang, J.; Li, Y.; Tao, H.; Xiong, H.; Lian, F.; Gao, J.; Ma, H.; Lu, T.; Zhang, D.; Ye,
588 X.; Ding, H.; Yue, L.; Zhang, Y.; Tang, H.; Zhang, N.; Yang, Y.; Jiang, H.; Chen, K.; Zhou, B.; Luo,
589 C., Discovery and biological evaluation of vinylsulfonamide derivatives as highly potent, covalent
590 TEAD autopalmitoylation inhibitors. *European Journal of Medicinal Chemistry* **2019**, 184, 111767.

591 20. Kaneda, A.; Seike, T.; Danjo, T.; Nakajima, T.; Otsubo, N.; Yamaguchi, D.; Tsuji, Y.;
592 Hamaguchi, K.; Yasunaga, M.; Nishiya, Y.; Suzuki, M.; Saito, J.-I.; Yatsunami, R.; Nakamura, S.;
593 Sekido, Y.; Mori, K. The novel potent TEAD inhibitor, K-975, inhibits YAP1/TAZ-TEAD protein-
594 protein interactions and exerts an anti-tumor effect on malignant pleural mesothelioma *Am J Cancer*
595 *Res* [Online], 2020, p. 4399-4415. PubMed. <http://europepmc.org/abstract/MED/33415007>

- 596 <https://europepmc.org/articles/PMC7783735>
- 597 <https://europepmc.org/articles/PMC7783735?pdf=render> (accessed 2020).
- 598 21. Zhang, T.; Hatcher, J. M.; Teng, M.; Gray, N. S.; Kostic, M., Recent Advances in Selective
- 599 and Irreversible Covalent Ligand Development and Validation. *Cell Chemical Biology* **2019**, 26 (11),
- 600 1486-1500.
- 601 22. (a) Resnick, E.; Bradley, A.; Gan, J.; Douangamath, A.; Krojer, T.; Sethi, R.; Geurink, P. P.;
- 602 Aimon, A.; Amitai, G.; Bellini, D.; Bennett, J.; Fairhead, M.; Fedorov, O.; Gabizon, R.; Gan, J.; Guo,
- 603 J.; Plotnikov, A.; Reznik, N.; Ruda, G. F.; Díaz-Sáez, L.; Straub, V. M.; Szommer, T.; Velupillai, S.;
- 604 Zaidman, D.; Zhang, Y.; Coker, A. R.; Dowson, C. G.; Barr, H. M.; Wang, C.; Huber, K. V. M.;
- 605 Brennan, P. E.; Ovaa, H.; von Delft, F.; London, N., Rapid Covalent-Probe Discovery by
- 606 Electrophile-Fragment Screening. *Journal of the American Chemical Society* **2019**, 141 (22), 8951-
- 607 8968; (b) Lu, W.; Kostic, M.; Zhang, T.; Che, J.; Patricelli, M. P.; Jones, L. H.; Chouchani, E. T.;
- 608 Gray, N. S., Fragment-based covalent ligand discovery. *RSC Chemical Biology* **2021**, 2, 354-367; (c)
- 609 Dubiella, C.; Pinch, B. J.; Koikawa, K.; Zaidman, D.; Poon, E.; Manz, T. D.; Nabet, B.; He, S.;
- 610 Resnick, E.; Rogel, A.; Langer, E. M.; Daniel, C. J.; Seo, H.-S.; Chen, Y.; Adelmant, G.; Sharifzadeh,
- 611 S.; Ficarro, S. B.; Jamin, Y.; Martins da Costa, B.; Zimmerman, M. W.; Lian, X.; Kibe, S.; Kozono, S.;
- 612 Doctor, Z. M.; Browne, C. M.; Yang, A.; Stoler-Barak, L.; Shah, R. B.; Vangos, N. E.; Geffken, E. A.;
- 613 Oren, R.; Koide, E.; Sidi, S.; Shulman, Z.; Wang, C.; Marto, J. A.; Dhe-Paganon, S.; Look, T.; Zhou,
- 614 X. Z.; Lu, K. P.; Sears, R. C.; Chesler, L.; Gray, N. S.; London, N., Sulfopin is a covalent inhibitor of
- 615 Pin1 that blocks Myc-driven tumors in vivo. *Nature Chemical Biology* **2021**, 17 (9), 954-963.
- 616 23. Yu, C.; Mannan, A. M.; Yvone, G. M.; Ross, K. N.; Zhang, Y.-L.; Marton, M. A.; Taylor, B.
- 617 R.; Crenshaw, A.; Gould, J. Z.; Tamayo, P.; Weir, B. A.; Tsherniak, A.; Wong, B.; Garraway, L. A.;
- 618 Shamji, A. F.; Palmer, M. A.; Foley, M. A.; Winckler, W.; Schreiber, S. L.; Kung, A. L.; Golub, T. R.,
- 619 High-throughput identification of genotype-specific cancer vulnerabilities in mixtures of barcoded
- 620 tumor cell lines. *Nature Biotechnology* **2016**, 34 (4), 419-423.
- 621 24. Ábrányi-Balogh, P.; Petri, L.; Imre, T.; Szijj, P.; Scarpino, A.; Hrast, M.; Mitrović, A.;
- 622 Fonovič, U. P.; Németh, K.; Barreteau, H.; Roper, D. I.; Horváti, K.; Ferenczy, G. G.; Kos, J.; Ilaš, J.;

623 Gobec, S.; Keserü, G. M., A road map for prioritizing warheads for cysteine targeting covalent
624 inhibitors. *European Journal of Medicinal Chemistry* **2018**, *160*, 94-107.

625 25. Pobbati, A. V.; Rubin, B. P., Protein-Protein Interaction Disruptors of the YAP/TAZ-TEAD
626 Transcriptional Complex. *Molecules* **2020**, *25* (24), 6001.

627 26. Kuljanin, M.; Mitchell, D. C.; Schweppe, D. K.; Gikandi, A. S.; Nusinow, D. P.; Bulloch, N.
628 J.; Vinogradova, E. V.; Wilson, D. L.; Kool, E. T.; Mancias, J. D.; Cravatt, B. F.; Gygi, S. P.,
629 Reimagining high-throughput profiling of reactive cysteines for cell-based screening of large
630 electrophile libraries. *Nature Biotechnology* **2021**, *39*, 630-641.

631 27. Kim, M.; Kim, T.; Johnson, Randy L.; Lim, D.-S., Transcriptional Co-repressor Function of
632 the Hippo Pathway Transducers YAP and TAZ. *Cell Reports* **2015**, *11* (2), 270-282.

633 28. Zhou, Y.; Zhou, B.; Pache, L.; Chang, M.; Khodabakhshi, A. H.; Tanaseichuk, O.; Benner, C.;
634 Chanda, S. K., Metascape provides a biologist-oriented resource for the analysis of systems-level
635 datasets. *Nature Communications* **2019**, *10* (1), 1523.

636 29. Karaman, R.; Halder, G., Cell Junctions in Hippo Signaling. *Cold Spring Harbor*
637 *Perspectives in Biology* **2018**, *10*:a028753.

638 30. Li, N.; Lu, N.; Xie, C., The Hippo and Wnt signalling pathways: crosstalk during neoplastic
639 progression in gastrointestinal tissue. *The FEBS Journal* **2019**, *286* (19), 3745-3756.

640 31. Mizuno, T.; Murakami, H.; Fujii, M.; Ishiguro, F.; Tanaka, I.; Kondo, Y.; Akatsuka, S.;
641 Toyokuni, S.; Yokoi, K.; Osada, H.; Sekido, Y., YAP induces malignant mesothelioma cell
642 proliferation by upregulating transcription of cell cycle-promoting genes. *Oncogene* **2012**, *31* (49),
643 5117-5122.

644 32. Liu, C.-Y.; Lv, X.; Li, T.; Xu, Y.; Zhou, X.; Zhao, S.; Xiong, Y.; Lei, Q.-Y.; Guan, K.-L.,
645 PP1 Cooperates with ASPP2 to Dephosphorylate and Activate TAZ*. *Journal of Biological*
646 *Chemistry* **2011**, *286* (7), 5558-5566.

647 33. (a) Cai, D.; Feliciano, D.; Dong, P.; Flores, E.; Gruebele, M.; Porat-Shliom, N.; Sukenik, S.;
648 Liu, Z.; Lippincott-Schwartz, J., Phase separation of YAP reorganizes genome topology for long-term
649 YAP target gene expression. *Nature cell biology* **2019**, *21* (12), 1578-1589; (b) Lu, Y.; Wu, T.;

Gutman, O.; Lu, H.; Zhou, Q.; Henis, Y. I.; Luo, K., Phase separation of TAZ compartmentalizes the transcription machinery to promote gene expression. *Nature Cell Biology* **2020**, 22 (4), 453-464.

EXPERIMENTAL DETAILS, METHODS AND CHARACTERIZATION DATA

1.1 Cloning

The stretch of residues 209-424 of human TEAD1, residues 220-450 of human TEAD2, residues 119-436 of human TEAD3, residues 216-434 of human TEAD4, were inserted into the pET28PP (N-terminal His 3C tag) vector.

1.2 Protein expression and purification

The N-terminal His tag construct of human TEAD1 (residues 209-424) was overexpressed in *E. coli* BL21 (DE3) and purified using affinity chromatography and size-exclusion chromatography. Briefly, cells were grown at 37°C in TB medium in the presence of 50 µg/ml of kanamycin to an OD of 0.8, cooled to 17°C, induced with 500 µM isopropyl-1-thio-D-galactopyranoside (IPTG), incubated overnight at 17°C, collected by centrifugation, and stored at -80°C. Cell pellets were lysed in buffer A (25 mM HEPES, pH 7.5, 200 mM NaCl, 5% glycerol, 7 mM mercapto-ethanol, and 20 mM Imidazole) using Microfluidizer (Microfluidics), and the resulting lysate was centrifuged at 30,000g for 40 min. Ni-NTA beads (Qiagen) were mixed with cleared lysate for 30 min and washed with buffer A. Beads were transferred to an FPLC-compatible column, and the bound protein was washed further with buffer A for 10 column volumes and eluted with buffer B (25 mM HEPES, pH 7.5, 200 mM NaCl, 5% glycerol, 7 mM mercapto-ethanol, and 400 mM Imidazole). The eluted sample was concentrated and purified further using a Superdex 200 16/600 column (Cytiva) in buffer C containing 20 mM HEPES, pH 7.5, 200 mM NaCl, 5% glycerol, 0.5 mM TCEP and 2 mM DTT. HRV 3C protease was added to TEAD1 containing fractions and incubated overnight at 4°C, followed by passing through Ni-NTA column to remove His-tag and 3C protease. The flow-through fractions of the second Ni-NTA column, containing cleaved TEAD1, was concentrated to ~9mg/mL and stored in -80°C. The N-terminal His

tag construct of human TEAD2 (residues 220-450), TEAD3 (residues 119-436) and TEAD4 (residues 216-434) were purified as TEAD1.

2 Mass spectrometry analysis

TEAD2 protein (5 µg) was treated with DMSO or a 10-fold molar excess of MYF-03-69 and analyzed by LC-MS using an HPLC system (Shimadzu, Marlborough, MA) interfaced to an LTQ ion trap mass spectrometer (ThermoFisher Scientific, San Jose, CA). Proteins were desalted for 4 minutes on column with 100% A, and eluted with an HPLC gradient (0-100% B in 1 minute; A=0.2M acetic acid in water; B=0.2M acetic acid in acetonitrile). The mass spectrometer was programmed to acquire full scan mass spectra (m/z 300-2000) in profile mode (spray voltage = 4.5 kV). Mass spectra were deconvoluted using MagTran software version 1.03b2¹. To analyze the site of modification, DMSO or MYF-03-69 treated proteins were first captured on SP3 beads² by adding an equal volume of acetonitrile and washed (2x with 70% acetonitrile, 1x with 100% acetonitrile). Beads were then resuspended with 100 mM ammonium bicarbonate containing 0.1% Rapigest (Waters, Milford, MA). Proteins were reduced with 10 mM DTT for 30 minutes at 56°C, alkylated with 22.5 mM iodoacetamide for 30 minutes at room temperature, and then digested overnight with trypsin at 37 °C. Rapigest was cleaved according to the manufacturer's instructions, and peptides were desalted by C18 and dried by vacuum centrifugation. Peptides were reconstituted in 50% acetonitrile, 1% formic acid, 100 mM ammonium acetate, and analyzed by CE-MS using a ZipChip CE-MS instrument and autosampler (908 devices, Boston, MA) interfaced to a QExactive HF mass spectrometer (ThermoFisher Scientific). Peptides were resolved at 500V/cm using an HR chip with a background electrolyte consisting of 50% acetonitrile with 1% formic acid. The mass spectrometer was operated in data dependent mode, and subjected the 5 most abundant ions in each MS scan (m/z 300-2000, 60K resolution, 3E6 target, 100 ms max fill time) to MS/MS (15K resolution, 1E5 target, 100 ms max fill time). Dynamic exclusion was enabled with a repeat count of 1 and an exclusion duration of 5 seconds. Raw mass spectrometry data files were converted to .mgf using multiplier³ and searched against a forward-reverse human refseq database using Mascot version 2.6.2. Search parameters specified fixed carbamidomethylation of cysteine, variable methionine oxidation, and

variable MYF-03-69 modification of cysteine. MYF-03-69 modified spectra were examined and figures prepared using mzStudio software⁴. Inhibitor related ions in MS/MS spectra were identified as described⁵.

3 Docking

Docking of MYF-01-37 was performed with covalent docking protocol from Schrodinger suite software (release 2019-02) with default parameters in TEAD2 structure (PDB code: 5HGU) exporting 5 poses per molecule. Both stereoisomers were docked. Top scoring pose was chosen to illustrate the binding pose. The pdb structure was processed and optimized with protein preparation protocol with default setting in Schrodinger suite.

4.1 Crystallization

Using Formulatrix NT8 and RockImager and ArtRobbins Phoenix liquid handlers, a 100 nL sample of 300 μ M TEAD1 that was preincubated for 1 hour with 600 μ M MYF-03-69 was dispensed in an equal volume of crystallization buffer (3M NH₄SO₄ and 0.1 M Tris pH 9.0) and incubated against 25 μ L of crystallization buffer in a 384-well hanging-drop vapor diffusion microtiter plate at 20 °C for three days.

4.2 Data collection and structure determination

Diffraction data were collected at beamline 24ID-E of the NE-CAT at the Advanced Photon Source (Argonne National Laboratory). Data sets were integrated and scaled using XDS⁶. Structures were solved by molecular replacement using the program Phaser⁷ and available search models from the PDB. Iterative manual model building and refinement using Phenix⁸ and Coot⁹ led to a model with excellent statistics.

5 Gel-based palmitoylation studies

1 μ M TEADs-YBD recombinant protein was incubated with inhibitors at the indicated concentrations at 37 °C for 2 h followed by the addition of palmitoyl alkyne-coenzyme A (Cayman chemical, no. 15968) in a total volume of 50 μ L. After 30 min reaction, 5 μ L 10%SDS were added and 5 μ L click reagents were added to start click reaction as previously reported¹⁰. After another 1 h, 4x loading

buffer were added to the reaction mixture and the samples subjected for western blot analysis. IRDye 800CW Streptavidin (LI-COR, no. 92632230) and His-Tag Mouse mAb (Cell Signaling, no. 2366S) was used for biotin detection and His-tag detection. The blots were imaged on Odyssey CLx Imager (LI-COR).

6.1 Sample preparation for SLC-ABPP

Samples for whole cysteine profiling were prepared as previously described¹¹. Briefly, frozen cell pellets from H226 cells were lysed using PBS (pH 7.4). Samples were further homogenized, and DNA was sheared using sonication with a probe sonicator (20 x 0.5 s pulses) at 4°C. Total protein was determined using a BCA assay and cell lysates were used immediately for each experiment. Depending on the experiment, 50 µg of total cell extract was aliquoted for each TMT channel for further downstream processing. Excess DBIA, along with disulfide bonds were quenched and reduced using 5 mM dithiothreitol (DTT) for 30 min in the dark at room temperature. Subsequently reduced cysteine residues were alkylated using 20 mM iodoacetamide for 45 min in the dark at room temperature. To facilitate removal of quenched DBIA and incompatible reagents, proteins were precipitated using chloroform/methanol. Briefly, to 100 µL of each sample, 400 µL of methanol were added, followed by 100 µL of chloroform with thorough vortexing. Next, 300 µL of HPLC grade water were added, and the samples were mixed to facilitate precipitation. Samples were centrifuged at maximum speed (14,000 rpm) for 3 min at room temperature, the aqueous top layer was removed, and the samples were washed additionally three times with 500 µL of methanol. Protein pellets were resolubilized in 200 mM 4-(2-hydroxyethyl)-1-piperazinepropanesulfonic acid (EPPS) at pH 8.5 and digested using LysC and trypsin (1:100, enzyme to protein ratios) overnight at 37°C using a ThermoMixer set to 1200 rpm. The next day samples were labeled with TMT reagents or stored at -80°C until further use.

6.2 TMT Labeling for SLC-ABPP

Digested peptides containing DBIA conjugated-cysteines were labeled using TMTPro 16-plex reagents as previously described¹². Briefly, peptides were labeled at a 1:2 ratio by mass (peptides to TMT reagents) for 1 hr with shaking at 1,200 rpm. To equalize protein loading, ~ 2 µg of each sample

were aliquoted, and a 60-min quality control analysis (ratio check) was performed using SPS-MS3. Excess TMT reagent was quenched with hydroxylamine (0.3% final concentration) for 15 min at room temperature. Next, samples were mixed 1:1 across all TMT channels and the pooled sample was dried using a Speedvac to ensure all acetonitrile was removed.

6.3 Cysteine peptide enrichment using streptavidin in SLC-ABPP

Pierce streptavidin magnetic beads were washed with PBS pH 7.4 prior to use. To each TMT labeled pooled sample, 100 µL of a 50% slurry of streptavidin beads were added. Samples were further topped up with 1 mL of PBS in a 2 mL Eppendorf tube. Samples and beads were incubated overnight at 4 °C to enrich for TMT-labeled DBIA conjugated cysteine peptides. Following enrichment, the beads were placed onto a magnetic rack and allowed to equilibrate for 5 min. Beads were washed to remove non-specific binding using the following procedure: 3 x 1 mL of PBS pH 7.4, 1 x 1 mL of PBS with 0.1% SDS pH 7.4 and finally 3 x 1 mL HPLC grade water. Beads were resuspended using a pipette between washes and placed on the magnet between each wash. To elute cysteine containing peptides, 500 µL of 50% acetonitrile with 0.1% trifluoroacetic acid (TFA) were added, and the beads were mixed at 1,000 rpm for 15 min at room temperature. Eluted peptides were transferred to a new tube, and the beads were additionally washed with 200 µL of 50% acetonitrile with 0.1% TFA and combined. Cysteine containing peptides were dried to completion using a Speedvac and were stored at -80°C.

6.4 Desalting cysteine-containing peptides in SLC-ABPP

TMT-labeled cysteine-containing peptides were resuspended using 200 µL of 1% formic acid (FA) and were desalted using StageTips as previously described^{12a}. Briefly, eight 18-gauge cores were packed into a 200 µL pipette tip and were passivated and equilibrated using the following solutions: 100 µL of 100% methanol, 70% acetonitrile 1% FA, and 5% acetonitrile 5% FA. Peptides were loaded and were washed with 0.1% FA and eluted using 150 µL of 70% acetonitrile 1% FA and dried to completion using a Speedvac. Enriched peptides samples were next resuspended in 5-10 µL of 5% acetonitrile 5% FA and between 50 and 100% of the sample was injected for analysis using LC-RTS-SPS-MS3.

6.5 Mass spectrometry and real-time searching in SLC-ABPP

All SLC-ABPP mass spectrometry data were acquired using an Orbitrap Fusion Eclipse mass spectrometer in-line with a Proxeon NanoLC-1200 UPLC system. Peptides were separated using an in-house 100 μ m capillary column packed with 35 cm of Accucore 150 resin (2.6 μ m, 150 Å) (ThermoFisher Scientific) using 210 min gradients from 4 to 24% acetonitrile in 0.125% formic acid per run. Eluted peptides were quantified using the synchronous precursor selection (SPS-MS3) method for TMT quantification. Briefly, MS1 spectra were acquired at 120K resolving power with a maximum of 50 ms ion injection in the Orbitrap. MS2 spectra were acquired by selecting the top 10 most abundant features via collisional induced dissociation (CID) in the ion trap using an automatic gain control (AGC) setting of 15K, quadrupole isolation width of 0.5 m/z and a maximum ion accumulation time of 50 ms. These spectra were passed in real time to the external computer for online database searching. Intelligent data acquisition (IDA) using real-time searching (RTS) was performed using Obiter¹³. Cysteine-containing peptides or whole proteome peptide spectral matches were analyzed using the Comet search algorithm (release_2019010) designed for spectral acquisition speed¹⁴. The same forward- and reversed-sequence human protein databases were used for both the RTS search and the final search (Uniprot). The RTS Comet functionality has been released and is available here: <http://cometms.sourceforge.net/>. Real-time access to spectral data was enabled by the Thermo Scientific Fusion API (<https://github.com/thermofisherlms/iapi>). Next, peptides were filtered using simple initial filters that included the following: not a match to a reversed sequence, maximum PPM error of <50, minimum XCorr of 0.5, minimum deltaCorr of 0.10 and minimum peptide length of 7. If peptide spectra matched to above criteria, an SPS-MS3 scan was performed using up to 10 *b*- and *y*-type fragment ions as precursors with an AGC of 200K for a maximum of 200 ms with a normalized collision energy setting of 55 (TMTPro 16-plex)^{12b}.

6.6 Mass spectrometry data analysis in SLC-ABPP

All acquired data were searched using the open-source Comet algorithm (release_2019010) using a previously described informatics pipeline¹⁵. Spectral searches were done using a custom FASTA-formatted database which included common contaminants, reversed sequences (Uniprot Human, 2014)

and the following parameters: 50 PPM precursor tolerance, fully tryptic peptides, fragment ion tolerance of 0.9 Da and a static modification by TMTPro (+304.2071 Da) on lysine and peptide N termini. Carbamidomethylation of cysteine residues (+57.021 Da) was set as a static modification while oxidation of methionine residues (+15.995 Da) and DBIA on cysteine residues (+239.262) was set as a variable modification. Peptide spectral matches were filtered to a peptide false discovery rate (FDR) of less than 1% using linear discriminant analysis employing a target-decoy strategy. Resulting peptides were further filtered to obtain a 1% protein FDR at the entire dataset level (including all plexes per cell line), and proteins were collapsed into groups. Cysteine-modified peptides were further filtered for site localization using the AScore algorithm with a cutoff of 13 ($p < 0.05$) as previously described^{15b}. Overlapping peptide sequences that were generated from different charge states, elution times and tryptic termini were grouped together into a single entry. A single quantitative value was reported, and only unique peptides were reported. Reporter ion intensities were adjusted to correct for impurities during synthesis of different TMT reagents according to the manufacturer's specifications. For quantification of each MS3 spectrum, a total sum signal-to-noise of all reporter ions of 160 (TMTPro) was required. Lastly, peptide quantitative values were normalized so that the sum of the signal for all proteins in each channel was equal to account for sample loading differences (column normalization).

6.7 SLC-ABPP competition ratio calculation

Cysteine site-specific engagement was assessed by the blockage of DBIA probe labeling. Peptides that showed >95% reduction in TMT intensities in the electrophile-treated samples were assigned a maximum ratio of 20 for graphing purposes with preserved ranking. TMT reporter ion sum-signal-to-noise for each SLC-ABPP experiment was used to calculate the competition ratios by dividing the control channel (DMSO) by the electrophile treated channel. Replicate measurements were averaged and reported as a single entry. To avoid false positives, sites with large coefficients of variation had the highest replicate CR values removed before averaging as previously described^{11, 16}.

7.1 Cell culture

HEK293T (ATCC, no. CRL-3216), NCI-H226 (ATCC, no. CRL-5826), MSTO-211H (ATCC, no. CRL-2081), NCI-H2452 (ATCC, no. CRL-5946) mesothelioma cells, MeT-5A (ATCC, no. CRL-9444) mesothelium cells, 94T778 cells (ATCC, no. CRL-3044), 93T449 cells (ATCC, no. 3043), MIA PaCa-2 cells (ATCC, no. CRM-CRL-1420), MM.1S (ATCC, no. CRL-2974) and SKHEP-1 cells (ATCC, no. HTB-52) were obtained from American Type Culture Collection and cultured as recommended. HuCCT1 cells were gifts from Bardeesy lab in Massachusetts General Hospital. PC9 cells were gifts from Pasi lab in Dana-Farber Cancer Institute. Cells were negative for mycoplasma using MycoAlert mycoplasma detection kit (LONZA, no. LT07-418).

7.2 Cell proliferation assay

For 2D adherent cell viability experiment, the cells were seeded at 384-well plate (Corning, no. 3570) at the density of 200 cells/well. The next day, compounds were added using Janus workstation (PerkinElmer). After 5 days treatment, the cell viability was measured by CellTiter-Glo kit (Promega, no. G7570) as the manufacturer recommended. For 3D spheroid assays, NCI-H226 and MSTO-211H cells were plated at the density of 200 cells/well in Ultra-low Attachment (ULA) plate (S-bio, no. MS-9384WZ) without or with 5% Matrigel matrix (Corning, no. 356231) respectively. The cell viability was measured using 3D CellTiter-Glo kit (Promega, no. G9681). The luminescent signal was collected on EnVision plate reader (PerkinElmer). The GR₅₀ values were calculated as previously described¹⁷.

7.3 Cell cycle analysis

Cells were plated in 6-well plate (Corning, no. 3506) and treated by MYF-03-69 at indicated concentrations. Then cells were harvested and fixed in cold 70% ethanol overnight. Next day, the samples were treated with 100 µg/mL RNAase A (Life Technologies, no. EN0531) and stained by 50 µg/mL propidium iodide solution (Life Technologies, no. P3566). After incubation at room temperature for 30 min, the samples were subjected to assessment using Guava flow cytometer. The data were further analyzed in Flowjo software.

8 PRISM screening and data analysis

Up to 931 barcoded cell lines in pools of 20-25 were thawed and plated into 384-well plates (1250 cells/well for adherent cell pools, 2000 cells/well for suspension or mixed suspension/adherent cell pools) containing compound (top concentration: 10 μ M, 8-point, threefold dilution). All conditions were tested in triplicate. Cells were lysed after 5 days of treatment and mRNA based Luminex detection of barcode abundance from lysates was carried out as described previously¹⁸. Luminex median fluorescence intensity (MFI) data was input to a standardized R pipeline (https://github.com/broadinstitute/prism_data_processing) to generate viability estimates relative to vehicle treatment for each cell line and treatment condition, and to fit dose-response curves from viability data. Correlation analysis was also performed in the R pipeline mentioned above.

9 TEAD reporter assays

The TEAD transcriptional vector (TBS-mCherry) was used to produce lentivirus as previously reported in HEK293T cells¹⁹. The virus was collected at 48 h and 72 h post transfection and concentrated using Lenti-X concentrator (Takara Bio, no. 631231). Then NCI-H226 cells were transduced using the concentrated virus in the presence of 8 μ g/mL polybrene (MilliporeSigma, no. TR1003G). The positively transduced cells were further sorted by GFP. For reporter assays, the selected cells were plated in black 384-well plate (Corning, no. 4514) at the density of 1,000 cells/well. The next day, compounds were added at indicated concentrations using Janus workstation. After 72 h incubation, the cells were stained by Hoechst 33342 (Life Technologies, no. 62249). The mCherry and Hoechst signals were read using Acumen high content imager (TTP Labtech).

For the luciferase reporter assay, NCI-H226 cells were transduced with 250 μ L TEAD luciferase reporter lentivirus (BPS Biosciences, no. 79833) in the presence of 8 μ g/mL polybrene (MilliporeSigma, no. TR1003G). Cells were then selected by 2 μ g/mL puromycin. The selected cells were seeded at the density of 1,000 cells/well and treated with compounds at indicated concentrations. After three days treatment, ONE-Glo™ luciferase assay kit (Promega, no. E6110) and CellTiter-Glo kit (Promega, no. G9681) were used according to manufacturer instructions. The TEAD transcriptional activity was then calculated by normalizing luciferase signal normalized with relative cell viability.

10 Lipid displacement assay

The HEK-293T cells were plated in six-well plate (Corning, no. 3506) and transfected with Myc-TEAD4 vector the next day. The Myc-TEAD4 was a gift from Kunliang Guan²⁰ (Addgene plasmid # 24638; <http://n2t.net/addgene:24638>; RRID: Addgene_24638). Then transfected cells were treated with 50 μ M alkyne palmitic acid (Click Chemistry Tools, cat. no. 1165-5) in the presence of TEAD inhibitors at indicated concentrations next day. After 24-hour incubations, cells were collected and lysed in RIPA buffer with proteasome inhibitor. The cleared supernatant was then subjected to click reactions using same procedures in the gel-based palmitoylation assay. Palmitoylation levels were detected by immunoblot using streptavidin antibody (LI-COR, no. 92632230).

11 RT-PCR studies

The cells were plated in six-well plate (Corning, no. 3506) and treated with compounds the next day. The total RNA was extracted using RNeasy Plus Mini Kit (Qiagen, no. 74134). Then 500 ng purified RNA was used to synthesize cDNA by SuperScript III First-Strand Synthesis System Kit (Life Technologies, no. 18080051). The following TaqMan probes were used for follow-up RT-PCR reactions: CTGF (Hs00170014_m1), CYR61 (Hs00155479_m1), GAPDH (Hs02786624_g1), BMF (Hs00372937_m1), IGFBP3 (Hs00181211_m1), KRT34 (Hs02569742_s1) and NPPB (Hs00173590_m1).

12 Co-immunoprecipitation (Co-IP) studies

The cells were plated in 10 cm dish (Corning, no. 430293) and treated with compounds at indicated concentrations the next day. The Co-IP experiments were conducted using Dynabeads Co-Immunoprecipitation Kit (Life Technologies, no. 14321D) according to the manufacturer instructions.

13.1 RNA-Seq study: Sample Treatment

The NCI-H226 cells were plated in the 10 cm dish and treated with MYF-03-69 at indicated concentrations with biological triplicates for 6 hours next day. The RNA was extracted using RNeasy plus mini kit (Qiagen, cat no.74134) according to the manufacturer instructions.

13.2 RNA-Seq study: Library Preparation and Sequencing

Libraries were prepared using Roche Kapa mRNA HyperPrep strand specific sample preparation kits from 200 ng of purified total RNA according to the manufacturer's protocol on a Beckman Coulter Biomek i7. The finished dsDNA libraries were quantified by Qubit fluorometer and Agilent TapeStation 4200. Uniquely dual indexed libraries were pooled in an equimolar ratio and shallowly sequenced on an Illumina MiSeq to further evaluate library quality and pool balance. The final pool was sequenced on an Illumina NovaSeq 6000 targeting 40 million 100bp read pairs per library at the Dana-Farber Cancer Institute Molecular Biology Core Facilities.

13.3 RNA-Seq study: Analysis

Sequenced reads were aligned to the UCSC hg19 reference genome assembly and gene counts were quantified using STAR (v2.7.3a)²¹. Differential gene expression testing was performed by DESeq2 (v1.22.1)²². RNAseq analysis was performed using the VIPER snakemake pipeline²³. KEGG pathway enrichment analysis was performed through metascape webportal²⁴.

13.4 RNA-Seq study: Datasets availability

These RNA-seq datasets were deposited to BioSample database under accession# as below:
SAMN19288936, SAMN19288937, SAMN19288938, SAMN19288939, SAMN19288940,
SAMN19288941, SAMN19288942, SAMN19288943, SAMN19288944, SAMN19288945,
SAMN19288946.

14.1 Efficacy study in NCI-H226 CDX model: Animals

Female 7-week-old NSG mice were purchased from The Jackson Laboratory (Bar Harbor, ME). Animals acclimated for at least 5 days before initiation of the study. All *in vivo* studies were conducted at Dana-Farber Cancer Institute with the approval of the Institutional Animal Care and Use Committee in an AAALAC accredited vivarium.


14.2 Efficacy study in NCI-H226 CDX model: *In vivo* studies

The NCI-H226 cells were grown in RPMI1640 media supplemented with 10% fetal bovine serum. Cells were harvested, and 5×10^6 cells with 50% Matrigel (Fisher Scientific) were implanted subcutaneously in the right flank of the NSG mice. Tumors were allowed to establish to an average of $141.3 \pm 24.9 \text{ mm}^3$ in size before randomization using Studylog software (San Francisco, CA) into

various treatment groups with 8-9 mice per group. MYF-03-176 was formulated as a suspension in 10% DMSO with 10% Tween 80 in water and dosed twice daily via oral gavage. Control treated mice received vehicle alone. Tumor volumes were determined from caliper measurements by using the formula, Tumor volume = (length × width²)/2. Tumor volumes and body weights were measured twice weekly. Mice were treated for 28 days, followed by measuring for re-growth of tumors.

15 Chemical synthesis and characterization

MYF-03-42



Step 1: Synthesis of *trans*-tert-butyl 3-hydroxy-4-(1H-pyrazol-1-yl)pyrrolidine-1-carboxylate (Compound 3)

To a suspension of 60% NaH (323mg, 8.1 mmol) in THF (20 mL) was added the solution of 1H-pyrazole (551 mg, 8.1 mmol) in THF (5 mL), the mixture was stirred at 0 °C under N₂ for 30 minutes, and then tert-butyl 6-oxa-3-azabicyclo[3.1.0] hexane-3-carboxylate (1.0 g, 5.4 mmol) was added. The resulting mixture was heated at 60°C under N₂ for 16 hours. After cooled down to room temperature the mixture was diluted with ethyl acetate (50 mL) and washed with water (50 mL). The organic layer was dried over anhydrous Na₂SO₄, filtered, concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 40% v/v) to afford compound **3** as solid (600 mg, yield 44%). LC-MS (ESI) m/z: 254 [M+H]⁺.

Step 2: Synthesis of *trans*-tert-butyl 3-(1H-pyrazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 5)

A mixture of compound **3** (300 mg, 1.18 mmol), 1-(bromomethyl)-4-(trifluoromethyl)benzene (283 mg, 1.18 mmol) and 60% NaH (47 mg, 1.18 mmol) in DMF (10 mL) was stirred at room temperature under N₂ for 2 hours. The mixture was diluted with water (50 mL) and extracted with ethyl acetate (50 mL x 2), the combined organic was washed with brine (50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to obtain compound **5** as yellow oil (400 mg, yield 82%). LC-MS (ESI) m/z: 356 [M-56+H]⁺.

Step 3: Synthesis of 1-(*trans*-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-3-yl)-1H-pyrazole (Compound 6)

A mixture of compound **5** (300 mg, 0.72 mmol) and TFA (3 mL) in DCM (10 mL) was stirred at room

temperature for 2 hours. The mixture was concentrated to leave crude compound **6** (250 mg) as yellow oil, which was used directly in the next step. LC-MS (ESI) m/z : 312[M+H]⁺.

Step 4: Synthesis of 1-(*trans*-3-(1H-pyrazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-1-yl)prop-2-en-1-one (MYF-03-42)

A mixture of compound **6** (70 mg, 0.22 mmol), acryloyl chloride (20 mg, 0.22 mmol), and TEA (44 mg, 0.44 mmol) in DCM (10 mL) was stirred at room temperature under N₂ for 2 hours. The mixture was concentrated, the residue was purified by prep-HPLC to afford **MYF-03-42** as yellow oil (50 mg, yield 62%). LC-MS (ESI) m/z : 366[M+H]⁺.


¹H-NMR (400 MHz, CD₃OD): δ (ppm) 7.74 (dd, J = 5.2, 2.4 Hz, 1H), 7.63 (d, J = 8.0 Hz, 2H), 7.56 (d, J = 1.6 Hz, 1H), 7.47 (d, J = 8.1 Hz, 2H), 6.66-6.59 (dd, J = 16.8, 10.4 Hz, 1H), 6.36-6.30 (m, 2H), 5.85 – 5.74 (m, 1H), 5.18 – 5.02 (m, 1H), 4.68 (d, J = 5.8 Hz, 2H), 4.51-4.42 (m, 1H), 4.28-4.08 (m, 2H), 4.01-3.93 (m, 1H), 3.83-3.65 (m, 1H).

MYF-02-111

MYF-02-111 was synthesized through the same route as **MYF-03-42** except 1-(bromomethyl)-3-(trifluoromethyl)benzene was used in **Step 2** instead of 1-(bromomethyl)-4-(trifluoromethyl)benzene.

MYF-02-111 was obtained as light yellow oil. LC-MS (ESI) m/z : 366[M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ 7.87 (ddd, J = 9.3, 2.3, 0.7 Hz, 1H), 7.69 – 7.54 (m, 4H), 7.53 – 7.48 (m, 1H), 6.61 (dd, J = 16.8, 10.3 Hz, 1H), 6.30 (dt, J = 3.4, 2.1 Hz, 1H), 6.17 (dt, J = 16.7, 2.5 Hz, 1H), 5.70 (td, J = 9.9, 2.3 Hz, 1H), 5.13 (ddt, J = 33.6, 7.8, 3.4 Hz, 1H), 4.66 (d, J = 4.0 Hz, 2H), 4.39 (ddd, J = 25.5, 4.1, 1.7 Hz, 1H), 4.19 – 4.09 (m, 0.5H), 4.07 – 3.87 (m, 1.5H), 3.86 – 3.70 (m, 1.5H), 3.52 (dd, J = 13.0, 4.0 Hz, 0.5H).

MYF-03-69



Step 1: Synthesis of (3*R*, 4*R*)-tert-butyl 3-azido-4-hydroxypyrrolidine-1-carboxylate (Compound 2)

A mixture of tert-butyl 6-oxa-3-azabicyclo[3.1.0]hexane-3-carboxylate (4 g, 21.6 mmol), TMSN₃ (2.664 g, 23.2 mmol) and chiral catalyst (1*S*,2*S*)-(-)-[1,2-cyclohexanediamino-N,N'-bis(3,5-di-*t*-

butylsalicylidene)]chromium(III) chloride (328 mg, 0.42 mmol) was stirred at rt under N₂ overnight. The reaction mixture was treated with MeOH (60 mL) and K₂CO₃ (1.788 g, 12.8 mmol) and continued to stir at rt for 5 hours. The reaction mixture was diluted with ethyl acetate (300 mL), washed with water (300 mL x 2), dried over anhydrous Na₂SO₄, concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 30% v/v) to obtain the compound **2** as clear oil (3.5g, 96% e.e., yield 71%). LC-MS (ESI) m/z: 129 [M+H-100]⁺.

Step 2: Synthesis of (3R, 4R)-tert-butyl 3-azido-4-(4-(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 4)

A mixture of compound **2** (3 g, 13.1 mmol), 1-(bromomethyl)-4-(trifluoromethyl)benzene (3.1 g, 13.1 mmol) and 60% NaH (0.6 g, 15.7 mmol) in DMF (20 mL) was stirred at 0 °C under N₂ for 6 hours. The reaction mixture was diluted with water (200 mL) and extracted with ethyl acetate (200 mL), the organic was washed with water (100 mL), dried over anhydrous Na₂SO₄, concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 20% v/v) to obtain compound **4** as oil (3.8g, yield 75%). LC-MS (ESI) m/z: 287[M+H-100]⁺.

Step 3: Synthesis of (3R, 4R)-tert-butyl 3-(4-(pyridin-3-yl)-1H-1,2,3-triazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 6)

A mixture of compound **4** (3.8 g, 9.8 mmol), 3-ethynylpyridine (1.216 g, 11.8 mmol), CuSO₄ (244 mg, 0.98 mmol) and, sodium L-ascorbate (388 mg, 1.96 mmol) in THF (20 mL), H₂O (20 mL) and ⁿBuOH (20 mL) was stirred at 70 °C under N₂ overnight. The reaction mixture was concentrated in vacuum, the residue was diluted with ethyl acetate (600 mL), washed with water (400 mL), dried over anhydrous Na₂SO₄, concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 90% v/v) to obtain compound **6** as yellow solid (4g, yield 83%). LC-MS (ESI) m/z: 490[M+H]⁺.

Step 4: Synthesis of 3-(1-((3R, 4R)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-3-yl)-1H-1,2,3-triazol-4-yl)pyridine (Compound 7)


A mixture of compound **6** (4 g, 8.1 mmol) and TFA (5 mL) in DCM (20 mL) was stirred at 0 °C under N₂ for 2 hours. The mixture was concentrated to leave the crude compound **7** (4 g, crude) as yellow oil, which was used directly for next step. LC-MS (ESI) m/z: 390[M+H]⁺.

Step 5: Synthesis of 1-((3R, 4R)-3-(4-(pyridin-3-yl)-1H-1,2,3-triazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-1-yl)prop-2-en-1-one (MYF-03-69)

A mixture of compound **7** (500 mg, 1.28 mmol), acryloyl chloride (115 mg, 1.28 mmol) and TEA (388 mg, 3.84 mmol) in DCM (20 mL) was stirred at rt for 2 hours. The mixture was concentrated and purified by prep-HPLC to obtain **MYF-03-69** as light yellow solid (300 mg, yield 52%). LC-MS

(ESI) m/z : 444 $[M+H]^+$. 1H NMR (400 MHz, CD_3OD) δ (ppm) 8.91 (d, J = 1.8 Hz, 1H), 8.50 (d, J = 4.9 Hz, 1H), 8.43 (d, J = 4.9 Hz, 1H), 8.18 (dt, J = 8.0, 1.8 Hz, 1H), 7.51 (d, J = 8.0 Hz, 2H), 7.48-7.38 (m, 3H), 6.60-6.49 (m, 1H), 6.23 (dd, J = 16.8, 1.9 Hz, 1H), 5.75-5.67 (m, 1H), 5.41 – 5.30 (m, 1H), 4.68 (d, J = 4.0 Hz, 2H), 4.60-4.49 (m, 1H), 4.32 – 4.16 (m, 1H), 4.15-3.87 (m, 2H), 3.85 – 3.63 (m, 1H).

MYF-03-162



Step 1: Synthesis of Compound 2'

A mixture of tert-butyl 6-oxa-3-azabicyclo[3.1.0]hexane-3-carboxylate (10 g, 54 mmol), NaN_3 (7g, 108 mmol) and NH_4Cl (2.8g, 54 mmol) in MeOH (120 mL) and H_2O (20 mL) was stirred at 65°C under N_2 overnight. The reaction mixture was concentrated in vacuum, the residue was extracted with ethyl acetate (300 mL x 3), the combined organic was washed with water (200 mL x 2), dried over anhydrous Na_2SO_4 , concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 20% v/v) to obtain compound 2' (11 g, yield 89%) as oil. LC-MS (ESI) m/z : 129 $[M+H-100]^+$.

Step 2: Synthesis of Compound 4'

A mixture of compound 2' (3g, 13.1 mmol), 1-(bromomethyl)-4-(trifluoromethyl)benzene (3.1g, 13.1mmol) and 60% NaH (0.6g, 15.7 mmol) in DMF (20 mL) was stirred at rt under N_2 protection for 6 hours. The reaction mixture was diluted with water (100 mL) and extracted with ethyl acetate (100 mL x 2), the combined organic was washed with water (100 mL), dried over anhydrous Na_2SO_4 , concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 30% v/v) to obtain compound 4' (3.8g, yield 75%) as oil. LC-MS (ESI) m/z : 287 $[M+H-$

100]⁺.

Step 3: Synthesis of 4-ethynylpiperidine (Compound 2)

A mixture of tert-butyl 4-ethynylpiperidine-1-carboxylate (1000 mg, 4.7 mmol) and TFA (0.5 mL) in DCM (1 mL) was stirred at rt under N₂ protection for 2 hours. The mixture was concentrated to leave the crude compound **2** (1.2 g) as white solid, which was used directly in the next step. LC-MS (ESI) m/z: 110 [M+H]⁺.

Step 4: Synthesis of *trans*-tert-butyl 3-(4-(piperidin-4-yl)-1H-1,2,3-triazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 3)

A mixture of compound **4'** (200 mg, 0.52 mmol), compound **2** (113 mg, 1.04 mmol), CuSO₄ (48 mg, 0.3 mmol) and sodium L-ascorbate (28 mg, 0.14 mmol) in THF (1 mL), H₂O (1 mL) and ⁿBuOH (1 mL) was stirred at 70 °C under N₂ overnight. The reaction mixture was concentrated in vacuum, the residue was diluted with water (50 mL) and extracted with ethyl acetate (20 mL x 2), the combined organic was washed with brine (20 mL), dried over anhydrous Na₂SO₄, concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 90% v/v) to obtain compound **3** as yellow solid (200 mg, yield 78%). LC-MS (ESI) m/z: 496[M+H]⁺.

Step 5: Synthesis of *trans*-tert-butyl 3-(4-(1-methylpiperidin-4-yl)-1H-1,2,3-triazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 4)

A mixture of compound **3** (100 mg, 0.2 mmol) and formaldehyde (10 mg, 0.3 mmol) and NaBH₄ (20 mg, 0.4 mmol) in MeOH (10 mL) was stirred at rt for 2 hours. The resulting mixture was concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 80% v/v) to obtain compound **4** as oil (70 mg, yield 69%). LC-MS (ESI) m/z: 510[M+H]⁺.

Step 6: Synthesis of 1-methyl-4-(1-(*trans*-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-3-yl)-1H-1,2,3-triazol-4-yl)piperidine (Compound 5)


A mixture of compound **4** (70 mg, 0.13 mmol) and TFA (0.5 mL) in DCM (1 mL) was stirred at rt under N₂ for 2 hours. The mixture was concentrated to leave the crude compound **5** (50 mg) as yellow oil, which was used directly in the next step. LC-MS (ESI) m/z: 410[M+H]⁺.

Step 7: Synthesis of 1-(*trans*-3-(4-(1-methylpiperidin-4-yl)-1H-1,2,3-triazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-1-yl)prop-2-en-1-one (Compound MYF-03-162)

To the mixture of compound **5** (50 mg, 0.12 mmol) and TEA (30 mg, 0.24 mmol) in DCM (10 mL) was added acryloyl chloride (20 mg, 0.12 mmol), the mixture was stirred at rt under N₂ for 2 hours, and then concentrated and purified by prep-HPLC to obtain compound **MYF-03-162** as white solid (13 mg, yield 23%). LC-MS (ESI) m/z: 464 [M+H]⁺; ¹H NMR (400 MHz, CD₃OD) δ (ppm) 7.89 (d,

$J = 3.6$ Hz, 1H), 7.63 (d, $J = 8.1$ Hz, 2H), 7.49(d, $J = 8.1$ Hz, 2H), 6.61 (dd, $J = 16.8, 10.4$ Hz, 1H), 6.31 (dd, $J = 16.8, 1.8$ Hz, 1H), 5.78 (ddd, $J = 10.4, 3.6, 1.9$ Hz, 1H), 5.39 – 5.27 (m, 1H), 4.73 (d, $J = 4.6$ Hz, 2H), 4.61-4.50 (m, 1H), 4.34 – 4.10 (m, 2H), 4.09-3.69 (m, 2H), 2.94 (d, $J = 11.6$ Hz, 2H), 2.80-2.68 (m, 1H), 2.31 (s, 3H), 2.24 – 2.12 (m, 2H), 2.08 – 1.98 (m, 2H), 1.80-1.65 (m, 2H).

MYF-03-139



Step 1: Synthesis of *trans*-tert-butyl 3-hydroxy-4-(4-(pyridin-3-yl)-1H-1,2,3-triazol-1-yl)pyrrolidine-1-carboxylate (Compound 3')

A mixture of tert-butyl compound **1'** (1 g, 4.38 mmol), 3-ethynylpyridine (451 mg, 4.38 mmol), CuSO₄ (654 mg, 2.6 mmol) and sodium L-ascorbate (257 mg, 1.3 mmol) in THF (3 mL), H₂O (3 mL) and ⁿBuOH (3 mL) was stirred at 70 °C under N₂ protection overnight. The reaction mixture was diluted with ethyl acetate (60mL), washed with water (40 mL), dried over anhydrous Na₂SO₄, concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 80% v/v) to obtain compound **3'** (650 mg, yield 45%) as yellow solid. LC-MS (ESI) m/z: 332[M+H]⁺.

Step 2: Synthesis of (4,4-difluorocyclohexyl)methyl trifluoromethanesulfonate (Compound 2)

A mixture of (4,4-difluorocyclohexyl)methanol (1 g, 6.6 mmol), Tf₂O (2.81 g, 9.9 mmol) and pyridine (1 mL) in DCM (20 mL) was stirred at rt under N₂ protection for 3 hours. The reaction mixture was diluted with ethyl acetate (100 mL), washed with water (100 mL x 2), dried over anhydrous Na₂SO₄, filtered and concentrated to leave the crude compound **2** (500 mg) as yellow oil, which was used directly in the next step. LC-MS (ESI) m/z: no MS.

Step 3: Synthesis of *trans*-tert-butyl 3-((4,4-difluorocyclohexyl)methoxy)-4-(4-(pyridin-3-yl)-1H-1,2,3-triazol-1-yl)pyrrolidine-1-carboxylate (Compound 3)

A mixture of compound **2** (200 mg, 0.7 mmol), compound **3'** (58 mg, 0.175 mmol) and ^tBuONa (25 mg, 0.26 mmol) in THF (5 mL) was stirred at 0 °C under N₂ for 2 hours. The reaction mixture was diluted with ethyl acetate (50 mL), washed with water (50 mL x 2), dried over anhydrous Na₂SO₄, concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 50% v/v) to obtain compound **3** as oil (100 mg, yield 30%). LC-MS (ESI) m/z: 464[M+H]⁺.


Step 4: Synthesis of 3-(1-(trans-4-((4,4-difluorocyclohexyl)methoxy)pyrrolidin-3-yl)-1H-1,2,3-triazol-4-yl)pyridine (Compound 4)

A mixture of compound **3** (50 mg, 0.1 mmol) and TFA (1 mL) in DCM (3 mL) was stirred at rt for 2 hours. The mixture was concentrated to leave the crude compound **4** (50 mg) as yellow oil, which was used directly in the next step. LC-MS (ESI) m/z: 364[M+H]⁺.

Step 5: Synthesis of 1-(trans-3-((4,4-difluorocyclohexyl)methoxy)-4-(4-(pyridin-3-yl)-1H-1,2,3-triazol-1-yl)pyrrolidin-1-yl)prop-2-en-1-one (Compound MYF-03-139)

A mixture of compound **4** (50 mg, 0.1 mmol), acryloyl chloride (10 mg, 0.1 mmol), and TEA (20 mg, 0.2 mmol) in DCM (5 mL) was stirred at rt under N₂ for 2 hours. The mixture was diluted with DCM (50 mL), washed with water (50 mL x 2), dried over anhydrous Na₂SO₄, concentrated and purified by prep-HPLC to obtain **MYF-03-139** as white solid (16 mg, yield 38%). LC-MS (ESI) m/z: 418[M+H]⁺; ¹H-NMR (400 MHz, CD₃OD) δ (ppm) 8.91 (d, *J* = 1.2 Hz, 1H), 8.55 – 8.40 (m, 2H), 8.18 (dt, *J* = 8.0, 1.9 Hz, 1H), 7.43 (dd, *J* = 8.0, 4.9 Hz, 1H), 6.55 (ddd, *J* = 16.8, 10.4, 2.5 Hz, 1H), 6.28 – 6.20 (m, 1H), 5.70 (ddd, *J* = 10.5, 4.7, 1.9 Hz, 1H), 5.29 – 5.18 (m, 1H), 4.43 – 4.32 (m, 1H), 4.27 – 4.16 (m, 1H), 4.07 – 3.37 (m, 5H), 1.91 (ddd, *J* = 13.9, 7.0, 3.5 Hz, 2H), 1.76 – 1.54 (m, 5H), 1.26 – 1.13 (m, 2H).

MYF-03-137



Step 1: Synthesis of (6-(trifluoromethyl)pyridin-3-yl)methanol (Compound 2)

A mixture of 6-(trifluoromethyl)nicotinaldehyde (170 mg, 1 mmol) and NaBH₄ (76 mg, 2 mmol) in MeOH (3 mL) was stirred at 0 °C for 3 hours. The reaction mixture was concentrated in vacuum, the residue was extracted with ethyl acetate (60 mL), washed with water (40 mL), dried over anhydrous

Na₂SO₄, filtered and concentrated to leave the crude product (150 mg, yield 84%) as oil, which was used directly in the next step. LC-MS (ESI) m/z: 178[M+H]⁺.

Step 2: Synthesis of 5-(bromomethyl)-2-(trifluoromethyl)pyridine (Compound 3)

A mixture of compound **2** (400 mg, 2.2 mmol) and 48% aqueous HBr solution (6 mL) was stirred at 110 °C overnight. The reaction mixture was concentrated in vacuum, the residue was diluted with ethyl acetate (60 mL), washed with water (40 mL), dried over anhydrous Na₂SO₄, concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 20% v/v) to obtain compound **3** as oil (200 mg, yield 38%). LC-MS (ESI) m/z: 240[M+H]⁺.

Step 3: Synthesis of *trans*-tert-butyl 3-(4-(pyridin-3-yl)-1H-1,2,3-triazol-1-yl)-4-((6-(trifluoromethyl)pyridin-3-yl)methoxy)pyrrolidine-1-carboxylate (Compound 5)

A mixture of compound **3** (100 mg, 0.4 mmol), compound **4** (140 mg, 0.4 mmol) and 60% NaH (32 mg, 0.8 mmol) in DMF (5 mL) was stirred at rt under N₂ overnight. The reaction mixture was diluted with water (100 mL) and extracted with ethyl acetate (20 mL x 2), the combined organic was washed with water (50 mL), dried over anhydrous Na₂SO₄, concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 90% v/v) to obtain compound **5** as solid (100 mg, yield 51%). LC-MS (ESI) m/z: 491[M+H]⁺.

Step 4: Synthesis of 5-((*trans*-4-(4-(pyridin-3-yl)-1H-1,2,3-triazol-1-yl)pyrrolidin-3-yl)oxy)methyl)-2-(trifluoromethyl)pyridine (Compound 6)


A mixture of compound **5** (100 mg, 0.2 mmol) and TFA (1 mL) in DCM (3 mL) was stirred at rt for 2 hours. The mixture was concentrated to leave the crude compound **6** (100 mg, crude) as yellow oil, which was used directly in the next step. LC-MS (ESI) m/z: 391

Step 5: Synthesis of 1-(*trans*-3-(4-(pyridin-3-yl)-1H-1,2,3-triazol-1-yl)-4-((6-(trifluoromethyl)pyridin-3-yl)methoxy)pyrrolidin-1-yl)prop-2-en-1-one (Compound MYF-03-137)

A mixture of compound **6** (50 mg, 0.125 mmol), acryloyl chloride (15 mg, 0.125 mmol) and TEA (25 mg, 0.25 mmol) in DCM (3 mL) was stirred at 0 °C for 2 hours. The mixture was concentrated and purified by prep-HPLC to obtain **MYF-03-137** as white solid (5 mg, yield 9%). LC-MS (ESI) m/z: 445. ¹H NMR (400 MHz, CD₃OD) δ (ppm) 9.04 (d, *J* = 2.2 Hz, 1H), 8.76 – 8.50 (m, 3H), 8.31 (dt, *J* = 7.9, 1.9 Hz, 1H), 8.03 (d, *J* = 8.2 Hz, 1H), 7.80 (d, *J* = 8.1 Hz, 1H), 7.56 (dd, *J* = 8.0, 4.9 Hz, 1H), 6.67 (ddd, *J* = 16.8, 10.4, 3.5 Hz, 1H), 6.36 (dd, *J* = 16.8, 1.7 Hz, 1H), 5.83 (ddd, *J* = 10.5, 5.1, 1.9 Hz, 1H), 5.58 – 5.44 (m, 1H), 4.97 (s, 2H), 4.81 – 4.68 (m, 1H), 4.48 – 4.03 (m, 3H), 4.02 – 3.80 (m, 1H).

1195

MYF-03-138



1196

Step 1: Synthesis of 4-((trimethylsilyl)ethynyl)pyridin-2(1H)-one (Compound 3)

To the solution of ethynyltrimethylsilane (0.595 g, 6.07 mmol) in DMF (40 mL) was added 4-bromopyridin-2(1H)-one (1 g, 5.8 mmol), PdCl₂(PPh₃)₂ (0.204 g, 0.29 mmol), CuI (55 mg, 0.29 mmol) and Et₃N (1.17 g, 11.6 mmol). The mixture was stirred at 90 °C under N₂ for 2 hours. After cooled down to rt the mixture was diluted with water (200 mL) and extracted with EtOAc (50 mL x 2), the combined organic was dried over anhydrous Na₂SO₄, concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 50% v/v) to obtain compound **3** as oil (500 mg, yield 45.4%). LC-MS (ESI) m/z: 192[M+H]⁺.

1205

Step 2: Synthesis of 4-ethynylpyridin-2(1H)-one (Compound 4)

To the solution of compound **3** (450 mg, 2.35 mmol) in MeOH (20 mL) was added KOH (263 mg, 4.70 mmol). The mixture was stirred at rt under N₂ for 2 hours. The resulted mixture was concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 90% v/v) to obtain compound **4** as oil (200 mg, yield 71.4%). LC-MS (ESI) m/z: 120[M+H]⁺.

1211

Step 3: Synthesis of *trans*-tert-butyl 3-(4-(2-oxo-1,2-dihydropyridin-4-yl)-1H-1,2,3-triazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 6)

To the solution of compound **5** (250 mg, 0.65 mmol) in THF (10mL), H₂O (10mL) and ⁿBuOH (10mL) was added compound **4** (116 mg, 0.97 mmol), CuSO₄ (15 mg, 0.065 mmol) and sodium L-ascorbate (26 mg, 0.13 mmol). The mixture was stirred at 70 °C under N₂ for 16 hours. The resulting mixture was concentrated and purified by flash column chromatography on silica gel (methanol in dichloromethane = 20% v/v) to obtain compound **6** as solid (200 mg, yield 60%). LC-MS (ESI) m/z: 506[M+H]⁺.

1220

Step 4: Synthesis of 4-(1-(*trans*-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-3-yl)-1H-1,2,3-triazol-4-yl)pyridin-2(1H)-one (Compound 7)


1222

To the solution of compound **6** (180 mg, 0.36 mmol) in DCM (10 mL) was added TFA (2 mL). The mixture was stirred at rt for 2 hours and concentrated in vacuum, the residue was adjusted to pH~8 with NaHCO₃ solution and extracted with EtOAc (50 mL x 3), the combined organics were washed with brine (100 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to leave crude compound **7** as oil (150 mg, crude). LC-MS (ESI) m/z: 406[M+H]⁺.

Step 5: Synthesis of 4-(1-(*trans*-1-acryloyl-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-3-yl)-1H-1,2,3-triazol-4-yl)pyridin-2(1H)-one (Compound MYF-03-138)

To the solution of compound **7** (130 mg, 0.32 mmol) in THF (10 mL) was added acryloyl chloride (29 mg, 0.32 mmol) and Et₃N (65 mg, 0.64 mmol). The mixture was stirred at 0°C for 1 hour, and then concentrated and purified by prep-HPLC to obtain compound **MYF-03-138** as solid (44 mg, yield 30.0%). LC-MS (ESI) m/z: 460[M+H]⁺. ¹H-NMR (400 MHz, DMSO-*d*₆) δ (ppm) 11.58 (br, 1H), 8.89 (d, *J* = 5.5 Hz, 1H), 7.70 (d, *J* = 8.0 Hz, 2H), 7.54 (d, *J* = 8.0 Hz, 2H), 7.46 (d, *J* = 6.6 Hz, 1H), 6.79 (s, 1H), 6.70 – 6.56 (m, 2H), 6.24 – 6.14 (m, 1H), 5.79-5.71 (m, 1H), 5.53 – 5.39 (m, 1H), 4.81 – 4.71 (m, 2H), 4.62-4.50 (m, 1H), 4.30-4.13 (m, 1H), 4.12-3.95 (m, 1H), 3.91-3.79 (m, 1H), 3.68-3.58 (m, 1H).

MYF-03-146



Step 1: Synthesis of 1-methyl-4-((trimethylsilyl)ethynyl)pyridin-2(1H)-one (Compound 3)

To the solution of 4-bromo-1-methylpyridin-2(1H)-one (**1**, 5.3 mmol) in DMF (20 mL) was added ethynyltrimethylsilane (0.55 g, 5.6 mmol), PdCl₂(PPh₃)₂ (0.21 g, 0.3 mmol), CuI (0.06 g, 0.3 mmol) and Et₃N (1.07 g, 10.6 mmol). The mixture was stirred at room temperature under N₂ for 2 hours. The resulting mixture was concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 50% v/v) to obtain compound **3** as oil (1.1 g, yield 99.9%). LC-MS (ESI) m/z: 206[M+H]⁺.

Step 2: Synthesis of 4-ethynyl-1-methylpyridin-2(1H)-one (Compound 4)

To the solution of compound **3** (1 g, 4.9 mmol) in MeOH (20 mL) was added K₂CO₃ (1.35 g, 9.8

mmol). The mixture was stirred at room temperature under N₂ for 2 hours. The resulting mixture was concentrated and purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 90% v/v) to obtain compound **4** as solid (400 mg, yield 56.1%). LC-MS (ESI) m/z: 134[M+H]⁺.

Step 3: Synthesis of *trans*-tert-butyl 3-(4-(1-methyl-2-oxo-1,2-dihydropyridin-4-yl)-1H-1,2,3-triazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 6)

To the solution of *trans*-tert-butyl 3-azido-4-(4-(trifluoromethyl)benzyloxy) pyrrolidine-1-carboxylate (350 mg, 0.91 mmol) in THF (5mL), H₂O (5mL) and ⁿBuOH (5mL) was added compound **4** (181 mg, 1.36 mmol), CuSO₄ (23 mg, 0.09 mmol) and sodium L-ascorbate (36 mg, 0.18 mmol). The mixture was stirred at 70 °C under N₂ for 16 hours. The resulting mixture was concentrated and purified by flash column chromatography on silica gel (methanol in dichloromethane=20% v/v) to obtain compound **6** as solid (300 mg, yield 63.5%). LC-MS (ESI) m/z: 520[M+H]⁺.


Step 4: Synthesis of 1-methyl-4-(1-(*trans*-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-3-yl)-1H-1,2,3-triazol-4-yl)pyridin-2(1H)-one (Compound 7)

To the solution of compound **6** (280 mg, 0.54 mmol) in DCM (10 mL) was added TFA (2 mL). The mixture was stirred at room temperature for 2 hours and concentrated in vacuum, the residue was adjusted to pH~8 with NaHCO₃ solution and extracted with EtOAc (50 mL x 3), the combined organics were washed with brine (100 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to leave the crude compound **7** as oil (200 mg, crude). LC-MS (ESI) m/z: 420[M+H]⁺.

Step 5: Synthesis of 4-(1-(*trans*-1-acryloyl-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-3-yl)-1H-1,2,3-triazol-4-yl)-1-methylpyridin-2(1H)-one (Compound MYF-03-146)

To the solution of compound **7** (180 mg, 0.43 mmol) in THF (10 mL) was added acryloyl chloride (39 mg, 0.43 mmol) and Et₃N (87 mg, 0.86 mmol). The mixture was stirred at 0°C under N₂ for 1 hour, and then concentrated and purified by prep-HPLC to obtain compound **MYF-03-146** as solid (215 mg, yield 95.5%). LC-MS (ESI) m/z: 474[M+H]⁺. ¹H-NMR (400 MHz, DMSO-*d*₆): δ (ppm) 8.89 (d, *J* = 5.8 Hz, 1H), 7.78 (d, *J* = 7.1 Hz, 1H), 7.70 (d, *J* = 8.2 Hz, 2H), 7.53 (d, *J* = 8.1 Hz, 2H), 6.85 (s, 1H), 6.76 – 6.53 (m, 2H), 6.19 (dd, *J* = 16.8, 2.3 Hz, 1H), 5.73 (ddd, *J* = 10.2, 5.1, 2.3 Hz, 1H), 5.47 (d, *J* = 25.4 Hz, 1H), 4.75 (s, 2H), 4.57 (d, *J* = 24.5 Hz, 1H), 4.31 – 3.57 (m, 4H), 3.44 (s, 3H).

MYF-03-135



Step 1: Synthesis of *trans*-tert-butyl 3-(4-(tetrahydro-2H-pyran-4-yl)-1H-1,2,3-triazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 3)

To the solution of compound 2 (300 mg, 0.78 mmol) in THF (10 mL), H₂O (10 mL) and ⁿBuOH (10 mL) was added 4-ethynyltetrahydro-2H-pyran (128 mg, 1.16 mmol), CuSO₄ (20 mg, 0.078 mmol) and sodium L-ascorbate (31 mg, 0.156 mmol). The mixture was stirred at 70 °C under N₂ for 16 hours. The resulting mixture was concentrated and purified by flash column chromatography on silica gel (Methanol in dichloromethane = 20% v/v) to obtain compound 3 as solid (300 mg, yield 77.9%). LC-MS (ESI) m/z: 497 [M+H]⁺.

Step 2: Synthesis of 4-(tetrahydro-2H-pyran-4-yl)-1-(*trans*-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-3-yl)-1H-1,2,3-triazole (Compound 4)


To the solution of compound 3 (280 mg, 0.56 mmol) in DCM (10 mL) was added TFA (2 mL). The mixture was stirred at room temperature for 2 hours and concentrated in vacuum, the residue was adjusted to pH~8 with NaHCO₃ solution and extracted with EtOAc (50 mL x 3), the combined organics were washed with brine (100 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to leave the crude compound 4 as oil (250 mg, crude). LC-MS (ESI) m/z: 397 [M+H]⁺.

Step 3: Synthesis of 1-(*trans*-3-(4-(tetrahydro-2H-pyran-4-yl)-1H-1,2,3-triazol-1-yl)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-1-yl)prop-2-en-1-one (Compound MYF-03-135)

To the solution of compound 4 (230 mg, 0.58 mmol) in THF (10 mL) was added acryloyl chloride (52 mg, 0.58 mmol) and Et₃N (117 mg, 1.16 mmol). The mixture was stirred at 0°C under N₂ for 1 hour, and then concentrated and purified by prep-HPLC to obtain compound **MYF-03-135** as solid (180 mg, yield 69.0%). LC-MS (ESI) m/z: 451[M+H]⁺. ¹H-NMR (400 MHz, DMSO-*d*₆): δ (ppm) 8.05 (d, *J* = 6.6 Hz, 1H), 7.70 (d, *J* = 7.6 Hz, 2H), 7.51 (d, *J* = 7.8 Hz, 2H), 6.66 – 6.57 (m, 1H), 6.18 (d, *J* = 16.8 Hz, 1H), 5.77 – 5.67 (m, 1H), 5.43 – 5.29 (m, 1H), 4.72 (d, *J* = 5.2 Hz, 2H), 4.50 (dd, *J* = 31.4, 5.0 Hz, 1H), 4.24 – 3.96 (m, 2H), 3.89 (d, *J* = 13.1 Hz, 2H), 3.82 (td, *J* = 13.1, 4.4 Hz, 1H), 3.58 (dd, *J* = 13.1, 3.3 Hz, 1H), 3.44 (t, *J* = 11.3 Hz, 2H), 2.93 (t, *J* = 10.6 Hz, 1H), 1.85 (d, *J* = 12.9 Hz, 2H), 1.61 (q, *J* = 11.4 Hz, 2H).

1315

MYF-03-176



1316

Step 1: Synthesis of (3R,4R)-tert-butyl 3-azido-4-hydroxypyrrolidine-1-carboxylate (Compound 3)

A mixture of compound **1** (4000mg, 21.6mmol), compound **2** (2664mg, 23.2mmol) and catalyst **3** (328mg, 0.42mmol) was stirred at rt for overnight under N₂ protection. The reaction mixture was treated with MeOH (60mL) and K₂CO₃ (1788mg, 12.8mmol) and continued to stirring for 5h. The reaction mixture was extracted with ethyl acetate (300mL x 3), and washed by water (300 mL x 2). The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by flash column chromatography on silica gel (ethyl acetate in petroleum ether = 20% v/v) to obtain 3.5 g product compound **3** as yellow oil (3.5g, yield 71%). LC-MS (ESI) m/z: 129[M+H]⁺.

Step 2: Synthesis of (3R,4R)-tert-butyl 3-azido-4-(4-(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 5)

A mixture of compound **3** (500mg, 2.19 mmol), compound **4** (524mg, 2.19mmol) and NaH (105mg, 2.62 mmol) in THF (10 mL) was stirred at rt for 6h under N₂ protection. The reaction mixture was monitored by LC-MS. The reaction mixture was extracted with ethyl acetate (100mL), and washed by water (50mL). The organic layer was dried over Na₂SO₄, was concentrated and purified by column chromatography on silica gel (ethyl acetate in petroleum ether = 30% v/v) to obtain 600mg product compound **5** as white oil (600mg, yield 95%). LC-MS (ESI) m/z: 287[M+ H]⁺.

Step 3: Synthesis of (3R,4R)-tert-butyl 3-amino-4-(4-(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 7)

A mixture of compound **5** (1000mg, 2.58 mmol), PPh₃ (814mg, 3.1mmol) and H₂O (930mg, 51.6 mmol) in THF (40 mL) was stirred at 70 °C for 5h under N₂ protection. The reaction mixture was monitored by LC-MS. The reaction mixture was extracted with ethyl acetate (300mL), and washed by water (200mL), The organic layer was dried over Na₂SO₄, concentrated and purified by p-HPLC to obtain 800mg product compound **7** as yellow oil (800mg, yield 86%). LC-MS (ESI) m/z: 261[M+H-100]⁺.

Step 4: Synthesis of (3R,4R)-tert-butyl 3-(pyrimidin-2-ylamino)-4-(4-

(trifluoromethyl)benzyloxy)pyrrolidine-1-carboxylate (Compound 9)

A mixture of compound **7** (600 mg, 1.6mmol), compound **8** (240 mg, 1.84mmol) and DIPEA (420mg, 3.24 mmol) in *n*BuOH (6 mL) was stirred at 70 °C for overnight under N₂ protection. The reaction mixture was monitored by LC-MS. The reaction mixture was concentrated and purified by p-HPLC to obtain 500mg product compound **9** as white oil (500mg, yield 71%). LC-MS (ESI) *m/z*: 439[M+H]⁺.

Step 5: Synthesis of N-((3R,4R)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-3-yl)pyrimidin-2-amine (Compound 10)

A mixture of compound **9** (400mg, 0.91 mmol) and TFA (1mL) in DCM (3 mL) was stirred at rt for 2h under N₂ protection. The mixture was concentrated to obtain 300mg crude of compound **10** as yellow oil. Used for next step (400mg, yield 97%). LC-MS (ESI) *m/z*: 339[M+H]⁺.

Step 6: Synthesis of 2-fluoro-1-((3R,4R)-3-(pyrimidin-2-ylamino)-4-(4-(trifluoromethyl)benzyloxy)pyrrolidin-1-yl)prop-2-en-1-one (MYF-03-176)

A mixture of compound **10** (200mg, 0.58 mmol), compound **11** (60mg, 0.69mmol) and HATU (256mg, 0.69 mmol) and DIEA (224mg, 1.74mmol) in DMF (5 mL) was stirred rt for overnight under N₂ protection. The reaction mixture was monitored by LC-MS. The reaction mixture was concentrated and purified by p-HPLC to obtain 150mg product **MYF-03-176** as white solid (150mg, yield 63%). LC-MS (ESI) *m/z*: 411[M+H]⁺. ¹H NMR (400 MHz, MeOD) δ 8.44 (s, 2H), 7.82 – 7.50 (m, 4H), 6.83 (dd, *J* = 8.3, 4.9 Hz, 1H), 5.50 (dt, *J* = 47.1, 3.3 Hz, 1H), 5.27 (dt, *J* = 16.5, 3.4 Hz, 1H), 4.82 (dd, *J* = 13.3, 8.8 Hz, 2H), 4.68 – 4.57 (m, 1H), 4.28 – 3.67 (m, 5H).

16 References

1. Zhang, Z.; Marshall, A. G., A universal algorithm for fast and automated charge state deconvolution of electrospray mass-to-charge ratio spectra. *Journal of the American Society for Mass Spectrometry* **1998**, 9 (3), 225-233.
2. Hughes, C. S.; Foehr, S.; Garfield, D. A.; Furlong, E. E.; Steinmetz, L. M.; Krijgsveld, J., Ultrasensitive proteome analysis using paramagnetic bead technology. *Molecular Systems Biology* **2014**, 10 (10), 757.
3. Alexander, W. M.; Ficarro, S. B.; Adelmant, G.; Marto, J. A., multipliez v2.0: A Python-based ecosystem for shared access and analysis of native mass spectrometry data. *PROTEOMICS* **2017**, 17 (15-16), 1700091.
4. Ficarro, S. B.; Alexander, W. M.; Marto, J. A., mzStudio: A Dynamic Digital Canvas for User-Driven Interrogation of Mass Spectrometry Data. *Proteomes* **2017**, 5 (3), 20.
5. Ficarro, S. B.; Browne, C. M.; Card, J. D.; Alexander, W. M.; Zhang, T.; Park, E.; McNally, R.; Dhe-Paganon, S.; Seo, H.-S.; Lamberto, I.; Eck, M. J.; Buhrlage, S. J.; Gray, N. S.; Marto, J. A., Leveraging Gas-Phase Fragmentation Pathways for Improved Identification and Selective Detection of Targets Modified by Covalent Probes. *Analytical Chemistry* **2016**, 88 (24), 12248-12254.

- 1379 6. Kabsch, W., Integration, scaling, space-group assignment and post-refinement. *Acta*
1380 *crystallographica. Section D, Biological crystallography* **2010**, 66 (Pt 2), 133-44.
- 1381 7. McCoy, A. J.; Grosse-Kunstleve, R. W.; Adams, P. D.; Winn, M. D.; Storoni, L. C.; Read, R.
1382 J., Phaser crystallographic software. *Journal of applied crystallography* **2007**, 40 (Pt 4), 658-674.
- 1383 8. Adams, P. D.; Afonine, P. V.; Bunkoczi, G.; Chen, V. B.; Davis, I. W.; Echols, N.; Headd, J.
1384 J.; Hung, L. W.; Kapral, G. J.; Grosse-Kunstleve, R. W.; McCoy, A. J.; Moriarty, N. W.; Oeffner, R.;
1385 Read, R. J.; Richardson, D. C.; Richardson, J. S.; Terwilliger, T. C.; Zwart, P. H., PHENIX: a
1386 comprehensive Python-based system for macromolecular structure solution. *Acta crystallographica.*
1387 *Section D, Biological crystallography* **2010**, 66 (Pt 2), 213-21.
- 1388 9. Emsley, P.; Cowtan, K., Coot: model-building tools for molecular graphics. *Acta*
1389 *crystallographica. Section D, Biological crystallography* **2004**, 60 (Pt 12 Pt 1), 2126-32.
- 1390 10. Lu, W.; Wang, J.; Li, Y.; Tao, H.; Xiong, H.; Lian, F.; Gao, J.; Ma, H.; Lu, T.; Zhang, D.; Ye,
1391 X.; Ding, H.; Yue, L.; Zhang, Y.; Tang, H.; Zhang, N.; Yang, Y.; Jiang, H.; Chen, K.; Zhou, B.; Luo,
1392 C., Discovery and biological evaluation of vinylsulfonamide derivatives as highly potent, covalent
1393 TEAD autopalmitoylation inhibitors. *European Journal of Medicinal Chemistry* **2019**, 184, 111767.
- 1394 11. Kuljanin, M.; Mitchell, D. C.; Schweppe, D. K.; Gikandi, A. S.; Nusinow, D. P.; Bulloch, N.
1395 J.; Vinogradova, E. V.; Wilson, D. L.; Kool, E. T.; Mancias, J. D.; Cravatt, B. F.; Gygi, S. P.,
1396 Reimagining high-throughput profiling of reactive cysteines for cell-based screening of large
1397 electrophile libraries. *Nature Biotechnology* **2021**, 39, 630-641.
- 1398 12. (a) Navarrete-Perea, J.; Yu, Q.; Gygi, S. P.; Paulo, J. A., Streamlined Tandem Mass Tag (SL-
1399 TMT) Protocol: An Efficient Strategy for Quantitative (Phospho)proteome Profiling Using Tandem
1400 Mass Tag-Synchronous Precursor Selection-MS3. *Journal of Proteome Research* **2018**, 17 (6), 2226-
1401 2236; (b) Li, J.; Van Vranken, J. G.; Pontano Vaite, L.; Schweppe, D. K.; Huttlin, E. L.; Etienne, C.;
1402 Nandhikonda, P.; Viner, R.; Robitaille, A. M.; Thompson, A. H.; Kuhn, K.; Pike, I.; Bomgarden, R.
1403 D.; Rogers, J. C.; Gygi, S. P.; Paulo, J. A., TMTpro reagents: a set of isobaric labeling mass tags
1404 enables simultaneous proteome-wide measurements across 16 samples. *Nature Methods* **2020**, 17 (4),
1405 399-404.
- 1406 13. Schweppe, D. K.; Eng, J. K.; Yu, Q.; Bailey, D.; Rad, R.; Navarrete-Perea, J.; Huttlin, E. L.;
1407 Erickson, B. K.; Paulo, J. A.; Gygi, S. P., Full-Featured, Real-Time Database Searching Platform
1408 Enables Fast and Accurate Multiplexed Quantitative Proteomics. *Journal of Proteome Research* **2020**,
1409 19 (5), 2026-2034.
- 1410 14. (a) Eng, J. K.; Hoopmann, M. R.; Jahan, T. A.; Egertson, J. D.; Noble, W. S.; MacCoss, M. J.,
1411 A Deeper Look into Comet—Implementation and Features. *Journal of The American Society for*
1412 *Mass Spectrometry* **2015**, 26 (11), 1865-1874; (b) Eng, J. K.; Jahan, T. A.; Hoopmann, M. R., Comet:
1413 An open-source MS/MS sequence database search tool. *PROTEOMICS* **2013**, 13 (1), 22-24.
- 1414 15. (a) McAlister, G. C.; Huttlin, E. L.; Haas, W.; Ting, L.; Jedrychowski, M. P.; Rogers, J. C.;
1415 Kuhn, K.; Pike, I.; Grothe, R. A.; Blethrow, J. D.; Gygi, S. P., Increasing the Multiplexing Capacity of

TMTs Using Reporter Ion Isotopologues with Isobaric Masses. *Analytical Chemistry* **2012**, *84* (17), 7469-7478; (b) Beausoleil, S. A.; Villén, J.; Gerber, S. A.; Rush, J.; Gygi, S. P., A probability-based approach for high-throughput protein phosphorylation analysis and site localization. *Nature Biotechnology* **2006**, *24* (10), 1285-1292; (c) Elias, J. E.; Gygi, S. P., Target-decoy search strategy for increased confidence in large-scale protein identifications by mass spectrometry. *Nature Methods* **2007**, *4* (3), 207-214; (d) Huttlin, E. L.; Jedrychowski, M. P.; Elias, J. E.; Goswami, T.; Rad, R.; Beausoleil, S. A.; Villén, J.; Haas, W.; Sowa, M. E.; Gygi, S. P., A Tissue-Specific Atlas of Mouse Protein Phosphorylation and Expression. *Cell* **2010**, *143* (7), 1174-1189.

16. Backus, K. M.; Correia, B. E.; Lum, K. M.; Forli, S.; Horning, B. D.; González-Páez, G. E.; Chatterjee, S.; Lanning, B. R.; Teijaro, J. R.; Olson, A. J.; Wolan, D. W.; Cravatt, B. F., Proteome-wide covalent ligand discovery in native biological systems. *Nature* **2016**, *534* (7608), 570-574.

17. Hafner, M.; Niepel, M.; Chung, M.; Sorger, P. K., Growth rate inhibition metrics correct for confounders in measuring sensitivity to cancer drugs. *Nature Methods* **2016**, *13* (6), 521-527.

18. Corsello, S. M.; Nagari, R. T.; Spangler, R. D.; Rossen, J.; Kocak, M.; Bryan, J. G.; Humeidi, R.; Peck, D.; Wu, X.; Tang, A. A.; Wang, V. M.; Bender, S. A.; Lemire, E.; Narayan, R.; Montgomery, P.; Ben-David, U.; Garvie, C. W.; Chen, Y.; Rees, M. G.; Lyons, N. J.; McFarland, J. M.; Wong, B. T.; Wang, L.; Dumont, N.; O'Hearn, P. J.; Stefan, E.; Doench, J. G.; Harrington, C. N.; Greulich, H.; Meyerson, M.; Vazquez, F.; Subramanian, A.; Roth, J. A.; Bittker, J. A.; Boehm, J. S.; Mader, C. C.; Tsherniak, A.; Golub, T. R., Discovering the anticancer potential of non-oncology drugs by systematic viability profiling. *Nature Cancer* **2020**, *1* (2), 235-248.

19. Mohseni, M.; Sun, J.; Lau, A.; Curtis, S.; Goldsmith, J.; Fox, V. L.; Wei, C.; Frazier, M.; Samson, O.; Wong, K.-K.; Kim, C.; Camargo, F. D., A genetic screen identifies an LKB1-MARK signalling axis controlling the Hippo-YAP pathway. *Nature Cell Biology* **2014**, *16* (1), 108-117.

20. Li, Z.; Zhao, B.; Wang, P.; Chen, F.; Dong, Z.; Yang, H.; Guan, K.-L.; Xu, Y., Structural insights into the YAP and TEAD complex. *Genes & Development* **2010**, *24* (3), 235-240.

21. Dobin, A.; Davis, C. A.; Schlesinger, F.; Drenkow, J.; Zaleski, C.; Jha, S.; Batut, P.; Chaisson, M.; Gingeras, T. R., STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **2012**, *29* (1), 15-21.

22. Love, M. I.; Huber, W.; Anders, S., Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **2014**, *15* (12), 550.

23. Cornwell, M.; Vangala, M.; Taing, L.; Herbert, Z.; Köster, J.; Li, B.; Sun, H.; Li, T.; Zhang, J.; Qiu, X.; Pun, M.; Jeselsohn, R.; Brown, M.; Liu, X. S.; Long, H. W., VIPER: Visualization Pipeline for RNA-seq, a Snakemake workflow for efficient and complete RNA-seq analysis. *BMC Bioinformatics* **2018**, *19* (1), 135.

24. Zhou, Y.; Zhou, B.; Pache, L.; Chang, M.; Khodabakhshi, A. H.; Tanaseichuk, O.; Benner, C.; Chanda, S. K., Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nature Communications* **2019**, *10* (1), 1523.

SUPPLEMENTARY FIGURES




Figure S1 | Chemical structures of known TEAD palmitate pocket binders. Covalent binders are labeled in blue.




Figure S2 | Rationale of covalent fragments design. (a) Binding model of FA in TEAD2 palmitate pocket. (b) Chemical structures of representative covalent fragments. This figure is related to **Figure 1a**.




Figure S3 | Original western blot image of palmitoylation assay in YBD protein of TEAD2. This figure is related to **Figure 1d-e**.




Figure S4 | Original western blot images of palmitoylation assays in YBD protein of TEAD1-4. This figure is related to **Figure 2e**.




Figure S5 | MYF-03-69 exhibited low reactivity towards the cysteine proteome. (a) Overview of the SLC-ABPP proteome-wide selectivity approach for profiling of MYF-03-69. NCI-H226 cells were treated with 0.5, 2, 10 or 25 μ M MYF-03-69 and DMSO over 3 hours in triplicates before cell lysis followed by labeling with DBIA probe. The SLC-ABPP approach was used to profile competition of MYF-03-69 in proteome cysteine labeling. Competition ratio (CR) was calculated in order to quantitatively assess labeling at every cysteine residue. (b) The 7 cysteine sites that were significantly labeled (i.e. exhibited >50% conjugation or CR>2) by 25 μ M of MYF-03-69 are in blue, all of which exhibited dose-dependent engagement.




Figure S6 | MYF-03-69 showed comparatively weak inhibition on YAP-TEAD downstream gene expression in normal mesothelium cell MeT-5A upon 6-hour and 24-hour treatment. This figure is related to Figure 4b-c.




Figure S7 | MYF-03-69 downregulated product of canonical YAP downstream gene AXL with minimal effect on TEAD stability. This figure is related to **Figure 4d**.




Figure S8 | The number of genes with significantly altered expression level decreased with lower concentration treatment in NCI-H226 cells. (a) 0.5 μM MYF-

03-69, (b) 0.1 μ M MYF-03-69. This figure is related to **Figure 4e-f**. Details are in **Supplementary Dataset 2**.




Figure S9 | (a) The calculated GR₅₀ values of MYF-03-69 in 5-day proliferation assay on mesothelioma cells. NCI-H226 and MSTO-211H are Hippo signaling defective mesothelioma cells. NCI-H2452 is Hippo signaling intact mesothelioma cell. MeT-5A is non-cancerous mesothelium cell. This figure is related to **Figure 5a**. (b) MYF-03-69-NC did not inhibit cell proliferation of NCI-H226 and MSTO-211H cells in 5-day treatment. This figure is related to **Figure 5a**. (c) Correlation analysis between compound PRISM sensitivity (log₂.AUC of each cell line) and dependency of certain gene (CRISPR knockout score for each cell line, from DepMap Public 20Q4 Achilles_gene_effect.csv dataset) across the PRISM cell line panel. The Pearson correlation coefficients (X axis) and associated p-values were computed. Positive correlations correspond to dependency correlating with increased sensitivity. The q-values (a corrected significance value accounting for false discovery rate) are computed from p-values using the Benjamini Hochberg algorithm. Associations with q-values above 0.1 are filtered out. Top 3 correlated genes are in red. Details are in **Supplementary Dataset 4**.




Figure S10 | (a) MYF-03-69, but not MYF-03-69-NC, inhibits YAP-TEAD transcriptional activity in NCI-H226 luciferase reporter cells. This figure is related to **Figure 6b-c**. (b) MYF-03-176 downregulates YAP target genes and upregulates a pro-apoptotic gene *BMF*. (c-d) Liver microsome stability and PK parameters comparison of MYF-03-176, MYF-03-69 and MYF-01-37.