

A phased, chromosome-scale genome of ‘Honeycrisp’ apple (*Malus domestica*)

Awais Khan¹□, Sarah B. Carey^{2,3}, Alicia Serrano¹, Huiting Zhang^{4,5}, Heidi Hargarten⁴, Haley Hale^{2,3}, Alex Harkess^{2,3}, Loren Honaas⁴□

¹Plant Pathology and Plant-Microbe Biology Section, Cornell University, Geneva, NY, 14456, USA

²Department of Crop, Soil, and Environmental Sciences, Auburn University, Auburn, AL, 36849, USA

³HudsonAlpha Institute for Biotechnology, Huntsville AL, 35806, USA

⁴USDA ARS Tree Fruit Research Lab, Wenatchee, WA 98801, USA

⁵Department of Horticulture, Washington State University, Pullman, WA USA

□ corresponding author(s) = awais.khan@cornell.edu; loren.honaas@usda.gov

Abstract

‘Honeycrisp’ is one of the most valuable apple cultivars grown in the United States and a popular breeding parent due to its superior fruit quality traits, high levels of cold hardiness, and disease resistance. However, it suffers from a number of physiological disorders and is susceptible to production and postharvest issues. Although several apple genomes have been sequenced in the last decade, there is still a substantial knowledge gap in understanding the genetic mechanisms underlying cultivar-specific traits. Here we present a fully phased, chromosome-level genome of ‘Honeycrisp’ apples, using PacBio HiFi, Omni-C, and Illumina sequencing platforms. Our genome assembly is by far the most contiguous among all the apple genomes. The sizes of the two assembled haplomes are 674 Mb and 660 Mb, with contig N50s of 32.8 Mb and 31.6 Mb, respectively. In total, 47,563 and 48,655 protein coding genes were annotated from each haplome, capturing 96.8-97.4% complete BUSCOs in the eudicot database, the most complete among all *Malus* annotations. A gene family analysis using seven *Malus* genomes shows that a vast majority of ‘Honeycrisp’ genes are assigned into orthogroups shared with other genomes, but it also reveals 121 ‘Honeycrisp’-specific orthogroups. We provide a valuable resource for understanding the genetic basis of horticulturally important traits in apples and other related tree fruit species, including at-harvest and postharvest fruit quality, abiotic stress tolerance, and disease resistance, all of which can enhance breeding efforts in Rosaceae.

Main Content

Background

Apples are the most consumed fruit in the United States (www.ers.usda.gov). The annual estimated total value of the US apple industry is \$21 billion, with five cultivars alone accounting for 2/3 of production (in order of proportion): ‘Gala’, ‘Red Delicious’, ‘Honeycrisp’, ‘Granny Smith’, and ‘Fuji’ (www.usapple.org). Of these, ‘Honeycrisp’ is by far the most valuable - it has roughly twice the value per pound of the next most valuable cultivar, ‘Fuji’ [1]. ‘Honeycrisp’ is appreciated by consumers, and therefore by the US apple industry, for its superior flavor and crisp juicy texture. Importantly, properly stored ‘Honeycrisp’ fruit is well-preserved for several months [2,3]. Additionally, this cultivar shows high levels of cold hardiness [4] and resistance to apple scab, the most economically important fungal disease of apples worldwide [5]. ‘Honeycrisp’ was bred at the University of Minnesota in the 1960s aiming to obtain cold hardy cultivars with high-quality fruit; it was released in 1991 [6] (Figure 1A). Recent genome-wide analysis (following the resolution of the ‘Honeycrisp’ pedigree [7,8]) showed that the genetic background of ‘Honeycrisp’ is distinct from other important apple cultivars in the US. This is highlighted by the success of ‘Honeycrisp’ as a source of interesting genetic diversity in apple breeding programs worldwide to enhance texture, storability, and improved disease resistance [3,5,7,9,10]. In fact, nine new cultivars derived from ‘Honeycrisp’ are already on the market.

Disease resistance, critical for sustainable apple production, has historically been less important due to a market dominated by modern cultivars bred primarily for fruit quality and intensive conventional production systems [11]. Most apple cultivars grown commercially in the US are susceptible to fungal diseases such as apple scab. In temperate and humid regions around the world, frequent applications of fungicides are necessary, contributing significantly to production costs, and to negative human health and environmental impacts [12]. ‘Honeycrisp’ is resistant to apple scab and importantly, this cultivar’s ability to retain crispness and firmness during storage is one of the most outstanding traits of ‘Honeycrisp’ fruit [13]. However, there are other production issues with ‘Honeycrisp’ that present challenges for apple growers (Figure 1E-G). ‘Honeycrisp’ needs a carefully designed nutrient management program during the growing season for optimal production and fruit quality, especially to limit the occurrence of the physiological disorder bitter pit [3]. ‘Honeycrisp’ trees also have greater tendency to develop zonal leaf chlorosis, a physiological disorder that reduces photosynthetic capacity [14]. However,

in the Pacific Northwest (PNW), where a large majority of the ‘Honeycrisp’ apples are grown in the US (www.nwhort.org) due in part to low disease pressure, postharvest issues during long-term storage pose substantial challenges to producers.

The total cullage of ‘Honeycrisp’ fruit is likely among the highest of apple cultivars due to its susceptibility to various postharvest physiological disorders, which have complex etiologies that are poorly understood, and include bitter pit, soft scald, soggy breakdown, and CO₂ injury [15–18]. Postharvest technologies have been developed and deployed to mitigate these disorders [19–21]. However, the efficacy of postharvest treatments can be affected by many factors such as pre-harvest orchard management and at-harvest fruit maturity, a key factor in the maintenance of postharvest apple fruit quality. Growers must balance the acquisition of certain fruit quality characteristics (*e.g.* size, color, flesh texture, and sugar content), while attempting to minimize risk for maturity-linked losses in quality that may occur in the supply chain [22]. This balancing act for maximizing at-harvest fruit quality and long-term cold storage potential in controlled atmospheres is especially difficult for ‘Honeycrisp’.

To maximize both our understanding of genetic mechanisms driving important ‘Honeycrisp’ traits and to assist tree fruit breeders, high quality genomes are required [23]. Indeed, in the last decade since ‘Golden Delicious’ was sequenced [24], a large number of genes and QTLs linked to fruit disease resistance, quality traits, and abiotic stress tolerance in apples have been identified [5,25,26]. Recent high-quality genomes of ‘Gala’, the double haploid ‘Golden Delicious’, and the triploid ‘Hanfu’ provide genomic resources for apple genetics and breeding [27–29]. These studies have identified targeted genomic regions for the development of diagnostic molecular markers to breed disease resistant apple cultivars with good fruit quality [30]. However, the fact remains that traditional apple breeding is a resource-intensive and time-consuming process [9,26,30] and there are still substantial gaps in our knowledge of genetic mechanisms involved in many important apple traits. In this manuscript we report a phased, chromosome-level genome assembly of the ‘Honeycrisp’ apple cultivar generated from PacBio HiFi and Dovetail Omni-C, plus a high-quality annotation – thus providing one of the best genome resources available for apples to date.

Methods

PacBio HiFi sequencing: Cuttings of dormant wood were collected from ‘Honeycrisp’ trees growing in the experimental orchard at Cornell AgriTech (Geneva, NY, USA). The cuttings were placed in water in the greenhouse until leaves began emerging from the buds, and thereafter placed in the dark for two days. Young, dark-adapted leaves were collected and shipped on dry ice to the DNA Sequencing and Genotyping Center at the University of Delaware (DL, USA) for DNA extraction and Single Molecule Real Time (SMRT) PacBio (Pacific BioSciences) sequencing.

High-molecular-weight (HMW) genomic DNA was extracted using a DNeasy Plant Mini Kit (Qiagen) according to the manufacturer's protocol. HMW genomic DNA was sheared to 15 kb fragments, and the HiFi library was prepared using SMRTbell Express Template Prep Kit 2.0 and the DNA/Polymerase Binding Kit 2.0 (Pacific Biosciences) according to the manufacturer's protocol. The sequencing library was size-selected using Sage Blue Pippin (Sage Sciences) to select fragment sizes of >10 kb to ensure removal of smaller fragments and adapter dimers. The library was sequenced on a PacBio Sequel II instrument in CCS/HiFi mode with two SMRT cells with 2 hours pre-extension and 30-hour movie times. Read length distribution and quality of all HiFi reads was assessed using Pave v0.1923 (<https://github.com/conchoecia/pave>).

To scaffold the genome using chromatin conformation sequencing, 1 g of flash-frozen young leaf material was harvested from ‘Honeycrisp’ trees at the Washington State University Sunrise Research Orchard near Rock Island, WA USA and shipped to the HudsonAlpha Institute for Biotechnology in Huntsville, AL USA. The sequencing library was prepared using the Dovetail Genomics Omni-C kit and was sequenced on an Illumina NovaSeq 6000 with PE150 reads. A subset of 1 million read pairs were used as input for Phase Genomics *hic_qc* to validate the overall quality of the library (https://github.com/phasegenomics/hic_qc).

Phased haplome assembly and scaffolding: The expected genome size, heterozygosity, and percent of repeats was assessed by generating 21-mer sequences from the raw HiFi data with Jellyfish v2.3.0 [31] and GenomeScope2 [32,33]. HiFi reads were assembled into contigs using hifiasm v0.16.1 [34,35], with the Hi-C integration mode that incorporated Dovetail Omni-C reads for phasing. Both haplomes of the assembly were scaffolded into chromosomes using the Juicer pipeline v1.6 [36], where the Omni-C reads were mapped separately to both hifiasm haplomes [35,37] with the parameter “-s none”. The Omni-C data was subset to ~100x coverage

and the 3D-DNA v201008 scaffolding pipeline [38] was run with options “--editor-saturation-centile 10 --editor-coarse-resolution 100000 --editor-coarse-region 400000 --editor-repeat-coverage 50”. Contact maps were manually edited using the Juicebox Assembly Tools (JBAT) v1.11.08 [36] to produce the expected 17 chromosomes per haplome. Contigs that contained assembled telomeres were correctly oriented to the terminal ends by searching for the TTTAGGG repeat (or the reverse complement CCCTAAA) using the analyze_genome function of GENESPACE [39]. The chromosomes were numbered and oriented using haplome A of the ‘Gala’ assembly [27]. Genome quality and completeness was assessed using benchmarking universal single-copy gene orthologs (BUSCO v5.2.2) [40] with the “eudicots_odb10” database. Haplome completeness was also assessed using Merqury v1.3 [41].

Transcriptome sequencing: To facilitate gene annotation, total RNA was isolated from various tissues harvested from ‘Honeycrisp’, ‘Red Delicious’, and ‘Granny Smith’ apple trees grown at the Washington State University (WSU) Sunrise Research Orchard near Rock Island, WA USA, ‘Gala’ and ‘WA38’ apple trees grown at the WSU and USDA-ARS Columbia View Research Orchard near Orondo, WA USA, and ‘D’Anjou’ pear trees grown at the WSU Tree Fruit Research and Extension Center Research Orchard in Wenatchee, WA USA using a modified CTAB/Chloroform extraction [42]. Total RNA was assessed for quality (RNA integrity number (RIN) ≥ 8) and purity (A260/280 > 1.8). Sources for all RNA are available in Table 3. 2 μ g of total RNA was used to construct Illumina TruSeq stranded libraries following manufacturers’ instructions. Libraries were sequenced on an Illumina NovaSeq 6000 with PE150 reads at the HudsonAlpha Institute for Biotechnology in Huntsville, AL USA.

Repeat analysis and gene annotation: Repetitive elements on both haplotypes were annotated using EDTA v2.0.0 [43] with flags “--genome, --anno 1, --sensitive=1”. To supplement *ab initio* gene predictions, extensive extrinsic gene annotation homology evidence is needed. Thus, we downloaded existing RNA-seq data for ‘Honeycrisp’ apples from NCBI using SRA toolkit v2.9.6-1 (SRX3408575, SRX5369275, SRX5369276, SRX5369290, SRX5369299, SRX5369300, SRX5369302, SRX8712695 and SRX8712718) [44–46], and combined with the RNA-seq data generated for this project (described above). We *de novo* assembled these two sets of RNA transcripts separately using Trinity v2.13.2 [47], where we used the flag --trimmomatic

to filter the reads for quality. Because the newly generated RNA-seq data were strand-specific, for these we also used the flag “--SS_lib_type RF”. We identified open reading frames using TransDecoder v5.5.0 [48]. Gene annotation was performed using BRAKER2 v2.1.6 [49], where we ran BRAKER2 twice, with RNA-seq data and protein databases run separately. For the RNA-seq run, we first filtered the data for adapters and quality using TRIMMOMATIC v0.39 [50] with leading and trailing values of 3, sliding window of 30, jump of 10, and a minimum remaining read length of 40. We next mapped these data to the genome using STAR v2.7.9a [51] and combined the BAM files using SAMtools [52]. For the homology-based annotation in BRAKER2, we used gene models from *Malus domestica* ‘Gala’ diploid v2, *M. sieversii* diploid v2 [27], *M. baccata* v1 [53], *M. domestica* ‘Golden Delicious’ double haploid v1 (GDDH13) [29], *Pyrus communis* ‘Barlett’ double haploid v2 [54], and our *de novo* assemblies, in addition to the viridiplantae OrthoDB [55]. We filtered the resulting AUGUSTUS [49] output for those that contained full hints (gene model support) and combined the two runs using TSEBRA v1.0.3 [56]. Finally, we removed any transcript/gene that had $\geq 90\%$ softmasking, *i.e.*, mainly repeat sequences. Genome annotation completeness of our genome and other *Malus* genomes were assessed using BUSCO v5.2.2 [40] with the “eudicots_odb10” database for comparative purposes.

The final ‘Honeycrisp’ gene sets from both haplomes were annotated with InterProScan v5.44-79.0 [57,58], including a search against all the available interpro databases and Gene Ontology (GO) [59,60] prediction. In addition, genes were searched against the 26Gv2.0 OrthoFinder v1.1.5 [61] gene family database using both BLASTp [62] and HMMscan [63] classification methods with the *GeneFamilyClassifier* tool from PlantTribes 2 (github.com/dePamphilis/PlantTribes/). This analysis provided additional functional annotation information that includes gene counts of scaffold taxa, superclusters at multiple clustering stringencies, and functional annotations that were pulled from various public genomic databases.

Comparative genomics: Similarities in lengths and structural variations between the two haplomes were determined by running MUMmer v4.0 [64] and Assemblytics [65]. To identify the shared and unique gene families among *Malus* species and cultivars, genes from the six publicly available *Malus* genomes (Table 5) were integrated into the aforementioned PlantTribes 2 gene model database (26Gv2.0) using the same method as described above. The overlapping

orthogroups (with at least 30 counts in the category) among the eight *Malus* annotations (including both haplomes from ‘Honeycrisp’) were calculated and visualized with an upset plot generated by TBtools v1.0986982 [66].

Results

A haplotype-phased chromosome-scale assembly

In total, nearly 55X coverage of PacBio HiFi reads and nearly 200X coverage of Dovetail Omni-C reads (Table 1) was generated. This included 2,543,518 HiFi reads with an average length of 14,655 bp and ~91% of reads $\geq 10,000$ bp. Two phased haplomes, haplome A (HAP1) and haplome B (HAP2, these two sets of terms will be used interchangeably in this manuscript), were assembled and validated by inspection of the Omni-C contact maps (Figure 2). Both haplomes are highly contiguous and of similar size. HAP1 is 674 Mb in length, contained in 473 contigs with a contig N_{50} of 32.8 Mb, whereas HAP2 is 660 Mb in length, contained in 215 contigs with a contig N_{50} of 31.6 Mb (Table 2). Zero miss-joins requiring manual breaks were identified in the assemblies. For HAP1, a total of 13 joins were made to build the final assembly into 17 chromosomes, with 95.4% of the assembled sequence contained in the 17 pseudomolecules representing chromosomes. A total of 19 joins were made for HAP2, with 98.2% of the assembled sequence in the 17 pseudomolecules. Based on the Merqury k-mer analysis (Figure 3), the HAP1 assembly had a k-mer completeness of 82.7% (Quality value (QV) 64.5), the HAP2 assembly 83% (QV 66.7), and the combined assemblies were 98.6% (QV 65.5) (Table 2). BUSCO completeness of HAP1 was 98.6% and HAP2 98.7%, suggesting high genome completeness for both haplomes, comparable or superior to other high quality apple genome assemblies (Table 5). The two haplomes are structurally similar to each other (Figure 4). Compared to the assembly statistics of previously published apple genomes, the current ‘Honeycrisp’ assemblies are the most contiguous to date (Table 5).

Genome annotation

The yield of Illumina transcriptome sequencing data of fruit, leaves, and flower tissues of apples and pear ranged from approximately 9 to 27 gigabases (Gb) in flowers and leaf buds respectively (Table 3). Nearly 62% of both haplomes were annotated as repetitive DNA, mostly comprised of Long Terminal Repeat (LTR) retrotransposons (Table 4). A total of 47,563 genes

were annotated in HAP1 and 48,655 in HAP2, slightly more than in other published *Malus* annotations (Table 5). Complete BUSCO scores of the protein annotations are 96.8% for HAP1 and 97.4% for HAP2, the highest completeness among all publicly available *Malus* genome annotations (Table 5). 72.85% and 68.88% of the predicted transcripts were annotated with Interpro terms, 68.58% and 64.94% with Pfam domains, and 51.04% and 48.76% with at least one GO terms in HAP1 and HAP2, respectively. In the PlantTribes 2 classification, 91.11% and 85.50% of the predicted transcripts from HAP1 and HAP2, respectively, were assigned to pre-computed orthogroups.

As the number of plant genomes are being generated at an unprecedented speed, we developed the following gene naming convention to avoid potential ambiguity. Maldo.hc.v1a1.ch10A.g00001.t1: Maldo for *Malus domestica*; hc for the cultivar, ‘Honeycrisp’; v1a1 indicating this is the first assembly and first annotation of this genome; ch10A identifies that the gene is annotated from chromosome 10 (versus from an unplaced scaffold, which will be indicated by “sc”) in haplome A (HAP1) (versus haplome B (HAP2)); g00001 is a five digit gene identifier; t1 represents a transcript number of the gene.

Gene family analysis

A gene family evaluation was performed using PlantTribes 2 and its 26Gv2-scaffold orthogroup database, which contains representative protein coding sequences from most major land plant lineages. A total of 11,263 unique orthogroups (OGs) were identified in all eight *Malus* annotations (including the two ‘Honeycrisp’ haplomes) investigated. ‘Honeycrisp’ transcripts were assigned to 10,351 and 10,367 orthogroups, similar to ‘Gala’ and GDDH13 (Table 5 and Figure 5). We further investigated orthogroups that are shared and unique in the eight *Malus* annotations. A vast majority (7,645) of orthogroups are shared by all the genomes, and a total of 9,279 orthogroups were shared among both ‘Honeycrisp’ haplomes and five other genomes (Figure 5). This comparison indicates that the ‘Honeycrisp’ annotation captured genes in virtually all the *Malus* gene families. In addition, we also found 54 orthogroups that are unique to ‘Honeycrisp’ (*i.e.*, shared by the two ‘Honeycrisp’ haplomes only) and 35 and 32 that are unique to each ‘Honeycrisp’ haplome (Figure 5). These orthogroups could provide valuable information in the molecular mechanisms underlying genotype-specific traits.

Re-use Potential

This fully phased, high-quality, chromosome-scale genome of ‘Honeycrisp’ apple will add to the toolbox for apple genetic research and breeding. It will enable genetic mapping, identification of genes, and development of molecular markers linked to disease, pest resistance, abiotic stress tolerance and adaptation, as well as horticulturally relevant harvest and postharvest fruit quality traits for use in apple breeding programs. Ultimately, the addition of high-quality genomic resources for ‘Honeycrisp’ can lead to enhanced orchard and supply chain management for many other apple cultivars, promoting future sustainability of the pome fruit industry.

Data Availability

The whole genome sequence data generated in this study have been deposited at the National Center for Biotechnology Information (NCBI) database under BioProject ID PRJNA791346. PacBio HiFi reads, and Hi-C reads are deposited in NCBI with the SRA accession number SAMN24287034 and SAMN29611953, respectively. Transcriptomic data generated in this study for genome annotation are deposited in NCBI with SRA accession number from SAMN29611954 to SAMN29611992. The Maldo.hc.v1a1 ‘Honeycrisp’ genome assembly, gene annotation, and functional annotation for both haplotypes can be accessed *via* the Genomic Database for Rosaceae (in progress) and the GigaScience GigaDB repository.

Declarations

List of abbreviations

BLAST: Basic Local Alignment Search Tool; bp: Base Pair; BUSCO: Benchmarking Universal Single Copy Orthologs; Gb: gigabases; GO: Gene Ontology; HMW: High-molecular-weight; JBAT: Juicebox Assembly Tools; LTR: Long Terminal Repeat; NCBI: National Centre for Biotechnology Information; OG: Orthogroup; QV: Quality Value; RIN: RNA Integrity Number; SMRT: Single Molecule Real Time; TRF: Tandem Repeat Finder

Competing interests

The authors declare that they have no competing interests.

Funding

This research was funded by Washington Tree Fruit Research Commission AP-19-103, United States Department of Agriculture, Agricultural Research Service and New York State's Apple Research and Development Program (ARDP).

Author Contribution

A.K., and L.H. conceptualized, designed, and managed the project. S.B.C., H.Z., A.H., and H.H, constructed DNA and RNA, and RNA-seq libraries for sequencing. S.B.C., and H.Z., performed genome and transcriptome sequence analysis and interpretation. S.B.C., H.Z., A.S., H.H., A.H., L.H., and A.K. drafted, revised, and finalized the manuscript. All authors read and approved the manuscript.

Acknowledgements

We acknowledge the help of Della Cobb-Smith and Jugpreet Singh for sample collection and preparation for genomic DNA extraction.

Figures and Tables

Figure 1: ‘Honeycrisp’ is a highly desirable apple cultivar, appreciated by consumers for its superior flavor, texture, and visual appeal (A). To fulfill high-market demand for ‘Honeycrisp’, growers need to optimize production traits, and address several diseases and physiological disorders of concern to ensure high levels of production and fruit quality, including zonal leaf chlorosis (B), fungal diseases like the bitter rot pathogen complex (*Colletotrichum gloeosporioides* and *C. acutatum*) (C) the black rot pathogen (*Botryosphaeria obtuse*) (D), as well as postharvest storage disorders like bitter pit (E) soft scald (F), and soggy breakdown (G).

Figure 2: Omni-C contact maps of the assembled chromosome-length scaffolds of 17 chromosomes of (A) Haplome A and (B) Haplome B, from the ‘Honeycrisp’ genome.

Figure 3: Histogram of k-mer multiplicity of sequence reads for (A) Haplome A and, (B) Haplome B of ‘Honeycrisp’ genome assemblies. K-mer multiplicity (x-axis) is plotted against k-mer counts (y-axis) to estimate the heterozygosity, copy numbers, sequencing depth, and completeness of a genome using Merquy v1.3 [41]. Colors in the plot represent the number of times each k-mer is found in the genome assembly.

Figure 4: Synteny comparison of ‘Honeycrisp’ Haplome 1 (HAP1), ‘Honeycrisp’ Haplome 2 (HAP2) from this study, and ‘Gala’ [27] genomes. GENESPACE [39] was used for synteny comparison.

Figure 5: The Honeycrisp genome captured a vast majority of *Malus* gene families. Black dots indicate presence of gene families and gray dots indicate absence. Yellow horizontal bars represent the number of orthogroups in each genome. The black vertical bars represent the number of orthogroups in each category. Genome abbreviations - HC: ‘Honeycrisp’ (this work); GDDH13: *Malus domestica* GDDH13; Gala_hap: *M. domestica* ‘Gala’ haploid; M.si_hap: *M. sieversii* haploid; M.sy_hap: *M. sylvestris* haploid; HFTH: *M. domestica* HFTH1; GDv1: *M. domestica* Golden Delicious v1.

Table 1: Overview of PacBio HiFi and Omni-C sequencing data generated for the ‘Honeycrisp’ genome assembly.

Library	Sequencing	Length (Nucleotides)	Number of reads
JNQN	Omni-C	150	951,241,272
HiFi-1	PacBio HiFi	14,881*	1,088,992
HiFi-2	PacBio HiFi	14,429*	1,454,526

* Average length

Table 2: Summary of ‘Honeycrisp’ genome assembly statistics.

Assembly	Length	# Contigs	Longest contig	N50	L50	QV	k-mer completeness (%)	BUSCO (%)
Honeycrisp Haplome A	674,476,353	473	55,653,390	32,818,622	9	64.5	82.7	98.6
Honeycrisp Haplome B	660,238,068	215	56,154,892	31,578,807	9	66.7	83	98.7
Combined						65.5	98.6	

Table 3: Yield of Illumina transcriptome sequencing of fruit, leaves, and flower tissues of apples and pear generated and used for genome annotation in this study.

Cultivar	Tissue	Reads	Yield (Gb)	Yield P20 (Gb)	Ave Read length	NCBI SRA
Honeycrisp	Fruitlet stage 1	45,773,784	13,823,682,768	13,069,822,280	142	SAMN29611971
	Fruitlet stage 2	35,618,706	10,756,849,212	10,227,275,771	143	SAMN29611972
	Budding leaves	81,448,971	24,597,589,242	22,769,634,770	139	SAMN29611973
	Expanding leaves	35,381,039	10,685,073,778	9,971,308,535	141	SAMN29611974
	Half-inch terminal buds	47,811,924	14,439,201,048	13,409,542,519	140	SAMN29611975
	Flower buds	45,822,773	13,838,477,446	13,175,876,315	144	SAMN29611976
	Open flowers	30,938,395	9,343,395,290	8,718,474,885	141	SAMN29611977
Gala	Fruitlet stage 1	80,440,219	24,292,946,138	22,928,129,883	142	SAMN29611954
	Fruitlet stage 2	32,475,136	9,807,491,072	9,284,944,973	143	SAMN29611955
	Budding leaves	30,368,057	9,171,153,214	8,508,033,713	140	SAMN29611956
	Expanding leaves	40,650,277	12,276,383,654	11,306,267,120	138	SAMN29611957
	Roots from tissue culture	35,324,786	10,668,085,372	9,940,132,737	140	SAMN29611958
	Quarter-inch terminal buds	37,532,631	11,334,854,562	10,634,379,784	141	SAMN29611959
	Flower buds	39,636,821	11,970,319,942	11,141,652,382	140	SAMN29611960
	Open flowers	34,363,075	10,377,648,650	9,775,838,818	142	SAMN29611961
Red Delicious	Fruitlet stage 2	27,319,955	8,250,626,410	7,682,200,349	140	SAMN29611962
Granny Smith	Fruitlet stage 1	29,426,606	8,886,835,012	8,335,731,187	141	SAMN29611963
	Fruitlet stage 2	72,205,133	21,805,950,166	20,663,261,900	143	SAMN29611964
	Budding leaves	57,244,195	17,287,746,890	16,179,280,911	141	SAMN29611965
	Expanding leaves	40,798,422	12,321,123,444	11,499,303,808	140	SAMN29611966
	Roots from tissue culture	32,493,822	9,813,134,244	9,207,784,729	141	SAMN29611967
	Quarter-inch terminal buds	30,394,263	9,179,067,426	8,512,945,196	140	SAMN29611968
	Flower buds	29,735,514	8,980,125,228	8,364,532,017	140	SAMN29611969
	Open flowers	34,303,317	10,359,601,734	9,603,420,430	140	SAMN29611970
WA 38	Fruitlet stage 1	45,284,208	13,675,830,816	12,831,991,620	141	SAMN29611978

	Fruitlet stage 2	25,486,256	7,696,849,312	7,261,195,330	142	SAMN29611979
	Budding leaves	39,339,589	11,880,555,878	11,017,185,994	140	SAMN29611980
	Expanding leaves	34,784,980	10,505,063,960	9,719,694,010	139	SAMN29611981
	Roots from tissue culture	33,935,508	10,248,523,416	9,426,506,860	138	SAMN29611982
	Quarter-inch terminal buds	88,677,165	26,780,503,830	24,913,194,030	140	SAMN29611983
	Flower buds	23,170,354	6,997,446,908	6,588,921,074	142	SAMN29611984
	Open flowers	35,274,250	10,652,823,500	9,941,466,644	141	SAMN29611985
d'Anjou	Fruitlet stage 1	89,462,306	27,017,616,412	25,459,693,894	142	SAMN29611986
	Fruitlet stage 2	48,481,031	14,641,271,362	13,921,844,851	143	SAMN29611987
	Budding leaves	29,823,484	9,006,692,168	8,442,259,663	141	SAMN29611988
	Expanding leaves	57,920,009	17,491,842,718	16,460,531,509	142	SAMN29611989
	Quarter-inch terminal buds	40,966,825	12,371,981,150	11,476,090,088	140	SAMN29611990
	Flower buds	29,183,231	8,813,335,762	8,264,473,671	141	SAMN29611991
	Open flowers	32,128,369	9,702,767,438	8,996,878,963	140	SAMN29611992

Table 4: Summary of repetitive element annotation in Haplome A and Haplome B of the ‘Honeycrisp’ genome assemblies.

Class		Haplome A	Haplome B
LTR			
	Copia	9.73%	9.60%
	Ty3	20.29%	17.80%
	unknown	14.89%	16.86%
TIR			
	CACTA	2.21%	1.95%
	Mutator	4.16%	4.25%
	PIF Harbinger	2.43%	2.60%
	Tc1_Mariner	0.15%	0.27%
	hAT	2.30%	2.31%
	polinton	--	0.01%
nonLTR			
	LINE_element	0.18%	0.17%
	unknown	0.09%	0.18%
nonTIR			
	helitron	2.95%	3.18%
repeat region		2.91%	2.78%
Total		62.43%	61.97%

Table 5: Comparison of genomic features and assembly statistics of current assembly of ‘Honeycrisp’ genome and previously published genomes of apples.

Genomes	‘Honeycrisp’	‘Gala’, <i>M. sieversii</i> , <i>M. sylvestris</i> (all Diploid)	HFTH1; ‘Hanfu’ (Triploid)	GDDH13; ‘Golden Delicious’ (Double haploid)	‘Golden Delicious’ (Diploid)
Reference	This work	Sun et al. 2021	Zhang et. 2019	Daccord et al. 2017	Velasco et al. 2010
Assembly					
Haploid genome size (Mb)	660-674	666-679	658.9	651	742
scaffold N50	31.6-32.8	6.1-21.8	6.99	5.5	16Kb
Complete BUSCO	98.6-98.7%	98.0-98.8%	98.6%	98.0%	82.0%

Annotation					
Protein-coding genes	47,563-48,655	44,691-44,847	44,677	42,140	57,386
Complete BUSCO	96.8-97.4%	94.6-95.4%	93.6%	96.1%	68%
Gene family					
Number of OG in 26Gv2	10,351-10,367	10,044-10,115	9,974	10,117	8,824

References

1. WSTFA Annual Crop Summary 2017-18: <https://wstfa.org/wstfa-assets/uploads/2017-18-Annual-Growers-Review.pdf>.
2. Yan D, Shi J, Ren X, Tao Y, Ma F, Li R, et al.. Insights into the aroma profiles and characteristic aroma of ‘Honeycrisp’ apple (*Malus × domestica*). *Food Chemistry*. 2020; doi: 10.1016/j.foodchem.2020.127074.
3. Chang H-Y, Vickers ZM, Tong CBS. The use of a combination of instrumental methods to assess change in sensory crispness during storage of a “Honeycrisp” apple breeding family. *Journal of Texture Studies*. 2018; doi: 10.1111/jtxs.12325.
4. Cline JA, Neilsen D, Brownlee RA, Norton D, Quamme HA. Cold Hardiness of New Apple Cultivars of Commercial Importance in Canada. *Journal of the American Pomological Society*. 2012.
5. Clark M, Luby J, Bradeen J, Bus V. Identification of Candidate Genes at Rvi19 and Rvi20, Two Apple Scab Resistance Loci in the ‘Honeycrisp’ Apple (*Malus x domestica*).
6. Luby JJ, Bedford DS. Honeycrisp Apple. Minnesota Agricultural Experiment Station; 1992.
7. Howard NP, van de Weg E, Bedford DS, Peace CP, Vanderzande S, Clark MD, et al.. Elucidation of the ‘Honeycrisp’ pedigree through haplotype analysis with a multi-family integrated SNP linkage map and a large apple (*Malus×domestica*) pedigree-connected SNP data set. *Hortic Res*. Nature Publishing Group; 2017; doi: 10.1038/hortres.2017.3.
8. Cabe PR, Baumgarten A, Onan K, Luby JJ, Bedford DS. Using Microsatellite Analysis to Verify Breeding Records: A study of ‘Honeycrisp’ and Other Cold-hardy Apple Cultivars. *HortScience*. American Society for Horticultural Science; 2005; doi: 10.21273/HORTSCI.40.1.15.
9. Teh SL, Kostick S, Brutcher L, Schonberg B, Barritt B, Evans K. Trends in Fruit Quality Improvement From 15 Years of Selection in the Apple Breeding Program of Washington State University. *Frontiers in Plant Science*. 122021.
10. Migicovsky Z, Gardner KM, Richards C, Thomas Chao C, Schwaninger HR, Fazio G, et al.. Genomic consequences of apple improvement. *Hortic Res*. 2021; doi: 10.1038/s41438-020-00441-7.
11. Papp D, Gao L, Thapa R, Olmstead D, Khan A. Field apple scab susceptibility of a diverse *Malus* germplasm collection identifies potential sources of resistance for apple breeding. *CABI Agriculture and Bioscience*. 2020; doi: 10.1186/s43170-020-00017-4.
12. Papp D, Singh J, Gadoury D, Khan A. New North American Isolates of *Venturia inaequalis* Can Overcome Apple Scab Resistance of *Malus floribunda* 821. *Plant Disease*. Scientific Societies; 2020; doi: 10.1094/PDIS-10-19-2082-RE.

- 373 13. Tong C, Krueger D, Vickers Z, Bedford D, Luby J, El-Shiekh A, et al.. Comparison of
374 Softening-related Changes during Storage of 'Honeycrisp' Apple, Its Parents, and 'Delicious'.
375 *Journal of the American Society for Horticultural Science*. American Society for Horticultural
376 Science; 1999; doi: 10.21273/JASHS.124.4.407.
- 377 14. Howard NP, Tillman J, Vanderzande S, Luby JJ. Genetics of zonal leaf chlorosis and genetic
378 linkage to a major gene regulating skin anthocyanin production (MdMYB1) in the apple (*Malus*
379 $\times$ *domestica*) cultivar Honeycrisp. *PLOS ONE*. Public Library of Science; 2019; doi:
380 10.1371/journal.pone.0210611.
- 381 15. Brooks C, Harley CP. Soft scald and soggy breakdown of apples. *J Agric Res*; 1934.
- 382 16. Rosenberger D, Schupp J, Iungerman K, Hoying S, Straub D, Cheng L. Honeycrisp:
383 Promising Profit Maker or Just Another Problem Child? *Fruit Quarterly*; 2001.
- 384 17. Rosenberger D, Cox KD. Preventing bitter rot in apples. *Scaffolds Fruit*; 2016.18. Contreras
385 C, Alsmairat N, Beaudry R. Prestorage Conditioning and Diphenylamine Improve Resistance to
386 Controlled-atmosphere-related Injury in 'Honeycrisp' Apples. *HortScience*. American Society
387 for Horticultural Science; 2014; doi: 10.21273/HORTSCI.49.1.76.
- 388 19. DeBrouwer EJ, Sriskantharajah K, El Kayal W, Sullivan JA, Paliyath G, Subramanian J. Pre-
389 harvest hexanal spray reduces bitter pit and enhances post-harvest quality in 'Honeycrisp' apples
390 (*Malus domestica* Borkh.). *Scientia Horticulturae*. 2020; doi: 10.1016/j.scienta.2020.109610.
- 391 20. Mattheis JP, Rudell DR, Hanrahan I. Impacts of 1-Methylcyclopropene and Controlled
392 Atmosphere Established during Conditioning on Development of Bitter Pit in 'Honeycrisp'
393 Apples. *HortScience*. American Society for Horticultural Science; 2017; doi:
394 10.21273/HORTSCI11368-16.
- 395 21. Watkins CB, Nock JF, Weis SA, Jayanty S, Beaudry RM. Storage temperature,
396 diphenylamine, and pre-storage delay effects on soft scald, soggy breakdown and bitter pit of
397 'Honeycrisp' apples. *Postharvest Biology and Technology*. 2004; doi:
398 10.1016/j.postharvbio.2003.11.003.
- 399 22. Watkins CB, Mattheis JP. "Apple" in Postharvest Physiological Disorders in Fruits and
400 Vegetables. Edited by Sergio Tonetto de Freitas and Sunil Pareek. Postharvest Physiological
401 Disorders in Fruits and Vegetables. Boca Raton: CRC Press; 2019; doi: 10.1201/b22001.
- 402 23. Varshney RK, Bohra A, Yu J, Graner A, Zhang Q, Sorrells ME. Designing Future Crops:
403 Genomics-Assisted Breeding Comes of Age. *Trends in Plant Science*. 2021; doi:
404 10.1016/j.tplants.2021.03.010.
- 405 24. Velasco R, Zharkikh A, Affourtit J, Dhingra A, Cestaro A, Kalyanaraman A, et al.. The
406 genome of the domesticated apple (*Malus* $\times$ *domestica* Borkh.). *Nat Genet*. 2010; doi:
407 10.1038/ng.654.
- 408 25. Honaas LA, Hargarten HL, Ficklin SP, Hadish JA, Wafula E, dePamphilis CW, et al.. Co-
409 expression networks provide insights into molecular mechanisms of postharvest temperature

410 modulation of apple fruit to reduce superficial scald. *Postharvest Biology and Technology*. 2019;
411 doi: 10.1016/j.postharvbio.2018.09.016.

412 26. Khan A, Schuyler SK. Breeding and genetics of disease resistance in temperate fruit trees:
413 challenges and new opportunities. *Theor Appl Genet*. 2022; doi: 10.1007/s00122-022-04093-0.

414 27. Sun X, Jiao C, Schwaninger H, Chao CT, Ma Y, Duan N, et al.. Phased diploid genome
415 assemblies and pan-genomes provide insights into the genetic history of apple domestication.
416 *Nat Genet*. Nature Publishing Group; 2020; doi: 10.1038/s41588-020-00723-9.

417 28. Zhang L, Hu J, Han X, Li J, Gao Y, Richards CM, et al.. A high-quality apple genome
418 assembly reveals the association of a retrotransposon and red fruit colour. *Nat Commun*. Nature
419 Publishing Group; 2019; doi: 10.1038/s41467-019-09518-x.

420 29. Daccord N, Celton J-M, Linsmith G, Becker C, Choisne N, Schijlen E, et al.. High-quality de
421 novo assembly of the apple genome and methylome dynamics of early fruit development. *Nat*
422 *Genet*. Nature Publishing Group; 2017; doi: 10.1038/ng.3886.

423 30. Singh J, Sun M, Cannon SB, Wu J, Khan A. An accumulation of genetic variation and
424 selection across the disease-related genes during apple domestication. *Tree Genetics & Genomes*.
425 2021; doi: 10.1007/s11295-021-01510-1.

426 31. Marçais G, Kingsford C. A fast, lock-free approach for efficient parallel counting of
427 occurrences of k-mers. *Bioinformatics*. 2011; doi: 10.1093/bioinformatics/btr011.

428 32. Ranallo-Benavidez TR, Jaron KS, Schatz MC. GenomeScope 2.0 and Smudgeplot for
429 reference-free profiling of polyploid genomes. *Nat Commun*. Nature Publishing Group; 2020;
430 doi: 10.1038/s41467-020-14998-3.

431 33. Vulture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, et al..
432 GenomeScope: fast reference-free genome profiling from short reads. *Bioinformatics*. 2017; doi:
433 10.1093/bioinformatics/btx153.

434 34. Cheng H, Jarvis ED, Fedrigo O, Koepfli K-P, Urban L, Gemmell NJ, et al.. Haplotype-
435 resolved assembly of diploid genomes without parental data. *Nat Biotechnol*. Nature Publishing
436 Group; 2022; doi: 10.1038/s41587-022-01261-x.

437 35. Cheng H, Concepcion GT, Feng X, Zhang H, Li H. Haplotype-resolved de novo assembly
438 using phased assembly graphs with hifiasm. *Nat Methods*. Nature Publishing Group; 2021; doi:
439 10.1038/s41592-020-01056-5.

440 36. Durand NC, Shamim MS, Machol I, Rao SSP, Huntley MH, Lander ES, et al.. Juicer
441 Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. *cels*. Elsevier;
442 2016; doi: 10.1016/j.cels.2016.07.002.

443 37. Li X, Kui L, Zhang J, Xie Y, Wang L, Yan Y, et al.. Improved hybrid de novo genome
444 assembly of domesticated apple (*Malus x domestica*). *GigaScience*. 2016; doi: 10.1186/s13742-
445 016-0139-0.

38. Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, et al.. De novo assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. *Science*. American Association for the Advancement of Science; 2017; doi: 10.1126/science.aal3327.
39. Lovell J, Sreedasyam A, Schranz E, Wilson M, Carlson J, Harkess A, et al.. GENESPACE: syntenic pan-genome annotations for eukaryote.
40. Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology and Evolution*. 2021; doi: 10.1093/molbev/msab199.
41. Rhie A, Walenz BP, Koren S, Phillippy AM. Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. *Genome Biology*. 2020; doi: 10.1186/s13059-020-02134-9.
42. Honaas L, Kahn E. A practical examination of RNA isolation methods for European pear (*Pyrus communis*). *BMC Research Notes*. 2017; doi: 10.1186/s13104-017-2564-2.
43. Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, et al.. Benchmarking transposable element annotation methods for creation of a streamlined, comprehensive pipeline. *Genome Biology*. 2019; doi: 10.1186/s13059-019-1905-y.
44. Pratas MI, Aguiar B, Vieira J, Nunes V, Teixeira V, Fonseca NA, et al.. Inferences on specificity recognition at the *Malus×domestica* gametophytic self-incompatibility system. *Sci Rep*. 2018; doi: 10.1038/s41598-018-19820-1.
45. Galimba KD, Bullock DG, Dardick C, Liu Z, Callahan AM. Gibberellic acid induced parthenocarpic ‘Honeycrisp’ apples (*Malus domestica*) exhibit reduced ovary width and lower acidity. *Hortic Res*. Nature Publishing Group; 2019; doi: 10.1038/s41438-019-0124-8.
46. Chang H-Y, Tong CBS. Identification of Candidate Genes Involved in Fruit Ripening and Crispness Retention Through Transcriptome Analyses of a ‘Honeycrisp’ Population. *Plants*. Multidisciplinary Digital Publishing Institute; 2020; doi: 10.3390/plants9101335.
47. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al.. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*. Nature Publishing Group; 2011; doi: 10.1038/nbt.1883.
48. Haas B, Papanicolaou A and others. TransDecoder (find Coding Regions within Transcripts). 2015; <https://github.com/TransDecoder/transdecoder.github.io>.
49. Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. BRAKER1: Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS. *Bioinformatics*. 2016; doi: 10.1093/bioinformatics/btv661.
50. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 2014; doi: 10.1093/bioinformatics/btu170.

482 51. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al.. STAR: ultrafast
483 universal RNA-seq aligner. *Bioinformatics*. 2013; doi: 10.1093/bioinformatics/bts635.

484 52. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al.. The Sequence
485 Alignment/Map format and SAMtools. *Bioinformatics*. 2009; doi:
486 10.1093/bioinformatics/btp352.

487 53. Chen X, Li S, Zhang D, Han M, Jin X, Zhao C, et al.. Sequencing of a Wild Apple (*Malus*
488 *baccata*) Genome Unravels the Differences Between Cultivated and Wild Apple Species
489 Regarding Disease Resistance and Cold Tolerance. *G3 (Bethesda)*. 2019; doi:
490 10.1534/g3.119.400245.

491 54. Linsmith G, Rombauts S, Montanari S, Deng CH, Celton J-M, Guérif P, et al.. Pseudo-
492 chromosome-length genome assembly of a double haploid “Bartlett” pear (*Pyrus communis* L.).
493 *GigaScience*. 2019; doi: 10.1093/gigascience/giz138.

494 55. Kriventseva EV, Kuznetsov D, Tegenfeldt F, Manni M, Dias R, Simão FA, Zdobnov EM.
495 OrthoDB v10: sampling the diversity of animal, plant, fungal, protist, bacterial and viral
496 genomes for evolutionary and functional annotations of orthologs. *Nucleic Acids Research*.
497 2019; doi: 10.1093/nar/gky1053.
498

499 56. Gabriel L, Hoff KJ, Brûna T, Borodovsky M, Stanke M. TSEBRA: transcript selector for
500 BRAKER. *BMC Bioinformatics*. 2021; doi: 10.1186/s12859-021-04482-0.

501 57. Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al.. The
502 InterPro protein families and domains database: 20 years on. *Nucleic Acids Res*. 2021; doi:
503 10.1093/nar/gkaa977.

504 58. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al.. InterProScan 5: genome-
505 scale protein function classification. *Bioinformatics*. 2014; doi: 10.1093/bioinformatics/btu031.

506 59. The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing
507 strong. *Nucleic Acids Res*. 2019; doi: 10.1093/nar/gky1055.


508 60. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al.. Gene Ontology:
509 tool for the unification of biology. *Nat Genet*. Nature Publishing Group; 2000; doi:
510 10.1038/75556.



511 61. Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome comparisons
512 dramatically improves orthogroup inference accuracy. *Genome Biology*. 2015; doi:
513 10.1186/s13059-015-0721-2.


514 62. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.. BLAST+:
515 architecture and applications. *BMC Bioinformatics*. 2009; doi: 10.1186/1471-2105-10-421.

516 63. Eddy SR. Accelerated Profile HMM Searches. *PLOS Computational Biology*. Public Library
517 of Science; 2011; doi: 10.1371/journal.pcbi.1002195.

- 518 64. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al.. Versatile and
519 open software for comparing large genomes. *Genome Biology*. 2004; doi: 10.1186/gb-2004-5-2-
520 r12.
- 521 65. Nattestad M, Schatz MC. Assemblytics: a web analytics tool for the detection of variants
522 from an assembly. *Bioinformatics*. 2016; doi: 10.1093/bioinformatics/btw369.
- 523 66. Chen C, Chen H, Zhang Y, Thomas HR, Frank MH, He Y, et al.. TBtools: An Integrative
524 Toolkit Developed for Interactive Analyses of Big Biological Data. *Mol Plant*. 2020; doi:
525 10.1016/j.molp.2020.06.009.

A**B****C****D****E****F****G**



(A)**(B)**