

## Faster detection of “darks” than “brights” by monkey superior colliculus neurons

Tatiana Malevich<sup>1, 2\*</sup>, Tong Zhang<sup>1, 2\*</sup>, Matthias P. Baumann<sup>1, 2\*</sup>, Amarender R. Bogadhi<sup>1, 2, 3</sup>,  
and Ziad M. Hafed<sup>1, 2</sup>

1. Werner Reichardt Center for Integrative Neuroscience,  
University of Tübingen, Tübingen, Germany 72076
2. Hertie Institute for Clinical Brain Research,  
University of Tübingen, Tübingen, Germany 72076
3. Central Nervous System Diseases Research, Boehringer Ingelheim Pharma GmbH & Co. KG,  
Biberach, Riß, Germany 88400

\* Co-first authors

### **Abbreviated title:**

## Visual processing of darks by superior colliculus

## Correspondence to:

Tatiana Malevich ([tatiana.malevich@cin.uni-tuebingen.de](mailto:tatiana.malevich@cin.uni-tuebingen.de))  
and

Ziad M. Hafed ([ziad.m.hafed@cin.uni-tuebingen.de](mailto:ziad.m.hafed@cin.uni-tuebingen.de))  
Werner Reichardt Center for Integrative Neuroscience  
Otfried-Müller Str. 25  
Tübingen, Germany, 72076

**Number of pages:** 53

**Number of figures:** 11

**Number of words in Abstract:** 247

Number of words in Introduction: 661

Number of words in Discussion: 143

**Conflict of interest statement:** The authors declare no competing interests.

**Acknowledgements:** We were funded by the Deutsche Forschungsgemeinschaft (DFG): (1) BO5681/1-1; (2) EXC307; (3) HA6749/3-1; (4) HA6749/4-1; and (5) SFB 1233, Robust Vision: Inference Principles and Neural Mechanisms, TP 11, project number: 276693517.

46 **Abstract**

47

48 Visual processing is segregated into ON and OFF channels as early as in the retina, and the  
49 superficial (output) layers of the primary visual cortex are dominated by neurons preferring  
50 dark stimuli. However, it is not clear how the timing of neural processing differs between  
51 “darks” and “brights” in general, especially in light of psychophysical evidence; it is also  
52 equally not clear how subcortical visual pathways that are critical for active orienting  
53 represent stimuli of positive (luminance increments) and negative (luminance decrements)  
54 contrast polarity. Here, we recorded from all visually-responsive neuron types in the  
55 superior colliculus (SC) of two male rhesus macaque monkeys. We presented a disc (0.51 deg  
56 radius) within the response fields (RF's) of neurons, and we varied, across trials, stimulus  
57 Weber contrast relative to a gray background. We also varied contrast polarity. There was a  
58 large diversity of preferences for darks and brights across the population. However,  
59 regardless of individual neural sensitivity, most neurons responded significantly earlier to  
60 dark than bright stimuli. This resulted in a dissociation between neural preference and visual  
61 response onset latency: a neuron could exhibit a weaker response to a dark stimulus than to  
62 a bright stimulus of the same contrast, but it would still have an earlier response to the dark  
63 stimulus. Our results highlight an additional candidate visual neural pathway for explaining  
64 behavioral differences between the processing of darks and brights, and they demonstrate  
65 the importance of temporal aspects in the visual neural code for orienting eye movements.

66

67

68

69 **Significance statement**

70

71 Objects in our environment, such as birds flying across a bright sky, often project shadows  
72 (or images darker than the surround) on our retina. We studied how primate superior  
73 colliculus (SC) neurons visually process such dark stimuli. We found that the overall  
74 population of SC neurons represented both dark and bright stimuli equally well, as  
75 evidenced by a relatively equal distribution of neurons that were either more or less  
76 sensitive to darks. However, independent of sensitivity, the great majority of neurons  
77 detected dark stimuli earlier than bright stimuli, evidenced by a smaller response latency for  
78 the dark stimuli. Thus, SC neural response latency can be dissociated from response  
79 sensitivity, and it favors the faster detection of dark image contrasts.

80

81

82

83 **Introduction**

84

85 Early visual processing is segregated into parallel pathways conveying information about  
86 either luminance increments or decrements in visual scenes (Hartline, 1938; Schiller et al.,  
87 1986). Such segregation starts in the retina and persists in the early retino-geniculate visual  
88 pathway (Hubel and Wiesel, 1961; Schiller et al., 1986). Interestingly, such segregation is also  
89 accompanied by asymmetries with which dark and bright stimuli are processed. For  
90 example, primate retinal ganglion cells possess asymmetric spatial and temporal properties  
91 depending on whether they are part of the ON or OFF pathway (Chichilnisky and Kalmar,  
92 2002). Similarly, in the primate lateral geniculate nucleus (LGN), neurons with OFF-center  
93 response fields (RF's) are more sensitive to their preferred stimuli (dark contrasts) than  
94 neurons with ON-center RF's experiencing bright contrasts (Jiang et al., 2015). OFF-center  
95 neurons also have higher spontaneous activity and more sustained responses during visual  
96 stimulation (Jiang et al., 2015). Ultimately, signals reach the primary visual cortex (V1),  
97 where ON/OFF asymmetries are amplified. For example, primate V1 is dominated by "black"  
98 responses, especially in the superficial cortico-cortical output layers (Yeh et al., 2009).

99

100 Asymmetries in the processing of dark versus bright stimuli might make ecological sense. For  
101 example, the incidence of dark contrasts in natural scenes is not necessarily uniform.  
102 Instead, there is a coincidence of dark contrasts with regions of low spatial frequency, high  
103 contrast, and far binocular disparities in natural images (Cooper and Norcia, 2015). As a  
104 result, rhesus macaque V1 neurons having far preferred binocular disparities tend to also  
105 prefer dark contrasts (Samonds et al., 2012). Similarly, in cat V1, there is a systematic  
106 contrast-dependent OFF-dominance, matching natural scene statistics (Liu and Yao, 2014),

107 and cat V1 neurons are more strongly driven by luminance decrements than increments at  
108 low spatial frequencies (Kremkow et al., 2014). Interestingly, cat studies revealed that ON  
109 and OFF domains in the LGN also exist in the V1 projections (Jin et al., 2008), with area  
110 centralis representations being dominated by dark preferences. Moreover, OFF-dominated  
111 LGN (Jin et al., 2011) and V1 (Komban et al., 2014) neurons respond earlier than ON-  
112 dominated ones. These last observations on OFF and ON channel timing are consistent with  
113 a large body of psychophysical literature for better and faster processing of dark stimuli (e.g.  
114 Komban et al., 2011; Komban et al., 2014).

115  
116 Having said that, whether monkey superior colliculus (SC) neurons differentially process dark  
117 stimuli remains unclear. In the mouse SC, the majority of superficial layer neurons prefer  
118 dark stimuli (De Franceschi and Solomon, 2018), consistent with the RF subfield structure of  
119 these neurons (Wang et al., 2010). Yet, it is not clear whether such dark preference still  
120 exists in the deeper SC layers, and whether it is accompanied by differences in visual  
121 response latencies. Moreover, differences in the ecological environments and  
122 neuroanatomical organizations of mice and other species do not trivially predict how  
123 primate SC neurons might behave with respect to luminance contrast polarity. Therefore, we  
124 exhaustively characterized all visually-responsive rhesus macaque monkey SC neurons (that  
125 is, also including intermediate and deeper layer neurons). We were particularly motivated by  
126 our recent observations of differential effects of contrast polarity on microsaccades  
127 (Malevich et al., 2021).

128  
129 In contrast to LGN, V1, and SC results from other species, we did not find a dominant  
130 preference for dark stimuli in the primate SC. Rather, there was significant diversity, with

131 approximately half of the neurons being more sensitive to bright stimuli. Moreover, at high  
132 contrasts, SC neurons tended to prefer bright rather than dark stimuli, with this trend  
133 disappearing for the lowest contrasts. Despite such diversity, what we did find was that the  
134 majority of SC neurons had significantly shorter visual response latencies to dark stimuli.  
135 Thus, there was a dissociation between visual response latency and visual response  
136 sensitivity, reminiscent of a similar dissociation that we observed in the case of spatial  
137 frequency tuning (Chen et al., 2018). Such a dissociation was sufficient to account for at least  
138 some saccadic reaction time dependencies on stimulus luminance polarity in our  
139 experiments.

140

141 **Materials and methods**

142

143

144 *Experimental animals and ethics approvals*

145 We recorded superior colliculus (SC) neural activity from two adult, male rhesus macaque

146 monkeys (M and A) aged 9 and 10 years, and weighing 9.5 kg and 10 kg, respectively. We

147 also measured saccadic reaction times from the same two animals plus a third one (F; aged

148 11 years and weighing 14 kg). The experiments were approved by ethics committees at the

149 regional governmental offices of the city of Tübingen.

150

151

152 *Laboratory setup and animal preparation*

153 The experiments were conducted in the same laboratory as that described in our recent

154 studies (Bogadhi et al., 2020; Bogadhi and Hafed, 2022). Briefly, the monkeys were seated in

155 a darkened booth approximately 72 cm from a calibrated and linearized CRT display

156 spanning approximately 31 deg horizontally and 23 deg vertically. For monkey F only, the

157 display was an LCD device running at 138 Hz (AOC AG273QX2700, 27"), as in (Malevich et al.,

158 2021). Data acquisition and stimulus control were managed by a custom-made system based

159 on PLDAPS (Eastman and Huk, 2012). The system integrated a DataPixx display control

160 device (VPixx Technologies, Inc.) with the Psychophysics Toolbox (Brainard, 1997; Pelli, 1997;

161 Kleiner et al., 2007) and an OmniPlex neural data processor (Plexon, Inc.).

162

163 The monkeys were prepared for behavioral training and electrophysiological recordings

164 earlier (Tian et al., 2018; Buonocore et al., 2019; Skinner et al., 2019; Malevich et al., 2020).

165 Specifically, each monkey was implanted with a head-holder, and monkeys M and A were  
166 also implanted with a scleral search coil in one eye. The search coil allowed tracking eye  
167 movements using the magnetic induction technique (Fuchs and Robinson, 1966; Judge et al.,  
168 1980), and the head-holder comfortably stabilized head position during the experiments.  
169 Eye movements in monkey F were recorded with a video-based eye tracker (EyeLink1000;  
170 desktop mount; 1 KHz sampling rate). For the present experiments, monkeys M and A also  
171 each had a recording chamber centered on the midline and tilted 38 deg posterior of  
172 vertical, allowing access to both the right and left SC (Bogadhi and Hafed, 2022).

173

174

175 *Behavioral tasks*

176 For the recording data in monkeys M and A, we employed a gaze fixation task in which we  
177 presented static disc of 0.51 deg radius within the visual response field (RF) of a recorded  
178 neuron. Each trial started with the onset of a black (0.11 cd/m<sup>2</sup>) fixation spot at screen  
179 center. After 550-800 ms of stable fixation on the spot, the disc appeared and remained on  
180 for at least ~500 ms. In each trial, the disc could have a Weber contrast of 5%, 10%, 20%,  
181 50%, or 100%. We defined Weber contrast as  $|I_s - I_b|/I_b$ , where  $I_s$  is the disc's luminance value  
182 and  $I_b$  is the gray background's luminance value. We often described the contrast as a  
183 percentage for convenience (e.g. 5% contrast). Importantly, across trials, the disc could have  
184 either positive or negative luminance polarity relative to the gray background, meaning that  
185  $I_s$  could be either higher (positive polarity) or lower (negative polarity) than  $I_b$ . The gray  
186 background had a luminance ( $I_b$ ) of 25.09 cd/m<sup>2</sup>. We collected approximately 50 trials per  
187 condition per neuron.

188

189 For some neurons in both monkeys (sometimes in the very same sessions as in the above  
190 task), we also ran an immediate orienting version of the stimulus polarity task. That is, at the  
191 time of disc onset, we extinguished the fixation spot (which was now white instead of black)  
192 simultaneously. This instructed the monkeys to generate an immediate orienting saccade  
193 towards the disc. We used this task to confirm that initial visual responses in the main task  
194 above were not dictated by the black fixation spot at display center, since the current task  
195 had a white fixation spot and showed similar observations (see Results), and also to obtain  
196 saccadic reaction time data for additional behavioral analyses (see below). Also note that,  
197 for neurophysiological analysis purposes, we only analyzed the initial visual response in this  
198 task. Saccade-related responses were deferred to another unrelated project focusing on SC  
199 motor bursts, and they are not described here. Finally, to reduce trial counts in this task, we  
200 only tested three contrast levels (10%, 50%, and 100%). We collected approximately 50 trials  
201 per condition per neuron.

202  
203 For exploring a potential behavioral consequence of faster detection of dark stimuli by SC  
204 neurons (which we describe in Results), we tested our three monkeys on the saccadic  
205 reaction time version of the task, which we just described above. For monkeys M and A, we  
206 analyzed reaction times from the same sessions as those collected during neurophysiological  
207 recordings. Stimulus locations were, thus, dictated by recorded neurons' response field (RF)  
208 locations. For monkey F, we ran behavior-only sessions. In this case, we randomly varied  
209 stimulus locations across 4 diagonals, with an eccentricity of 8.9 deg. We analyzed a total of  
210 457-773 saccades per condition per monkey for our behavioral reaction time analyses.

211

212

213 *Neurophysiological procedures*

214 For most experiments, we recorded SC neurons using linear electrode arrays inserted across  
215 the SC depth (24 channel V Probes; Plexon, Inc.). For some experiments, we also used single  
216 tungsten electrodes, in which we targeted and isolated individual neurons online during the  
217 experiments. In all cases, including the single electrode sessions, we also performed offline  
218 sorting to re-isolate neurons for inclusion in the data analysis pipeline (Pachitariu et al.,  
219 2016). Sorting and general data analysis pipeline details were similar to those described  
220 recently (Bogadhi and Hafed, 2022).

221

222 Before collecting data from our main tasks, we first identified the SC by running RF mapping  
223 tasks. These included delayed and memory-guided saccades (Chen et al., 2015; Chen and  
224 Hafed, 2017). The mapping tasks allowed us to select the stimulus location for our main  
225 experiments, and also to confirm that our neurons possessed visual responses (or stimulus-  
226 triggered inhibition). For the simultaneous recordings of multiple neurons with electrode  
227 arrays, we picked a disc location that we felt lay within the RF's of most neurons that we  
228 could identify online. This was possible given that our electrodes were penetrating the SC  
229 surface at a quasi-orthogonal angle, meaning that the RF's at different depths generally had  
230 similar locations. Also note that for all analyses, we were always interested in comparing  
231 responses to bright and dark discs at the very same location. That is, our comparison of  
232 interest was the luminance polarity at a given RF location for a given neuron. In separate  
233 experiments, we mapped RF's with positive and negative luminance polarity spots, but these  
234 data will be described in detail separately. For the present purposes, suffice it to say that all  
235 RF's had sensitivity to both black and white targets at their center, justifying our current  
236 comparison of response sensitivity at a single given RF location per neuron.

237

238 Across all experiments, we recorded from neurons with extrafoveal eccentricities (e.g. 2.1-  
239 20 deg preferred eccentricity across the population), meaning that we often presented  
240 stimuli far from the fixation spot.

241

242

243 *Eye movement data analysis*

244 We detected saccades and microsaccades as described previously (Chen and Hafed, 2013;  
245 Bellet et al., 2019). We used the detections for two primary purposes. First, in the recording  
246 tasks, we excluded all trials in which there were microsaccades occurring within an interval  
247 from -50 ms to +50 ms relative to stimulus onset. This allowed us to measure baseline visual  
248 responses that were not modulated by the known influences of microsaccades on SC activity  
249 (Hafed and Krauzlis, 2010; Chen et al., 2015; Chen and Hafed, 2017). We also performed the  
250 same filtering for behavioral analyses of saccadic reaction times. Second, for the behavioral  
251 analyses, we used saccade detection to measure saccadic reaction times towards the dark  
252 and bright stimuli.

253

254 To identify a response saccade and subsequently analyze its reaction time, we required that  
255 it had a latency of 50 to 500 ms from stimulus onset, and that it was directed towards the  
256 stimulus (this latter criterion was easy to achieve because we used computer-controlled  
257 reward windows around the target to allow rewarding the monkeys based on successful  
258 saccade generation towards the target). In all neural and behavioral analyses, we also  
259 excluded trials with blinks or other movement artifacts near stimulus onset. Statistically, we  
260 were interested in whether contrast or luminance polarity affected saccadic reaction times.

261 Therefore, we performed a 1-way non-parametric ANOVA (Kruskal-Wallis test) in each  
262 monkey testing for the effect of contrast (d.f.: 2), when collapsing across luminance  
263 polarities, on the monkey's reaction times. Similarly, we also performed a Kruskal-Wallis test  
264 exploring the effect of luminance polarity (d.f.: 1), when collapsing across contrasts, on  
265 reaction times.

266

267

268 *Neural data analysis*

269 We analyzed a total of 221 SC neurons (109 from monkey M and 112 from monkey A) from  
270 the fixation task. We also analyzed 225 neurons from the immediate saccade version of the  
271 task (113 from monkey M and 112 from monkey A). Ninety of the neurons in the second task  
272 (all in monkey A) were also recorded from the fixation variant of the task.

273

274 The bulk of our analyses was on neurons exhibiting a positive visual response to stimulus  
275 onset (that is, an increase in firing rate relative to baseline shortly after stimulus onset). We,  
276 therefore, first tested for the presence of a positive visual response (or burst) after stimulus  
277 appearance. In each neuron, we defined a baseline interval as the final 50 ms before  
278 stimulus onset. We then defined a visual response interval as the time interval 10-200 ms  
279 after stimulus onset. Across all repetitions of a given stimulus condition (e.g. 100% contrast;  
280 positive polarity), we measured average firing rate in the response interval and statistically  
281 compared it to average firing rate in the baseline interval. If the response interval firing rate,  
282 across repetitions of a given condition, was statistically significantly larger (one-tailed, paired  
283 t-test;  $p < 0.025$ ) than baseline firing rate, and if this significance occurred for both polarity  
284 conditions (dark and bright) and with absolute Weber contrasts of 50% and 100%, then we

285 considered the neuron to have a positive visual response to stimulus onset. We did not  
286 include lower contrast trials (whether positive or negative polarity) in assessing for the  
287 presence of visual responses because some neurons, even when having strong visual bursts  
288 for high contrast stimuli, did not respond to such lower contrasts. Across our population in  
289 the fixation task, we had a total of 172 neurons (92 from monkey M and 80 from monkey A)  
290 exhibiting visual bursts after stimulus onset with the above criteria. For the immediate  
291 saccade version of the task, we only analyzed neurons with a positive visual burst in the  
292 interval 10-180 ms after stimulus onset; this resulted in a total of 213 neurons (109 from  
293 monkey M and 104 from monkey A).

294

295 For a subset of neurons, stimulus onset caused a transient decrease in firing rate from  
296 baseline, rather than an increase. We performed analyses of these neurons as well, from the  
297 fixation task only. To assess the neurons as having a transient decrease in firing rate that was  
298 time-locked to stimulus onset, we repeated the same procedure above, but we now checked  
299 for a statistically significant decrease in firing rate in the response interval, rather than an  
300 increase. We analyzed 15 neurons with transiently decreasing firing rates immediately after  
301 stimulus onset (9 from monkey M and 6 from monkey A).

302

303 To obtain contrast sensitivity curves from the neurons with visual bursts, we measured the  
304 peak value of the average firing rate curve in a response interval after stimulus onset. Since  
305 visual response latency in the SC varies with stimulus contrast (Li and Basso, 2008; Marino et  
306 al., 2012; Marino et al., 2015), we tailored the measurement interval for each contrast as  
307 follows: 15-105 ms after stimulus onset for 100% contrast; 20-110 ms after stimulus onset  
308 for 50% contrast; 20-115 ms after stimulus onset for 20% contrast; 35-125 ms after stimulus

309 onset for 10% contrast; and 45-135 ms after stimulus onset for 5% contrast. Note that we  
310 used the same measurement intervals for all neurons and also for both positive and negative  
311 polarity stimuli. Even though we found a difference in response latency between positive  
312 and negative polarity stimuli (as described in Results below), our measurement intervals  
313 were large enough to encompass (and exceed) any such latency differences. Therefore, our  
314 estimates of contrast sensitivity for brights and darks were not biased by using similar  
315 measurement intervals for both types of stimuli (especially because we were searching for  
316 only the peak firing rate). After measuring firing rates in the above intervals for each  
317 contrast, we plotted the measured firing rates as a function of absolute contrast. We then fit  
318 contrast sensitivity curves using the following equation:

319

320 
$$f(c) = R \frac{c^N}{C_{50}^N + c^N} + B \quad (\text{Equation 1})$$

321  
322 where  $f$  is the estimated firing rate,  $c$  is stimulus contrast,  $C_{50}$  is semi-saturation contrast,  $R$   
323 is the dynamic range of the response,  $N$  is the sensitivity/slope of the contrast sensitivity  
324 curve, and  $B$  is the baseline firing rate (which we just measured across all trials from the  
325 same baseline interval mentioned above; final 50 ms before stimulus onset). We then  
326 compared the fit parameters  $R$ ,  $C_{50}$ , and  $N$  for either bright or dark stimuli to assess whether  
327 there were differences in contrast sensitivity between them in the SC. We did this by  
328 computing parameter modulation indices as a function of luminance contrast polarity. For  
329 example, to compare how  $R$  was modulated by luminance polarity, we calculated the  $R$   
330 parameter for bright stimuli minus the  $R$  parameter for dark stimuli, and we divided this  
331 difference by the sum of  $R$  values for bright and dark stimuli. This gave us a value between -1  
332 and +1. We then plotted histograms of parameter modulation indices across the population.

333

334 For assessing the time course of changes of contrast sensitivity in the sustained interval long

335 after stimulus onset (that is, after the initial visual burst), we again obtained fits of equation

336 1 but now based on measurements after the initial visual burst. To do so, we exploited the

337 fact that our firing rate estimates were already averaging across time (because of a

338 convolution of spike times with a gaussian of  $\sigma$  40 ms). Therefore, for each time after 80 ms

339 after stimulus onset and before 220 ms, we used instantaneous average firing rate as a

340 measure to input to the fit of equation 1 for a given contrast. This allowed us to obtain time

341 courses of semi-saturation contrast ( $C50$ ), sensitivity/slope ( $N$ ), and dynamic range ( $R$ ) during

342 the sustained interval long after stimulus onset. Note that even though the intervals that we

343 chose for the sustained response analysis slightly overlapped with the intervals that we

344 picked up for the initial visual burst analyses mentioned above, the latter analyses were

345 performed on the peak values of the average firing rates, which definitely belonged to the

346 earlier phases of the neural responses (and typically occurred earlier than 80 ms); that is, the

347 initial visual burst intervals were just ranges meant to catch the peak response. Also note

348 that the above contrast sensitivity fits were only performed on the fixation version of the

349 task because we could obtain a longer period of sustained response than in the immediate

350 saccade version of the task.

351

352 For estimating visual response latency in both tasks, we measured the firing rate of a given

353 neuron in a baseline interval (final 50 ms before stimulus onset) across all trials. Then, for

354 each condition (e.g. bright luminance polarity; 20% contrast), we marched forward in time

355 after stimulus onset until 300 ms (typically, the algorithm converged on a visual burst much

356 earlier, of course). As soon as the lower bound of the 95% confidence interval around the

357 average firing rate of the neuron across trial repetitions exceeded the average baseline  
358 activity (and continued to do so for at least 30 ms), we flagged the time as the response  
359 latency of the neuron. Whenever this algorithm failed to detect the response latency in at  
360 least one of the luminance polarities for a given stimulus contrast level, we excluded the  
361 neuron from further analysis in that contrast level. This explains the varying numbers of  
362 neurons reported in some figures (e.g. the different panels of Fig. 4 in Results). We then  
363 compared such response latency across contrasts and stimulus polarities. Note that we  
364 focused on relative latency differences across luminance polarities in our analyses. This is  
365 important to note because firing rate estimates (in our case, convolution of spike times with  
366 a Gaussian kernel) necessarily blurs the exact response onset times of the neurons.

367 However, our approach of estimating response latencies described above still captured the  
368 latency differences that we were interested in documenting, and it simplified the detection  
369 of visual response latencies for neurons with non-zero baseline firing rates.

370

371 To statistically test for differences in latencies between luminance polarities at a given  
372 contrast level, we used non-parametric permutation tests on the pairwise mean latency  
373 differences, with 10000 permutations. That is, we obtained the permutation distribution by  
374 shuffling the polarity labels of the latencies for 10000 times while maintaining their pairwise  
375 relationship and calculating their pairwise difference. Monte Carlo p-values were obtained  
376 by assessing the probability of getting larger than or equal to absolute latency differences in  
377 the permutation distribution than the absolute latency difference of the original data. We  
378 ran the tests separately for each monkey to ensure that our pooling of data in figures for  
379 visualization purposes was justified.

380

381 We also used a similar approach to test for statistically significant effects of upper versus  
382 lower visual field RF location on the latency differences between luminance polarities. This  
383 time, we obtained the latency differences between the responses to bright and dark stimuli,  
384 and then subtracted this measurement for the upper visual field neurons from the  
385 measurement for the lower visual field neurons. We defined lower and upper visual field  
386 neurons based on the location of the stimulus (which was placed close to the location of the  
387 RF hotspot location). Thus, negative values in the final measurement would indicate a larger  
388 difference between dark and bright stimuli in the upper visual field than in the lower visual  
389 field. After that, we ran permutation tests by shuffling the labels of the upper and lower  
390 visual field neurons for 10000 times. To assess the significance of the results, we calculated  
391 the Monte Carlo p-value. The same procedure was applied to assess the absolute values of  
392 sensitivity differences (see next paragraph) between dark and bright stimuli in the upper and  
393 lower visual fields.

394

395 To compare visual response latency to sensitivity, we also measured peak firing rate in the  
396 initial visual response interval (as defined for each contrast above) of the neuron. First, to  
397 test whether there was an effect of luminance polarity on sensitivity, we used permutation  
398 tests in the same way as we did for the latency analysis described above, but this time on  
399 the pairwise mean peak response differences, separately for each monkey and contrast  
400 level. We then checked whether there was a dissociation between response latency and  
401 sensitivity (i.e. response strength) for black and white stimuli, as we previously saw for  
402 spatial frequency stimuli (Chen et al., 2018). We did so by sorting the neurons according to  
403 the difference in response latencies between brights and darks, and then checking whether  
404 the same sorting applied to the difference in response sensitivities. Further, we pooled the

405 data across monkeys and calculated Pearson's correlation coefficients between differences  
406 in peak visual responses and differences in visual response latencies, separately for each of  
407 the contrast levels (see Results).

408  
409 For the immediate saccade version of the task, we only analyzed initial visual bursts (50-130  
410 ms after stimulus onset) and not sustained intervals. This was because the response saccade  
411 occurred too soon after the initial visual bursts. We assessed both response sensitivity and  
412 response latency (as described above) to confirm that we got similar results to those from  
413 the fixation task.

414  
415 For the neurons with transient decreases in firing rate, we assessed response latency in a  
416 similar way to the neurons with visual bursts, but we looked for statistically significant  
417 decreases in firing rate after stimulus onset, rather than increases.

418  
419 In some figures, for illustration and visualization purposes, we elected to show example  
420 population firing rates from individual monkeys. For example, we did this in Fig. 8A, B in  
421 Results. To obtain such population firing rates, we obtained the normalized average firing  
422 rate of each neuron, per monkey and condition. That is, for each neuron, we found the peak  
423 visual response in the interval 0-100 ms after stimulus onset for the 100% contrast stimuli,  
424 regardless of the stimulus polarity. Then, for each contrast and polarity, we normalized the  
425 neuron's average firing rate by that peak visual response value. This resulted in a series of  
426 average normalized curves for the neuron across conditions. After that, we averaged all of  
427 the normalized firing rate curves of each monkey's neurons in a given condition. This gave us

428 a population summary of responses, maintaining the relative changes in responses across  
429 conditions. We used a similar approach in Fig. 11B, D.

430

431

432 *Experimental design and statistical analyses*

433 We recorded neurons in an unbiased manner by collecting data in parallel (with linear  
434 electrode arrays) in most sessions and then sorting the neurons offline. This allowed us to  
435 minimize sampling bias. In each variant of the task, we also analyzed >80 neurons per  
436 monkey. This provided a large enough sample to assess the reliability of our interpretations.

437 Within each neuron, we ensured collecting approximately 35-50 repetitions per condition  
438 (after filtering out bad trials and so on) to allow robust within-neuron statistics. Similarly, in  
439 our behavioral analyses, we collected thousands of saccades. In all cases, we randomly  
440 interleaved stimulus presentations across trials, to avoid any blocking effects.

441

442 We provided descriptive statistics in all figures, showing numbers of observations and  
443 measures of variability. Also, in most of our critical analyses (e.g. Figs. 4-6 in Results), we  
444 showed the full distributions of data points that we had.

445

446 Since the replicate of interest was neurons, our numbers of sampled neurons were  
447 sufficient. The use of two monkeys in recording was valuable to increase neuron counts, and  
448 to also demonstrate repeatability across individuals. Our results were highly similar in the  
449 two animals (e.g. Fig. 8A, B in Results). When they did differ, we showed each individual  
450 monkey's results separately (e.g. Fig. 11 in Results), and this was highly useful for us to

451 interpret the behavioral results. Moreover, we collected behavior from a third monkey

452 exactly to improve our interpretation of our individual monkey behavioral phenomena.

453

454 All statistical tests are reported and justified in Results at appropriate points in the text. As

455 stated above, we statistically analyzed each monkey's data individually, confirming that each

456 monkey showed the same effects (unless otherwise stated; for example, in Fig. 11 in

457 Results).

458

459 **Results**

460

461 We investigated how monkey superior colliculus (SC) neurons respond to dark and bright  
462 visual stimuli. In our primary task, the monkeys fixated while we presented a small disc that  
463 was either higher or lower in luminance than the gray background of the display. We varied  
464 the contrast of the disc from the background luminance, and we assessed contrast  
465 sensitivity curves separately for positive and negative luminance polarities. We first analyzed  
466 the neurons that exhibited visual bursts (that is, increases in firing rate) after stimulus onset,  
467 and we investigated visual burst strength, visual burst latency, as well as sustained response  
468 dynamics for dark and bright stimuli. The results for these neurons are described next,  
469 followed by an analysis of a smaller number of neurons for which stimulus onsets caused  
470 transient decreases in firing rates, rather than increases.

471

472

473 *Diverse preferences for darks and brights across SC neurons*

474 We first asked whether neurons tended to be more sensitive to darks or brights across the  
475 population. For each recorded neuron, we plotted firing rate as a function of time from  
476 stimulus onset, and we assessed the strength of the visual burst as a function of luminance  
477 contrast polarity. Figure 1A-C shows the responses of three example neurons (from the  
478 same monkey, A) to a 100% contrast stimulus. The black lines show responses to the  
479 negative polarity stimulus (darker than background), and the light gray lines show responses  
480 to the positive polarity stimulus (brighter than background). In all cases, the negative and  
481 positive polarity stimuli were of the same size and presented at the same location. They also  
482 had the same absolute Weber contrast, and their presentation sequence was randomly

483 counterbalanced across trials. As can be seen, there was a diversity of neural preferences  
484 across the three neurons: neuron 1 (Fig. 1A) was more sensitive to the positive polarity  
485 stimulus than to the negative polarity stimulus; neuron 2 was, more or less, equally sensitive  
486 to the two stimuli (Fig. 1B); and neuron 3 was clearly more sensitive to the dark stimulus  
487 (Fig. 1C). We also plotted full contrast sensitivity curves for the same neurons (Fig. 1D-F) by  
488 relating peak visual response strength to stimulus contrast (Methods). Consistent with Fig.  
489 1A-C, there was a diversity of preferences for darks and brights across the three neurons in  
490 their full contrast sensitivity curves.

491  
492 These observations held across the population of 172 neurons that we analyzed. For each  
493 neuron, we fit a contrast sensitivity function (equation 1; see Fig. 1D-F for examples) by  
494 optimizing three parameters characterizing how the neuron altered its visual response with  
495 Weber contrast:  $R$  reflected the dynamic range of the response,  $C50$  characterized the semi-  
496 saturation contrast of the neuron, and  $N$  characterized the steepness of the contrast  
497 sensitivity curve (slope parameter). We performed such a fit for either positive or negative  
498 luminance polarity stimuli. We then obtained a parameter modulation index, describing, for  
499 each neuron, to what extent each parameter of the fit was different between positive and  
500 negative luminance polarity stimuli. For example, for dynamic range (parameter  $R$  in  
501 equation 1), we obtained the  $R$  value for bright stimuli minus the  $R$  value for dark stimuli in  
502 each neuron, and we then divided this difference by the sum of  $R$  values for the two stimulus  
503 types (Methods). This gave us an index in which 1 meant that the neuron responded  
504 maximally only to bright stimuli and -1 meant that the neuron responded maximally only to  
505 dark stimuli. An  $R$  parameter modulation index value of 0, instead, indicated equal visual  
506 response dynamic ranges for bright and dark stimuli. We then plotted histograms of the

507 parameter modulation indices across the population. As can be seen in Fig. 2, all three  
508 modulation indices of the contrast sensitivity function fits had distributions straddling 0, and  
509 with large diversity across the population. Some neurons clearly preferred bright stimuli,  
510 others clearly preferred dark stimuli, and yet others were equally sensitive to darks and  
511 brights (near 0 in the histograms of Fig. 2). The vertical lines in Fig. 2 indicate the mean  
512 (solid) and median (dashed) parameter modulation index values across neurons, and they  
513 were all close to 0. Approximately half of the neurons were more sensitive to bright stimuli  
514 (whether in terms of dynamic range, semi-saturation contrast, or slope of the contrast  
515 sensitivity function), and the other half were more sensitive to dark stimuli.

516

517 Therefore, in the SC, we noticed a substantial diversity of sensitivity preferences for darks  
518 and brights across the population (unlike in LGN and V1). This suggests that stimuli of both  
519 positive and negative luminance polarities can indeed be represented well by SC neural  
520 populations.

521

522

523 *Earlier detection of darks by SC neurons, regardless of preference*

524 Unlike response sensitivity, for which we saw diverse preferences for brights and darks (Figs.  
525 1, 2), SC neurons exhibited systematically shorter visual response latencies for dark stimuli,  
526 independently of their visual response strengths at a given contrast. Consider, for example,  
527 the same three neurons of Fig. 1A-C. In each of them, visual responses occurred earlier for  
528 the dark stimuli than for the bright stimuli, as can be visually assessed from the spike rasters  
529 and the firing rate density plots below them. This happened even for neuron 1, which  
530 preferred bright stimuli (Fig. 1A). It also happened at different contrast levels (Fig. 3), even

531 though stimulus contrast expectedly modulated the response strength and latency of each  
532 neuron. For example, at 20% contrast, all three neurons from Fig. 1 still responded earlier to  
533 dark than bright stimuli, despite the weakened and delayed visual responses relative to the  
534 100% contrast conditions. Thus, at each contrast level, there was an apparent dissociation  
535 between visual response sensitivity and visual response latency in these three neurons, not  
536 unlike what we recently observed when we presented different spatial frequencies to SC  
537 neurons (Chen et al., 2018).

538

539 To investigate this dissociation further, we estimated, for each neuron, the onset of the  
540 visual burst as the first time point at which the lower bound of the 95% confidence interval  
541 of the neuron's average firing rate was elevated for a prolonged period of time above its  
542 baseline activity (Methods). Even though estimating neural response latencies from firing  
543 rate measures like we did might blur the actual absolute values of the response latencies,  
544 due to convolution kernels with spike times, this approach was still sufficient to capture the  
545 latency differences across luminance polarities that we were interested in (Methods).

546 Therefore, for each contrast, we subtracted each neuron's visual response latency for dark  
547 stimuli from its visual response latency for bright stimuli, and we sorted the neurons  
548 according to this difference. An example of such sorting can be seen in the top panel of Fig.  
549 4A for the 100% contrast stimuli. Note how a majority of neurons (76.7%; 128 out of 167)  
550 had an earlier visual response for dark stimuli (evidenced by a positive latency difference in  
551 the figure). This is in contrast to the diversity of preferences for darks and brights seen in  
552 Figs. 1-3. In fact, with the very same sorting of the neurons as in the top panel, we next  
553 plotted (bottom panel of Fig. 4A) the same neurons' differences in peak visual burst  
554 strengths between darks and brights (Methods). The neurons were no longer as properly

555 ordered as in the top panel, suggesting that the latency effect in the top panel was not  
556 trivially explained by a systematic difference in response sensitivity between darks and  
557 brights. For example, both neurons 126 and 156 (highlighted in Fig. 4A with small diagonal  
558 arrows) possessed clearly stronger responses for bright stimuli than dark stimuli (positive  
559 difference in the bottom panel), but they both had a later response latency for bright stimuli  
560 (positive difference in the top panel). Therefore, visual responses to dark stimuli still  
561 occurred earlier than visual responses to bright stimuli even when neurons preferred bright  
562 stimuli.

563  
564 We also made similar observations for the other stimulus contrasts that we tested (Fig. 4B-  
565 E). Note that for each panel in Fig. 4, we indicated the total number of neurons included into  
566 each analysis, which varied across panels (that is, across contrast levels). This happened  
567 because some neurons may not have met our inclusion criteria for estimating visual  
568 response latencies, resulting in slightly different neuron counts across the different panels  
569 (Methods). For example, for the particularly low contrast stimuli (e.g. 5% and 10%), some  
570 neurons did not exhibit any significant visual bursts at all (Methods), so they were not  
571 included in the figure. Having said that, in all contrasts, there was a majority of neurons  
572 responding earlier to dark than bright stimuli (top row in each panel of Fig. 4) regardless of  
573 the relative strengths of their visual responses (bottom row).

574  
575 We confirmed this observation statistically with permutation tests, conducted on pairwise  
576 latency differences separately for each contrast level and for each monkey (Methods). In  
577 monkey M, there were significantly longer latencies for bright stimuli in all contrasts (100%  
578 and 50% contrasts: mean differences = 3.18 ms and 3.56 ms, respectively, Monte Carlo p-

579 values < 0.0001; 20% contrast: mean difference = 1.62 ms, Monte Carlo p-value = 0.0448;  
580 10% contrast: mean difference = 2.03 ms, Monte Carlo p-value = 0.0152; and 5% contrast:  
581 mean difference = 4.5 ms, Monte Carlo p-value < 0.001). In monkey A, latencies were  
582 significantly longer for bright stimuli in 100%, 50%, and 5% contrasts (mean differences =  
583 2.97 ms, 4.31 ms, and 4.91 ms, respectively; Monte Carlo p-values: < 0.0001, < 0.0001, and  
584 0.0059, respectively); no significant differences were found in 20% and 10% contrasts (mean  
585 differences = 1.39 ms and 0.27 ms, respectively; Monte Carlo p-values: 0.0548 and 0.7587,  
586 respectively), but the same trends were still there (also see Fig. 5E). Thus, faster detection of  
587 dark than bright stimulus contrasts is a general property of SC neurons.

588

589 Of course, our results do not deny that high visual response sensitivity is normally associated  
590 with short visual response latencies. For example, with the sorting of neurons shown in Fig. 4  
591 based on their response latency differences (top row), there was still a hint of an additional  
592 trend: neurons with a smaller latency difference between dark and bright stimuli tended to  
593 be the neurons preferring bright stimuli (bottom row). For example, compare the first and  
594 last quartiles in the bottom panel of Fig. 4A: more bright-preferring neurons occurred in the  
595 first quartile (having 0 or negative latency differences) than in the last quartile (having  
596 positive latency differences). This suggests that there were divergent forces influencing  
597 visual response latency: a neuron strongly preferring bright stimuli might have had its high  
598 response strength for bright stimuli (at a given contrast level) counterbalance the normally  
599 earlier detection of dark stimuli. Indeed, when we evaluated response latency as a function  
600 of both stimulus contrast (a proxy for visual response strength in the neurons) and  
601 luminance polarity, we found that both factors clearly influenced the neurons' visual  
602 response latencies. This is shown in Fig. 5A, B for an example neuron, and in Fig. 5C for the

603 population. In Fig. 5A, high contrast stimuli evoked stronger and, therefore, earlier visual  
604 responses than low contrast stimuli, as expected (Boehnke and Munoz, 2008; Marino et al.,  
605 2012; Marino et al., 2015; Hafed and Chen, 2016; Chen et al., 2018). With high contrast dark  
606 stimuli, the same neuron exhibited even earlier visual bursts than for high contrast bright  
607 stimuli (Fig. 5B). Across the population, cumulative histograms of estimated visual response  
608 latencies (Fig. 5C), as well as their averages and standard errors of the mean (Fig. 5D),  
609 revealed that increasing stimulus contrast systematically decreased response latencies, as  
610 expected (Boehnke and Munoz, 2008; Marino et al., 2012; Marino et al., 2015; Chen et al.,  
611 2018), but also that visual response latencies were always systematically shorter, at a given  
612 contrast, for dark than bright stimuli (consistent with Fig. 4; also see Fig. 6 below). This  
613 polarity effect on response latencies had an order of magnitude of a few milliseconds  
614 difference between dark and bright response latencies (Fig. 5E), similar to results in the cat  
615 LGN (Jin et al., 2011) and V1 (Komban et al., 2014). Therefore, both stimulus contrast (a  
616 proxy for response sensitivity) and stimulus polarity (conferring a temporal advantage for  
617 darks) dictated our SC neurons' visual response latencies.

618

619

620 *Interaction between stimulus contrast and the processing of darks and brights  
621 by SC neurons*

622 The results of Fig. 5E were particularly intriguing to us, in the sense that the lowest contrast  
623 stimuli (5%) were associated with a seemingly bigger effect of visual response latency  
624 difference between darks and brights than the more visible 10% and 20% contrast targets.  
625 One possibility could be that at 5% contrast, there were fewer bright-preferring neurons.  
626 We, therefore, next asked whether there was an interaction between luminance contrast

627 level and the preference of neurons for darks or brights. To do so, we replotted the data  
628 above as scatter plots of visual response latency for brights versus darks in Fig. 6A-E and as  
629 scatter plots of visual response sensitivity for brights versus darks in Fig. 6F-K. The latency  
630 plots confirmed our earlier observations that there was systematically faster detection of  
631 dark contrasts across all contrasts. For sensitivity, there was an interaction between contrast  
632 level and SC population preference. At high contrasts (e.g. Fig. 6F, G), the population was  
633 biased toward preferring bright stimuli (despite responding faster for dark stimuli), whereas  
634 at 5% contrast (Fig. 6K), this bias disappeared and tended to be in the opposite direction;  
635 neuron 1 in Figs. 1D, 3A also demonstrates this effect: even though the neuron preferred  
636 brights in its plateau firing rate of the contrast sensitivity curve, its (weak) response at 5%  
637 contrast was still higher for dark targets. Thus, there was an interaction between contrast  
638 level and sensitivity to darks in our SC neurons. While this pattern is different from natural  
639 image statistics (Cooper and Norcia, 2015) and cat V1 properties (Liu and Yao, 2014), in the  
640 sense that we found more bright-preferring than dark-preferring neurons at high contrast, it  
641 does suggest that the larger latency effect magnitude in Fig. 5E at 5% contrast might have  
642 been driven by a larger number of dark-preferring neurons at this contrast level.

643  
644 Statistically, we confirmed that there were contrast-dependent sensitivity preference  
645 differences between brights and darks. We applied the same pairwise latency difference  
646 procedure described above, but now to pairwise peak visual response differences  
647 (Methods). In monkey M, SC visual responses to brights were significantly stronger than  
648 responses to darks in the 100%, 50%, and 20% contrast conditions (mean differences = 11.69  
649 spikes/s, 8.92 spikes/s, and 4.26 spikes/s, respectively; Monte Carlo p-values: < 0.0001, <  
650 0.001, and 0.0187, respectively); the differences were not significant for 10% and 5%

651 contrasts, and their trends were in the opposite direction (mean differences = -0.09 spikes/s  
652 and -2.35 spikes/s, respectively; Monte Carlo p-values: 0.9611 and 0.1712, respectively). In  
653 monkey A, the neurons were significantly more sensitive for brights in all but the lowest  
654 contrast (100%, 50%, 20%, 10%, and 5% contrasts: mean differences = 6.73 spikes/s, 7.35  
655 spikes/s, 6.52 spikes/s, 6.92 spikes/s, and 0.81 spikes/s, respectively; Monte Carlo p-values:  
656 0.0106, 0.002, 0.0028, < 0.001, and 0.8652 respectively).

657

658 In the same vein, for each neuron, we plotted the visual response latency difference against  
659 the peak response difference in Fig. 6L-P. We used the same conventions as in Fig. 4: a  
660 positive response latency difference indicating a faster response to dark stimuli, and a  
661 positive peak response difference meaning higher sensitivity to bright stimuli. If the results  
662 of Figs. 4, 5 were solely determined by response sensitivity at each contrast level, then all  
663 neurons should have occupied the shaded quadrants of these plots. In contrast, only a  
664 minority of neurons occupied these quadrants, particularly at high contrast. For example, in  
665 Fig. 6L, even neurons with >50 spikes/s difference in peak sensitivity in favor of bright stimuli  
666 were still significantly faster to detect dark stimuli. Interestingly, at 5% contrast, there were  
667 significantly fewer bright-preferring neurons, again providing a plausible explanation for the  
668 relatively large effect size in response latency seen in Fig. 5E at 5% contrast versus 10% and  
669 20% contrast.

670

671

672

673

674 *Interaction between visual field location and the faster detection of darks by SC*

675 *neurons*

676 The above results indicate that there is faster detection of dark than bright stimuli by SC  
677 neurons, in general. However, it is also known that SC visual responses preferentially process  
678 the upper visual field (Hafed and Chen, 2016), consistent with the notion that eye  
679 movements support orienting towards or away from extra-personal stimuli largely occupying  
680 the upper visual field (Previc, 1990). If that is indeed the case, then it might be expected that  
681 differential temporal processing of dark versus bright stimuli might be magnified in the SC's  
682 upper visual field representation. For example, birds of prey, or other threats, across a  
683 daylight sky would normally cast dark contrasts on retinal images, and they need to be  
684 detected efficiently by SC neurons. We, therefore, also asked whether the results of Fig. 4  
685 could depend on the visual field locations of our recorded neurons.

686

687 We repeated the analyses of Fig. 4, but this time after separating neurons based on upper  
688 and lower visual field RF locations. The results are shown in Fig. 7 (for the highest contrast  
689 stimuli only, for simplicity). There was indeed a larger latency difference between dark and  
690 bright stimulus responses in the upper visual field neurons than in the lower visual field  
691 neurons (top panel). That is, the latency advantage for dark stimuli was magnified in the case  
692 of upper visual field SC neurons. We tested this observation statistically by using a  
693 permutation test with 10000 shuffles. Here we should note that although we pooled the  
694 data of both monkeys for visualization purposes in Fig. 7, we performed the statistical  
695 procedures only on data collected from monkey A. This was because there was a strongly  
696 unbalanced sampling of neurons in the upper and lower visual fields in monkey M (68 and 21  
697 upper and lower visual field neurons in this monkey, versus a more balanced distribution of

698 24 and 33 neurons in monkey A). In monkey A, there was a significant difference between  
699 upper and lower visual field effects (mean difference = -4.09 ms, Monte Carlo p-value =  
700 0.0099, Methods). Therefore, a known visual response latency advantage for the upper  
701 visual field in SC neurons (Hafed and Chen, 2016) was accompanied, at least in one monkey,  
702 by a larger difference between dark and bright stimulus responses. This result is consistent  
703 with the ecological likelihood of dark contrasts in natural environments (Liu and Yao, 2014;  
704 Cooper and Norcia, 2015), and also with the role of the SC's visual processing machinery in  
705 supporting the sampling of extra-personal visual space by orienting eye movements (Previc,  
706 1990; Hafed and Chen, 2016; Fracasso et al., 2022).

707

708 Note also that the same dissociation between response latency and response sensitivity  
709 occurred in Fig. 7 as in Fig. 4: the bottom panel in Fig. 7 shows that with the same ordering  
710 of the neurons as in the top panel, response sensitivity was not systematically ordered in  
711 either the upper or lower visual fields, consistent with the results of Fig. 4. Interestingly, the  
712 absolute value of the difference in response strength between dark and bright stimuli was  
713 also higher in the upper visual field neurons than in the lower visual field neurons (monkey  
714 A; mean difference = -13.3 spikes/s; Monte Carlo p-value = 0.0166, permutation test). This  
715 suggests that both latency differences (top panel) and absolute values of sensitivity  
716 differences (bottom panel) between dark and bright stimuli were amplified in the upper  
717 visual field representation of the SC, adding to a growing body of evidence of visual field  
718 asymmetries in the primate SC (Hafed and Chen, 2016; Hafed, 2021; Fracasso et al., 2022).

719

720

721 *Independence of the faster SC detection of darks from the luminance polarity at*  
722 *fixation*

723 Finally, we wondered whether the black fixation spot at display center (Methods) might have  
724 dictated our results above. Such an effect would be unlikely because our neurons were  
725 extra-foveal; our stimuli were, therefore, generally far from the fixation spot (Methods).  
726 However, to unambiguously rule such an effect out, we repeated the same experiment with  
727 two slight modifications. First, the fixation spot was now white instead of black (Methods). If  
728 the black fixation spot was the reason for the faster detection of dark stimuli in the results of  
729 Figs. 1, 3-7 above, then this effect should be altered with a white fixation spot. Second, the  
730 fixation spot was now removed at the same time as stimulus onset, allowing the monkeys to  
731 generate immediate, visually-guided saccades. We analyzed 213 neurons recorded with this  
732 variant of the task (81 were also recorded in the original fixation task). We will describe  
733 saccadic reaction times as a function of contrast and luminance polarity in more detail  
734 below. However, for now, our aim was to replicate the visual burst results shown above. For  
735 each neuron, we normalized the neuron's average firing rate by the peak visual response for  
736 100% contrast stimuli in the interval 0-100 ms after stimulus onset (Methods). We then  
737 averaged all of the normalized firing rate curves of each monkey's neurons (Fig. 8A, B). We  
738 separated the neurons of each monkey in this analysis to demonstrate the repeatability of  
739 our results across the animals, and also because subsequent saccadic behavior later in the  
740 trials differed between them, as we clarify in more detail below.

741  
742 Both animals had clear visual responses in the task, consistent with the results of the fixation  
743 variant (Figs. 1, 3-7). Most importantly, these responses were also clearly still occurring  
744 earlier for dark stimuli than for bright stimuli (Fig. 8A, B). To summarize these results on an

745 individual neuron basis, we replicated the same analyses of Fig. 4 (Fig. 8C-E; we combined  
746 the neurons of both monkeys here because of how similarly they behaved in Fig. 8A, B). We  
747 did this for all three stimulus contrast levels that we tested in this variant of the task  
748 (Methods). For all contrasts, most neurons still detected dark contrasts earlier than bright  
749 contrasts, irrespective of neural sensitivity (Fig. 8C-E), just as in Figs. 1, 3-7. Permutation  
750 tests run on the latency differences confirmed this observation for all contrasts (in monkey  
751 M, 100% and 50% contrasts: mean differences = 4.97 ms and 4.42 ms, respectively, Monte  
752 Carlo p-values < 0.0001; 10% contrast: mean difference = 1.62 ms, Monte Carlo p-value =  
753 0.0185; in monkey A, 100% and 50% contrasts: mean differences = 3.71 ms and 3.35 ms,  
754 respectively; Monte Carlo p-values < 0.0001; 10% contrasts: mean difference = 2.52 ms;  
755 Monte Carlo p-value < 0.001). Note that the effect sizes were also of the same order of  
756 magnitude as those shown in Fig. 5E. Therefore, the results of Figs. 1-7 were not trivially  
757 caused by the use of a black fixation spot at screen center. Moreover, the results still  
758 persisted in a more reflexive behavioral task, in which prolonged fixation was not enforced  
759 in the face of a salient eccentric stimulus onset.

760

761

762 *Different temporal dynamics of firing rates in the sustained interval for darks*  
763 *and brights*

764 The results so far have focused on initial visual bursts. However, with prolonged fixation (as  
765 in our primary task of Figs. 1-7), we also observed significant differences in SC neural  
766 response dynamics in the sustained interval (long after stimulus onset) for bright and dark  
767 stimuli. In particular, bright stimuli were generally associated with secondary elevations of  
768 firing rate above those of dark stimuli. To illustrate this, Fig. 9A, B shows the responses of

769 four example neurons to high contrast stimuli (100%). Two neurons are from monkey A (Fig.  
770 9A), and two neurons are from monkey M (Fig. 9B). In the first three neurons (neurons 5-7 in  
771 Fig. 9A, B), after the initial visual bursts, bright stimuli evoked stronger sustained activity  
772 than dark stimuli (see pink intervals highlighting the sustained interval). The stimuli were still  
773 present within the RF's of the neurons in all cases, but there was an altered response  
774 dynamic after the initial visual bursts, particularly for bright stimuli. Even the fourth neuron  
775 (neuron 8 in Fig. 9B), which showed relatively weak sustained activity, still showed a  
776 subdued secondary peak in firing rate after the initial visual burst for bright stimuli (also see  
777 Fig. 11 below for more details on monkey M's secondary bursts for bright stimuli).

778

779 To characterize this altered dynamic of neural responses as a function of time in more detail,  
780 we took each firing rate curve after 80 ms from stimulus onset (that is, after the initial visual  
781 bursts). We then estimated contrast sensitivity curves at each time point. Each time sample  
782 of a firing rate curve is already a kind of average over some discrete measurement interval  
783 (due to the convolution of spike times with a gaussian kernel to generate firing rates).  
784 Therefore, we took each sample of the firing rate curve of a neuron in the sustained interval,  
785 and we used it to fit contrast sensitivity curves from equation 1 at each time point. This gave  
786 us a series of contrast sensitivity curves as a function of time. We then plotted the time  
787 courses of the parameters  $R$ ,  $C50$ , and  $N$  of the fits during the sustained interval, and we did  
788 this for either bright or dark stimuli. The results across the entire population of neurons are  
789 shown in Fig. 9C. As can be seen, all parameters were varying differently between darks and  
790 brights in the interval around approximately 100-200 ms after stimulus onset (that is, during  
791 sustained visual response intervals), consistent with the example neurons of Fig. 9A, B. The  
792 biggest effect was in the  $R$  parameter, which was stronger for brights than darks, suggesting

793 higher sustained firing rates for brights after the ends of the initial visual bursts. Both  $C50$   
794 and  $N$  gradually changed in a manner that was consistent with higher thresholds and  
795 shallower contrast sensitivity functions in the sustained interval. That is, the contrast  
796 sensitivity of the neurons was generally the highest in the initial visual burst intervals, and it  
797 gradually degraded in sustained intervals (other than the  $R$  parameter elevation for bright  
798 stimuli). This makes sense given that sustained intervals were generally associated with  
799 much lower firing rates than in the initial visual burst intervals (and were therefore less likely  
800 to be strongly differentially modulated by stimulus properties). In any case, during the  
801 sustained interval, and unlike in the initial phases of SC visual responses, there was a  
802 generalized elevation of firing rates for bright stimuli compared to dark stimuli for all  
803 contrasts. As we show later, this effect was strong enough in monkey M, to the extent that it  
804 appeared to dominate this monkey's saccadic reaction time patterns in the immediate,  
805 visually-guided saccade version of the task.

806

807

808 *Earlier detection of dark stimuli also by inhibited SC neurons*

809 In all of the above analyses, we focused solely on neurons exhibiting positive visual  
810 responses (that is, increases in firing rates above baseline). However, with our offline neuron  
811 sorting pipelines (Methods), we also isolated a fewer number of neurons that exhibited  
812 transient decreases in activity after stimulus onset rather than increases. These neurons  
813 were obtained from similar recording sites to those from which we isolated neurons with  
814 visual bursts (we used linear electrode arrays primarily orthogonal to the SC surface;  
815 Methods). The neurons were, therefore, from similar topographic locations to those  
816 associated with the neurons reported in Figs. 1-9. When we analyzed these inhibited

817 neurons in more detail, we found that their transient, stimulus-induced decreases in firing  
818 rates were still sensitive to luminance polarity. For example, in Fig. 10A, B, we show two  
819 example neurons from one of our electrode penetrations in monkey M. The two neurons  
820 showed classic visual responses (to black stimuli); moreover, their RF's (shown in the insets  
821 for data collected with the presentation of black small spots during fixation) were spatially  
822 localized and overlapping with each other. From the very same electrode penetration, Fig.  
823 10C shows a third sample neuron that was recorded simultaneously with the two other  
824 neurons; it was thus in the same SC topographic region as the two neurons of Fig. 10A, B.  
825 The neuron of Fig. 10C was inhibited instead. Most interestingly, this neuron clearly  
826 "responded" to a high contrast dark stimulus earlier than to a bright stimulus of the same  
827 contrast, with the only difference from the results of Figs. 1-9 being that the response in this  
828 case was a transient reduction from baseline activity rather than an increase. Across the  
829 population of such inhibited neurons (n=15 neurons), we repeated the same latency  
830 analyses of Fig. 4 above. That is, we assessed the relative time of "response" between bright  
831 and dark contrasts (Methods). As can be seen from Fig. 10D, the majority of such inhibited  
832 neurons also reacted to dark stimuli earlier than to bright stimuli, just like with the neurons  
833 possessing visual bursts. Similar observations were also made for the lower contrast stimuli.  
834 Interestingly, all 15 inhibited neurons had their "response" to stimulus onset slightly later  
835 than classic visual bursts in other neurons (compare the visual bursts in Fig. 10A, B to the  
836 inhibition time in Fig. 10C; the inhibition occurred slightly later than the bursts). Therefore,  
837 even inhibited neurons in the SC detected dark contrasts faster than bright contrasts.  
838  
839  
840

841 *Saccadic reaction times can be significantly shorter for dark stimuli*

842 Finally, prior work has demonstrated a tight relationship between SC visual response  
843 properties and saccadic reaction times (Boehnke and Munoz, 2008; Marino et al., 2012;  
844 Marino et al., 2015; Hafed and Chen, 2016; Chen et al., 2018). Specifically, both visual  
845 response sensitivity (Boehnke and Munoz, 2008; Marino et al., 2012; Marino et al., 2015;  
846 Hafed and Chen, 2016; Chen et al., 2018) and visual response latency (Chen et al., 2018) can  
847 predict such reaction times. Therefore, given the faster response latencies of SC neurons for  
848 dark stimuli that we found, we wondered whether this effect was sufficient to be associated  
849 with faster saccadic reaction times to such stimuli (even when response sensitivity was, on  
850 average, similar for darks and brights, as shown in Fig. 2; or even slightly favoring brights at  
851 high contrasts, as shown in Fig. 6). We tested our two monkeys and a third one on the  
852 immediate visually-guided saccade task; we used the same sessions as in Fig. 8 for monkeys  
853 M and A, and we ran separate behavior-only sessions for monkey F. The monkeys simply  
854 generated a saccade as soon as the target appeared (the fixation spot also disappeared at  
855 target onset, as mentioned above for Fig. 8 and in Methods). We measured saccadic reaction  
856 times and plotted them as a function of stimulus contrast and stimulus luminance polarity.

857

858 All monkeys showed faster reaction times for higher contrast stimuli, as expected (Marino et  
859 al., 2012; Marino et al., 2015) ( $p < 3 \times 10^{-103}$  in each monkey individually, Kruskal-Wallis test  
860 exploring the effect of contrast on reaction time, when collapsing across luminance  
861 polarities). Interestingly, two out of the three monkeys (A and F) also showed consistently  
862 faster reaction times for the darker stimuli, like with the SC visual bursts. These results are  
863 shown in Fig. 11A, C, E; monkeys A and F were faster to react to dark stimuli at all contrasts.  
864 Monkey M, on the other hand, had faster reaction times for the bright stimuli (Fig. 11C). All

865 of these results (of luminance polarity effects on reaction times) were significant ( $p < 2.5 \times 10^{-8}$ )  
866 in each monkey individually, Kruskal-Wallis test exploring the relationship between  
867 luminance polarity and reaction time, when collapsing across contrasts); the effect sizes  
868 (shown for each condition in Fig. 11A, C, E) were also substantial. In addition, the effect sizes  
869 in monkeys A and F were of the same order of magnitude as the effect sizes of the visual  
870 response latency differences between darks and brights seen in Fig. 5E.

871  
872 We were particularly intrigued by the discrepancy in the reaction times of monkey M with  
873 respect to dark and bright stimuli. On the one hand, it might suggest that SC visual response  
874 latency (e.g. Figs. 4, 5) is not the only determinant of saccadic reaction times, which is  
875 indeed plausible. For example, we earlier found that SC visual response latency and visual  
876 response sensitivity together provided a better correlate of reaction times than either  
877 parameter alone (Chen et al., 2018). Therefore, since about half of the neurons in our  
878 population were more sensitive to bright stimuli anyway (Fig. 2), despite the faster detection  
879 of darks, it could be that this particular monkey's reaction times were more dictated by SC  
880 visual response sensitivity than by visual response latency. On the other hand, it could  
881 additionally be the case that the later elevation of responses for bright stimuli that we saw in  
882 Fig. 9 was more pronounced in this monkey, potentially suggesting stronger top-down  
883 control for bright stimuli. In that case, bright stimuli could be preferentially processed by this  
884 monkey. Indeed, in a previous behavioral study in which we investigated the properties of  
885 saccadic inhibition as a function of luminance contrast polarity, this monkey reacted  
886 differently to full field white versus black visual flashes from the two other monkeys in the  
887 very initial oculomotor response to flash onset, again reacting faster for bright than dark  
888 flashes (Malevich et al., 2021) (see their Fig. 3). Therefore, we decided to check how this

889 monkey's neurons, in particular, reacted to white stimuli long after the initial visual bursts,  
890 and we were able to do so from our fixation variant of the task.

891  
892 We plotted each monkey's population visual responses for dark and bright stimuli in the  
893 fixation variant of the task, allowing us to explore the longer sustained interval. These results  
894 are shown in Fig. 11B, D. Even though monkey M's neurons still detected dark stimuli earlier  
895 than bright stimuli in the initial visual response period (consistent with all of our results  
896 shown earlier), this monkey's elevation of sustained visual activity for the bright stimuli (e.g.  
897 Fig. 9) was particularly pronounced when compared to monkey A (note the secondary peak  
898 in population firing rate for bright stimuli in Fig. 11D for monkey M, which was stronger than  
899 the same peak in monkey A). We also even saw hints of this secondary elevation in Fig. 8B in  
900 the immediate, visually-guided saccade variant of the task, with a sharper elevation for  
901 bright stimuli right after the initial visual burst and leading up to the saccade-related burst;  
902 however, of course, in this task, this sharper elevation for brights was harder to properly  
903 analyze in the saccade task because of how quickly the motor burst came.

904  
905 Therefore, the results of Fig. 11 suggest that saccadic reaction times can indeed be faster for  
906 dark than bright stimuli, consistent with the faster detection of dark stimuli by SC neurons,  
907 and that even violations of such an observation (as in the case of monkey M) are still related  
908 to the SC visual responses (in this case, the sustained responses after the initial visual bursts  
909 subside).

910  
911 In all, our results in this study indicate that SC neurons robustly detect dark stimuli faster  
912 than bright stimuli; that sustained visual responses in the SC instead favor bright stimuli; and

913 that saccadic reaction times can reflect the faster detection of dark stimuli in the SC's initial  
914 visual bursts and/or the later elevation for bright stimuli.

915

916

917 **Discussion**

918

919 We evaluated the sensitivity of monkey SC neurons to luminance contrast polarity. We  
920 found that there was a diversity of preferences for darks and brights across the population  
921 (Figs. 1, 2). However, irrespective of preference (as defined by visual neural sensitivity), most  
922 neurons detected dark contrasts earlier than bright contrasts (Figs. 3-8, 10). Such earlier  
923 detection of dark stimuli was correlated with faster reaction times for such stimuli in 2 out of  
924 3 monkeys (Fig. 11). And, even in the third monkey, this monkey's observed opposite  
925 reaction time effect could be related to sustained modulations of SC neural activity, which  
926 exhibited a strong secondary elevation particularly for bright stimuli after the initial visual  
927 bursts (Figs. 9, 11).

928

929 Our results demonstrate that the primate SC does not necessarily exhibit identical ON/OFF  
930 sensitivity asymmetries for brights and darks as LGN and V1, refuting the idea that the  
931 primate SC simply inherits its visual properties from V1. For example, V1 neurons mostly  
932 prefer dark contrasts (Yeh et al., 2009), unlike in our SC population, and it would be  
933 interesting to further investigate whether deep V1 layers, projecting to the SC, violate this  
934 property or not. In fact, at high contrasts, our SC neurons significantly preferred bright,  
935 rather than dark, stimuli even while having faster response latencies to the dark ones (Fig.  
936 6A, F, L).

937

938 Our results are additionally interesting because they add to a growing literature  
939 demonstrating that the primate SC is as visual a brain structure as the SC in other species,  
940 like mice, in which the SC is the primary recipient of retinal projections and, indeed, a

941 primary visual structure. Consistent with this idea, the monkey SC receives a large amount of  
942 cortical visual input (Kadoya et al., 1971; Fries, 1984; Lui et al., 1995; Lock et al., 2003;  
943 Cerkevich et al., 2014), in addition to direct retinal input (Perry and Cowey, 1984). Thus, the  
944 SC in primates should be viewed as being even more visual than, say, the mouse SC. Such a  
945 rich visual nature of the primate SC matters a great deal for orienting responses, consistent  
946 with how SC visual responses can be linked to various aspects of saccadic behavior, like  
947 reaction time (Boehnke and Munoz, 2008; Marino et al., 2012; Marino et al., 2015; Hafed  
948 and Chen, 2016; Chen et al., 2018) and landing accuracy (Hafed and Chen, 2016). Such a link  
949 to saccadic behaviors was also clearly still evident in our current study (e.g. Fig. 11). In the  
950 future, it would be important to relate trial-to-trial variability in saccadic reaction times to  
951 trial-to-trial variability in SC, LGN, and V1 visual responses, as was done previously in V1 (Lee  
952 et al., 2010), to better appreciate the different functional specializations that exist in early  
953 visual responses that occur in multiple brain areas at approximately the same time.

954

955 Our observation that the primate SC can represent dark contrasts well (e.g. Fig. 2) is also  
956 consistent with earlier observations that SC neurons detect dark “shadows” (Humphrey,  
957 1968; Cynader and Berman, 1972; Updyke, 1974). We are additionally particularly intrigued  
958 by the earlier response latencies for dark stimuli that we observed (e.g. Fig. 4), as well as by  
959 the altered temporal dynamics of responses during the sustained interval long after stimulus  
960 onsets (e.g. Fig. 9). These observations could potentially be used to further interpret earlier  
961 reports in the literature about SC visual and visual-motor modulations. For example, in  
962 investigating color-related responses in the SC, White and colleagues used a high contrast  
963 black target as the comparison stimulus to the colored ones (White et al., 2009). Because of  
964 that black stimulus, we predict that the latency differences that these authors observed

965 relative to colored targets were slightly amplified than what they would have observed had  
966 they used a white target as the reference non-colored stimulus. Similarly, Churan and  
967 colleagues investigated how SC visual RF's were modified around the time of saccades  
968 (Churan et al., 2011). They found dramatically different effects depending on whether the  
969 saccades were made across a gray background or across a dark background. It is intriguing to  
970 consider whether (and how) our sustained response effects, amplifying responses for bright  
971 stimuli (e.g. Fig. 9), could be related to their observations.

972

973 The fact that SC neurons can be strongly sensitive to dark contrasts is also interesting with  
974 respect to spatial frequency tuning in SC neurons. In recent work, we found that SC neurons  
975 can be sensitive to minute phase shifts of spatial frequency gratings, as small as 1 minute of  
976 arc in amplitude (Hafed et al., 2022). It would be fruitful, in light of these observations and  
977 the current work, to investigate RF subfield structure in more detail, for example, to study  
978 phase tuning in SC neurons. Indeed, both the current work and these recent results motivate  
979 a detailed mapping of RF's with both bright and dark stimuli, to assess asymmetries beyond  
980 just visual response sensitivity and visual response latency. Indeed, prior work with reverse  
981 correlation techniques has suggested that there may be informative observations to be  
982 made about RF's mapped with bright versus dark stimuli (Churan et al., 2012). In the near  
983 future, we hope to report SC RF maps for bright and dark stimuli in detail.

984

985 The altered long term temporal dynamics of firing rates as a function of luminance polarity  
986 that we observed (e.g. Fig. 9) also motivate modeling how these dynamics emerge. In V1,  
987 various stimulus factors, like contrast, alter not only the initial visual bursts (as might be  
988 expected), but also the sustained responses. Moreover, such alterations can be modeled

989 using variants of linear/nonlinear filters and divisive normalization (Groen et al., 2021). It  
990 would be valuable to investigate such models in the SC, and to relate them to asymmetries  
991 in ON and OFF channels in both the LGN (Jin et al., 2011) and V1 (Komban et al., 2014).

992  
993 Related to this, we would like to investigate, in the near future, how scene statistical  
994 regularities, with respect to orienting eye movements, can allow further expansion of our  
995 upper versus lower visual field analyses of Fig. 7. In these analyses, we were motivated by  
996 the theoretical framework of (Previc, 1990), in which he predicted that SC neurons should  
997 over-represent the upper visual field of retinal images because eye movements are relevant  
998 for sampling extra-foveal visual space. Thus, in the analyses of Fig. 7, we were driven by our  
999 earlier discoveries of significant asymmetries between upper and lower visual field SC  
1000 neurons and saccadic performance (Hafed and Chen, 2016; Hafed and Goffart, 2020; Hafed,  
1001 2021; Fracasso et al., 2022). Indeed, we found that dark versus bright asymmetries were  
1002 amplified in the upper visual field (Fig. 7), and this is ecologically sensible. For example, birds  
1003 in the sky normally cast shadows on the retina. However, it would be even more intriguing  
1004 to go even deeper when assessing such visual field anisotropies. For example, one could  
1005 consider studying SC binocularity in more detail, to investigate whether neurons preferring  
1006 far disparities would be more prevalent in the upper visual field representation of the SC or  
1007 not. And, if so, would these far-preferring neurons also prefer more dark contrasts, like in  
1008 the case of V1 (Samonds et al., 2012)? This is important to consider, especially given how our  
1009 neurons seemed to prefer bright stimuli in a contrast-dependent manner (Fig. 6F-K) that was  
1010 the opposite of what might be predicted from natural scene statistics (Cooper and Norcia,  
1011 2015).

1012

1013 Finally, we found a small subset of neurons, in the same topographic location as our bursting  
1014 neurons, that were inhibited by stimulus onset (Fig. 10). Interestingly, these neurons still  
1015 “detected” dark contrasts (by their transient inhibition of firing rate) earlier than bright  
1016 contrasts. It would be important to investigate whether such neurons contribute to saccadic  
1017 inhibition (Reingold and Stampe, 2002; Buonocore and McIntosh, 2008; Hafed and  
1018 Ignashchenkova, 2013), which we recently found to also depend on stimulus luminance  
1019 polarity (Malevich et al., 2021). Our initial intuition with regard to saccadic inhibition,  
1020 described in detail in our theoretical proposal elsewhere (Hafed et al., 2021), is that  
1021 structures beyond the SC are critical for this phenomenon. However, this does not deny the  
1022 potential involvement of SC neurons (particularly those neurons that are transiently  
1023 inhibited by stimulus onsets), and future research should investigate the mechanisms of  
1024 saccadic inhibition in much more detail, including recording SC neurons with full or localized  
1025 flashes of different luminance polarities like in psychophysics. Critical in those studies would  
1026 be to quantitatively assess whether the small latency differences in “inhibition” versus  
1027 excitatory visual “bursts” that we observed in Fig. 10 are consistent with the timing  
1028 properties of saccadic inhibition or not.

1029

1030

1031 **References**

1032

1033 Bellet ME, Bellet J, Nienborg H, Hafed ZM, Berens P (2019) Human-level saccade detection  
1034 performance using deep neural networks. *J Neurophysiol* 121:646-661.

1035 Boehnke SE, Munoz DP (2008) On the importance of the transient visual response in the  
1036 superior colliculus. *Curr Opin Neurobiol* 18:544-551.

1037 Bogadhi AR, Hafed ZM (2022) Express detection and discrimination of visual objects by  
1038 primate superior colliculus neurons. *BioRxiv*.

1039 Bogadhi AR, Buonocore A, Hafed ZM (2020) Task-irrelevant visual forms facilitate covert and  
1040 overt spatial selection. *J Neurosci* 40:9496-9506.

1041 Brainard DH (1997) The Psychophysics Toolbox. *Spatial vision* 10:433-436.

1042 Buonocore A, McIntosh RD (2008) Saccadic inhibition underlies the remote distractor effect.  
1043 *Exp Brain Res* 191:117-122.

1044 Buonocore A, Skinner J, Hafed ZM (2019) Eye Position Error Influence over "Open-Loop"  
1045 Smooth Pursuit Initiation. *J Neurosci* 39:2709-2721.

1046 Cerkevich CM, Lyon DC, Balaram P, Kaas JH (2014) Distribution of cortical neurons projecting  
1047 to the superior colliculus in macaque monkeys. *Eye and brain* 2014:121-137.

1048 Chen CY, Hafed ZM (2013) Postmicrosaccadic enhancement of slow eye movements. *The  
1049 Journal of neuroscience : the official journal of the Society for Neuroscience* 33:5375-  
1050 5386.

1051 Chen CY, Hafed ZM (2017) A neural locus for spatial-frequency specific saccadic suppression  
1052 in visual-motor neurons of the primate superior colliculus. *J Neurophysiol* 117:1657-  
1053 1673.

1054 Chen CY, Ignashchenkova A, Thier P, Hafed ZM (2015) Neuronal Response Gain  
1055 Enhancement prior to Microsaccades. *Curr Biol* 25:2065-2074.

1056 Chen CY, Hoffmann KP, Distler C, Hafed ZM (2019) The Foveal Visual Representation of the  
1057 Primate Superior Colliculus. *Curr Biol* 29:2109-2119 e2107.

1058 Chen CY, Sonnenberg L, Weller S, Witschel T, Hafed ZM (2018) Spatial frequency sensitivity  
1059 in macaque midbrain. *Nat Commun* 9:2852.

1060 Chichilnisky EJ, Kalmar RS (2002) Functional asymmetries in ON and OFF ganglion cells of  
1061 primate retina. *J Neurosci* 22:2737-2747.

1062 Churan J, Guitton D, Pack CC (2011) Context dependence of receptive field remapping in  
1063 superior colliculus. *J Neurophysiol* 106:1862-1874.

1064 Churan J, Guitton D, Pack CC (2012) Spatiotemporal structure of visual receptive fields in  
1065 macaque superior colliculus. *J Neurophysiol* 108:2653-2667.

1066 Cooper EA, Norcia AM (2015) Predicting cortical dark/bright asymmetries from natural  
1067 image statistics and early visual transforms. *PLoS Comput Biol* 11:e1004268.

1068 Cynader M, Berman N (1972) Receptive-field organization of monkey superior colliculus. *J  
1069 Neurophysiol* 35:187-201.

1070 De Franceschi G, Solomon SG (2018) Visual response properties of neurons in the superficial  
1071 layers of the superior colliculus of awake mouse. *J Physiol* 596:6307-6332.

1072 Eastman KM, Huk AC (2012) PLDAPS: A Hardware Architecture and Software Toolbox for  
1073 Neurophysiology Requiring Complex Visual Stimuli and Online Behavioral Control.  
1074 *Front Neuroinform* 6:1.

1075 Fracasso A, Buonocore A, Hafed ZM (2022) Peri-saccadic visual sensitivity is higher in the  
1076 upper visual field. *BioRxiv*.

1077 Fries W (1984) Cortical projections to the superior colliculus in the macaque monkey: a  
1078 retrograde study using horseradish peroxidase. *J Comp Neurol* 230:55-76.

1079 Fuchs AF, Robinson DA (1966) A method for measuring horizontal and vertical eye  
1080 movement chronically in the monkey. *J Appl Physiol* 21:1068-1070.

1081 Groen IIA, Pinantoni G, Montenegro S, Flinker A, Devore S, Devinsky O, Doyle W, Dugan P,  
1082 Friedman D, Ramsey N, Petridou N, Winawer J (2021) Temporal dynamics of neural  
1083 responses in human visual cortex. *BioRxiv*.

1084 Hafed ZM (2021) Superior colliculus saccade motor bursts do not dictate movement  
1085 kinematics. *BioRxiv*.

1086 Hafed ZM, Krauzlis RJ (2010) Microsaccadic suppression of visual bursts in the primate  
1087 superior colliculus. *J Neurosci* 30:9542-9547.

1088 Hafed ZM, Ignashchenkova A (2013) On the dissociation between microsaccade rate and  
1089 direction after peripheral cues: microsaccadic inhibition revisited. *J Neurosci*  
1090 33:16220-16235.

1091 Hafed ZM, Chen CY (2016) Sharper, Stronger, Faster Upper Visual Field Representation in  
1092 Primate Superior Colliculus. *Curr Biol* 26:1647-1658.

1093 Hafed ZM, Goffart L (2020) Gaze direction as equilibrium: more evidence from spatial and  
1094 temporal aspects of small-saccade triggering in the rhesus macaque monkey. *J  
1095 Neurophysiol* 123:308-322.

1096 Hafed ZM, Chen C-Y, Khademi F (2022) Superior colliculus visual neural sensitivity at the  
1097 lower limit of natural self-induced image displacements. *BioRxiv*.

1098 Hafed ZM, Yoshida M, Tian X, Buonocore A, Malevich T (2021) Dissociable Cortical and  
1099 Subcortical Mechanisms for Mediating the Influences of Visual Cues on Microsaccadic  
1100 Eye Movements. *Front Neural Circuits* 15:638429.

1101 Hartline HK (1938) The response of single optic nerve fibers of the vertebrate eye to  
1102 illumination of the retina. *American Journal of Physiology* 121:400-415.

1103 Hubel DH, Wiesel TN (1961) Integrative action in the cat's lateral geniculate body. *J Physiol*  
1104 155:385-398.

1105 Humphrey NK (1968) Responses to visual stimuli of units in the superior colliculus of rats and  
1106 monkeys. *Exp Neurol* 20:312-340.

1107 Jiang Y, Purushothaman G, Casagrande VA (2015) The functional asymmetry of ON and OFF  
1108 channels in the perception of contrast. *J Neurophysiol* 114:2816-2829.

1109 Jin J, Wang Y, Lashgari R, Swadlow HA, Alonso JM (2011) Faster thalamocortical processing  
1110 for dark than light visual targets. *J Neurosci* 31:17471-17479.

1111 Jin JZ, Weng C, Yeh CI, Gordon JA, Ruthazer ES, Stryker MP, Swadlow HA, Alonso JM (2008)  
1112 On and off domains of geniculate afferents in cat primary visual cortex. *Nat Neurosci*  
1113 11:88-94.

1114 Judge SJ, Richmond BJ, Chu FC (1980) Implantation of magnetic search coils for  
1115 measurement of eye position: an improved method. *Vision Res* 20:535-538.

1116 Kadoya S, Massopust LC, Jr., Wolin LR (1971) Striate cortex-superior colliculus projection in  
1117 squirrel monkey. *Exp Neurol* 32:98-110.

1118 Kleiner M, Brainard D, Pelli DG (2007) What's new in Psychtoolbox-3? (Abstract). *Perception*  
1119 36.

1120 Komban SJ, Alonso JM, Zaidi Q (2011) Darks are processed faster than lights. *J Neurosci*  
1121 31:8654-8658.

1122 Komban SJ, Kremkow J, Jin J, Wang Y, Lashgari R, Li X, Zaidi Q, Alonso JM (2014) Neuronal  
1123 and perceptual differences in the temporal processing of darks and lights. *Neuron*  
1124 82:224-234.

1125 Kremkow J, Jin J, Komban SJ, Wang Y, Lashgari R, Li X, Jansen M, Zaidi Q, Alonso JM (2014)  
1126 Neuronal nonlinearity explains greater visual spatial resolution for darks than lights.  
1127 Proc Natl Acad Sci U S A 111:3170-3175.

1128 Lee J, Kim HR, Lee C (2010) Trial-to-trial variability of spike response of V1 and saccadic  
1129 response time. J Neurophysiol 104:2556-2572.

1130 Li X, Basso MA (2008) Preparing to move increases the sensitivity of superior colliculus  
1131 neurons. The Journal of neuroscience : the official journal of the Society for  
1132 Neuroscience 28:4561-4577.

1133 Liu K, Yao H (2014) Contrast-dependent OFF-dominance in cat primary visual cortex  
1134 facilitates discrimination of stimuli with natural contrast statistics. Eur J Neurosci  
1135 39:2060-2070.

1136 Lock TM, Baizer JS, Bender DB (2003) Distribution of corticotectal cells in macaque. Exp Brain  
1137 Res 151:455-470.

1138 Lui F, Gregory KM, Blanks RH, Giolli RA (1995) Projections from visual areas of the cerebral  
1139 cortex to pretectal nuclear complex, terminal accessory optic nuclei, and superior  
1140 colliculus in macaque monkey. J Comp Neurol 363:439-460.

1141 Malevich T, Buonocore A, Hafed ZM (2020) Rapid stimulus-driven modulation of slow ocular  
1142 position drifts. eLife 9.

1143 Malevich T, Buonocore A, Hafed ZM (2021) Dependence of the stimulus-driven microsaccade  
1144 rate signature in rhesus macaque monkeys on visual stimulus size and polarity. J  
1145 Neurophysiol 125:282-295.

1146 Marino RA, Levy R, Munoz DP (2015) Linking express saccade occurrence to stimulus  
1147 properties and sensorimotor integration in the superior colliculus. J Neurophysiol  
1148 114:879-892.

1149 Marino RA, Levy R, Boehnke S, White BJ, Itti L, Munoz DP (2012) Linking visual response  
1150 properties in the superior colliculus to saccade behavior. Eur J Neurosci 35:1738-  
1151 1752.

1152 Pachitariu M, Steinmetz NA, Kadir SN, Carandini M, Harris KD (2016) Fast and accurate spike  
1153 sorting of high-channel count probes with KiloSort. Advances in Neural Information  
1154 Processing Systems (NIPS 2016) 29.

1155 Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming numbers  
1156 into movies. Spatial vision 10:437-442.

1157 Perry VH, Cowey A (1984) Retinal ganglion cells that project to the superior colliculus and  
1158 pretectum in the macaque monkey. Neuroscience 12:1125-1137.

1159 Previc FH (1990) Functional Specialization in the Lower and Upper Visual-Fields in Humans -  
1160 Its Ecological Origins and Neurophysiological Implications. Behavioral and Brain  
1161 Sciences 13:519-575.

1162 Reingold EM, Stampe DM (2002) Saccadic inhibition in voluntary and reflexive saccades. J  
1163 Cogn Neurosci 14:371-388.

1164 Robinson DA (1972) Eye movements evoked by collicular stimulation in the alert monkey.  
1165 Vision Res 12:1795-1808.

1166 Samonds JM, Potetz BR, Lee TS (2012) Relative luminance and binocular disparity  
1167 preferences are correlated in macaque primary visual cortex, matching natural scene  
1168 statistics. Proc Natl Acad Sci U S A 109:6313-6318.

1169 Schiller PH, Sandell JH, Maunsell JH (1986) Functions of the ON and OFF channels of the  
1170 visual system. Nature 322:824-825.

1171 Skinner J, Buonocore A, Hafed ZM (2019) Transfer function of the rhesus macaque  
1172 oculomotor system for small-amplitude slow motion trajectories. *J Neurophysiol*  
1173 121:513-529.  
1174 Tian X, Yoshida M, Hafed ZM (2018) Dynamics of fixational eye position and microsaccades  
1175 during spatial cueing: the case of express microsaccades. *J Neurophysiol* 119:1962-  
1176 1980.  
1177 Updyke BV (1974) Characteristics of unit responses in superior colliculus of the *Cebus*  
1178 monkey. *J Neurophysiol* 37:896-909.  
1179 Wang L, Sarnaik R, Rangarajan K, Liu X, Cang J (2010) Visual receptive field properties of  
1180 neurons in the superficial superior colliculus of the mouse. *J Neurosci* 30:16573-  
1181 16584.  
1182 White BJ, Boehnke SE, Marino RA, Itti L, Munoz DP (2009) Color-related signals in the  
1183 primate superior colliculus. *J Neurosci* 29:12159-12166.  
1184 Yeh CI, Xing D, Shapley RM (2009) "Black" responses dominate macaque primary visual  
1185 cortex v1. *J Neurosci* 29:11753-11760.  
1186


1187

1188

1189

1190

1191 **Figure legends**




1192

1193 **Figure 1 Diversity of preferences, in terms of visual burst strength, for dark and bright stimuli by SC neurons.**  
1194 **(A-C)** Visual responses of three example neurons from monkey A to a 100% contrast stimulus appearing within  
1195 their response fields (RF's). Black indicates that the stimulus was negative in luminance polarity (darker than the  
1196 background); gray indicates that the stimulus was brighter than the background. Each row of tick marks indicates  
1197 a single trial, and each tick mark indicates the time of an action potential. The firing rate plots below the raster  
1198 plots summarize the neurons' firing rates. Neuron 1 had a higher sensitivity to bright stimuli, whereas neuron 2  
1199 was equally sensitive to dark and bright stimuli. Neuron 3, on the other hand, clearly preferred dark stimuli. Note  
1200 that response latency (that is, when the stimulus-evoked action potentials first appeared) was always shorter for  
1201 dark stimuli (see subsequent analyses). Error bars denote s.e.m. across trials. **(D-F)** For each neuron, we  
1202 measured peak average firing rate after stimulus onset (individual symbols), and we plotted it as a function of  
1203 stimulus Weber contrast. We also fit the data with continuous curves (Methods). Neuron 1 had higher contrast  
1204 sensitivity for bright stimuli, evidenced by the higher plateau firing rate at maximal contrast. Neuron 2 plateaued  
1205 at the same firing rate for both dark and bright stimuli, and neuron 3 was more sensitive to dark stimuli. Dashed  
1206 vertical lines indicate  $C_{50}$ , the semi-saturation contrast of each neuron (Methods).

1207


1208



1209


1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226

**Figure 2 Diversity of contrast sensitivity curve parameters for dark and bright stimuli across SC neurons. (A)** For each neuron, we compared the neuron's visual response dynamic range (parameter  $R$  in equation 1; Methods) between dark and bright stimuli. A modulation index value of 1 indicates maximal responsiveness to only bright stimuli, and a value of -1 indicates maximal responsiveness only to dark stimuli. Most neurons responded to both stimuli (note the rarity of  $+/-1$  modulation index values), but with varying degrees of sensitivity; some neurons clearly preferred dark stimuli, whereas others clearly preferred bright stimuli (similar to the examples in Fig. 1). The vertical lines show mean (solid) and median (dashed) modulation index values across the population, and they were both close to 0: across the population, SC visual responses were equally sensitive to dark and bright stimuli. **(B)** Similar observations for the semi-saturation contrast ( $C_{50}$ ) parameter. Neurons with positive modulation indices in this case are neurons with higher semi-saturation contrasts for bright stimuli (that is, they were less sensitive to bright than dark stimuli). Again, the population average and median semi-saturation contrasts were largely similar between darks and brights (vertical lines), but with large variability across individual neurons. **(C)** Similar observations for the slope parameter of the contrast sensitivity curves. These results are consistent with the example neurons of Fig. 1.




1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238

**Figure 3 Earlier visual bursts for dark than bright stimuli at different stimulus contrasts. (A-C)** Visual responses of the three example neurons of Fig. 1 at each of our tested contrasts (different columns). Black curves indicate dark stimuli, and gray curves indicate bright stimuli. Reducing contrast expectedly weakened and delayed visual responses for both dark and bright stimuli (compare firing rates across columns; also see Fig. 1D-F). Note, however, that at each contrast level, visual bursts occurred slightly earlier for dark than bright stimuli, even for neuron 1, which was more sensitive to bright stimuli. Also note that at the weakest contrast level (5%), both neuron 1 (A) and neuron 3 (C) were more sensitive to light decrements than light increments. We quantify these observations in subsequent figures and analyses. Error bars denote s.e.m. across trials.



1239  
1240


1241 **Figure 4 Faster detection of darks than brights by monkey superior colliculus neurons.** (A) For the highest  
1242 stimulus contrast, we estimated visual response latency (Methods) separately for dark and bright stimuli in each  
1243 neuron. We then subtracted, for each neuron, the visual response latency for dark stimuli from the visual  
1244 response latency for bright stimuli (positive would indicate earlier responses for dark stimuli). We then sorted all  
1245 neurons based on this difference (top panel). In the bottom panel, we used the very same sorting, but we now  
1246 plotted the difference in peak visual response strength between bright and dark stimuli (Methods). Most neurons  
1247 had a shorter visual response latency for dark than bright stimuli (top panel; vertical line shows the sorted neuron  
1248 index at which response latency differences flipped sign from negative to positive). This happened independently  
1249 of response strength; the bottom panel (with the same sorting) did not show a systematic ordering. For example,  
1250 the neurons highlighted with diagonal arrows preferred bright stimuli (bottom panel) but still detected dark  
1251 stimuli earlier (top panel). (B-E) Similar results for lower contrasts. Of course, with lower and lower contrasts,  
1252 the earlier detection of darks was less and less prevalent (see the crossover points in the top row). However, this  
1253 was because lower contrasts were already associated with delayed and weakened visual responses (see Fig. 5).  
1254 Also note that the lower row shows a decreasing likelihood of bright-preferring neurons as contrast level  
1255 decreases, suggesting a contrast-dependent processing of darks and brights in SC neurons (see Fig. 6).  
1256  
1257



1258  
1259

1260 **Figure 5 Interaction between contrast and luminance polarity in SC visual response latencies.** (A) Visual  
1261 responses of an example SC neuron from monkey M for high (100%) and low (5%) contrast bright stimuli (that  
1262 is, with luminance higher than the background luminance). Each row of tick marks represents a single trial  
1263 repetition, and each tick mark indicates the time of an action potential. The neuron responded earlier to the high  
1264 contrast stimulus. (B) In the same neuron, the visual responses to a dark 100% contrast stimulus (that is, with  
1265 luminance darker than the background luminance) occurred even earlier than the responses to a bright 100%  
1266 contrast stimulus. Thus, both contrast and luminance polarity affected the neuron's visual response latency. (C)  
1267 Cumulative histograms of our estimates of visual response latency across all of our neurons, for both stimulus  
1268 contrast (different colors) and stimulus luminance contrast polarity (solid versus dashed lines). High contrasts  
1269 were associated with earlier visual response latencies in the SC. In addition, at each contrast, dark stimuli were  
1270 systematically associated with earlier visual response latencies. (D) Average visual response latencies for brights  
1271 and darks across contrast levels, demonstrating consistently faster responses for dark stimuli. (E) Average  
1272 differences in visual response latencies between responses for bright and dark stimuli per contrast level. All  
1273 contrast levels were associated with faster detection of dark than bright stimuli. The effect increased in strength  
1274 with increasing contrast from 10% to 100%. At the 5% contrast condition, the effect was the strongest, likely  
1275 because there were more dark-preferring neurons than at higher contrasts (see the bottom row of Fig. 4 and Fig.  
1276 6K, P). Error bars in D, E denote s.e.m. across neurons.


1277  
1278



1279  
1280


1281 **Figure 6 Contrast-dependent SC processing of dark and bright stimuli.** (A-E) Scatterplots of response latencies  
1282 to dark versus bright stimuli at each contrast level. The proportion of neurons responding faster to darks  
1283 (above the unity line) increased with increasing contrast level. The proportion of neurons responding faster to  
1284 darks was also substantially larger at the lowest contrast level (E). (F-K) Scatterplots of peak visual responses to  
1285 darks versus brights at each contrast level. With increasing contrast, the proportion of neurons responding  
1286 stronger to brights (above the unity line) also increased. (L-P) Correlations between differences in visual  
1287 response latencies and differences in peak visual responses for brights and darks. The differences were  
1288 obtained in the same way as in Fig. 4: a positive response latency difference indicates a faster response to dark  
1289 stimuli, and a positive peak response means higher sensitivity to bright stimuli. The gray quadrants indicate the  
1290 regions where preferences in terms of sensitivity and latency coincided: neurons in the upper left quadrant  
1291 responded both stronger and faster for brights, and neurons in the lower right quadrant responded both  
1292 stronger and faster for darks. There were weak negative correlations between response latency and sensitivity  
1293 differences across contrasts, consistent with a known relationship between response latency and sensitivity.  
1294 Critically, however, earlier responses to darks were not dictated by sensitivity preferences; if this was the case,  
1295 most neurons would have occupied the shaded gray quadrants. Instead, at the highest contrast levels (e.g. L,  
1296 M, N), the majority of neurons, which responded earlier for darks, responded stronger for brights (occupying  
1297 the upper right quadrants), suggesting that dark stimuli were potent in expediting visual bursts despite the  
1298 bursts being non-preferred by the neurons.

1299  
1300



1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314

**Figure 7 Magnification of response latency and sensitivity differences between dark and bright stimuli in the upper visual field.** We repeated the same analyses of Fig. 4A, but now separating the neurons according to whether they represented upper or lower visual field locations. The top panel shows that the faster detection of dark stimuli by SC neurons that we saw in Fig. 4 was amplified for upper visual field neurons. The lower panel again shows that the effects of the top panel were dissociated from visual response sensitivity (all figure conventions are similar to Fig. 4). Interestingly, the lower panel shows that the absolute value of visual response sensitivity difference (between brights and darks) was also higher for upper visual field neurons than for lower visual field neurons (compare the dynamic ranges of the two curves). Therefore, both visual response latency and visual response sensitivity effects, in terms of luminance contrast polarity, were magnified in the upper visual field.




1315  
1316

1317 **Figure 8 Faster detection of darks than brights in the SC, independently of central fixation spot appearance.**  
1318 **(A, B)** We repeated the same experiment as in Figs. 1-7, but this time with a white fixation spot rather than a  
1319 black fixation spot (Methods). Also, here, we removed the fixation spot at stimulus onset, to allow monkeys to  
1320 generate visually-guided saccades to the appearing stimuli. In each panel, we averaged each monkey's stimulus-  
1321 aligned firing rates (after normalizing each neuron's firing rate curve to its peak visual response at high contrast;  
1322 Methods). The numbers of neurons are shown in each panel, and error bars denote s.e.m. across neurons. As  
1323 can be seen, visual responses still occurred earlier for dark stimuli than for bright stimuli in this task, and in both  
1324 monkeys. Note that at around 100 ms from stimulus onset, there was an elevation of firing rates, which was the  
1325 beginning of the saccade bursts for the triggered eye movements. Nonetheless, in monkey M, the elevation  
1326 looked to be sharper than in monkey A, an observation that we discuss in more detail in Figs. 9, 11 below with  
1327 respect to saccadic reaction times. **(C-E)** We replicated the analyses of Fig. 4 for the three contrasts that we  
1328 tested in this task variant. The same conclusions were reached. Most neurons detected dark stimuli earlier than  
1329 bright stimuli in their visual response latencies (top row), and this effect was dissociated from individual neuron  
1330 sensitivity to either darks or brights (bottom row). All other conventions are similar to Fig. 4.

1331

1332




1333 **Figure 9 Preference for bright stimuli in later sustained intervals of visual neural SC responses. (A)** Two example  
1334 neurons from monkey A showing how later sustained responses (after the initial visual bursts) were elevated for  
1335 bright more than dark stimuli. Responses for 100% contrast stimuli are shown, but similar observations were also  
1336 made across contrasts (see **C**). **(B)** Two example neurons from monkey M showing similar observations. Note  
1337 how neuron 8 had a weaker sustained response for bright stimuli, but it still exhibited a secondary burst (of small  
1338 amplitude) for bright stimuli (also see Fig. 11D for this monkey's population response summary in the sustained  
1339 interval). The pink rectangles denote our interval of choice when analyzing sustained visual responses. **(C)** In such  
1340 interval, we obtained millisecond-by-millisecond fits of contrast sensitivity curves for bright and dark stimuli. The  
1341 dynamic range parameter of equation 1,  $R$ , showed a clear and significant elevation for bright stimuli relative to  
1342 dark stimuli across the population (top panel). This was also the case in each monkey individually. The middle  
1343 and lower panels show that the thresholds (middle panel) and slopes (lower panel) of contrast sensitivity curves  
1344 were getting progressively worse in the sustained interval (relative to initial visual bursts), as expected, but with  
1345 little differences between dark and bright stimuli. Error bars in all panels denote s.e.m. (across trials in **A, B**, and  
1346 across neurons in **C**).  
1347

1348

1349

1350



1351

1352

1353 **Figure 10 Faster detection of darks than brights even in SC neurons inhibited by stimulus onset. (A, B)** Firing rates of two example neurons from a single linear electrode array penetration into the SC of monkey M. Both neurons had visual responses to dark stimuli in the upper left quadrant, with their RF's (obtained by presenting small black spots at different locations; insets) being well localized in space, consistent with the SC topographic representation (Robinson, 1972; Chen et al., 2019). **(C)** A third neuron recorded simultaneously with the neurons in **A, B**. This neuron was inhibited by stimulus onset (also see the RF map in the inset). Nonetheless, the inhibition was still stimulus-dependent: there was earlier inhibition for dark than bright stimuli, consistent with our earlier results (e.g. Figs. 1, 4-8). Error bars denote s.e.m. across trials in **A-C**. **(D)** Replication of the analysis of Fig. 4A (top) for all neurons that were inhibited by stimulus onset. Now, we estimated visual response latency by checking when the neural activity was significantly decreased from baseline. Most neurons were still modulated earlier by dark than bright stimuli, consistent with our earlier results for visual bursts (e.g. Figs. 1, 4-8). All other conventions are similar to Fig. 4A.

1354

1355

1356

1357

1358

1359

1360

1361


1362

1363

1364

1365

1366



1367  
1368

1369 **Figure 11 Relationship between saccadic reaction times and SC visual response properties for darks and brights**  
1370 **in the SC. (A)** Saccadic reaction times as a function of stimulus contrast (x-axis) and luminance polarity (different  
1371 lines) in monkey A from the saccade version of our task. Error bars denote s.e.m. across trials. Reaction times  
1372 were significantly shorter for dark than bright stimuli at all contrasts, suggesting a potential role for the earlier  
1373 SC visual responses for dark stimuli in triggering earlier saccades for such stimuli. **(B)** This monkey's neural  
1374 responses during the fixation task showed clearly earlier visual bursts for dark stimuli, with equal visual burst  
1375 strengths for darks and brights (consistent with Fig. 2). The figure was obtained similarly to Fig. 8A, B. Note also  
1376 that the population sustained response (from the peak of the visual burst onward) was larger for brights than  
1377 darks (consistent with Fig. 9). Error bars denote s.e.m. across neurons. **(C)** Monkey M showed the opposite  
1378 reaction time effects from Monkey A. **(D)** In this monkey's neurons, the secondary burst for bright stimuli in the  
1379 fixation variant of the task was particularly prominent (compare to the monkey A neural responses). This suggests  
1380 that in this monkey, this secondary preference for bright stimuli might have dominated the monkey's reaction  
1381 times in the saccade task. **(E)** We tested a third monkey behaviorally, and we replicated the monkey A results.  
1382 Therefore, in 2 out of the 3 monkeys, saccadic reaction times were earlier for dark than bright stimuli, consistent  
1383 with the neural results of Figs. 1, 4-8, 10.

1384