

# <sup>1</sup> Fat2 polarizes the WAVE complex in *trans* to align <sup>2</sup> cell protrusions for collective migration

<sup>3</sup> Audrey M. Williams<sup>1</sup>, Seth Donoughe<sup>1</sup>, Edwin Munro<sup>1,2,3</sup>, Sally Horne-Badovinac<sup>1,2,\*</sup>

<sup>4</sup>

## <sup>5</sup> Abstract

<sup>6</sup> For a group of cells to migrate together, each cell must couple the polarity of its migratory machinery with that  
<sup>7</sup> of the other cells in the cohort. Although collective cell migrations are common in animal development, little is  
<sup>8</sup> known about how protrusions are coherently polarized among groups of migrating epithelial cells. We address  
<sup>9</sup> this problem in the collective migration of the follicular epithelial cells in *Drosophila melanogaster*. In this epithel-  
<sup>10</sup> lium, the cadherin Fat2 localizes to the trailing edge of each cell and promotes the formation of lamellipodia at  
<sup>11</sup> the leading edge of the cell behind. We show that Fat2 performs this function by acting in *trans* to restrict WAVE  
<sup>12</sup> complex activity to one long-lived region along each cell's leading edge. Without Fat2, the WAVE complex distri-  
<sup>13</sup> bution expands around the cell perimeter and fluctuates over time, resulting in reduced, unpolarized protrusive  
<sup>14</sup> activity. We further show that Fat2's influence is very local, with sub-micron-scale puncta of Fat2 concentrat-  
<sup>15</sup> ing the WAVE complex in corresponding puncta just across the leading-trailing cell-cell interface. These findings  
<sup>16</sup> demonstrate that a *trans* interaction between Fat2 and the WAVE complex creates stable regions of protrusive  
<sup>17</sup> activity in each cell and aligns the cells' protrusions across the epithelium for directionally persistent collective  
<sup>18</sup> migration.

## <sup>19</sup> Introduction

<sup>20</sup> Collective cell migration is essential for a variety of morphogenetic processes in animals<sup>1-4</sup>. As with individual  
<sup>21</sup> cell migrations, adherent collective migrations are driven by the concerted action of cell protrusions, contractile  
<sup>22</sup> actomyosin networks, and adhesions to a substrate<sup>2,5,6</sup>. To move forward, individual cells polarize these structures  
<sup>23</sup> along a migratory axis, and to move persistently in one direction, they need to maintain that polarity stably over  
<sup>24</sup> time<sup>7</sup>. Collective cell migrations introduce a new challenge: to move together, the group of migrating cells must  
<sup>25</sup> be polarized in the same direction<sup>7</sup>. Otherwise, they would exert forces in different directions and move less  
<sup>26</sup> efficiently, separate, or fail to migrate altogether.

<sup>27</sup> The epithelial follicle cells of the *Drosophila melanogaster* ovary are a powerful experimental system in which to  
<sup>28</sup> investigate how local interactions among migrating cells establish and maintain group polarity. Follicle cells are

---

<sup>\*</sup>Corresponding author. E-mail: [shorne@uchicago.edu](mailto:shorne@uchicago.edu)

<sup>1</sup>Department of Molecular Genetics and Cell Biology, University of Chicago, Chicago, IL, USA

<sup>2</sup>Committee on Development, Regeneration, and Stem Cell Biology, University of Chicago, Chicago, IL, USA

<sup>3</sup>Institute for Biophysical Dynamics, University of Chicago, Chicago, IL, USA

29 arranged in a continuous, topologically closed monolayer epithelium that forms the outer cell layer of the ellip-  
30 soidal egg chamber—the organ-like structure that gives rise to the egg<sup>8</sup> (Fig. 1). The apical surfaces of follicle cells  
31 adhere to a central germ cell cluster, and their basal surfaces face outward and adhere to a surrounding basement  
32 membrane extracellular matrix. The follicle cells migrate along this stationary basement membrane, resulting in  
33 rotation of the entire cell cluster<sup>9</sup>. As the cells migrate, they secrete additional basement membrane proteins<sup>9</sup>.  
34 The coordination of migration with secretion causes the cells to produce a basement membrane structure that  
35 channels tissue growth along one axis<sup>9–12</sup>. Follicle cell migration lasts for roughly two days, and the migration  
36 direction—and resulting direction of egg chamber rotation—is stable throughout<sup>13,14</sup>. The edgeless geometry  
37 of the epithelium means cells are not partitioned into “leader” and “follower” roles, and there is no open space,  
38 chemical gradient, or other external guidance cue to dictate the migration direction. Instead, this feat of stable  
39 cell polarization and directed migration is accomplished through local interactions between the migrating cells  
40 themselves<sup>14,15</sup>.

41 Follicle cell migration is driven, in part, by lamellipodial protrusions that extend from the leading edge of each  
42 cell<sup>10,16</sup>. Lamellipodia are built by the WASP family verprolin homolog regulatory complex (WAVE complex)<sup>17,18</sup>,  
43 which is a protein assembly composed of five subunits: SCAR/WAVE, Abi, Sra1/Cyfip, Hem/Nap1, and HSPC300<sup>19</sup>.  
44 The WAVE complex adds branches to actin filaments by activating the Actin-related proteins-2/3 complex (Arp2/3)  
45 and elongates existing filaments, building the branched actin network that pushes the leading edge forward<sup>20–22</sup>.

46 The follicle cells align their lamellipodial protrusions across the tissue, a form of planar polarity<sup>10,16</sup>. The atypical  
47 cadherin Fat2 is required both for this planar polarity and for collective migration to occur<sup>23–25</sup>. Fat2 is planar  
48 polarized to the trailing edge of each cell<sup>24</sup>, where it promotes the formation of protrusions at the leading edge  
49 of the cell immediately behind<sup>15</sup>. Interestingly, in addition to migration depending on polarized Fat2 activity,  
50 Fat2’s planar polarity also depends on epithelial migration<sup>15</sup>. It is not known how Fat2 regulates lamellipodia or  
51 cell polarity, or how these processes influence one another. We hypothesized that Fat2 acts as a coupler between  
52 tissue planar polarity and cell protrusion by polarizing WAVE complex activity to the leading edge of each cell.  
53 To test this, we used genetic mosaic analysis and quantitative imaging of fixed and live tissues to dissect Fat2’s  
54 contributions to protrusivity and protrusion polarity at cell and tissue scales.

55 We show that Fat2 signals in *trans*, entraining WAVE complex activity to one long-lived region along each cell’s  
56 leading edge. Without Fat2, the WAVE complex accumulates transiently at different regions around the cell  
57 perimeter, and cell protrusivity is reduced and unpolarized. The interaction between Fat2 and the WAVE complex  
58 is non-cell-autonomous but very local, with sub-micron-scale puncta of Fat2 along the trailing edge concentrating  
59 the WAVE complex just across the cell-cell interface, at the tips of filopodia embedded within the lamellipodium.  
60 These findings demonstrate how an intercellular interaction between Fat2 and the WAVE complex promotes cell  
61 protrusivity, stabilizes regions of protrusive activity along the cell perimeter, and aligns protrusions across the  
62 epithelium by coupling leading and trailing edges. Fat2-WAVE complex interaction thereby stabilizes the planar  
63 polarity of protrusions for directionally persistent collective migration.

## 64 Results

### 65 Fat2 increases and polarizes protrusions at the basal surface of the follicular epithelium

66 Recent work has shown that Fat2 regulates migration of the follicular epithelium by polarizing F-actin-rich pro-  
67 trusions; specifically, Fat2 at the trailing edge of each cell causes protrusions to form at the leading edge of the  
68 cell behind it, and without Fat2, protrusions are reduced or lost<sup>15,26</sup>. Beyond this qualitative description, it is not  
69 known how Fat2 modulates cell protrusion. Because protrusion is a dynamic process of spatially coordinated cell  
70 extension and retraction, we used live imaging and fluorescent labeling of the plasma membrane to obtain a more  
71 detailed, time-resolved view of protrusions and their distribution around cells. We acquired timelapse movies of  
72 the basal surface of control and *fat2*<sup>N103-2</sup> epithelia (a null allele, hereafter referred to as *fat2*) and saw that there  
73 was substantial protrusive activity in *fat2* epithelia, but the protrusions lacked the clear polarized distribution of  
74 control epithelia (Fig. 2A; Movie 1).

75 To quantify the extent and distribution of protrusions in these epithelia, we developed methods to segment mem-  
76 brane extensions and measure their lengths and orientations (Fig. S1). We briefly summarize our quantification  
77 approach here and include a more detailed explanation in the Methods and Materials. First, we measured the av-  
78 erage lengths of membrane extensions from all cell-cell interfaces (Fig. 2B). The distribution of measured lengths  
79 was unimodal, with no natural division between protrusive and non-protrusive interfaces. Therefore, to estab-  
80 lish an empirically-grounded cutoff between these categories, we recorded timelapse movies of control epithelia  
81 treated with the Arp2/3 inhibitor CK-666, which are non-migratory and almost entirely non-protrusive<sup>16</sup>. We  
82 used measurements from CK-666-treated epithelia to set a cutoff for the minimum length of a protrusion: any  
83 edges with membrane extensions longer than the 98th percentile of those in CK-666-treated epithelia were con-  
84 sidered *protrusive* for subsequent analysis.

85 Using this quantification approach, we first asked how the amount of protrusions was affected by the loss of Fat2.  
86 We found that the protrusivity of *fat2* epithelia was lower than that of control on average, but highly variable,  
87 with overlap between the protrusivity distributions of both untreated and CK-666-treated epithelia (Fig. 2B,C;  
88 S2A,B; Movie 1). As a complementary method, we also measured protrusivity via F-actin labeling in fixed and live  
89 tissues, using *abi*-RNAi-expressing epithelia as a nearly non-protrusive benchmark. The results largely paralleled  
90 those seen with membrane labeling (Fig. S3A-D; Movie 2); however, F-actin labeling showed a larger disparity  
91 in protrusivity between *fat2* and control epithelia (Figs. 2C; S3C). Images of follicle cell protrusions visualized  
92 by F-actin staining are dominated by fluorescence from filopodia, so the appearance of lower protrusivity of *fat2*  
93 epithelia as measured with an F-actin label may indicate that filopodia are disproportionately reduced by loss of  
94 Fat2. Altogether, these data show that *fat2* epithelia are less protrusive than control, but do retain some protrusive  
95 activity.

96 These results raised an important question—if some *fat2* epithelia have levels of membrane protrusivity compa-  
97 rable to that of control epithelia, then why do all *fat2* epithelia fail to migrate<sup>13,15,24</sup>? We hypothesized that the  
98 mispolarization of protrusions across the tissue contributes to *fat2* migration failure. In control epithelia, the  
99 majority of protrusions were polarized in the direction of migration, orthogonally to the egg chamber’s anterior-  
100 posterior axis (Fig. 2A,D; S2C; Movie 3). In contrast, in *fat2* epithelia, protrusions were fairly uniformly dis-

101 tributed in all directions or biased in two opposite directions (Fig. 2A,D; S2C; Movie 3). Where an axial bias was  
102 present, the axis was inconsistent between egg chambers. We also confirmed this finding using F-actin labeling of  
103 protrusions. To compare the planar polarity of F-actin protrusions between control and *fat2* epithelia, we mea-  
104 sured F-actin enrichment at cell-cell interfaces as a function of the angle of the interface with respect to the egg  
105 chamber's anterior-posterior axis. We again saw that protrusions were planar-polarized in control epithelia and  
106 unpolarized in *fat2* epithelia (Fig. S3A,E,F). These data show that Fat2 is required to polarize protrusions in a  
107 common direction across the epithelium.

108 Because Fat2 regulates both follicle cell migration and planar polarity, and migration and planar polarity are inter-  
109 dependent<sup>15,16,23,24</sup>, the unpolarized protrusions of *fat2* epithelia could be a cause or a consequence of inability  
110 of *fat2* epithelia to migrate. To distinguish between these possibilities, we exploited the fact that small groups of  
111 *fat2* cells can be carried along by neighboring non-mutant, migratory cells<sup>24</sup>, allowing us to evaluate polarity of  
112 protrusions from *fat2* cells in a migratory context. We generated *fat2* mosaic tissues that had sufficiently small  
113 fractions of mutant cells that the tissue as a whole still migrated, and found that *fat2* cells in these tissues were  
114 often protrusive, but their protrusions were not polarized in the direction of migration (Fig. S4; Movie 4). This  
115 demonstrates that Fat2 does not simply polarize protrusions indirectly by maintaining tissue-wide migration.  
116 Rather, Fat2 is required at the scale of groups of cells to polarize those cells' protrusions in alignment with the  
117 direction of collective migration.

## 118 **Fat2 increases and polarizes the WAVE complex at the basal surface of the follicular epithelium**

119 Follicle cell protrusions are built by the WAVE complex<sup>16</sup>, so we hypothesized that Fat2 polarizes protrusions  
120 by polarizing the WAVE complex's distribution. To visualize the WAVE complex in living tissue and at normal  
121 expression levels, we used CRISPR/Cas9 to endogenously tag the WAVE complex subunit Sra1 with eGFP (here-  
122 after: Sra1-GFP). We confirmed that Sra1-GFP flies are viable and fertile when the tagged allele is homozygous,  
123 Sra1-GFP localizes to follicle cell leading edges like other WAVE complex labels<sup>16,26</sup>, its localization depends on  
124 WAVE complex subunit Abi, and F-actin protrusions appear normal (Figs. 3A,B; S5A-C). Migration was slower  
125 when Sra1-GFP was present in two copies (Fig. S5D; Movie 5), so we performed all subsequent experiments with  
126 one copy of Sra1-GFP.

127 With an endogenous WAVE complex label in hand, we investigated how Fat2 affects WAVE complex localization.  
128 Previous work has shown that WAVE complex levels are reduced at the basal surface of follicle cells lacking Fat2<sup>26</sup>.  
129 Consistent with this result, we found that Sra1-GFP levels were lower along cell-cell interfaces at the basal surface  
130 of *fat2* epithelia than of control epithelia (Fig. S6A-C). Planar polarity of Sra1-GFP across the epithelium was  
131 also lost in the absence of Fat2 (Fig. S6D,E). Fat2 acts non-cell-autonomously to cause protrusions to form at  
132 the leading edge of the cell just behind<sup>15</sup> (Fig. 3C), so we next tested the hypothesis that Fat2 localizes the WAVE  
133 complex to the leading edge in the same non-cell-autonomous pattern. We did this using *fat2* mosaic epithelia, in  
134 which we could measure Sra1-GFP levels at leading-trailing interfaces shared by control and *fat2* cells. We found  
135 that Sra1-GFP levels were normally-enriched along the leading edges of *fat2* cells if control cells were present  
136 immediately ahead, showing that Sra1 can still localize to the leading edge of cells lacking Fat2. Conversely, Sra1-  
137 GFP levels were reduced along the leading edge of control cells if *fat2* cells were immediately ahead (Fig. 3D,E). We

138 also observed a corresponding non-autonomous pattern of membrane protrusion polarity in timelapse movies  
139 of *fat2* mosaic epithelia (Fig. S4; Movie 4). We conclude that Fat2 acts non-cell-autonomously to localize the  
140 WAVE complex to leading edges, resulting in tissue-wide planar polarization of protrusive activity, and thereby  
141 in collective cell migration.

142 We next asked if loss of the WAVE complex from the leading edge in the absence of Fat2 meant that the protein  
143 was redistributed to other cell surfaces. To test this, we compared the level of Sra1-GFP at the medial basal surfaces  
144 of control and *fat2* cells in mosaic epithelia (see diagram in Fig. 3F). We found that Sra1-GFP levels were slightly  
145 increased in *fat2* cells compared to control cells in the same tissue (Fig. 3D,F). There was not a statistically signifi-  
146 cant increase in Sra1-GFP levels at medial basal cell surfaces of epithelia composed entirely of control or *fat2* cells,  
147 although the trend was the same (Fig. S6C). These data demonstrate that Fat2 concentrates the WAVE complex  
148 at the leading edge, and that without Fat2, the WAVE complex becomes distributed more broadly across the basal  
149 surface.

150 **Fat2 stabilizes a region of WAVE complex enrichment and protrusivity in *trans***

151 In individually migrating cells, the excitable dynamics of the WAVE complex and its regulators enable it to form  
152 transient zones of enrichment along the cell perimeter even in the absence of a directional signal<sup>7,27,28</sup>. Although  
153 the planar-polarized distribution of the WAVE complex across the epithelium was lost in *fat2* mutant tissue,  
154 we wondered (1) whether the WAVE complex could still form regions of enrichment in individual cells and (2)  
155 whether these WAVE complex-enriched regions were active and responsible for templating unpolarized protru-  
156 sions. To evaluate the WAVE complex distribution along the edges of individual cells, we generated entirely *fat2*  
157 mutant epithelia in which patches of cells expressed Sra1-GFP. At cell-cell interfaces along Sra1-GFP expression  
158 boundaries, we found that the boundary cells often had cortical regions devoid of Sra-GFP (Fig 4A). This ob-  
159 servation shows that the WAVE complex is not uniformly localized around the cortex and can form regions of  
160 enrichment without Fat2. We also saw that Sra1-GFP enrichment coincided with the presence of F-actin protru-  
161 sions (Fig. 4A), indicating that the WAVE complex in these regions is active. To confirm that the WAVE complex  
162 builds the protrusions in *fat2* epithelia, we co-imaged Sra1-GFP and a membrane label, and found that Sra1-GFP  
163 was enriched at the tips of membrane protrusions (Fig. S7A; Movie 6). These data indicate that the WAVE com-  
164 plex can still accumulate and build protrusions in the absence of Fat2, tissue-wide planar polarity, and collective  
165 cell migration.

166 A striking feature of migrating follicle cells is the stable polarization of their protrusive leading edges. It is not  
167 known whether Fat2 contributes to the stabilization of protrusive regions in addition to positioning them. If  
168 so, the positions of protrusive regions of *fat2* epithelia should fluctuate more than those of control epithelia, in  
169 addition to being less well-polarized at the tissue level. To see if this is the case, we acquired timelapse movies  
170 of Sra1-GFP and monitored its distribution along cell perimeters over time. In control epithelia, Sra1-GFP was  
171 strongly enriched along leading-trailing interfaces relative to side interfaces over the 20-minute timelapse. Side  
172 interfaces were mostly devoid of Sra1-GFP, except for infrequent Sra1-GFP accumulations that persisted for sev-  
173 eral minutes (Fig. 4B-D; Movies 7; 8). In contrast, in *fat2* epithelia, the regions of greatest Sra1-GFP enrichment  
174 along the cell perimeter changed substantially over the 20-minute timelapse and multiple Sra1-GFP-enriched re-

175 gions were often present simultaneously in individual *fat2* cells. Sra1-GFP accumulated in these regions, typically  
176 spreading outward along the membrane as it did so, and then dissipated. These events had a duration that was  
177 comparable to the transient accumulations of Sra1-GFP at side interfaces in control cells (Fig. 4B-D; Movies 7;  
178 8). Because all cell-cell interfaces in *fat2* epithelia and side interfaces in control epithelia lack Fat2, this suggests  
179 a several-minutes timescale over which regions of WAVE complex enrichment can persist without stabilization  
180 by Fat2. Live imaging of Sra1-GFP in *fat2* mosaic epithelia yielded similar information—Sra1-GFP enrichment  
181 fluctuated more at interfaces between *fat2* cells than interfaces between control cells despite both being in a mi-  
182 gratory tissue (Fig. S7B,C; Movie 9). To see if Fat2’s role stabilizing the WAVE complex distribution translates to  
183 a role stabilizing protrusive regions, we compared the stability of regions of membrane protrusion between cells  
184 in control or *fat2* epithelia. In control cells, we found that protrusions extended and retracted from one leading  
185 edge region, and changes to the direction of cell protrusion were rare (Fig. 4E; Movie 10). In contrast, the edges  
186 of cells in *fat2* epithelia undergoing protrusion often shifted substantially over the 20 minutes of the timelapse.  
187 Together, these observations show that, in addition to polarizing protrusive activity to the leading edge, Fat2 sta-  
188 bilizes the distribution of WAVE complex activity for repeated cycles of protrusion from one long-lived protrusive  
189 region (Fig 4F).

## 190 **Fat2 and the WAVE complex colocalize across leading-trailing cell-cell interfaces**

191 Finally, we explored how Fat2 recruits the WAVE complex across the cell-cell interface. To constrain the set of  
192 possible mechanisms, we assessed the spatial scale of their interaction. Fat2 has a punctate distribution along  
193 each cell’s trailing edge<sup>15,24</sup>, so we asked whether Fat2 recruits the WAVE complex locally to these sites, or recruits  
194 it more broadly to the entire interface. We evaluated the colocalization between Fat2 and the WAVE complex  
195 along leading-trailing interfaces, visualizing Fat2 with an endogenous 3xeGFP tag (Fat2-3xGFP) and the WAVE  
196 complex with mCherry-tagged Abi under control of the ubiquitin promoter (Abi-mCherry). Like Fat2-3xGFP,  
197 Abi-mCherry formed puncta, and Abi-mCherry and Fat2-3xGFP puncta colocalized significantly more than  
198 Abi-mCherry and uniformly-distributed E-cadherin-GFP (Spearman’s  $r = 0.71 \pm 0.04$  vs.  $0.49 \pm 0.07$ ; Figs 5A-E;  
199 S8A,B). In timelapse movies, Fat2-3xGFP and Abi-mCherry puncta moved together through cycles of protrusion  
200 extension and retraction (Figs. 5B; Movie 11). Short-lived Abi-mCherry accumulations formed infrequently at  
201 cell sides away from Fat2, similar to the Sra1-GFP side accumulations we described earlier (Figs. 4D; S9; Movies  
202 7; 12). Together, these findings suggest that Fat2 recruits the WAVE complex locally, at the scale of individual  
203 puncta, with the WAVE complex occasionally “escaping” Fat2-dependent concentration at the leading edge.

204 If Fat2 puncta locally recruit the WAVE complex, changing the distribution of Fat2 puncta should cause cor-  
205 responding changes to the distribution of the WAVE complex. To test this, we examined follicle cells express-  
206 ing an endogenous Fat2 truncation that lacks the intracellular domain (Fat2<sup>ΔICD</sup>-3xGFP), which distributes  
207 more broadly around the cell perimeter than wild-type Fat2<sup>15,29</sup>, but remains punctate. The distribution of  
208 Abi-mCherry expanded around the cell perimeter in the Fat2<sup>ΔICD</sup>-3xGFP background (Fig. 5A) as was previ-  
209 ously reported for protrusions<sup>15</sup>. Despite their altered distributions, Abi-mCherry puncta colocalized just as well  
210 with Fat2<sup>ΔICD</sup>-3xGFP puncta as with Fat2-3xGFP puncta (Spearman’s  $r = 0.71 \pm 0.04$  vs.  $0.71 \pm 0.05$ ; Figs. 5E;  
211 S8B). From these data we conclude that Fat2 controls the distribution of the WAVE complex by concentrating  
212 the WAVE complex in adjacent puncta. These findings also demonstrate that the Fat2 intracellular domain is

213 dispensable for Fat2-WAVE complex interaction in collectively migrating follicle cells.

214 Ena-dependent filopodia are embedded within and grow from the lamellipodia<sup>16</sup>. The WAVE complex inter-  
215 acts with Ena and is required for the filopodia to form<sup>16,30</sup>, so we asked whether the distribution of Fat2-WAVE  
216 complex puncta is related to the distribution of filopodia. Labeling filopodia tips with a GFP-tagged Ena trans-  
217 gene (GFP-Ena) and comparing the localization of Abi-mCherry and F-actin to either GFP-Ena or to Fat2-  
218 3xGFP, we found that the sites of highest Fat2-3xGFP and Abi-mCherry enrichment coincided with filopodia  
219 tips (Figs. 5C,D,F; S8C,D). Fluorescence intensity profiles along filopodia lengths showed that Fat2-3xGFP and  
220 Abi-mCherry were enriched just ahead of the F-actin-rich region (Fig. 5F,G). Fat2-3xGFP was shifted slightly  
221 forward from Abi-mCherry, consistent with the separation of Fat2-3xGFP and Abi-mCherry fluorophores by a  
222 cell-cell interface (Figs. 5F,G; S8D,E). This analysis demonstrates a stereotyped organization in which Fat2 and  
223 the WAVE complex are concentrated along with Ena near the tips of the filopodia, with Fat2 at the trailing edge  
224 across the cell-cell interface from the leading edge components.

225 We considered two explanations for the close spatial relationship between Fat2 puncta, WAVE complex puncta,  
226 and filopodia. Fat2 could recruit the WAVE complex locally to puncta, and WAVE complex puncta shape the  
227 distribution of filopodia. Alternatively, Fat2 could recruit the WAVE complex to the leading edge, but their colo-  
228 calization in puncta be a secondary effect of the filopodia, perhaps caused by the known interaction between Ena  
229 and Abi<sup>30</sup> or by deformation of the leading-trailing interface. To rule out the second possibility, we measured  
230 colocalization between Fat2-3xGFP and Abi-mCherry in *ena*-RNAi-expressing epithelia, in which filopodia are  
231 strongly depleted<sup>16</sup> (Fig. S8F). Despite the loss of filopodia, both Fat2-3xGFP and Abi-mCherry remained punc-  
232 tate, and their colocalization was only slightly reduced (Spearman's  $r = 0.71 \pm 0.04$  vs.  $0.65 \pm 0.03$ , Fig. S8A,B).  
233 We therefore rule out Ena or the filopodia themselves as required mediators of the spatial relationship between  
234 Fat2 and the WAVE complex, and infer that Fat2-WAVE complex colocalization is indicative of Fat2 recruitment  
235 of the WAVE complex locally to these sites.

236 Altogether, we propose that Fat2 acts locally, at the scale of individual Fat2 puncta, to concentrate the WAVE  
237 complex in adjacent puncta across the cell-cell interface. Because Fat2 puncta are distributed along the trailing  
238 edge, this has the broader effect of stabilizing a region of WAVE complex enrichment at the leading edge.

## 239 Discussion

240 This work demonstrates that a *trans* interaction between the atypical cadherin Fat2 and the WAVE complex can  
241 stabilize WAVE complex polarity for directed cell migration. Fat2, localized to the trailing edge of each cell, re-  
242 cruits the WAVE complex to the leading edge of the cell behind, just across their shared interface. By concentrating  
243 WAVE complex activity in a restricted region, Fat2 strongly biases lamellipodia and filopodia to form at these lead-  
244 ing edge sites, stably polarizing overall cell protrusive activity to one cell side. Because the Fat2-WAVE complex  
245 signaling system is deployed at each leading-trailing interface in a planar-polarized manner, it both polarizes pro-  
246 trusions within individual cells and aligns these individual cell polarities across the epithelium. This allows the  
247 cells to exert force in a common direction and achieve a highly coordinated collective cell migration.

248 While the molecular players differ, local coupling of leading and trailing edges through asymmetric interactions

249 across their shared interface is a recurring motif in studies of epithelial collective cell migrations. In an epithelial cell culture model of collective migration, asymmetric pulling forces across cell-cell interfaces polarize Rac1  
250 activity and cell protrusion<sup>31</sup>. In another model, one cell's lamellipodium is stabilized by confinement under  
251 the trailing edge of the cell ahead, reinforcing interface asymmetry<sup>32</sup>. In an endothelial collective cell migration  
252 model, asymmetric membrane “fingers” containing VE-cadherin extend from the trailing edge and are engulfed  
253 by the leading edge of the cell behind, whose movement they help guide<sup>33</sup>. These types of leading-trailing edge  
254 coupling systems could operate together with longer-range cues to reinforce the planar polarity of cells' migratory  
255 structures. In migrations with a closed topology and no extrinsic directional cues, such as that of the follicle cells,  
256 local polarity coupling may be especially critical for collective migration.

258 Our development of new computational tools to segment and quantify membrane protrusion dynamics in a  
259 collectively-migrating epithelium has led to new insights into how Fat2 regulates protrusions. We found that  
260 without Fat2, protrusivity was reduced, and the distribution of remaining protrusions expanded around the en-  
261 tire cell periphery. Therefore, Fat2 not only promotes protrusion at the leading edge, but also restricts protrusion  
262 to that edge. Analysis of *fat2* mosaic epithelia revealed that Fat2 acts locally to enforce this restriction—even in  
263 the context of a globally planar-polarized, migratory epithelium, cells lacking input from Fat2 in the cell ahead  
264 were unable to polarize their protrusions in the direction of migration. Although we focused on the length and  
265 orientation of protrusions in this study, our approach could be extended to other protrusion traits such as shape,  
266 lifetime, or elongation rate, and/or to other tissues, making it broadly applicable to studies of epithelial collective  
267 migrations.

268 Excitable WAVE complex dynamics underlie lamellipodial protrusion<sup>27,34,35</sup>, and these dynamics were especially  
269 apparent in our data in contexts where Fat2 was absent from the cell-cell interface—either at interfaces between  
270 cells in *fat2* epithelia, or at protrusive side interfaces we observed with low frequency in control cells. In both  
271 contexts, the WAVE complex accumulated at an edge region, spread laterally along the membrane, and then dissi-  
272 pated. This corresponded with the initiation, growth, and collapse of a protrusion. Where Fat2 was present, the  
273 WAVE complex distribution along the cell perimeter stayed more constant and WAVE complex levels fluctuated  
274 in place, but did not appear to spread laterally. In the absence of Fat2, the WAVE complex also had an expanded  
275 distribution around lateral cell edges and across the basal surface. These findings suggests that Fat2 acts by se-  
276 questering the WAVE complex, reducing its ability to accumulate elsewhere and thus restricting protrusions to a  
277 single leading edge. Moreover, the reduced protrusivity of *fat2* epithelia may be due to their broader and more  
278 diffuse WAVE complex distribution, which could reduce the frequency with which the WAVE complex crosses  
279 a threshold of enrichment necessary to initiate a protrusion. The sporadic side-facing protrusions in wild-type  
280 cells show that the Fat2 signaling system does not exert perfect control over the distribution of WAVE complex  
281 activity, but this level of control is sufficient to stably polarize cell protrusive activity for days-long, highly directed  
282 migration.

283 Given that follicle cells can form WAVE complex-dependent protrusions without Fat2, how does Fat2 shape the  
284 WAVE complex's distribution and dynamics in wild-type cells? WAVE complex activity is often entrained by di-  
285 rectional cues from the environment<sup>34,36–38</sup>, and we hypothesize that Fat2 is acting as a similar directional cue  
286 in follicle cells. The WAVE complex is activated by recruitment to the plasma membrane<sup>19,39,40</sup>. Positive reg-

287 regulators of WAVE complex accumulation include active Rac, phosphatidylinositol (3,4,5)-triphosphate (PIP<sub>3</sub>),  
288 membrane-localized proteins that directly bind the WAVE complex, and the WAVE complex itself<sup>17,35,39,41–46</sup>.  
289 We propose that Fat2 promotes WAVE complex accumulation within a stable region by acting through one or  
290 more of these positive regulators, thereby controlling the site where the WAVE complex excitation threshold is  
291 crossed and a protrusion is formed. Under this model, in the absence of Fat2, this site selection instead becomes  
292 more stochastic and therefore long-lasting protrusive regions cannot form.

293 Fat2 acts at the trailing edge of each cell to recruit the WAVE complex in *trans*, so there must be one or more  
294 transmembrane proteins at the leading edge of each cell that bridge this interaction. Previous work has shown that  
295 the receptor tyrosine phosphatase Lar is part of this bridge—Fat2 recruits Lar to each follicle cell’s leading edge<sup>15</sup>,  
296 and in Lar’s absence both WAVE complex levels and cell protrusions are reduced<sup>15,26</sup> (Fig. S8G). However, the  
297 WAVE complex that persists at the leading edges of *lar* cells still colocalizes with Fat2 (Fig. S8A,B). Therefore,  
298 there must be at least one other transmembrane protein that works alongside Lar to mediate the Fat2-WAVE  
299 complex interaction. Identifying the missing leading edge protein(s) will be important to fully understand how  
300 Fat2 shapes WAVE complex activity.

301 Fat2 is localized in puncta along each cell’s trailing edge<sup>15,24</sup>, and we show here that these puncta correspond 1:1  
302 with regions of high WAVE complex enrichment just across the leading-trailing cell-cell interface. Fat2’s punctate  
303 distribution and its levels along cell-cell interfaces are unaffected by loss of the WAVE complex<sup>15</sup>, indicating that  
304 Fat2 puncta shape the distribution of the WAVE complex and protrusions, not the reverse. We further show that  
305 the puncta sit at the tips of filopodia that form within the lamellipodial actin network. Filopodia are a prominent  
306 feature of the long-lived protrusive regions that form in wild-type epithelia, but appear to be disproportionately  
307 reduced in the short-lived, fluctuating protrusive regions that form in *fat2* epithelia. We therefore propose that  
308 by concentrating the WAVE complex and/or stabilizing its distribution, Fat2 also facilitates filopodia formation.  
309 It should be noted, however, that the filopodia are dispensable for collective follicle cell migration<sup>16</sup>, so the reason  
310 these structures form remains to be determined.

311 Why, and how, is Fat2 localized in puncta? Cadherins commonly form puncta, though the causes and functions  
312 of this organization vary<sup>47–49</sup>. For example, Flamingo (or mammalian Celsr1), an atypical cadherin and central  
313 component of the core planar cell polarity pathway, is stabilized by clustering, and this clustering is important  
314 for its planar polarization<sup>50–52</sup>. In future work, it will be important to determine how Fat2 assembles in puncta,  
315 and whether this local clustering is important for its polarization to trailing edges or its effect on the organization  
316 of leading edges. More broadly, it will be critical to determine how Fat2 achieves its trailing edge localization, a  
317 necessary step in the polarization of the tissue.

### 318 **Author Contributions**

319 A.M.W. and S.H.-B. conceived of the study. A.M.W. designed experiments with critical input from all authors.  
320 A.M.W. generated new reagents and performed experiments. A.M.W. and S.D. wrote data analysis software and  
321 analyzed data. A.M.W., S.D., E.M., and S.H.-B. interpreted data. A.M.W. prepared figures. A.M.W. and S.H.-B.  
322 wrote the manuscript with editorial input from all authors.

### 323 **Acknowledgements**

324 We thank members of the Horne-Badovinac and Munro labs, Allison Zajac, Sherzod Tokamov, Ellie Heckscher,  
325 Michael Glotzer, and Carmen Williams for feedback throughout the study and comments on the manuscript.  
326 This work was supported by NIH R01 GM126047 to S.H.B., NIH R01 HD88831 to E.M., NIH T32 HD055164  
327 to A.M.W., and postdoctoral fellowships from the Chicago Fellows Program and Jane Coffin Childs Memorial  
328 Fund for Medical Research to S.D..

329 **Figures**







Figure 1: **Introduction to egg chamber rotation.** **A**, Diagram of a stage 6 egg chamber in cross-section. Anterior is left, posterior right. **B**, Three-dimensional diagram of an egg chamber with the anterior half shown. Arrows indicate the migration of follicle cells along the basement membrane and the resulting rotation of the egg chamber around its anterior-posterior axis. **C**, Diagram of three follicle cells. Their apical surfaces adhere to the germ cells and their basal surfaces adhere to the basement membrane. The dashed line represents the basal imaging plane used throughout this study except where indicated.




**Figure 2: Fat2 increases and polarizes follicle cell protrusivity.** **A**, Timelapse frames of control, *fat2*, and CK-666-treated epithelia labeled with a membrane dye. Middle row shows segmented edges. Protrusive edges, defined as edges with average membrane extension lengths longer than the 98th percentile of those of CK-666-treated epithelia, are shown in red. Non-protrusive edges are white. Bottom row shows arrows indicating the orientation of each protrusion overlaid on labeled cell membrane. Arrows originate at protrusion bases and have lengths proportional to protrusion lengths. See related Movies 1, 3. **B**, Histogram showing the distribution of average membrane extension lengths. The 98th percentile length threshold for CK-666-treated epithelia is indicated. **C**, Plot showing the ratio of protrusive to total edges. The protrusivity of *fat2* epithelia is variable, with a distribution overlapping with control and CK-666-treated epithelia. Welch's ANOVA ( $W(2,8.5)=17$ ,  $p=0.0011$ ) with Dunnet's T3 multiple comparisons test; n.s.  $p=0.07$ ,  $*p=0.03$ ,  $**p=0.006$ . **D**, Polar histograms of the distribution of protrusion orientations in control and *fat2* epithelia. Anterior is left, posterior is right, and in control epithelia images were flipped as needed so that migration is always oriented downward. Control protrusions point predominantly in the direction of migration, whereas *fat2* protrusions are less polarized. Associated with Figs. S1, S2, S3, S4; Movies 1, 3, 2, 4.



**Figure 3: Fat2 in each cell concentrates the WAVE complex at the leading edge of the cell behind.** **A**, Diagram showing Fat2 localization at the trailing edge and WAVE complex at the leading edge of the basal surface of follicle cells. The WAVE complex subunits referenced in this paper listed. **B**, Images of an Sra1-GFP mosaic epithelium with phalloidin-stained F-actin, showing Sra1-GFP enrichment at leading edges (filled arrows) and not trailing edges (open arrows). **C**, Images of a *fat2* mosaic epithelium with phalloidin-stained F-actin. Filled arrows indicate leading edges of *fat2* cells behind control cells, where protrusions are present. Open arrows indicate leading edges of control cells behind *fat2* cells, where protrusions are reduced. **D**, Images of a *fat2* mosaic epithelium expressing Sra1-GFP. Filled arrows indicate leading edges of *fat2* cells behind control cells. Open arrows indicate leading edges of control cells behind *fat2* cells. **E,F**, Quantification of Sra1-GFP mean fluorescence intensity in *fat2* mosaic epithelia along leading-trailing interfaces (E) or medial basal surfaces (F). Diagrams to the left of plots show the measured regions with respect to control (cyan) and *fat2* (gray) cells. The genotype(s) of cells in each measured category are shown below the x-axis. **E**, Sra1-GFP is reduced at the leading edge of cells of any genotype behind *fat2* cells. Bars indicate mean  $\pm$  SD. One-way ANOVA ( $F(3,57)=10.40$ ,  $p<0.0001$ ) with post-hoc Tukey's test; n.s. (left to right)  $p=0.96$ ,  $0.90$ ,  $**p<0.01$ . **F**, Sra1-GFP is slightly increased at the medial basal surface of *fat2* cells. Lines connect measurements from the same egg chamber. Paired t-test;  $**p<0.01$ . Associated with Figs. S5, S6; Movie 5.



**Figure 4: Fat2 stabilizes a region of WAVE complex enrichment and protrusivity.** **A**, Images of phalloidin-stained F-actin and mosaically-expressed Sra1-GFP in an entirely *fat2* mutant epithelium. Filled and open arrows indicate genotype boundary interfaces with and without Sra1-GFP enrichment, respectively. Sra1-GFP enrichment is heterogeneous, and interfaces with Sra1-GFP enrichment have more F-actin protrusions. **B**, Timelapse frames of Sra1-GFP in control and *fat2* epithelia. Top row shows Sra1-GFP with arrows indicating regions of Sra1-GFP enrichment; bottom row shows Sra1-GFP and outlines of cell perimeters used to make kymographs. Laser intensity and brightness display settings differ between genotypes. See related Movies 7, 8. **C**, Diagram of cell perimeter unrolling for kymograph generation. Red represents planar-polarized Sra1 as distributed before and after unrolling. **D**, Kymographs of Sra1-GFP fluorescence intensity along cell perimeter outlines exemplified in (C). The *y*-axis length of regions of high Sra1-GFP enrichment reports their stability over time. Control cells have Sra1-GFP regions along leading-trailing interfaces that are stable over 20 minutes. In *fat2* cells, Sra1-GFP-enriched regions are less stable. The arrow indicates a transient accumulation of Sra1-GFP at a control cell side. These occur occasionally, and their stability is similar to Sra1-GFP regions in *fat2* cells. **E**, Timelapse frames of control and *fat2* epithelia with a membrane dye. Top row shows the interfaces and protrusions of one cell and its neighbors. Segmented membrane extensions originating from the center cell (red) are overlaid in the bottom row. Arrows indicate sites of membrane protrusion. The position of protrusions in the *fat2* cell changes more than in the control cell. See related Movie 10. **F**, Diagram showing the proposed role of Fat2 stabilizing a region of WAVE complex enrichment and protrusivity. Without Fat2, WAVE complex-enriched, protrusive regions are reduced and more transient. Associated with Fig. S7; Movies 6, 7, 8, 10, 9.



**Figure 5: Fat2 colocalizes with the WAVE complex across leading-trailing cell-cell interfaces.** **A**, Images of cells expressing Abi-mCherry and endogenous full-length Fat2-3xGFP or endogenous Fat2-3xGFP lacking the intracellular domain (Fat2<sup>ΔICD</sup>), used to assess colocalization. **B**, Timelapse frames showing the leading-trailing interfaces of two cells expressing Fat2-3xGFP and Abi-mCherry, showing their colocalization over time. See related Movie 11. **C**, Image showing the leading-trailing interface region used in (D); it is also an example of a region used in (E). **D**, Line scan showing the fluorescence intensity of Fat2-3xGFP, Abi-mCherry, and F-actin (phalloidin) along the leading-trailing interfaces of the two cells in (C), showing their corresponding peaks of enrichment. **E**, Plot of Spearman's correlation coefficients of Fat2-3xGFP or Fat2<sup>ΔICD</sup>-3xGFP and Abi-mCherry showing no significant difference in colocalization. Bars indicate mean  $\pm$  SD. One-way ANOVA ( $F(5,81)=44.86$ ,  $p=0.0164$  with Fig. S8B) with post-hoc Tukey's test; n.s.  $p>0.99$ . **F**, Image showing the distribution of Fat2-3xGFP, Abi-mCherry, and F-actin (phalloidin) at the leading-trailing interface and along the boxed filopodium. **G**, Plot showing fluorescence intensity of traces of F-actin, Abi-mCherry, and Fat2-3xGFP showing their relative sites of enrichment along the length of filopodia. Lines and shaded regions indicate mean  $\pm$  SD.  $n=74$  protrusions (used for SD), 18 cells, 1 cell/egg chamber. **H**, Diagram of proposed organization of Fat2, the WAVE complex, and F-actin along the leading-trailing interface based on the present data and previously published work<sup>15,16,24</sup>. Fat2 puncta at the trailing edge colocalize with WAVE complex puncta at the leading edge, ahead of filopodia embedded within the lamellipodium. Associated with Figs. S8, S9; Movies 11, 12.

## 330 Methods and Materials

### 331 Materials, data, and code availability

332 New plasmids and *Drosophila* lines reported in this paper are available upon request to the corresponding au-  
333 thor. The code necessary to reproduce core aspects of data analysis, along with example datasets, are available  
334 at [https://github.com/a9w/Fat2\\_polarizes\\_WAVE](https://github.com/a9w/Fat2_polarizes_WAVE). Additional data and code are available upon request to the  
335 corresponding author.

### 336 *Drosophila* sources, care, and genetics

337 The sources and references of all stocks used in this study are listed in Supplemental Table 1 and the genotypes of  
338 *Drosophila* used in each experiment and associated figure panels are listed in Supplemental Table 2. *Drosophila*  
339 were raised at 25°C and fed cornmeal molasses agar food. Females 0-3 days post-eclosion were aged on yeast with  
340 males prior to dissection. In most cases, they were aged for 2-3 days at 25°C. Temperatures and yeasting times used  
341 for each experiment are reported in Supplemental Table 3. In all RNAi experiments, *traffic jam*>Gal4 (*tj*>Gal4)<sup>53</sup>  
342 was used to drive RNAi expression in follicle cells and not in germ cells. *Sra1*-GFP and *fat2* mosaic epithelia were  
343 generated using the Flp/FRT method<sup>54,55</sup>, using FRT82B and FRT80B recombination sites, respectively. In both  
344 cases *tj*>Gal4 was used to drive expression of UAS>Flp recombinase.

### 345 Generation of *Sra1*-GFP

346 Endogenous *Sra1* was tagged C-terminally with enhanced GFP (GFP) following the general approaches described  
347 by Gratz *et al.* (2013) and Gratz *et al.* (2014)<sup>56,57</sup>. The guide RNA target sequence 5'-GCTTAAATGCATCC  
348 CTTTCCGGG-3' was chosen with flyCRISPR Target Finder<sup>57</sup>. The underlined sequence was cloned into the  
349 pU6-BbsI-chiRNA plasmid, and the bold sequence is the adjacent PAM motif. For homologous recombination,  
350 homology arms approximately 2 kb long flanking the insertion target site were amplified from genomic DNA  
351 from the y1 M{nos-Cas9.P}ZH-2A w\* (nanos-Cas9)<sup>58</sup> background. GFP was amplified from the pTWG plas-  
352 mid. A linker with sequence encoding the amino acids 'GSGGSGGS' was added to the N-terminal side of GFP.  
353 Homology arms, linker, and GFP were inserted into donor plasmid pDsRed-attP, which contains 3xP3-DsRed  
354 flanked by loxP sites for insertion screening and subsequent removal. The linker-GFP insertion was made imme-  
355 diately before the *Sra1* stop codon. Guide and homologous recombination plasmids were injected by Genetivi-  
356 sion Inc. into the nanos-Cas9 background. F1 males were screened for 3xP3-DsRed and then 3xP3-DsRed was  
357 excised by crossing to Cre-expressing flies (MKRS hsFLP/TM6b Cre).

### 358 Egg chamber dissection

359 Ovaries were dissected into live imaging media (Schneider's *Drosophila* medium with 15% fetal bovine serum and  
360 200 µg/mL insulin) in a spot plate using 1 set of Dumont #55 forceps and 1 set of Dumont #5 forceps. Ovarioles  
361 were removed from the ovary and from ovariole muscle sheathes with forceps. For live imaging, egg chambers  
362 older than the egg chamber to be imaged were removed from the ovariole strands by cutting through the stalk  
363 with a 27-gauge hypodermic needle. For fixed imaging, egg chambers older than stage 9 were removed prior  
364 to fixation. Removal of older egg chambers allows more compression of the imaged egg chamber between the

365 slide and coverslip so that the basal surface of a field of cells can be imaged in a single plane. For a more detailed  
366 description and movies of dissection, see Cetera *et al.* (2016)<sup>59</sup>.

### 367 **Live imaging sample preparation**

368 Following dissection, ovarioles were transferred to a fresh well of live imaging media. For membrane staining,  
369 CellMask Orange or Deep Red plasma membrane stain (Thermo Fisher Scientific, Waltham, MA, 1:500) was  
370 added and ovarioles incubated for 15 minutes, followed by a wash in live imaging media to remove excess stain  
371 before mounting. Ovarioles were then transferred to a glass slide with 20  $\mu$ L of live imaging media. For CK-666  
372 treatment, following plasma membrane staining, ovarioles were transferred to live imaging media with 750  $\mu$ M  
373 CK-666 (Millipore Sigma, St. Louis, MO) and then mounted in the same media. Glass beads with diameter 51  
374  $\mu$ m were added to support the 22x22 mm #1.5 coverslip and limit egg chamber compression. Coverslip edges  
375 were sealed with melted petroleum jelly to prevent evaporation while imaging. Samples were checked for damage  
376 using the membrane stain or other fluorescent markers as indicators, and excluded if damage was observed. Slides  
377 were used for no more than 1 hour.

### 378 **Immunostaining and F-actin staining**

379 Following dissection, ovarioles were fixed in 4% EM-grade formaldehyde in PBT (phosphate buffered saline +  
380 0.1% Triton X-100) and then washed 3x5 minutes in PBT at room temperature. Egg chambers were incubated  
381 with primary antibodies in PBT overnight at 4°C (anti-Scar, 1:200) or for 2 hours at room temperature (anti-  
382 Discs Large, 1:20) while rocking. Ovarioles were then washed 3x5 minutes in PBT and incubated in secondary  
383 antibody diluted 1:200 in PBT for two hours at room temperature while rocking. F-actin staining was performed  
384 using either TRITC phalloidin (Millipore Sigma, 1:250) or Alexa Fluor 647 phalloidin (Thermo Fisher Scientific,  
385 1:50). If TRITC phalloidin was the only stain or antibody used, it was added directly to the fixation media for  
386 15 minutes of staining concurrent with fixation. Otherwise, TRITC phalloidin was added for 15-30 minutes at  
387 room temperature as the final staining step. Alexa Fluor 647 phalloidin staining was performed for two hours at  
388 room temperature while the sample was rocking, concurrent with secondary antibody staining where applicable.  
389 Ovarioles were then washed 3x5 minutes in PBT and mounted in 40  $\mu$ L SlowFade Diamond antifade on a slide  
390 using a 22x50 mm #1.5 coverslip, sealed with nail polish, and stored at 4°C until imaged.

### 391 **Microscopy**

#### 392 **Laser scanning confocal microscopy**

393 Laser scanning confocal microscopy was used for all fixed imaging and for live imaging of membrane-dyed egg  
394 chambers. Imaging was performed with a Zeiss LSM 800 upright laser scanning confocal with a 40x/1.3 NA EC  
395 Plan-NEOFLUAR oil immersion objective or a 63x/1.4 NA Plan-APOCHROMAT oil immersion objective,  
396 diode lasers (405, 488, 561, and 640 nm), and GaAsP detectors. The system was controlled with Zen 2.3 Blue  
397 acquisition software (Zeiss). Imaging was performed at room temperature. All images show the basal surface of  
398 stage 6-7 egg chambers except for Fig. S5A, bottom row, which shows follicle cells in cross-section. Cross-section  
399 images were used for egg chamber staging throughout. Laser scanning confocal microscopy was used to acquire

400 the data in Figs. 2; 3B-F; 4A,E; 5A,C-G; S1; S2; S3A-C,E,F; S4; S5; S6; S7A; S8; Movies 1; 3; 4; 5; 6; 10.

401 **TIRF microscopy**

402 Near-TIRF microscopy was used to visualize Fat2-GFP, Sra1-GFP, Abi-mCherry, and F-Tractin-tdTomato<sup>60</sup> dy-  
403 namics at the basal surface. Near-TIRF imaging was performed with a Nikon ECLIPSE-Ti inverted microscope  
404 with Ti-ND6-PFS Perfect Focus Unit, solid-state 50 mW 481 and 561 nm Sapphire lasers (Coherent technology),  
405 motorized TIRF illuminator, laser merge module (Spectral Applied Research), Nikon CFI 100x Apo 1.45 NA  
406 oil immersion TIRF objective with 1.5x intermediate magnification, and Andor iXon3 897 electron-multiplying  
407 charged-coupled device (EM-CCD) camera. Image acquisition was controlled using MetaMorph software. For  
408 two color imaging, frames were collected for each color consecutively with the TIRF illumination angle adjusted  
409 in between. Imaging was performed at room temperature. For display, movies were corrected for bleaching using  
410 the histogram matching method in Fiji (ImageJ)<sup>61,62</sup>. TIRF microscopy was used to capture the data in Figs.  
411 4B,D; 5B; S3D; S7B,C; S9; Movies 2; 7; 8; 9; 11; 12.

412 **Cell and protrusion segmentation from timelapses of cell membrane**

413 Protrusions from timelapse datasets of the follicle cell basal surface stained with CellMask Orange (see Live imag-  
414 ing sample preparation) were segmented with the Python scikit-image and scipy libraries (Fig. S1)<sup>63,64</sup>. First, each  
415 cell was segmented and tracked, with manual corrections to cell-cell interface placements made using napari<sup>65</sup>.  
416 Next, a watershed-based approach was used to segment the regions of high fluorescence intensity at the interface  
417 of each pair of neighboring cells. This segmented shape encompasses the cell-cell interface and any associated  
418 protrusions from either neighboring cell. Last, to assign protrusions to the cell from which they originated, the  
419 segmented region was divided in two by the shortest path between its bounding vertices that lay entirely within  
420 the region. This approximates the position of the interface between the cells, and in subsequent steps we will  
421 call this line “the interface”. Each of the two resulting protrusion shapes was assigned as originating from the cell  
422 on the opposite side of the interface, because protrusions extend from one cell and overlap the other. Using this  
423 approach, all of the protrusive structures that emerge from one cell, and that overlap a single neighboring cell, are  
424 grouped together as a single segmented region for subsequent analysis.

425 **Measurement of membrane protrusivity, protrusion length, and protrusion orientation**

426 After cell edges and associated protrusions were segmented, they were categorized as either protrusive or non-  
427 protrusive and their lengths and orientations using Python scikit-fmm, scikit-image, and scipy libraries. We use  
428 the term “membrane extensions” to refer to the cell edge shapes before the protrusive ones have been identified.

429 To measure the length of a membrane extension, we used two different metrics, each of which provides a single  
430 length value per cell edge. In one, we calculated the “average length” of a membrane extension as the membrane  
431 extension’s area divided by the length of the interface it extended across. As an alternate length measurement,  
432 we calculated its “longest length.” To do so, we first found its “tip”, defined as the pixel within the segmented  
433 region farthest from any point along the interface. We then found its “base”, the pixel along the interface that  
434 was closest to the tip. We defined membrane extension longest length as the length of the shortest path between

435 base and tip that lay entirely within the membrane extension. To categorize membrane extensions as *protrusive*  
436 or *non-protrusive* throughout the study, we used the “average length” metric. We measured the average length dis-  
437 tribution in CK-666-treated epithelia, which are nearly non-protrusive and so provided a measure of the width  
438 of the cell-cell interface alone. For all conditions, we categorized a membrane extension as protrusive if its average  
439 length was greater than the 98th percentile of length of CK-666-treated epithelia. We then defined the protru-  
440 sivity of an entire epithelium as the ratio of protrusive to total cell edges in the field of view. We also report two  
441 alternate measurements of the protrusivity of an epithelium. In one, We calculate epithelial protrusivity as above,  
442 but substitute the longest length as our length measurement (Fig. S2A). In a second, cutoff-independent epithe-  
443 lial protrusivity measurement, we report the epithelium-mean average membrane extension length (Fig. S2B).  
444 Swarm plots of each of these analyses were generated using GraphPad Prism 9 (GraphPad, San Diego, CA), as  
445 were all other swarm plots.

446 For analysis of protrusion orientation, we included only the membrane extensions categorized as protrusive ac-  
447 cording to the “average length” metric. We defined a protrusion’s orientation as the orientation of the vector from  
448 its base to its tip. Polar histograms, generated in Python with matplotlib<sup>66</sup>, show the distribution of protrusion  
449 orientations. In these plots, bar area is proportional to the number of protrusions in the corresponding bin.

#### 450 **Quantification of F-actin and Sra1-GFP cell-cell interface and non-interface basal surface fluo- 451 rescence**

452 Cells and cell-cell interfaces were segmented as described above. Cells and interfaces in contact with the tissue  
453 border or image border were excluded from analysis. For interface fluorescence intensity, interfaces were dilated  
454 by 5 pixels, and mean fluorescence intensity calculated from within this region. Non-interface basal surface fluo-  
455 rescence intensity was calculated as the mean of the remaining (non-interface) tissue surface. For F-actin cell-cell  
456 interface enrichment measurements, the overall brightness of the phalloidin staining varied between epithelia  
457 independent of genotype. To control for this variation we subtracted the mean intensity of the epithelium’s non-  
458 interface basal surface from its mean interface intensity measurement. This value, the degree of F-actin interface  
459 enrichment, was used as a proxy for F-actin protrusivity.

#### 460 **Quantification of F-actin and Sra1-GFP planar polarity**

461 As a simple planar polarity measurement, we quantified mean F-actin (phalloidin) or Sra1-GFP fluorescence in-  
462 tensity along each cell-cell interface as a function of the interface’s orientation. To do this, cells and cell-cell in-  
463 terfaces were segmented as described above. For interface angle measurements, the angular distance between the  
464 line defined by the interface-bounding vertices and the anterior-posterior (horizontal) axis was calculated. For  
465 interface fluorescence intensity measurements, interface regions were identified as segmented interfaces dilated  
466 by 5 pixels. Vertices, dilated by 10 pixels, were excluded from interface regions. Mean fluorescence intensity was  
467 calculated within each interface region, and background (the mean non-interface basal surface fluorescence in-  
468 tensity of all cells in the image) was subtracted. To calculate the leading-trailing interface enrichment of F-actin  
469 or Sra1-GFP for each egg chamber, interface fluorescence intensities were averaged for all interfaces with angles  
470 between 0° and 10° (leading-trailing interfaces), and between 80° and 90° (side interfaces). The leading-trailing

471 interface enrichment is the ratio of these numbers.

## 472 **Autonomy analysis in mosaic epithelia**

473 Egg chambers were stained with Alexa Fluor 647 phalloidin to mark protrusions, which indicate migration direc-  
474 tion, and to determine whether egg chambers were planar polarized. We analyzed only S6-7 egg chambers with  
475 mixtures of control and *fat2* cells that had global stress fiber alignment orthogonal to the anterior-posterior axis,  
476 indicating global planar polarity. Since migration is required to maintain planar polarity<sup>16</sup>, this also indicates  
477 that the epithelium was migratory. Looking at phalloidin and genotype markers only, we drew 10 pixel-wide seg-  
478 mented lines along leading-trailing interface boundaries of different genotype combinations in Fiji. Lines were  
479 drawn along all visible, in-focus *fat2*-control and control-*fat2* boundaries and a similar number of control-control  
480 and *fat2*-*fat2* boundaries. A boundary category was excluded if there were fewer than 3 usable interfaces to mea-  
481 sure. Mean Sra1-GFP fluorescence intensities were calculated for each interface type in each egg chamber. For  
482 a diagram of this method, see Barlan *et al.* (2017)<sup>15</sup>. To quantify non-interface basal surface fluorescence, we  
483 drew polygonal regions of the basal surface of control and *fat2* cells, excluding cells immediately behind those of  
484 a different genotype. Egg chambers were excluded if there were fewer than 3 usable cells of either genotype. Mean  
485 Sra1-GFP fluorescence intensities were calculated within these polygonal regions for all control cells and all *fat2*  
486 cells in an egg chamber.

## 487 **Quantification of migration rate**

488 Egg chambers were dissected, dyed with CellMask Orange, and mounted for live imaging as described above.  
489 Several ovarioles were mounted on each slide, with each ovariole terminating in a S6-7 egg chamber. Timelapse  
490 imaging was performed for 30 minutes with frames acquired every 30 seconds. Multi-point acquisition was used  
491 to obtain movies of up to 5 egg chambers simultaneously. To generate a kymograph, a line was drawn along the  
492 axis of migration at the center of the anterior-posterior egg chamber axis in Fiji. In these kymographs, cell-cell  
493 interfaces are visible as lines, and their slope gives a measurement of cell migration rate. Egg chamber migration  
494 rates were calculated from the average of 4 cell interface slopes from each kymograph. Egg chambers that clearly  
495 slowed down over the course of the timelapse, visible as curvature in the interface lines in the kymographs, were  
496 excluded. For an illustration of this method, see Barlan *et al.*<sup>15</sup>.

## 497 **Cell perimeter kymograph generation and interpretation**

498 To visualize the distribution of Sra1-GFP along cell-cell interfaces over time, we generated kymographs of cell  
499 perimeters from timelapses of Sra1-GFP-expressing epithelia obtained using near-TIRF microscopy. Perimeters  
500 were drawn manually in Fiji in each frame with the pencil tool, and then these perimeters were used to generate  
501 kymographs in Python. Perimeters were thinned to 1 pixel and then perimeter pixels were sequenced with Python  
502 scikit-image and scipy libraries. Kymographs were generated with matplotlib. Kymograph rows were constructed  
503 by linearizing the perimeters from each frame, starting with the pixel directly above the cell centroid (the center of  
504 the trailing edge in control cells) and continuing counter-clockwise. Each row shows the fluorescence intensity of  
505 the perimeter pixels in sequence. Cell perimeter lengths varied between frames, so kymograph row lengths varied  
506 and were aligned to their center position.

507 At the spatial and temporal resolution of the timelapses and corresponding kymographs, we cannot evaluate dif-  
508 ferences in the dynamics the puncta-scale WAVE complex accumulations highlighted in Fig. 5. Instead, we focus  
509 on the “region”-scale distribution of Sra1-GFP, and the stability of that distribution over time. The regions we  
510 refer to here are approximately the length of a cell-cell interface, with variation. Because the kymographs are gen-  
511 erated from epithelia in which all cells express Sra1-GFP, we need additional information to identify the cell to  
512 which a region of Sra1-GFP enrichment belongs. We infer that Sra1-GFP is predominantly at leading edges in  
513 polarized, migratory epithelia based on the Sra1-GFP distribution in epithelia with mosaic Sra1-GFP expression  
514 (Fig. 3B). Based on consistent correlation between Sra1-GFP enrichment and the presence of protrusions (Fig.  
515 3B, 4A, S7A), and its known role building lamellipodia as part of the WAVE complex<sup>17,18,41</sup>, we also infer that  
516 regions of Sra1-GFP enrichment belong to the cell that is protruding outward regardless of genotype. Our in-  
517 terpretations of Sra1-GFP enrichment patterns in movies and corresponding kymographs are made with these  
518 assumptions.

### 519 Colocalization of proteins along the leading-trailing interface

520 Data used for colocalization analysis were collected with 63x/1.4 NA Plan-APOCHROMAT oil immersion ob-  
521 jective to minimize chromatic aberration. Linescans were generated in Fiji by manually drawing a 10 pixel-wide  
522 segmented line along rows of leading-trailing interfaces at the follicle cell basal surface. At least 20 leading-trailing  
523 interfaces were included per egg chamber. For the  $\text{Fat2}^{\Delta\text{ICD}}$  condition, in which the distribution of Fat2 expands  
524 beyond leading-trailing interfaces, we measured colocalization either along randomly oriented interfaces (Fig. 5E)  
525 or leading-trailing interfaces (Fig. S8B) and obtained very similar results. Fluorescence intensities along the lines-  
526 cans were obtained with the PlotProfile function, which averages pixel intensities along the width of the line and  
527 reports a list of averaged values along the line’s length. Spearman’s correlation coefficients were calculated for  
528 each egg chamber in Python with the `scipy.stats` module. Failure to exactly follow leading-trailing interfaces and  
529 cusps in the segmented lines will artificially inflate the measured correlation, so we used correlation between E-  
530 cadherin-GFP<sup>67</sup> and Abi-mCherry as a negative control that is also subject to this inflation. Abi-mCherry and  
531 E-cadherin-GFP are slightly displaced from each other (anticorrelated) along the length of protrusions (the width  
532 of the linescans), but averaging across the line width collapses this displacement, resulting in measured intensity  
533 signals that are roughly uncorrelated. Spearman’s correlation coefficients  $\pm$  standard deviation are reported in  
534 the text. Linescans of leading-trailing interfaces were plotted using the fluorescence intensities from along the  
535 leading-trailing interfaces of two cells. Intensities from each fluorophore were rescaled between 0 and 1 and plot-  
536 ted with matplotlib in Python.

### 537 Protrusion profile generation

538 Viewing only the F-actin channel in Fiji, we drew 1 pixel-wide lines down the length of F-actin bundles at the  
539 leading edge. Fluorescence intensities along these lines were obtained for all fluorophores with the Fiji PlotProfile  
540 function. In Python, these traces were aligned to the pixel with highest  $\text{Fat2-3xGFP}$  or  $\text{Ena-GFP}$  intensity (Fig.  
541 5G, S8F). To calculate standard deviation, all traces were first rescaled individually so that their values ranged  
542 between 0 and 1. To plot “protrusion profiles,” the mean fluorescence was determined for each fluorophore at  
543 each pixel position, and then average values were rescaled between 0 and 1. Plots of protrusion profiles were

544 generated with matplotlib.

545 **Movie generation**

546 Migration motion was subtracted from several timelapse movies of migratory cells or epithelia for ease of visual-  
547 ization. Motion subtraction was performed using the Fiji MultiStackReg plugin “translation” transformation<sup>68</sup>  
548 (control condition in Movies 1; 6; 8) or by aligning to the centroid of a tracked cell in each frame using the scikit-  
549 image library (Movies 7; 10). Labels were added to movies in Fiji and then exported as uncompressed .avi files.  
550 These were encoded as 1080p30 .mp4 files with H.264 (x264) video encoder using HandBrake 1.4.

551 **Reproducibility and statistical analysis**

552 Visibly damaged egg chambers were excluded from all analyses. Each experiment was performed at least two in-  
553 dependent times, and results confirmed to be qualitatively consistent. Each experiment included egg chambers  
554 pooled from multiple flies. Experiments and analysis were not randomized or performed blinded. Sample sizes  
555 were not predetermined using a statistical method. The number of biological replicates (n), statistical tests per-  
556 formed, and their significance can be found in figures or figure legends. Based on visual inspection, all data on  
557 which statistical tests were performed followed an approximately normal distribution, so tests assuming normalcy  
558 were used. Alpha was set to 0.05 for all statistical tests. Paired statistical tests were used for comparisons of cells  
559 of different genetic conditions within mosaic epithelia, except if all epithelia did not have all genetic conditions  
560 represented, in which case an unpaired test was used so that all samples could still be included. All t-tests were  
561 two-tailed. One-sample t-tests were used when comparing a distribution of ratios to a null expectation of one. A  
562 one-way ANOVA was used when multiple pairs of conditions were compared, with the exception of plots in Figs.  
563 2C, S2A, and S3C, for which the variance did not appear consistent between conditions, so Welch’s ANOVA was  
564 used instead. For post-hoc comparison tests, all pairs of conditions present in the corresponding plot were com-  
565 pared using post-hoc Tukey’s multiple comparisons test with the following exception: the data plotted in Figs.  
566 5E and S8B were analyzed together, and all conditions were compared to Fat2-Abi and E-cadherin-Abi only, and  
567 in Fig. S6C only data from the same region (total, interface, or non-interface) was compared. For these, Šidák’s  
568 multiple comparisons tests were used. For Welch’s ANOVA, Dunnet’s T3 multiple comparisons tests were used.  
569 P-values reported for all post-hoc tests were adjusted for multiple comparisons. All statistical tests except for the  
570 calculation of Spearman’s correlation coefficients were performed in GraphPad Prism 9.

571 **References**

- 572 1. Friedl, P. & Gilmour, D. Collective Cell Migration in Morphogenesis, Regeneration and Cancer. *Nature Reviews Molecular Cell Biology* **10**, 445–457 (July 2009).
- 573 2. Scarpa, E. & Mayor, R. Collective Cell Migration in Development. *Journal of Cell Biology* **212**, 143–155 (Jan. 2016).
- 574 3. Norden, C. & Lecaudey, V. Collective Cell Migration: General Themes and New Paradigms. *Current Opinion in Genetics & Development. Developmental Mechanisms, Patterning and Evolution* **57**, 54–60 (Aug. 2019).
- 575 4. Perez-Vale, K. Z. & Peifer, M. Orchestrating Morphogenesis: Building the Body Plan by Cell Shape Changes and Movements. *Development* **147**, dev191049 (Sept. 2020).
- 576 5. Bodor, D. L., Pönisch, W., Endres, R. G. & Paluch, E. K. Of Cell Shapes and Motion: The Physical Basis of Animal Cell Migration. *Developmental Cell* **52**, 550–562 (Mar. 2020).
- 577 6. Buttenschön, A. & Edelstein-Keshet, L. Bridging from Single to Collective Cell Migration: A Review of Models and Links to Experiments. *PLOS Computational Biology* **16**, e1008411 (Dec. 2020).
- 578 7. Stock, J. & Pauli, A. Self-Organized Cell Migration across Scales – from Single Cell Movement to Tissue Formation. *Development* **148** (Apr. 2021).
- 579 8. Duhart, J. C., Parsons, T. T. & Raftery, L. A. The Repertoire of Epithelial Morphogenesis on Display: Progressive Elaboration of Drosophila Egg Structure. *Mechanisms of Development* **148**, 18–39 (Dec. 2017).
- 580 9. Haigo, S. L. & Bilder, D. Global Tissue Revolutions in a Morphogenetic Movement Controlling Elongation. *Science* **331**, 1071–1074 (Feb. 2011).
- 581 10. Gutzeit, H. O., Eberhardt, W. & Gratwohl, E. Laminin and Basement Membrane-Associated Microfilaments in Wild-Type and Mutant Drosophila Ovarian Follicles. *Journal of Cell Science* **100**, 781–788 (Dec. 1991).
- 582 11. Isabella, A. J. & Horne-Badovinac, S. Rab10-Mediated Secretion Synergizes with Tissue Movement to Build a Polarized Basement Membrane Architecture for Organ Morphogenesis. *Developmental Cell* **38**, 47–60 (July 2016).
- 583 12. Crest, J., Diz-Muñoz, A., Chen, D.-Y., Fletcher, D. A. & Bilder, D. Organ Sculpting by Patterned Extracellular Matrix Stiffness. *eLife* **6** (ed Spradling, A. C.) e24958 (June 2017).
- 584 13. Chen, D.-Y., Crest, J. & Bilder, D. A Cell Migration Tracking Tool Supports Coupling of Tissue Rotation to Elongation. *Cell Reports* **21**, 559–569 (Oct. 2017).
- 585 14. Stedden, C. G., Menegas, W., Zajac, A. L., Williams, A. M., Cheng, S., Özkan, E. & Horne-Badovinac, S. Planar-Polarized Semaphorin-5c and Plexin A Promote the Collective Migration of Epithelial Cells in Drosophila. *Current Biology* **29**, 908–920.e6 (Mar. 2019).
- 586 15. Barlan, K., Cetera, M. & Horne-Badovinac, S. Fat2 and Lar Define a Basally Localized Planar Signaling System Controlling Collective Cell Migration. *Developmental Cell* **40**, 467–477.e5 (Mar. 2017).
- 587 16. Cetera, M., Ramirez-San Juan, G. R., Oakes, P. W., Lewellyn, L., Fairchild, M. J., Tanentzapf, G., Gardel, M. L. & Horne-Badovinac, S. Epithelial Rotation Promotes the Global Alignment of Contractile Actin Bundles during Drosophila Egg Chamber Elongation. *Nature Communications* **5**, 5511 (Nov. 2014).
- 588 17. Miki, H., Suetsugu, S. & Takenawa, T. WAVE, a Novel WASP-family Protein Involved in Actin Reorganization Induced by Rac. *The EMBO Journal* **17**, 6932–6941 (Dec. 1998).
- 589 18. Miki, H., Yamaguchi, H., Suetsugu, S. & Takenawa, T. IRSp53 Is an Essential Intermediate between Rac and WAVE in the Regulation of Membrane Ruffling. *Nature* **408**, 732–735 (Dec. 2000).
- 590 19. Chen, Z., Borek, D., Padrick, S. B., Gomez, T. S., Metlagel, Z., Ismail, A. M., Umetani, J., Billadeau, D. D., Otwinowski, Z. & Rosen, M. K. Structure and Control of the Actin Regulatory WAVE Complex. *Nature* **468**, 533–538 (Nov. 2010).

616 20. Machesky, L. M., Mullins, R. D., Higgs, H. N., Kaiser, D. A., Blanchoin, L., May, R. C., Hall, M. E. & Pollard, T. D. Scar, a WASp-related Protein, Activates Nucleation of Actin Filaments by the Arp2/3 Complex. *Proceedings of the National Academy of Sciences* **96**, 3739–3744 (Mar. 1999).

617 21. Bieling, P., Hansen, S. D., Akin, O., Li, T.-D., Hayden, C. C., Fletcher, D. A. & R. Dyche, M. WH2 and Proline-Rich Domains of WASP-family Proteins Collaborate to Accelerate Actin Filament Elongation. *The EMBO Journal* **37**, 102–121 (Jan. 2018).

618 22. Mullins, R. D., Bieling, P. & Fletcher, D. A. From Solution to Surface to Filament: Actin Flux into Branched Networks. *Biophysical Reviews* **10**, 1537–1551 (Dec. 2018).

619 23. Viktorinová, I., König, T., Schlichting, K. & Dahmann, C. The Cadherin Fat2 Is Required for Planar Cell Polarity in the Drosophila Ovary. *Development* **136**, 4123–4132 (Dec. 2009).

620 24. Viktorinová, I. & Dahmann, C. Microtubule Polarity Predicts Direction of Egg Chamber Rotation in Drosophila. *Current Biology* **23**, 1472–1477 (Aug. 2013).

621 25. Horne-Badovinac, S. Fat-like Cadherins in Cell Migration—Leading from Both the Front and the Back. *Current Opinion in Cell Biology. Cell Dynamics* **48**, 26–32 (Oct. 2017).

622 26. Squarr, A. J., Brinkmann, K., Chen, B., Steinbacher, T., Ebnet, K., Rosen, M. K. & Bogdan, S. Fat2 Acts through the WAVE Regulatory Complex to Drive Collective Cell Migration during Tissue Rotation. *Journal of Cell Biology* **212**, 591–603 (Feb. 2016).

623 27. Weiner, O. D., Marganski, W. A., Wu, L. F., Altschuler, S. J. & Kirschner, M. W. An Actin-Based Wave Generator Organizes Cell Motility. *PLoS Biology* **5**, e221 (Aug. 2007).

624 28. Iglesias, P. A. & Devreotes, P. N. Biased Excitable Networks: How Cells Direct Motion in Response to Gradients. *Current Opinion in Cell Biology. Cell Regulation* **24**, 245–253 (Apr. 2012).

625 29. Aurich, F. & Dahmann, C. A Mutation in Fat2 Uncouples Tissue Elongation from Global Tissue Rotation. *Cell Reports* **14**, 2503–2510 (Mar. 2016).

626 30. Chen, X. J., Squarr, A. J., Stephan, R., Chen, B., Higgins, T. E., Barry, D. J., Martin, M. C., Rosen, M. K., Bogdan, S. & Way, M. Ena/VASP Proteins Cooperate with the WAVE Complex to Regulate the Actin Cytoskeleton. *Developmental Cell* **30**, 569–584 (Sept. 2014).

627 31. Das, T., Safferling, K., Rausch, S., Grabe, N., Boehm, H. & Spatz, J. P. A Molecular Mechanotransduction Pathway Regulates Collective Migration of Epithelial Cells. *Nature Cell Biology* **17**, 276–287 (Mar. 2015).

628 32. Jain, S., Cachoux, V. M. L., Narayana, G. H. N. S., de Beco, S., D’Alessandro, J., Cellerin, V., Chen, T., Heuzé, M. L., Marcq, P., Mège, R.-M., Kabla, A. J., Lim, C. T. & Ladoux, B. The Role of Single-Cell Mechanical Behaviour and Polarity in Driving Collective Cell Migration. *Nature Physics* **16**, 802–809 (July 2020).

629 33. Hayer, A., Shao, L., Chung, M., Joubert, L.-M., Yang, H. W., Tsai, F.-C., Bisaria, A., Betzig, E. & Meyer, T. Engulfed Cadherin Fingers Are Polarized Junctional Structures between Collectively Migrating Endothelial Cells. *Nature Cell Biology* **18**, 1311–1323 (Dec. 2016).

630 34. Xiong, Y., Huang, C.-H., Iglesias, P. A. & Devreotes, P. N. Cells Navigate with a Local-Excitation, Global-Inhibition-Biased Excitable Network. *Proceedings of the National Academy of Sciences* **107**, 17079–17086 (Oct. 2010).

631 35. Graziano, B. R. & Weiner, O. D. Self-Organization of Protrusions and Polarity during Eukaryotic Chemo-taxis. *Current Opinion in Cell Biology* **30**, 60–67 (Oct. 2014).

632 36. Millius, A., Dandekar, S. N., Houk, A. R. & Weiner, O. D. Neutrophils Establish Rapid and Robust WAVE Complex Polarity in an Actin-Dependent Fashion. *Current Biology: CB* **19**, 253–259 (Feb. 2009).

633 37. Huang, C.-H., Tang, M., Shi, C., Iglesias, P. A. & Devreotes, P. N. An Excitable Signal Integrator Couples to an Idling Cytoskeletal Oscillator to Drive Cell Migration. *Nature Cell Biology* **15**, 1307–1316 (Nov. 2013).

634 38. Hayashi, S., Inoue, Y., Kiyonari, H., Abe, T., Misaki, K., Moriguchi, H., Tanaka, Y. & Takeichi, M. Protocadherin-17 Mediates Collective Axon Extension by Recruiting Actin Regulator Complexes to Interaxonal Contacts. *Developmental Cell* **30**, 673–687 (Sept. 2014).

664 39. Oikawa, T., Yamaguchi, H., Itoh, T., Kato, M., Ijuin, T., Yamazaki, D., Suetsugu, S. & Takenawa, T. Pt-  
665 dIns(3,4,5)P 3 Binding Is Necessary for WAVE2-induced Formation of Lamellipodia. *Nature Cell Biology*  
666 **6**, 420–426 (May 2004).

667 40. Lebensohn, A. M. & Kirschner, M. W. Activation of the WAVE Complex by Coincident Signals Controls  
668 Actin Assembly. *Molecular Cell* **36**, 512–524 (Nov. 2009).

669 41. Steffen, A., Rottner, K., Ehinger, J., Innocenti, M., Scita, G., Jürgen, W. & Stradal, T. E. Sra-1 and Nap1  
670 Link Rac to Actin Assembly Driving Lamellipodia Formation. *The EMBO Journal* **23**, 749–759 (Feb.  
671 2004).

672 42. Sossey-Alaoui, K., Li, X., Ranalli, T. A. & Cowell, J. K. WAVE3-mediated Cell Migration and Lamellipodia  
673 Formation Are Regulated Downstream of Phosphatidylinositol 3-Kinase\*. *Journal of Biological Chemistry*  
674 **280**, 21748–21755 (June 2005).

675 43. Weiner, O. D., Rentel, M. C., Ott, A., Brown, G. E., Jedrychowski, M., Yaffe, M. B., Gygi, S. P., Cant-  
676 ley, L. C., Bourne, H. R. & Kirschner, M. W. Hem-1 Complexes Are Essential for Rac Activation, Actin  
677 Polymerization, and Myosin Regulation during Neutrophil Chemotaxis. *PLOS Biology* **4**, e38 (Jan. 2006).

678 44. Nakao, S., Platek, A., Hirano, S. & Takeichi, M. Contact-Dependent Promotion of Cell Migration by the  
679 OL-protocadherin–Nap1 Interaction. *Journal of Cell Biology* **182**, 395–410 (July 2008).

680 45. Namekata, K., Harada, C., Taya, C., Guo, X., Kimura, H., Parada, L. F. & Harada, T. Dock3 Induces  
681 Axonal Outgrowth by Stimulating Membrane Recruitment of the WAVE Complex. *Proceedings of the Na-  
682 tional Academy of Sciences* **107**, 7586–7591 (Apr. 2010).

683 46. Chen, B., Brinkmann, K., Chen, Z., Pak, C. W., Liao, Y., Shi, S., Henry, L., Grishin, N. V., Bogdan, S. &  
684 Rosen, M. K. The WAVE Regulatory Complex Links Diverse Receptors to the Actin Cytoskeleton. *Cell*  
685 **156**, 195–207 (Jan. 2014).

686 47. Truong Quang, B.-A., Mani, M., Markova, O., Lecuit, T. & Lenne, P.-F. Principles of E-Cadherin Supramolec-  
687 ular Organization In Vivo. *Current Biology* **23**, 2197–2207 (Nov. 2013).

688 48. Rubinstein, R., Goodman, K. M., Maniatis, T., Shapiro, L. & Honig, B. Structural Origins of Clustered  
689 Protocadherin-Mediated Neuronal Barcoding. *Seminars in Cell & Developmental Biology. Spectraplakins,  
690 Versatile Roles in Physiology and Pathology* **69**, 140–150 (Sept. 2017).

691 49. Li, J. X. H., Tang, V. W., Boateng, K. A. & Brieher, W. M. Cadherin Puncta Are Interdigitated Dynamic  
692 Actin Protrusions Necessary for Stable Cadherin Adhesion. *Proceedings of the National Academy of Sciences*  
693 **118** (June 2021).

694 50. Strutt, H., Warrington, S. J. & Strutt, D. Dynamics of Core Planar Polarity Protein Turnover and Stable  
695 Assembly into Discrete Membrane Subdomains. *Developmental Cell* **20**, 511–525 (Apr. 2011).

696 51. Cho, B., Pierre-Louis, G., Sagner, A., Eaton, S. & Axelrod, J. D. Clustering and Negative Feedback by En-  
697 docytosis in Planar Cell Polarity Signaling Is Modulated by Ubiquitylation of Prickle. *PLOS Genetics* **11**,  
698 e1005259 (May 2015).

699 52. Stahley, S. N., Basta, L. P., Sharan, R. & Devenport, D. Celsr1 Adhesive Interactions Mediate the Asym-  
700 metric Organization of Planar Polarity Complexes. *eLife* **10** (eds Weis, W. I. & Akhmanova, A.) e62097  
701 (Feb. 2021).

702 53. Hayashi, S., Ito, K., Sado, Y., Taniguchi, M., Akimoto, A., Takeuchi, H., Aigaki, T., Matsuzaki, F., Nak-  
703 agoshi, H., Tanimura, T., Ueda, R., Uemura, T., Yoshihara, M. & Goto, S. GETDB, a Database Compiling  
704 Expression Patterns and Molecular Locations of a Collection of Gal4 Enhancer Traps. *genesis* **34**, 58–61  
705 (2002).

706 54. Golic, K. G. & Lindquist, S. The FLP Recombinase of Yeast Catalyzes Site-Specific Recombination in the  
707 Drosophila Genome. *Cell* **59**, 499–509 (Nov. 1989).

708 55. Golic, K. G. Site-Specific Recombination Between Homologous Chromosomes in Drosophila. *Science* **252**,  
709 958–961 (May 1991).

710 56. Gratz, S. J., Cummings, A. M., Nguyen, J. N., Hamm, D. C., Donohue, L. K., Harrison, M. M., Wildonger,  
711 J. & O'Connor-Giles, K. M. Genome Engineering of *Drosophila* with the CRISPR RNA-Guided Cas9  
712 Nuclease. *Genetics* **194**, 1029–1035 (Aug. 2013).

713 57. Gratz, S. J., Ukkenn, F. P., Rubinstein, C. D., Thiede, G., Donohue, L. K., Cummings, A. M. & O'Connor-  
714 Giles, K. M. Highly Specific and Efficient CRISPR/Cas9-Catalyzed Homology-Directed Repair in *Drosophila*.  
715 *Genetics* **196**, 961–971 (Apr. 2014).

716 58. Port, F., Chen, H.-M., Lee, T. & Bullock, S. L. Optimized CRISPR/Cas Tools for Efficient Germline and  
717 Somatic Genome Engineering in *Drosophila*. *Proceedings of the National Academy of Sciences* **111**, E2967–  
718 E2976 (July 2014).

719 59. Cetera, M., Lewellyn, L. & Horne-Badovinac, S. in *Drosophila: Methods and Protocols* (ed Dahmann, C.)  
720 215–226 (Springer, New York, NY, 2016).

721 60. Spracklen, A. J., Fagan, T. N., Lovander, K. E. & Tootle, T. L. The Pros and Cons of Common Actin  
722 Labeling Tools for Visualizing Actin Dynamics during *Drosophila* Oogenesis. *Developmental Biology* **393**,  
723 209–226 (Sept. 2014).

724 61. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden,  
725 C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P. & Car-  
726 dona, A. Fiji: An Open-Source Platform for Biological-Image Analysis. *Nature Methods* **9**, 676–682 (July  
727 2012).

728 62. Schindelin, J., Rueden, C. T., Hiner, M. C. & Eliceiri, K. W. The ImageJ Ecosystem: An Open Platform  
729 for Biomedical Image Analysis. *Molecular Reproduction and Development* **82**, 518–529 (2015).

730 63. Van der Walt, S., Schönberger, J. L., Nunez-Iglesias, J., Boulogne, F., Warner, J. D., Yager, N., Gouillart, E.  
731 & Yu, T. Scikit-Image: Image Processing in Python. *PeerJ* **2**, e453 (June 2014).

732 64. Virtanen, P. *et al.* SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*  
733 **17**, 261–272 (Mar. 2020).


734 65. napari contributors. *Napari: A Multi-Dimensional Image Viewer for Python* 2019.

735 66. Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering* **9**, 90–95  
736 (May 2007).


737 67. Huang, J., Zhou, W., Dong, W., Watson, A. M. & Hong, Y. Directed, Efficient, and Versatile Modifications  
738 of the *Drosophila* Genome by Genomic Engineering. *Proceedings of the National Academy of Sciences* **106**,  
739 8284–8289 (May 2009).

740 68. Thevenaz, P., Ruttmann, U. & Unser, M. A Pyramid Approach to Subpixel Registration Based on Inten-  
741 sity. *IEEE Transactions on Image Processing* **7**, 27–41 (Jan. 1998).


742 **Supplemental Figures**




**Figure S1: Method used to segment and measure membrane protrusions.** Top row shows an example of a pair of neighboring cells in which one cell is protruding across their shared interface. Bottom row shows a case in which both cells are protruding across the interface. **A**, Cell interfaces and protrusions were labeled with a membrane dye and timelapses of the basal surface were collected. **B**, Cells were automatically segmented with a watershed-based method, and segmentation errors were hand-corrected. **C**, The bright interface region between each pair of neighboring cells was identified using a watershed-based method. This region includes the interface and any membrane protrusions that extend across it. **D**, An enlargement of the boxed regions of (C). **E**, The interface region was divided into two parts by the shortest path from vertex to vertex within the region, which approximates the true cell-cell interface position. The two resulting regions were then assigned to the cell from which they each extended. The area of these regions and the length of the interface between them was used to define average membrane extension length (as described in Methods). **F**, The tip and base of each region were identified, and then used to measure lengths and orientations (see Methods). Associated with Fig. 2.




**Figure S2: Membrane extension length and protrusion orientation in individual egg chambers.** **A**, Plot showing the ratio of protrusive to total edges, with protrusivity defined in terms of membrane extension *longest length* (see Methods). In agreement with the *average length* definition, the protrusivity of *fat2* epithelia is variable, with a distribution overlapping with control and CK-666-treated epithelia. Welch's ANOVA ( $W(2,8.74)=19.3$ ,  $p=0.0006$ ) with Dunnet's T3 multiple comparisons test; n.s.  $p=0.16$ ,  $*p=0.02$ ,  $**p=0.0024$ . **B**, Plot showing the mean membrane extension lengths of control, *fat2*, and CK-666-treated egg chambers, with membrane extension length defined as *average length* (see Methods). With this cutoff-independent protrusivity measurement, the protrusivity of *fat2* egg chambers is intermediate between control and CK-666, with a wider distribution that overlaps both. Welch's ANOVA ( $W(2,7.31)=14.75$ ,  $p=0.0027$ ) with Dunnet's T3 multiple comparisons test; n.s.  $p=0.069$ ,  $*p=0.042$ ,  $**p=0.0036$ . **C**, Polar histograms showing the distribution of membrane protrusion orientations in individual control and *fat2* egg chambers. Anterior is left, posterior is right, and images were flipped as needed so that migration is downward for control epithelia, in which membrane protrusions are biased in the direction of migration. In *fat2* epithelia, protrusions have varying levels of axial bias and little or no vectorial bias. Associated with Fig. 2.




**Figure S3: Actin protrusions are reduced and unpolarized without Fat2 and further reduced without the WAVE complex.** **A**, Images showing phalloidin staining of F-actin in control, *fat2*, and *abi-RNAi*-expressing epithelia. Bottom row shows the same images with displayed brightness increased. From dataset quantified in C,E,F. **B**, Examples of segmented cell-cell interfaces or medial basal surfaces overlayed on F-actin. **C**, Plot of the difference in F-actin fluorescence intensity between cell interfaces and medial basal surfaces shows that while *fat2* and *abi-RNAi* epithelia have less F-actin interface enrichment than control epithelia, F-actin interface enrichment remains higher in *fat2* epithelia than *abi-RNAi* epithelia. Welch's ANOVA ( $W(2, 19.84)=94.68$ ,  $p<0.0001$ ) with Dunnet's T3 multiple comparisons test; \* $p=0.033$ , \*\*\*\* $p<0.0001$ . **D**, Frames from timelapse movies of control, *fat2*, and *abi-RNAi* epithelia with F-actin labeled with F-Tractin-tdTomato. As with phalloidin staining, the protrusivity of *fat2* epithelia is intermediate between that of control and *abi-RNAi* epithelia. Brightness display settings vary between genotypes to correct for variability in F-Tractin-tdTomato expression levels. See related Movie 2. **E**, Plot of F-actin fluorescence intensity at cell interfaces as a function of interface angular distance from horizontal. Gray bars below the x-axis represent interface angles. **F**, Plot of the F-actin fluorescence ratio between leading-trailing (0-10°) and side (80-90°) interfaces, a measure of F-actin enrichment along leading-trailing interfaces. F-actin is enriched at leading-trailing interfaces in control, but not *fat2*, egg chambers. Control-*fat2* comparisons: unpaired t-test, \*\*\*\* $p<0.0001$ . Comparison between *fat2* and one (dashed line, the expectation if there is no enrichment): one sample t-test, n.s.  $p=0.88$ . **C,E,F**, Bars (C,F) or lines and shaded regions (E) indicate mean  $\pm$  SD. Associated with Fig. 2; Movie 2.




**Figure S4: Fat2 acts locally across the cell interface to orient membrane protrusions.** **A**, Timelapse frame of a *fat2* mosaic epithelium with cell membrane labeled, used to evaluate protrusion orientations in control or *fat2* cells within a migratory context. Boxes indicate examples of leading-trailing interfaces between neighbor pairs with each possible combination of genotypes. See related Movie 4. **B**, Larger images of the interfaces boxed in (A), showing that protrusions are misoriented when *fat2* cells are ahead of the interface regardless of the genotype of the cell behind the interface. Arrows point in the direction of protrusion. Associated with Fig. 2; Movie 4.




**Figure S5: Evaluation of endogenous Sra1-GFP functionality.** **A**, Images comparing the localization of markers of WAVE complex subunits: Sra1-GFP, Scar antibody, and Abi-mCherry, at the basal surface (top row) and in cross-section (bottom row). **B**, Images of Sra1-GFP localization in control and *abi*-RNAi-expressing epithelia. Sra1-GFP is dispersed in the absence of Abi. **C**, Images showing phalloidin-stained F-actin in epithelia with wild-type Sra1, one or two copies of Sra1-GFP, or expressing *sra1*-RNAi, used to assess the appearance of protrusions in each condition. **D**, Plot of migration speed of epithelia with wild-type Sra1 or one or two copies of Sra1-GFP. Migration speed is reduced when both Sra1 copies are GFP-tagged. One-way ANOVA ( $F(2,49)=18.37$ ,  $p<0.0001$ ) with post-hoc Tukey's test; n.s. 0.66, \*\*\*\* $p<0.0001$ . See related Movie 5. Associated with Fig. 3; Movie 5.



**Figure S6: Fat2 concentrates the WAVE complex at cell-cell interfaces and polarizes it across the epithelium.** A, Images of Sra1-GFP at the basal surface in control and *fat2* epithelia. B, Examples of segmented cell-cell interfaces or medial basal surfaces overlaid on Sra1-GFP images. C, Plot of mean Sra1-GFP fluorescence intensity across the entire basal surface (total), at cell-cell interfaces, and at medial basal surfaces in control and *fat2* epithelia. One-way ANOVA ( $F(5,111)=63.22$ ,  $p<0.0001$ ) with post-hoc Šidák's test; n.s. (left to right)  $p=0.67$ ,  $0.64$ ,  $***p<0.0001$ . D, Plot of Sra1-GFP fluorescence at cell-cell interfaces as a function of interface angular distance from horizontal. Gray bars below the  $x$ -axis represent interface angles. E, Plot of the Sra1-GFP fluorescence intensity ratio between leading-trailing (0-10°) and side (80-90°) interfaces, a measure of Sra1-GFP enrichment along leading-trailing interfaces. Sra1-GFP is enriched at leading-trailing interfaces in control, but not *fat2*, epithelia. Control-*fat2* comparison: unpaired t-test,  $***p<0.0001$ . Comparison between *fat2* and one (dashed line, the expectation if there is no enrichment): one sample t-test, n.s.  $p=0.052$ . Bars (C,E) or lines and shaded regions (D) indicate mean  $\pm$  SD. Associated with Fig. 3.



**Figure S7: Fat2 stabilizes domains of WAVE complex enrichment locally across the cell-cell interface.** **A**, Timelapse frames showing pairs of cell interfaces from control or *fat2* epithelia expressing Sra1-GFP and labeled with a membrane dye. Arrows indicate membrane protrusions. Sra1-GFP is enriched at protrusion tips in both control and *fat2* epithelia. See related Movie 6. **B**, Timelapse frame of a *fat2* mosaic epithelium in which all cells express Sra1-GFP, used to evaluate Sra1-GFP dynamics in control or *fat2* cells within a migratory context. Boxes indicate a leading-trailing interface between two control cells (blue) or *fat2* cells (green). See related Movie 9. **C**, Larger images of the interfaces boxed in (B), taken 9.8 minutes apart. Sra1-GFP is initially enriched along both interfaces. It remains enriched in the control interface throughout, but loses enrichment along the *fat2* interface. Associated with Fig. 4; Movies 6, 9.



**Figure S8: Ena and Lar are not required for colocalization between Fat2 and the WAVE complex.** **A**, Images of cells expressing Fat2-3xGFP and Abi-mCherry in control, *lar*, and *ena*-RNAi backgrounds (top 3 rows) or E-cadherin-GFP and Abi-mCherry (bottom row, negative control for colocalization measurements). **B**, Plot of Spearman's correlation coefficients of Abi-mCherry and Fat2-3xGFP (or Fat2 $^{\Delta ICD}$ -3xGFP) (gray background) or E-cadherin-GFP (white background) along leading-trailing interfaces show that Fat2 and Abi colocalize in all four conditions more strongly than E-cadherin and Abi. Bars indicate mean  $\pm$  SD. One-way ANOVA ( $F(5,81)=44.86$ ,  $p=0.0164$ ) with post-hoc Tukey's test; n.s. (left to right)  $p>0.99$ ,  $p=0.41$ ,  $**p<0.0046$ ,  $***p<0.0001$ . **A,B**, Control, Fat2-3xGFP, and Abi-mCherry images and Spearman's coefficients are also in Fig. 5A,E. **C**, Line scan of GFP-Ena, Abi-mCherry, and F-actin (phalloidin) fluorescence intensity along a leading-trailing interface region, showing their corresponding peaks of enrichment. **D**, Image showing the GFP-Ena, Abi-mCherry, and F-actin (phalloidin) at the leading edge and in the boxed filopodium. **E**, Plot of mean fluorescence intensity of F-actin, Abi-mCherry, and GFP-Ena along the length of filopodia showing their relative distribution. Lines and shaded regions indicate mean  $\pm$  SD.  $n=54$  filopodia (used for SD), 39 cells from 2 egg chambers. **F**, Images of F-actin (phalloidin) and cell interfaces (anti-Discs Large) in control, *ena*-RNAi, and *abi*-RNAi backgrounds. Expression of *ena*-RNAi strongly depletes filopodia, and *abi*-RNAi expression removes both filopodia and lamellipodia. **G**, Plot of mean fluorescence intensity of Abi-mCherry along leading-trailing interfaces in control epithelia or similarly-oriented interfaces in *lar* epithelia, some of which are non-migratory. Associated with Fig. 5.




Figure S9: **The WAVE complex occasionally accumulates at side-facing interfaces away from Fat2.** Timelapse frames of a side-facing cell-cell interface from an epithelium expressing Abi-mCherry and Fat2-3xGFP. Arrows indicate a site of transient Abi-mCherry accumulation, protrusion, and dissipation with no corresponding Fat2-3xGFP enrichment. See related Movie 12. Associated with Fig. 5; Movie 12.

743 **Movie captions**

**Movie 1: Membrane protrusivity of control, *fat2*, and CK-666-treated epithelia.** Control, *fat2*, and CK-666-treated epithelia labeled with a membrane dye. Bottom row shows segmented edges. Protrusive edges, defined as ones with membrane extensions longer than the 98th percentile of those of CK-666-treated epithelia, are red. Non-protrusive edges are white. The control field of view moves to follow the cells as they migrate. Acquired with laser scanning confocal microscope. Associated with Fig. 2A.

**Movie 2: F-actin protrusivity and protrusion polarity of control and *fat2* epithelia.** Control, *fat2*, and *abi*-RNAi epithelia with F-actin labeled with F-Tractin-tdTomato. The protrusivity of *fat2* epithelia is intermediate between that of control and *abi*-RNAi epithelia. Brightness display settings vary between genotypes to correct for variability in F-Tractin-tdTomato expression levels. Acquired with TIRF microscope. Associated with Fig. S3D.

**Movie 3: Protrusion orientation in control and *fat2* epithelia.** Control and *fat2* epithelia are labeled with a membrane dye, and arrows indicating the orientation of protrusions are overlaid. Arrows originate at protrusion bases and have lengths proportional to protrusion lengths. Acquired with laser scanning confocal microscope. Associated with Fig. 2A.

**Movie 4: Membrane protrusion in a *fat2* mosaic epithelium.** A *fat2* mosaic epithelium with cell membrane labeled, used to evaluate protrusion orientations in control or *fat2* cells within a migratory context. Boxes indicate examples of leading-trailing interfaces between neighbor pairs with each possible combination of genotypes. Acquired with laser scanning confocal microscope. Associated with Fig. S4.

**Movie 5: Migration of epithelia with endogenously-tagged Sra1-GFP.** Epithelia with unlabeled Sra1 (Control), one copy of Sra1-GFP and one unlabeled Sra1, or two copies of Sra1-GFP, imaged at the mid-plane between apical and basal cell surfaces. Representative of timelapse movies used to measure migration speed. Acquired with laser scanning confocal microscope. Associated with Fig. S5D.

**Movie 6: Sra1 enrichment at protrusion tips in control and *fat2* epithelia.** Pairs of cell edges from control or *fat2* epithelia expressing Sra1-GFP and labeled with a membrane dye. Sra1-GFP is enriched at protrusion tips in both control and *fat2* epithelia. The control movie field of view moves to follow the cells as they migrate. Associated with Fig. S7A.

**Movie 7: WAVE complex-enriched domain dynamics in control and *fat2* cells.** Cells from control and *fat2* epithelia expressing Sra1-GFP. Laser intensity and brightness display settings differ between genotypes. Used to evaluate the stability of domains of Sra1-GFP accumulation. The control field of view moves to follow the cell as it migrates. Associated with Fig. 4B,D.

**Movie 8: WAVE complex-enriched domain dynamics in control and *fat2* epithelia.** Fields of cells from control and *fat2* epithelia expressing Sra1-GFP. The control movie field of view moves to follow the cells as they migrate. Laser intensity and brightness display settings differ between genotypes. Used to evaluate the stability of domains of Sra1-GFP accumulation. Wider view of the epithelia shown in Fig. 4B,D.

Movie 9: **WAVE complex-enriched domain dynamics in a *fat2* mosaic epithelium.** A *fat2* mosaic epithelium in which all cells express Sra1-GFP, used to evaluate WAVE complex-enriched domain dynamics in control or *fat2* cells within a migratory context. Associated with Fig. S7B,C.

Movie 10: **Dynamics of protrusive domains in control and *fat2* cells.** Top row shows the interfaces and membrane protrusions of one cell and its neighbors, labeled with a membrane dye. The segmented membrane extensions originating from the centered cell are overlaid in the bottom row. The control field of view moves to follow the cell as it migrates. Associated with Fig. 4E.

Movie 11: **Colocalization of puncta of Fat2 and the WAVE complex along leading-trailing interfaces.** The leading-trailing interfaces of two cells expressing Fat2-3xGFP and Abi-mCherry, used to compare the distributions of Fat2 and the WAVE complex over time. The Fat2-3xGFP channel is offset 2 pixels downward so puncta positions can be more easily compared. Associated with Fig. 5B.

Movie 12: **WAVE complex accumulation at side-facing protrusions away from Fat2.** A side-facing cell-cell interface from an epithelium expressing Abi-mCherry and Fat2-3xGFP. Arrows indicate a site of transient Abi-mCherry accumulation, protrusion, and dissipation with no corresponding Fat2-3xGFP enrichment, in contrast to the colocalized Fat2-3xGFP and Abi-mCherry on the leading-trailing interfaces above and below. Associated with Fig. S9.

Supp. Table 1: Key resources

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
gene ( <i>Drosophila melanogaster</i> )	Abi	NA	FLYB:FBgn0020510	FlyBase Name: Abelson interacting protein
gene ( <i>Drosophila melanogaster</i> )	Dlg	NA	FLYB:FBgn0001624	FlyBase Name: discs large 1
gene ( <i>Drosophila melanogaster</i> )	E-cadherin	NA	FLYB:FBgn0003391	FlyBase Name: shotgun
gene ( <i>Drosophila melanogaster</i> )	Ena	NA	FLYB:FBgn0000578	FlyBase Name: enabled
gene ( <i>Drosophila melanogaster</i> )	Fat2 (Kug)	NA	FLYB:FBgn0261574	FlyBase Name: kugelei
gene ( <i>Drosophila melanogaster</i> )	Lar	NA	FLYB:FBgn0000464	FlyBase Name: Leukocyte-antigen-related-like
gene ( <i>Drosophila melanogaster</i> )	Scar	NA	FLYB:FBgn0041781	FlyBase Name: SCAR
gene ( <i>Drosophila melanogaster</i> )	Sra1 (CYFIP)	NA	FLYB:FBgn0038320	FlyBase Name: Cytoplasmic FMR1 interacting protein
genetic reagent ( <i>Drosophila melanogaster</i> )	Abi-mCherry or ubi>Abi-mCherry	Bloomington Drosophila Stock Center; FLYB:FBrlf0227194 (S. Huelsmann)	FLYB:FBst0058729; BDSC:58729	FlyBase Symbol: P{Ubi-mCherry.Abi}3
genetic reagent ( <i>Drosophila melanogaster</i> )	abi-RNAi	National Institute of Genetics, Japan	FLYB:FBtp0079430; NIG:9749R	
genetic reagent ( <i>Drosophila melanogaster</i> )	E-cadherin-GFP	Bloomington Drosophila Stock Center; PMID:19429710	FLYB:FBst0060584; BDSC:60584	FlyBase Genotype: y[1] w*; Ti[T]shg[GFP]
genetic reagent ( <i>Drosophila melanogaster</i> )	GFP-Ena or ubi>GFP-Ena	Bloomington Drosophila Stock Center; FLYB:FBrlf0208868 (S. Nowotarski & M. Peiger)	FLYB:FBst0028798; BDSC:28798	FlyBase Genotype: w*; P{Ubi-GFP.ena}3
genetic reagent ( <i>Drosophila melanogaster</i> )	ena-RNAi	Vienna Drosophila Resource Center	FLYB:FBst0464896; VDRC:43058	
genetic reagent ( <i>Drosophila melanogaster</i> )	Fat2-3xGFP FRT80B	Laboratory of S. Horne-Badovinac; PMID:28292425	FLYB:FBal0326664	FlyBase Symbol: kug[3xGFP]
genetic reagent ( <i>Drosophila melanogaster</i> )	Fat2[ΔICD]-3xGFP FRT80B	Laboratory of S. Horne-Badovinac; PMID:28292425	FLYB:FBal0326665	FlyBase Symbol: kug[ΔICD.3xGFP]
genetic reagent ( <i>Drosophila melanogaster</i> )	fat2 or fat2[N103-2] FRT80B	Laboratory of Sally Horne-Badovinac; PMID:22413091	FLYB:FBal0267777	FlyBase Symbol: kug[N103-2]
genetic reagent ( <i>Drosophila melanogaster</i> )	UAS>Flp	Bloomington Drosophila Stock Center; PMID:9584125	FLYB:FBst0004539; BDSC:4539	FlyBase Genotype: y[1] w[*]; P{UAS-FLP.D}JD1
genetic reagent ( <i>Drosophila melanogaster</i> )	FRT80B	Bloomington Drosophila Stock Center; PMID:8404527	FLYB:FBti0002073	FlyBase Symbol: P{neoFRT}80B
genetic reagent ( <i>Drosophila melanogaster</i> )	UAS>F-Tractin-tdTomato	Bloomington Drosophila Stock Center; FLYB:FBrlf0226873 (T. Tootle); PMID:24995797	FLYB:FBst0058989; BDSC:58989	FlyBase Genotype: w*; P{UASp-F-Tractin.tdTomato}15A/SM6b; MKRS/TM2
genetic reagent ( <i>Drosophila melanogaster</i> )	ubi>GFP-NLS (3L) FRT80B	Bloomington Drosophila Stock Center; FLYB:FBrlf0108530 (D. Bilder & N. Perrimon)	FLYB:FBst0001620; BDSC:1620	FlyBase Genotype: w*; P{Ubi-GFP.D}61EF P{neoFRT}80B
genetic reagent ( <i>Drosophila melanogaster</i> )	lar[13.2] FRT40A	Bloomington Drosophila Stock Center; PMID:8598047	FLYB:FBst0008774; BDSC:8774	
genetic reagent ( <i>Drosophila melanogaster</i> )	lar[bola1]	Bloomington Drosophila Stock Center; PMID:11688569	FLYB:FBst0091654; BDSC:91654	
genetic reagent ( <i>Drosophila melanogaster</i> )	MKRS hsFLP/TM6b, Cre	Bloomington Drosophila Stock Center	FLYB:FBst0001501; BDSC:1501	y[1] w[67c23]; MKRS, P{hsFLP}86E/TM6B, P{Crew} DH2, Tb[1]
genetic reagent ( <i>Drosophila melanogaster</i> )	nanos-Cas9	Bloomington Drosophila Stock Center; FLYB:FBrlf0223952 (F. Port & S. Bullock); PMID:25002478	FLYB:FBst0054591; BDSC:54591	FlyBase Genotype: y[1] M{nos-Cas9.P}ZH-2A w*
genetic reagent ( <i>Drosophila melanogaster</i> )	ubi>mRFP-NLS (3L) FRT80B	Bloomington Drosophila Stock Center; FLYB:FBrlf0210705 (J. Lipsick)	FLYB:FBti0129786; BDSC:30852	FlyBase Genotype: w1118; P{Ubi-mRFP.nls}3L P{neoFRT}80B

Supp. Table 1: Key resources

745				
genetic reagent (Drosophila melanogaster)	FRT82b ubi>mRFP-NLS (3R)	Bloomington Drosophila Stock Center; FLYB:FBref0210705 (J. Lipsick)	FLYB:FBst0030555; BDSC:30555	FlyBase Genotype: w1118; P{neoFRT}82B P{Ubi-mRFP.nls}3R
genetic reagent (Drosophila melanogaster)	Sra1-GFP	this paper		
genetic reagent (Drosophila melanogaster)	Sra1-GFP FRT80B	this paper		
genetic reagent (Drosophila melanogaster)	sra1-RNAi	Bloomington Drosophila Stock Center; PMID:21460824	FLYB:FBst0038294; BDSC:38294	FlyBase Genotype: y[1] sc <sup>-</sup> v[1] sev[21]; P{TRiP.HMS01754} attP2
genetic reagent (Drosophila melanogaster)	TJ>Gal4	National Institute of Genetics, Japan; PMID:12324948	FLYB:FBtp0089190; DGRC:104055	FlyBase Symbol: P{tj-GAL4.U}
genetic reagent (Drosophila melanogaster)	w1118	Bloomington Drosophila Stock Center	FLYB:FBal0018186	
antibody	Discs Large; Dlg	Developmental Studies Hybridoma Bank	DSHB:4F3; RRID:AB_528203	
antibody	Scar	Developmental Studies Hybridoma Bank	AB_2618386	1:200
antibody	Alexa Fluor™ 647, donkey anti-mouse secondary	Thermo Fisher Scientific	Cat:A31571; RRID:AB_162542	1:300, 3 hrs at room temp
chemical compound, drug	CellMask™ Orange Plasma Membrane Stain	Thermo Fisher Scientific	Cat:C10045	1:250, 15 min
chemical compound, drug	CellMask™ Deep Red Plasma Membrane Stain	Thermo Fisher Scientific	Cat:C10046	1:250, 15 min
chemical compound, drug	TRITC Phalloidin	Millipore Sigma	Cat:1951	1:300, 15 min at room temp
chemical compound, drug	Alexa Fluor™ 647 phalloidin	Thermo Fisher Scientific	Cat:A22287	1:50, 3 hrs at room temp
chemical compound, drug	CK-666, Arp2/3 complex inhibitor	Millipore Sigma	Cat:553502	750 µM
chemical compound, drug	Formaldehyde, 16%, methanol free, ultra pure	Polysciences	Cat:18814-10	
chemical compound, drug	Recombinant human insulin	Millipore Sigma	Cat:12643	
recombinant DNA reagent	plasmid: pU6-BbsI-chiRNA	Addgene	Addgene:45946; RRID:Addgene_45946	PMID:23709638
recombinant DNA reagent	plasmid: pU6 chiRNA Sra1 C-term	this paper		CRISPR chiRNA construct for generation of Sra1-GFP
recombinant DNA reagent	plasmid: pDsRed-attP	Addgene	Addgene:51019	Vector used to make pDsRed-attP Sra1-GFP HR
recombinant DNA reagent	plasmid: pTWG	Drosophila Genome Resource Center	DGRC:1076	source of enhanced GFP for generation of Sra1-GFP
recombinant DNA reagent	plasmid: pDsRed-attP Sra1-GFP HR	this paper		CRISPR homologous recombination construct for generation of Sra1-GFP
software, algorithm	Zen Blue	Zeiss		
software, algorithm	MetaMorph	Molecular Devices		
software, algorithm	FIJI (ImageJ)	PMID:22743772		
software, algorithm	GraphPad Prism 9 for Mac	GraphPad Software		
software, algorithm	Microsoft Excel for Mac, version 16.47	Microsoft		
software, algorithm	Python 3	Python Software Foundation		<a href="https://www.python.org">https://www.python.org</a>
software, algorithm	imageio	imageio contributors		<a href="https://imageio.readthedocs.io/">https://imageio.readthedocs.io/</a>
software, algorithm	matplotlib	The Matplotlib Development team		<a href="https://matplotlib.org/">https://matplotlib.org/</a>
software, algorithm	napari	napari contributors		<a href="https://napari.org/">https://napari.org/</a>
software, algorithm	numpy	numpy contributors		<a href="https://numpy.org/">https://numpy.org/</a>
software, algorithm	pims	pims contributors		<a href="http://soft-matter.github.io/pims/v0.5/">http://soft-matter.github.io/pims/v0.5/</a>
software, algorithm	pandas	pandas contributors		<a href="https://pandas.pydata.org/">https://pandas.pydata.org/</a>
software, algorithm	scikit-image	scikit-image development team		<a href="https://scikit-image.org/">https://scikit-image.org/</a>

Supp. Table 1: Key resources

746 software, algorithm	scikit-fmm	scikit-fmm contributors		<a href="https://pythonhosted.org/scikit-fmm/">https://pythonhosted.org/scikit-fmm/</a>
software, algorithm	scipy	scipy contributors		<a href="https://scipy.org/">https://scipy.org/</a>

Supp. Table 2. Genotypes of experimental females

Figure	Panel	Name	Genotype
2 <sup>747</sup>	A-C	Control	w1118
		<i>fat2</i>	w <sup>+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B
		CK-666	w1118
	D	Control	w1118
		<i>fat2</i>	w <sup>+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B
3	B	Sra1-GFP mosaic	w; tj>Gal4DGRC:104055, UAS>Flp <sup>BDSC:4539/+</sup> ; FRT82B Sra1-GFP/FRT82B ubi>mRFP-NLS <sup>BDSC:30555</sup>
	C	<i>fat2</i> mosaic	w; tj>Gal4DGRC:104055, UAS>Flp <sup>BDSC:4539/+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B/ubi>GFP-NLS FRT80B <sup>BDSC:1620</sup>
	D-F	<i>fat2</i> mosaic + Sra1	w; tj>Gal4DGRC:104055, UAS>Flp <sup>BDSC:4539/+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B Sra1-GFP/ubi>mRFP-NLS FRT80B <sup>BDSC:30852</sup>
4	A	Sra1-GFP mosaic + <i>fat2</i>	w; tj>Gal4DGRC:104055, UAS>Flp <sup>BDSC:4539/+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B FRT82B Sra1-GFP/ <i>fat2</i> <sup>N103-2</sup> FRT80B FRT82B
	B,D	Control	w <sup>+</sup> ; Sra1-GFP/+
		<i>fat2</i>	w <sup>+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B Sra1-GFP/ <i>fat2</i> <sup>N103-2</sup> FRT80B
	E	Control	w1118
		<i>fat2</i>	w; <i>fat2</i> <sup>N103-2</sup> FRT80B
5	A,E	Fat2 + Abi	w <sup>+</sup> ; ubi>Abi-mCherry <sup>BDSC:58729</sup> , Fat2-3xGFP FRT80B/ Fat2-3xGFP FRT80B
		Fat2 <sup>ΔICD</sup> + Abi	w <sup>+</sup> ; ubi>Abi-mCherry <sup>BDSC:58729</sup> , Fat2 <sup>ΔICD</sup> -3xGFP FRT80B/Fat2-3xGFP FRT80B
	B	Fat2 + Abi	w <sup>+</sup> ; ubi>Abi-mCherry <sup>BDSC:58729</sup> , Fat2-3xGFP FRT80B/ Fat2-3xGFP FRT80B
	C,D,F,G	Fat2 + Abi + F-actin	w <sup>+</sup> ; ubi>Abi-mCherry <sup>BDSC:58729</sup> , Fat2-3xGFP FRT80B/ Fat2-3xGFP FRT80B
S1	Top row	Protrusion in 1 direction	w1118
	Bottom row	Protrusion in both directions	w; <i>fat2</i> <sup>N103-2</sup> FRT80B
S2	A	Control	w1118
		<i>fat2</i>	w <sup>+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B
		CK-666	w1118 (treated with 750 μM CK-666)
	B	Control	w1118
		<i>fat2</i>	w <sup>+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B
S3	A-C	Control	w; tj>Gal4DGRC:104055/+
		<i>fat2</i>	w; tj>Gal4DGRC:104055/+; <i>fat2</i> <sup>N103-2</sup> FRT80B
		<i>abi</i> -RNAi	w; tj>Gal4DGRC:104055/+; UAS> <i>abi</i> -RNAi <sup>IG:9749R-3/+</sup>
	D	Control	w; tj>Gal4DGRC:104055/UAS>F-Tractin-tdTomato <sup>BDSC:58989</sup>
		<i>fat2</i>	w; tj>Gal4DGRC:104055/UAS>F-Tractin-tdTomato <sup>BDSC:58989</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B
		<i>abi</i> -RNAi	w; tj>Gal4DGRC:104055/UAS>F-Tractin-tdTomato <sup>BDSC:58989</sup> ; UAS> <i>abi</i> -RNAi <sup>IG:9749R-3/+</sup>
	E,F	Control	w1118
		<i>fat2</i>	w <sup>+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B
S4	A,B	<i>fat2</i> mosaic	w; tj>Gal4DGRC:104055, UAS>Flp <sup>BDSC:4539/+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B/ubi>GFP-NLS FRT80B <sup>BDSC:1620</sup>
S5	A	Sra1-GFP	w <sup>+</sup> ; Sra1-GFP
		anti-SCAR	w1118
		ubi>Abi-mCherry	w; ubi>Abi-mCherry <sup>BDSC:58729/+</sup>
	B	Control	w; tj>Gal4DGRC:104055/+
		<i>abi</i> -RNAi	w; tj>Gal4DGRC:104055/+; UAS> <i>abi</i> -RNAi <sup>IG:9749R-3/+</sup>
	C	Control	w1118
		Sra1-GFP x1	w <sup>+</sup> ; Sra1-GFP/+

Supp. Table 2. Genotypes of experimental females

748	D	Sra1-GFP x2	w;; Sra1-GFP
		<i>sra1</i> -RNAi	w; tj>Gal4DGRC:104055/+; UAS> <i>sra1</i> -RNAi <sup>BDSC:38294/+</sup>
S6	A,C-E	Control	w1118
		Sra1-GFP x1	w;; Sra1-GFP/+
		Sra1-GFP x2	w;; Sra1-GFP
S7	A	Control	w;; Sra1-GFP/+
		<i>fat2</i>	w;; <i>fat2</i> <sup>N103-2</sup> FRT80B Sra1-GFP/ <i>fat2</i> <sup>N103-2</sup> FRT80B
	B,C	<i>fat2</i> mosaic + Sra1	w; tj>Gal4DGRC:104055, UAS>Flp <sup>BDSC:4539/+</sup> ; <i>fat2</i> <sup>N103-2</sup> FRT80B Sra1-GFP/ubi>mRFP-NLS FRT80B <sup>BDSC:30852</sup>
S8	A,B	Control Fat2 + Abi	w;; ubi>Abi-mCherry <sup>BDSC:58729</sup> , Fat2-3xGFP FRT80B/ Fat2-3xGFP FRT80B
		<i>ena</i> -RNAi, Fat2 + Abi	w; tj>Gal4DGRC:104055/UAS> <i>ena</i> -RNAi <sup>VDRC:43058</sup> ; ubi>Abi-mCherry <sup>BDSC:58729</sup> , Fat2-3xGFP FRT80B/Fat2-3xGFP FRT80B
		<i>lar</i> Fat2 + Abi	w; <i>larbola1</i> <sup>BDSC:91654</sup> / <i>lar</i> <sup>13.2</sup> <sup>BDSC8774</sup> FRT40A; ubi>Abi-mCherry <sup>BDSC:58729</sup> , Fat2-3xGFP FRT80B/Fat2-3xGFP FRT80B
		Control E-cadherin + Abi	w; Shg-GFP <sup>BDSC:60584/+</sup> ; ubi>Abi-mCherry <sup>BDSC:58729/+</sup>
	C-E	Ena + Abi + F-actin	w; ubi>GFP-Ena <sup>BDSC:28798</sup> /ubi>Abi-mCherry <sup>BDSC:58729</sup>
	F	Control	w; tj>Gal4DGRC:104055/+
		<i>ena</i> -RNAi	w; tj>Gal4DGRC:104055/UAS- <i>ena</i> -RNAi <sup>VDRC:43058</sup>
		<i>abi</i> -RNAi	w; tj>Gal4DGRC:104055/+; UAS- <i>abi</i> -RNAi <sup>NIG:9749R-3/+</sup>
	G	Control	w;; ubi>Abi-mCherry <sup>BDSC:58729</sup> , Fat2-3xGFP FRT80B/ Fat2-3xGFP FRT80B
		<i>lar</i>	w; <i>larbola1</i> <sup>BDSC:91654</sup> / <i>lar</i> <sup>13.2</sup> <sup>BDSC8774</sup> FRT40A; ubi>Abi-mCherry <sup>BDSC:58729</sup> , Fat2-3xGFP FRT80B/Fat2-3xGFP FRT80B
S9		Fat2 + Abi	w;; ubi>Abi-mCherry <sup>BDSC:58729</sup> , Fat2-3xGFP FRT80B/ Fat2-3xGFP FRT80B

Supp. Table 3: *Drosophila* culture conditions for experiments

749

Figure	Panel	Days on yeast	Temp (°C)
2	A-D	2-3	25
3	B	7	25
	C	3	25
	D-F	3	25
4	A	5	25
	B,D	2-3	25
	E	2-3	25
5	A,E	2-3	25
	B	2-3	25
	C,D,F,G	2-3	25
S1	A-F	2-3	25
S2	A,B	2-3	25
S3	A-C,E,F	2-3	29
	D	2-3	29
S4	A,B	3	25
S5	A	2-3	25
	B	3	29
	C	3	29
	D-F	2-3	25
S6	A-E	2-3	25
S7	A	2-3	25
	B,C	3	25
S8	A,B,G	3	29
	C-E	2-3	25
	F	3	29
S9		2-3	25