

1 **Title** Sex-biased expression is associated with chromatin state in *D. melanogaster* and *D. simulans*

2 Adalena V. Nanni^{1,2}, Natalie Martinez¹, Rita Graze³, Alison Morse^{1,2}, Jeremy R. B. Newman²,
3 Vaibhav Jain¹, Srna Vlaho⁴, Sarah Signor⁵, Sergey V. Nuzhdin⁴, Rolf Renne^{1,2}, Lauren M.
4 McIntyre^{1,2}

5 1)Department of Molecular Genetics and Microbiology, University of Florida, Gainesville, FL

6 2) University of Florida Genetics Institute, University of Florida, Gainesville, FL, USA

7 3) Department of Biological Sciences, Auburn University, Auburn, AL, USA

8 4) Department of Biological Sciences, University of Southern California, Los Angeles, CA, USA

9 5) Department of Biological Sciences, North Dakota State University, Fargo, ND, USA

10 **Abstract**

11 We propose a new model for the association of chromatin state and sex-bias in expression. We
12 hypothesize enrichment of open chromatin in the sex where we see expression bias (OS) and closed
13 chromatin in the opposite sex (CO). In this study of *D. melanogaster* and *D. simulans* head tissue,
14 sex-bias in expression is associated with H3K4me3 (open mark) in males for male-biased genes
15 and in females for female-biased genes in both species. Sex-bias in expression is also largely
16 conserved in direction and magnitude between the two species on the X and autosomes. In male-
17 biased orthologs, the sex-bias ratio is more divergent between species if both species have
18 H3K27me2me3 marks in females compared to when either or neither species has H3K27me2me3
19 in females. H3K27me2me3 marks in females are associated with male-bias in expression on the
20 autosomes in both species, but on the X only in *D. melanogaster*. In female-biased orthologs the
21 relationship between the species for the sex-bias ratio is similar regardless of the H3K27me2me3
22 marks in males. Female-biased orthologs are more similar in the ratio of sex-bias than male-biased
23 orthologs and there is an excess of male-bias in expression in orthologs that gain/lose sex-bias.
24 There is an excess of male-bias in sex-limited expression in both species suggesting excess male-
25 bias is due to rapid evolution between the species. The X chromosome has an enrichment in male-
26 limited H3K4me3 in both species and an enrichment of sex-bias in expression compared to the
27 autosomes.

28 **Introduction**

29 Chromatin accessibility is known to be important for multiple levels of gene regulation, as well as
30 in large scale modifications of expression such as in dosage compensation of sex chromosomes.
31 H3K4me3 is an open chromatin mark correlated with activate expression (Santos-Rosa, et al. 2002;
32 Schneider, et al. 2004) and closed chromatin marks H3K27me2 and H3K27me3, together referred
33 to as H3K27me2me3, are correlated with silenced expression (Wang, et al. 2008; Juan, et al. 2016).
34 These three marks act together with other histone modifications, DNA methylation, and chromatin
35 factors, in the establishment and modification of chromatin accessibility (reviewed in Boros 2012).
36 and the extent to which epigenetics influences behavior is an emerging paradigm explored in
37 several systems (e.g., Spannhoff, et al. 2011; Sun, et al. 2015; Elliott, et al. 2016; Sun, et al. 2016;

38 Opachaloemphan, et al. 2018; Qin, et al. 2018; Bludau, et al. 2019) including in plastic behavioral
39 traits such as foraging (Anreiter, et al. 2017; Anreiter and Sokolowski 2019).

40 Differences between the X chromosome and autosomes in the evolution of gene expression may
41 be due to changes in regulation of chromatin conformation. There is evidence in third instar larvae
42 for an enrichment of open chromatin marks on the X chromosome compared to the autosomes in
43 both males and females of *D. miranda* and *D. melanogaster*, as well as more open chromatin in
44 males compared to females on the X chromosome and more closed chromatin in females compared
45 to males (Brown and Bachtrog 2014). Furthermore, Brown, et al. (Brown, et al. 2020) also
46 demonstrated that the male Y chromosome has a genome-wide effect on heterochromatin factors,
47 leading to a heterochromatin sink effect. Specifically, the male X chromosome and autosomes are
48 more open compared to the female due to a sequestering of closed chromatin factors on the Y
49 chromosome (Henikoff 1996; Francisco and Lemos 2014).

50
51 Chromatin remodeling genes have known roles in sex determination and may be important for
52 overall regulation of sex differences in gene expression. For example, the sex determination gene
53 *fru* aids in recruiting of histone deacetylase (HDAC) *Rpd3* and heterochromatin protein 1A (HP1a)
54 encoded by *Su(var)205* (Ito, et al. 2012) to genes associated with male courtship behaviors. The
55 expression of *fru* decreases with mutation of a histone demethylase *kdm4* (Lorbeck, et al. 2010),
56 resulting in the male chain mating phenotype, also found in *fru* mutants (Ito, et al. 1996). Sexually
57 dimorphic chromatin modifications such as H3K9me2 (associated with closed chromatin) and
58 H4K16ac (associated with open chromatin) have been reported (Brown and Bachtrog 2014).

59
60 In *D. melanogaster*, sexually dimorphic chromatin accessibility is stage and cell-type specific
61 (Palmateer, et al. 2021). For example, in *fru-P1*-expressing neurons of 1-day old adults the TSS of
62 genes are enriched for H3K4me3 in males compared to females, while the reverse is true in 10- to
63 12-day old adults. These sex differences at the TSS were not observed in *elav*-expressing neurons
64 (Palmateer, et al. 2021) supporting their role in directing sex-specific behaviors (Ito, et al. 1996;
65 Ryner, et al. 1996; Demir and Dickson 2005; Manoli, et al. 2005; Stockinger, et al. 2005; Goldman
66 and Arbeitman 2007; reviewed in Yamamoto and Koganezawa 2013).

67 Morphological and behavioral differences between males and females are common in sexually
68 reproducing organisms (Hedrick and Temeles 1989). *Drosophila* species have diverged in sexually
69 dimorphic morphology and behaviors (Sturtevant 1920; Ewing and Bennet-Clark 1968; Cobb, et
70 al. 1989; Chakir, et al. 2002; Kopp 2011; Arthur, et al. 2013). For example, there has been
71 relatively rapid diversification in reproductive behaviors such as the male courtship song (e.g.,
72 Ritchie, et al. 1999; Markow and O'Grady 2005; Latschev and Billeter 2014; Anholt, et al. 2020).
73 However, while there is strong evidence that links sex differences in expression with sex
74 dimorphism, the underlying mechanisms of species differences in sex dimorphism are poorly
75 understood. A study by Graze, et al. 2012 (Graze, et al. 2012) found that genes with *D. simulans*-
76 biased alleles in interspecific hybrids were enriched for genes associated with the GO term “H3K4
77 methyltransferase activity” and *D. melanogaster*-biased alleles to be enriched for genes associated
78 with the GO term “H3K9 methyltransferase activity”. H3K9 methylation is correlated with closed
79 chromatin and silenced expression (reviewed in Boros 2012; Kimura 2013), similar to H3K27

80 methylation. This finding led us to hypothesize that there may be divergence in chromatin patterns
81 between the species.

82 Sex-biased gene expression in brain, eye, and antennal genes has been shown to be associated with
83 sexually dimorphic behavior and sensory perception (Landry, et al. 2007; Kopp, et al. 2008; Shiao,
84 et al. 2015; reviewed in Anholt, et al. 2020). Sex-biased gene expression, or a gene with greater
85 expression in one sex over the other, has been previously shown to be rapidly evolving (e.g.,
86 Ellegren and Parsch 2007; Zhang, et al. 2007; Harrison, et al. 2015). Studying the evolution of
87 gene expression regulation within a relatively short period of evolutionary time, such as between
88 two closely related species, allows for identification of the sets of genes, mechanisms and
89 processes contributing to speciation and the evolution of species differences. *D. melanogaster* and
90 *D. simulans* have diverged relatively recently in evolutionary history (~5 million years ago)
91 (Tamura, et al. 2004) and *D. melanogaster* and *D. simulans* have diverged in many sexually
92 dimorphic phenotypes, including courtship behavior (Cobb, et al. 1989). Additionally, hybrid
93 studies of *D. melanogaster* and *D. simulans* have suggested divergence of sex-biased expression
94 regulatory mechanisms between the species (Ranz, et al. 2004). In combination with the wealth of
95 resources available for Drosophila as a model organism, the comparison of *D. melanogaster* and
96 *D. simulans* provides an exceptionally tractable model in which to explore the relationship between
97 chromatin marks and sex difference in expression, in an evolutionary context. To this end, we
98 assess the relationship between sex-biased expression and chromatin accessibility within each
99 species, as well as how this relationship evolves in two closely related species.

100 **Results**

101 We assayed males and females in the sister species, *D. melanogaster* and *D. simulans*, for gene
102 expression (n=48; 2 sexes x 2 genotypes x 2 species x 6 replicates), and chromatin (n=24; 2 sexes
103 x 1 genotypes x 2 species x 6 replicates). For each sample ChIP for the open chromatin mark,
104 H3K4me3, and closed chromatin marks, H3K27me2me3 and input were collected. We compared
105 the two sexes within each species, trends of sex-bias between species, and one-to-one orthologous
106 loci between species for gene expression and chromatin and evaluated the relationship between
107 sex-bias in gene expression and chromatin status. Within *D. melanogaster*, 2,556 genes on the X
108 chromosome and 14,114 genes on the autosomes were examined, and in *D. simulans*, 2,305 genes
109 on the X and 12,504 genes on the autosomes. There were 1,840 X and 10,097 autosomal one-to-
110 one orthologs used to compare species gene-to-gene. We performed extensive quality control of
111 the data (See Supplementary Materials Sections 4-7). For example, to evaluate whether genome
112 quality affected the results all analyses were also performed with both species mapped to *D.*
113 *melanogaster* (FlyBase r6.17) and both species mapped to *D. simulans* (FlyBase r2.02). While
114 there were a few genes with consistent map bias, there was no evidence that genome quality
115 impacted mapping (mapping rates were similar between species) and no trends reported were
116 affected by the choice to map each species to its own genome rather than mapping both to one of
117 the two genomes (Supplementary Materials Section 5.3).

118 Exonic regions were separated into non-overlapping exonic features where alternative
119 donor/acceptor sites were quantified separately from shared exonic regions, in order to capture the
120 potential sex-specific exonic features in each gene (Newman 2018). Non-overlapping exonic
121 features were quantified as $C_{is} = (\sum(d_{ijs})/N_i) \times (Q/U_s)$, where d is the depth of reads at
122 nucleotide j of feature i , N is the length of the feature, U_s is the upper quartile of $(\sum(d_{ijs})/N_i)$
123 values in sample s , and Q is the median of all U_s values within the given species (Bullard, et al.
124 2010; Dillies, et al. 2013).

125 If all detected features were detected in only one sex, the gene was labeled as sex-limited. There
126 were 770 genes (~6% of expressed genes) determined to be sex-limited in *D. melanogaster* (569
127 in males, 201 in females) and 547 genes (~4% of expressed genes) in *D. simulans* (352 in males,
128 195 in females) (Supplementary File 1, Supplementary File 2, *flag_sex_limited*=1). Differential
129 expression analyses were performed separately for each exonic feature detected in both sexes of
130 each species. Genes were considered sex-biased in expression if at least one exonic feature was
131 statistically significantly differentially expressed between sexes. Genes with both significantly
132 male- and female-biased exonic features were designated “Male-biased and Female-biased” and
133 are expected in genes that are sex-specifically alternatively spliced, such as the sex determination
134 gene *dsx* (Supplementary Figure 1).

135 ChIP samples were compared to input controls for genomic features (Transcription start sites, 5',
136 3' UTR's, exonic features and introns). Genomic features were considered detected above the input
137 control in H3K4me3 (DAI) if $C_{K4,is} > C_{Input,is}$, in more than 50% of the replicates for that species-
138 sex combination, and as $C_{K27,is} > C_{Input,is}$, for H3K27me3me4. ChIP data were found to be high
139 quality and conform with general expectations for detection of the marks (Supplementary
140 Materials Sections 7.1-7.2). A gene was considered as having a mark if at least one exonic feature
141 in the gene was DAI. A gene was considered sex-limited (male/female) when marks were detected
142 in only one sex.

143 **Genes that diverge in sex-bias between the species are more likely to be male-biased than 144 female-biased**

145 Gene expression in head tissues was measured in independent replicates of males and females for
146 each species ($n = 48$, 2 species x 2 sexes x 2 genotypes x 6 replicates). In addition to the excess of
147 male-limited expression compared to female limited expression, there is an excess of male-biased
148 expression compared to female-biased expression observed in both *D. melanogaster* (2723 male-
149 biased vs. 2185 female-biased, Binomial $p < 0.0001$) and *D. simulans* (2160 male-biased vs. 1873
150 female-biased, Binomial $p < 0.0001$) is statistically significantly different in orthologous genes
151 where sex-bias is gained/lost between species (Table 1, rows 13 vs. 14 and 16 vs. 17,
152 Supplementary Figure 2C). Although the number of male biased orthologs was greater than the
153 number of female biased orthologs (Table 1, rows 1 vs. 2), there was no statistically significant
154 excess of male-bias based on our threshold of $p < 0.001$. For orthologs unbiased in *D. melanogaster*
155 and sex-biased in *D. simulans*, more genes were male-biased than female-biased (Binomial $p =$
156 0.0001, Table 1, Supplementary Figure 2C). Similarly, for orthologs unbiased in *D. simulans* and

157 sex-biased in *D. melanogaster*, we found an excess of male-biased expression compared to female-
158 biased expression (Binomial $p < 0.0001$, Table 1, Supplementary Figure 2C).


		<i>D. melanogaster</i>	<i>D. simulans</i>	Orthologs
1	Male-biased	2723	2160	1154
2	Female-biased	2185	1873	1038
3	Male- and Female-biased	142	100	10
4	Unbiased	6666	7410	3816
5	Reversal	Male	Female	113
6	Reversal	Female	Male	70
7	Gain/Loss	Male	Male and Female	42
8	Gain/Loss	Female	Male and Female	16
9	Gain/Loss	Male and Female	Male	31
10	Gain/Loss	Male and Female	Female	35
11	Gain/Loss	Male	Unbiased	1049
12	Gain/Loss	Female	Unbiased	872
13	Gain/Loss	Male and Female	Unbiased	53
14	Gain/Loss	Unbiased	Male	657
15	Gain/Loss	Unbiased	Female	525
16	Gain/Loss	Unbiased	Male and Female	27
17	Expressed	11716	11543	9508

$\left. \right\} p = 0.014$
 $\left. \right\} p < 0.0001$
 $\left. \right\} p \approx 0.0001$

159 **Table 1 – Sex bias in expression.** The observed pattern of sex-bias is listed in column 2, all
160 definitions are mutually exclusive. In rows 1-4 the number of genes following the pattern in
161 column 1 for each species are given in columns 3 and 4 and the number of orthologous that have
162 the same pattern in both species are in the right-most column. Larger numbers in *D. simulans* and
163 *D. melanogaster* columns reflect observations for genes for which no one-to-one ortholog was
164 identified or the one-to-one ortholog was not expressed in both sexes of both species. There are a
165 total of 5050 sex-biased genes in *D. melanogaster*, 4133 in *D. simulans*, and 2202 orthologs with
166 the same sex-bias observed in both species. Rows 5-17 are the observed patterns of sex bias where
167 the two species diverge. Binomial test probabilities are indicated to the right of the table for the
168 comparison of male-biased vs. female-biased for consistent and species-specific sex-biased gene
169 classifications. P-values are in black if below the significant threshold of $p = 0.001$ and gray if
170 above the threshold. Reversal of sex-bias is rare, only two percent (183 / 9508) of orthologs. Genes
171 on chromosome 4 and on scaffolds, as well as those that change location are omitted. Values of
172 the X and autosomes separately for each category are listed in Supplementary Table 1.

173 **Male-biased orthologs are less constrained than female-biased orthologs**

174 The estimated sex-ratio in orthologs is strikingly concordant for females ($\beta_{1f} = 0.6428$; Pearson's
175 $r = 0.69$; T-test $H_0: \beta_1 = 0, p < 0.0001$) and males ($\beta_1 = 0.4663$; Pearson's $r = 0.49$; T-test
176 $H_0: \beta_{1m} = 0, p < 0.0001$). Male-biased orthologs, are less concordant than female-biased
177 orthologs ($H_0: \beta_{1m} = \beta_{1f}, p < 0.0001$) (Figure 1A). In addition, since the regression coefficients
178 of both male-biased orthologs and female-biased orthologs are significantly less than 1 (T-test
179 $H_0: \beta_1 \geq 1, p < 0.0001$ for both male-biased and female-biased orthologs), there is evidence for a
180 larger sex-bias in *D. melanogaster* compared to *D. simulans*. Orthologs with gains/losses in sex-
181 bias show large variation in the magnitude of the ratio of sex-bias compared to those with male or
182 female bias (Figure 1B).



183 **Figure 1 – Sex bias ratios across orthologs.** (Panel A) For the orthologous genes where sex bias
184 is in the same direction between the two species, we examined the relationship between the
185 observed ratio of sex bias in *D. melanogaster* (X-axis) and *D. simulans* (Y-axis). The $Y=X$ line is
186 in gray. To compare female-biased and male-biased genes on the same plot we calculated the sex-
187 bias ratio here as $(1 - \frac{\hat{f}}{\hat{m}})$ for male-biased orthologs (blue dots), and $(1 - \frac{\hat{m}}{\hat{f}})$ for female-biased
188 orthologs (red dots); where \hat{f} is the average UQ normalized expression across female samples and
189 \hat{m} is the average UQ normalized expression across male samples. A value close to 1 indicates
190 extreme sex-bias, while a value close to 0 indicates low sex-bias. A linear regression of the *D.*
191 *melanogaster* estimate on the *D. simulans* estimate was calculated for female-biased (red) or male-
192 biased (blue) orthologs separately. The ellipses represent the 95th percentile of the observed data.
193 (Panel B) Gains and losses in sex-bias. *D. melanogaster* sex-bias ratios (X-axis) compared to *D.*
194 *simulans* sex-bias ratios (Y-axis). In order to visually separate the male and female bias, we
195 calculated the sex-bias ratio as $(\frac{\hat{m}}{\hat{f}} - 1)$ if $\hat{m} > \hat{f}$, and as $(1 - \frac{\hat{f}}{\hat{m}})$ if $\hat{f} > \hat{m}$. Orthologs significant
196 for male-bias in one species are colored blue, and those significant for female-bias in one species
197 are colored red. The solid ellipses represent the 95th percentile of the observed statistically
198 significant species-specific sex bias in *D. melanogaster*. The dashed ellipses represent 95th
199 percentile of the observed statistically significant species-specific sex bias in *D. simulans*.

200 Orthologs with no significant sex-bias in either species are plotted in gray. Orthologs with reversal
201 of sex bias are potted in black (n=183, 2% of all orthologs).

202 **Male-bias in orthologs is associated with signatures of positive selection**

203 The comparative genomics database, flyDIVas (Clark, et al. 2007; Stanley and Kulathinal 2016)
204 provides gene-level estimates of divergence with nonsynonymous (dN) to synonymous
205 substitution (dS) rates (dN/dS) and tests of positive selection using PAML (Yang 1997) for the
206 melanogaster subgroup (*D. melanogaster*, *D. simulans*, *D. sechellia*, *D. yakuba*, *D. erecta*),
207 melanogaster group (melanogaster subgroup and *D. ananassae*), and the 12 Drosophila species
208 (melanogaster group and *D. pseudoobscura*, *D. persimilis*, *D. willistoni*, *D. mojavensis*, *D. virilis*,
209 and *D. grimshawi*). These different group allow for evaluation of selection across these three levels
210 of phylogenetic depth; however, the number of orthologous loci does decline as the distance from
211 melanogaster increases and the tests at the 12 genome level are to be thought of as suggestive
212 (Stanley and Kulathinal 2016). The null hypotheses tested are codon-based tests of positive
213 Darwinian selection based on dN/dS (ω) ratios estimated by PAML model M0 (Yang 1997). Three
214 nested pairs of site-specific models are available on flyDIVas: 1) model M1a (neutral) vs. M2a
215 (positive selection), 2) model M7 (beta-distributed) vs. M8 (beta+ ω >1) (Yang 1997), and 3) model
216 M8 (beta+ ω >1) vs. model M8a (beta+ ω =1) (Swanson, et al. 2003; Wong, et al. 2004). This is a
217 set of 9 tests of association (Supplementary Table 2). We used $p < 0.001$ as the significance
218 threshold and find that only the *D. melanogaster*-*D. simulans* male-biased orthologs are
219 significantly enriched for positive selection. This is a consistent inference for 8 of the 9 tests with
220 the exception being test M8 vs. M8a in the 12 species comparison (χ^2 : $p = 0.10$) (Supplementary
221 Table 2). There were 6 genes where reversals in sex bias occurred where individual genes showed
222 signatures of positive selection (*Exn*, *yin*, *SNF4A γ* , *Esyt2*, *DIP- η* , and *milt*).

223 **Sex-bias is enriched on the X chromosome**

224 Sex-biased expression in orthologs is enriched on the X chromosome compared to the autosomes
225 (Fisher's exact test: overall sex-biased $p < 0.0001$, male-biased $p < 0.0001$, female-biased $p <$
226 0.0001). No significant enrichment on the X is observed in the genes with gains/losses of sex
227 bias between the species according to our threshold of $p < 0.001$ (Supplementary Figure 3),
228 including orthologs sex-biased in *D. melanogaster* and unbiased in *D. simulans* (Fisher's exact
229 test: overall sex-biased $p = 0.002$, male-biased $p = 0.05$, female-biased $p = 0.03$) and orthologs
230 sex-biased in *D. simulans* and unbiased in *D. melanogaster* (Fisher's exact test: overall sex-
231 biased $p = 0.43$, male-biased $p = 0.42$, female-biased $p = 0.54$). Although, there were more
232 orthologous genes with sex-biased expression in *D. melanogaster* compared to *D. simulans*
233 (Table 1, rows 13-15 vs. 16-18; McNemar: $p < 0.0001$). These results are robust to map bias
234 (See Supplementary Materials Section 5.3).


235 **Genic chromatin marks are conserved**

236 The marginal frequencies for both open and closed chromatin differ from 0.5. In order to evaluate
237 the agreement in marks, we use a chance corrected measure of agreement (kappa, κ) to account

238 for these differences in the marginal frequencies (Fleiss 1981) (Supplementary Figure 4). A $\kappa = 1$
239 indicates perfect agreement, and negative values indicate less agreement than expected by chance
240 and a value of 0 indicates no agreement. Agreement between sexes and species is high for both
241 chromatin marks (Supplementary Table 3). Within each species, agreement between males and
242 females for the presence of H3K4me3 was high (*D. melanogaster* $\kappa = 0.73$, *D. simulans* $\kappa = 0.68$),
243 as well as for H3K27me2me3 (*D. melanogaster* $\kappa = 0.58$, *D. simulans* $\kappa = 0.54$). Additionally,
244 agreement between the species is high for H3K4me3 in males ($\kappa = 0.67$) and females ($\kappa = 0.73$)
245 and for H3K27me2me3 in males ($\kappa = 0.52$) and females ($\kappa = 0.54$). There is a set of 7,714 orthologs
246 (~65% of all one-to-one orthologs) with H3K4me3 marks present in both sexes and both species
247 and 1,817 (~16%) orthologs with H3K27me2me3 present in both sexes and both species. These
248 common marks make up ~76% of all genes with H3K4me3 and ~30% of genes with
249 H3K27me2me3. However, as expected, there are very few genes with both marks. The agreement
250 between genes with H3K4me3 and H3K27me2me3 is negative for both sexes and both species
251 indicating that these marks coincide less frequently than expected by chance (Supplementary Table
252 3).

253 Sex-limited chromatin accessibility diverges

254 We find there is more open chromatin in *D. simulans* compared to *D. melanogaster* (McNemar: p
255 < 0.0001) and more closed chromatin in *D. melanogaster* compared to *D. simulans* (McNemar: p
256 < 0.0001). While agreement between species is high in marks overall, there is low agreement when
257 marks are sex-limited (H3K4me3 κ : 0.05-0.30, H3K27me2me3 κ : 0.05-0.15, Supplementary
258 Table 3). However, there are more genes with conserved male-limited H3K4me3 than female-
259 limited H3K4me3 on the X (Binomial p < 0.0001, Figure 2B) and autosomes (Binomial p < 0.0001,
260 Figure 2C), while there are nearly equal numbers of genes with conserved male-limited and
261 female-limited H3K27me2me3 on both the X (Binomial p = 0.002, Figure 2B) and autosomes
262 (Binomial p = 0.83, Figure 2C). Interestingly, on the autosomes, sex-limited H3K4me3 shows
263 more genes with male-limited marks in *D. simulans* compared to *D. melanogaster* (McNemar: p
264 < 0.0001, Figure 2A) and female-limited marks are more prevalent in *D. melanogaster* compared
265 to *D. simulans* (McNemar: p < 0.0001, Figure 2A).



266 **Figure 2 – Chromatin marks in *D. melanogaster* and *D. simulans*.** The number of orthologs
 267 ($n=12,083$) with male-limited, female-limited, or marks in both sexes indicated in blue, red, and
 268 purple respectively. Most marks are detected in both sexes. Panel A) In *D. melanogaster*, 1,882
 269 and 10,121 genes are on the X and autosomes respectively, and 1,841 and 10,105 for *D. simulans*
 270 X and autosomes. There are 1,840 genes on the X of both species, 10,097 genes on the autosomes
 271 of both species, 7 genes on the X of *D. melanogaster* and autosomes of *D. simulans*, and 1 gene
 272 on the X of *D. simulans* and autosomes of *D. melanogaster*. The differences between the presence
 273 of marks in males compared to females was evaluated using McNemar test (McNemar 1947) with
 274 p-values for each test indicated in black for significant (< 0.001) and gray otherwise. Genes on the
 275 X (Panel B) and autosomes (Panel C) with conserved male-limited and female-limited chromatin
 276 marks (where both species are sex-limited in the same direction for a given chromatin mark in a
 277 gene) are indicated in blue or red for male-limited and female-limited respectively, and open box
 278 for the H3K4me3 and filled box for H3K27me2me3. P-values for binomial tests between the
 279 number of genes with male-limited and female-limited are indicated in black for significant ($<$
 280 0.001) and gray otherwise.

281 Chromosomal bias in chromatin


282 There is a higher proportion of genes with open chromatin marks detected in both species on the
 283 X chromosome compared to the autosomes (Supplementary Figure 5, Supplementary Figure

284 6D). We observe more male-limited than female-limited H3K4me3 on the X chromosome of *D.*
285 *simulans* (McNemar: $p < 0.0001$) and *D. melanogaster* (McNemar: $p < 0.0001$) and conserved
286 male-limited H3K4me3 marks are enriched on the X compared to the autosomes (χ^2 : $p < 0.0001$)
287 while conserved female-limited H3K4me3 marks have no chromosomal bias (Fisher exact: $p =$
288 0.08). Concomitantly, genes with conserved presence of female-limited H3K27me2me3 marks are
289 enriched on the X compared to the autosomes (χ^2 : $p = 0.0004$) and no chromosomal bias is
290 observed for male-limited H3K27me2me3 marks (χ^2 : $p = 0.35$).

291 **Sex-biased expression is associated with open chromatin**

292 We propose a model, “Open in Same sex and/or Closed in Opposite” (OS-CO), as an expectation
293 of chromatin accessibility states for genes with sex-biased expression (Figure 3). We expect
294 chromatin in male-biased genes to have i) open chromatin marks in males, and/or ii) closed
295 chromatin marks in females. We test this expectation by comparing the chromatin state in male-
296 biased genes to genes without male bias using Fisher exact test (Fisher 1934). Under the null
297 hypothesis that chromatin is independent of sex-bias there should be no difference in the
298 proportion of genes with open chromatin in males in these two groups. Similarly, we compare the
299 presence of open chromatin marks in females between female biased genes and non-female biased
300 genes. In both species, open chromatin marks in females are more likely to occur in female-biased
301 genes relative to non-female-biased genes (*D. melanogaster* χ^2 : $p < 0.0001$; *D. simulans* χ^2 : $p <$
302 0.0001; Figure 4A) and open chromatin marks in males are enriched in genes with male expression
303 bias compared to genes without male bias in expression (*D. melanogaster* χ^2 : $p < 0.0001$; *D.*
304 *simulans* χ^2 : $p < 0.0001$; Figure 4B).

305 When comparing the X and autosomes separately, female-biased genes showed the same pattern
306 of association with chromatin marks as in the combined set of genes across the genome (Figure
307 4C). Genes with male-biased expression were enriched for female closed chromatin on the
308 autosomes in both *D. melanogaster* and *D. simulans*, but on the X chromosome the chromatin
309 pattern was divergent between the two species. In *D. melanogaster* there was an enrichment for
310 male-biased expression with female closed chromatin on the X, whereas in *D. simulans* there was
311 not (Figure 4D). Outside of this divergence on the X, the expression and chromatin association
312 patterns are remarkably similar between the species on both the X and autosomes, with the most
313 striking differences observed between the sexes.



314 **Figure 3 – “Open in Same and/or Closed in Opposite” (OS-CO): a model for chromatin**
315 **accessibility patterns for sex-biased expression.** Representation of gene expression categories

316 between males and females on the X chromosome (X) and autosomes (A). Unbiased (X1, A1)

317 genes are defined as those without statistical evidence of differential expression. Female-biased

318 (X2, A2) genes are those with at least one exon with statistical evidence towards female expression.

319 Male-biased (X3, A3) genes are similarly defined towards male expression. Female-biased (X2,

320 A2) expression patterns are expected to have open chromatin marks (H3K4me3) in females and/or

321 closed chromatin marks (H3K27me2me3) in males. The mirror pattern is expected for male-biased



322 (X3, A3) expression patterns are expected to have open chromatin marks (H3K4me3) in males

323 and/or closed chromatin marks (H3K27me2me3). Not all sex-biased genes are expected to have



324 these patterns as there are other chromatin marks and regulatory factors that may influence

325 expression.



A.





B.



C.



D.



■ H3K4me3
● H3K27me2me3


326 **Figure 4 – Sex-biased expression is associated with chromatin marks.** The Y-axis of each graph
 327 represents the percent of expressed female-biased (solid red), non-female-biased (hatched red),
 328 male-biased (solid blue), or non-male-biased (hatched blue) genes within each species with the
 329 indicated chromatin (cartoon representations below each set of bars). Consistent with the model
 330 presented in Figure 4, (Panel A) Female-biased genes (solid red) are enriched for H3K4me3 (open)
 331 chromatin when compared to non-female-biased genes (hatched red) in both species. (Panel B)
 332 Male-biased genes (solid blue) are enriched for male open chromatin and female H3K27me2me3
 333 (closed) chromatin when compared to non-male-biased genes (hatched blue) in both species. The
 334 model in Figure 4 was also evaluated for X and autosomes separately. (Panel C) Female-biased
 335 genes (solid red) are enriched for open chromatin when compared to non-female-biased genes
 336 (hatched red) on both the X and autosomes of both species. (Panel D) Male-biased genes (solid
 337 blue) are enriched for male open chromatin and female closed chromatin when compared to non-
 338 male-biased genes (hatched blue) on both the X and autosomes of *D. melanogaster*. *D. simulans*
 339 shows the same pattern on the autosomes. On the X chromosome, male-bias genes are enriched
 340 for open chromatin in males but not for closed chromatin in females, showing a divergence in the
 341 regulatory pattern between the two species. There were 11,716 (nx=1,919, na=9,797) genes
 342 expressed in *D. melanogaster* and 9,902 genes expressed in *D. simulans* (nx=1,893, na=9,650)

343 evaluated for sex-biased expression and chromatin presence. Each set of female-biased (male-
344 biased) and non-female-biased (non-male-biased) genes were tested for enrichment of the
345 indicated chromatin mark using Fisher exact test (Fisher 1934) with the alternative expectation
346 that the indicated chromatin marks would be more likely in genes with female-biased (male-biased)
347 expression. Significant p-values ($p < 0.001$) are black and p-values above the significance
348 threshold are gray.

349 There are 34 genes on the X chromosome with male-biased expression in both species but female
350 closed chromatin only in *D. melanogaster* and not *D. simulans*. These genes are contributing to
351 the different patterns of chromatin mark usage observed on the X chromosomes of the two species
352 in Figure 4D. These 34 genes include the well described *D. melanogaster*-*D. simulans* hybrid
353 incompatibility gene, *Hmr* (Hutter and Ashburner 1987; Barbash, et al. 2003) that also has been
354 associated with heterochromatin factors (Satyaki, et al. 2014). Genes associated with habituation
355 [*wcy*, (Lugtenberg, et al. 2016)] and behavior [*Adar*, (Palladino, et al. 2000); *norpA*, (Pick and
356 Strauss 2005)] were also observed in this set of 34 genes. Further study of the *D. melanogaster*-
357 specific sex-biased genes with divergent chromatin regulation may reveal insights into sex-
358 dependent gene expression evolution and the role chromatin accessibility may play in the evolution
359 of these genes.

360 **Sex-biased orthologs have conserved presence of open chromatin**

361 In both species, the vast majority sex-biased orthologs have open chromatin in the sex with greater
362 expression (Figure 4, Supplementary Figure 7) consisted with our model (Figure 3). Male-biased
363 orthologs are significantly enriched for conserved open marks in males (~89% of male-biased
364 orthologs vs ~76% of unbiased orthologs; χ^2 : $p < 0.0001$). Similarly, female-biased orthologs are
365 significantly enriched for conserved open marks in females (~90% of female-biased orthologs vs
366 ~73% of unbiased orthologs; χ^2 : $p < 0.0001$). In addition, the agreement for H3K4me3 marks
367 within species in male-biased orthologs (*D. melanogaster* 0.63; *D. simulans* 0.60) and female
368 biased orthologs (*D. melanogaster* 0.65; *D. simulans* 0.60), is lower than for unbiased orthologs
369 (*D. melanogaster* 0.71; *D. simulans* 0.66). When male-biased orthologs have conserved H3K4me3
370 marks in males and no H3K27me2me3 mark in females, the sex-bias ratio is more similar (Figure
371 5A; $\beta_1 = 0.5023$) than when both species have a female H3K27me2me3 mark (Figure 5C; $\beta_1 =$
372 0.1795; β_{1A} vs. β_{1C} : $p < 0.0001$). When the female mark is not conserved between the species
373 (present in either species the sex-bias ratio is more conserved than when both species have the
374 female mark (Figure 5B; $\beta_1 = 0.4099$; β_{1A} vs. β_{1B} : $p \approx 0.1590$) (for all combinations of chromatin
375 marks for male-biased orthologs see Supplementary Figure 8). In contrast, the sex-ratio for female-
376 biased orthologs does not change with the male H3K27me2me3 chromatin mark (Figure 4, Figure
377 5D-F, Supplementary Figure 9).



378 **Figure 5 - Male and female biased orthologs.** Estimated ratio of sex-bias in *D. melanogaster*
379 (X-axis) and *D. simulans* (Y-axis), with the $y=x$ line in gray. To compare female-biased and male-
380 biased genes on the same scale, (0,1) we plotted $(1 - \frac{\hat{f}}{\hat{m}})$ for male-biased orthologs (blue dots),
381 where \hat{f} is average UQ normalized expression across female samples and \hat{m} is average UQ
382 normalized expression across male, and $(1 - \frac{\hat{m}}{\hat{f}})$ for female-biased orthologs (red dots). Genes are
383 separated by chromatin presence within the species. For male-biased orthologs: (Panel A)
384 conserved presence of male H3K4me3 and absence of female H3K27me2me3, (Panel B)
385 conserved presence of male H3K4me3 and female H3K27me2me3 in one species only, and (Panel
386 C) conserved presence of male H3K4me3 and female H3K27me2me3. For female-biased
387 orthologs: (Panel D) conserved presence of female H3K4me3 and absence of male H3K27me2me3,
388 (Panel E) conserved presence of female H3K4me3 and male H3K27me2me3 in one species only,
389 and (Panel F) conserved presence of female H3K4me3 and male H3K27me2me3. A red or blue
390 line indicates the linear regression calculated for the conserved female-biased or conserved male-
391 biased genes respectively using the least-squares method. Regression coefficients for each panel
392 are as follows: Panel A $\beta_1 = 0.5023$, Panel B $\beta_1 = 0.4099$, Panel C $\beta_1 = 0.1795$, Panel D $\beta_1 = 0.6021$,
393 Panel E $\beta_1 = 0.6507$, and Panel F $\beta_1 = 0.5935$.

394 Discussion

395 We propose a new model of how chromatin state is associated with sex-bias in expression. This
396 model hypothesizes that in genes with male-biased expression we expect to see an excess of open
397 chromatin in males compared to genes without male-bias; and in genes with female-biased
398 expression we expect to see an excess of open chromatin in females compared to genes without

399 female-bias. That is, we expect to observe open chromatin in the sex where we see expression bias
400 (OS). We also hypothesize that sex-bias in expression toward one sex may be associated with
401 closed chromatin in the opposite sex (CO). While, the open chromatin mark H3K4me3 is
402 correlated with active expression (Santos-Rosa, et al. 2002; Schneider, et al. 2004) and the closed
403 chromatin marks H3K27me2 and H3K27me3, together referred to as H3K27me2me3, are
404 correlated with silenced expression (Wang, et al. 2008; Juan, et al. 2016); these marks do not act
405 independently to affect chromatin accessibility. In *Drosophila* embryos, expression variation has
406 been found to be more predictive of the open chromatin mark H3K4me3 rather than the reverse
407 (Floc'hlay, et al. 2021), supporting the hypothesis that H3K4me3 does not induce transcription but
408 is instead deposited as a result of active transcription (reviewed in Howe, et al. 2017). Other histone
409 modifications, DNA methylation, and chromatin factors are involved in the establishment and
410 plasticity of chromatin accessibility (reviewed in Boros 2012). It is likely that our observations
411 using these marks does not completely reflect final active or repressed states of expression
412 resulting from the chromatin state as a whole, as we assayed only 2 of the many possible marks.
413 Our study does not demonstrate a causal relationship between chromatin accessibility and sex-
414 biased expression, nor do we claim to provide a comprehensive survey of chromatin accessibility.
415 Rather, our findings likely reflect the role of different regulators that impact chromatin states. Even
416 with broad limitations with respect to the suite of marks assessed, the OS component of the model
417 holds broadly. *Genes with sex-biased expression are more likely to have H3K4me2 marks in the*
418 *sex with greater expression in *D. melanogaster* and *D. simulans*, for both sexes, on both X and*
419 *autosomes compared to unbiased genes.*

420 The direction of sex-bias in expression agreed between the two species much more frequently than
421 expected by chance (male-bias: $\kappa = 0.41$, $p < 0.0001$; female bias: $\kappa = 0.45$, $p < 0.0001$). This
422 agreement in presence/absence of sex-bias between *D. melanogaster* and *D. simulans* may be due
423 to the short evolutionary time and the maintenance of the ancestral state where the sex-bias in the
424 common ancestor is random. Consistent with drift, the proportion of orthologs with male-bias is
425 not different from those with female-bias at our threshold ($p < 0.001$), although the number of
426 male biased orthologs is greater than the number of female biased orthologs. Under the null
427 hypothesis that the direction of bias is random, we would also expect to see approximately an even
428 number of gains/losses in transitions between the two species from unbiased to male- or female-
429 biased. In a binomial test, the null hypothesis of equal probability for male/female gain/loss ($p=0.5$)
430 is rejected for both transitions from unbiased genes in *D. melanogaster* to sex biased genes in *D.*
431 *simulans* (~55% male-biased, Binomial $p < 0.0001$) and unbiased genes in *D. simulans* to sex-
432 biased genes in *D. melanogaster* (~56% male-biased, Binomial $p \approx 0.0001$). There is also more
433 male-bias than female-bias in sex limited expression ($p < 0.0001$ for both species).

434 Sex-bias is conserved in magnitude, as well as direction. Intriguingly, sex-bias ratios for
435 expression are more similar between the species in females than males, suggesting there may be
436 either less constraint in males, or potentially a difference in selection between the sexes. While
437 there is no evidence in female-biased orthologs for the CO portion of our model, in male-biased
438 orthologs, the magnitude of sex-bias is affected by the presence of the female closed chromatin
439 marks providing some support for this hypothesis.

440 The excess of male-bias in sex-limited gene expression in both species, coupled with a significant
441 excess of male-bias in orthologs in the gain/loss of sex-bias, and less conservation in the magnitude
442 of the sex-bias ratio suggests that there is a possibility that the male-biased genes are evolving
443 faster. Male-biased genes have been shown to be evolving faster than other genes in comparisons
444 between *D. melanogaster* and *D. simulans* (Meiklejohn, et al. 2003) with overall higher rates of
445 evolution in male-biased genes observed in gonadal tissue (Perry, et al. 2014; Whittle and Extavour
446 2019) as well as whole body or somatic tissue (Ranz, et al. 2003; Zhang, et al. 2004; Connallon
447 and Knowles 2005; Ellegren and Parsch 2007).

448 In previous studies of *D. melanogaster*, head and brain tissues have been reported to have more
449 male-biased than female-biased expression (Chang, et al. 2011; Catalan, et al. 2012; Newell, et al.
450 2016; Palmateer, et al. 2021) with enrichment for male-biased genes on the X chromosome
451 compared to the autosomes (Goldman and Arbeitman 2007; Chang, et al. 2011; Catalan, et al.
452 2012; Meisel, et al. 2012a; Huylmans and Parsch 2015). Whole body tissue has been observed to
453 have more female-biased expression than male-biased expression (Ranz, et al. 2003; McIntyre, et
454 al. 2006; Wayne, et al. 2007; Graze, et al. 2014; Allen, et al. 2017) and enrichment for female-
455 biased genes on the X compared to the autosomes (Ranz, et al. 2003; McIntyre, et al. 2006; Wayne,
456 et al. 2007; Meisel, et al. 2012a; Graze, et al. 2014). We do find some evidence of positive selection
457 in male-biased orthologs. However, we cannot exclude the possibility that we observe an excess
458 in male-bias due to a relaxation of constraints in this specific tissue. The smaller slope in the
459 comparison between the species of the magnitude of sex-bias ratios in male-biased orthologs
460 compared to female-biased orthologs supports the relaxation of constraint hypothesis.

461 As terminal transcription factors of the sex determination pathway, *dsx* and *fru* have male- and
462 female-specific isoforms (Supplementary Figure 1). *Dsx* contributes to the regulation of sexual
463 dimorphism in the brain of both sexes (Rideout, et al. 2007; Kimura, et al. 2008; Rideout, et al.
464 2010; Arbeitman, et al. 2016), and is conserved among *Drosophila* species (Shukla and Nagaraju
465 2010). Although female-biased orthologs were not enriched for genes regulated by *dsx* (Arbeitman,
466 et al. 2016) ($\chi^2: p = 0.664$), male-biased orthologs were enriched for genes regulated by *dsx* ($\chi^2: p$
467 < 0.0001). *Fru*, is highly conserved in sex-specific splicing across insects (Salvemini, et al. 2010).
468 *Fru^M* is associated with chromatin remodeling factors (Lorbeck, et al. 2010; Ito, et al. 2012).
469 Additionally, the *fru* gene itself may be regulated by pheromone-induced chromatin remodeling at
470 the *fru* promotor in specific neurons (Zhao, et al. 2020) and *fru* expression decreases with mutation
471 in histone demethylase *Kdm4A* (Lorbeck, et al. 2010). Expression of *Fru^M* has been shown to affect
472 the establishment of closed chromatin marks in male neurons (Brovkina, et al. 2021) resulting in
473 the repression of genes that lead to sex-specific phenotypes (Ito, et al. 2016; Sato, et al. 2020;
474 reviewed in Goodwin and Hobert 2021). Palmateer et al. show overall differences in chromatin
475 patterns within genes enriched in *fru-P1* TRAP experiments (Palmateer, et al. 2021). Consistent
476 with the potential role of *Fru^M* as an activator of expression in males (Vernes 2014) is the excess
477 of open chromatin in males compared to female-limited open chromatin for both species in this
478 study on the X and for *D. simulans* on the autosomes. The male-specific *Fru^M* protein is a primary
479 regulator of sex dimorphism in the *Drosophila* brain (Ito, et al. 1996; Ryner, et al. 1996; Kido and
480 Ito 2002; Demir and Dickson 2005; Manoli, et al. 2005; Stockinger, et al. 2005; Rideout, et al.

481 2007; Kimura, et al. 2008; von Philipsborn, et al. 2011); and sex dimorphism has been shown to
482 be directed by *fru*. We hypothesized that the conservation in the male-specific Fru^M contributes to
483 conservation in male-biased expression. There were 1,771 and 729 genes identified as regulated
484 by *fru* in *D. melanogaster* males and females respectively (Dalton, et al. 2013). Male-biased
485 orthologs were enriched for genes regulated by the Fru^M protein in *D. melanogaster* males (χ^2 : *p*
486 < 0.0001) and female-biased orthologs were depleted for signatures of Fru^M (χ^2 : *p* = 0.002).

487 The faster-X hypothesis predicts that genes on the X chromosome evolve faster than those on the
488 autosomes (Haldane 1924b, a; Charlesworth, et al. 1987). When considering the unique properties
489 of the X, in combination with sex-differential effects of alleles, there is increased efficiency of
490 positive selection for X-linked alleles that are recessive and male-beneficial, or dominant and
491 female-beneficial (Wu and Davis 1993; Wu, et al. 1996). In the context of the evolution of sex-
492 biased genes, and in combination with other unique properties of the X, this may result in
493 preferential accumulation of sex-biased genes on the X chromosome over evolutionary time (Rice
494 1984; Charlesworth, et al. 1987; Oliver and Parisi 2004; Ellegren and Parsch 2007). In comparison,
495 the faster-male theory, a possible explanation of Haldane's rule (Haldane 1922; Wu and Davis
496 1993; Turelli and Orr 1995; reviewed in Schilthuizen, et al. 2011), predicts faster evolution of
497 genes related to male reproduction, regardless of location (reviewed in Schilthuizen, et al. 2011).
498 These are not mutually exclusive ideas. We observe an enrichment of genes with sex-biased
499 expression on the X chromosome compared to the autosomes in both species (Supplementary
500 Figure 2A). This is consistent with previous studies in the brain and is predicted by models of
501 sexually antagonistic evolution followed by gain of sex-biased or sex-limited expression (Rice
502 1984; Khodursky, et al. 2020).

503 Differences between the X chromosome and autosomes in the evolution of gene expression may
504 be due to changes in regulation of chromatin conformation associated with the X. Consistent with
505 this hypothesis, there were a higher proportions of genes with open chromatin marks detected in
506 both species on the X chromosome compared to the autosomes and conserved male-limited
507 H3K4me3 marks are enriched on the X compared to the autosomes.

508 Association between chromatin and male- and female-biased expression may be related to
509 evolutionary dynamics between the sexes. Sexual conflicts arise when the optima for a specific
510 trait differ between the sexes and therefore selection differs between the sexes. These conflicts can
511 come in two forms: interlocus and intralocus conflict (reviewed in Rice and Holland 1997;
512 Chapman, et al. 2003; Tregenza, et al. 2006; Bonduriansky and Chenoweth 2009; Cox and
513 Calsbeek 2009; Schenkel, et al. 2018). Intralocus conflict occurs when the optimal fitness of a
514 shared trait/locus is different between males and females, with different alleles favored in males
515 and females. It has been argued that the degree of observed sexual dimorphism can signify the
516 extent to which intralocus sexual conflict has been fully or partially resolved (Cox and Calsbeek
517 2009). In the whole fly, a small proportion (8.5%) of sex-biased genes have evidence of current
518 sexually antagonistic selection (Innocenti and Morrow 2010), indicating that in the majority of
519 cases, any sex-biased expression observed in this study that is associated with intralocus conflict
520 resolution would be expected to result from a history of partially or fully resolved intralocus

521 conflict, rather than ongoing intralocus conflict. We note that we find no association between the
522 fitness associated genes reported by Innocenti and Morrow (Innocenti and Morrow 2010) and the
523 observed conserved/diverged sex-biased orthologs reported here.

524 The findings that when female closed chromatin marks are absent in both species the male sex-
525 bias ratio is more similar between the species than when there is a mark in only one species,
526 suggests that the closed chromatin marks may play a role in resolving the ongoing sexual conflict
527 in males. The divergence in the degree of male-bias is associated with female H3K27me2me3
528 marks which are predicted to reduce expression in females. It is possible that this reflects a
529 mechanism of resolving cases of intralocus conflict in which expression of an allele in females has
530 deleterious effects. However, while male-biased genes on the autosomes show potential
531 suppression of expression in females, female-biased genes (both X and autosomal) in both species
532 lack the association of closed chromatin marks in males. This may suggest that genes with female
533 biased expression either i) don't involve deleterious effects in males, ii) involve genes that are
534 important for male fitness and are incompatible with closed marks and gene silencing, or iii) do
535 not involve resolution of intralocus sexual conflict. These overall patterns suggest specific testable
536 hypotheses regarding the role of activation and repression via chromatin modifications in the
537 resolution of intralocus sexual conflict for future experiments.

538 **Methods**

539

540 **Experimental Design**

541 Isogenic male and female *D. melanogaster* (DGRP r153 and r301) (Mackay, et al. 2012) and *D.*
542 *simulans* (Winters lines sz11 and sz12) (Signor 2017) flies were raised on standard Bloomington
543 recipe medium at 25C with a 12-h light/dark cycle. There were 2 sexes and 2 genotypes for each
544 species with 6 replicates for a total of 48 samples. Half of the samples were exposed to ethanol.
545 Samples were flash frozen in liquid nitrogen and freeze dried (Supplementary Figure 10).

546 For RNA-seq, 12 heads from each sample were collected. mRNA purification, cDNA synthesis
547 and dual index barcoding library preparation were carried out by Rapid Genomics (Gainesville,
548 FL, <http://rapid-genomics.com>). Individual libraries (n=48) were pooled in equimolar ratios as
549 estimated by Qubit and sequenced on a total of 7 Illumina lanes at Rapid Genomics (paired-end
550 2x100 3 lanes with HiSeq 3000 and paired-end 2x150 2 lanes with HiSeq X and 2 lanes with
551 NovaSeq 6000). External RNA Control Consortium (ERCC) spike-in control was used to evaluate
552 the quality of all RNA-seq sequencing libraries (Jiang, et al. 2011). After the first lane, read counts
553 of each library were used to confirm the pooling strategy.

554 For ChIP-seq, a target number of ~200 heads from each sample of *D. melanogaster* r301 and *D.*
555 *simulans* sz11 were collected (2 species x 6 replicates x 2 sexes x 1 genotype = 24 samples). Each
556 sample was used to assay histone marks H3K4me3 (open chromatin), H3K27me2me3 (closed
557 chromatin), and input. (3 antibodies/input x 24 samples = 72 assays). One r301 female untreated

558 sample contained ~175 heads and one 2 sz11 male ethanol treated sample contained ~120 heads,
559 and one sz11 ethanol treated female sample contained 50 heads. A full protocol for the ChIP
560 (Supplementary File 3, developed by NM and RR) is available in Supplementary File 1. ChIP
561 samples were indexed, pooled, and sequenced on one lane of an Illumina HiSeq2500 (paired-end
562 2x100) at the University of Florida, ICBR (Gainesville, FL, <https://biotech.ufl.edu/>).

563 **Genome Annotations**

564 All genome and annotation versions used were from FlyBase release FB2017_04
565 (<http://www.flybase.org>) *D. melanogaster* FlyBase r6.17 and *D. simulans* FlyBase r2.02. The
566 FlyBase gene OrthoDB ortholog report (Waterhouse, et al. 2013) (Supplementary File 4) was used
567 to identify one-to-one orthologous gene pairs (one gene in *D. melanogaster* associating with one
568 gene in *D. simulans*, and vice versa).

569 We created BED files for both genic features (exons, exonic features, TSS +/- 150 bp, 5' UTR,
570 3'UTR, and introns) and intergenic features (defined as the non-genic features greater than 50 bp
571 in length) for each reference from the relevant GFF annotation file. We note that in areas where
572 there were overlapping exons (where intron/exon boundaries vary by transcript), alternative donor
573 and acceptor sites were defined as exonic and tracked as separate features in downstream analyses
574 (Newman, et al. 2018). Counts of each unique feature type are in Supplementary Table 4. We note
575 that there are fewer genic features annotated in *D. simulans* compared to *D. melanogaster*.

576 **RNA-seq and ChIP-seq**

577 All results were consistent with reasonable quality data (Yang, et al. 2014) albeit with some shorter
578 sequences and higher duplication rates typically associated with libraries run on the NovaSeq 6000
579 in some of the RNA-seq runs.

580 Sequencing adapters were removed from both RNA-seq and ChIP-seq reads using Cutadapt
581 version 2.1 (Martin 2011) with a max error rate of 0.1 and a minimum overlap of 3 nt. Forward
582 and reverse reads were merged using BBMerge (Bushnell, et al. 2017). Reads less than 14bp + 50%
583 original read length were not considered further. Identical reads were identified (fastqSplitDups.py)
584 and removed. The resulting processed reads consisted of i) merged reads ('single-end'), ii)
585 unmerged reads without a proper pair ('single-end'), and iii) unmerged reads with proper pairs
586 (paired-end).

587 Processed RNA-seq reads and all ChIP reads were aligned to the corresponding genome reference
588 (*D. melanogaster* reads mapped to *D. melanogaster* FlyBase r6.17 and *D. simulans* reads mapped
589 to *D. simulans* FlyBase r.202) using BWA-MEM v0.7.15 (Li 2013) as single-end or paired-end
590 with default parameters. To determine if there was any systematic reference bias processed RNA-
591 seq reads from *D. melanogaster* samples were mapped to the *D. simulans* FlyBase r.202 genome,
592 and *D. simulans* samples were mapped to the *D. melanogaster* FlyBase r6.17 genome. A small

593 bias was observed towards mapping to the *D. simulans* genome and in both species, female
594 samples tended to have, on average, slightly higher mapping rates in the ChIP experiment.
595 Sensitivity to mapping bias was examined and results are described in detail in (Supplementary
596 Materials Section 5.3).

597 RNA-seq feature detection

598 A feature was considered detected by RNA-seq if at least one read was present in more than 50%
599 of the replicates for a species-sex combination (e.g., present in at least 7 of the 12 female or male
600 replicates for a given species). The number of detected features for each species-sex combination
601 is summarized in Supplementary Table 5. There are fewer features in *D. simulans* and despite the
602 slightly higher mapping rates found in *D. simulans* samples, there are slightly fewer features
603 detected in *D. simulans* samples compared to *D. melanogaster* across all feature types except for
604 3'UTR. The 3'UTR features has a higher proportion detection in *D. simulans* compared to *D.*
605 *melanogaster*, suggesting there may be a systematic bias in the 3'UTR regions of the two species
606 of either an over-annotation of these regions in *D. melanogaster* or an under-annotation in *D.*
607 *simulans*. There do not seem to be many missing genes in the *D. simulans* annotation because there
608 are not more detected features in *D. simulans* intronic and intergenic features compared to *D.*
609 *melanogaster*. In fact, there is a lower proportion of detected intronic and intergenic features in *D.*
610 *simulans* samples compared to *D. melanogaster* samples. Exonic feature detection was similar
611 between the species, with a slightly higher detection rates in *D. melanogaster* males. A feature was
612 considered sex-limited if the feature was detected in only one of the 2 sexes. Approximately 4%
613 of exonic features were sex-limited in *D. melanogaster* samples (2,530 in males, 1,195 in females)
614 and *D. simulans* (1,801 in males, 1,506 in females).

615 For the gene expression analysis, exonic regions were separated into non-overlapping exonic
616 features where alternative donor/acceptor sites were quantified separately from shared exonic
617 regions, in order to capture the potential sex-specific structures in the gene (Newman, et al. 2018).
618 Genes were defined as detected if at least one exonic feature was detected for either sex. There are
619 a similar number but proportionally more genes detected in *D. simulans* (11,543 out of 15,385,
620 ~75%) compared to *D. melanogaster* (11,716 out of 17,737, ~66%) indicating that there were no
621 large quality differences in the *D. simulans* genome compared to the *D. melanogaster* samples to
622 the *D. melanogaster* genome despite the differences in annotation.

623 To compare genes across *D. melanogaster* and *D. simulans*, we focus on annotated orthologs from
624 the OrthoDB ortholog report (Waterhouse, et al. 2013) to identify one-to-one orthologous gene
625 pairs (one gene in *D. melanogaster* associating with one gene in *D. simulans*, and vice versa)
626 (Supplementary File 4). There are 14,241 orthologous gene pairs between the species, 12,083 of
627 which are one-to-one orthologs. Genes on chromosome 4, the Y chromosome, and scaffolds of
628 either species were excluded from further analysis. There were 7 genes on the X chromosome of
629 *D. melanogaster* with orthologs on autosomes of *D. simulans*, and 1 gene on the X of *D. simulans*
630 with an ortholog on an autosome of *D. melanogaster*. These 8 genes were also excluded. The
631 remaining 11,937 one-to-one orthologous genes on the X (n = 1,840) and autosomes (n = 10,097)
632 of both species were carried forward.

633 RNA-seq Differential Expression

634 For each species, exonic features were quantified as $C_{is} = (\sum(d_{ijs})/N_i) \times (Q/U_s)$, where d is
635 the depth of reads at nucleotide j of feature i , N is the length of the feature, U_s is the upper quartile
636 of $(\sum(d_{ijs})/N_i)$ values in sample s , and Q is the median of all U_s values within the given species
637 (Bullard, et al. 2010; Dillies, et al. 2013) (Supplementary File 5). Distributions of upper quartile
638 values across exonic features were evaluated for each sample mapped to the genome of the sample
639 species (Supplementary Figure 11). Median upper quartile values and associated distributions were
640 strikingly similar across all samples in both species except for one *D. simulans* sz12 male replicate,
641 which was removed from further analysis.

642 For each species separately, differential expression between males and females was evaluated for
643 exonic features detected in both sexes. We used the linear fixed effect model $Y_{xp} = \mu + g_x + \varepsilon_{xp}$,
644 where Y is the log-transformed UQ normalized C_{is} values for the x th sex ($x = male, female$),
645 p th replicate ($p = 1, 2, \dots, 12$). We accounted for potential heteroscedasticity of variance between
646 the sexes (Graze, et al. 2012) and used the Kenward-Roger adjustment for the degrees of freedom
647 (Kenward and Roger 1997). Normality of residuals was tested using the Shapiro-Wilk test (Shapiro
648 and Wilk 1965). Fold-change ratios were calculated for each exonic feature i , $r_i =$
649 $(\sum(f_{ip})/k)/(\sum(m_{il})/n)$, where f_{ij} is the UQ normalized C_{is} for exonic region i in female
650 replicate $p = 1 \dots k$ total female replicates, and m_{il} is the UQ normalized C_{is} for exonic region i
651 in male replicate $l = 1 \dots n$ total male replicates. Exonic features were classified as male-biased
652 (or female-biased) if the nominal p-value was less than or equal to 0.05 and the fold-change less
653 than (or greater than) 1.

654 ChIP-seq Feature Detection

655 While peak calling is a common method of ChIP-seq analysis; it is highly dependent on the
656 algorithm used and the parameters selected (Yang, et al. 2014), especially for ChIP marks that are
657 predicted to show broad peaks such as certain histone modifications (Park 2009; Pepke, et al. 2009;
658 Dahl, et al. 2016). To have a consistent method for evaluation and comparison of ChIP results
659 across different marks and between males and females, and to compare ChIP results directly to the
660 RNA-sea results in *cis*, we use ChIP-seq reads to quantified features based on the annotations of
661 the reference genomes (Katz, et al. 2010; Anders, et al. 2012; Zhang, et al. 2012; Yang, et al. 2014;
662 Newman, et al. 2018). By focusing on features rather than MACS2 peaks, many more detections
663 above input control are identified at the feature-level and at the gene-level (See Supplementary
664 Methods Section 7.1 for detailed results from MACS2).

665 A feature was considered detected above the input control in H3K4me3/H3K27me3me4 (DAI) if
666 $C_{K4,is} > C_{Input,is}$, in more than 50% of the replicates for that species-sex combination A gene was
667 considered as having a mark if at least one exonic feature in the gene was DAI. A gene was
668 considered male-limited (or female-limited) if only sex-limited exonic features were identified in
669 both treatments. The agreement between histone marks for males and females, as well as between

670 H3K4me3 and H3K27me2me3 marks within each sex, was estimated using Cohen's kappa (Fleiss
671 1981) rather than simple agreement in order to account for marginal frequencies and provide a
672 more accurate assessment of the relationship between sexes and the marks (Supplementary Figure
673 4).

674 **Chromatin and expression**

675 Histone modifications change the availability of chromatin for transcription (Santos-Rosa, et al.
676 2002; Schneider, et al. 2004; Wang, et al. 2008; Juan, et al. 2016); therefore, we examine the
677 impact of chromatin marks on expression. When sex-biased expression is observed, this may be
678 due to open marks in the sex with the higher expression, or closed marks in the other sex.
679 Specifically, if there is male-biased expression, are there open (H3K4me3) marks in males or
680 closed (H3K27me2me3) marks in females for that gene, and if there is female-biased expression,
681 are there open (H3K4me3) marks in females or closed (H3K27me2me3) marks in males (Figure
682 3). As chromatin marks in males do not influence expression in females, or vice versa, the
683 appropriate statistical comparison is not a test of general association between expression and
684 chromatin marks between the sexes.

685 For males, the presence/absence of the chromatin marks, H3K4me3 and H3K27me2me3, was
686 compared to presence/absence of gene expression in males and evaluated for agreement using
687 Cohen's kappa coefficients (Fleiss 1981) (Supplementary Table 3). Females were examined
688 separately in the same manner. For genes with detected expression in both sexes, the
689 presence/absence of sex bias in males was compared to the presence/absence of male H3K4me3
690 marks using Fisher exact test (Fisher 1934) with the alternative expectation that male open
691 chromatin marks would be more likely in male-biased expression. For genes with sex-biased
692 expression in males, the presence/absence of H3K27me2me3 marks in females was tested using
693 Fisher exact test (Fisher 1934) with the alternative expectation that female closed chromatin marks
694 would be more likely in genes with male-biased expression. Tests were similarly performed for
695 the presence/absence of sex bias in females compared to the presence/absence of female H3K4me3
696 and presence/absence of male H3K27me2me3 using Fisher exact test (Fisher 1934).

697 **List enrichment**

698 Genes with sex-biased gene expression conserved between *D. melanogaster* and *D. simulans* in
699 this study were compared to genes identified in previous studies of sex-biased expression in *D.*
700 *melanogaster* head tissue (Chang, et al. 2011) using Pearson's Chi-square (χ^2) test (Pearson 1900).
701 Additionally, conserved male-biased (or female-biased) genes were compared to genes previously
702 identified as male-biased (or female-biased) in *D. melanogaster* head tissue and in *fru-P1*-
703 expressing neurons (Newell, et al. 2016) using Pearson's Chi-square (χ^2) test (Pearson 1900).
704 Based on the extensive knowledge of the sex-specifically spliced Drosophila sex determination
705 gene *fru* (Ryner, et al. 1996; Heinrichs, et al. 1998; reviewed in Salvemini, et al. 2010), we
706 expected *fru* to play a role in conserved sex-biased expression. Genes with male-biased and

707 female-biased expression conserved between *D. melanogaster* and *D. simulans* in this study were
708 compared to genes regulated by the Fru^M protein in *D. melanogaster* males (Dalton, et al. 2013)
709 using Pearson's Chi-square (χ^2) test (Pearson 1900).

710 Divergence of the targets of the terminal sex determination genes may contribute to the divergence
711 of sex-biased expression between the species. To evaluate this, species-specific sex-biased genes
712 identified in this study were compared to genes in a study of *dsx* regulation in *dsx* null females
713 and *dsx* pseudomales of *D. melanogaster* (Arbeitman, et al. 2016) and to genes observed to be
714 regulated downstream of *fru* in *D. melanogaster* males (Dalton, et al. 2013) using Pearson's Chi-
715 square (χ^2) test (Pearson 1900). To validate the patterns of open and closed chromatin in males and
716 females, gene-level presence of open (H3K4me3) and closed (H3K27me2me3) chromatin marks
717 in *D. melanogaster* males and females found in this study were compared to previous observations
718 of H3K4me3 and H3K27me3 marks in *D. melanogaster* male and female (*elav*-expressing)
719 neurons (Palmateer, et al. 2021) using Pearson's Chi-square (χ^2) test (Pearson 1900). Tests of
720 agreement between these datasets were carried out for males and females separately using Cohen's
721 kappa coefficients (Fleiss 1981) (Supplementary Table 3).

722 To evaluate if the patterns of the chromatin marks in the head tissue described here are consistent
723 with patterns of chromatin marks in neurons known to direct male and female reproductive
724 behaviors (Demir and Dickson 2005; Manoli, et al. 2005; Stockinger, et al. 2005; Kvitsiani and
725 Dickson 2006), the genes we detected with open (or closed) chromatin marks were compared to
726 genes with H3K4me3 (or H3K27me3) marks in *D. melanogaster* male and female *fru-P1*-
727 expressing neurons (Palmateer, et al. 2021) using Pearson's Chi-square (χ^2) test (Pearson 1900).
728 We also compared genes with male-limited and female-limited open (or closed) chromatin to the
729 genes with H3K4me3 (or H3K27me3) marks in *D. melanogaster* male and female *fru-P1*-
730 expressing neurons (Palmateer, et al. 2021) using Pearson's Chi-square (χ^2) test (Pearson 1900).
731 Tests of agreement of the comparable marks between head tissue and *fru-P1*-expressing neurons
732 were also evaluated for males and females separately using Cohen's kappa coefficients (Fleiss
733 1981) (Supplementary Table 3).

734 **Data availability**

735 Raw short-read data from the RNA-seq and ChIP-seq experiments are available under SRA
736 BioProject accession PRJNA737411. RNA-seq and ChIP-seq mapped read count summary
737 (Supplementary Table 6) and RNA-seq UQ normalization factors (Supplementary File 5) are
738 provided in the supplement. Analyzed data are provided as supplementary files for i) *D. melanogaster*
739 gene-level chromatin and expression variables (Supplementary File 1), ii) *D. melanogaster*
740 feature-level level chromatin and expression variables (Supplementary File 6), iii)
741 *D. simulans* gene-level chromatin and expression variables (Supplementary File 2), iv) *D. simulans*
742 feature-level chromatin and expression variables (Supplementary File 7), and v) *D. melanogaster*-
743 *D. simulans* orthologous gene chromatin and expression variables (Supplementary File 8). Further
744 detail of methods can be found in Supplementary Materials and documentation of all analyses and

745 comparisons as well as scripts are on github (https://github.com/McIntyre-Lab/papers/tree/master/nanni_chip_rna_2022).

747 **Acknowledgments**

748 Michelle Arbeitman for helpful discussion and sharing of pre-publication results. HiPerGator high-
749 performance supercomputer at the University of Florida. The University of Florida Genetics
750 Institute. Funding provided by National Institute of General Medical Sciences R01GM128193
751 (LMM, SVN) and National Institute of Dental and Craniofacial Research R01DE026707 (RR).

752

753 **References**

754 Anders S, Reyes A, Huber W. 2012. Detecting differential usage of exons from RNA-seq data.
755 Genome Research 22:2008-2017.

756 Anholt RRH, O'Grady P, Wolfner MF, Harbison ST. 2020. Evolution of Reproductive Behavior.
757 Genetics 214:49-73.

758 Anreiter I, Kramer JM, Sokolowski MB. 2017. Epigenetic mechanisms modulate differences in
759 Drosophila foraging behavior. Proceedings of the National Academy of Sciences of the United
760 States of America 114:12518-12523.

761 Anreiter I, Sokolowski MB. 2019. The *foraging* Gene and Its Behavioral Effects: Pleiotropy and
762 Plasticity. Annu Rev Genet 53:373-392.

763 Arbeitman MN, New FN, Fear JM, Howard TS, Dalton JE, Graze RM. 2016. Sex Differences in
764 Drosophila Somatic Gene Expression: Variation and Regulation by doublesex. G3-Genes
765 Genomes Genetics 6:1799-1808.

766 Arthur BJ, Sunayama-Morita T, Coen P, Murthy M, Stern DL. 2013. Multi-channel acoustic
767 recording and automated analysis of Drosophila courtship songs. Bmc Biology 11.

768 Barbash DA, Siino DF, Tarone AM, Roote J. 2003. A rapidly evolving MYB-related protein
769 causes species isolation in Drosophila. Proc Natl Acad Sci U S A 100:5302-5307.

770 Bludau A, Royer M, Meister G, Neumann ID, Menon R. 2019. Epigenetic Regulation of the Social
771 Brain. Trends in Neurosciences 42:471-484.

772 Bonduriansky R, Chenoweth SF. 2009. Intralocus sexual conflict. Trends in Ecology & Evolution
773 24:280-288.

774 Boros I. 2012. Histone modification in Drosophila. Briefings in Functional Genomics 11:319-331.

775 Brovkina MV, Duffie R, Burtis AEC, Clowney EJ. 2021. Fruitless decommissions regulatory
776 elements to implement cell-type-specific neuronal masculinization. Plos Genetics 17.

777 Brown EJ, Bachtrog D. 2014. The chromatin landscape of *Drosophila*: comparisons between
778 species, sexes, and chromosomes. *Genome Research* 24:1125-1137.

779 Brown EJ, Nguyen AH, Bachtrog D. 2020. The *Drosophila* Y Chromosome Affects
780 Heterochromatin Integrity Genome-Wide. *Molecular Biology and Evolution* 37:2808-2824.

781 Bullard JH, Purdom E, Hansen KD, Dudoit S. 2010. Evaluation of statistical methods for
782 normalization and differential expression in mRNA-Seq experiments. *Bmc Bioinformatics* 11.

783 Bushnell B, Rood J, Singer E. 2017. BBMerge - Accurate paired shotgun read merging via overlap.
784 *Plos One* 12.

785 Chakir M, Chafik A, Gibert P, David JR. 2002. Phenotypic plasticity of adult size and
786 pigmentation in *Drosophila*: thermosensitive periods during development in two sibling species.
787 *Journal of Thermal Biology* 27:61-70.

788 Chang P, Dunham J, Nuzhdin S, Arbeitman M. 2011. Somatic sex-specific transcriptome
789 differences in *Drosophila* revealed by whole transcriptome sequencing. *Bmc Genomics* 12.

790 Chapman T, Arnqvist G, Bangham J, Rowe L. 2003. Sexual conflict. *Trends in Ecology &*
791 *Evolution* 18:41-47.

792 Charlesworth B, Coyne JA, Barton NH. 1987. The relative rates of evolution of sex chromosomes
793 and autosomes. *American Naturalist* 130:113-146.

794 Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, Kaufman TC, Kellis M,
795 Gelbart W, Iyer VN, et al. 2007. Evolution of genes and genomes on the *Drosophila* phylogeny.
796 *Nature* 450:203-218.

797 Cobb M, Burnet B, Blizzard R, Jallon JM. 1989. Courtship in *Drosophila sechellia*: its structure,
798 functional aspects, and relationship to those of other members of the *Drosophila melanogaster*
799 species subgroup. *Journal of Insect Behavior* 2:63-89.

800 Connallon T, Knowles LL. 2005. Intergenomic conflict revealed by patterns of sex-biased gene
801 expression. *Trends in Genetics* 21:495-499.

802 Cox RM, Calsbeek R. 2009. Sexually Antagonistic Selection, Sexual Dimorphism, and the
803 Resolution of Intralocus Sexual Conflict. *American Naturalist* 173:176-187.

804 Dahl JA, Jung I, Aanes H, Greggains GD, Manaf A, Lerdrup M, Li GQ, Kuan S, Li B, Lee AY, et
805 al. 2016. Broad histone H3K4me3 domains in mouse oocytes modulate maternal-to-zygotic
806 transition. *Nature* 537:548-+.

807 Dalton JE, Fear JM, Knott S, Baker BS, McIntyre LM, Arbeitman MN. 2013. Male-specific
808 Fruitless isoforms have different regulatory roles conferred by distinct zinc finger DNA binding
809 domains. *Bmc Genomics* 14.

810 Demir E, Dickson BJ. 2005. fruitless splicing specifies male courtship behavior in *Drosophila*.
811 *Cell* 121:785-794.

812 Dillies MA, Rau A, Aubert J, Hennequet-Antier C, Jeanmougin M, Servant N, Keime C, Marot G,
813 Castel D, Estelle J, et al. 2013. A comprehensive evaluation of normalization methods for Illumina
814 high-throughput RNA sequencing data analysis. *Brief Bioinform* 14:671-683.

815 Ellegren H, Parsch J. 2007. The evolution of sex-biased genes and sex-biased gene expression.
816 *Nature Reviews Genetics* 8:689-698.

817 Elliott E, Manashirov S, Zwang R, Gil S, Tsoory M, Shemesh Y, Chen A. 2016. Dnmt3a in the
818 Medial Prefrontal Cortex Regulates Anxiety-Like Behavior in Adult Mice. *Journal of*
819 *Neuroscience* 36:730-740.

820 Ewing AW, Bennet-Clark HC. 1968. The Courtship Songs of *Drosophila*. In. p. 288-+.

821 Fisher RA. 1934. Statistical methods for research workers. Edinburgh: Oliver and Boyd.

822 Fleiss JL. 1981. Statistical methods for rates and proportions. Second edition: Wiley, John and
823 Sons, Inc., New York, N.Y.

824 Floc'hlay S, Wong ES, Zhao BQ, Viales RR, Thomas-Chollier M, Thieffry D, Garfield DA,
825 Furlong EEM. 2021. Cis-acting variation is common across regulatory layers but is often buffered
826 during embryonic development. *Genome Research* 31.

827 Francisco FO, Lemos B. 2014. How do y-chromosomes modulate genome-wide epigenetic States:
828 genome folding, chromatin sinks, and gene expression. *Journal of genomics* 2:94-103.

829 Goldman TD, Arbeitman MN. 2007. Genomic and functional studies of *Drosophila* sex hierarchy
830 regulated gene expression in adult head and nervous system tissues. *Plos Genetics* 3:2278-2295.

831 Goodwin SF, Hobert O. 2021. Molecular Mechanisms of Sexually Dimorphic Nervous System
832 Patterning in Flies and Worms. *Annual Review of Cell and Developmental Biology*, Vol 37
833 37:519-547.

834 Graze RM, Novelo LL, Amin V, Fear JM, Casella G, Nuzhdin SV, McIntyre LM. 2012. Allelic
835 Imbalance in *Drosophila* Hybrid Heads: Exons, Isoforms, and Evolution. *Molecular Biology and*
836 *Evolution* 29:1521-1532.

837 Haldane JBS. 1924a. A mathematical theory of natural and artificial selection. *Trans Phil Soc*
838 Cambridge U K 23:19-41.

839 Haldane JBS. 1924b. A mathematical theory of natural and artificial selection - Part II - The
840 influence of partial self-fertilisation, inbreeding, assortative mating, and selective fertilisation on
841 the composition of mendelian populations, and on natural selection. *Proceedings of the Cambridge*
842 *Philosophical Society-Biological Sciences* 1:158-163.

843 Haldane JBS. 1922. Sex ratio and unisexual sterility in hybrid animals. *Journal of Genetics* 12:101-
844 109.

845 Harrison PW, Wright AE, Zimmer F, Dean R, Montgomery SH, Pointer MA, Mank JE. 2015.
846 Sexual selection drives evolution and rapid turnover of male gene expression. *Proceedings of the*
847 *National Academy of Sciences of the United States of America* 112:4393-4398.

848 Hedrick AV, Temeles EJ. 1989. The evolution of sexual dimorphism in animals: Hypotheses and
849 tests. *Trends in Ecology & Evolution* 4:136-138.

850 Heinrichs V, Ryner LC, Baker BS. 1998. Regulation of sex-specific selection of fruitless 5' splice
851 sites by transformer and transformer-2. *Molecular and Cellular Biology* 18:450-458.

852 Henikoff S. 1996. Dosage-dependent modification of position-effect variegation in *Drosophila*.
853 *Bioessays* 18:401-409.

854 Howe FS, Fischl H, Murray SC, Mellor J. 2017. Is H3K4me3 instructive for transcription
855 activation. *Bioessays* 39.

856 Hutter P, Ashburner M. 1987. Genetic rescue of inviable hybrids between *Drosophila*
857 *melanogaster* and its sibling species. *Nature* 327:331-333.

858 Innocenti P, Morrow E. 2010. The Sexually Antagonistic Genes of *Drosophila melanogaster*. *Plos*
859 *Biology* 8.

860 Ito H, Fujitani K, Usui K, ShimizuNishikawa K, Tanaka S, Yamamoto D. 1996. Sexual orientation
861 in *Drosophila* is altered by the satori mutation in the sex-determination gene fruitless that encodes
862 a zinc finger protein with a BTB domain. *Proceedings of the National Academy of Sciences of the*
863 *United States of America* 93:9687-9692.

864 Ito H, Sato K, Koganezawa M, Ote M, Matsumoto K, Hama C, Yamamoto D. 2012. Fruitless
865 Recruits Two Antagonistic Chromatin Factors to Establish Single-Neuron Sexual Dimorphism.
866 *Cell* 149:1327-1338.

867 Ito H, Sato K, Kondo S, Ueda R, Yamamoto D. 2016. Fruitless Represses *robot* Transcription to
868 Shape Male-Specific Neural Morphology and Behavior in *Drosophila*. *Current Biology* 26:1532-
869 1542.

870 Jiang L, Schlesinger F, Davis CA, Zhang Y, Li R, Salit M, Gingeras TR, Oliver B. 2011. Synthetic
871 spike-in standards for RNA-seq experiments. *Genome Res* 21:1543-1551.

872 Juan AH, Wang S, Ko KD, Zare H, Tsai PF, Feng XS, Vivanco KO, Ascoli AM, Gutierrez-Cruz
873 G, Krebs J, et al. 2016. Roles of H3K27me2 and H3K27me3 Examined during Fate Specification
874 of Embryonic Stem Cells. *Cell Reports* 17:1369-1382.

875 Katz Y, Wang ET, Airoldi EM, Burge CB. 2010. Analysis and design of RNA sequencing
876 experiments for identifying isoform regulation. *Nature Methods* 7:1009-U1101.

877 Kenward MG, Roger JH. 1997. Small sample inference for fixed effects from restricted maximum
878 likelihood. *Biometrics* 53:983-997.

879 Khodursky S, Svetec N, Durkin SM, Zhao L. 2020. The evolution of sex-biased gene expression
880 in the *Drosophila* brain. *Genome Research* 30:874-884.

881 Kido A, Ito K. 2002. Mushroom bodies are not required for courtship behavior by normal and
882 sexually mosaic *Drosophila*. *Journal of Neurobiology* 52:302-311.

883 Kimura H. 2013. Histone modifications for human epigenome analysis. *Journal of Human
884 Genetics* 58:439-445.

885 Kimura K, Hachiya T, Koganezawa M, Tazawa T, Yamamoto D. 2008. *Fruitless* and *doublesex*
886 coordinate to generate male-specific neurons that can initiate courtship. *Neuron* 59:759-769.

887 Kopp A. 2011. *Drosophila* sex combs as a model of evolutionary innovations. *Evolution &
888 Development* 13:504-522.

889 Kopp A, Barmina O, Hamilton AM, Higgins L, McIntyre LM, Jones CD. 2008. Evolution of gene
890 expression in the *Drosophila* olfactory system. *Molecular Biology and Evolution* 25:1081-1092.

891 Kvitsiani D, Dickson BJ. 2006. Shared neural circuitry for female and male sexual behaviours in
892 *Drosophila*. *Current Biology* 16:R355-R356.

893 Landry CR, Castillo-Davis CI, Ogura A, Liu JS, Hartl DL. 2007. Systems-level analysis and
894 evolution of the phototransduction network in *Drosophila*. *Proceedings of the National Academy
895 of Sciences of the United States of America* 104:3283-3288.

896 Langley CH, Stevens K, Cardeno C, Lee YCG, Schrider DR, Pool JE, Langley SA, Suarez C,
897 Corbett-Detig RB, Kolaczkowski B, et al. 2012. Genomic Variation in Natural Populations of
898 *Drosophila melanogaster*. *Genetics* 192:533-.

899 Leturney M, Billeter JC. 2014. Neurogenetics of female reproductive behaviors in *Drosophila
900 melanogaster*. *Adv Genet* 85:1-108.

901 Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

902 Lorbeck MT, Singh N, Zervos A, Dhatta M, Lapchenko M, Yang C, Elefant F. 2010. The histone
903 demethylase *DmelKdm4A* controls genes required for life span and male-specific sex
904 determination in *Drosophila*. *Gene* 450:8-17.

905 Lugtenberg D, Reijnders MR, Fenckova M, Bijlsma EK, Bernier R, van Bon BW, Smeets E,
906 Vulfo-van Silfhout AT, Bosch D, Eichler EE, et al. 2016. *De novo* loss-of-function mutations in
907 WAC cause a recognizable intellectual disability syndrome and learning deficits in *Drosophila*.
908 *Eur J Hum Genet* 24:1145-1153.

909 Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, Casillas S, Han Y, Magwire
910 MM, Cridland JM, et al. 2012. The *Drosophila melanogaster* Genetic Reference Panel. *Nature*
911 482:173-178.

912 Manoli DS, Foss M, Villella A, Taylor BJ, Hall JC, Baker BS. 2005. Male-specific fruitless
913 specifies the neural substrates of *Drosophila* courtship behaviour. *Nature* 436:395-400.

914 Markow TA, O'Grady PM. 2005. Evolutionary genetics of reproductive behavior in *Drosophila*:
915 connecting the dots. *Annu Rev Genet* 39:263-291.

916 Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
917 2011 17:3.

918 McNemar Q. 1947. Note on the sampling error of the difference between correlated proportions
919 or percentages. *Psychometrika* 12:153-157.

920 Meiklejohn CD, Parsch J, Ranz JM, Hartl DL. 2003. Rapid evolution of male-biased gene
921 expression in *Drosophila*. *Proceedings of the National Academy of Sciences of the United States*
922 of America

923 Newell NR, New FN, Dalton JE, McIntyre LM, Arbeitman MN. 2016. Neurons That Underlie
924 *Drosophila melanogaster* Reproductive Behaviors: Detection of a Large Male-Bias in Gene
925 Expression in fruitless-Expressing Neurons. *G3 (Bethesda)* 6:2455-2465.

926 Newman J, Concannon P, Tardaguila M, Conesa A, McIntyre L. 2018. Event Analysis: Using
927 Transcript Events To Improve Estimates of Abundance in RNA-seq Data. *G3-Genes Genomes*
928 *Genetics* 8:2923-2940.

929 Oliver B, Parisi M. 2004. Battle of the Xs. *Bioessays* 26:543-548.

930 Opachaloemphan C, Yan H, Leibholz A, Desplan C, Reinberg D. 2018. Recent Advances in
931 Behavioral (Epi) Genetics in Eusocial Insects. *Annual Review of Genetics*, Vol 52 52:489-510.

932 Palladino MJ, Keegan LP, O'Connell MA, Reenan RA. 2000. A-to-I pre-mRNA editing in
933 *Drosophila* is primarily involved in adult nervous system function and integrity. *Cell* 102:437-449.

934 Palmateer CM, Moseley SC, Ray S, Brovero SG, Arbeitman MN. 2021. Analysis of cell-type-
935 specific chromatin modifications and gene expression in *Drosophila* neurons that direct
936 reproductive behavior. *Plos Genetics* 17.

937 Park PJ. 2009. ChIP-seq: advantages and challenges of a maturing technology. *Nature Reviews*
938 *Genetics* 10:669-680.

939 Pearson K. 1900. On the Criterion that a given System of Deviations from the Probable in the Case
940 of a Correlated System of Variables is such that it can be reasonably supposed to have arisen from
941 Random Sampling. *Philosophical Magazine* 50:157-175.

942 Pepke S, Wold B, Mortazavi A. 2009. Computation for ChIP-seq and RNA-seq studies. *Nature Methods* 6:S22-S32.

944 Perry JC, Harrison PW, Mank JE. 2014. The Ontogeny and Evolution of Sex-Biased Gene Expression in *Drosophila melanogaster*. *Molecular Biology and Evolution* 31:1206-1219.

946 Pick S, Strauss R. 2005. Goal-driven behavioral adaptations in gap-climbing *Drosophila*. *Curr Biol* 15:1473-1478.

948 Qin LY, Mal K, Wang ZJ, Hu ZH, Matas E, Wei J, Yan Z. 2018. Social deficits in Shank3-deficient mouse models of autism are rescued by histone deacetylase (HDAC) inhibition. *Nature Neuroscience* 21:564-+.

951 Ranz JM, Castillo-Davis CI, Meiklejohn CD, Hartl DL. 2003. Sex-dependent gene expression and evolution of the *Drosophila* transcriptome. *Science* 300:1742-1745.

953 Ranz JM, Namgyal K, Gibson G, Hartl DL. 2004. Anomalies in the expression profile of interspecific hybrids of *Drosophila melanogaster* and *Drosophila simulans*. *Genome Research* 14:373-379.

956 Rice WR. 1984. Sex chromosomes and the evolution of sexual dimorphism. *Evolution* 38:735-742.

958 Rice WR, Holland B. 1997. The enemies within: intergenomic conflict, interlocus contest evolution (ICE), and the intraspecific Red Queen. *Behavioral Ecology and Sociobiology* 41:1-10.

960 Rideout EJ, Billeter JC, Goodwin SF. 2007. The sex-determination genes *fruitless* and *doublesex* specify a neural substrate required for courtship song. *Current Biology* 17:1473-1478.

962 Rideout EJ, Dornan AJ, Neville MC, Eadie S, Goodwin SF. 2010. Control of sexual differentiation and behavior by the *doublesex* gene in *Drosophila melanogaster*. *Nature Neuroscience* 13:458-U479.

965 Ritchie MG, Halsey EJ, Gleason JM. 1999. *Drosophila* song as a species-specific mating signal and the behavioural importance of Kyriacou & Hall cycles in *D-melanogaster* song. *Animal Behaviour* 58:649-657.

968 Ryner LC, Goodwin SF, Castrillon DH, Anand A, Villella A, Baker BS, Hall JC, Taylor BJ, Wasserman SA. 1996. Control of male sexual behavior and sexual orientation in *Drosophila* by the *fruitless* gene. *Cell* 87:1079-1089.

971 Salvemini M, Polito C, Saccone G. 2010. *fruitless* alternative splicing and sex behaviour in insects: an ancient and unforgettable love story? *Journal of Genetics* 89:287-299.

973 Santos-Rosa H, Schneider R, Bannister AJ, Sherriff J, Bernstein BE, Emre NCT, Schreiber SL, Mellor J, Kouzarides T. 2002. Active genes are tri-methylated at K4 of histone H3. *Nature* 419:407-411.

976 Sato K, Ito H, Yamamoto D. 2020. *teiresias*, a Fruitless target gene encoding an immunoglobulin-
977 superfamily transmembrane protein, is required for neuronal feminization in *Drosophila*.
978 *Communications Biology* 3.

979 Satyaki PR, Cuykendall TN, Wei KH, Brideau NJ, Kwak H, Aruna S, Ferree PM, Ji S, Barbash
980 DA. 2014. The Hmr and Lhr hybrid incompatibility genes suppress a broad range of
981 heterochromatic repeats. *PLoS Genet* 10:e1004240.

982 Schenkel MA, Pen I, Beukeboom LW, Billeter JC. 2018. Making sense of intralocus and interlocus
983 sexual conflict. *Ecology and Evolution* 8:13035-13050.

984 Schilthuizen M, Giesbers M, Beukeboom LW. 2011. Haldane's rule in the 21st century. *Heredity*
985 107:95-102.

986 Schneider R, Bannister AJ, Myers FA, Thorne AW, Crane-Robinson C, Kouzarides T. 2004.
987 Histone H3 lysine 4 methylation patterns in higher eukaryotic genes. *Nature Cell Biology* 6:73-77.

988 Shapiro SS, Wilk MB. 1965. An analysis of variance test for normality (complete samples). In. p.
989 591-&.

990 Shiao MS, Chang JM, Fan WL, Lu MYJ, Notredame C, Fang S, Kondo R, Li WH. 2015.
991 Expression Divergence of Chemosensory Genes between *Drosophila sechellia* and Its Sibling
992 Species and Its Implications for Host Shift. *Genome Biology and Evolution* 7:2843-2858.

993 Shukla JN, Nagaraju J. 2010. Doublesex: a conserved downstream gene controlled by diverse
994 upstream regulators. *Journal of Genetics* 89:341-356.

995 Signor S. 2017. Population genomics of Wolbachia and mtDNA in *Drosophila simulans* from
996 California. *Sci Rep* 7:13369.

997 Spannhoff A, Kim YK, Raynal NJM, Gharibyan V, Su MB, Zhou YY, Li J, Castellano S, Sbardella
998 G, Issa JPJ, et al. 2011. Histone deacetylase inhibitor activity in royal jelly might facilitate caste
999 switching in bees. *Embo Reports* 12:238-243.

1000 Stanley CE, Kulathinal RJ. 2016. flyDIVaS: A Comparative Genomics Resource for *Drosophila*
1001 Divergence and Selection. *G3-Genes Genomes Genetics* 6:2355-2363.

1002 Stockinger P, Kvitsiani D, Rotkopf S, Tirian L, Dickson BJ. 2005. Neural circuitry that governs
1003 *Drosophila* male courtship behavior. *Cell* 121:795-807.

1004 Sturtevant AH. 1920. Genetic studies on *drosophila simulans*. I. Introduction. Hybrids with
1005 *Drosophila melanogaster*. *Genetics* 5:488-500.

1006 Sun HS, Damez-Werno DM, Scobie KN, Shao NY, Dias C, Rabkin J, Koo JW, Korb E, Bagot RC,
1007 Ahn FH, et al. 2015. ACF chromatin-remodeling complex mediates stress-induced depressive-like
1008 behavior. *Nature Medicine* 21:1146-+.

1009 Sun WJ, Poschmann J, del Rosario RCH, Parikshak NN, Hajan HS, Kumar V, Ramasamy R,
1010 Belgard TG, Elangovan B, Wong CCY, et al. 2016. Histone Acetylome-wide Association Study
1011 of Autism Spectrum Disorder. *Cell* 167:1385-+.

1012 Swanson WJ, Nielsen R, Yang Q. 2003. Pervasive adaptive evolution in mammalian fertilization
1013 proteins. *Mol Biol Evol* 20:18-20.

1014 Tamura K, Subramanian S, Kumar S. 2004. Temporal patterns of fruit fly (*Drosophila*) evolution
1015 revealed by mutation clocks. *Molecular Biology and Evolution* 21:36-44.

1016 Tregenza T, Wedell N, Chapman T. 2006. Introduction. Sexual conflict: a new paradigm?
1017 *Philosophical Transactions of the Royal Society B-Biological Sciences* 361:229-234.

1018 Turelli M, Orr HA. 1995. The dominance theory of Haldane's Rule. *Genetics* 140:389-402.

1019 Vernes SC. 2014. Genome wide identification of *Fruitless* targets suggests a role in upregulating
1020 genes important for neural circuit formation. *Scientific Reports* 4.

1021 von Philipsborn AC, Liu TX, Yu JY, Masser C, Bidaye SS, Dickson BJ. 2011. Neuronal Control
1022 of *Drosophila* Courtship Song. *Neuron* 69:509-522.

1023 Wang ZB, Zang CZ, Rosenfeld JA, Schones DE, Barski A, Cuddapah S, Cui KR, Roh TY, Peng
1024 WQ, Zhang MQ, et al. 2008. Combinatorial patterns of histone acetylations and methylations in
1025 the human genome. *Nature Genetics* 40:897-903.

1026 Waterhouse R, Tegenfeldt F, Li J, Zdobnov E, Kriventseva E. 2013. OrthoDB: a hierarchical
1027 catalog of animal, fungal and bacterial orthologs. *Nucleic Acids Research* 41:D358-D365.

1028 Whittle CA, Extavour CG. 2019. Selection shapes turnover and magnitude of sex-biased
1029 expression in *Drosophila* gonads. *Bmc Evolutionary Biology* 19.

1030 Wong WS, Yang Z, Goldman N, Nielsen R. 2004. Accuracy and power of statistical methods for
1031 detecting adaptive evolution in protein coding sequences and for identifying positively selected
1032 sites. *Genetics* 168:1041-1051.

1033 Wu CI, Davis AW. 1993. Evolution of Postmating Reproductive Isolation - The Composite Nature
1034 of Haldane's Rule and Its Genetic Bases. *American Naturalist* 142:187-212.

1035 Wu CI, Johnson NA, Palopoli MF. 1996. Haldane's rule and its legacy: Why are there so many
1036 sterile males? *Trends Ecol Evol* 11:281-284.

1037 Yamamoto D, Koganezawa M. 2013. Genes and circuits of courtship behaviour in *Drosophila*
1038 males. *Nature Reviews Neuroscience* 14:681-692.

1039 Yang Y, Fear J, Hu J, Haecker I, Zhou L, Renne R, Bloom D, McIntyre LM. 2014. Leveraging
1040 biological replicates to improve analysis in ChIP-seq experiments. *Computational and structural*
1041 *biotechnology journal* 9:e201401002-e201401002.

1042 Yang ZH. 1997. PAML: a program package for phylogenetic analysis by maximum likelihood.
1043 Computer Applications in the Biosciences 13:555-556.

1044 Zhang Y, Sturgill D, Parisi M, Kumar S, Oliver B. 2007. Constraint and turnover in sex-biased
1045 gene expression in the genus *Drosophila*. *Nature* 450:233-U232.

1046 Zhang YJ, Lameijer EW, t Hoen PAC, Ning ZM, Slagboom PE, Ye K. 2012. PASSion: a pattern
1047 growth algorithm-based pipeline for splice junction detection in paired-end RNA-Seq data.
1048 *Bioinformatics* 28:479-486.

1049 Zhang Z, Hambuch TM, Parsch J. 2004. Molecular evolution of sex-biased genes in *Drosophila*.
1050 *Molecular Biology and Evolution* 21:2130-2139.


1051 Zhao L, Begun DJ. 2017. Genomics of parallel adaptation at two timescales in *Drosophila*. *Plos*
1052 *Genetics* 13.

1053 Zhao SH, Deanhardt B, Barlow GT, Schleske PG, Rossi AM, Volka PC. 2020. Chromatin-based
1054 reprogramming of a courtship regulator by concurrent pheromone perception and hormone
1055 signaling. *Science Advances* 6.
1056


Supplementary Figures: pages 1-10

Supplementary Tables: pages 11-16


Supplementary Files: page 17



Supplementary Figure 1 – Drosophila sex determination hierarchy, XX females (A) and XY males (B) adapted from Figure 1 in Fear, et al. 2015. Transcripts are italicized and proteins are bold. Solid arrows are genetic interactions (e.g., splicing, transcription) and dashed arrows are protein translation. The X within a circle represents no productive protein product. (A1) Spf45 → Sxl (Lallena, et al. 2002), (A2) Snf → Sxl (Flickinger and Salz 1994), (A3) vir → Sxl (Hilfiker, et al. 1995), (A4) vir → tra (Hilfiker, et al. 1995), (A5) fl(2)d → Sxl (Granadino, et al. 1990), (A6) Sxl → Sxl (Cline 1978; Bell, et al. 1988; Lallena, et al. 2002), (A7) Sxl → Tra (Sosnowski, et al. 1989; Inoue, et al. 1990), (A8) Sxl → Msl-2 (Bashaw and Baker 1997; Kelley, et al. 1997; Gebauer, et al. 1998), (A9) fl(2)d → Tra (Granadino, et al. 1996), (A10) Tra → Dsx^F (Inoue, et al. 1992), (A11) Tra2 → Dsx^F (Inoue, et al. 1992), (A12) Dsx^F → Yps (Burtis, et al. 1991; Coschigano and Wensink 1993; An and Wensink 1995; Erdman, et al. 1996), (A13) Her → Yps (Li and Baker 1998), (A14) ix → Yps (Garrett-Engel, et al. 2002), (A15) Tra → Fru^M (Ryner, et al. 1996; Heinrichs, et al. 1998), (A16) Tra2 → FruM (Ryner, et al. 1996; Heinrichs, et al. 1998), (B1) default splicing of *sxl* transcripts results in no functional protein (Bell, et al. 1988), (B2) Msl-2 protein produced (Bashaw and Baker 1995; Kelley, et al. 1995; Zhou, et al. 1995), (B3) default splicing of *tra* transcripts results in no functional protein (Boggs, et al. 1987), (B4) Fru^M protein produced (Ryner, et al. 1996; Heinrichs, et al. 1998), (B5) default splicing of *dsx* transcripts in XY individuals results in Dsx^M protein (Burtis and Baker 1989), (B6) Dsx^M represses expression of Yps (Coschigano and Wensink 1993).



Supplementary Figure 2 – Excess of male-bias in *D. melanogaster* and *D. simulans* is due to divergent male-biased expression. (Panel A) Relative percent of expressed genes for *D. melanogaster* on the X (X, 1,919) and autosomes (A, 9,797) and for *D. simulans* on the X (X, 1,893) and autosomes (A, 9,650). (Panel B) Relative percent of orthologous genes expressed in *D. melanogaster* X (1,599) and autosomes (8,206) and *D. simulans* X (1,583) and autosomes (8,327). Chromosome 4 is excluded from the autosomes. Note that total numbers of orthologous genes differ between the species due to differences in chromosomal assignments. The number of genes in each category is printed over the box with male-bias (blue), female-bias (red), and both male- and female-biased (purple). Sex-limited genes (expressed in only one sex) are excluded. P-values for differences in male-biased and female-biased expression are reported. (Panel C) Conservation and divergence of sex-biased expression for expressed orthologous genes (n=9,508, with a consistent X/autosome chromosomal assignment between the species). Dots below the histogram are solid for the combination of factors reported as the number of genes in the bar plot.



Supplementary Figure 3 – X vs. autosomes of orthologs with conserved and divergent sex-biased expression. Expression of orthologous genes in the head for both sexes and both species on the X ($n_x=1,529$ genes) and autosomes ($n_a=7,979$). (Panel A) Genes that are conserved in their sex bias ($n_x=541$, $n_a=1,968$) are more likely to be on the X (35% on X vs. 24% on autosomes; χ^2 : $p < 0.0001$), while those that are unbiased are more likely to be on the autosomes (28% on X vs. 42% on autosomes; χ^2 : $p < 0.0001$). (Panel B) Genes divergent in sex bias ($n_x=558$, $n_a=2,625$) have no significant chromosomal bias for either *D. simulans*-specific sex-biased genes (12% on X vs. 12% on autosomes; χ^2 : $p = 0.86$) or *D. melanogaster*-specific sex-biased genes (23% on X vs. 20% on autosomes; χ^2 : $p = 0.003$). Connected black dots indicate the category plotted in the two bars above. The Y-axis is of the percentage of the total number of genes on the X or autosomes in each of the four categories. Chromosome 4 is excluded from the autosomes. X vs. autosome tests were performed using Pearson's Chi-square (χ^2) test (Pearson 1900) with a significance threshold of $p < 0.001$.

		Variable 1		Total
		0	1	
Variable 2	0	A_O	B_O	$A_O + B_O$
	1	C_O	D_O	$C_O + D_O$
Total	$A_O + C_O$	$B_O + D_O$	$A_O + B_O + C_O + D_O$	

$$\text{Simple agreement: } \frac{(A_O + D_O)}{(A_O + B_O + C_O + D_O)}$$


$$\text{Kappa: } k = \frac{\frac{(A_O + D_O)}{(A_O + B_O + C_O + D_O)} - (A_E + D_E)}{1 - (A_E + D_E)}$$

		Variable 1		Total
		0	1	
Variable 2	0	A_E	B_E	$A_E + B_E$
	1	C_E	D_E	

$$\text{Expected} = \frac{(\text{row total}) * (\text{column total})}{(\text{total observed})}$$

$$\text{e.g., } A_E = \frac{(A_O + B_O) * (A_O + C_O)}{(A_O + B_O + C_O + D_O)}$$


Supplementary Figure 4 – Measurements of Agreement. Given the table of observed values A_O , B_O , C_O , and D_O , the expected values indicated on the right can be calculated (A_E , B_E , C_E , and D_E). The formulas for calculating simple agreement and Cohen's Kappa agreement (Fleiss 1981) are also presented. Cohen's Kappa values correct for marginal frequencies, for when there is an imbalance between the variables tested.



Supplementary Figure 5 – Chromatin marks in males and females. (Panel A) Genes on the X or autosomes (denoted as A) of *D. melanogaster* FlyBase reference r6.17 ($n_X=2,556$; $n_A=14,114$; $n_X + n_A = 16,670$) and *D. simulans* FlyBase reference r2.02 ($n_X=2,305$; $n_A=12,504$; $n_X + n_A = 14,809$) with the number of genes with H3K4me3 (top) or H3K27me2me3 (bottom) male-limited, female-limited, or detected in both sexes indicated in blue, red, and purple respectively. Note that chromosome 4 is not included in the autosomes. (Panel B) Similar to Panel A, but with selecting for the one-to-one orthologs between *D. melanogaster* and *D. simulans* ($n=12,083$), excluding genes on chromosome 4 or unmapped scaffolds from further analysis (80 genes in *D. melanogaster* and 137 genes in *D. simulans*), resulting in 1,882 and 10,121 genes are on the *D. melanogaster* X and autosomes respectively, and 1,841 and 10,105 for *D. simulans* X and autosomes.


A.

Presence of H3K4me3 vs. H3K27me2me3




B.

Presence of sex-limited H3K4me3 vs. H3K27me2me3




C.

Presence of Marks X vs. A




D.

Sex-limited X vs. A




E.

Sex-limited *D. melanogaster* vs. *D. simulans*



F.


Male-limited vs. Female-limited




Supplementary Figure 6 –Tests for differential chromatin. H3K4me3 and H3K27me2me3 marks not present (black), present only in females (female-limited, red), present only in males (male-limited, blue), or present in both males and females (purple) on the X chromosome (X) and autosomes (A) for one-to-one orthologous genes of *D. melanogaster* and *D. simulans*. Chromosome 4 is excluded from the autosomes. The number of genes for each group is indicated. The total genes evaluated for are 1,882 and 10,121 for *D. melanogaster* X and autosomes respectively, and 1,841 and 10,105 for *D. simulans* X and autosomes. Tests are performed as follows. Panel A compares the presence of H3K4me3 vs. H3K27me2me3 marks in males/females within each species and chromosomal location. Panel B compares the presence of sex-limited H3K4me3 vs. H3K27me2me3 marks within each species and chromosomal location. Panels C-F compare within H3K4me3 or H3K27me2me3 marks separately. Panel C compares the presence of chromatin marks in either sex on the X vs. the autosomes within each species.

Panel D compares the presence of male-limited or female-limited marks on the X vs. the autosomes within each species. Panel E compares the proportion of male-limited or female-limited marks between *D. melanogaster* and *D. simulans* within each chromosomal location. Panel F compares male-limited vs. female-limited within each species and chromosomal location. All tests of X vs. autosomes are evaluated using Pearson's Chi-square (χ^2) test (Pearson 1900). Differences between species, sexes, and H3K4me3 vs. H3K27me2me3 are evaluated using McNemar's test of homogeneity (McNemar 1947).


A.





B.



C.




D.




Supplementary Figure 7 – Sex-biased expression is associated with chromatin marks in subset of orthologs. The Y-axis of each graph represents the percent of expressed female-biased (solid red), non-female-biased (hatched red), male-biased (solid blue), or non-male-biased (hatched blue) genes with a one-to-one ortholog within each species with the indicated chromatin (cartoon representations below each set of bars). Consistent with the model presented in Figure 4, (Panel A) Female-biased genes (solid red) are enriched for H3K4me3 (open) chromatin when compared to non-female-biased genes (hatched red) in both species. (Panel B) Male-biased genes (solid blue) are enriched for male open chromatin and female H3K27me2me3 (closed) chromatin when compared to non-male-biased genes (hatched blue) in both species. The model in Figure 4 was also evaluated for X and autosomes separately. (Panel C) Female-biased genes (solid red) are enriched for open chromatin when compared to non-female-biased genes (hatched red) on both the X and autosomes of both species. (Panel D) Male-biased genes (solid blue) are enriched for male open chromatin and female closed chromatin when compared to non-male-biased genes (hatched blue) on both the X and autosomes of both species.

(hatched blue) on both the X and autosomes of *D. melanogaster*. *D. simulans* shows the same pattern on the autosomes. On the X chromosome, male-bias genes are enriched for open chromatin in males but not for closed chromatin in females, showing a divergence in the regulatory pattern between the two species. There were 11,937 orthologous genes evaluated, 9,747 ($n_x=1,562$, $n_A=8,182$) genes expressed in *D. melanogaster* and Y genes expressed in *D. simulans* ($n_x=1,582$, $n_A=8,320$). Each set of female-biased (male-biased) and non-female-biased (non-male-biased) genes were tested for enrichment of the indicated chromatin mark using Fisher exact test (Fisher 1934) with the alternative expectation that the indicated chromatin marks would be more likely in genes with female-biased (male-biased) expression. Significant p-values ($p < 0.001$) are black and p-values above the significance threshold are gray.




Supplementary Figure 8. Plotted on the interval of (0,1) is the value $(1 - \frac{\hat{f}}{\hat{m}})$ for male-biased orthologs (blue dots), where \hat{f} is average UQ normalized expression across female samples and \hat{m} is average UQ normalized expression across male samples. *D. melanogaster* is on the X-axis and *D. simulans* on the Y axis. Each male-biased ortholog is plotted based on the sex bias ratio observed in each species, and placed in the box corresponding to the chromatin observed in *D. melanogaster* and *D. simulans*. Chromatin of *D.*

melanogaster is indicated at the top of each column of plots and chromatin of *D. simulans* is indicated at the left of each row of plots. Plots along the diagonal from the top left to the bottom right are genes where the observed chromatin is the same between the species. For each row (*D. simulans*) and column (*D. melanogaster*) the presence of H3K4me3 in males is indicated by a blue “YES” next to “Male K4” or a gray “NO” if it is not present. Similarly for the presence of H3K27me2me3 in females indicated by a blue “YES” next to “Female K27” if present, or a gray “NO” otherwise. Linear regression estimates are calculated for plots with at least 25 genes and plotted as a blue line.



Supplementary Figure 9. Plotted on the interval of $(0,1)$ is the value $(1 - \frac{\hat{m}}{\hat{f}})$ for female-biased orthologs (red dots), where \hat{f} is average UQ normalized expression across female samples and \hat{m} is average UQ normalized expression across male samples. *D. melanogaster* is on the X-axis and *D. simulans* on the Y axis. Each female-biased ortholog is plotted based on the sex bias ratio observed in each species, and placed in the box corresponding to the chromatin observed in *D. melanogaster* and *D. simulans*. Chromatin of *D. melanogaster* is indicated at the top of each column of plots and chromatin of *D. simulans* is indicated at the left of each row of plots. Plots along the diagonal from the top left to the bottom right are genes where the observed chromatin is the same between the species. For each row (*D. simulans*) and column (*D.*

melanogaster) the presence of H3K4me3 in females is indicated by a red “YES” next to “Female K4” or a gray “NO” if it is not present. Similarly for the presence of H3K27me2me3 in males indicated by a red “YES” next to “Male K27” if present, or a gray “NO” otherwise. Linear regression estimates are calculated for plots with at least 25 genes and plotted as a blue line.



Supplementary Figure 10 – Experimental Design. For RNA-seq there were a total of 48 samples (2 species x 2 genotypes x 2 sexes x 6 replicates). For ChIP-seq there were a total of 24 samples (2 species x 1 genotype x 2 sexes x 6 replicates) used for assaying chromatin (3 antibody/inputs per sample). Note that half of the replicates were exposed to ethanol (EtOH) and are included as additional data.



Supplementary Figure 11 – Distributions of RNA-seq expression values after UQ normalization. Upper quartile (UQ) normalization distributions per sample for (Panel A) *D. melanogaster* and (Panel B) *D. simulans* samples excluding the *D. simulans* sz12 male replicate that was removed due to a low median UQ relative to the rest of the samples.

Supplementary Tables

		<i>D. melanogaster</i> (X, A)	<i>D. simulans</i> (X, A)	Orthologs (X, A)
1	Male-biased	2723 (539, 2184)	2160 (433, 1727)	1154 (235, 919)
2	Female-biased	2185 (449, 1736)	1873 (398, 1475)	1038 (215, 823)
3	Male- and Female-biased	142 (38, 104)	100 (26, 74)	10 (2, 8)
4	Sex-biased	5050 (1026, 4024)	4133 (857, 3276)	2202 (452, 1750)
5	Unbiased	6666 (893, 5773)	7410 (1036, 6374)	3816 (430, 3386)
6	Switch			3490 (647, 2843)
7	Reversal	Male	Female	113 (37, 76)
8	Reversal	Female	Male	70 (17, 53)
9	Gain/Loss	Male	Male and Female	42 (12, 30)
10	Gain/Loss	Female	Male and Female	16 (4, 12)
11	Gain/Loss	Male and Female	Male	31 (11, 20)
12	Gain/Loss	Male and Female	Female	35 (8, 27)
13	Gain/Loss	Male	Unbiased	1049 (188, 861)
14	Gain/Loss	Female	Unbiased	872 (161, 711)
15	Gain/Loss	Male and Female	Unbiased	53 (12, 41)
16	Gain/Loss	Unbiased	Male	657 (108, 549)
17	Gain/Loss	Unbiased	Female	525 (84, 441)
18	Gain/Loss	Unbiased	Male and Female	27 (5, 22)
19	Expressed	11716 (1919, 9797)	11543 (1893, 9650)	9508 (1529, 7979)

Supplementary Table 1 – Number of genes showing different patterns of expression bias. The number of genes on the X and autosomes (excluding chromosome 4) for each pattern of expression bias for *D. melanogaster* and *D. simulans* head tissue (individual counts of X and autosomes are in parentheses). Sex-biased genes are the sum of male-biased (Male), female-biased (Female), and male- and female-biased (Male and Female) genes. Expression bias of orthologous of the species are indicated in the right-most column. Conserved expression bias, where both species are classified as the same category within the orthologous gene pair, are included in rows 1-5, followed by rows 6-18 with diverged expression bias, where the gene pair is assigned different expression categories between *D. melanogaster* and *D. simulans*. Binomial test probabilities are indicated to the right of the table for the comparison of male-biased vs. female-biased for conserved and species-specific sex-biased genes. Significant p-values are in black if below the significant threshold of $p = 0.001$ and gray if above the threshold.

Phylogeny	<i>D. melanogaster</i> Subgroup			<i>D. melanogaster</i> Group			12 Species			Total
Model	M1a vs. M2a	M7 vs. M8	M8 vs. M8a	M1a vs. M2a	M7 vs. M8	M8 vs. M8a	M1a vs. M2a	M7 vs. M8	M8 vs. M8a	
Conserved Male-biased Expression	66	83	4	44	67	47	13	104	20	1154
Conserved Male-biased Expression (Male H3K4me3 Both Species)	59	75	4	39	62	41	13	99	19	1027
Conserved Male-biased Expression (Male H3K4me3 <i>D. melanogaster</i> only)	2	2	0	2	1	2	0	1	1	26
Conserved Male-biased Expression (Male H3K4me3 <i>D. simulans</i> only)	5	5	0	3	3	3	0	4	0	62
Conserved Female-biased Expression	17	22	0	13	25	13	2	51	4	1038
Divergent Sex-biased Expression	96	128	6	62	105	68	10	223	23	3490
<i>D. melanogaster</i> -specific Sex-biased Expression	49	60	2	29	47	33	5	121	11	1974
<i>D. simulans</i> -specific Sex-biased Expression	29	43	2	18	39	21	3	72	8	1209
Reversal of Sex-biased Expression	6	9	2	5	7	5	0	14	2	183
Female-biased Expression in One Species	27	39	0	17	33	21	3	79	6	1397
Male-biased Expression in One Species	47	59	3	26	50	30	5	107	13	1706
Conserved Presence of Male H3K4me3	187	249	13	132	226	141	33	467	58	8462
Conserved Presence of Female H3K4me3	185	241	12	128	216	139	30	449	56	8022
Conserved Presence of Male H3K27me2me3	111	137	3	71	109	73	14	162	16	2687
Conserved Presence of Female H3K27me2me3	111	139	6	64	104	70	15	160	14	2762

Supplementary Table 2 – Enrichment of genes with positive selection. Summary of enrichment tests performed between genes with evidence of positive selection from flyDIVAs (Stanley and Kulathinal 2016; Clark 2007) and genes with conserved/diverged expression or conserved presence of chromatin marks described in this study. The number of genes with evidence of positive selection for the 3 phylogenetic levels (*D. melanogaster* subgroup, *D. melanogaster* group, and 12 species) and 3 models tested (M1a vs. M2a, M7 vs. M8, and M8 vs. M8a) in flyDIVAs is provided for each group. Gene numbers in red are those that were significantly enriched (χ^2 : $p < 0.001$) for genes with positive selection. More detailed descriptions of the models tested can be found in Table 2 of the PAML manual (<http://abacus.gene.ucl.ac.uk/software/pamlDOC.pdf>). Briefly, M1a vs. M2a compares nearly neutral evolution and positive selection, M7 vs. M8 compares where dN/dS (ω) varies according to a beta distribution vs. a beta distribution plus a discrete ω class where $\omega > 1$ (positive selection), and M8 vs. M8a which compares where ω varies according to a beta distribution with a discrete ω class where $\omega > 1$ vs. a beta distribution with $\omega=1$.

Comparison	Feature	Description	All	X	Autosomes
<i>D. melanogaster</i> vs. <i>D. simulans</i>	Gene	Male H3K4me3	0.67	0.65	0.67
		Female H3K4me3	0.73	0.75	0.72
		Male H3K27me2me3	0.52	0.45	0.54
		Female H3K27me2me3	0.54	0.55	0.54
		Male-limited H3K4me3	0.19	0.30	0.16
		Female-limited H3K4me3	0.07	0.05	0.07
		Male-limited H3K27me2me3	0.08	0.05	0.09
		Female-limited H3K27me2me3	0.09	0.15	0.08
	3' UTR	<i>D. melanogaster</i> H3K4me3	0.63	-	-
		<i>D. simulans</i> H3K4me3	0.54	-	-
		<i>D. melanogaster</i> H3K27me2me3	0.31	-	-
		<i>D. simulans</i> H3K27me2me3	0.27	-	-
<i>Male</i> vs. <i>Female</i>	5' UTR	<i>D. melanogaster</i> H3K4me3	0.72	-	-
		<i>D. simulans</i> H3K4me3	0.68	-	-
		<i>D. melanogaster</i> H3K27me2me3	0.29	-	-
		<i>D. simulans</i> H3K27me2me3	0.29	-	-
	Exon	<i>D. melanogaster</i> H3K4me3	0.63	-	-
		<i>D. simulans</i> H3K4me3	0.58	-	-
		<i>D. melanogaster</i> H3K27me2me3	0.34	-	-
		<i>D. simulans</i> H3K27me2me3	0.30	-	-
	Intron	<i>D. melanogaster</i> H3K4me3	0.58	-	-
		<i>D. simulans</i> H3K4me3	0.55	-	-
		<i>D. melanogaster</i> H3K27me2me3	0.36	-	-
		<i>D. simulans</i> H3K27me2me3	0.31	-	-
<i>H3K4me3</i> vs. <i>H3K27me2me3</i>	TSS (300bp Windows)	<i>D. melanogaster</i> H3K4me3	0.74	-	-
		<i>D. simulans</i> H3K4me3	0.68	-	-
		<i>D. melanogaster</i> H3K27me2me3	0.39	-	-
		<i>D. simulans</i> H3K27me2me3	0.30	-	-
	Intergenic	<i>D. melanogaster</i> H3K4me3	0.64	-	-
		<i>D. simulans</i> H3K4me3	0.69	-	-
		<i>D. melanogaster</i> H3K27me2me3	0.55	-	-
		<i>D. simulans</i> H3K27me2me3	0.59	-	-
	Gene	<i>D. melanogaster</i> H3K4me3	0.73	0.67	0.74
		<i>D. simulans</i> H3K4me3	0.68	0.63	0.69
		<i>D. melanogaster</i> H3K27me2me3	0.58	0.53	0.59
		<i>D. simulans</i> H3K27me2me3	0.54	0.49	0.54
<i>H3K4me3</i> vs. <i>H3K27me2me3</i>	3' UTR	<i>D. melanogaster</i> Males	-0.10	-	-
		<i>D. simulans</i> Males	-0.08	-	-
		<i>D. melanogaster</i> Females	-0.11	-	-
		<i>D. simulans</i> Females	-0.12	-	-
	5' UTR	<i>D. melanogaster</i> Males	-0.09	-	-
		<i>D. simulans</i> Males	-0.07	-	-

	<i>D. melanogaster</i> Females	-0.11	-	-
	<i>D. simulans</i> Females	-0.10	-	-
Exon	<i>D. melanogaster</i> Males	-0.16	-	-
	<i>D. simulans</i> Males	-0.10	-	-
	<i>D. melanogaster</i> Females	-0.19	-	-
	<i>D. simulans</i> Females	-0.14	-	-
	<i>D. melanogaster</i> Males	-0.15	-	-
Intron	<i>D. simulans</i> Males	-0.08	-	-
	<i>D. melanogaster</i> Females	-0.16	-	-
	<i>D. simulans</i> Females	-0.12	-	-
	<i>D. melanogaster</i> Males	-0.20	-	-
TSS (300bp Windows)	<i>D. simulans</i> Males	-0.10	-	-
	<i>D. melanogaster</i> Females	-0.22	-	-
	<i>D. simulans</i> Females	-0.14	-	-
	<i>D. melanogaster</i> Males	-0.31	-	-
Intergenic	<i>D. simulans</i> Males	-0.21	-	-
	<i>D. melanogaster</i> Females	-0.30	-	-
	<i>D. simulans</i> Females	-0.21	-	-
	<i>D. melanogaster</i> Males	-0.22	-0.11	-0.25
Genes	<i>D. simulans</i> Males	-0.12	-0.05	-0.13
	<i>D. melanogaster</i> Females	-0.26	-0.28	-0.26
	<i>D. simulans</i> Females	-0.22	-0.22	-0.22
	H3K4me3 in <i>D. melanogaster</i> Males	0.28	-	-
Head tissue vs. <i>elav</i> -expressing neurons	H3K4me3 in <i>D. melanogaster</i> Females	0.37	-	-
	H3K27me2me3 in <i>D. melanogaster</i> Males	0.26	-	-
	H3K27me2me3 in <i>D. melanogaster</i> Females	0.42	-	-
	H3K4me3 in <i>D. melanogaster</i> Males	0.27	-	-
Head tissue vs. <i>fru</i> - <i>P1</i> -expressing neurons	H3K4me3 in <i>D. melanogaster</i> Females	-0.04	-	-
	H3K27me2me3 in <i>D. melanogaster</i> Males	0.33	-	-
	H3K27me2me3 in <i>D. melanogaster</i> Females	0.32	-	-
	H3K4me3 in <i>D. melanogaster</i> Females	-	-	-

Supplementary Table 3 – Summary of Kappa values for the indicated comparisons.

Cohen's Kappa values (Fleiss 1981) indicating chance corrected agreement of the comparison described the “Comparison” column for the feature described in the “Feature” column and group in the “Description” column. Kappa values are presented for all chromosomes (X and autosomes combined) for all comparisons, as well as X chromosomes and autosomes separately for several indicated comparisons. Chromosome 4 is excluded from the autosomes.

Feature Type	<i>D. melanogaster</i>	<i>D. simulans</i>
Genes	17737	15385
Transcripts	35254	26261
TSS (300bp Windows)	22893	21069
5'UTR	28479	25081
3'UTR	21600	16231
Exonic Features	87473	79405
Intronic Features	44769	47236
Intergenic Features	11356	16174

Supplementary Table 4 – Number of annotated genomic features in *D. melanogaster* and *D. simulans*.

Counts of features within *D. melanogaster* and *D. simulans* genome annotation files. 5' UTR and 3'UTR were determined for each transcript using the references described in the Genome Annotation section of the Methods. A transcription start site (TSS) was defined as a 300 bp region, 150 bp upstream and downstream from each annotated transcript start. In *D. melanogaster* there were three pairs of genes where the members in each pair had the same start position but opposite strands: i) *bug* (FBgn0034050) and *Diap2* (FBgn0015247), ii) *lncRNA:CR44456* (FBgn0265649) and *lncRNA:CR44455* (FBgn0265648), and iii) *CR43482* (FBgn0263493) and *CR43483* (FBgn0263494). Event analysis (Newman, et al. 2018) was used to determine exonic and intronic features. Intergenic features were defined by subtracting the genic features from the entire genome with a length greater than 50 bp.

Species	Feature Type	# Detected in Males	# Detected in Females	# Detected in Either Sex	# Detected in Both Sexes
<i>D. melanogaster</i>	3UTR	14505 (67.15%)	14280 (66.11%)	14700 (68.06%)	14085 (65.21%)
	5UTR	18640 (65.45%)	18198 (63.9%)	19066 (66.95%)	17772 (62.4%)
	TSS	15761 (68.85%)	15323 (66.93%)	16161 (70.59%)	14923 (65.19%)
	Exonic	69373 (79.31%)	68038 (77.78%)	70568 (80.67%)	66843 (76.42%)
	Intergenic	6000 (52.84%)	5633 (49.6%)	6260 (55.13%)	5373 (47.31%)
	Intronic	29555 (66.02%)	28483 (63.62%)	30576 (68.3%)	27462 (61.34%)
<i>D. simulans</i>	3UTR	11820 (72.82%)	11717 (72.19%)	12032 (74.13%)	11505 (70.88%)
	5UTR	16108 (64.22%)	16054 (64.01%)	16653 (66.4%)	15509 (61.84%)
	TSS	13806 (65.53%)	13712 (65.08%)	14291 (67.83%)	13227 (62.78%)
	Exonic	61777 (77.8%)	61482 (77.43%)	63283 (79.7%)	59976 (75.53%)
	Intergenic	6769 (41.85%)	6616 (40.91%)	7144 (44.17%)	6241 (38.59%)
	Intronic	30010 (63.53%)	30013 (63.54%)	31531 (66.75%)	28492 (60.32%)

Supplementary Table 5 – Summary of features detected by RNA-seq. The number (percent) of features detected in males (irrespective of females), in females (irrespective of males), in either males or females (union), and in both males and females (intersection) for each species mapped to the associated reference

genome. Percent (in parentheses) is calculated by dividing the number detected by the total number for each feature type (see Supplementary Table 4 for feature totals).

A.								
Species	<i>D. melanogaster</i>							
Genome	<i>D. melanogaster</i> FB r6.17			<i>D. simulans</i> FB r2.02				
Sex	Male		Female	Male		Female		
Mean # mapped reads per replicate	16,368,252		16,479,280	16,877,562		16,915,308		
Mean % mapped reads per replicate	91.37%		92.76%	94.35%		95.22%		
Species								
Genome	<i>D. melanogaster</i> FB r6.17			<i>D. simulans</i> FB r2.02				
Sex	Male		Female	Male		Female		
Mean # mapped reads per replicate	14,774,793		15,598,854	15,498,853		16,423,578		
Mean % mapped reads per replicate	90.44%		94.61%	94.83%		94.61%		
B.								
Species	<i>D. melanogaster</i>							
Sex	Male			Female				
ChIP/Input	Input	H3K4me3	H3K27me2me3	Input	H3K4me3	H3K27me2me3		
Mean # mapped reads per replicate	10,915,497	13,688,173	12,666,807	12,239,391	13,817,377	14,380,575		
Mean % mapped reads per replicate	78.82%	86.08%	76.38%	81.83%	87.73%	78.70%		
Species								
Sex	Male			Female				
ChIP/Input	Input	H3K4me3	Input	H3K4me3	Input	H3K4me3		
Mean # mapped reads per replicate	10,435,104	14,728,518	10,435,104	14,728,518	10,435,104	14,728,518		
Mean % mapped read per replicate	80.25%	92.01%	80.25%	92.01%	80.25%	92.01%		

Supplementary Table 6 – Summary of read mapping counts and percentages. (A) RNA-seq mapped reads. All RNA-seq samples were mapped to both the *D. melanogaster* FlyBase 6.17 genome and the *D. simulans* FlyBase r.202 genome. The mean number and percent of processed reads across replicates after mapping to the indicated genome are given. (B) ChIP-seq mapped reads. All ChIP-seq samples were mapped to the associated reference genome based on the species of the sample (*D. melanogaster* FlyBase 6.17 genome or *D. simulans* FlyBase r.202). The mean number and percent of mapped processed reads across replicates for the indicated ChIP mark or input control are given.

Supplementary Files:

Supplementary File 1 - Gene-level expression and chromatin accessibility results for *D. melanogaster*. All column variables are defined in Supplementary File 9.

Supplementary File 2 – Gene-level expression and chromatin accessibility results for *D. simulans*. All column variables are defined in Supplementary File 9.

Supplementary File 3 – ChIP-seq protocol

Supplementary File 4 – Orthologous gene pairs of *D. melanogaster* to *D. simulans* selected from FlyBase OrthoDB report (Waterhouse, et al. 2013) in release 2017_04 (dmel_orthologs_in_drosophila_species_fb_2017_04.tsv.gz, downloaded 4/17/19). The original FlyBase file was modified to have individual columns for coordinates, +/- values for strand (compared to 1/-1), and “Dsim\” removed from Ortholog_GeneSymbol elements.

Supplementary File 5 – Upper quartile values used in for RNA-seq quantification.

Supplementary File 6 – Feature-level expression and chromatin accessibility results for *D. melanogaster*.

Supplementary File 7 – Feature-level expression and chromatin accessibility results for *D. simulans*.

Supplementary File 8 – Gene-level expression and chromatin accessibility results for *D. melanogaster* and *D. simulans* orthologs as identified by the FlyBase OrthoDB report (Waterhouse, et al. 2013). All column variables are defined in Supplementary File 10.

Supplementary File 9 – Gene-level variable definitions for species result files (Supplementary Files 1, 2).

Supplementary File 10 – Gene-level variable definitions for the *D. melanogaster* and *D. simulans* ortholog result file (Supplementary File 8).

Supplementary File 11 – For all gene numbers called out in the main text, the descriptions and the flags needed to identify those genes in Supplementary Files 1 or 8.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *Journal of Molecular Biology* 215:403-410.

Amrein H, Axel R. 1997. Genes expressed in neurons of adult male *Drosophila*. *Cell* 88:459-469.

An WQ, Wensink PC. 1995. Three protein binding sites form an enhancer that regulates sex-and fat body-specific transcription of *Drosophila* yolk protein genes. *Embo Journal* 14:1221-1230.

Bashaw GJ, Baker BS. 1995. The msl-2 dosage compensation gene of *Drosophila* encodes a putative DNA-binding protein whose expression is sex specifically regulated by Sex-lethal. *Development* 121:3245-3258.

Bashaw GJ, Baker BS. 1997. The regulation of the *Drosophila* msl-2 gene reveals a function for Sex-lethal in translational control. *Cell* 89:789-798.

Bell LR, Maine EM, Schedl P, Cline TW. 1988. Sex-lethal, a *Drosophila* sex determination switch gene, exhibits sex-specific RNA splicing and sequence similarity to RNA binding proteins. *Cell* 55:1037-1046.

Boggs RT, Gregor P, Idriss S, Belote JM, McKeown M. 1987. Regulation of sexual differentiation in *D. melanogaster* via alternative splicing of RNA from the transformer gene. *Cell* 50:739-747.

Burtis KC, Baker BS. 1989. *Drosophila doublesex* Gene Controls Somatic Sexual Differentiation by Producing Alternatively Spliced mRNAs Encoding Related Sex-Specific Polypeptides. *Cell* 56:997-1010.

Burtis KC, Coschigano KT, Baker BS, Wensink PC. 1991. The doublesex proteins of *Drosophila melanogaster* bind directly to a sex-specific yolk protein gene enhancer. *Embo Journal* 10:2577-2582.

Byron K. 2010. Mining roX1 RNA in *Drosophila* genomes using covariance models.

Cline TW. 1978. Two closely linked mutations in *Drosophila melanogaster* that are lethal to opposite sexes and interact with daughterless. *Genetics* 90:683-698.

Coschigano KT, Wensink PC. 1993. Sex-specific transcriptional regulation by the male and female doublesex proteins of *Drosophila*. *Genes & Development* 7:42-54.

Erdman SE, Chen HJ, Burtis KC. 1996. Functional and genetic characterization of the oligomerization and DNA binding properties of the *Drosophila* doublesex proteins. *Genetics* 144:1639-1652.

Fleiss JL. 1981. Statistical methods for rates and proportions. Second edition: Wiley, John and Sons, Inc., New York, N.Y.

Flickinger TW, Salz HK. 1994. The *Drosophila* sex determination gene *snf* encodes a nuclear protein with sequence and functional similarity to the mammalian U1A snRNP protein. *Genes & Development* 8:914-925.

Garrett-Engel CM, Siegal ML, Manoli DS, Williams BC, Li H, Baker BS. 2002. *intersex*, a gene required for female sexual development in *Drosophila*, is expressed in both sexes and functions together with *doublesex* to regulate terminal differentiation. *Development* 129:4661-4675.

Gebauer F, Merendino L, Hentze MW, Valcarcel J. 1998. The *Drosophila* splicing regulator *sex-lethal* directly inhibits translation of male-specific-lethal 2 mRNA. *Rna-a Publication of the Rna Society* 4:142-150.

Gorman M, Franke A, Baker BS. 1995. Molecular characterization of the male-specific lethal-3 gene and investigations of the regulation of dosage compensation in *Drosophila*. *Development* 121:463-475.

Granadino B, Campuzano S, Sanchez L. 1990. The *Drosophila melanogaster* *fl(2)d* gene is needed for the female-specific splicing of *Sex-lethal* RNA. *Embo Journal* 9:2597-2602.

Granadino B, Penalva LOF, Sanchez L. 1996. The gene *fl(2)d* is needed for the sex-specific splicing of transformer pre-mRNA but not for double-sex pre-mRNA in *Drosophila melanogaster*. *Molecular and General Genetics* 253:26-31.

Heinrichs V, Ryner LC, Baker BS. 1998. Regulation of sex-specific selection of *fruitless* 5' splice sites by transformer and transformer-2. *Molecular and Cellular Biology* 18:450-458.

Hilfiker A, Amrein H, Dubendorfer A, Schneiter R, Nothiger R. 1995. The gene *virilizer* is required for female-specific splicing controlled by *Sxl*, the master gene for sexual development in *Drosophila*. *Development* 121:4017-4026.

Inoue K, Hoshijima K, Higuchi I, Sakamoto H, Shimura Y. 1992. Binding of the *Drosophila* transformer and transformer-2 proteins to the regulatory elements of *doublesex* primary transcript for sex-specific RNA processing. *Proceedings of the National Academy of Sciences of the United States of America* 89:8092-8096.

Inoue K, Hoshijima K, Sakamoto H, Shimura Y. 1990. Binding of the *Drosophila* *Sex-lethal* gene product to the alternative splice site of transformer primary transcript. *Nature* 344:461-463.

Kelley RL, Solovyeva I, Lyman LM, Richman R, Solovyev V, Kuroda MI. 1995. Expression of *Msl-2* causes assembly of dosage compensation regulators on the X-chromosomes and female-lethality in *Drosophila*. *Cell* 81:867-877.

Kelley RL, Wang JW, Bell L, Kuroda MI. 1997. Sex lethal controls dosage compensation in *Drosophila* by a non-splicing mechanism. *Nature* 387:195-199.

Kuroda MI, Kernan MJ, Kreber R, Ganetzky B, Baker BS. 1991. The maleless Protein Associates with the X Chromosome to Regulate Dosage Compensation in *Drosophila*. *Cell* 66:935-947.

Lallena MJ, Chalmers KJ, Llamazares S, Lamond AI, Valcarcel J. 2002. Splicing regulation at the second catalytic step by sex-lethal involves 3' splice site recognition by SPF45. *Cell* 109:285-296.

Li H, Baker BS. 1998. hermaphrodite and doublesex function both dependently and independently to control various aspects of sexual differentiation in *Drosophila*. *Development* 125:2641-2651.

McNemar Q. 1947. Note on the sampling error of the difference between correlated proportions or percentages. *Psychometrika* 12:153-157.

Meller VH, Wu KH, Roman G, Kuroda MI, Davis RL. 1997. roX1 RNA paints the X chromosome of male *Drosophila* and is regulated by the dosage compensation system. *Cell* 88:445-457.

Newman J, Concannon P, Tardaguila M, Conesa A, McIntyre L. 2018. Event Analysis: Using Transcript Events To Improve Estimates of Abundance in RNA-seq Data. *G3-Genes Genomes Genetics* 8:2923-2940.

Palmateer CM, Moseley SC, Ray S, Brovero SG, Arbeitman MN. 2021. Analysis of cell-type-specific chromatin modifications and gene expression in *Drosophila* neurons that direct reproductive behavior. *Plos Genetics* 17.

Park SW, Kang YIe, Sypula JG, Choi J, Oh H, Park Y. 2007. An evolutionarily conserved domain of roX2 RNA is sufficient for induction of H4-Lys16 acetylation on the *Drosophila* X chromosome. *Genetics* 177:1429-1437.

Pearson K. 1900. On the Criterion that a given System of Deviations from the Probable in the Case of a Correlated System of Variables is such that it can be reasonably supposed to have arisen from Random Sampling. *Philosophical Magazine* 50:157-175.

Quinn JJ, Zhang QFC, Georgiev P, Ilik IA, Akhtar A, Change HY. 2016. Rapid evolutionary turnover underlies conserved lncRNA-genome interactions. *Genes & Development* 30:191-207.

Ryner LC, Goodwin SF, Castrillon DH, Anand A, Villella A, Baker BS, Hall JC, Taylor BJ, Wasserman SA. 1996. Control of male sexual behavior and sexual orientation in *Drosophila* by the fruitless gene. *Cell* 87:1079-1089.

Sosnowski BA, Belote JM, McKeown M. 1989. Sex-Specific Alternative Splicing of RNA from the transformer Gene Results from Sequence-Dependent Splice Site Blockage. *Cell* 58:449-459.

Waterhouse R, Tegenfeldt F, Li J, Zdobnov E, Kriventseva E. 2013. OrthoDB: a hierarchical catalog of animal, fungal and bacterial orthologs. *Nucleic Acids Research* 41:D358-D365.

Zhou SB, Yang YF, Scott MJ, Pannuti A, Fehr KC, Eisen A, Koonin EV, Fouts DL, Wrightsman R, Manning JE, et al. 1995. Male-specific lethal 2, a dosage compensation gene of *Drosophila*, undergoes sex-specific regulation and encodes a protein with a RING finger and a metallothionein-like cysteine cluster. *Embo Journal* 14:2884-2895.