

Feeding Neurons Integrate Metabolic and Reproductive States in Mice

Megan G. Massa¹, Rachel L. Scott¹, Alexandra L. Cara¹, Laura R. Cortes¹, Norma P. Sandoval¹, Jae W. Park¹, Sahara Ali¹, Leandro M. Velez³, Bethlehem Tesfaye¹, Karen Reue², J. Edward van Veen¹, Marcus Seldin³, and Stephanie M. Correa¹

¹Department of Integrative Biology and Physiology, University of California, Los Angeles, CA

²Department of Human Genetics, David Geffen School of Medicine at University of California, Los Angeles, CA

³Department of Biological Chemistry, School of Medicine, University of California, Irvine, CA

Abstract

Trade-offs between metabolic and reproductive processes are important for survival, particularly in mammals that gestate their young. Puberty and reproduction, as energetically taxing life stages, are often gated by metabolic availability in animals with ovaries. How the nervous system coordinates these trade-offs is an active area of study. We identify somatostatin neurons of the tuberal nucleus (TN^{SST}) as a node of the feeding circuit that alters feeding in a manner sensitive to metabolic and reproductive states in mice. Whereas chemogenetic activation of TN^{SST} neurons increased food intake across sexes, selective ablation decreased food intake only in female mice during proestrus. Interestingly, this ablation effect was only apparent in animals with a low body mass. Fat transplantation and bioinformatics analysis of TN^{SST} neuronal transcriptomes revealed white adipose as a key modulator of the effects of TN^{SST} neurons on food intake. Together, these studies point to a mechanism whereby TN^{SST} hypothalamic neurons modulate feeding by responding to varying levels of circulating estrogens differentially based on energy stores. This research provides insight into how neural circuits integrate reproductive and metabolic signals, and illustrates how gonadal steroid modulation of neuronal circuits can be context-dependent and gated by metabolic status.

Introduction

The homeostatic processes of metabolism and reproduction are mutually dependent on one another. Reproductive milestones, including pregnancy and quiescence, result in major metabolic shifts (Baz et al., 2016; Mauvais-Jarvis, 2017; Palmer and Clegg, 2015), and metabolic status can gate reproductive function in menstrual and estrual mammals. Pubertal onset has been shown to require a critical threshold of body fat (Frisch, 1972; Frisch and Revelle, 1970), and increasing adiposity is associated with a decreasing age of pubertal onset in individuals with ovaries in particular (Solorzano and McCartney, 2010). During reproductive years, undernutrition can acutely disrupt menstrual cyclicity (Nelson et al., 1985) via hypogonadotropic hypogonadism (Muñoz and Argente, 2002) and result in fewer successful pregnancies (Ball et al., 1947). Rodent models have been used to investigate the effects of sex variables on metabolic tissues. Estradiol has been shown to regulate key metabolic processes such as adiposity (Palmer and Clegg, 2015), thermogenic capacity & locomotion (Correa et al., 2015; Martínez de Morentin et al., 2014; Musatov et al., 2007; Palmer and Clegg, 2015; Xu et al., 2011), and feeding (Massa and Correa, 2020). Sex chromosome complement has been demonstrated to affect fat deposition and energy output (Chen et al., 2012). Few studies have investigated the mechanisms by which both metabolic and reproductive status reciprocally interact to modulate behavior.

A candidate region for the seat of this interaction is the mediobasal hypothalamus, a region typically thought to be comprised of the arcuate nucleus, ventromedial nucleus, and median eminence. Positioned near the third ventricle and partially outside of the blood brain barrier (Ciofi, 2011), the mediobasal hypothalamus is situated as a nexus region with the ability to sample circulating homeostatic hormones and relay relevant information to other regions of the brain. Importantly, this region is sensitive to and vital for both reproductive and metabolic homeostasis. For instance, in addition to the

role of kisspeptin neurons in the arcuate nucleus as key modulators of estrogen-mediated negative feedback in the hypothalamic-pituitary-ovarian axis (Mittelman-Smith et al., 2012; Smith et al., 2005), they also require epigenetic de-repression for menarche (Lomniczi and Ojeda, 2016; Wright et al., 2021). The arcuate nucleus is also home to the canonical homeostatic feeding neurons – appetitive agouti-related peptide (AgRP)/neuropeptide Y (NPY) neurons and satiety-related proopiomelanocortin (POMC) neurons – which can detect and respond to sensory and metabolic cues including leptin, ghrelin, and insulin (Burnett et al., 2016; Chen et al., 2015; Oldfield et al., 2016). The ventromedial nucleus of the hypothalamus, and in particular the ventrolateral subregion, is important for various functions across sexes, including mating behavior (reviewed in (Kammel and Correa, 2019)). Distinct, estrogen-sensitive cellular populations in the ventrolateral ventromedial nucleus contribute to various aspects of metabolism, including locomotion (Correa et al., 2015; Krause et al., 2021; Narita et al., 2016) and thermoregulation (van Veen & Kammel et al., 2020). Given the substantive neuronal heterogeneity within a small brain region, it is unsurprising that these populations form complex functional interactions. For example, simulation of starvation via chronic chemogenetic activation of appetitive arcuate AgRP neurons has been demonstrated to acutely disrupt estrous cyclicity (Padilla et al., 2017), providing a neuronal contributor to the link between metabolic state and reproductive function.

The tuberal nucleus (TN) straddles the mediobasal hypothalamus and lateral hypothalamic area. It is an understudied region marked by expression of the neuropeptide somatostatin (SST) which has been mostly characterized in rats. Studies have found the TN to be closely related to the ventrolateral region of the ventromedial hypothalamus (Canteras et al., 1994) (though in a recent mouse study, the TN was considered a part of the lateral hypothalamic area; Mickelsen et al., 2019), and suggest the region may also be estrogen sensing (Canteras et al., 1994; Pfaff and Keiner, 1973; Simerly et al., 1990). Recent studies investigating this region in mice have found the TN to promote feeding behavior through canonical homeostatic feeding circuitry (Luo et al., 2018) and learning/hedonic feeding in males (Mohammad et al., 2021). However, this region is an excellent prospective candidate for integrating metabolic and reproductive cues to affect feeding based on its anatomical location, possible sensitivity to circulating reproductive hormones, detection of metabolic hormones such as ghrelin (Luo et al., 2018), and promotion of feeding behavior. Indeed, whole-body knockouts of SST exhibit weight gain that is exacerbated by sex category and high fat diet (Luque et al., 2016).

Here, we use mice to interrogate the role of SST neurons of the tuberal nucleus (TN^{SST}) in integrating metabolic and reproductive cues to affect feeding. TN^{SST} neurons exhibited differential control of feeding in female and male mice (defined by anogenital distance at weaning and postmortem inspection of the gonads), with neuronal ablation decreasing food intake only in females. This effect was primarily due to a decrease in food intake during proestrus, when circulating ovarian hormones are at high concentrations, and was only observed in animals with a low body mass. To determine whether adiposity could mediate the effect of body mass on food intake, fat transplantation experiments were performed. Increased white adipose tissue was sufficient to induce TN^{SST} neuronal modulation of food intake. Together, these data reveal a context dependent role for TN^{SST} neurons in the regulation of food intake, by which TN^{SST} neurons tune feeding behavior in response to metabolic and reproductive states.

Results

Chemogenetic activation of TN^{SST} neurons increases food intake in female and male mice

To test the role of TN^{SST} neurons across sexes, an AAV expressing a Cre-dependent Gq-coupled hM3Dq (Krashes et al., 2011) was stereotactically injected to the TN of *Sst-Cre* mice (Figure 1A-C). Overall, activation of TN^{SST} neurons using the small molecule ligand clozapine-N-oxide (CNO) increased daytime food intake over a four-hour testing period in both females and males (Figure 1D). Control animals without expression of hM3Dq-mCherry confirmed no effect of CNO alone, while within-subjects comparisons of animals expressing hM3Dq in TN^{SST} neurons indicated an increase in feeding upon CNO-induced cellular activation. There was a significant interaction between hM3Dq

88 presence and treatment (saline v. CNO), both with time (treatment*hm3Dq*time, $F(2,105)=3.2964$,
89 $p=0.0409$) and without time (treatment*hm3Dq, $F(1,105)=35.2054$, $p<0.0001$). The effect of neuronal
90 activation (genotype-by-treatment interaction) was most prominent across sexes at 4 hours post-CNO
91 injection (females: $F(1,12)=10.0208$, $p=0.0081$; males: $F(1,9)=12.1521$, $p=0.0069$), though males also
92 exhibited a significant activation-by-treatment interaction at 2 hours post-CNO administration
93 ($F(1,9)=8.6957$, $p=0.0163$). Post-hoc within-subjects analyses of animals bilaterally transduced with
94 hm3Dq-mCherry indicated a significant increase in food intake during activation by CNO as compared




Figure 1: Transient activation of TN^{SST} neurons increased food intake across sexes. (A) Schematic of experimental paradigm. AAVs encoding hm3Dq-mCherry or GFP within Flip-Excision (FLEX) cassettes were injected bilaterally into the tubular nucleus (TN) of *Sst-Cre* or wildtype mice. Created with BioRender.com. (B) Fluorescent images of mCherry (magenta) and cFOS (yellow) in the TN of *Sst-Cre* mice injected with h3MDq-mCherry show mCherry expression and more cFOS positive cells 90 minutes after CNO injection regardless of animal sex (overall effect of hm3Dq-mCherry presence, $F(1,21)=73.634$, $p<0.0001$; no interaction effect, $F(1,21)=0.939$, $p=0.3435$). Dotted line indicates boundaries of the TN. (C) Quantification of cFOS+ cells in the TN. Activation of TN^{SST} neurons in both female and male mice leads to higher food intake within the four-hour daytime testing period (left column, interaction between hm3Dq-mCherry presence and treatment: $F(2,105)=3.2964$, $p=0.0409$). (D) Within sex, only males exhibited an effect of activation and treatment over time ($F(2,45)=3.2793$, $p=0.0468$), though both females and males exhibited an effect of hm3Dq presence and treatment independent of time ($F(1,60)=12.7928$, $p=0.0007$ and $F(1,45)=25.1794$, $p<0.0001$, respectively). Males also exhibited a significant hm3Dq-by-treatment interaction at 2 hours post-CNO ($F(1,9)=8.69$, $p=0.0163$). CNO did not affect food intake in wild-type control mice (right column). Mean $\pm$ SEM, * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$. F Control $n=6$; M Control $n=5$, F hm3Dq $n=8$; M hm3Dq $n=6$.

to treatment with saline control (females overall: $t(31)=2.8486$, $p=0.007732$; males at 2 hours: $t(5)=2.9701$, $p=0.0311$; males at 4 hours: $t(5)=3.3263$, $p=0.0286$). Thus, activating TN^{SST} neurons elicits feeding across sexes.

Caspase ablation of TN^{SST} decreases food intake only in females

To determine if permanent inactivation of TN^{SST} alters feeding across sexes, an AAV expressing a Cre-dependent modified caspase virus (taCasp3-TEVp; (Yang et al., 2013) was stereotactically delivered to the TN of *Sst-Cre* mice (Figure 2A). Bilateral elimination of *Sst* expression was validated by *in situ* hybridization (Figure 2B). Mice were subjected to two 96-hour food assays along with a battery of other metabolic tests. Final food intake, accounting for spillage, is depicted as an average over 24 hours (Figure 2C). ANOVA revealed an overall effect of sex where males consume more food than females, as expected ($F(1,38)=14.1896$, $p=0.0006$).

Interestingly, the effect of TN^{SST} neuron ablation differed by sex (sex-by-ablation interaction: $F(1,38)=4.6852$, $p=0.0368$), an effect which post-hoc t-tests revealed to be carried by a decrease in food intake specifically in females ($t(15.671)=-3.0561$, $p=0.007686$, Figure 2C). This sex difference was not previously reported, but collapsing these data across sex results in an overall decrease in food intake with TN^{SST} neuron ablation ($t(39.86)=-2.2713$, $p=0.0286$), consistent with previous studies (Luo et al., 2018).

The female specific effect of TN^{SST} neuron ablation is modulated by estrous cycle stage. There was a significant interaction between TN^{SST} neuron ablation and estrous stage ($F(1,45)=8.1581$, $p=0.0065$) which was predominantly due to a decrease in food intake specifically during the night of




Figure 2: Caspase ablation of TN^{SST} neurons decreased food intake only in females. (A) Schematic of experimental paradigm. AAVs encoding taCasp3-TEVp or GFP within Flip-Excision (FLEX) cassettes were injected bilaterally into the tuberal nucleus (TN) of *Sst-Cre* or wildtype mice. Created with BioRender.com. (B) Representative brightfield images of *Sst* transcript expression in the TN of caspase-ablated and control animals. Dotted line indicates boundary of the TN. (C) Permanent TN^{SST} neuronal ablation decreases average daily food intake in females but not males. F Control $n=10$; M Control $n=11$; F Ablated $n=11$; M Ablated $n=10$. (D) This decrease in food intake is detected only in the night of proestrus. Proestrus: Control $n=6$, Ablated $n=8$; Estrus: Control $n=7$, Ablated $n=7$; Metestrus: Control $n=10$, Ablated $n=9$; Diestrus: Control $n=10$, Ablated $n=9$. Mean $\pm$ SEM; between subjects: * $p<0.05$, ** $p<0.01$; within subjects: # $p<0.10$.

proestrus ($t(5.902)=-2.6044$, $p=0.04104$, Figure 2D). Within-subjects analysis of mice in the neuronal ablation group suggested that nighttime food intake during proestrus was also slightly lower than consumption during metestrus ($t(7)=-1.976$, $p=0.0887$) and diestrus ($t(7)=-2.3276$, $p=0.0528$), although these effects across the cycle did not reach statistical significance.

Despite effects on food intake, no other metabolic measures were altered by TN^{SST} neuron ablation (see Table 2 for statistical results). TN^{SST} neuron ablation did not affect telemetry measures of activity/movement (Supp Fig 1A) or core body temperature (Supp Fig 1B), nor response to fasting glucose tolerance test (Supp Fig 1C). TN^{SST} neuron ablation did not affect body mass (Supp Fig 1D), suggesting that the selective decrease in food intake during proestrus was not sufficient to alter body mass.

Body mass, specifically adiposity, influences effect of TN^{SST} neuronal ablation on food intake

There was an overall interaction between body mass and estrous phase ($F(1,150)=3.9433$, $p=0.04889$) and between body mass and ablation ($F(1,150)=16.9924$, $p<0.0001$; see Table 2 for full results). Analyzing the relationship between body mass and food intake within each estrous stage revealed significant negative correlations between body mass and food intake in wild-type animals during nights of proestrus ($r^2=0.3263$; $F(1,17)=8.235$, $p=0.01062$; Figure 3A) and metestrus ($r^2=0.2801$; $F(1,16)=6.227$, $p=0.0239$; Supp Fig 2), stages with higher relative circulating estradiol levels (Handelsman et al., 2020). There were no correlations between food intake and body mass in estrus and diestrus, stages with lower relative circulating estradiol levels (Figure 3A, Supp Fig 2, Table 2). TN^{SST}




Figure 3: The effect of TN^{SST} neuron ablation in proestrus depends on body mass or adiposity. (A) Regression analysis within proestrus (top panel) and estrus (bottom panel) across all ovary-intact animals reveals an interaction between body mass and nightly food intake in females. Significant negative correlations in wildtype animals are seen in the high estradiol phase of proestrus but not in caspase ablated females. Control $n=19$, Ablated $n=21$ across stages. (B) Representative brightfield images of inguinal and gonadal white adipose tissue (iWAT & gWAT, respectively) in high and low body weight (BW) females. (C) Terminal iWAT (left) and gWAT (right) adipocyte size both positively correlate with starting body weight regardless of TN^{SST} neuron ablation (Interaction of slopes: iWAT $F(1,22)=0.2337$, $p=0.6336$, gWAT $F(1,18)=0.4552$, $p=0.5085$; difference in y-intercept: iWAT $F(1,22)=0.1417$, $p=0.7103$, gWAT $F(1,18)=0.000$, $p=0.9971$). Linear regression lines represent compiled data across ablation status. iWAT: Control $n=12$, Ablated $n=14$; gWAT: Control $n=11$, Ablated $n=11$. Mean $\pm$ 95% CI, $^*p<0.05$, $^{**}p<0.01$

neuron ablation uncoupled this relationship with body mass in high estradiol stages (proestrus: $r^2=0.00029$, $F(1,19)=0.05492$, $p=0.08172$; metestrus: $r^2=0.09921$, $F(1,19)=2.093$, $p=0.1643$), suggesting that this neuronal population is required for the body mass-dependent modulation of food intake during these stages.

Given the known influence of fat mass on feeding, adiposity was examined post-mortem. Post-mortem adipocyte size of subcutaneous inguinal and visceral perigonadal white adipose tissue (iWAT and gWAT, respectively) positively correlated with the body mass determined at the onset of feeding assays (iWAT: $F(1,22)=4.3346$, $p=0.04919$; gWAT: $F(1,18)=14.9872$, $p=0.001119$; Figure 3B&C) regardless of TN^{SST} neuron ablation (ANCOVA revealed no significant interaction of slopes or difference in y-intercept, see Table 2). Body mass accounted for a larger percentage of variation in visceral gWAT adipocyte size ($r^2=0.4481$; $F(1,20)=16.24$, $p=0.0006557$) than it did for subcutaneous iWAT ($r^2=0.1623$; $F(1,24)=4.649$, $p=0.0413$), suggesting a possible increased contribution of visceral adiposity to the effect of TN^{SST} neuron ablation.

TN^{SST} neurons are sensitive to estrogens and adiposity signals

To test if TN^{SST} neurons can respond to estrogens and/or signals from white adipose tissue, we profiled the transcriptome of fluorescently labeled TN^{SST} neurons using flow cytometry followed by bulk RNA sequencing (Flow-Seq; Figure 4A). Transcriptomic analysis of isolated TN^{SST} neurons




Figure 4: A subset of TN^{SST} neurons is sensitive to estrogens. (A) Schematic of FlowSeq analysis (also applies to Figure 5 genetic input). AAV-flex-tdTomato was stereotactically injected into the tubular nucleus (TN) of *Sst-Cre* mice. Lateral hypothalamic region was grossly dissected and enzymatically dissociated before segregating *Sst*⁺ red neurons via flow cytometry. Resultant cells were sent for Bulk RNA sequencing. Created using BioRender.com. (B) Numerous genes are more differentially expressed between females and males (black dots), including canonical Y-associated genes and *Esr1* (purple dot). (C) Representative fluorescent images from *Esr1-Cre* mice showing colocalization of *Sst* (magenta), *Esr1::tdTomato* (yellow), and DAPI nuclear stain (cyan). White arrows indicate *Sst-Esr1* double labeled cells. (D) Quantification of colocalization confirms *Esr1/Sst* co-expression but reveals no sex difference. Mean $\pm$ SEM.

uncovered numerous differentially expressed genes between females and males, with the gene for estrogen receptor alpha, *Esr1*, being more highly expressed in females ($W\chi^2=6.736$, adj $p=2.356 \times 10^{-8}$; Figure 4B). However, we were unable to detect ER α immunoreactivity in the TN using standard antibodies (Millipore Sigma 06-935). Instead, we confirmed expression from the *Esr1* locus through the injection of a Cre-dependent tdTomato reporter into the TN of female and male *Esr1-Cre* mice. Subsequent colocalization of tdTomato with *Sst* via *in situ* hybridization revealed co-expression of *Esr1* in approximately 10% *Sst*-expressing cells across sexes (Figure 4C&D), confirming that at least a subset of TN^{SST} neurons is sensitive to estrogens.

To determine if TN^{SST} neurons communicate with adipose tissue or *vice versa*, we used a co-correlation analysis method based on genetic variation. High expressing genes from TN^{SST} neurons (based on counts > glial fibrillary acidic protein, GFAP, expression) were used as the “target” pathways, where human orthologues within the GTEx database (Lonsdale et al., 2013) and subjected to cross-tissue genetic co-correlational analyses (Seldin et al., 2018; Velez et al., 2022) (Figure 5A). As previous data indicated that TN^{SST} neuron responsivity to metabolic cues might be localized to periods of higher circulating estradiol (Figure 2D) and TN^{SST} may be able to directly sense circulating estradiol levels (Figure 4), individuals in the GTEx database were binned into groups with either “high” or “low” circulating estradiol levels via weighted aggregation of pan-tissue z-scores corresponding to estrogen responsive gene expression (Sup Fig 3). Binning individuals into groups with indicators of “low” or “high” estrogen signaling revealed weakly associated groups as per biweight midcorrelation (Langfelder and Horvath, 2008); bicor coefficient = -0.32, $p = 0.0023$), suggesting that strong cross-tissue interactions differed depending on estrogen signaling status. Next, genetic co-correlation analyses from adipose (subcutaneous & omental), skeletal muscle, stomach, and small intestine to hypothalamic highly expressed genes were conducted. A lack of significant co-correlations with small intestine resulted in this tissue being omitted from the rest of analyses. Given that subsets of strong cross-tissue correlations remained for the other tissues to highly-expressed hypothalamic DEG orthologues, relevant pathways which might contribute to signaling were examined accordingly (Figure 5). Significant interactions for co-correlations between estrogen signaling group and tissue were observed across tissues for all secreted proteins (Kruskal-Wallis test for interaction between estrogen category + tissue $p=1.7 \times 10^{-218}$; Figure 5B), known ligands (Kruskal-Wallis interaction $p=9.9 \times 10^{-20}$; Figure 5C), peptide hormones (Kruskal-Wallis interaction $p=1.8 \times 10^{-13}$, data not shown), and feeding behavior pathways (Kruskal-Wallis interaction $p=0.0023$; Figure 5D). In addition, several pathways showed specificity in strength of co-correlations from one tissue to another. For example, individuals in the higher inferred estrogen signaling group exhibited higher co-correlations between TN^{SST} and adipose within secreted proteins ($p<2.2 \times 10^{-16}$), ligand ($p=0.025$), and feeding behavior pathways ($p=0.042$) as compared to individuals in the low estrogen signaling group. Interestingly, this relationship was reversed for all secreted proteins in skeletal muscle, with individuals in the low estrogen signaling group exhibiting higher co-correlations ($p=4.2 \times 10^{-12}$). These results indicate increased communication between adipose and TN^{SST} neurons during periods of high estrogen signaling, and a switch to skeletal muscle communication when estrogen signaling is low. Across all gene sets, individuals with inferred low estrogen signaling exhibited higher co-correlations with stomach as compared to individuals with high estrogen signaling (all secreted proteins: $p<2.2 \times 10^{-16}$; ligands: $p=1.2 \times 10^{-5}$; peptide hormones: $p=1.8 \times 10^{-4}$; and feeding behavior: $p=0.029$). Together, these human genetic co-correlation data indicate that TN^{SST} neurons modulate feeding pathways through preferential communication with adipose when estrogen signaling is high and stomach hormones when estrogen signaling is decreased. These observations are in line with known responsivity of TN^{SST} neurons to stomach peptide hormone and known regulator of feeding ghrelin (Luo et al., 2018), but indicate that this communication pathway may be more salient when estradiol levels or

200 estrogen signaling is low. Further, these analyses of human data may suggest that a similar a similar
201 integration of metabolic cues alongside reproductive hormones in humans as well as mice.




Figure 5: TN^{SST} neurons display increased hormonal pathway co-correlations with white adipose tissue in individuals with inferred high estrogen signaling. (A) Schematic overview of co-correlation analysis. High expressing TN^{SST} genes from mouse Flow-Seq experiments were co-correlated across various peripheral metabolic tissues across high and low estradiol groups identified in the GTEx database. Created with BioRender.com. (B) Inferred estradiol levels affected co-correlation pathways relevant to all secreted proteins. High estradiol individuals showed increased co-correlations in adipose tissue whereas those with lower estradiol showed increases in skeletal muscle and stomach communication. (C) Similar trends in adipose and stomach co-correlations across estradiol groupings were seen for ligand pathways. (D) Co-correlations across tissues showed differential impact on genes associated with feeding pathways. Individuals with higher estradiol showed increased co-correlations within these pathways with adipose tissue and decreased co-correlations with stomach as compared to those with lower estradiol levels.

202 To test the causal, directional relationship between fat and TN^{SST} neurons in the modulation of
203 food intake, caspase ablation studies were repeated in combination with fat transplantation.
204 Approximately 1.5 weeks following transplantation of ~0.8 g subcutaneous fat (Figure 6A), recipient
205 mice exhibited significantly increased body mass ($F(1,25) = 28.3184$, $p < 0.0001$; Figure 6B), raw fat
206 mass ($F(1,25) = 34.2342$, $p < 0.0001$; Figure 6B), and percent fat mass ($F(1,25) = 30.0008$, $p < 0.0001$;

Figure 6C) and no interaction with TN^{SST} neuronal ablation in any case. Thus, fat transplantation increased adiposity similarly across neuronal ablation groups.

Fat transplant also increased food intake in general ($F(1,25)=52.524$, $p<0.0001$), and the effect of TN^{SST} neuronal ablation was affected by fat transplant ($F(1,25)=26.660$, $p<0.0001$; Figure 6D). *Post-hoc* t-tests revealed that TN^{SST} neuronal ablation significantly decreased food intake in sham transplant animals ($t(11.646)=-2.917$, $p=0.01327$) but significantly increased food intake in animals receiving fat transplant ($t(4.8.2536)=4.8427$, $p=0.001175$), similar to the previous relationship with body mass in proestrus (Figure 3A). However, we were unable to detect a significant interaction with fat transplantation and ablation status over the estrous cycle (Figure 6E), possibly due to high variability in nighttime food intake in sham-transplanted mice. Together, these findings indicate that fat transplantation masks the proestrus-specific effect of TN^{SST} neuron ablation and reveal a role for fat mass in modulating the function of TN^{SST} neurons within the feeding circuit.




Figure 6: Fat transplantation modulates the effect of TN^{SST} neuron ablation. (A) Representative schematic of fat transplants. Four deposits of 0.2 g each were placed in the dorsal subcutaneous region. Created with BioRender.com. (B) Fat transplant increases raw body mass (left axis, $p<0.0001$) and fat mass (right axis, $p<0.001$) regardless of TN^{SST} neuronal ablation. (C) This translates to an overall increase in adiposity ($p<0.0001$). (D) Fat transplantation reverses the effect of TN^{SST} ablation, significantly increasing daily food intake compared to non-ablated controls. (E) This effect of fat transplant (top panel) seems to be due to a lack of effect during proestrus, though data were underpowered to detect the effect of estrous stage in sham controls (bottom panel). Mean $\pm$ SEM; within adiposity group: * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$; between adiposity group: ### $p<0.001$. Sham: Control $n=10$, Ablated $n=5$; Transplant: Control $n=7$, Ablated $n=7$. Sham: Proestrus Control $n=10$ & Ablated $n=5$, Estrus Control $n=9$ & Ablated $n=5$, Metestrus Control $n=10$ & Ablated $n=5$, Diestrus Control $n=10$ & Ablated $n=4$; Fat Transplant: Proestrus Control $n=6$ & Ablated $n=5$, Estrus Control $n=6$ & Ablated $n=6$, Metestrus Control $n=6$ & Ablated $n=6$, Diestrus Control $n=6$ & Ablated $n=6$.

Discussion

These data suggest that TN^{SST} neurons are a locus in the brain that mediates metabolic and reproductive tradeoffs. While activation of TN^{SST} neurons increases food intake across sexes, permanent inactivation by ablation during adulthood results in decreased food intake only in females during the proestrus phase. This effect depends on body mass, as this effect is apparent only in lighter animals. In wildtype mice, body mass inversely correlates with food intake on the night of proestrus, but TN^{SST} neuron ablation uncouples this effect. Further analysis reveals that white adipose tissue abundance is a significant contributing factor to this effect. Not only does post-mortem adipocyte size correlate with body mass in neuron ablation experiments, but fat transplantation studies confirm that TN^{SST} neuron ablation only decreases food intake in lean animals compared to their fat transplanted counterparts. This interaction between cycling adipokines and gonadal hormones may be mediated by the direct effects of these circulating molecules on TN^{SST} neurons, as these cells show some estrogen sensitivity via co-expression analyses. Furthermore, co-correlations between the hypothalamus and adipose tissue in humans and fat transplantation experiments in mice point to the importance of secreted proteins and ligands, suggesting TN^{SST} may detect and respond to adipokines. Future studies are needed to confirm and dissect the mechanisms of these cellular effects.

What adipokine factor is possibly being detected by the TN^{SST} remains to be determined. Leptin positively correlates with overall adiposity (Fontana and Della Torre, 2016), and it has been known to play a crucial role in reproductive responsiveness to metabolic condition, namely as the permissive signal required for pubertal onset (Chehab et al., 1996; Cheung et al., 1997). Adiponectin, an adipokine that negatively correlates with visceral fat mass in mammals (Fontana and Della Torre, 2016), has long been shown to downregulate reproduction through direct impacts on the hypothalamus (Rodriguez-Pacheco et al., 2007). Resistin is correlated with higher overall adiposity (Yang et al., 2012), exhibits numerous interactions with the hypothalamic-pituitary-gonadal axis (Mathew et al., 2018; Nogueiras et al., 2003; Tsatsanis et al., 2015), and is down-regulated during the fertile periods of the mouse estrous cycle (Gui et al., 2004).

Regardless of adipokine contributor, this trade-off paradigm provides a plausible explanation for the varied effects of estradiol on food intake in mice. While endogenous fluctuations and experimental manipulations of estradiol consistently reveal that estrogens decrease food intake in rats and guinea pigs (Asarian and Geary, 2013, 2002; Clegg et al., 2007; Eckel, 2011), the mouse literature is less definitive (Eckel, 2011; Geary et al., 2001; Naaz et al., 2002; Petersen, 1976; Witte et al., 2010). Instead, the more consistent phenotype in mice is a decrease in energy expenditure following estradiol depletion (Correa et al., 2015; Musatov et al., 2007; Xu et al., 2011). In light of this study, it is possible that the effects of estradiol on feeding across mouse studies, as observed by either endogenous estrous cycle fluctuations or ovariectomy manipulation, could be confounded by body mass and adiposity. Thus, factors like age at time of experiment, differences in fat distributions between species or strains, diet, or ovariectomy and time from ovariectomy to estradiol replacement might present confounds based on changes to fat and/or lean mass.

How circulating estrogen levels contribute to this circuit also requires further investigation. Our human GTEx analyses shows that co-correlations between TN^{SST} genes and that in peripheral tissue shifts from predominantly adipose-based to skeletal muscle- and stomach-based depending on evident estrogen signaling (Figure 5). This suggests that higher estrogen levels may increase communication between TN^{SST} neurons and white adipocyte depots, particularly as it relates to regulation of feeding behavior (Figure 5D). While this could be due to the actions of circulating estrogens on white adipose tissue itself (reviewed in (Hevener et al., 2015; Palmer and Clegg, 2015)), it is also possible that estrogens directly act on TN^{SST} neurons to increase their sensitivity and/or responsiveness to adipokines. Indeed, TN^{SST} neurons exhibit estrogen sensitivity (Figure 4), though future studies would be needed to test for a possible direct effect.

Alternatively, fluctuating hormone levels might be detected elsewhere in the brain and impact TN^{SST} neuronal modulation of feeding through integration at the circuit level. TN^{SST} neurons project to many estrogen-sensitive nodes or nodes receiving direct input from estrogen-responsive regions, including the bed nucleus of the stria terminalis, parabrachial nucleus, and central amygdala (Luo et al., 2018). This circuit-wide integration of estradiol is a known mechanism of action for the gonadal hormone, with estrogens acting on many circuit nodes to coordinate behavioral output in a variety of cases, including reward/addiction (Becker and Chartoff, 2019) and thermoregulation (Zhang et al., 2021). It is therefore probable that the effects of estradiol on feeding function similarly, as the anorexigenic effects of estradiol have been localized to numerous feeding nodes such as the hypothalamic arcuate nucleus (Roepke et al., 2010, 2007; Santollo et al., 2011; Todd L Stincic et al., 2018; Todd L. Stincic et al., 2018) and the nucleus of the solitary tract of the brainstem (Asarian and Geary, 2006; Maske et al., 2017).

In all, this study adds to the growing literature interrogating the contributions of TN^{SST} neurons to feeding behavior. Central SST (originally named growth hormone inhibiting hormone, GHIH, in the central nervous system, (Painson and Tannenbaum, 1991) had long been known to affect food intake through somatostatin receptor 2 (SSTR2; (Beranek et al., 1999; Campbell et al., 2017; Danguir, 1988; Lin et al., 1989; Andreas Stengel et al., 2010b, 2010a; A Stengel et al., 2010; Stengel et al., 2015, 2013, 2011; Tachibana et al., 2009). This effect was seemingly localized to the tuberal nucleus, when TN^{SST} neurons were found to integrate into the melanocortin feeding system, though the effect of these neurons on feeding was attributed to γ -aminobutyric acid (GABA) release as opposed to direct SST effects (Luo et al., 2018). Subsequently, TN^{SST} neurons were also found to contribute to food context learning in males¹ (Mohammad et al., 2021), indicating that the TN may straddle homeostatic and hedonic feeding mechanisms (Massa and Correa, 2020). This paper adds to this growing literature by not only delineating an apparent sex difference but also context dependence in TN^{SST} neuronal modulation of food intake.

We further speculate that TN^{SST} neurons serve as a nexus of integration and a mediator of reproductive and metabolic tradeoffs within the feeding circuit. In cycling rodents, fertile periods during the estrous cycle are accompanied by alterations to metabolic output, including a decrease in food intake (Asarian and Geary, 2013, 2002; Brobeck et al., 1947; Eckel, 2011), increase in locomotion (Brobeck et al., 1947; Kent et al., 1991; Sanchez-Alavez et al., 2011; Steiner et al., 1982), and increased core body temperature (Kent et al., 1991; Sanchez-Alavez et al., 2011). These changes are hypothesized to suppress energy intake and promote active mate-seeking behavior and sexual receptivity. This study identifies TN^{SST} neurons as possible mediators of such a trade-off, actively promoting energy intake during fertile periods when metabolic reserves may be insufficient to support reproduction.

Materials & Methods

Mice

Female (defined as having small anogenital distance at weaning and presence of ovaries at time of death) and male (defined as having large anogenital distance at weaning and presence of testes postmortem) mice expressing the *Sst-Cre* driver transgene (JAX stock no. 013044, *Sst^{tm2.1(cre)Zjh}/J*) were maintained on a C57BL/6J genetic background. Heterozygotes (*Sst-Cre/+*) and/or wildtype littermates (+/+) were used for all studies. Genotypes were determined as per JAX protocol 28317. Female and male mice expressing the *Esr1-Cre* driver transgene (JAX stock no. 017911, B6N.129S6(Cg)-*Esr1^{tm1.1(cre)And}/J*) were maintained on a C57BL/6J genetic background. Heterozygotes (*Esr1-Cre/+*) were used for colocalization studies. Genotypes were determined as per primers from JAX protocol 27213. Experiments were performed on cycling females and gonadally-intact males unless otherwise stated. Mice were maintained on a 12:12 light cycle, with *ad libitum* access to food and water (unless

¹ In all external papers discussed, no definitions for sex category were ever provided. In mice, we assume that sexes were defined using anogenital distance.

otherwise specified), under controlled humidity conditions, and in single-housed cages with non-caloric paper bedding to ensure accurate food intake assessment. All studies were carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. UCLA is AALAS accredited, and the UCLA Institutional Animal Care and Use Committee (IACUC) approved all animal procedures.

Estrous cycle staging

Vaginal lavages were performed on females daily, between ZT 0 and ZT 4, using 30 μ L of standard phosphate buffered saline (PBS). Samples were deposited onto slides and allowed to dry prior to staining. Males were subjected to similar handling during this time to ensure roughly equivalent handling stress. Giemsa staining was carried out to visualize cellular composition of the vaginal cavity. Stock Giemsa stain was prepared at least one week in advance of use. An 18.5% solution of Giemsa powder (Fisher G146-10) in glycerin was heated to 60°C and cooled before diluting 9:14 with 100% methanol. Stock was diluted 1:30 in PBS before use, shaking vigorously before stain. Slides were incubated for one hour at room temperature. Prior to staining, slides were briefly fixed in 100% methanol. Staging was assessed via light microscopy as in (Cora et al., 2015), and stages were assigned using the behavioral method (Becker et al., 2005), with morning samples indicating the prior night's estrous stage. This staging method was confirmed by core body temperature waveform alignment (Sanchez-Alavez et al., 2011).

Surgical Procedures

Mice received analgesics (0.074 mg/kg buprenorphine two times daily, 7.11 mg/kg carprofen one time daily) on the day of and one day post-surgery. Mice were anaesthetized with 3% isoflurane and maintained within a range of 1.25-2.5%. AAVs were bilaterally injected into the TN of adult mice (coordinates relative to Bregma: A-P -1.65 mm, lateral $\pm$ 0.75, D-V -5.45; scaled when Bregma-Lambda distance was not equivalent to 4.2 mm) at a rate of 5 nL/s using a glass-pulled needle. See Table 1 for titers and injection volumes. Controls consisted of both wildtype animals injected with the experimental virus (virus controls) and Cre positive animals injected with cell-filling GFP (genotype controls). Ovariectomy surgeries included complete removal of gonads from adult mice. Gonadectomies occurred immediately prior to stereotaxic viral injections within the same surgical period. In telemetry experiments, G2 eMitters (Starr Life Sciences) were implanted intraperitoneally on the same day as viral injection. Experiments were conducted following at least two weeks recovery from surgical proceedings.

Caspase ablation experiments

Gross movement and core body temperature were passively measured every other week for eight weeks using VitalView software (Starr Life Sciences). Body weight was measured every week. Food assay was performed when mice were not on telemetry pads. At ZT 0.5 on the start day of the experiment, 2/3 of the non-caloric paper bedding was removed. A pre-measured amount of food was delivered, and mouse body weight measured. Food in hopper was weighed at ZT 0.5 and ZT 11.5 every day until experiment conclusion. After 96 hours, food and all bedding were collected to account for food spillage. For some experiments, 4-5 hour fasted glucose tolerance tests were performed prior to sacrifice. In ovariectomy experiments, two food assays were performed back-to-back, non-fasted resting glucose levels were collected, body composition was measured via NMR, and indirect calorimetry was performed in Oxymax metabolic chambers (Columnbus Instruments) at room temperature. Upon experiment completion, all brains were collected using RNase-free conditions. Inguinal white adipose tissue (iWAT) and gonadal white adipose tissue (gWAT) were collected for histology analyses.

Transient activation food intake assay

Clozapine-n-oxide (CNO; MilliporeSigma #0832) was used to activate TN^{SST} neurons in *Sst-Cre* animals expressing hM3Dq-mCherry. Stock solution of 20 mg/mL in DMSO was stored at -20°C and

diluted to a working solution of 0.03 mg/mL in sterile saline also stored at -20°C. Saline control (0.15% DMSO) or CNO (10 µL/g body weight, dose of 0.3 mg/kg) working solution were administered IP in a counterbalanced design. Experiments were completed in duplicate replicate trials. Mice were transferred to experimental room at least 15 minutes prior to experimentation. Experiments were begun between ZT 2-3 and terminated between ZT 6-7. Following injection, food intake was measured at 0.5, 1, 2, and 4 hr. Vaginal lavage was performed on female mice after experiment conclusion to prevent stress interference with food intake. All mice were injected with CNO 90 minutes prior to sacrifice to enable neuronal activation validation via cFOS immunohistochemistry.

Fat Transplantations

Donor fat was taken from various visceral (i.e., periuterine perigonadal, retroperitoneal, and omental) depots of wildtype female C57BL/6J mice and implanted into female mice recently stereotactically injected under standard surgical conditions. Four depots of 0.15-0.25g were placed subcutaneously on the dorsal surface through a single incision mid-back, for a final transplantation total of 0.6-0.9g of white adipose. Fat for each depot was divided into at least three individual pieces to promote vascularization. The visceral-to-subcutaneous paradigm was used due to the deleterious metabolic effects of this graft (Tran et al., 2008). Food intake was assayed 2-3 weeks following transplantation to allow for sufficient angiogenesis (Gavrilova et al., 2000) and graft stabilization without endogenous fat depot compensation (Rooks et al., 2004). Upon sacrifice, grafts were examined to confirm tissue was not necrotic.

Histology

In situ hybridization (ISH) and immunostaining (IHC)

RNA probe generation was accomplished as in (van Veen & Kammel et al., 2020). Briefly, *Sst* sense and antisense probes were transcribed using a DIG or FITC RNA labeling kit (Roche) and purified with RNA Clean & Concentrator (Zymo Research). PCR products were amplified using Allen Brain Institute-derived reference primer sequences and cloned into pCR 2.1 TOPO (Invitrogen). Plasmid DNA was then isolated from bacterial cultures (ZymoPURE II Plasmid Midiprep kit), linearized, and purified (Zymo DNA Clean & Concentrator). Validation of caspase ablation was carried out on 35µm-thick coronal slices via chromogen ISH protocol as per (van Veen & Kammel et al., 2020). Validation of hM3Dq targeting and activation was accomplished by visualization of native mCherry expression and IHC stain for cFOS. Briefly, slides were blocked and incubated with rabbit anti-cFOS (1:200, Synaptic Systems # 226003, RRID: 2231974) primary antibody overnight at 4°C. The next day, sections were incubated for 1 hour at room temperature with goat anti-rabbit Alexa Fluor 488 secondary (1:500, Thermo Fisher Scientific # A11034, RRID: AB_2576217) and counterstained with DAPI. For colocalization experiments, *Esr1-Cre* mice were bilaterally injected with 400 µl AAV2-flex-tdTomato into the TN coordinates. Native tdTomato fluorescence destroyed by combined ISH protocol was recovered by rabbit anti-DsRed (1:1000, Takara Bio Clontech # 632496, RRID: AB_10013483) antibody and switched to the green channel using an Alexa Fluor 488 secondary. Dual *Sst* ISH & tdTomato IHC protocol was accomplished via TSA amplification. Briefly, 35 µm sections were fixed, permeabilized with Triton X-100, and acetylated before overnight ISH probe incubation at 65°C. The next day, tissue was then washed, blocked with Blocking Reagent (MilliporeSigma 11096176001Roche) and heat inactivated sheep serum, and incubated with anti-DsRed overnight at 4°C. The final day, tissue was washed before ISH signal was developed with the TSA Plus Cyanine 5 System (Akoya Biosciences # NEL745001KT). Slides were then stripped of horseradish peroxidase and blocked with normal goat serum before incubating with goat anti-rabbit Alexa Fluor 488 (1:400) for 2 hours at room temperature.

Adipocyte size quantification

Inguinal and white adipose tissue was collected post-mortem and drop-fixed in 4% paraformaldehyde (PFA) for at least 18 hours. Tissue was then washed in PBS before being stored in PBS at 4°C until

tissue analysis. For histological processing, tissue was placed in tissue processing cassettes and submerged in 70% ethanol before being embedded in paraffin, sectioned at 4 μ M, and stained with hematoxylin & eosin (H&E) by the UCLA Translational Pathology Core Laboratory. Three regions of interest per tissue-type per mouse were imaged by light microscopy at 20x magnification. Adipocyte area was quantified using a custom pipeline in CellProfiler. Inclusion parameters were cell diameters of 100-300 pixel units and a global threshold strategy with minimum cross-entropy.

Colocalization analysis

Sst and *Esr1* co-expression was determined using CellProfiler (version 4.2.1). First, a contour was drawn around a matched section of the TN using anatomical landmarks (i.e., shape of arcuate nucleus and third ventricle). For each hemisphere, DAPI-stained nuclei were detected and intensity thresholding was used to determine *Sst*⁺ cells. Incorrectly labeled cells were manually erased or added. *Sst*⁺ cells were then filtered based on *Esr1::tdTomato* signal intensity. Counts were made of total *Sst*⁺ cells, as well as *Sst*⁺/*Esr1*⁺ and *Sst*⁺/*Esr1*⁻ cells. The counts were averaged across the two hemispheres, and percent was calculated as $([Sst+/Esr1+] / \text{Total SST}) \times 100$.

Bioinformatics Analysis

Sst-Cre female and male mice were bilaterally stereotactically injected with AAV expressing Cre-dependent tdTomato (See Table 1). Following at least two weeks for viral expression, animals were sacrificed and TN was microdissected under fluorescent illumination. Dissected TN was dissociated using a papain-based enzymatic process (Worthington Biochemical) and then TN^{SST} neurons were enriched and collected via flow cytometry. Cells were sorted from debris and doublets were excluded by gating on forward-scatter and side-scatter profiles. Live nucleated cells were then selected by DAPI-negative (live) and DRAQ5-positive (nucleated) staining. Finally, tdTomato-positive cells were selected based on relatively high levels of red fluorescence (as in van Veen & Kammel et al., 2020). RNA was isolated from 500-2500 cells by RNeasy Micro kit (Qiagen). Cells were then submitted for bulk RNA sequencing. Single-end reads (~10 million unique reads per mouse) were assembled to the mouse transcriptome (version mm10) using kallisto (version 0.46.2). Differentially expressed genes and normalized read counts were identified using DESeq2 Galaxy Version 2.11.40.6+galaxy1. Volcano plots were produced by the custom R function “deseq_volcano_plot_gs()” available through the following package: <http://github.com/jevanveen/ratplots>. Raw reads of the RNA sequencing data were also examined for hypothalamus-peripheral tissue co-correlations across stomach, small intestine, skeletal muscle, visceral fat, and subcutaneous fat as hypothalamic reads using the GTEx database as previously described (Seldin et al., 2018; Velez et al., 2022). In addition, estrogen-responsive genes used to infer “low” vs “high estrogen signaling were gathered from: https://www.gsea-msigdb.org/gsea/msigdb/cards/HALLMARK_ESTROGEN_RESPONSE_EARLY.html. To clarify the analysis, estrogen signaling binning per individual and subsequent cross-tissue correlations with mouse DEG orthologues, all processed datasets, scripts used to analyze, and detailed walk-through is available at: <https://github.com/Leandromvelez/sex-specific-endocrine-signals>.

Statistical Analyses

All statistics were carried out in R. Sex differences were determined by interaction terms between genotype and sex (caspase ablation experiments) or genotype, treatment, and sex (chemogenetic experiments). In caspase ablation and fat transplantation experiments, animals meeting both the criteria of outlier by Cook’s distance, as well as “miss” (no hit or unilateral hit as defined by more than 5% of targeted cells still present) were excluded. For fat transplantation studies, only sham animals with <10% fat mass at the beginning of the feeding assay were included. All data were checked and transformed, if necessary, to meet normalcy criteria.

Acknowledgements


This research was supported by the UCLA Life Sciences Division (NIH R01 AG066821), the NIH Specialized Centers of Research Excellence (SCORE) grant U54 DK120342, and the National Center for Advancing Translational Science (NCATS) under the UCLA Clinical and Translational Science Institute grant UL1TR001881. MGM was supported by an NSF GRFP (DGE-2034835), Dissertation Year Fellowship from the UCLA Graduate Division, and the UCLA Kenneth I. Shine Fellowship. SA was supported by the CARE program of the UCLA Undergraduate Research Center (NIH NIGMS IMSD GM055052; PI: Hasson). MS was supported by NIH grant DK130640. The authors also wish to thank Drs. A. Arnold, K. Wassum, and E. Hsiao for helpful feedback. MGM would also like to thank all members of the Correa lab who ever assisted with timed food assays.

Competing Interests


The authors have no competing interests to declare.

Experiment	Virus	Depositor & Procurement	Titer (vg/mL)	Volume (nL)	Citation
Caspase ablation	AAV2-flex-taCasp3-TEVp	Nirao Shah & Jim Wells, UNC Vector Core	$1-8 \times 10^{12}$	200-250	(Yang et al., 2013)
Transient activation	AAV8-hSyn-hM3D(Gq)-mCherry	Brian Roth, Addgene viral prep # 50474-AAV8	$\geq 4 \times 10^{12}$	150-200	(Krashes et al., 2011)
Fluorescent localization	AAV2-FLEX-tdTomato	Edward Boyden, Addgene plasmid #28306	$\geq 5 \times 10^{12}$	200 for Flow-Seq; 400 for <i>Esr1-Cre</i>	
Fluorescent controls	AAV8-Syn-FLEX-Mac-GFP	Edward Boyden, Addgene plasmid #58852	1:5 dilution of stock	Matching volume to experimental animals	


Table 1: List of viral vectors used.



Supplementary Figure 1 (companion to Figure 2): TN^{SST} neuronal ablation does not affect any other metabolic measures studied. Telemetry measures of (A) activity and (B) core body temperature are unaffected by TN^{SST} neuronal ablation. (C) Fasting glucose tolerance test is also unaffected by TN^{SST} neuron ablation. (D) Despite changes to food intake, ablation does not affect body weight over time. Mean $\pm$ SEM; *, $p < 0.05$; ***, $p < 0.001$. M Control $n = 9$; M Ablated $n = 9$; F Control $n = 5$; F Ablated $n = 7$. Pro: Proestrus, Est: Estrus, Met: Metestrus, Di: Diestrus.



Supplementary Figure 2 (companion to Figure 3). (A) Regression analysis of food intake and body mass across all ovary-intact animals in metestrus (left panel) and diestrus (right panel) reveals an interaction between body mass and nightly food intake in females. Significant negative correlations in wildtype animals (black line, round black dots) are seen in the higher estradiol phase of metestrus but not in caspase ablated females (cyan line, square cyan points). Metestrus: Control n=18, Ablated n=21; Diestrus: Control n=19, Ablated n=20.



Supplementary Figure 3 (companion to Figure 5). (A&B) Z-scores of estrogen signaling genes (y-axis, https://www.gsea-msigdb.org/gsea/msigdb/cards/HALLMARK_ESTROGEN_RESPONSE_EARLY.html) across indicated tissues (x-axis) in GTEx female (no Y chromosome, A) and male (Y chromosome present, B) individuals. (C&D) Based on the distributions of these scores across relevant metabolic tissues, individuals were segregated into categories of “low” (<0, left of blue line) or “high” (>0, right of blue line) estrogen signaling and used for cross-tissue genetic correlations.

References

- Asarian L, Geary N. 2013. Sex differences in the physiology of eating. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **305**:R1215–R1267. doi:10.1152/ajpregu.00446.2012
- Asarian L, Geary N. 2006. Modulation of appetite by gonadal steroid hormones. *Philosophical Transactions of the Royal Society B: Biological Sciences* **361**:1251–1263. doi:10.1098/rstb.2006.1860
- Asarian L, Geary N. 2002. Cyclic Estradiol Treatment Normalizes Body Weight and Restores Physiological Patterns of Spontaneous Feeding and Sexual Receptivity in Ovariectomized Rats. *Hormones and Behavior* **42**:461–471. doi:10.1006/HBEH.2002.1835
- Ball ZB, Barnes RH, Visscher MB. 1947. The Effects of Dietary Caloric Restriction on Maturity and Senescence, with Particular Reference to Fertility and Longevity. *American Journal of Physiology-Legacy Content* **150**:511–519. doi:10.1152/AJPLEGACY.1947.150.3.511
- Baz B, Riveline J-P, Gautier J-F. 2016. Endocrinology of Pregnancy: Gestational diabetes mellitus: definition, aetiological and clinical aspects. *European journal of endocrinology* **174**:R43–51. doi:10.1530/EJE-15-0378
- Becker JB, Arnold AP, Berkley KJ, Blaustein JD, Eckel LA, Hampson E, Herman JP, Marts S, Sadée W, Steiner M, Taylor J, Young E. 2005. Strategies and Methods for Research on Sex Differences in Brain and Behavior. *Endocrinology* **146**:1650–1673. doi:10.1210/en.2004-1142
- Becker JB, Chartoff E. 2019. Sex differences in neural mechanisms mediating reward and addiction. *Neuropsychopharmacology* **44**:166–183. doi:10.1038/s41386-018-0125-6
- Beranek L, Hajdu I, Gardi J, Taishi P, Obál F, Krueger JM. 1999. Central administration of the somatostatin analog octreotide induces captopril-insensitive sleep responses. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* **277**:R1297–R1304. doi:10.1152/ajpregu.1999.277.5.R1297
- Brobeck JR, Wheatland M, Strominger JL. 1947. Variations in regulation of energy exchange associated with estrus, diestrus and pseudopregnancy in rats. *Endocrinology* **40**:65–72. doi:10.1210/endo-40-2-65
- Burnett CJ, Li C, Webber E, Tsaousidou E, Xue SY, Brüning JC, Krashes MJ. 2016. Hunger-Driven Motivational State Competition. *Neuron* **92**:187–201. doi:10.1016/J.NEURON.2016.08.032
- Campbell JN, Macosko EZ, Fenselau H, Pers TH, Lyubetskaya A, Tenen D, Goldman M, Verstegen AMJ, Resch JM, McCarroll SA, Rosen ED, Lowell BB, Tsai LT. 2017. A molecular census of arcuate hypothalamus and median eminence cell types. *Nature Neuroscience*. doi:10.1038/nn.4495
- Canteras NS, Simerly RB, Swanson LW. 1994. Organization of projections from the ventromedial nucleus of the hypothalamus: A Phaseolus vulgaris-Leucoagglutinin study in the rat. *Journal of Comparative Neurology* **348**:41–79. doi:10.1002/CNE.903480103
- Chehab FF, Lim ME, Lu R. 1996. Correction of the sterility defect in homozygous obese female mice by treatment with the human recombinant leptin. *Nature Genetics* **12**:318–320. doi:10.1038/ng0396-318
- Chen X, McClusky R, Chen J, Beaven SW, Tontonoz P, Arnold AP, Reue K. 2012. The Number of X Chromosomes Causes Sex Differences in Adiposity in Mice. *PLoS Genetics* **8**:e1002709. doi:10.1371/journal.pgen.1002709
- Chen Y, Lin Y-C, Kuo T-W, Knight ZA. 2015. Sensory Detection of Food Rapidly Modulates Arcuate Feeding Circuits. *Cell* **160**:829–841. doi:10.1016/J.CELL.2015.01.033
- Cheung CC, Thornton JE, Kuijper JL, Weigle DS, Clifton DK, Steiner RA. 1997. Leptin Is a Metabolic Gate for the Onset of Puberty in the Female Rat. *Endocrinology* **138**:855–858. doi:10.1210/ENDO.138.2.5054
- Ciofi P. 2011. The arcuate nucleus as a circumventricular organ in the mouse. *Neuroscience Letters* **487**:187–190. doi:10.1016/J.NEULET.2010.10.019

- Clegg DJ, Brown LM, Zigman JM, Kemp CJ, Strader AD, Benoit SC, Woods SC, Mangiaracina M, Geary N. 2007. Estradiol-Dependent Decrease in the Orexigenic Potency of Ghrelin in Female Rats. *Diabetes* **56**:1051–1058. doi:10.2337/DB06-0015
- Cora MC, Kooistra L, Travlos G. 2015. Vaginal Cytology of the Laboratory Rat and Mouse:Review and Criteria for the Staging of the Estrous Cycle Using Stained Vaginal Smears. *Toxicologic Pathology*. doi:10.1177/0192623315570339
- Correa SM, Newstrom DW, Warne JP, Flandin P, Cheung CC, Lin-Moore AT, Pierce AA, Xu AW, Rubenstein JL, Ingraham HA. 2015. An Estrogen-Responsive Module in the Ventromedial Hypothalamus Selectively Drives Sex-Specific Activity in Females. *Cell Reports* **10**:62–74. doi:10.1016/j.celrep.2014.12.011
- Danguir J. 1988. Food intake in rats is increased by intracerebroventricular infusion of the somatostatin analogue SMS 201–995 and is decreased by somatostatin antiserum. *Peptides* **9**:211–213. doi:10.1016/0196-9781(88)90030-7
- Eckel LA. 2011. The ovarian hormone estradiol plays a crucial role in the control of food intake in females. *Physiology and Behavior* **104**:517–524. doi:10.1016/j.physbeh.2011.04.014
- Fontana R, Della Torre S. 2016. The Deep Correlation between Energy Metabolism and Reproduction: A View on the Effects of Nutrition for Women Fertility. *Nutrients* 2016, Vol 8, Page 87 **8**:87. doi:10.3390/NU8020087
- Frisch RE. 1972. Weight at Menarche: Similarity for Well-Nourished and Undernourished Girls at Differing Ages, and Evidence for Historical Constancy. *Pediatrics* **50**.
- Frisch RE, Revelle R. 1970. Height and weight at menarche and a hypothesis of critical body weights and adolescent events. *Science* **169**:397–399. doi:10.1126/SCIENCE.169.3943.397
- Gavrilova O, Marcus-Samuels B, Graham D, Kim JK, Shulman GI, Castle AL, Vinson C, Eckhaus M, Reitman ML. 2000. Surgical Implantation of Adipose Reverses Diabetes Lipoatrophic Mice. *J Clin Invest* **105**:271–78. doi:10.1172/JCI7901
- Geary N, Asarian L, Korach KS, Pfaff DW, Ogawa S. 2001. Deficits in E2-Dependent Control of Feeding, Weight Gain, and Cholecystokinin Satiation in ER- α Null Mice. *Endocrinology* **142**:4751–4757. doi:10.1210/endo.142.11.8504
- Gui Y, Silha JV, Murphy LJ. 2004. Sexual dimorphism and regulation of resistin, adiponectin, and leptin expression in the mouse. *Obesity Research* **12**:1481–1491. doi:10.1038/oby.2004.185
- Handelsman DJ, Gibson E, Davis S, Golebiowski B, Walters KA, Desai R. 2020. Ultrasensitive serum estradiol measurement by liquid chromatography-mass spectrometry in postmenopausal women and mice. *Journal of the Endocrine Society* **4**:1–12. doi:10.1210/jendso/bvaa086
- Hevener AL, Clegg DJ, Mauvais-Jarvis F. 2015. Impaired estrogen receptor action in the pathogenesis of the metabolic syndrome. *Molecular and Cellular Endocrinology* **418**:306–321. doi:10.1016/J.MCE.2015.05.020
- Kammel LG, Correa SM. 2019. Selective sexual differentiation of neuron populations may contribute to sex-specific outputs of the ventromedial nucleus of the hypothalamus. *Journal of Neuroendocrinology*. doi:10.1111/jne.12801
- Kent S, Hurd M, Satinoff E. 1991. Interactions between body temperature and wheel running over the estrous cycle in rats. *Physiology and Behavior* **49**:1079–1084. doi:10.1016/0031-9384(91)90334-K
- Krashes MJ, Koda S, Ye C, Rogan SC, Adams AC, Cusher DS, Maratos-Flier E, Roth BL, Lowell BB. 2011. Rapid, reversible activation of AgRP neurons drives feeding behavior in mice. *The Journal of clinical investigation* **121**:1424–8. doi:10.1172/JCI46229
- Krause WC, Rodriguez R, Gegenhuber B, Matharu N, Rodriguez AN, Padilla-Roger AM, Toma K, Herber CB, Correa SM, Duan X, Ahituv N, Tollkuhn J, Ingraham HA. 2021. Oestrogen engages brain MC4R signalling to drive physical activity in female mice. *Nature* **599**:131–135. doi:10.1038/s41586-021-04010-3

- Langfelder P, Horvath S. 2008. WGCNA: An R package for weighted correlation network analysis. *BMC Bioinformatics* **9**:1–13. doi:10.1186/1471-2105-9-559/FIGURES/4
- Lin MT, Uang WN, Ho LT, Chuang J, Fan LJ. 1989. Somatostatin: a hypothalamic transmitter for thermoregulation in rats. *Pflügers Archiv European Journal of Physiology* **413**:528–532. doi:10.1007/BF00594185
- Lomniczi A, Ojeda SR. 2016. The Emerging Role of Epigenetics in the Regulation of Female Puberty. *Endocrine Development* **29**:1–16. doi:10.1159/000438840
- Lonsdale J, Thomas J, Salvatore M, Phillips R, Lo E, Shad S, Hasz R, Walters G, Garcia F, Young N, Foster B, Moser M, Karasik E, Gillard B, Ramsey K, Sullivan S, Bridge J, Magazine H, Syron J, Fleming J, Siminoff L, Traino H, Mosavel M, Barker L, Jewell S, Rohrer D, Maxim D, Filkins D, Harbach P, Cortadillo E, Berghuis B, Turner L, Hudson E, Feenstra K, Sobin L, Robb J, Branton P, Korzeniewski G, Shive C, Tabor D, Qi L, Groch K, Nampally S, Buia S, Zimmerman A, Smith A, Burges R, Robinson K, Valentino K, Bradbury D, Cosentino M, Diaz-Mayoral N, Kennedy M, Engel T, Williams P, Erickson K, Ardlie K, Winckler W, Getz G, DeLuca D, MacArthur D, Kellis M, Thomson A, Young T, Gelfand E, Donovan M, Meng Y, Grant G, Mash D, Marcus Y, Basile M, Liu J, Zhu J, Tu Z, Cox NJ, Nicolae DL, Gamazon ER, Im HK, Konkashbaev A, Pritchard J, Stevens M, Flutre T, Wen X, Dermitzakis ET, Lappalainen T, Guigo R, Monlong J, Sammeth M, Koller D, Battle A, Mostafavi S, McCarthy M, Rivas M, Maller J, Rusyn I, Nobel A, Wright F, Shabalina A, Feolo M, Sharopova N, Sturcke A, Paschal J, Anderson JM, Wilder EL, Derr LK, Green ED, Struwing JP, Temple G, Volpi S, Boyer JT, Thomson EJ, Guyer MS, Ng C, Abdallah A, Colantuoni D, Insel TR, Koester SE, Little AR, Bender PK, Lehner T, Yao Y, Compton CC, Vaught JB, Sawyer S, Lockhart NC, Demchok J, Moore HF. 2013. The Genotype-Tissue Expression (GTEx) project. *Nat Genet* **45**:580–585. doi:10.1038/ng.2653
- Luo SX, Huang J, Li Q, Mohammad H, Lee C-Y, Krishna K, Kok AM-Y, Tan YL, Lim JY, Li H, Yeow LY, Sun J, He M, Grandjean J, Sajikumar S, Han W, Fu Y. 2018. Regulation of feeding by somatostatin neurons in the tuberal nucleus. *Science* **361**:76–81. doi:10.1126/science.aar4983
- Luque RM, Cordoba-Chacon J, Pozo-Salas AI, Porteiro B, De Lecea L, Nogueiras R, Gahete MD, Castaño JP. 2016. Obesity- and gender-dependent role of endogenous somatostatin and cortistatin in the regulation of endocrine and metabolic homeostasis in mice. *Scientific Reports* **6**:37992. doi:10.1038/srep37992
- Martínez de Morentin PB, González-García I, Martins L, Lage R, Fernández-Mallo D, Martínez-Sánchez N, Ruíz-Pino F, Liu J, Morgan DA, Pinilla L, Gallego R, Saha AK, Kalsbeek A, Fliers E, Bisschop PH, Diéguez C, Nogueiras R, Rahmouni K, Tena-Sempere M, López M. 2014. Estradiol regulates brown adipose tissue thermogenesis via hypothalamic AMPK. *Cell metabolism* **20**:41–53. doi:10.1016/j.cmet.2014.03.031
- Maske CB, Jackson CM, Terrill SJ, Eckel LA, Williams DL. 2017. Estradiol modulates the anorexic response to central glucagon-like peptide 1. *Hormones and Behavior* **93**:109–117. doi:10.1016/j.yhbeh.2017.05.012
- Massa MG, Correa SM. 2020. Sexes on the brain: Sex as multiple biological variables in the neuronal control of feeding. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1866**:165840. doi:10.1016/j.bbadis.2020.165840
- Mathew H, Castracane VD, Mantzoros C. 2018. Adipose tissue and reproductive health. *Metabolism* **86**:18–32. doi:10.1016/J.METABOL.2017.11.006
- Mauvais-Jarvis F. 2017. Menopause, Estrogens, and Glucose Homeostasis in Women In: Mauvais-Jarvis F, editor. Sex and Gender Factors Affecting Metabolic Homeostasis, Diabetes and Obesity. pp. 217–227. doi:10.1007/978-3-319-70178-3
- Mickelsen LE, Bolisetty M, Chimileski BR, Fujita A, Beltrami EJ, Costanzo JT, Naparstek JR, Robson P, Jackson AC. 2019. Single-cell transcriptomic analysis of the lateral hypothalamic area reveals

- 652 molecularly distinct populations of inhibitory and excitatory neurons. *Nature Neuroscience*
- 653 **22**:642–656. doi:10.1038/s41593-019-0349-8
- 654 Mittelman-Smith MA, Williams H, Krajewski-Hall SJ, Lai J, Ciofi P, McMullen NT, Rance NE. 2012.
- 655 Arcuate Kisspeptin/Neurokinin B/Dynorphin (KNDy) Neurons Mediate the Estrogen
- 656 Suppression of Gonadotropin Secretion and Body Weight. *Endocrinology* **153**:2800–2812.
- 657 doi:10.1210/EN.2012-1045
- 658 Mohammad H, Senol E, Graf M, Lee C-Y, Li Q, Liu Q, Yeo XY, Wang M, Laskaratos A, Xu F, Luo
- 659 SX, Jung S, Augustine GJ, Fu Y. 2021. A neural circuit for excessive feeding driven by
- 660 environmental context in mice. *Nature Neuroscience* 2021 **24**:8 **24**:1132–1141.
- 661 doi:10.1038/s41593-021-00875-9
- 662 Muñoz MT, Argente J. 2002. Anorexia nervosa in female adolescents: endocrine and bone mineral
- 663 density disturbances. *European Journal of Endocrinology* **147**:275–286.
- 664 Musatov S, Chen W, Pfaff DW, Mobbs CV, Yang X-J, Clegg DJ, Kaplitt MG, Ogawa S. 2007.
- 665 Silencing of estrogen receptor alpha in the ventromedial nucleus of hypothalamus leads to
- 666 metabolic syndrome. *Proceedings of the National Academy of Sciences of the United States of*
- 667 *America* **104**:2501–6. doi:10.1073/pnas.0610787104
- 668 Naaz A, Zakroczyński M, Heine P, Taylor J, Saunders P, Lubahn D, Cooke PS. 2002. Effect of
- 669 Ovariectomy on Adipose Tissue of Mice in the Absence of Estrogen Receptor Alpha (ER α): a
- 670 Potential Role for Estrogen Receptor Beta (ER β). *Hormone and Metabolic Research* **34**:758–
- 671 763. doi:10.1055/s-2002-38259
- 672 Narita K, Murata T, Matsuoka S. 2016. The ventromedial hypothalamus oxytocin induces locomotor
- 673 behavior regulated by estrogen. *Physiology & Behavior* **164**:107–112.
- 674 doi:10.1016/j.physbeh.2016.05.047
- 675 Nelson JF, Gosden RG, Felicio LS. 1985. Effect of Dietary Restriction on Estrous Cyclicity and
- 676 Follicular Reserves in Aging C57BL/6J Mice. *Biology of Reproduction* **32**:515–522.
- 677 doi:10.1095/BIOLREPROD32.3.515
- 678 Nogueiras R, Gualillo O, Caminos JE, Casanueva FF, Diéguez C. 2003. Regulation of resistin by
- 679 gonadal, thyroid hormone, and nutritional status. *Obesity Research* **11**:408–414.
- 680 doi:10.1038/oby.2003.55
- 681 Oldfield BJ, Mirabella PN, Stefanidis A. 2016. Neuroanatomy of Feeding Pathways In: Dickson SL,
- 682 Mercer JG, editors. *Neuroendocrinology of Appetite*. pp. 1–23.
- 683 Padilla SL, Qiu J, Nestor CC, Zhang C, Smith AW, Whiddon BB, Rønnekleiv OK, Kelly MJ, Palmiter
- 684 RD. 2017. AgRP to Kiss1 neuron signaling links nutritional state and fertility. *Proceedings of the*
- 685 *National Academy of Sciences of the United States of America* **114**:2413–2418.
- 686 doi:10.1073/pnas.1621065114
- 687 Painson J-C, Tannenbaum GS. 1991. Sexual Dimorphism of Somatostatin and Growth Hormone-
- 688 Releasing Factor Signaling in the Control of Pulsatile Growth Hormone Secretion in the Rat*.
- 689 *Endocrinology* **128**:2858–2866. doi:10.1210/endo-128-6-2858
- 690 Palmer BF, Clegg DJ. 2015. The sexual dimorphism of obesity. *Molecular and Cellular Endocrinology*
- 691 **402**:113–119. doi:10.1016/J.MCE.2014.11.029
- 692 Petersen S. 1976. The temporal pattern of feeding over the oestrous cycle of the mouse. *Animal*
- 693 *Behaviour* **24**:939–955. doi:10.1016/S0003-3472(76)80023-1
- 694 Pfaff D, Keiner M. 1973. Atlas of estradiol-concentrating cells in the central nervous system of the
- 695 female rat. *Journal of Comparative Neurology* **151**:121–157. doi:10.1002/cne.901510204
- 696 Rodriguez-Pacheco F, Martinez-Fuentes AJ, Tovar S, Pinilla L, Tena-Sempere M, Dieguez C, Castaño
- 697 JP, Malagon MM. 2007. Regulation of Pituitary Cell Function by Adiponectin. *Endocrinology*
- 698 **148**:401–410. doi:10.1210/EN.2006-1019
- 699 Roepke TA, Bosch MA, Rick EA, Lee B, Wagner EJ, Seidlova-Wuttke D, Wuttke W, Scanlan TS,
- 700 Rønnekleiv OK, Kelly MJ. 2010. Contribution of a Membrane Estrogen Receptor to the

- 701 Estrogenic Regulation of Body Temperature and Energy Homeostasis. *Endocrinology* **151**:4926–
702 4937. doi:10.1210/en.2010-0573
- 703 Roepke TA, Malyala A, Bosch MA, Kelly MJ, Rønnekleiv OK. 2007. Estrogen Regulation of Genes
704 Important for K⁺ Channel Signaling in the Arcuate Nucleus. *Endocrinology* **148**:4937–4951.
705 doi:10.1210/en.2007-0605
- 706 Rooks C, Bennet T, Bartness TJ, Harris RBS. 2004. Compensation for an increase in body fat caused by
707 donor transplants into mice. *Am J Physiol Regul Integr Comp Physiol* **286**:1149–1155.
708 doi:10.1152/ajpregu.00634.2003.-Rodents
- 709 Sanchez-Alavez M, Alboni S, Conti B. 2011. Sex- and age-specific differences in core body temperature
710 of C57Bl/6 mice. *Age (Dordrecht, Netherlands)* **33**:89–99. doi:10.1007/s11357-010-9164-6
- 711 Santollo J, Torregrossa A-M, Eckel LA. 2011. Estradiol acts in the medial preoptic area, arcuate
712 nucleus, and dorsal raphe nucleus to reduce food intake in ovariectomized rats. *Hormones and*
713 *Behavior* **60**:86–93. doi:10.1016/J.YHBEH.2011.03.009
- 714 Seldin MM, Koplev S, Rajbhandari P, Vergnes L, Rosenberg GM, Meng Y, Pan C, Phuong TMN,
715 Gharakhanian R, Che N, Mäkinen S, Shih DM, Civelek M, Parks BW, Kim ED, Norheim F,
716 Chella Krishnan K, Hasin-Brumshtein Y, Mehrabian M, Laakso M, Drevon CA, Koistinen HA,
717 Tontonoz P, Reue K, Cantor RM, Björkegren JLM, Lusis AJ. 2018. A Strategy for Discovery of
718 Endocrine Interactions with Application to Whole-Body Metabolism. *Cell Metabolism* **27**:1138-
719 1155.e6. doi:10.1016/J.CMET.2018.03.015/ATTACHMENT/1AA459EB-A64D-4330-A459-
720 F54C4F7105DD/MMC2.XLSX
- 721 Simerly RB, Swanson LW, Chang C, Muramatsu M. 1990. Distribution of androgen and estrogen
722 receptor mRNA-containing cells in the rat brain: An in situ hybridization study. *The Journal of*
723 *Comparative Neurology* **294**:76–95. doi:10.1002/cne.902940107
- 724 Smith JT, Cunningham MJ, Rissman EF, Clifton DK, Steiner RA. 2005. Regulation of Kiss1 Gene
725 Expression in the Brain of the Female Mouse. *Endocrinology* **146**:3686–3692.
726 doi:10.1210/EN.2005-0488
- 727 Solorzano CMB, McCartney CR. 2010. Obesity and the pubertal transition in girls and boys.
728 *Reproduction* **140**:399–410. doi:10.1530/REP-10-0119
- 729 Steiner M, Katz Richard J, Carroll Bernard J, Katz R J, Carroll B J. 1982. Detailed Analysis of Estrous-
730 Related Changes in Wheel Running and Self-Stimulation, Physiology & Behavior. Pergamon
731 Press and Brain Research Publ.
- 732 Stengel Andreas, Coskun T, Goebel M, Wang L, Craft L, Alsina-Fernandez J, Rivier J, Tache Y. 2010a.
733 Central Injection of the Stable Somatostatin Analog ODT8-SST Induces a Somatostatin 2
734 Receptor-Mediated Orexigenic Effect: Role of Neuropeptide Y and Opioid Signaling Pathways
735 in Rats. *Endocrinology* **151**:4224–4235. doi:10.1210/en.2010-0195
- 736 Stengel A, Coskun T, Goebel-Stengel M, Craft LS, Alsina-Fernandez J, Wang L, Rivier J, Taché Y.
737 2011. Chronic injection of pansomatostatin agonist ODT8-SST differentially modulates food
738 intake and decreases body weight gain in lean and diet-induced obese rats. *Regulatory Peptides*
739 **167**:201–208. doi:10.1016/J.REGPEP.2011.01.006
- 740 Stengel Andreas, Goebel M, Wang L, Rivier J, Kobelt P, Mönnikes H, Taché Y. 2010b. Activation of
741 brain somatostatin 2 receptors stimulates feeding in mice: analysis of food intake microstructure.
742 *Physiology & behavior* **101**:614–22. doi:10.1016/j.physbeh.2010.09.009
- 743 Stengel A, Goebel M, Wang L, Rivier J, Kobelt P, Monnikes H, Taché Y. 2010. Selective central
744 activation of somatostatin receptor 2 increases food intake, grooming behavior and rectal
745 temperature in rats. *Journal of physiology and pharmacology : an official journal of the Polish*
746 *Physiological Society* **61**:399–407.
- 747 Stengel A, Karasawa H, Taché Y. 2015. The role of brain somatostatin receptor 2 in the regulation of
748 feeding and drinking behavior. *Hormones and Behavior* **73**:15–22.
749 doi:10.1016/J.YHBEH.2015.05.009

- Stengel A, Rivier J, Taché Y. 2013. Central actions of somatostatin-28 and oligosomatostatin agonists to prevent components of the endocrine, autonomic and visceral responses to stress through interaction with different somatostatin receptor subtypes. *Current pharmaceutical design* **19**:98–105.
- Stincic Todd L, Grachev P, Bosch MA, Rønnekleiv OK, Kelly MJ. 2018. Estradiol Drives the Anorexigenic Activity of Proopiomelanocortin Neurons in Female Mice. *eNeuro* **5**:ENEURO.0103-18.2018. doi:10.1523/ENEURO.0103-18.2018
- Stincic Todd L., Rønnekleiv OK, Kelly MJ. 2018. Diverse actions of estradiol on anorexigenic and orexigenic hypothalamic arcuate neurons. *Hormones and Behavior* **104**:146–155. doi:10.1016/J.YHBEH.2018.04.001
- Tachibana T, Cline MA, Sugahara K, Ueda H, Hiramatsu K. 2009. Central administration of somatostatin stimulates feeding behavior in chicks. *General and Comparative Endocrinology* **161**:354–359. doi:10.1016/J.YGCEN.2009.01.022
- Tran TT, Yamamoto Y, Gesta S, Kahn CR. 2008. Beneficial Effects of Subcutaneous Fat Transplantation on Metabolism. *Cell Metabolism* **7**:410–420. doi:10.1016/j.cmet.2008.04.004
- Tsatsanis C, Dermitzaki E, Avgoustinaki P, Malliaraki N, Mytaras V, Margioris AN. 2015. The impact of adipose tissue-derived factors on the hypothalamic-pituitary-gonadal (HPG) axis. *Hormones* **14**:549–562. doi:10.14310/HORM.2002.1649
- van Veen & Kammel J& LG, Bunda PC, Shum M, Reid MS, Massa MG, Arneson DV, Park JW, Zhang Z, Joseph AM, Hrcir H, Liesa M, Arnold AP, Yang X, Correa SM. 2020. Hypothalamic oestrogen receptor alpha establishes a sexually dimorphic regulatory node of energy expenditure. *Nature Metabolism* **2**:351–363. doi:10.1038/s42255-020-0189-6
- Velez LM, Van C, Moore T, Zhou Z, Johnson C, Hevener AL, Seldin MM. 2022. Genetic variation of putative myokine signaling is dominated by biological sex and sex hormones. *eLife* **11**:e76887. doi:10.7554/eLife.76887
- Witte MM, Resuehr D, Chandler AR, Mehle AK, Overton JM. 2010. Female mice and rats exhibit species-specific metabolic and behavioral responses to ovariectomy. *General and Comparative Endocrinology* **166**:520–528. doi:10.1016/j.ygcen.2010.01.006
- Wright H, Aylwin CF, Toro CA, Ojeda SR, Lomniczi A. 2021. Polycomb represses a gene network controlling puberty via modulation of histone demethylase Kdm6b expression. *Scientific Reports* **2021 11:1** **11**:1–17. doi:10.1038/s41598-021-81689-4
- Xu Y, Nedungadi TP, Zhu L, Sobhani N, Irani BG, Davis KE, Zhang X, Zou F, Gent LM, Hahner LD, Khan SA, Elias CF, Elmquist JK, Clegg DJ. 2011. Distinct hypothalamic neurons mediate estrogenic effects on energy homeostasis and reproduction. *Cell metabolism* **14**:453–65. doi:10.1016/j.cmet.2011.08.009
- Yang CF, Chiang MC, Gray DC, Prabhakaran M, Alvarado M, Juntti SA, Unger EK, Wells JA, Shah NM. 2013. Sexually Dimorphic Neurons in the Ventromedial Hypothalamus Govern Mating in Both Sexes and Aggression in Males. *Cell* **153**:896–909. doi:10.1016/j.cell.2013.04.017
- Yang Z-H, Miyahara H, Takeo J, Katayama M. 2012. Diet high in fat and sucrose induces rapid onset of obesity-related metabolic syndrome partly through rapid response of genes involved in lipogenesis, insulin signalling and inflammation in mice. *Diabetology & Metabolic Syndrome* **4**:32. doi:10.1186/1758-5996-4-32
- Zhang Z, Divittorio JR, Joseph AM, Correa SM. 2021. The Effects of Estrogens on Neural Circuits That Control Temperature. *Endocrinology* **162**:1–12. doi:10.1210/ENDOCR/BQAB087