

1 **Conservation and Expansion of Transcriptional Factor Repertoire in the**
2 ***Fusarium oxysporum* Species Complex**

3

4 Houlin Yu¹, He Yang¹, Sajeet Haridas², Richard D. Hayes², Hunter Lynch¹,
5 Sawyer Andersen¹, Gengtan Li¹, Domingo Martínez-Soto¹, Shira Milo-Cochavi¹,
6 Dilay Hazal Ayhan¹, Yong Zhang¹, Igor V. Grigoriev^{2,3}, Li-Jun Ma^{1†}

7

8 ¹Department of Biochemistry and Molecular Biology, University of Massachusetts
9 Amherst, Amherst, MA 01003, USA.

10 ²Department of Energy Joint Genome Institute, Lawrence Berkeley National
11 Laboratory, University of California Berkeley, Berkeley, CA 94720, USA.

12 ³Department of Plant and Microbial Biology, University of California Berkeley,
13 Berkeley, CA 94598, USA

14

15 [†]Correspondence: Li-Jun Ma (lijun@biochem.umass.edu).

16

17 Running Head: TFome Analysis of *Fusarium oxysporum* Species Complex

18

19

ABSTRACT

The *Fusarium oxysporum* species complex (FOSC) includes both plant and human pathogens that cause devastating plant vascular wilt diseases and threaten public health. Each *F. oxysporum* genome comprises core chromosomes (CCs) for housekeeping functions and accessory chromosomes (ACs) that contribute to host-specific adaptation. This study inspected global transcription factor profiles (TFomes) and their potential roles in coordinating CCs and ACs functions to accomplish host-specific pathogenicity. Remarkably, we found a clear positive correlation between the sizes of TFome and proteome of an organism, and FOSC TFomes are larger due to the acquisition of ACs. Among a total of 48 classified TF families, 14 families involved in transcription/translation regulations and cell cycle controls are highly conserved. Among 30 FOSC expanded families, Zn2-C6 and Znf_C2H2 are most significantly expanded to 671 and 167 genes per family, including well-characterized homologs of Ftf1 (Zn2-C6) and PacC (Znf_C2H2) involved in host-specific interactions. Manual curation of characterized TFs increased the TFome repertoires by 3%, including a disordered protein Ren1. Expression profiles revealed a steady expression of conserved TF families and specific activation of AC TFs. Functional characterization of these TFs could enhance our understanding of transcriptional regulation involved in FOSC cross-kingdom interactions, disentangle species-specific adaptation, and identify targets to combat diverse diseases caused by this group of fungal pathogens.

KEYWORDS

Fusarium oxysporum species complex; transcription factors; TFome; accessory chromosome; conservation; expansion

INTRODUCTION

The fungal species complex of *Fusarium oxysporum* (FOSC) has been used as a model to study cross-kingdom fungal pathogenesis. Members within FOSC can cause devastating fusarium wilt diseases among economically important crops (Ma et al. 2013; Ma 2014; Michielse and Rep 2009; Ploetz 2015; Edel-Hermann and Lecomte 2019; Pegg et al. 2019; Yang et al. 2020; Dean et al. 2012; Rahman et al. 2021; Viljoen et al. 2020; Halpern et al. 2018) and is listed among the top five most important plant pathogens that have a direct impact on the global economy and food security (Dean et al. 2012). With strong host specificity, plant pathogenic *F. oxysporum* strains are further grouped as *formae speciales* (Armstrong and Armstrong 1981). For instance, tomato pathogens are named *F. oxysporum* f.sp. *lycopersici*, cotton pathogens *F. oxysporum* f.sp. *vasinfectum* (Halpern et al. 2018), and banana pathogen *F. oxysporum* f.sp. *cubense* (Viljoen et al. 2020). Recently, members within FOSC have also been reported to be responsible for fusariosis, the top emerging opportunistic mycosis (Ma et al. 2013; Yang et al. 2020), and fusarium keratitis, one of the major causes of cornea infections in the developing world and the leading cause of blindness among fungal keratitis patients (Kredics et al. 2015; Hassan et al. 2016).

Comparative genomics studies on this cross-kingdom pathogen revealed that the FOSC genomes, both human and plant pathogens, are compartmented into two components: the core chromosomes (CCs) and accessory chromosomes (ACs). While CCs are conserved and vertically inherited to execute essential housekeeping functions, horizontally transmitted ACs are lineage- or strain-specific and related to fungal adaptation and pathogenicity, conferred by virulent factors such as SIX (Secreted in Xylem) proteins (Ma et al. 2013; Yang et al. 2020; Rep et al. 2004; Yu et al. 2023). ACs and CCs must coordinate their gene expression to coexist within the same genome.

A few characterized transcription factors (TFs) coordinate the crosstalk between CCs and ACs, two compartments. One intriguing example is the cross-regulation

among *F. oxysporum* transcription factors Sge1 (SIX Gene Expression 1), Ftf1, and effector genes. Sge1 is a highly conserved, CC-encoding TF. By name definition, Sge1 regulates the expression of SIX proteins (Michielse et al. 2009; van der Does et al. 2016). AC-encoding Ftf1 proteins (Ftf1 and its AC homologs) and a CC-encoding Ftf2 (Ftf1 CC homolog) are reported in the reference genome of *F. oxysporum* f.sp. *lycopersici* FoI4287 (van der Does et al. 2016). Constitutive expression of either *Ftf1* genes or *Ftf2* induced the expression of effector genes (van der Does et al. 2016). Furthermore, It was documented that DNA binding sites of Sge1 and Ftf1 are enriched among the cis-regulatory elements of *in planta* transcriptionally up-regulated genes (van der Does et al. 2016). Another CCs and ACs cross-talking example is the alkaline pH-responsive transcription factor PacC/Rim1p reported in *F. oxysporum* clinical strains (Zhang et al. 2020). In addition to the full-length *PacC* ortholog (*PacC_O*), located on a CC, the clinical isolate NRRL32931 genome encodes three truncated *PacC* homologs, named *PacC_a*, *PacC_b*, and *PacC_c* in ACs (Zhang et al. 2020).

To thoroughly understand the coordination of the crosstalk between genome compartments and their contribution to the cross-kingdom fungal pathogenesis, this study compared the repertoire of TFs (*i.e.*, TFome) among 15 *F. oxysporum* and 15 other ascomycete fungal genomes, which was organized into 48 families based on the InterPro classification of proteins. Remarkably, we discovered a strong positive correlation ($y = 0.07264x - 190.9$, $r^2 = 0.9361$) between the number of genes (x) and TFome size (y) of an organism. Primarily due to the acquisition of ACs, we observed increased TFome sizes among FOSC genomes. Fourteen out of 48 families involved in transcription/translation regulations and cell cycle controls are highly conserved. Thirty, accounting for $\frac{3}{4}$ of all families, are expanded in various degrees among FOSC genomes. Unique TF expansions driven by ACs include members of Zn2-C6 fungal-type (Zn2-C6) and Zinc Finger C2H2 (Znf_C2H2) families. This comparative study first highlights the conserved regulatory mechanisms of *F. oxysporum*, which are essential for variability and plant colonization. With the foundation established by functional conservation,

this study further emphasizes potential modifications of existing regulatory pathways by acquiring additional TFs. In combination with existing expression data, this study may provide clues to the fine-tuning of networks in the environmental adaptation of this group of diverse organisms to engage in complex cross-kingdom interactions with different hosts.

MATERIALS AND METHODS

Generation of fungal TFomes

The annotation pipeline is briefly summarized in Figure S1A-B. The fungal proteomes of 30 strains were downloaded from the JGI MycoCosm portal (Grigoriev et al. 2014). Protein annotation was performed using InterProScan/5.38-76.0 (<https://www.ebi.ac.uk/interpro/search/sequence/>) (Jones et al. 2014). Annotations of proteins putatively serve as TFs were filtered out using a table containing InterPro terms related to transcriptional regulatory functions summarized by literature (Park et al. 2008; Shelest 2017), with further addition by manual curation (Table S1). Orthologous analysis to probe orthologs of functionally validated TFs (Table S3-4 and Table 3) in *Fusarium* was done with OrthoFinder 2.5.4 (<https://github.com/davidemms/OrthoFinder>) (Emms and Kelly 2019).

RNA-seq analysis

The RNA-seq datasets were previously described (Guo et al. 2021; Redkar et al. 2022) and deposited by those authors to the NCBI Short Read Archive with accession number GSE87352 and to the ArrayExpress database at EMBL-EBI (www.ebi.ac.uk/arrayexpress) under accession number E-MTAB-10597, respectively. For data reprocessing, reads were mapped to reference genomes of *Arabidopsis* [annotation version Araport11 (Cheng et al. 2017)], *Fo5176* (Fokkens et al. 2021), *Fo47* (Wang et al. 2020) and *Fo4287* (Ma et al. 2010) using HISAT2 version 2.0.5 (Kim et al. 2019). Mapped reads were used to quantify the transcriptome by StringTie version 1.3.4 (Pertea et al. 2015), at which step TPM

(transcript per million) normalization was applied. Normalized read counts were first averaged per condition and then transformed by log2 (normalized read count + 1) and Z-scaled, then visualized in heatmap (version 1.0.12).

Genome partition

The genome partition results for chromosome-level assemblies were retrieved from previous reports for Fo4287 (Ma et al. 2010), Fo115 (Zhang 2019), Fo5176 (Fokkens et al. 2021), and Fo47 (Wang et al. 2020). Fo47 has a clear genome partition with 11 core chromosomes and one accessory chromosome, therefore serving as the reference for the genome partition of other *F. oxysporum* genomes. mummer/3.22 was applied to align scaffolds of genome assemblies against 11 core chromosomes of the reference genome Fo47 using default parameters. The scaffolds aligned to the core chromosomes of Fo47 with a coverage larger than 5% were annotated as core scaffolds. The rest of the scaffolds were partitioned as accessory scaffolds. Genes residing on core and accessory scaffolds were annotated as core and accessory genes, respectively.

Phylogenetics analysis

Protein sequences were aligned via MAFFT/7.313 (Katoh and Standley 2013). Then the iqtree/1.6.3 (Minh et al. 2020; Nguyen et al. 2015) was run on the sequence alignment to generate the phylogeny (by maximum likelihood method and bootstrapped using 1000 replicates) (Hoang et al. 2018) and was then visualized via the Interactive Tree of Life (Letunic and Bork 2021), producing the phylogram. OrthoFinder 2.5.4 (Emms and Kelly 2019) was used for orthogroup determination. To build a species phylogram, randomly selected 500 conserved proteins (single-copy orthologs) were aligned first. Then the alignment was concatenated, and phylogeny was determined and visualized using the above methods.

RESULTS

1. FOSC TFome expansion resulted from the acquisition of ACs




Figure 1. Phylogeny of fungal genomes included in this study. Both left and right phylograms were constructed by concatenated alignment of randomly selected 500 single-copy orthologous proteins, followed by the maximum likelihood method with 1000 bootstraps. Left shows a phylogram of FOSC (represented by the reference genome Fo4287) together with the other 15 ascomycetes. The right shows a phylogram of members within FOSC, rooted by *F. verticillioides* (not shown). Only bootstrap values not equal to 100 are shown.

We compared 30 ascomycete fungal genomes (Figure 1 and Table 1), including 15 strains within the FOSC, nine sister species close to *F. oxysporum*, two yeast genomes (*Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*), four other filamentous fungal species (*Neurospora crassa*, *Aspergillus nidulans*, *Aspergillus acristatus*, and *Magnaporthe oryzae*). To maintain consistency, the protein sequences for all these genomes were retrieved from the MycoCosm portal (Grigoriev et al. 2014).

Table 1. Fungal genomes used in this study

Fungal species or strains	MycoCosm identifier	Genome Size (MB)	No. of genes	TFome Size	Host	Reference
<i>Saccharomyces cerevisiae</i>	Sacce1	12.07	6575	284		(Goffeau et al. 1996)
<i>Schizosaccharomyces pombe</i>	Schpo1	12.61	5134	228		(Wood et al. 2002)
<i>Aspergillus nidulans</i>	Aspnid1	30.48	10680	635		(Galagan et al. 2005)
<i>Aspergillus acristatus</i>	Aspacri1	32.59	11221	666		(Vesth et al. 2018)
<i>Neurospora crassa</i>	Neucr2	41.04	9730	447		(Galagan et al. 2003)
<i>Magnaporthe oryzae</i>	Magor1	40.49	12673	520	Rice	(Dean et al. 2005)
<i>Fusarium solani</i>	Fusso1	52.93	17656	1137	broad hosts	(Mesny et al. 2021)
<i>F. pseudograminearum</i>	Fusps1	36.33	12395	627	Wheat	(Gardiner et al. 2012)
<i>F. graminearum</i>	Fusgr1	36.45	13321	608	Wheat	(Cuomo et al. 2007)
<i>F. venenatum</i>	Fusven1	37.45	12845	802		(Mesny et al. 2021)
<i>F. tricinctum</i>	Fustr1	43.69	14106	925	Broad hosts	(Mesny et al. 2021)
<i>F. verticillioides</i>	Fusve2	41.78	15869	917	Corn	(Ma et al. 2010)
<i>F. fujikuroi</i>	Fusfu1	43.83	14813	901	Broad hosts	(Wiemann et al. 2013)
<i>F. redolens</i>	Fusre1	52.56	17051	1098	Broad hosts	(Mesny et al. 2021)
<i>F. commune</i>	Fusco1	48.37	15731	1012	Broad hosts	(Mesny et al. 2021)
<i>F. oxysporum</i> f.sp. <i>cubense</i> (II5)	FoxII5	49.43	16048	1047	Banana	(Zhang 2019)
<i>F. oxysporum</i> f. sp. <i>radicis-lycopersici</i> (CL57)	Fusoxrad1	49.36	18238	1151	Tomato	(Delulio et al. 2018)
<i>F. oxysporum</i> Fo47 (Fo47)	FusoxFo47_2	50.36	16207	1082		(Wang et al. 2020)
<i>F. oxysporum</i> f. sp. <i>lycopersici</i> (MN25)	Fusoxlyc1	48.64	17931	1119	Tomato	(Delulio et al. 2018)
<i>F. oxysporum</i> NRRL26365 (NRRL26365)	Fox26365_1	48.46	16047	1036	Human	(Yang 2020)
<i>F. oxysporum</i> f. sp. <i>melonis</i> (FoMelon)	Fusoxmel1	54.03	19661	1219	Melon	(Ma et al. 2014)
<i>F. oxysporum</i> f. sp. <i>lycopersici</i> (Fol4287)	Fusox2	61.36	20925	1292	Tomato	(Ma et al. 2010)
<i>F. oxysporum</i> NRRL32931 (NRRL32931)	Fusox32931	47.91	17280	1072	Human	(Zhang et al. 2020)
<i>F. oxysporum</i> MRL8996 (MRL8996)	FoxMRL8996	50.07	16631	1057	Human	(Zhang et al. 2020)
<i>F. oxysporum</i> f. sp. <i>matthiolae</i> (PHW726)	FoxPHW726_1	57.22	17996	1157	Brassica	(Yu et al. 2020)
<i>F. oxysporum</i> f. sp. <i>vasinfectum</i> (FoCotton)	Fusoxvas1	52.91	19143	1189	Cotton	(Delulio et al. 2018)
<i>F. oxysporum</i> f. sp. <i>pisi</i> (HDV247)	Fusoxpis1	55.19	19623	1229	Pea	(Williams et al. 2016)
<i>F. oxysporum</i> f. sp. <i>raphani</i> (PHW815)	Fusoxrap1	53.5	19306	1132	Brassica	(Delulio et al. 2018)
<i>F. oxysporum</i> f. sp. <i>conglutinans</i> (PHW808)	Fusoxcon1	53.58	19854	1142	Brassica	(Delulio et al. 2018)
<i>F. oxysporum</i> Fo5176 (Fo5176)	FoxFo5176	67.98	19130	1236	Arabidopsis	(Fokkens et al. 2021)

177

178 To have a comprehensive TFome annotation, we started with reported
179 InterProScan (IPR) terms associated with fungal transcriptional regulation (Park
180 et al. 2008; Shelest 2017) and curated a mapping with updated IPR classification
181 (interproscan version: 5.38-76.0) (Blum et al. 2021). In addition, we searched the
182 IPR classification of protein families and obtained all other terms related to the
183 transcriptional regulation activity. This resulted in 234 TF-related IPR terms
184 (Table S1). Since most of the terms are initially defined in the mammalian
185 systems, it was not surprising to see that overall, our fungal genomes are only
186 associated with 71 IPR terms out of the total 234 TF-related IPR terms (Table
187 S1, Materials and Methods, and Figure S1A-B for annotation pipeline). After
188 filtering out 13 and 10 terms for redundancy (two terms describing the identical

189 domain) and minimal presentation (< 4 among the 30 genomes), respectively,
 190 this comparative TFome study focused on the rest 48 IPR terms, which
 191 represented a total of 27967 TFs (Table S1-S2). Notably, 12 out of 48 terms
 192 were not reported to be affiliated with fungal transcriptional regulation by either
 193 Park et al. 2008 or Shelest 2017 (Table S1), adding values to our manual IPR
 194 term search.

195 Comparing the total number of genes in a genome (x) and the total number of
 196 TFs within that genome (y), we observed a strong positive correlation ($y =$
 197 $0.07264x - 190.9$, $r^2 = 0.9361$) (Figure 2A). Among all genomes included in this
 198 study, FOSC TFomes are the largest, with an average of 1144 TFs per genome
 199 (Figure 2A, Table 1). After partitioning each FOSC genome into core and
 200 accessory regions (see Materials and Methods for details), we observed a
 201 positive correlation between the number of TFs encoded in the accessory
 202 chromosomal region of each strain (defined as accessory TFs hereafter) with the
 203 size of accessory genomes (Mb) ($y = 17.239x + 3.553$) (Figure S2), suggesting
 204 that accessory chromosomes contribute directly to the expanded TFome.

205




Figure 2. TFome conservation and variation among ascomycete fungi: baseline description. (A) There is a positive correlation between the number of genes and TFome size of an organism. JGI fungal genome identifiers were used as labels. (B) Histogram illustrates the distribution of expansion indexes among different families. (C) Average number of TFs of two most drastically expanded families (Znf_C2H2 and Zn2-C6) within each genome set. Genome Set 1 (G1) includes two yeast genomes (*S. cerevisiae* and *S. pombe*). Genome Set 2 (G2) includes four filamentous fungal species (*N. crassa*, *A. nidulans*, *A. acristatulus*, and *M. oryzae*). Genome Set 3 (G3) includes nine sister species close to *F. oxysporum*. Genome Set 4 (G4) includes 15 FOSC genomes.

206 To understand genome regulation among FO SC, we developed an expansion
207 index score using two yeast lineages as the baseline (El_y):

$$208 \quad El_y = \frac{\text{Average number of TFs in FO SC} + 1}{\text{Average number of TFs in yeasts} + 1}$$

209

Table 2. Expansion Index (El_y) of 48 TF families
Asterisk indicates the families without a presence in yeasts

IPR	Term	El_y
Group 1		
IPR000814	TBP	1
IPR003228	TFIID_TAF12	1
IPR004595	TFIIH_C1-like	1
IPR006809	TAFII28	1
IPR042225	Ncb2	1
IPR008570	Vps25	1
IPR008895	Vps72/YL1	1
IPR007196	CNOT1	1
IPR005612	CBF	1
IPR001289	NFYA	1
IPR018004	APSES-type HTH	1
IPR003150	RFX	1
IPR033896	MADS_MEF2-like	1
IPR018501	DDT	1
Group 2		
IPR006856	MATalpha_HMGbox	0.8
IPR039515	NOT4	0.9
IPR033897	MADS_SRF-like	0.95
IPR000232	HSF	0.98
Group 3		
IPR003163	Tscript_reg_HTH_APSES-type	1.04
IPR001766	Fork_head	1.05
IPR011016	Znf_RING-CH	1.11
IPR001965	Znf_PHD	1.11
IPR009071	HMG_box	1.12
IPR004181	Znf_MIZ	1.24
IPR001606	ARID	1.25
IPR000679	Znf_GATA	1.3
IPR001005	SANT/Myb	1.32
IPR000818	TEA/ATTS	1.33
IPR003120	Ste12	1.33
IPR003958	CBFA_NFYB	1.35
IPR001083	Cu_fist	1.37
IPR000967	Znf_NFX1	1.4
IPR006565	Bromodomain	1.52
IPR001387	Cro/C1-type_HTH	1.6
IPR001841	Znf_RING	1.64
IPR000571	Znf_CCCH	1.74
IPR001878	Znf_CCHC	1.83
IPR010666	Znf_GRF	2
IPR018060	HTH_AraC*	2
IPR001356	Homeobox	2.28
IPR007604	CP2*	2.73
IPR007396	PAI2	3.42
IPR024061	NDT80	3.47
IPR011598	bHLH	3.48
IPR007889	HTH_Psq*	3.53
IPR013087	Znf_C2H2	4.15
IPR004827	bZIP	5.8
IPR001138	Zn2-C6	15.09

211

Based on this index value, we classified TF families into three major groups (Table 2, Table S1). Group 1 contains 14 TF families with an expansion score of 1, indicating high conservation. Group 2 includes four families with an index score of less than 1, reflecting some level of gene family contraction. Group 3 contains 30 families with an expansion index greater than 1, indicating gene expansion.

2. Conserved TF families that are primarily associated with general/global transcription factors

About 30% of the TF families, fourteen, are associated with strong orthologous conservation in all genomes we included in this study (Figure 2B; Table 2; Table S1). Because most of these conserved TF families are single-copied TF families, these 30% conserved TF families only account for less than 2% of the total TFomes. Based on a detailed study on *S. cerevisiae* and other model organisms, these TF families are involved in transcription/translation regulation and cell cycle controls.

2.1. Transcription/Translation regulation

Either TF families are related to transcription initiation and elongation, including TATA box-binding protein (TBP), TBP-associated factors (TAFs), and RNA polymerase II elongation regulator Vps25. CCAAT-Binding Factors (CBFs) are related to ribosomal biogenesis. These families overall play conserved roles in general transcriptional and translational regulation across *Ascomycota*.

Transcription initiation TBP is one of the most conserved TF families, TBP binds directly to the TATA box to define the transcription start and initiate transcription facilitated by all three RNA polymerases. The function of TBP is so conserved as the yeast homolog can complement *TBP* mutations in humans (Yamaguchi et al. 2001; Roberts and Winston 1996).

Transcription positive/negative regulators, including **TAF12** and **TAF128**, are parts of the transcription factor TF_{II}D complex. Interacting with TBP, TAFs form

the TF_{II}D complex and positively participate in the assembly of the transcription preinitiation complex (Green 2000). Similarly, **TF_{II}H** works synergistically with TF_{II}D to promote the transcription (Fribourg et al. 2000). In contrast, Negative cofactor 2 (**Ncb2**) inhibits the preinitiation complex assembly (Goppelt et al. 1996). Other factors include the **CNOT1**, a global regulator involved in transcription initiation and RNA degradation (Chalabi Hagkarim and Grand 2020), and **Vps72/YL1** that contributes to transcriptional regulation through chromatin remodeling as reported in the yeast (Liang et al. 2016; Latrick et al. 2016).

Transcription elongation: **Vps25** is a subunit of the ESCRT-II complex, which binds to RNA polymerase II elongation factor to exert transcriptional control in mammalian systems (Kamura et al. 2001).

Translational regulation: CCAAT box is a common cis-acting element found in the promoter and enhancer regions of genes in the eukaryotes (Vuorio et al. 1990; Becker et al. 1991). **CBFs** are necessary for the 60S ribosomal subunit biogenesis and therefore involved in the translational control (Milkereit et al. 2001; Fromont-Racine et al. 2003; Edskes et al. 1998). This family, including Noc3, Noc4, and Mak21 in *S. cerevisiae*, has three members in each genome, and a clear single-copy orthologous relationship can be observed for each member (Figure S3A).

2.2. Cell cycle control

Five TF families are related to cell cycle control, including cell cycle progression, DNA repair, and machinery/cell integrity maintenance.

APSES-type HTH represents a family of fungal TFs involved in cell-cycle control and is crucial to the development (Xin et al. 2020). Every genome maintains four copies of genes encoding APSES-type HTH (Figure S3B), and they form single-copy orthologs in all genomes except yeasts. Genes in Clade 1, including StuA homologs, are targets of the cyclic AMP (cAMP)-dependent protein kinase A (PKA) signal transduction pathway and were reported to be involved in dimorphic switch (Pan and Heitman 2000; Gimeno and Fink 1994), fungal spore development and the production of secondary metabolites (Lysøe et al. 2011).

Genes in Clade 2 and Clade 3 include *S. cerevisiae* Swi4 and Swi6, which form a protein complex and regulate genes essential during cell cycle progression from G1 to S phase (Koch et al. 1993), as well as meiosis (Son et al. 2016b). Genes in Clade 4 include homologs of *S. pombe* Bqt4 that connect telomeres to the nuclear envelope (Chikashige et al. 2009). Since this family of TFs is highly conserved across ascomycetes, similar functions can be proposed in *F. oxysporum*.

DTT, represented by the *S. cerevisiae* homolog Itc1, is a subunit of ATP-dependent Isw2p-Itc1p chromatin remodeling complex and is required for repression of early meiotic gene expression during the mitotic growth (Sugiyama and Nikawa 2001).

RFX represents a family of fungal TFs involved in DNA repair. Each strain encodes two orthologous copies, except *F. venenatum* encodes two copies within the RFX1 clade (Figure S3C). A major transcriptional repressor of DNA-damage-regulated genes in *S. cerevisiae*, Rfx1, is involved in DNA damage repair and replication checkpoint pathways (Lubelsky et al. 2005). In *F. graminearum*, Rfx1 is essential for maintaining the genome integrity (Min et al. 2014). The other copy, Rsc9 in *S. cerevisiae*, is a chromatin structure-remodeling complex RSC involved in transcription regulation and nucleosome positioning (Cairns et al. 1996; Hsu et al. 2003).

NFYA can bind to the CCAAT box. All strains maintain one copy of this family. The yeast homolog Hap2 induces the expression of mitochondrial electron transport genes (Olesen et al. 1991). *F. verticillioides* NFYA Hap2 is essential for fungal growth and the virulence on maize stalks (Ridenour and Bluhm 2014). The conservativeness suggests the functional importance of these TFs across *Ascomycota*, possibly linked to cellular machinery control (e.g. mitochondrial electron transport chain).

MADS MEF2-like family includes *S. cerevisiae* Rlm1, a component of the protein

kinase C-mediated MAP kinase pathway involved in maintaining cell integrity (Jung et al. 2002). Rlm1 has a paralog from the whole genome duplication in *S. cerevisiae*, and all filamentous fungi encode one copy. In *F. verticillioides*, Mef2 plays a vital role in the sexual development (Ortiz and Shim 2013).

3. Minimal gene family contractions in FOSC partially caused by whole genome duplication in yeast

Four TF families, MATalpha_HMGbox, NOT4, MADS_SRF-like, and HSF (Heat Shock Factor), have an expansion score of less than 1, reflecting some level of gene family contraction among members of FOSC compared to the two yeast genomes (Figure S4).

MATalpha_HMGbox is a TF family that includes *S. cerevisiae* mating type protein alpha 1, a transcription activator that activates mating-type alpha-specific genes (Martin et al. 2010). All *F. oxysporum* Mat1-1 type strains contain this TF, but Mat1-2 strains do not. The contraction reflects the heterothallic mating strategy, even though sexual reproduction has not been observed in FOSC (Arie et al. 2000).

NOT4 is a component of the multifunctional CCR4-NOT complex, a global transcriptional repressor of the RNA polymerase II transcription (Albert et al. 2002). This TF family remains a single copy in most genomes but is lost in some filamentous fungal genomes, including *A. nidulans*, *F. redolens*, *F. oxysporum* strains NRRL26365, MRL8666, and PHW726. It remains to be discovered why this gene is lost in some of these strains.

The contractions of the other two TF families, **MADS SRF-like** and **HSF**, are primarily caused by the whole genome duplication in yeast. In both cases, some degree of expansion was found in FOSC compared to other filamentous fungi (Figure S4).




Figure 3. Evolutionary trajectory of heat shock factors (HSFs) suggesting genome expansion and adaptation. (A) Phylograms of HSFs were constructed by maximum likelihood method with 1000 bootstraps. Branches of *Fusarium* HSFs were colored in yellow. Accessory HSFs of FOSC are shared in red. (B) Number of accessory HSFs in some FOSC genomes. (C) Expression of *HSF* genes during plant colonization (hpi indicates hours post inoculation), compared to axenic growth. Transcriptome data was previously described in Guo et al. 2021. See Materials and Methods for details of data reprocessing and visualization.

336

337 **MADS SRF-like** is important for microconidium production and virulence in host
338 plants, as reported in *M. oryzae* (Ding et al. 2020), and is essential for
339 transcriptional regulation of growth-factor-inducible genes (Messenguy and
340 Dubois 2003). The average copy number of phytopathogenic FOSC strains is
341 2.73, and the Fo5176 genome has the highest copy number of 6, while most
342 other genomes only contain a single copy (Table S1).

343 **HSF** is a family of transcription factors that activate the production of many heat
344 shock proteins that prevent or mitigate protein misfolding under abiotic/biotic
345 stresses (Feder and Hofmann 1999). All non-FOSC filamentous fungi have three
346 copies, while members of FOSC show expansion (e.g., Fo47: 4, Fo4287: 5, I15:
347 4, HDV274: 4, and Fo5176: 4) (Figure 3A-B). Interestingly, all expanded HSFs
348 are phylogenetically close to Hsf1, which cluster together with the Hsf1 paralog of

Fusarium solani, suggesting their horizontal transfer origin (Figure 3A). We then examined the *Hsf1* expression during the plant colonization (Guo et al. 2021). We found that the core copies of *Hsf1* of both strains Fo47 and Fo5176 were up-regulated during plant colonization. In contrast, the *Hsf1* accessory copies of these two strains were under opposite regulations, with Fo47 one being up-regulated and Fo5176 one being down-regulated post infection (Figure 3C), suggesting distinct regulatory adaptations after expansion. Here we noted that transcriptome data could be powerful in understanding the functional importance of TFs (see Section 6 for systematic analysis). In filamentous fungi, there are experimental reports for the other two clades. *Sfl1* is essential for vegetative growth, conidiation, sexual reproduction, and pathogenesis, as shown in *M. oryzae* (Li et al. 2011); *Skn7* is a regulator of the oxidative stress response and is essential for pathogenicity in *F. graminearum* (Jiang et al. 2015). Not surprisingly, both genes of Fo5176 and Fo47 were upregulated during plant colonization (Figure 3C).

4. Significant TFome expansion in FOSC driven by a small number of exceedingly expanded TF families

4.1. Gain-of-function among filamentous ascomycete fungi

Three TF families, CP2 ($El_y = 2.73$), HTH_AraC ($El_y = 2$), and HTH_Psq ($El_y = 3.53$), are absent in both yeast genomes, suggesting a gain of function among filamentous ascomycete fungi (Table S1). **CP2** has been studied in animal and fungal kingdoms with a function related to differentiation and development (Paré et al. 2012). Both **HTH_AraC** and **HTH_Psq** belong to helix-turn-helix (HTH) superfamily. First reported in bacteria, **HTH_AraC** is a positive regulator associated with the arabinose operon regulatory protein AraC (Schleif 2010; Gallegos et al. 1993; Bustos and Schleif 1993). **HTH_Psq**, as part of the eukaryotic Pipsqueak protein family, reported in vertebrates, insects, nematodes, and fungi, regulates the cell death (Siegmund and Lehmann 2002). Most FOSC genomes have a single copy of HTH_AraC, while the count of proteins containing

the HTH_Psq ranges from 0 to 9 in the FOSC and ranges from 0 to 3 in other *Fusarium* relatives. Since the **HTH_Psq** domain also exists in transposases (Siegmund and Lehmann 2002), and ACs in FOSC are transposon-rich, it remains to be studied whether proteins containing the Psq domain are *bona fide* TFs.

4.2. Seven exceedingly expanded TF families

Among others, seven TF families have expansion indexes greater than 2 (Table 2 and Figure 2B). Because of their drastic expansion, these seven families overall account for more than 75% of the total TFome. These families include Zn2-C6 ($EI_y = 15.09$), bZIP ($EI_y = 5.80$), and Znf_C2H2 ($EI_y = 4.15$), Homeobox ($EI_y = 2.28$), PAI2 ($EI_y = 3.42$), NDT80 ($EI_y = 3.47$), and bHLH ($EI_y = 3.48$). All seven families show gradual expansion, reflected by the average copy number increment (FOSC > non-FOSC *Fusarium* > non-*Fusarium* filamentous fungi > yeasts, table S1). Furthermore, Zn2-C6 (44 in yeasts versus 671 in FOSC) and Znf_C2H2 (40 in yeasts versus 167 in FOSC) have the most drastic number increment along the evolutionary trajectory (Figure 2C and Table S1). Based on both high expansion index and large number increment, we considered Zn2-C6 and Znf_C2H2 as the most significantly expanded families.

The large copy number makes it hard to interpret functions from the protein domain annotation. Here we describe a couple of TFs reported in *F. oxysporum* and other systems and will introduce orthologous analysis to further survey the functionally validated TFs in the later section.

Zn2-C6, a fungal family TF (MacPherson et al. 2006), has the most significant expansion, reaching over 600 members among FOSC genomes and accounting for more than half of the total TFome. This group of TFs can form a homodimer and bind to the specific palindromic DNA sequence through direct contact with the major groove of the double-stranded DNA molecules (MacPherson et al. 2006). The versatility of this group of TFs can be achieved by domain shuffling and by changing the nucleotide binding specificity. In addition to the well-

documented Ftf1 (Niño-Sánchez et al. 2016; van der Does et al. 2016; Ramos et al. 2007; Zuriegat et al. 2021; Zhao et al. 2020), five additional TFs within this family have been characterized in *F. oxysporum*, including Ctf1 (Rocha et al. 2008), Ctf2 (Rocha et al. 2008), Fow2 (Imazaki et al. 2007), XlnR (Calero-Nieto et al. 2007) and Ebr1 (Jonkers et al. 2014). They are involved in the development, metabolism, stress response, and pathogenicity.

Znf_C2H2 is the most common DNA-binding motif found in the eukaryotic transcription factors (Fedotova et al. 2017). Five *F. oxysprum* TFs have been reported: *Czf1* (Yun et al. 2019), *Con7-1* (Ruiz-Roldán et al. 2015), *PacC* (Caracuel et al. 2003; Zhang et al. 2020), *ZafA* (López-Berges 2020) and *St12* (Asunción García-Sánchez et al. 2010; Risipail and Di Pietro 2009). Particularly, *PacC* was linked to the pathogenicity of both plant and human host (Zhang et al. 2020; Caracuel et al. 2003).

Other five families include **bZIP**, **Homeobox**, **PAI2**, **Ndt80** and **bHLH**. **bZIP** domain contains a region for sequence-specific DNA binding followed by a leucine zipper region required for dimerization (Bader and Vogt 2006). Three *F. oxysporum* bZIP TFs have been reported, including Atf1 (Li et al. 2013), Hapx (López-Berges et al. 2012), and MeaB (López-Berges et al. 2010), all of which are important for fungal pathogenicity. **Homeobox** is a DNA binding motif with a helix-turn-helix structure. In *S. pombe*, *Phx1* is a transcriptional coactivator that plays a role in yeast fission. In *M. oryzae*, *Hox* plays roles in the conidiation and appressorium development (Kim et al. 2009). **PAI2** is involved in the negative regulation of protease synthesis and sporulation of the *Bacillus subtilis* (Honjo et al. 1990). **Ndt80** is essential for completing meiosis in *S. cerevisiae* (Pierce et al. 2003; Tsuchiya et al. 2014) and *Ustilago maydis* (Doyle et al. 2016). It also promotes the expression of sporulation genes that are essential for the fulfillment of meiotic chromosome segregation (Hepworth et al. 1998). **bHLH** proteins form a large superfamily of transcriptional regulators found in almost all eukaryotes and function in critical developmental processes (Jones 2004). *F. graminearum* Gra2 is involved in the biosynthesis of phytotoxin gramillan (Bahadoor et al. 2018). *P. digitatum* encoding SreA is required for anti-fungal resistance and full

virulence in citrus fruits (Liu et al. 2015).

4.3. other families

Other 20 TF families (expanded but with $El_y \leq 2$) account for 20% of the TFome; on average, each of these 20 families contains 9.6 copies in each genome examined (Table S1). These TFs are involved in chromatin remodeling and pheromone response, among other functions.

Four TF families are functionally linked to chromatin remodeling, including Bromodomain ($El_y = 1.52$), CBFA_NFYB ($El_y = 1.35$), Znf_RING-CH ($El_y = 1.11$), and ARID ($El_y = 1.25$). **Bromodomain** containing Spt7 is a crucial part of the SAGA complex in yeast. The SAGA complex is required to transcribe many genes in the genome. The bromodomain that is part of this subunit can recognize acetylated lysines of histones and eventually lead it to a more chromatin unwinding (Donczew et al. 2020). **CBFA_NFYB** is found in the proteins (e.g., *S. cerevisiae* Dls1) that regulate RNA polymerase II transcription through controlling chromatin accessibility (e.g., telomeric silencing) (Iida and Araki 2004). **Znf_RING-CH** has a functional connection to chromatin modification (e.g., *S. cerevisiae* Rkr1) (Braun et al. 2007). **ARID** is a 100 amino acid motif found in many eukaryotic TFs (Iwahara 2002). *S. cerevisiae* Swi1 plays a role in chromatin remodeling and is required to transcribe a diverse set of genes, including HO and Ty retrotransposons (Breedon and Nasmyth 1987; Hirschhorn et al. 1992).

Ste12 is a family of TFs that regulate fungal development and pathogenicity (Risipail and Di Pietro 2010). These TFs are found only in the fungal kingdom. Ste12 binds to the DNA sequence that mediates pheromone response. It is involved in haploid mating and pseudohyphae formation in the diploid (Gancedo 2001). *F. oxysporum* Ste12 controls invasive growth and virulence downstream

of the Fmk1-mediated MAPK cascade (Risipail and Di Pietro 2009). Except for *S. pombe* (missing one), every genome encodes one copy.

Among others, **Znf_NFX1** domain is found in the NK-X1, a repressor of the human disease-associated gene HLA-DRA (Song et al. 1994). **HMG_box** (high mobility group box) in *S. cerevisiae*, Spp41, is involved in negative expression regulation of spliceosome components (Maddock et al. 1994); Nhp6a is required for the fidelity of some tRNA genes (Braglia et al. 2007); Ixr1 is a transcriptional repressor that regulates hypoxic genes (Vizoso-Vázquez et al. 2012). One example of **Znf_GATA** is Fep1, a transcription factor that represses the expression of particular iron transporter genes under a high iron concentration (Kim et al. 2016). *S. cerevisiae* Mbf1, belonging to **Cro/C1-type HTH**, is a transcriptional coactivator (Takemaru et al. 1997).

5. Orthologous survey of TF families that were manually curated

To further understand expanded TFs and their impacts on transcriptional regulation, we curated a list of 102 TFs reported in literature focusing on *F. oxysporum*, *F. graminearum*, and other phytopathogenic fungi (Table S3 and examples as described in the previous section). Compared to this list of curated TFs using Orthofinder, we define 80 orthologous groups among *Fusarium* genomes (Table S4). 62 out of the 80 orthogroups have been identified using the above IPR-annotated pipeline, which enables the dissection of vastly expanded and high copy number TF families such as Zn2-C6 and Znf_C2H2, which are further mapped to 27 orthologous groups, including 17 in Zn2C6, 9 in Znf_C2H2, and 1 containing both Znf_C2H2 and Zn2-C6 domains (Table S4).

This effort also results in additional annotation to 18 TF families (Table S4), accounting for 32 genes per genome (3% of average *Fusarium* TFome size). These newly annotated TFs include homologs of those without domain annotation, e.g., disordered proteins *F. oxysporum* Ren1 (Ohara et al. 2004) and

M. oryzae Som1 (Yan et al. 2011), and homologs of those with noncanonical TF domains such as **Ankyrin_rpt** and **WD40_repeat**.

We then directly compared *F. oxysporum* with its *Fusarium* relatives to calculate the expansion index as follows:

$$El_f = \frac{\text{Average number of TFs in FOSC} + 1}{\text{Average number of TFs in FOSC sister species} + 1}$$

The El_f ranged, with the highest score being 3.54 (Fug, AreA_GATA) and the lowest being 0.5 (Fox1, Fork_head) (Table S4). Among these 80 orthogroups, 36 groups show high conservation ($El_f = 1$) as they are single-copy orthologs across *Fusarium*, among which ten were functionally validated in *F. oxysporum* (Table S4). 24 groups have gene contraction in *F. oxysporum* ($El_f < 1$). A total of 20 groups are expanded in *F. oxysporum* ($El_f > 1$, Table 3, Table S4), including five groups Fug1 (AreA_GATA, $El_f = 3.54$), Cos1 (Znf_C2H2, $El_f = 2.8$), Ftf1/Ftf2 (Zn2-C6, $El_f = 2.7$), Ebr1/Ebr2 (Zn2-C6, $El_f = 2.5$) and Ren1 (disordered, $El_f = 2$), with an El_f value equal or greater than 2. We also identified PacC ($El_f = 1.57$) as the second most expanded group within the highly expanded Znf_C2H2 family. We will further discuss these six groups (highlighted in bold, Table 3).

519

Table 3. Ortholog copy number and expansion index (*E_I*) of characterized and expanded TFs in *F. oxysporum*

TF	Reported species	References	Family	Overlap	Average_ Fo	Average_ non- Fo	<i>E_I</i>
Ftf1/Ftf2	<i>F. oxysporum</i>	(Niño-Sánchez et al. 2016)	Zn2-C6	Yes	4.80	1.11	2.75
Ebr1/Ebr2	<i>F. oxysporum</i>	(Jonkers et al. 2014)	Zn2-C6	Yes	5.27	1.56	2.45
Znf1	<i>M. oryzae</i>	(Yue et al. 2016)	Zn2-C6	Yes	6.47	2.78	1.98
Ctf2	<i>F. oxysporum</i>	(Bravo-Ruiz et al. 2013)	Zn2-C6	Yes	2.93	1.33	1.69
Fow2	<i>F. oxysporum</i>	(Imazaki et al. 2007)	Zn2-C6	Yes	2.07	1.00	1.53
Dep6	<i>A. brassicicola</i>	(Wight et al. 2009)	Zn2-C6	Yes	0.93	0.67	1.16
Pf2	<i>A. brassicicola</i>	(Jones et al. 2019)	Zn2-C6	Yes	1.20	1.00	1.10
Art1	<i>F. verticillioides</i>	(Oh et al. 2016)	Zn2-C6	Yes	1.00	0.89	1.06
Clta1	<i>C. lindemuthianum</i>	(Dufresne et al. 2000)	Zn2-C6	Yes	1.07	1.00	1.03
Fhs1	<i>F. graminearum</i>	(Son et al. 2016a)	Zn2-C6	Yes	1.07	1.00	1.03
Cos1	<i>M. oryzae</i>	(Li et al. 2013)	Znf_C2H2	Yes	1.80	0.00	2.80
PacC	<i>F. oxysporum</i>	(Caracuel et al. 2003)	Znf_C2H2	Yes	2.13	1.00	1.57
Fug1	<i>F. verticillioides</i>	(Ridenour and Bluhm 2017)	AreA_GATA	No	7.27	1.33	3.54
Ren1	<i>F. oxysporum</i>	(Ohara et al. 2004)	disordered	No	3.00	1.00	2.00
Tri10	<i>F. graminearum</i>	(Jiang et al. 2016)	Fun_TF	No	1.13	0.33	1.60
Ltf1	<i>B. cinerea</i>	(Schumacher et al. 2014)	Znf_GATA	Yes	4.00	2.44	1.45
Ndt80	<i>U. maydis</i>	(Doyle et al. 2016)	NDT80	Yes	1.73	1.11	1.29
Hap3p	<i>F. verticillioides</i>	(Ridenour and Bluhm 2014)	CBFA_NFYB	Yes	1.33	1.00	1.17
Sod1	<i>F. oxysporum</i>	(Wang et al. 2021)	SOD_Cu_Zn	No	1.47	1.22	1.11
Prf1	<i>F. oxysporum</i>	(Mendoza-Mendoza et al. 2009)	HMG_box	Yes	1.07	1.00	1.03

520

521 Both **Ftf1/Ftf2** and **Ebr1/Ebr2** belong to the Zn2-C6 family and contribute directly
522 to the fungal virulence (Michielse et al. 2009; van der Does et al. 2016; Ramos et
523 al. 2007). Deletion of accessory copy Ftf1 reduced the pathogenicity of *F.*
524 *oxysporum* f. sp. *phaseoli* (Ramos et al. 2007), highlighting the direct functional
525 involvement of AC TF in virulence. In *Fol*, deletion of either Ftf1 (AC encoding) or
526 Ftf2 (CC encoding) reduced the virulence towards the host (de Vega-Bartol et al.
527 2011; Niño-Sánchez et al. 2016). Constitutive expression of either *Ftf1* or *Ftf2*
528 induced the expression of effector genes (van der Does et al. 2016). The core
529 copy Ftf2 is conserved among all *Fusarium* species, and the AC copy Ftf1 is only
530 found in *F. oxysporum* and *Fusarium redolens* (Figure 4). **Ebr1** and paralogues
531 are responsible for virulence and general metabolism. In *F. oxysporum*, Ebr1 is

found as multiple homologs, whereas in *F. graminearum*, it is seen as a single copy (Jonkers et al. 2014). In *F. oxysporum*, three paralogous copies, Ebr2, Ebr3, and Ebr4, are encoded in ACs and regulated by core copy Ebr1. The importance of the core paralog has been shown by the reduced pathogenicity and growth defects when it was knocked out (Jonkers et al. 2014). It is worth noting that the *Ebr2* coding sequence driven by an *Ebr1* promoter was able to rescue the *Ebr1* knockout mutation, indicating some functional redundancy of this family.

Both **Cos1** and **PacC** belong to the Znf_C2H2 family. Mutation to *M. oryzae* *Cos1* resulted in developmental failure of the conidiophores (Li et al. 2013). Furthermore, mutation to *Cos1* aggravated the plant infection of leaf blades and sheaths, indicating a negative role in the pathogenicity (Zhou et al. 2009). *PacC* is an important pH-responsive TF in *F. oxysporum* (Caracuel et al. 2003; Zhang et al. 2020). *PacC* homologs are expanded in clinical strains (average accessory copy number 3.7) of FOSC, compared to non-clinical strains (average accessory copy number 0.5), while all *Fusarium* relatives' genomes examined only contain a single copy of core *PacC*. Our previous study revealed that in *F. oxysporum* clinical strains, the expression of one expanded *PacC* gene on ACs was induced and the protein localized in the nucleus at mammalian physiological pH (7.4), indicating a potential role in host adaptation (Zhang et al. 2020). Interestingly, the induction of AC-encoding *PacC* genes was CC-encoding *PacC* gene-dependent, as the induction disappeared in the CC-encoding *PacC* knockout mutant, further supporting a cross-talking between core and accessory TFs (Yang 2020). Similar to *EBR1*, the expression of AC *PacC* genes is much lower than that of the CC *PacC* gene, and knockouts of one AC *PacC* gene affected a small subset of genes compared with the CC *PacC* knockout, which has a broader effect on cellular processes (Yang 2020).

Fug1 has a role in pathogenicity (maize kernel colonization) and fumonisin biosynthesis in *F. verticillioides* (Ridenour and Bluhm 2017). In addition, the deletion of *Fug1* increased sensitivity to the antimicrobial compound 2-

benzoxazolinone and to hydrogen peroxide, which indicates that Fug1 plays a role in mitigating stresses associated with the host defense (Ridenour and Bluhm 2017). Neither core copies nor accessory copies of these two genes were experimentally examined in FOSC. **Ren1** is a disordered protein with no IPR functional domain. The expansion score $E_{I_f} = 2$ suggests a unique expansion among FOSC. However, the only reported study on its function is in *F. oxysporum f. sp. melonis* regulating the development of the conidiation (Ohara et al. 2004).

6. Transcriptome analysis to probe the essential TFs during host colonization

To understand the functional importance of FOSC TFs, we take advantage of two recently reported transcriptomics datasets (Redkar et al. 2022; Guo et al. 2021), including pathogenic interactions (Fo5176 infecting Arabidopsis and Fo4287 infecting tomato) and endophytic interactions (Fo47 colonizing Arabidopsis) (Supplemental Dataset).

We first asked what proportion of genes was expressed in conserved and expanded categories (Table S5). We found that almost all genes (58 out of 60) within the conserved category (Group 1) were consistently expressed (TPM > 1 across all conditions), supporting their general roles in controlling life processes. Within the expanded category (Group 3), the proportion of genes being consistently expressed ranges from 41% to 59% for core TFs, and ranges from 5% to 16% for accessory TFs. With a less strict filter (TPM > 1 at minimum 1 condition), we found that all genes within the conserved category were expressed. Within the expanded category, the proportion of genes being expressed accounts for 93% of core TFs across all strains and ranges from 49% to 67% for accessory TFs. When we compared genes being consistently expressed versus genes being expressed at a minimum one condition, the more dramatic number increase for the expanded category (especially when we only consider the accessory TFs) highlighted that the expanded category, especially

the accessory TFs, are more likely to be conditional expressed, further supporting their role in niche adaptation.


With the goal of examining the expression and probing important core and accessory TFs, we aimed to develop filtering parameters. Since most validated TFs were reported in the reference Fol4287 strain, we first reviewed for the reported TFs, both core, and accessory, the expression pattern during Fol4287 infecting tomatoes (Table S6). Out of 27 TFs encoded on the core genome, 18 show up-regulation (defined by at least three out of four *in planta* conditions show up-regulation compared to the axenic growth) during plant colonization from 1 day post-inoculation (dpi) to 7 dpi, consistent with their reported roles in pathogenicity. The accessory *Ftf1* has been exclusively demonstrated to play essential functions in fungal pathogenicity (Niño-Sánchez et al. 2016). Eight of 10 accessory *Ftfs* were upregulated using the same criteria during plant colonization. Our results illustrate the power of using transcriptome data to probe the functionally important players during plant colonization/infection.

We further developed strict criteria to filter important TFs from TFome (Figure S5), by which half (nine) of previously described upregulated core TFs meet the 'core' criteria, and all eight up-regulated accessory *Ftfs* meet the 'accessory' criteria (Table S6). We then apply such measures to the transcriptome of all the TFomes of Fol4287, Fo5176, and Fo47 to probe two types of TFs: 1) The conserved core TFs related to plant colonization; 2) Expanded accessory TFs related to host-specific pathogenicity.

Fol4287, Fo5176, and Fo47 upregulated 95, 62, and 44 core TFs during plant colonization. Among them, ten copies are highly conserved (Table S7), as they are single-copy orthologs across all 15 *F. oxysporum* strains. Two out of ten were previously reported, *Fow2* and *Sfl1*. *Fow2*, Zn2C6 TF, is required for full virulence but not hyphal growth and conidiation in *F. oxysporum* f. sp. *melonis* (Imazaki et al. 2007). The downstream targets of *Fow2* remain unknown in *F. oxysporum*, thus meriting further analysis. *Sfl1* has been described in the

previous section and is essential for vegetative growth, conidiation, sexual reproduction, and pathogenesis, as shown in *M. oryzae* (Li et al. 2011). The functions of FoSfl1 remain to be validated.

Fol4287, Fo5176, and Fo47 upregulated 29, 34, and 9 accessory TFs. *Ftf1* and *Ren1* are particularly interesting (Figure 4 and Table S8). Though *Ftfs* have been shown to play an essential role in pathogenicity in Fol, whether this pathway is restricted to the same strain remains a question. Compared to Fol4287 which contains ten accessory *Ftfs* and eight were upregulated during plant colonization, Fo5176 includes six copies of accessory *Ftfs*, but only one copy was upregulated. Interestingly, eight upregulated Fol4287 and one upregulated Fo5176 *Ftfs* are clustered together (Figure 4). The unique expansion with regulatory adaptation (*i.e.*, fine-tuned expression regulation) seems to be restricted to Fol4287 but not another pathogenic strain, Fo5176, when they infect the hosts. Among Fo5176 expanded TFs, we identified *Ren1*. Compared to Fol4287 which encodes only one accessory *Ren1* that was not upregulated during plant colonization, Fo5176 encodes seven accessory copies, among which two were upregulated (Figure 4). Though functional validation is needed, the strain-specific expansion followed by fine-tuned expression regulation when infecting host species exists and likely contributes to the host-specific pathogenicity.



DISCUSSION

For a soilborne pathogen with strong host specificity like FOSC, the adjustment of growth and cell cycle control in response to environmental cues is likely essential for survival. At the same time, expanded families likely contribute to the enhanced functions related to niche adaptation. TFs transmit external and internal signals and regulate complex cellular signaling responses to the sensed stimuli. Transmitted through the soil and vascular wounds of plants causing vascular wilt (Gordon 2017), *F. oxysporum* must adapt to stresses encountered both outside and inside its host. Therefore, it is not surprising to see that genomes of FOSC have larger TFome than other fungi included in the study. The expansion of TFs among FOSC resulted in a positive correlation between the total number of proteins and the size of the fungal TFome, which was also observed before (Shelest 2017).

A total of 14 TF families that control the global transcriptional event, such as TBP, are highly conserved within the ascomycete fungal lineages. Conserved regulatory mechanisms revealed through this study suggest that the plant colonization process could be a common process among FOSC strains regardless of their host-specific pathogenesis. The notion was also supported by recent studies that highlighted the ability of FOSC as a root colonizer facilitated by the conserved genomics components (Martínez-Soto et al. 2022; Redkar et al. 2022).

In contrast to these stable TFs, 30 families are expanded in various degrees and most significant expansions occurred in Zn2-C6 and Znf_C2H2 TF families among FOSC genomes. The number of Zn2-C6 TFs increases significantly (with the highest expansion score) and makes up most of the TFs (56.7%) found within the FOSC TFome. For example, *Ftf1*, a TF belonging to the Zn2-C6 and involved in the tomato pathogenicity, is most significantly expanded (10 copies of accessory *Ftfs*) in the tomato pathogen Fol4287 genome, and the expression of eight out of 10 was induced during plant colonization. The continuous expansion

suggests the functional importance of these understudied TFs, further supported by the genetic studies (Niño-Sánchez et al. 2016; van der Does et al. 2016; Ramos et al. 2007; Zuriegat et al. 2021; Zhao et al. 2020) and their induction during host invasion revealed by our RNA-seq data.

Unique expansion of some TFs, driven by ACs, may provide a clue to host-specific interactions. Acquiring additional TFs will modify existing regulatory pathways. No question, this will require the fine-tuning of existing networks for this group of organisms to successfully adapt to different hosts under diverse environments. A previous survey of kinome (the complete set of protein kinases encoded in an organism's genome) among FOSC and other Ascomycetes revealed a positive correlation between the size of the kinome and the size of the genome (Delulio et al. 2018), exactly the same as we reported here for TFomes. As kinases and TFs are key regulators that modulate all important signaling pathways and are essential for the proper functions of almost all molecular and cellular processes. Strong correlations among kinome and TFome suggest an ordered, instead of chaotic, recruitment and establishment of ACs among FOSC genomes.

This realization further emphasizes the importance of additional functional studies. Reverse genetics is a powerful tool in defining the functional importance of a TF. For example, TF Ren1, a disordered protein, was identified by genetic and molecular characterization (Ohara et al. 2004). This TF is most significantly expanded (seven copies of accessory *Rens*) in the Arabidopsis pathogen Fo5176 genome, and two of them were induced during plant colonization. Experiments such as chromatin immunoprecipitation sequencing (CHIP-Seq) and DNA affinity purification sequencing (DAP-seq) to profile the cis-regulatory elements globally are high throughput approaches to define specific binding sites (cis-regulatory elements) of TFs. DAP-seq was used successfully to profile the Cistrome for the entire TFome of the bacterial organism (Baumgart et al. 2021), holding the promise for a better understanding of transcriptional regulation in the fungal model *F. oxysporum*. TFs can function individually or with other proteins in

a complex, and can act as an activator that promotes transcription or a repressor that blocks the recruitment of RNA polymerase. Therefore, defining specific functions of these identified binding sites through DAP-seq can be difficult. Gene regulatory networks based on gene co-expression and other phenotypic and multi-omics data as reported in *Fusarium* (Guo et al. 2016, 2020) can add more resolution to these complex regulatory processes. However, the ultimate understanding of the regulatory roles of each TF will come from careful molecular and biochemical characterization.

A systematic understanding of transcriptional regulation is essential to get the fine-tuned footprint of the gene regulatory network. Our study not only offered a comprehensive look at the regulation from the evolutionary perspective, but also provided an easily implemented computational pipeline to compare TFs and other functional groups in fungi. A better understanding of their functions would not only inform *Fusarium* biology but also could be extrapolated to other filamentous fungi and complex basidiomycetes.

SUPPLEMENTAL MATERIALS

Table S1. TF-type DNA-binding domains used to filter TFome

Table S2. TFome annotations across 30 genomes

Table S3. Curated TFs in phytopathogens

Table S4. Ortholog copy number of characterized TFs of phytopathogens across 30 genomes

Table S5. Number of TFs being expressed

Table S6. Orthologs of reported TFs in Fol4287

Table S7. Highly conserved TFs that are constantly upregulated during plant colonization

Table S8. Strain-specific accessory TFs that are upregulated during plant colonization

735
 736 Figure S1. TFome annotation pipelines
 737 Figure S2. ACs contribute to the FOSC TFome expansion
 738 Figure S3. Phylograms of three conserved families
 739 Figure S4. Minimal gene family contractions in FOSC partially caused by whole
 740 genome duplication in yeast
 741 Figure S5. The pipelines to probe the functional important TFs by RNA-seq data
 742
 743 Supplemental Dataset. Normalized read counts of RNA-seq datasets
 744

745 **FUNDING**

746 This project was supported by the Natural Science Foundation (IOS-165241), the
 747 National Institutes of Health (R01EY030150), and the USDA National Institute of
 748 Food and Agriculture (MAS00496). The work
 749 (proposal:10.46936/10.25585/60001360) conducted by the U.S. Department of
 750 Energy Joint Genome Institute (<https://ror.org/04xm1d337>), a DOE Office of
 751 Science User Facility, is supported by the Office of Science of the U.S.
 752 Department of Energy operated under Contract No. DE-AC02-05CH11231. H.Y.
 753 is also supported by Lotta M. Crabtree Fellowship and Constantine J. Gilgut
 754 Fellowship. S.M.C. is also supported by the Vaadia-BARD Postdoctoral
 755 Fellowship.

756

757 **ACKNOWLEDGMENTS**

758 We thank the Massachusetts Green High-Performance Computing Center for
 759 providing high-performance computing capacity.

760

761 **AUTHOR CONTRIBUTIONS**

Conceptualization, H.Yu and L-J.M.; Methodology, Analysis and Interpretation, all authors; Resources, S.H., R.H. and I.G.; Writing – H.Yu and L-J.M. with input from all authors; Supervision, L-J.M.; Funding Acquisition, L-J.M.

CONFLICTS OF INTERESTS

The authors declare no conflict of interest. The sponsors had no role in the design, execution, interpretation, or writing of the study.

REFERENCES

- Armstrong, G.M., and Armstrong, J.K. (1981). *Formae speciales and races of Fusarium oxysporum causing wilt disease*. In *Fusarium: Disease, Biology, and Taxonomy*, P.E. Nelson, T.A. Toussoun, and R.J. Cook, eds (University Park, PA: Pennsylvania State University Press), pp. 391–399.
- Albert, T. K., Hanzawa, H., Legtenberg, Y. I. A., de Ruwe, M. J., van den Heuvel, F. A. J., Collart, M. A., Boelens, R., and Timmers, H. Th. M. 2002. Identification of a ubiquitin-protein ligase subunit within the CCR4-NOT transcription repressor complex. *The EMBO Journal*. 21:355–364
- Arie, T., Kaneko, I., Yoshida, T., Noguchi, M., Nomura, Y., and Yamaguchi, I. 2000. Mating-type genes from asexual phytopathogenic ascomycetes *Fusarium oxysporum* and *Alternaria alternata*. *Mol Plant Microbe Interact*. 13:1330–1339
- Asunción García-Sánchez, M., Martín-Rodrigues, N., Ramos, B., de Vega-Bartol, J. J., Perlin, M. H., and Díaz-Mínguez, J. M. 2010. *fos12*, the *Fusarium oxysporum* homolog of the transcription factor *Ste12*, is upregulated during plant infection and required for virulence. *Fungal Genetics and Biology*. 47:216–225
- Bader, A. G., and Vogt, P. K. 2006. Leucine Zipper Transcription Factors: bZIP Proteins. Pages 964–967 in: *Encyclopedic Reference of Genomics and Proteomics in Molecular Medicine*, Springer, Berlin, Heidelberg.
- Bahadoor, A., Brauer, E. K., Bosnich, W., Schneiderman, D., Johnston, A., Aubin, Y., Blackwell, B., Melanson, J. E., and Harris, L. J. 2018. Gramillin A and B: Cyclic Lipopeptides Identified as the Nonribosomal Biosynthetic Products of *Fusarium graminearum*. *J. Am. Chem. Soc*. 140:16783–16791
- Baumgart, L. A., Lee, J. E., Salamov, A., Dilworth, D. J., Na, H., Mingay, M., Blow, M. J., Zhang, Y., Yoshinaga, Y., Daum, C. G., and O’Malley, R. C. 2021. Persistence and plasticity in bacterial gene regulation. *Nat Methods*. 18:1499–1505
- Becker, D. M., Fikes, J. D., and Guarente, L. 1991. A cDNA encoding a human CCAAT-binding protein cloned by functional complementation in yeast. *Proc. Natl. Acad. Sci. U.S.A*. 88:1968–1972

799 Blum, M., Chang, H.-Y., Chuguransky, S., Grego, T., Kandasaamy, S., Mitchell, A.,
800 Nuka, G., Paysan-Lafosse, T., Qureshi, M., Raj, S., Richardson, L., Salazar, G.
801 A., Williams, L., Bork, P., Bridge, A., Gough, J., Haft, D. H., Letunic, I.,
802 Marchler-Bauer, A., Mi, H., Natale, D. A., Necci, M., Orengo, C. A.,
803 Pandurangan, A. P., Rivoire, C., Sigrist, C. J. A., Sillitoe, I., Thanki, N., Thomas,
804 P. D., Tosatto, S. C. E., Wu, C. H., Bateman, A., and Finn, R. D. 2021. The
805 InterPro protein families and domains database: 20 years on. *Nucleic Acids*
806 *Research*. 49:D344–D354

807 Braglia, P., Dugas, S. L., Donze, D., and Dieci, G. 2007. Requirement of Nhp6 Proteins
808 for Transcription of a Subset of tRNA Genes and Heterochromatin Barrier
809 Function in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 27:1545–1557

810 Braun, M. A., Costa, P. J., Crisucci, E. M., and Arndt, K. M. 2007. Identification of Rkr1,
811 a Nuclear RING Domain Protein with Functional Connections to Chromatin
812 Modification in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 27:2800–2811

813 Bravo-Ruiz, G., Ruiz-Roldán, C., and Roncero, M. I. G. 2013. Lipolytic system of the
814 tomato pathogen *Fusarium oxysporum* f. sp. *lycopersici*. *Mol Plant Microbe*
815 *Interact*. 26:1054–1067

816 Breeden, L., and Nasmyth, K. 1987. Cell cycle control of the yeast HO gene: Cis- and
817 Trans-acting regulators. *Cell*. 48:389–397

818 Bustos, S. A., and Schleif, R. F. 1993. Functional domains of the AraC protein. *Proc.*
819 *Natl. Acad. Sci. U.S.A.* 90:5638–5642

820 Cairns, B. R., Lorch, Y., Li, Y., Zhang, M., Lacomis, L., Erdjument-Bromage, H.,
821 Tempst, P., Du, J., Laurent, B., and Kornberg, R. D. 1996. RSC, an Essential,
822 Abundant Chromatin-Remodeling Complex. *Cell*. 87:1249–1260

823 Calero-Nieto, F., Di Pietro, A., Roncero, M. I. G., and Hera, C. 2007. Role of the
824 Transcriptional Activator XlnR of *Fusarium oxysporum* in Regulation of
825 Xylanase Genes and Virulence. *MPMI*. 20:977–985

826 Caracuel, Z., Roncero, M. I. G., Espeso, E. A., González-Verdejo, C. I., García-Maceira,
827 F. I., and Di Pietro, A. 2003. The pH signalling transcription factor PacC controls
828 virulence in the plant pathogen *Fusarium oxysporum*: PacC controls virulence in
829 *Fusarium*. *Molecular Microbiology*. 48:765–779

830 Chalabi Hagkarim, N., and Grand, R. J. 2020. The Regulatory Properties of the Ccr4–Not
831 Complex. *Cells*. 9:2379

832 Cheng, C.-Y., Krishnakumar, V., Chan, A. P., Thibaud-Nissen, F., Schobel, S., and
833 Town, C. D. 2017. Araport11: a complete reannotation of the *Arabidopsis*
834 *thaliana* reference genome. *Plant J*. 89:789–804

835 Chikashige, Y., Yamane, M., Okamasa, K., Tsutsumi, C., Kojidani, T., Sato, M.,
836 Haraguchi, T., and Hiraoka, Y. 2009. Membrane proteins Bqt3 and -4 anchor
837 telomeres to the nuclear envelope to ensure chromosomal bouquet formation.
838 *Journal of Cell Biology*. 187:413–427

839 Cuomo, C. A., Guldener, U., Xu, J.-R., Trail, F., Turgeon, B. G., Di Pietro, A., Walton, J.
840 D., Ma, L.-J., Baker, S. E., Rep, M., Adam, G., Antoniw, J., Baldwin, T., Calvo,
841 S., Chang, Y.-L., DeCaprio, D., Gale, L. R., Gnerre, S., Goswami, R. S.,
842 Hammond-Kosack, K., Harris, L. J., Hilburn, K., Kennell, J. C., Kroken, S.,
843 Magnuson, J. K., Mannhaupt, G., Mauceli, E., Mewes, H.-W., Mitterbauer, R.,
844 Muehlbauer, G., Munsterkotter, M., Nelson, D., O'Donnell, K., Ouellet, T., Qi,

845 W., Quesneville, H., Roncero, M. I. G., Seong, K.-Y., Tetko, I. V., Urban, M.,
846 Waalwijk, C., Ward, T. J., Yao, J., Birren, B. W., and Kistler, H. C. 2007. The
847 *Fusarium graminearum* Genome Reveals a Link Between Localized
848 Polymorphism and Pathogen Specialization. *Science*. 317:1400–1402

849 Dean, R. A., Talbot, N. J., Ebbole, D. J., Farman, M. L., Mitchell, T. K., Orbach, M. J.,
850 Thon, M., Kulkarni, R., Xu, J.-R., Pan, H., Read, N. D., Lee, Y.-H., Carbone, I.,
851 Brown, D., Oh, Y. Y., Donofrio, N., Jeong, J. S., Soanes, D. M., Djonovic, S.,
852 Kolomiets, E., Rehmeier, C., Li, W., Harding, M., Kim, S., Lebrun, M.-H.,
853 Bohnert, H., Coughlan, S., Butler, J., Calvo, S., Ma, L.-J., Nicol, R., Purcell, S.,
854 Nusbaum, C., Galagan, J. E., and Birren, B. W. 2005. The genome sequence of
855 the rice blast fungus *Magnaporthe grisea*. *Nature*. 434:980–986

856 Dean, R., Van Kan, J. A. L., Pretorius, Z. A., Hammond-Kosack, K. E., Di Pietro, A.,
857 Spanu, P. D., Rudd, J. J., Dickman, M., Kahmann, R., Ellis, J., and Foster, G. D.
858 2012. The Top 10 fungal pathogens in molecular plant pathology: Top 10 fungal
859 pathogens. *Molecular Plant Pathology*. 13:414–430

860 DeLulio, G. A., Guo, L., Zhang, Y., Goldberg, J. M., Kistler, H. C., and Ma, L.-J. 2018.
861 Kinome Expansion in the *Fusarium oxysporum* Species Complex Driven by
862 Accessory Chromosomes A.P. Mitchell, ed. *mSphere*. 3:e00231-18,
863 /msphere/3/3/mSphere231-18.atom

864 Ding, Z., Xu, T., Zhu, W., Li, L., and Fu, Q. 2020. A MADS-box transcription factor
865 FoRlm1 regulates aerial hyphal growth, oxidative stress, cell wall biosynthesis
866 and virulence in *Fusarium oxysporum* f. sp. *cubense*. *Fungal Biology*. 124:183–
867 193

868 van der Does, H. C., Fokkens, L., Yang, A., Schmidt, S. M., Langereis, L., Lukasiewicz,
869 J. M., Hughes, T. R., and Rep, M. 2016. Transcription Factors Encoded on Core
870 and Accessory Chromosomes of *Fusarium oxysporum* Induce Expression of
871 Effector Genes E.H. Stukenbrock, ed. *PLoS Genet*. 12:e1006401

872 Donczew, R., Warfield, L., Pacheco, D., Erijman, A., and Hahn, S. 2020. Two roles for
873 the yeast transcription coactivator SAGA and a set of genes redundantly regulated
874 by TFIID and SAGA. *eLife*. 9:e50109

875 Doyle, C. E., Kitty Cheung, H. Y., Spence, K. L., and Saville, B. J. 2016. Unh1, an
876 *Ustilago maydis* Ndt80-like protein, controls completion of tumor maturation,
877 teliospore development, and meiosis. *Fungal Genetics and Biology*. 94:54–68

878 Dufresne, M., Perfect, S., Pellier, A.-L., Bailey, J. A., and Langin, T. 2000. A GAL4-like
879 Protein Is Involved in the Switch between Biotrophic and Necrotrophic Phases of
880 the Infection Process of *Colletotrichum lindemuthianum* on Common Bean. *Plant*
881 *Cell*. 12:1579–1590

882 Edel-Hermann, V., and Lecomte, C. 2019. Current Status of *Fusarium oxysporum*
883 *Formae Speciales* and Races. *Phytopathology*. 109:512–530

884 Edskes, H. K., Ohtake, Y., and Wickner, R. B. 1998. Mak21p of *Saccharomyces*
885 *cerevisiae*, a Homolog of Human CAATT-binding Protein, Is Essential for 60 S
886 Ribosomal Subunit Biogenesis. *Journal of Biological Chemistry*. 273:28912–
887 28920

888 Emms, D. M., and Kelly, S. 2019. OrthoFinder: phylogenetic orthology inference for
889 comparative genomics. *Genome Biol*. 20:238

- 890 Feder, M. E., and Hofmann, G. E. 1999. HEAT-SHOCK PROTEINS, MOLECULAR
891 CHAPERONES, AND THE STRESS RESPONSE: Evolutionary and Ecological
892 Physiology. *Annu. Rev. Physiol.* 61:243–282
- 893 Fedotova, A. A., Bonchuk, A. N., Mogila, V. A., and Georgiev, P. G. 2017. C2H2 Zinc
894 Finger Proteins: The Largest but Poorly Explored Family of Higher Eukaryotic
895 Transcription Factors. *Acta Naturae.* 9:47–58
- 896 Fokkens, L., Guo, L., Dora, S., Wang, B., Ye, K., Sánchez-Rodríguez, C., and Croll, D.
897 2021. A Chromosome-Scale Genome Assembly for the *Fusarium oxysporum*
898 Strain Fo5176 To Establish a Model Arabidopsis-Fungal Pathosystem. :7
- 899 Fribourg, S., Kellenberger, E., Rogniaux, H., Poterszman, A., Van Dorsselaer, A.,
900 Thierry, J.-C., Egly, J.-M., Moras, D., and Kieffer, B. 2000. Structural
901 Characterization of the Cysteine-rich Domain of TFIIF p44 Subunit. *Journal of*
902 *Biological Chemistry.* 275:31963–31971
- 903 Fromont-Racine, M., Senger, B., Saveanu, C., and Fasiolo, F. 2003. Ribosome assembly
904 in eukaryotes. *Gene.* 313:17–42
- 905 Galagan, J. E., Calvo, S. E., Borkovich, K. A., Selker, E. U., Read, N. D., Jaffe, D.,
906 FitzHugh, W., Ma, L.-J., Smirnov, S., Purcell, S., Rehman, B., Elkins, T., Engels,
907 R., Wang, S., Nielsen, C. B., Butler, J., Endrizzi, M., Qui, D., Ianakiev, P., Bell-
908 Pedersen, D., Nelson, M. A., Werner-Washburne, M., Selitrennikoff, C. P.,
909 Kinsey, J. A., Braun, E. L., Zelter, A., Schulte, U., Kothe, G. O., Jedd, G.,
910 Mewes, W., Staben, C., Marcotte, E., Greenberg, D., Roy, A., Foley, K., Naylor,
911 J., Stange-Thomann, N., Barrett, R., Gnerre, S., Kamal, M., Kamvysselis, M.,
912 Mauceli, E., Bielke, C., Rudd, S., Frishman, D., Krystofova, S., Rasmussen, C.,
913 Metzenberg, R. L., Perkins, D. D., Kroken, S., Cogoni, C., Macino, G.,
914 Catcheside, D., Li, W., Pratt, R. J., Osmani, S. A., DeSouza, C. P. C., Glass, L.,
915 Orbach, M. J., Berglund, J. A., Voelker, R., Yarden, O., Plamann, M., Seiler, S.,
916 Dunlap, J., Radford, A., Aramayo, R., Natvig, D. O., Alex, L. A., Mannhaupt, G.,
917 Ebbole, D. J., Freitag, M., Paulsen, I., Sachs, M. S., Lander, E. S., Nusbaum, C.,
918 and Birren, B. 2003. The genome sequence of the filamentous fungus *Neurospora*
919 *crassa*. *Nature.* 422:859–868
- 920 Galagan, J. E., Calvo, S. E., Cuomo, C., Ma, L.-J., Wortman, J. R., Batzoglou, S., Lee,
921 S.-I., Baştürkmen, M., Spevak, C. C., Clutterbuck, J., Kapitonov, V., Jurka, J.,
922 Scazzocchio, C., Farman, M., Butler, J., Purcell, S., Harris, S., Braus, G. H.,
923 Draht, O., Busch, S., D’Enfert, C., Bouchier, C., Goldman, G. H., Bell-Pedersen,
924 D., Griffiths-Jones, S., Doonan, J. H., Yu, J., Vienken, K., Pain, A., Freitag, M.,
925 Selker, E. U., Archer, D. B., Peñalva, M. Á., Oakley, B. R., Momany, M., Tanaka,
926 T., Kumagai, T., Asai, K., Machida, M., Nierman, W. C., Denning, D. W.,
927 Caddick, M., Hynes, M., Paoletti, M., Fischer, R., Miller, B., Dyer, P., Sachs, M.
928 S., Osmani, S. A., and Birren, B. W. 2005. Sequencing of *Aspergillus nidulans*
929 and comparative analysis with *A. fumigatus* and *A. oryzae*. *Nature.* 438:1105–
930 1115
- 931 Gallegos, M.-T., Michán, C., and Ramos, J. L. 1993. The XylS/AraC family of
932 regulators. *Nucl Acids Res.* 21:807–810
- 933 Gancedo, J. M. 2001. Control of pseudohyphae formation in *Saccharomyces cerevisiae*.
934 *FEMS Microbiology Reviews.* 25:107–123

935 Gardiner, D. M., McDonald, M. C., Covarelli, L., Solomon, P. S., Rusu, A. G., Marshall,
936 M., Kazan, K., Chakraborty, S., McDonald, B. A., and Manners, J. M. 2012.
937 Comparative Pathogenomics Reveals Horizontally Acquired Novel Virulence
938 Genes in Fungi Infecting Cereal Hosts. *PLOS Pathogens*. 8:e1002952

939 Gimeno, C. J., and Fink, G. R. 1994. Induction of Pseudohyphal Growth by
940 Overexpression of PHD1, a *Saccharomyces cerevisiae* Gene Related to
941 Transcriptional Regulators of Fungal Development. *MOL. CELL. BIOL.* 14:13

942 Goffeau, A., Barrell, B. G., Bussey, H., Davis, R. W., Dujon, B., Feldmann, H., Galibert,
943 F., Hoheisel, J. D., Jacq, C., Johnston, M., Louis, E. J., Mewes, H. W., Murakami,
944 Y., Philippsen, P., Tettelin, H., and Oliver, S. G. 1996. Life with 6000 Genes.
945 *Science*. 274:546–567

946 Goppelt, A., Stelzer, G., Lottspeich, F., and Meisterernst, M. 1996. A mechanism for
947 repression of class II gene transcription through specific binding of NC2 to TBP-
948 promoter complexes via heterodimeric histone fold domains. *The EMBO Journal*.
949 15:3105–3116

950 Gordon, T. R. 2017. *Fusarium oxysporum* and the *Fusarium* Wilt Syndrome. *Annu. Rev.*
951 *Phytopathol.* 55:23–39

952 Green, M. R. 2000. TBP-associated factors (TAF II s): multiple, selective transcriptional
953 mediators in common complexes. *Trends in Biochemical Sciences*. 25:59–63

954 Grigoriev, I. V., Nikitin, R., Haridas, S., Kuo, A., Ohm, R., Otiilar, R., Riley, R.,
955 Salamov, A., Zhao, X., Korzeniewski, F., Smirnova, T., Nordberg, H., Dubchak,
956 I., and Shabalov, I. 2014. MycoCosm portal: gearing up for 1000 fungal genomes.
957 *Nucl. Acids Res.* 42:D699–D704

958 Guo, L., Ji, M., and Ye, K. 2020. Dynamic network inference and association
959 computation discover gene modules regulating virulence, mycotoxin and sexual
960 reproduction in *Fusarium graminearum*. *BMC Genomics*. 21:179

961 Guo, L., Yu, H., Wang, B., Vescio, K., DeJulio, G. A., Yang, H., Berg, A., Zhang, L.,
962 Edel-Hermann, V., Steinberg, C., Kistler, H. C., and Ma, L.-J. 2021.
963 Metatranscriptomic Comparison of Endophytic and Pathogenic *Fusarium* –
964 *Arabidopsis* Interactions Reveals Plant Transcriptional Plasticity. *MPMI*.
965 34:1071–1083

966 Guo, L., Zhao, G., Xu, J., Kistler, H. C., Gao, L., and Ma, L. 2016. Compartmentalized
967 gene regulatory network of the pathogenic fungus *Fusarium graminearum*. *New*
968 *Phytol.* 211:527–541

969 Halpern, H. C., Bell, A. A., Wagner, T. A., Liu, J., Nichols, R. L., Olvey, J., Woodward,
970 J. E., Sanogo, S., Jones, C. A., Chan, C. T., and Brewer, M. T. 2018. First Report
971 of *Fusarium* Wilt of Cotton Caused by *Fusarium oxysporum* f. sp. *vasinfectum*
972 Race 4 in Texas, U.S.A. *Plant Disease*. 102:446–446

973 Hassan, A. S., Al-Hatmi, A. M. S., Shobana, C. S., van Diepeningen, A. D., Kredics, L.,
974 Vágvolgyi, C., Homa, M., Meis, J. F., de Hoog, G. S., Narendran, V.,
975 Manikandan, P., and IHFK Working Group. 2016. Antifungal Susceptibility and
976 Phylogeny of Opportunistic Members of the Genus *Fusarium* Causing Human
977 Keratomycosis in South India. *Med. Myco.* 54:287–294

978 Hepworth, S. R., Friesen, H., and Segall, J. 1998. NDT80 and the Meiotic Recombination
979 Checkpoint Regulate Expression of Middle Sporulation-Specific Genes in
980 *Saccharomyces cerevisiae*. *Mol Cell Biol.* 18:5750–5761

981 Hirschhorn, J. N., Brown, S. A., Clark, C. D., and Winston, F. 1992. Evidence that
982 SNF2/SWI2 and SNF5 activate transcription in yeast by altering chromatin
983 structure. *Genes & Development*. 6:2288–2298

984 Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., and Vinh, L. S. 2018.
985 UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Molecular Biology*
986 *and Evolution*. 35:518–522

987 Honjo, M., Nakayama, A., Fukazawa, K., Kawamura, K., Ando, K., Hori, M., and
988 Furutani, Y. 1990. A novel *Bacillus subtilis* gene involved in negative control of
989 sporulation and degradative-enzyme production. *Journal of Bacteriology*.
990 172:1783–1790

991 Hsu, J., Huang, J., Meluh, P. B., and Laurent, B. C. 2003. The Yeast RSC Chromatin-
992 Remodeling Complex Is Required for Kinetochores Function in Chromosome
993 Segregation. *Mol Cell Biol*. 23:3202–3215

994 Iida, T., and Araki, H. 2004. Noncompetitive Counteractions of DNA Polymerase ϵ and
995 ISW2/yCHRA1 for Epigenetic Inheritance of Telomere Position Effect in
996 *Saccharomyces cerevisiae*. *Mol Cell Biol*. 24:217–227

997 Imazaki, I., Kurahashi, M., Iida, Y., and Tsuge, T. 2007. Fow2, a Zn(II)₂Cys₆-type
998 transcription regulator, controls plant infection of the vascular wilt fungus
999 *Fusarium oxysporum*. *Mol Microbiol*. 63

1000 Iwahara, J. 2002. The structure of the Dead ringer-DNA complex reveals how AT-rich
1001 interaction domains (ARIDs) recognize DNA. *The EMBO Journal*. 21:1197–1209

1002 Jiang, C., Zhang, C., Wu, C., Sun, P., Hou, R., Liu, H., Wang, C., and Xu, J.-R. 2016.
1003 TRI6 and TRI10 play different roles in the regulation of deoxynivalenol (DON)
1004 production by cAMP signalling in *Fusarium graminearum*. *Environmental*
1005 *Microbiology*. 18:3689–3701

1006 Jiang, C., Zhang, S., Zhang, Q., Tao, Y., Wang, C., and Xu, J.-R. 2015. *FgSKN 7* and
1007 *FgATF 1* have overlapping functions in ascospore germination, pathogenesis and stress
1008 responses in *Fusarium graminearum*: Overlapping functions between *FgSKN7*
1009 and *FgATF1*. *Environ Microbiol*. 17:1245–1260

1010 Jones, D. A. B., John, E., Rybak, K., Phan, H. T. T., Singh, K. B., Lin, S.-Y., Solomon,
1011 P. S., Oliver, R. P., and Tan, K.-C. 2019. A specific fungal transcription factor
1012 controls effector gene expression and orchestrates the establishment of the
1013 necrotrophic pathogen lifestyle on wheat. *Sci Rep*. 9:15884

1014 Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H.,
1015 Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas,
1016 A., Scheremetjew, M., Yong, S.-Y., Lopez, R., and Hunter, S. 2014. InterProScan
1017 5: genome-scale protein function classification. *Bioinformatics*. 30:1236–1240

1018 Jones, S. 2004. An overview of the basic helix-loop-helix proteins. *Genome Biology*. :6

1019 Jonkers, W., Xayamongkhon, H., Haas, M., Olivain, C., van der Does, H. C., Broz, K.,
1020 Rep, M., Alabouvette, C., Steinberg, C., and Kistler, H. C. 2014. *EBR1* genomic
1021 expansion and its role in virulence of *Fusarium* species: *EBR1* in virulence of
1022 *Fusarium* species. *Environ Microbiol*. 16:1982–2003

1023 Jung, U. S., Sobering, A. K., Romeo, M. J., and Levin, D. E. 2002. Regulation of the
1024 yeast Rlm1 transcription factor by the Mpk1 cell wall integrity MAP kinase:
1025 Reporters for cell wall integrity signalling. *Molecular Microbiology*. 46:781–789

1026 Kamura, T., Burian, D., Khalili, H., Schmidt, S. L., Sato, S., Liu, W. J., Conrad, M. N.,
1027 Conaway, R. C., Conaway, J. W., and Shilatifard, A. 2001. Cloning and
1028 characterization of ELL-associated proteins EAP45 and EAP20. a role for yeast
1029 EAP-like proteins in regulation of gene expression by glucose. *J Biol Chem.*
1030 276:16528–16533

1031 Katoh, K., and Standley, D. M. 2013. MAFFT Multiple Sequence Alignment Software
1032 Version 7: Improvements in Performance and Usability. *Molecular Biology and*
1033 *Evolution.* 30:772–780

1034 Kim, D., Paggi, J. M., Park, C., Bennett, C., and Salzberg, S. L. 2019. Graph-based
1035 genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat*
1036 *Biotechnol.* 37:907–915

1037 Kim, H.-J., Lee, K.-L., Kim, K.-D., and Roe, J.-H. 2016. The iron uptake repressor Fep1
1038 in the fission yeast binds Fe-S cluster through conserved cysteines. *Biochemical*
1039 *and Biophysical Research Communications.* 478:187–192

1040 Kim, S., Park, S.-Y., Kim, K. S., Rho, H.-S., Chi, M.-H., Choi, J., Park, J., Kong, S.,
1041 Park, J., Goh, J., and Lee, Y.-H. 2009. Homeobox Transcription Factors Are
1042 Required for Conidiation and Appressorium Development in the Rice Blast
1043 Fungus *Magnaporthe oryzae* G.P. Copenhaver, ed. *PLoS Genet.* 5:e1000757

1044 Koch, C., Moll, T., Neuberg, M., Ahorn, H., and Nasmyth, K. 1993. A Role for the
1045 Transcription Factors Mbp1 and Swi4 in Progression from G1 to S Phase.
1046 *Science.* 261:1551–1557

1047 Kredics, L., Narendran, V., Shobana, C. S., Vágvolgyi, C., Manikandan, P., and Indo-
1048 Hungarian Fungal Keratitis Working Group. 2015. Filamentous fungal infections
1049 of the cornea: a global overview of epidemiology and drug sensitivity. *Mycoses.*
1050 58:243–260

1051 Latrick, C. M., Marek, M., Ouararhni, K., Papin, C., Stoll, I., Ignatyeva, M., Obri, A.,
1052 Ennifar, E., Dimitrov, S., Romier, C., and Hamiche, A. 2016. Molecular basis and
1053 specificity of H2A.Z–H2B recognition and deposition by the histone chaperone
1054 YL1. *Nat Struct Mol Biol.* 23:309–316

1055 Letunic, I., and Bork, P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for
1056 phylogenetic tree display and annotation. *Nucleic Acids Research.* 49:W293–
1057 W296

1058 Li, G., Zhou, X., Kong, L., Wang, Y., Zhang, H., Zhu, H., Mitchell, T. K., Dean, R. A.,
1059 and Xu, J.-R. 2011. MoSfl1 Is Important for Virulence and Heat Tolerance in
1060 *Magnaporthe oryzae* A. Idnurm, ed. *PLoS ONE.* 6:e19951

1061 Li, X., Han, X., Liu, Z., and He, C. 2013. The function and properties of the
1062 transcriptional regulator COS1 in *Magnaporthe oryzae*. *Fungal Biology.* 117:239–
1063 249

1064 Liang, X., Shan, S., Pan, L., Zhao, J., Ranjan, A., Wang, F., Zhang, Z., Huang, Y., Feng,
1065 H., Wei, D., Huang, L., Liu, X., Zhong, Q., Lou, J., Li, G., Wu, C., and Zhou, Z.
1066 2016. Structural basis of H2A.Z recognition by SRCAP chromatin-remodeling
1067 subunit YL1. *Nat Struct Mol Biol.* 23:317–323

1068 Liu, J., Yuan, Y., Wu, Z., Li, N., Chen, Y., Qin, T., Geng, H., Xiong, L., and Liu, D.
1069 2015. A Novel Sterol Regulatory Element-Binding Protein Gene (*sreA*) Identified
1070 in *Penicillium digitatum* Is Required for Prochloraz Resistance, Full Virulence
1071 and *erg11* (*cyp51*) Regulation R.A. Cramer, ed. *PLoS ONE.* 10:e0117115

1072 López-Berges, M. S. 2020. ZafA-mediated regulation of zinc homeostasis is required for
1073 virulence in the plant pathogen *Fusarium oxysporum*. *Molecular Plant Pathology*.
1074 21:244–249

1075 López-Berges, M. S., Capilla, J., Turrà, D., Schaffner, L., Matthijs, S., Jöchl, C.,
1076 Cornelis, P., Guarro, J., Haas, H., and Di Pietro, A. 2012. HapX-Mediated Iron
1077 Homeostasis Is Essential for Rhizosphere Competence and Virulence of the
1078 Soilborne Pathogen *Fusarium oxysporum*. *The Plant Cell*. 24:3805–3822

1079 López-Berges, M. S., Rispaill, N., Prados-Rosales, R. C., and Di Pietro, A. 2010. A
1080 Nitrogen Response Pathway Regulates Virulence Functions in *Fusarium*
1081 *oxysporum* via the Protein Kinase TOR and the bZIP Protein MeaB. *Plant Cell*.
1082 22:2459–2475

1083 Lubelsky, Y., Reuven, N., and Shaul, Y. 2005. Autorepression of Rfx1 Gene Expression:
1084 Functional Conservation from Yeast to Humans in Response to DNA Replication
1085 Arrest. *Mol Cell Biol*. 25:10665–10673

1086 Lysøe, E., Pasquali, M., Breakspear, A., and Kistler, H. C. 2011. The Transcription
1087 Factor FgStuAp Influences Spore Development, Pathogenicity, and Secondary
1088 Metabolism in *Fusarium graminearum*. *MPMI*. 24:54–67

1089 Ma, L.-J. 2014. Horizontal chromosome transfer and rational strategies to manage
1090 *Fusarium* vascular wilt diseases: Translating genomics for *Fusarium* vascular
1091 wilts. *Molecular Plant Pathology*. 15:763–766

1092 Ma, L.-J., van der Does, H. C., Borkovich, K. A., Coleman, J. J., Daboussi, M.-J., Di
1093 Pietro, A., Dufresne, M., Freitag, M., Grabherr, M., Henrissat, B., Houterman, P.
1094 M., Kang, S., Shim, W.-B., Woloshuk, C., Xie, X., Xu, J.-R., Antoniw, J., Baker,
1095 S. E., Bluhm, B. H., Breakspear, A., Brown, D. W., Butchko, R. A. E., Chapman,
1096 S., Coulson, R., Coutinho, P. M., Danchin, E. G. J., Diener, A., Gale, L. R.,
1097 Gardiner, D. M., Goff, S., Hammond-Kosack, K. E., Hilburn, K., Hua-Van, A.,
1098 Jonkers, W., Kazan, K., Kodira, C. D., Koehrsen, M., Kumar, L., Lee, Y.-H., Li,
1099 L., Manners, J. M., Miranda-Saavedra, D., Mukherjee, M., Park, G., Park, J.,
1100 Park, S.-Y., Proctor, R. H., Regev, A., Ruiz-Roldan, M. C., Sain, D.,
1101 Sakthikumar, S., Sykes, S., Schwartz, D. C., Turgeon, B. G., Wapinski, I., Yoder,
1102 O., Young, S., Zeng, Q., Zhou, S., Galagan, J., Cuomo, C. A., Kistler, H. C., and
1103 Rep, M. 2010. Comparative genomics reveals mobile pathogenicity chromosomes
1104 in *Fusarium*. *Nature*. 464:367–373

1105 Ma, L.-J., Geiser, D. M., Proctor, R. H., Rooney, A. P., O'Donnell, K., Trail, F.,
1106 Gardiner, D. M., Manners, J. M., and Kazan, K. 2013. *Fusarium* Pathogenomics.
1107 *Annu. Rev. Microbiol*. 67:399–416

1108 Ma, L.-J., Shea, T., Young, S., Zeng, Q., and Kistler, H. C. 2014. Genome Sequence of
1109 *Fusarium oxysporum* f. sp. *melonis* Strain NRRL 26406, a Fungus Causing Wilt
1110 Disease on Melon. *Genome Announcements*. 2:e00730-14, 2/4/e00730-14

1111 MacPherson, S., Laroche, M., and Turcotte, B. 2006. A Fungal Family of
1112 Transcriptional Regulators: the Zinc Cluster Proteins. *Microbiol Mol Biol Rev*.
1113 70:583–604

1114 Maddock, J. R., Weidenhammer, E. M., Adams, C. C., Lunz, R. L., and Woolford, J. L.
1115 1994. Extragenic suppressors of *Saccharomyces cerevisiae* *prp4* mutations
1116 identify a negative regulator of PRP genes. *Genetics*. 136:833–847

1117 Martin, T., Lu, S.-W., Tilbeurgh, H. van, Ripoll, D. R., Dixelius, C., Turgeon, B. G., and
1118 Debuchy, R. 2010. Tracing the Origin of the Fungal α 1 Domain Places Its
1119 Ancestor in the HMG-Box Superfamily: Implication for Fungal Mating-Type
1120 Evolution. PLOS ONE. 5:e15199

1121 Martínez-Soto, D., Yu, H., Allen, K. S., and Ma, L.-J. 2022. Differential colonization of
1122 the plant vasculature between endophytic versus pathogenic *Fusarium oxysporum*
1123 strains. MPMI.

1124 Mendoza-Mendoza, A., Eskova, A., Weise, C., Czajkowski, R., and Kahmann, R. 2009.
1125 Hap2 regulates the pheromone response transcription factor prf1 in *Ustilago*
1126 maydis. Molecular Microbiology. 72:683–698

1127 Mercier, A., Watt, S., Bähler, J., and Labbé, S. 2008. Key Function for the CCAAT-
1128 Binding Factor Php4 To Regulate Gene Expression in Response to Iron
1129 Deficiency in Fission Yeast. Eukaryot Cell. 7:493–508

1130 Mesny, F., Miyauchi, S., Thiergart, T., Pickel, B., Atanasova, L., Karlsson, M., Hüttel,
1131 B., Barry, K. W., Haridas, S., Chen, C., Bauer, D., Andreopoulos, W., Pangilinan,
1132 J., LaButti, K., Riley, R., Lipzen, A., Clum, A., Drula, E., Henrissat, B., Kohler,
1133 A., Grigoriev, I. V., Martin, F. M., and Hacquard, S. 2021. Genetic determinants
1134 of endophytism in the Arabidopsis root mycobiome. Nat Commun. 12:7227

1135 Messenguy, F., and Dubois, E. 2003. Role of MADS box proteins and their cofactors in
1136 combinatorial control of gene expression and cell development. Gene. 316:1–21

1137 Michielse, C. B., and Rep, M. 2009. Pathogen profile update: *Fusarium oxysporum*.
1138 Molecular Plant Pathology. 10:311–324

1139 Michielse, C. B., van Wijk, R., Reijnen, L., Manders, E. M. M., Boas, S., Olivain, C.,
1140 Alabouvette, C., and Rep, M. 2009. The Nuclear Protein Sge1 of *Fusarium*
1141 *oxysporum* Is Required for Parasitic Growth B.J. Howlett, ed. PLoS Pathog.
1142 5:e1000637

1143 Milkereit, P., Gadai, O., Podtelejnikov, A., Trumtel, S., Gas, N., Petfalski, E., Tollervey,
1144 D., Mann, M., Hurt, E., and Tschochner, H. 2001. Maturation and Intranuclear
1145 Transport of Pre-Ribosomes Requires Noc Proteins. Cell. 105:499–509

1146 Min, K., Son, H., Lim, J. Y., Choi, G. J., Kim, J.-C., Harris, S. D., and Lee, Y.-W. 2014.
1147 Transcription Factor RFX1 Is Crucial for Maintenance of Genome Integrity in
1148 *Fusarium graminearum*. Eukaryot Cell. 13:427–436

1149 Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von
1150 Haeseler, A., and Lanfear, R. 2020. IQ-TREE 2: New Models and Efficient
1151 Methods for Phylogenetic Inference in the Genomic Era. Molecular Biology and
1152 Evolution. 37:1530–1534

1153 Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., and Minh, B. Q. 2015. IQ-TREE: A
1154 Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood
1155 Phylogenies. Molecular Biology and Evolution. 32:268–274

1156 Niño-Sánchez, J., Casado-Del Castillo, V., Tello, V., De Vega-Bartol, J. J., Ramos, B.,
1157 Sukno, S. A., and Díaz Mínguez, J. M. 2016. The *FTF* gene family regulates
1158 virulence and expression of SIX effectors in *Fusarium oxysporum*. Molecular
1159 Plant Pathology. 17:1124–1139

1160 Oh, M., Son, H., Choi, G. J., Lee, C., Kim, J.-C., Kim, H., and Lee, Y.-W. 2016.
1161 Transcription factor ART1 mediates starch hydrolysis and mycotoxin production

1162 in *Fusarium graminearum* and *F. verticillioides*. *Molecular Plant Pathology*.
1163 17:755–768

1164 Ohara, T., Inoue, I., Namiki, F., Kunoh, H., and Tsuge, T. 2004. *REN1* Is Required for
1165 Development of Microconidia and Macroconidia, but Not of Chlamydospores, in
1166 the Plant Pathogenic Fungus *Fusarium oxysporum*. *Genetics*. 166:113–124

1167 Olesen, J. T., Fikes, J. D., and Guarente, L. 1991. The *Schizosaccharomyces pombe*
1168 homolog of *Saccharomyces cerevisiae* HAP2 reveals selective and stringent
1169 conservation of the small essential core protein domain. 11:9

1170 Ortiz, C. S., and Shim, W.-B. 2013. The role of MADS-box transcription factors in
1171 secondary metabolism and sexual development in the maize pathogen *Fusarium*
1172 *verticillioides*. *Microbiology*. 159:2259–2268

1173 Pan, X., and Heitman, J. 2000. Sok2 Regulates Yeast Pseudohyphal Differentiation via a
1174 Transcription Factor Cascade That Regulates Cell-Cell Adhesion. *Mol Cell Biol*.
1175 20:8364–8372

1176 Paré, A., Kim, M., Juarez, M. T., Brody, S., and McGinnis, W. 2012. The Functions of
1177 Grainy Head-Like Proteins in Animals and Fungi and the Evolution of Apical
1178 Extracellular Barriers J.E. Stajich, ed. *PLoS ONE*. 7:e36254

1179 Park, J., Park, J., Jang, S., Kim, S., Kong, S., Choi, J., Ahn, K., Kim, J., Lee, S., Kim, S.,
1180 Park, B., Jung, K., Kim, S., Kang, S., and Lee, Y.-H. 2008. FTFD: an informatics
1181 pipeline supporting phylogenomic analysis of fungal transcription factors.
1182 *Bioinformatics*. 24:1024–1025

1183 Pegg, K. G., Coates, L. M., O'Neill, W. T., and Turner, D. W. 2019. The Epidemiology
1184 of *Fusarium* Wilt of Banana. *Front. Plant Sci*. 10:1395

1185 Perteua, M., Perteua, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T., and Salzberg,
1186 S. L. 2015. StringTie enables improved reconstruction of a transcriptome from
1187 RNA-seq reads. *Nat Biotechnol*. 33:290–295

1188 Pierce, M., Benjamin, K. R., Montano, S. P., Georgiadis, M. M., Winter, E., and
1189 Vershon, A. K. 2003. Sum1 and Ndt80 Proteins Compete for Binding to Middle
1190 Sporulation Element Sequences That Control Meiotic Gene Expression. *Mol Cell*
1191 *Biol*. 23:4814–4825

1192 Ploetz, R. C. 2015. *Fusarium* Wilt of Banana. *Phytopathology*. 105:1512–1521

1193 Rahman, M. Z., Ahmad, K., Siddiqui, Y., Saad, N., Hun, T. G., Mohd Hata, E., Rashed,
1194 O., Hossain, M. I., and Kutawa, A. B. 2021. First Report of *Fusarium* wilt disease
1195 on Watermelon Caused by *Fusarium oxysporum* f. sp. *niveum* (FON) in Malaysia.
1196 *Plant Dis*.

1197 Ramos, B., Alves-Santos, F. M., García-Sánchez, M. A., Martín-Rodrigues, N., Eslava,
1198 A. P., and Díaz-Mínguez, J. M. 2007. The gene coding for a new transcription
1199 factor (ftf1) of *Fusarium oxysporum* is only expressed during infection of
1200 common bean. *Fungal Genetics and Biology*. 44:864–876

1201 Redkar, A., Sabale, M., Schudoma, C., Zechmann, B., Gupta, Y. K., López-Berges, M.
1202 S., Venturini, G., Gimenez-Ibanez, S., Turrà, D., Solano, R., and Di Pietro, A.
1203 2022. Conserved secreted effectors contribute to endophytic growth and multihost
1204 plant compatibility in a vascular wilt fungus. *The Plant Cell*. :koac174

1205 Rep, M., Van Der Does, H. C., Meijer, M., Van Wijk, R., Houterman, P. M., Dekker, H.
1206 L., De Koster, C. G., and Cornelissen, B. J. C. 2004. A small, cysteine-rich
1207 protein secreted by *Fusarium oxysporum* during colonization of xylem vessels is

1208 required for I-3-mediated resistance in tomato: Cysteine-rich protein required for
1209 I-3-mediated resistance. *Molecular Microbiology*. 53:1373–1383

1210 Ridenour, J. B., and Bluhm, B. H. 2014. The HAP complex in *Fusarium verticillioides* is
1211 a key regulator of growth, morphogenesis, secondary metabolism, and
1212 pathogenesis. *Fungal Genetics and Biology*. 69:52–64

1213 Ridenour, J. B., and Bluhm, B. H. 2017. The novel fungal-specific gene *FUG1* has a role
1214 in pathogenicity and fumonisin biosynthesis in *Fusarium verticillioides*.
1215 *Molecular Plant Pathology*. 18:513–528

1216 Risipail, N., and Di Pietro, A. 2009. *Fusarium oxysporum* Ste12 Controls Invasive
1217 Growth and Virulence Downstream of the Fmk1 MAPK Cascade. *MPMI*.
1218 22:830–839

1219 Risipail, N., and Di Pietro, A. 2010. The homeodomain transcription factor Ste12:
1220 Connecting fungal MAPK signaling to plant pathogenicity. *Communicative &*
1221 *Integrative Biology*. 3:327–332

1222 Roberts, S. M., and Winston, F. 1996. SPT20/ADA5 encodes a novel protein functionally
1223 related to the TATA-binding protein and important for transcription in
1224 *Saccharomyces cerevisiae*. *Mol Cell Biol*. 16:3206–3213

1225 Rocha, A. L. M., Di Pietro, A., Ruiz-Roldán, C., and Roncero, M. I. G. 2008. Ctf1, a
1226 transcriptional activator of cutinase and lipase genes in *Fusarium oxysporum* is
1227 dispensable for virulence. *Molecular Plant Pathology*. 9:293–304

1228 Ruiz-Roldán, C., Pareja-Jaime, Y., González-Reyes, J. A., and G. Roncero, M. I. 2015.
1229 The Transcription Factor Con7-1 Is a Master Regulator of Morphogenesis and
1230 Virulence in *Fusarium oxysporum*. *MPMI*. 28:55–68

1231 Schleif, R. 2010. AraC protein, regulation of the l-arabinose operon in *Escherichia coli* ,
1232 and the light switch mechanism of AraC action. *FEMS Microbiol Rev*. 34:779–
1233 796

1234 Schumacher, J., Simon, A., Cohrs, K. C., Viaud, M., and Tudzynski, P. 2014. The
1235 Transcription Factor BcLTF1 Regulates Virulence and Light Responses in the
1236 Necrotrophic Plant Pathogen *Botrytis cinerea*. *PLOS Genetics*. 10:e1004040

1237 Shelest, E. 2017. Transcription Factors in Fungi: TFome Dynamics, Three Major
1238 Families, and Dual-Specificity TFs. *Front. Genet*. 8:53

1239 Siegmund, T., and Lehmann, M. 2002. The *Drosophila* Pipsqueak protein defines a new
1240 family of helix-turn-helix DNA-binding proteins. *Dev Genes Evol*. 212:152–157

1241 Son, H., Fu, M., Lee, Y., Lim, J. Y., Min, K., Kim, J.-C., Choi, G. J., and Lee, Y.-W.
1242 2016a. A novel transcription factor gene *FHS1* is involved in the DNA damage
1243 response in *Fusarium graminearum*. *Sci Rep*. 6:21572

1244 Son, M., Lee, Y., and Kim, K.-H. 2016b. The Transcription Cofactor Swi6 of the
1245 *Fusarium graminearum* Is Involved in *Fusarium Graminearum* Virus 1 Infection-
1246 Induced Phenotypic Alterations. *The Plant Pathology Journal*. 32:281–289

1247 Song, Z., Krishna, S., Thanos, D., Strominger, J. L., and Ono, S. J. 1994. A novel
1248 cysteine-rich sequence-specific DNA-binding protein interacts with the conserved
1249 X-box motif of the human major histocompatibility complex class II genes via a
1250 repeated Cys-His domain and functions as a transcriptional repressor. *J Exp Med*.
1251 180:1763–1774

1252 Sugiyama, M., and Nikawa, J.-I. 2001. The *Saccharomyces cerevisiae* Isw2p-Itc1p
1253 Complex Represses *INO1* Expression and Maintains Cell Morphology. J
1254 Bacteriol. 183:4985–4993

1255 Takemaru, K., Li, F.-Q., Ueda, H., and Hirose, S. 1997. Multiprotein bridging factor 1
1256 (MBF1) is an evolutionarily conserved transcriptional coactivator that connects a
1257 regulatory factor and TATA element-binding protein. Proc. Natl. Acad. Sci.
1258 U.S.A. 94:7251–7256

1259 Tsuchiya, D., Yang, Y., and Lacefield, S. 2014. Positive Feedback of NDT80 Expression
1260 Ensures Irreversible Meiotic Commitment in Budding Yeast M. Lichten, ed.
1261 PLoS Genet. 10:e1004398

1262 de Vega-Bartol, J. J., Martín-Dominguez, R., Ramos, B., García-Sánchez, M.-A., and
1263 Díaz-Minguez, J. M. 2011. New Virulence Groups in *Fusarium oxysporum* f. sp.
1264 *phaseoli* : The Expression of the Gene Coding for the Transcription Factor *ftf1*
1265 Correlates with Virulence. Phytopathology. 101:470–479

1266 Vesth, T. C., Nybo, J. L., Theobald, S., Frisvad, J. C., Larsen, T. O., Nielsen, K. F., Hoof,
1267 J. B., Brandl, J., Salamov, A., Riley, R., Gladden, J. M., Phatale, P., Nielsen, M.
1268 T., Lyhne, E. K., Kogle, M. E., Strasser, K., McDonnell, E., Barry, K., Clum, A.,
1269 Chen, C., LaButti, K., Haridas, S., Nolan, M., Sandor, L., Kuo, A., Lipzen, A.,
1270 Hainaut, M., Drula, E., Tsang, A., Magnuson, J. K., Henrissat, B., Wiebenga, A.,
1271 Simmons, B. A., Mäkelä, M. R., de Vries, R. P., Grigoriev, I. V., Mortensen, U.
1272 H., Baker, S. E., and Andersen, M. R. 2018. Investigation of inter- and
1273 intraspecies variation through genome sequencing of *Aspergillus* section *Nigri*.
1274 Nat Genet. 50:1688–1695

1275 Viljoen, A., Mostert, D., Chiconela, T., Beukes, I., Fraser, C., Dwyer, J., Murray, H.,
1276 Amisse, J., Matabuana, E. L., Tazan, G., Amugoli, O. M., Mondjana, A., Vaz, A.,
1277 Pretorius, A., Bothma, S., Rose, L. J., Beed, F., Dusunceli, F., Chao, C.-P., and
1278 Molina, A. B. 2020. Occurrence and spread of the banana fungus *Fusarium*
1279 *oxysporum* f. sp. *cubense* TR4 in Mozambique. South African Journal of Science.
1280 116:1–11

1281 Vizoso-Vázquez, Á., Lamas-Maceiras, M., Becerra, M., González-Siso, M. I., Rodríguez-
1282 Belmonte, E., and Cerdán, M. E. 2012. *Ixr1p* and the control of the
1283 *Saccharomyces cerevisiae* hypoxic response. Appl Microbiol Biotechnol. 94:173–
1284 184

1285 Vuorio, T., Maity, S. N., and de Crombrughe, B. 1990. Purification and molecular
1286 cloning of the “A” chain of a rat heteromeric CCAAT-binding protein. Sequence
1287 identity with the yeast HAP3 transcription factor. Journal of Biological
1288 Chemistry. 265:22480–22486

1289 Wang, B., Yu, H., Jia, Y., Dong, Q., Steinberg, C., Alabouvette, C., Edel-Hermann, V.,
1290 Kistler, H. C., Ye, K., Ma, L.-J., and Guo, L. 2020. Chromosome-Scale Genome
1291 Assembly of *Fusarium oxysporum* Strain Fo47, a Fungal Endophyte and
1292 Biocontrol Agent. MPMI. :MPMI-05-20-0116

1293 Wang, Q., Pokhrel, A., and Coleman, J. J. 2021. The Extracellular Superoxide Dismutase
1294 *Sod5* From *Fusarium oxysporum* Is Localized in Response to External Stimuli
1295 and Contributes to Fungal Pathogenicity. Front Plant Sci. 12:608861

1296 Wiemann, P., Sieber, C. M. K., Barga, K. W. von, Studt, L., Niehaus, E.-M., Espino, J.
1297 J., Huß, K., Michielse, C. B., Albermann, S., Wagner, D., Bergner, S. V.,

- 1298 Connolly, L. R., Fischer, A., Reuter, G., Kleigrew, K., Bald, T., Wingfield, B.
1299 D., Ophir, R., Freeman, S., Hippler, M., Smith, K. M., Brown, D. W., Proctor, R.
1300 H., Münsterkötter, M., Freitag, M., Humpf, H.-U., Güldener, U., and Tudzynski,
1301 B. 2013. Deciphering the Cryptic Genome: Genome-wide Analyses of the Rice
1302 Pathogen *Fusarium fujikuroi* Reveal Complex Regulation of Secondary
1303 Metabolism and Novel Metabolites. *PLOS Pathogens*. 9:e1003475
- 1304 Wight, W. D., Kim, K.-H., Lawrence, C. B., and Walton, J. D. 2009. Biosynthesis and
1305 role in virulence of the histone deacetylase inhibitor depudecin from *Alternaria*
1306 *brassicicola*. *Mol Plant Microbe Interact*. 22:1258–1267
- 1307 Williams, A. H., Sharma, M., Thatcher, L. F., Azam, S., Hane, J. K., Sperschneider, J.,
1308 Kidd, B. N., Anderson, J. P., Ghosh, R., Garg, G., Lichtenzveig, J., Kistler, H. C.,
1309 Shea, T., Young, S., Buck, S.-A. G., Kamphuis, L. G., Saxena, R., Pande, S., Ma,
1310 L.-J., Varshney, R. K., and Singh, K. B. 2016. Comparative genomics and
1311 prediction of conditionally dispensable sequences in legume-infecting *Fusarium*
1312 *oxysporum* formae speciales facilitates identification of candidate effectors. *BMC*
1313 *Genomics*. 17:191
- 1314 Wood, V., Gwilliam, R., Rajandream, M.-A., Lyne, M., Lyne, R., Stewart, A., Sgouros,
1315 J., Peat, N., Hayles, J., Baker, S., Basham, D., Bowman, S., Brooks, K., Brown,
1316 D., Brown, S., Chillingworth, T., Churcher, C., Collins, M., Connor, R., Cronin,
1317 A., Davis, P., Feltwell, T., Fraser, A., Gentles, S., Goble, A., Hamlin, N., Harris,
1318 D., Hidalgo, J., Hodgson, G., Holroyd, S., Hornsby, T., Howarth, S., Huckle, E.
1319 J., Hunt, S., Jagels, K., James, K., Jones, L., Jones, M., Leather, S., McDonald, S.,
1320 McLean, J., Mooney, P., Moule, S., Mungall, K., Murphy, L., Niblett, D., Odell,
1321 C., Oliver, K., O’Neil, S., Pearson, D., Quail, M. A., Rabinowitsch, E.,
1322 Rutherford, K., Rutter, S., Saunders, D., Seeger, K., Sharp, S., Skelton, J.,
1323 Simmonds, M., Squares, R., Squares, S., Stevens, K., Taylor, K., Taylor, R. G.,
1324 Tivey, A., Walsh, S., Warren, T., Whitehead, S., Woodward, J., Volckaert, G.,
1325 Aert, R., Robben, J., Grymonprez, B., Weltjens, I., Vanstreels, E., Rieger, M.,
1326 Schäfer, M., Müller-Auer, S., Gabel, C., Fuchs, M., Fritz, C., Holzer, E., Moestl,
1327 D., Hilbert, H., Borzym, K., Langer, I., Beck, A., Lehrach, H., Reinhardt, R.,
1328 Pohl, T. M., Eger, P., Zimmermann, W., Wedler, H., Wambutt, R., Purnelle, B.,
1329 Goffeau, A., Cadieu, E., Dréano, S., Gloux, S., Lelaure, V., Mottier, S., Galibert,
1330 F., Aves, S. J., Xiang, Z., Hunt, C., Moore, K., Hurst, S. M., Lucas, M., Rochet,
1331 M., Gaillardin, C., Tallada, V. A., Garzon, A., Thode, G., Daga, R. R., Cruzado,
1332 L., Jimenez, J., Sánchez, M., del Rey, F., Benito, J., Domínguez, A., Revuelta, J.
1333 L., Moreno, S., Armstrong, J., Forsburg, S. L., Cerrutti, L., Lowe, T., McCombie,
1334 W. R., Paulsen, I., Potashkin, J., Shpakovski, G. V., Ussery, D., Barrell, B. G.,
1335 and Nurse, P. 2002. The genome sequence of *Schizosaccharomyces pombe*.
1336 *Nature*. 415:871–880
- 1337 Xin, C., Zhang, J., Nian, S., Wang, G., Wang, Z., Song, Z., and Ren, G. 2020. Analogous
1338 and Diverse Functions of APSES-Type Transcription Factors in the
1339 Morphogenesis of the Entomopathogenic Fungus *Metarhizium rileyi* N.-Y. Zhou,
1340 ed. *Appl Environ Microbiol*. 86:e02928-19
- 1341 Yamaguchi, Y., Narita, T., Inukai, N., Wada, T., and Handa, H. 2001. SPT Genes: Key
1342 Players in the Regulation of Transcription, Chromatin Structure and Other
1343 Cellular Processes. *Journal of Biochemistry*. 129:185–191

- 1344 Yan, X., Li, Y., Yue, X., Wang, C., Que, Y., Kong, D., Ma, Z., Talbot, N. J., and Wang,
1345 Z. 2011. Two Novel Transcriptional Regulators Are Essential for Infection-
1346 related Morphogenesis and Pathogenicity of the Rice Blast Fungus *Magnaporthe*
1347 *oryzae*. *PLOS Pathogens*. 7:e1002385
- 1348 Yang, H. 2020. ACCESSORY GENES CONTRIBUTE TO REWIRING THE
1349 TRANSCRIPTIONAL NETWORK IN *FUSARIUM OXYSPORUM*. Doctoral
1350 Dissertations.
- 1351 Yang, H., Yu, H., and Ma, L.-J. 2020. Accessory Chromosomes in *Fusarium oxysporum*.
1352 *Phytopathology*®. :PHYTO-03-20-006
- 1353 Yu, H., Ayhan, D. H., Diener, A. C., and Ma, L.-J. 2020. Genome Sequence of *Fusarium*
1354 *oxysporum* f. sp. *matthiolae* , a Brassicaceae Pathogen. *MPMI*. 33:569–572
- 1355 Yu, H., Ayhan, D. H., Martínez-Soto, D., Cochavi, S. M., and Ma, L.-J. 2023. Accessory
1356 Chromosomes of the *Fusarium oxysporum* Species Complex and Their
1357 Contribution to Host Niche Adaptation. Pages 371–388 in: *Plant Relationships,*
1358 *The Mycota*. B. Scott and C. Mesarich, eds. Springer International Publishing,
1359 Cham.
- 1360 Yue, X., Que, Y., Xu, L., Deng, S., Peng, Y., Talbot, N. J., and Wang, Z. 2016. ZNF1
1361 Encodes a Putative C2H2 Zinc-Finger Protein Essential for Appressorium
1362 Differentiation by the Rice Blast Fungus *Magnaporthe oryzae*. *Mol Plant Microbe*
1363 *Interact*. 29:22–35
- 1364 Yun, Y., Zhou, X., Yang, S., Wen, Y., You, H., Zheng, Y., Norvinyeku, J., Shim, W.-B.,
1365 and Wang, Z. 2019. *Fusarium oxysporum* f. sp. *lycopersici* C2H2 transcription
1366 factor *FolCzfl* is required for conidiation, fusaric acid production, and early host
1367 infection. *Curr Genet*. 65:773–783
- 1368 ZHANG, Y. 2019. EVOLUTION OF THE PATHOGENIC *FUSARIUM OXYSPORUM*
1369 THROUGH THE LENS OF COMPARATIVE GENOMICS. Doctoral
1370 Dissertations.
- 1371 Zhang, Y., Yang, H., Turra, D., Zhou, S., Ayhan, D. H., DeIulio, G. A., Guo, L., Broz,
1372 K., Wiederhold, N., Coleman, J. J., Donnell, K. O., Youngster, I., McAdam, A. J.,
1373 Savinov, S., Shea, T., Young, S., Zeng, Q., Rep, M., Pearlman, E., Schwartz, D.
1374 C., Di Pietro, A., Kistler, H. C., and Ma, L.-J. 2020. The genome of opportunistic
1375 fungal pathogen *Fusarium oxysporum* carries a unique set of lineage-specific
1376 chromosomes. *Commun Biol*. 3:1–12
- 1377 Zhao, S., An, B., Guo, Y., Hou, X., Luo, H., He, C., and Wang, Q. 2020. Label free
1378 proteomics and systematic analysis of secretome reveals effector candidates
1379 regulated by *SGE1* and *FTF1* in the plant pathogen *Fusarium oxysporum* f. sp.
1380 *cubense* tropical race 4. *BMC Genomics*. 21:275
- 1381 Zhou, Z., Li, G., Lin, C., and He, C. 2009. *Conidiophore Stalk-less1* Encodes a Putative
1382 Zinc-Finger Protein Involved in the Early Stage of Conidiation and Mycelial
1383 Infection in *Magnaporthe oryzae*. *MPMI*. 22:402–410
- 1384 Zuriegat, Q., Zheng, Y., Liu, H., Wang, Z., and Yun, Y. 2021. Current progress on
1385 pathogenicity-related transcription factors in *Fusarium oxysporum*. *Mol Plant*
1386 *Pathol*. 22:882–895