

Not only age affects cardiovascular parameters, salivary biomarkers, and their correlation, but the level of physical conditioning changes this behavior in the elderly

Phabloo José de Venâncio de Camargos^(ORCID 0000-0002-1568-9053)¹, Allisson Benatti Justino^(ORCID 0000-0003-3616-6611)⁴, Luis Carlos Oliveira Gonçalves^(ORCID 0000-0001-5368-1194)^{1,2,*}, Rafael Joviano Souza de Barros^(ORCID 0000-0003-1038-3888)³, Miguel Junior Sordi Bortolini^(ORCID 0000-0003-0778-1164)⁵, Foued Salmen Espindola^(ORCID 0000-0002-6937-1411)⁴, Aníbal Monteiro de Magalhães Neto^(ORCID 0000-0002-4887-5936)²

¹Graduate Program of Basic and Applied Immunology and Parasitology, Federal University of Mato Grosso-MT, Brazil;

²Graduate Program of Physical Education, Federal University of Mato Grosso-MG, Brazil;

³Physician at the Municipal Health Department of Barra do Garças-MT, Brazil;

⁴Graduate Program in Genetics and Biochemistry, Federal University of Uberlândia-MG, Brazil;

⁵Graduate Program in Health Sciences in the Western Amazon, Federal University of Acre-AC, Brazil.

Corresponding author: Gonçalves LCO, Graduate Program in Physical Education, Federal University of Mato Grosso, Brazil, Tel: 5521 964451022; **Email:** luisogoncalves@yahoo.com.br

ABSTRACT

To compare the physical conditioning, hemodynamic, and salivary biomarkers between elderly athletes and the physically active elderly. 14 men: EA (n = 8) and PAE (n = 6). Collection times (T0; TE; T5; T15). A negative correlation was found between SF and cardiovascular parameters, BL, and STP in both groups, but this was almost double among PAE. For HR and SBP, there was a faster recovery in EA. The EA increase was correlated with SBP, while for PAE it was correlated with HR. BL showed an increase in TE, reaching 481% in EA and 639% in PAE. SNO showed a similar increase for the groups at the TE, but at T5, while EA already showed a reduction, PAE saw a 94% increase, with a slower decay for this group at T15. The SF presented the negative $\Delta\%$ was almost double in PAE, with a quick recovery already at T5 for EA and levels still negative at all times for PAE. For SIgA-s, there was an increase of 37% in EA and only 7% at the TE in PAE; 41% in EA and 15% in PAE for T5; and 26% in EA and 14% in PAE at T15. SA showed a higher peak in EA (TE) and less acute in PAE (T5) but there was a decrease among both at T15. STP increased by 126% in EA and 438% in PAE, already showing a return at T5 for EA, but increasing by 213% in PAE. Negative levels were reached at T15 for EA but levels remained high in PAE. Levels of physical conditioning affect cardiovascular parameters, salivary biomarkers, and their correlation within the over-60s.

Key words: Aging. Health. Data mining. Sportomics.

EA – Elderly Athletes; PAE - Physically Active Elderly; BL - Blood Lactate; STP – Salivary Total Protein; SA - Salivary Amylase; IgA-s – Salivary Immunoglobulin A; SF - Salivary Flow; SNO - Salivary Nitric Oxide; DP - Double Product; SBP – Systolic Blood Pressure; DBP - Diastolic Blood Pressure; HR – Heart Rate.

46 **1. Introduction**

47 It is widely discussed by the scientific community that sarcopenia generated by physical
48 inactivity added to mental depression over the years can affect the quality of life of the
49 elderly.¹

50 This makes clear the need for regular exercise, minimizing the loss of bone and muscle
51 mass, providing better balance, and reducing the chances of falls, representing a
52 significant problem at this age.²

53 Therefore, using a sport method that reproduces natural conditions of the sport practiced
54 in a controlled and intentional way aims to understand the immunometabolic differences
55 generated in different populations. In the case of older people who regularly exercise
56 but with different levels of physical conditioning, it is essential. for the knowledge of
57 geriatricians and gerontologists.³⁻⁵

58 The main objective of this work was to evaluate and compare the physical conditioning,
59 hemodynamic, blood, and salivary biomarkers between elderly considered active and
60 athletes before, at the end, and during the recovery period of an exercise test on a
61 treadmill.

62 **2. Method**

63 *2.1 Participants*

64 Thirty men over the age of 60, each of whom were assiduous competitors and regular
65 exercisers with a minimum experience of 20 years in street racing events, were selected.
66 They were first examined by a general dentist who used the Plaque Control Index (PCI),
67 Bleeding Index (SI), and Simplified Oral Hygiene Index (OHIS) and checked for
68 xerostomia symptoms. The participants in good oral health moved on to the next phase
69 of the experiment. Those who were not were treated by the same dentist but did not take
70 any further part in the study.

71 The selected individuals underwent a careful anamnesis of general health conditions,
72 training routine, and competitions. The final sample of elderly people comprised 14
73 men divided into two groups: elderly athletes (EA: n = 8) and the physically active
74 elderly (PAE: n = 6).

The subjects' characteristics were as follows: **EA** 63.1 ± 6.2 years, weight 68.4 ± 8.1 kg, height 170.1 ± 1.9 cm, body mass index 20.5 ± 1.0 kg/m², resting heart rate 67.7 ± 7.0 bpm, maximal heart rate 154.7 ± 5.2 bpm, VO₂ max 32.8 ± 5.5 ml/kg/min). **PAE** 63.8 ± 4.0 years, weight 70.4 ± 3.1 kg, height 168.5 ± 2.8 cm, body mass index 23.5 ± 1.5 kg/m², resting heart rate 76.7 ± 1.5 bpm, maximal heart rate 147.0 ± 10.2 bpm, VO₂ max 24.4 ± 7.0 ml/kg/min).

2.2 Experimental Design

In the 48 hours before the exercise test, the participants did not perform any physical exercise. They ate their last meal at least two hours before the test. They were instructed to thoroughly clean their mouths and drink water ad libitum the day before the test. No licit or illicit stimulant substances (i.e., coffee, guarana, teas, and other xanthines, thermogenic, or central nervous system stimulants) or anything containing dyes could be ingested during this period.

For adaptation purposes, all participants had to perform two training sessions on different days. The stress test was carried out on a treadmill (Ecafix®, Brazil). The original Bruce protocol was used. Heart rate was monitored throughout the exercise test by 12-lead electrocardiogram using Ergo PC 13 for Windows. Systolic blood pressure (SBP) and diastolic blood pressure (DBP) were measured using the auscultatory method, with a sphygmomanometer and stethoscope (Tycos®, USA): $([2 \times \text{diastolic pressure}] + \text{systolic pressure})/3$.

2.3 Procedures and Measurements

Salivation was stimulated by chewing gum (Cadbury Adams Brasil Ind®) weighing 1.5g. Chewing took place naturally and personally without instructions on speed, strength, and frequency, though the participants were instructed to chew the gum on only one side of the mouth. On the other side, the Tubo Salivette® device was used for quick and safe saliva collection. The participants were given a chewing gum tablet to carry during street training sessions.

Saliva was placed in pre-cooled (4°C) mini-tubes. All samples were processed, and immediately after collection, centrifuged at 12,000 g; the pellet was discarded and the supernatant was frozen at 20°C until the day of the analysis.

105 The stress test was performed on a treadmill (Ecafix®, Brazil). The original Bruce
106 protocol was used.⁶ Heart rate was monitored throughout the exercise test by 12-lead
107 electrocardiogram using Ergo PC 13 for Windows. Systolic blood pressure (SBP) and
108 diastolic blood pressure (DBP) were measured using the auscultatory method, with a
109 sphygmomanometer and stethoscope (Tycos®, USA): $[(2 \times \text{diastolic pressure}] +$
110 $\text{systolic pressure})/3$. All the above-mentioned physiological parameters were evaluated
111 at rest (T0), during the exercise test, at the maximum peak of the exercise (TE), and on
112 the stands five and 15 minutes after the end of the test (T5 and T15). All stress tests
113 were performed in a private cardiology clinic in the presence of a physician. All
114 participants had to perform two training sessions on different days to adapt. The
115 exercise test was interrupted when any of these criteria were identified: elevation of
116 diastolic blood pressure (DBP) > 120mm/Hg in normotensive individuals and >
117 140mm/Hg in primary hypertensive individuals; elevation of systolic blood pressure
118 (SBP) > 260mm/Hg; a fall sustained SBP; clinical manifestations of typical severe chest
119 pain; ST-segment depression > 3mm; ST-segment elevation > 2 mm in the lead without
120 a q wave; complex ventricular arrhythmia; sustained supraventricular tachycardia onset;
121 atrial tachycardia; atrial fibrillation; atrial block (second and third degree ventricular);
122 signs of left ventricular failure; and the failure of monitoring and recording systems⁷.

123 Total salivary protein was measured by the biuret method using a standard laboratory
124 kit (UCFS DIASYS® cat. n° 1 0210 99 10 021 Germany) through two
125 spectrophotometric readings, one primary at 604 nm and the other at 700 nm
126 (Autoanalyser Architeet c8000, Abbot®, IL, USA).

127 The analysis of alpha-amylase activity was performed by a kinetic assay using the
128 CNPG kit (Pro Biotec, Ind. Com. Diagnóstico para Saúde, Uberlândia, MG, Brazil).
129 Assays were performed at room temperature (28°C) on microplates with ten microL of
130 saliva diluted 100 times in saline and 300 microL of CNPG buffer. The microplates
131 were read at 405 nm on a microplate reader (Amershan Biosciences, GE, Upsala,
132 Sweden) programmed to give the results unit of salivary amylase activity/mL of saliva
133 (U/mL). Total salivary protein was measured by the biuret method according to a
134 standard laboratory kit (UCFS DIASYS® cat. n° 1 0210 99 10 021 Germany) through
135 two spectrophotometric readings, one primary at 604nm and the other at 700nm
136 (Autoanalyser Architeet c8000, Abbot®, IL, USA).

137 The analysis of alpha-amylase activity was performed by a kinetic assay using the
138 CNPG kit (Pro Biotec, Ind. Com. Diagnóstico para Saúde, Uberlândia, MG, Brazil).
139 Assays were performed at room temperature (28°C) in microplates with ten microL of
140 saliva diluted 100 times in saline and 300 microL of CNPG buffer. The microplates
141 were read at 405nm on a microplate reader (Amershan Biosciences, GE, Upsala.
142 Sweden) programmed to give the results unit of salivary amylase activity/mL of saliva
143 (U/mL).

144 Total salivary protein was measured by the biuret method according to a standard
145 laboratory kit (UCFS DIASYS® cat. n° 1 0210 99 10 021 Germany) through two
146 spectrophotometric readings, one primary at 604nm and the other at 700nm
147 (Autoanalyser Architect c8000, Abbot®, IL, USA).

148 The analysis of alpha-amylase activity was performed by a kinetic assay using the
149 CNPG kit (Pro Biotec, Ind. Com. Diagnóstico para Saúde, Uberlândia, MG, Brazil).
150 Assays were performed at room temperature (28°C) in microplates with ten microL of
151 saliva diluted 100 times in saline and 300 microL of CNPG buffer. The microplates
152 were read at 405nm on a microplate reader (Amershan Biosciences, GE, Upsala.
153 Sweden) programmed to give the results unit of salivary amylase activity/mL of saliva
154 (U/mL). of a 250 µM NaNO₃ solution to verify the linearity of the reaction and
155 calculate a conversion factor of absorbance values into concentration (line equation).
156 The absorbance of the tests was quantified at 548 nm in a microplate spectrophotometer
157 (Synergy Microplate Reader) with analysis by the BioTek Gen5 Data Analysis Software
158 (BioTek instruments, Winooski, Vermont, USA).

159 The analysis of total IgA used the ELISA test according to laboratory routine. First,
160 polystyrene plates (Maxi-Sorp, Nunc, Wohlen) were sensitized with anti-human IgA
161 antibody (Sigma Chemical, Buchs), diluted at an optimal concentration in carbonate
162 buffer, 0.06M (pH 9.6) for 12 hours at 4°C. Then, the plates were washed and blocked
163 with a specific buffer of sodium dihydrochloride.

164 Orthophenylenediamine (OPD). Saliva samples were diluted from 1:2 in 1% BSA-PBS-
165 T and incubated for 1 hour at room temperature. After washing, biotinylated anti-IgA
166 conjugate labeled with peroxidase diluted in the concentration to be used was added.
167 The enzyme-substrate H₂O₂ + OPD (chromogen buffer) was incubated at room
168 temperature for 1 hour. The results were expressed in ELISA indices (IE) for individual

plate analysis. Optical density (OD) values were determined in an ELISA reader (Titertek Multiskan Plus, Flow Laboratories, USA) at 405 nm. After obtaining the results of total IgA, it was divided by the value of total protein in saliva, and the value of specific IgA was found.

Blood collection was performed in the left ear lobe of each volunteer. The first drop of blood was discarded to avoid contamination, with lactate eliminated in the sweat produced by the sweat glands. Then 25 microL of blood was collected in heparinized and calibrated capillaries. Blood lactate was analyzed by the electroenzymatic method.

2.4 Statistical Analysis

Initially, descriptive statistics were performed on the data, with measurements of position (mean, median, mode, and percentiles) and dispersion (amplitude, variance, standard deviation, and standard error).

Afterward, the univariate analysis of these data was performed using the Shapiro-Wilk normality test (because the sample was smaller than 30 individuals). The equal variance test would be applied if the Shapiro-Wilk test presented a result indicating normal distribution ($P > 0.05$). For results with $P > 0.05$, the paired T-Student test would follow; if $P \leq 0.05$, the paired T-Student test would follow the non-parametric Mann-Witney test. If the Shapiro-Wilk test presented a result indicating non-normal distribution ($P \leq 0.05$), the non-parametric Mann-Witney test would be applied directly.

Still, in the phase of the univariate analysis, the analysis of repeated measures ANOVA One Way dependent was performed because they were the same individuals in different conditions and moments.

So, for a better interpretation of the data, the individuals were divided into two groups according to their sex. Then, the calculation of percentage variation was applied:

$$\Delta\% = \frac{(Final\ value - Initial\ value)}{Initial\ value} \times 100$$

Cohen's equations⁸ were used to calculate the effect size for all variables to obtain Cohen d and r values:

7

$$d = \frac{M1-M2}{\sqrt{\frac{SD1^2+SD2^2}{2}}}$$

$$r = \frac{d}{\sqrt{(D^2)+4}}$$

195

196 Where, M represent the means of observations and SD their respective standard
197 deviations.

198 **Table 1.** Values of effect size.

Effect size	Small	Medium	Large
Cohen r	0.10	0.30	0.50
Cohen d	0.20	0.50	0.80

199 Source: ⁸

200

201 Next, multivariate data analysis was performed using data mining and machine learning
202 techniques.

203 In this phase, in order to seek a bivariate measure between the data, because the
204 observations contain quantitative values, the Pearson and Spearman correlation tests
205 were applied, with the Spearman correlation being used for a visual analysis using the
206 heat map strategy and the Pearson test as an initial measure for the following machine
207 learning analyses.

208 As exploratory models of machine learning: CLUSTER - Classical Clustering
209 (Agglomerative Hierarchical Method) and Nearest neighbor (single linkage);
210 ORDINATION – Principal component Analysis (PCA) and Correspondence Analysis
211 (CA).

212 The Z score was previously applied because the observations contained non similar
213 measurement units.


214 SigmaPlot 14.5 (Academic Perpetual License - Single User – ESD Systat® USA) and,
215 Past 4.03 (Free version for Windows) were used to carry out the different statistical tests
216 and produce the graphs. Finally, the correlation coefficients were presented using heat
217 maps ⁵.

218 **3. Results**

219 A holistic and integrated analysis was carried out to avoid traditional dogmas and
220 paradoxes.⁵ A Spearman rank correlation coefficient strategy was adopted so that the
221 findings could be plotted on heat maps for better visualization.

222

223



224 **Figure 1.** Heat map with holistic analysis (A and C – EA / B and D – PAE).

225 A negative correlation was found between salivary flow and cardiovascular parameters,
226 blood lactate, and total salivary proteins in both groups, but with correlation coefficients
227 of almost double for the elderly only active the elderly athletes, as the heat maps show
228 (Figs. 1 and 3).




Figure 2. The double product of the elderly athletes is more closely correlated with systolic blood pressure, while the double product is more closely correlated with heart rate.


Also, to facilitate the visualization of the behavioral data in both groups, Table 2 presents the values of the size of the Cohen effect and the percentage variations concerning T0. As was expected, heart rate and systolic blood pressure recovered faster among EA. The opposite was the case for diastolic blood pressure and the double product.

Table 2. The Cohen's effect size (r) for significant salivary, blood, and cardiovascular biomarkers between groups and (the percentage variation) of T0.

Parameters	Groups	TE	T5	T15
Heart Rate	EA	0.93 (125%)	0.64 (27%)	0.51 (-5%)
	PAE	0.99 (150%)	0.79 (34%)	0.79 (31%)
Systolic Blood Pressure	EA	0.83 (53%)	0.17 (5%)	0.04 (1%)
	PAE	0.76 (54%)	0.17 (-3%)	0.68 (-15%)
Diastolic Blood Pressure	EA	0.65 (26%)	0.06 (2%)	0.06 (-2%)
	PAE	0.69 (18%)	0.13 (-2%)	0.52 (-9%)
Double Product	EA	0.91 (242%)	0.57 (35%)	0.40 (21%)
	PAE	0.96 (297%)	0.60 (27%)	0.22 (8%)
Salivary Nitric Oxide	EA	0.16 (22%)	0.08 (8%)	0.03 (-3%)
	PAE	0.17 (34%)	0.28 (94%)	0.11 (27%)
Salivary Flow	EA	0.62 (-28%)	0.14 (5%)	0.40 (14%)
	PAE	0.97 (-54%)	0.93 (-38%)	0.70 (-17%)
Salivary Immunoglobulin A	EA	0.63 (37%)	0.64 (41%)	0.45 (26%)
	PAE	0.24 (7%)	0.52 (15%)	0.51 (14%)
Salivary Amylase	EA	0.48 (130%)	0.17 (26%)	0.02 (3%)
	PAE	0.28 (68%)	0.39 (100%)	0.01 (2%)
Salivary Total Protein	EA	0.49 (126%)	0.02 (-1%)	0.21 (-16%)
	PAE	0.78 (438%)	0.59 (213%)	0.37 (47%)

Blood Lactate	EA	0.84 (481%)	0.64 (398%)	0.58 (353%)
	PAE	0.91 (639%)	0.80 (495%)	0.55 (196%)

242 Regarding the latter double product, it was possible to observe an increase in both
 243 groups, but EA generated an intriguing finding: the increase was more closely
 244 correlated with the SBP (CC 0.854; $p = 0.0000002$), while for the PAE, the highest
 245 correlation was with HR (CC 0.872; $p = 0.0000002$; Figure 2).



246 **Figure 3.** The salivary flow of both groups is negatively
 247 correlated with the double product. This correlation is
 248 almost double in the elderly athletes; in the first group it is
 249 caused by SBP and in the second by HR.
 250
 251

252 As was expected, blood lactate showed an acute increase in time to exhaustion, reaching
 253 an increase of 481% in EA and 639% in PAE. However, what was not expected was the
 254 faster recovery of this metabolite in the PAE group, with a similar correlation between
 255 groups for DP and BL (Figure 4).





Figure 4. Blood lactate is positively correlated with double product in both groups.

The behavior of salivary biomarkers is shown in Figure 5 and Table 2 and their correlations with cardiovascular parameters in Figure 6, with very different designs.






Figure 5. The behavior of salivary biomarkers (A – SNO; B – IgA-s; C – SA; D – STP).

Dendrogram with Euclidian distance of Salivary Biomarkers with nearest neighbor clustering plot: single linkage (Fig 7A) confirmed that the double product of the physically active group had a behavior mainly altered by heart rate, while in the group of athletes by systolic blood pressure. Regarding salivary proteins, the active group presented behavior more similar to diastolic blood pressure, while the group of athletes in the group of athletes was more similar to salivary amylase. The similarity between the lactate of both groups made it clear that the exercise intensity was high and similar, and network plot with Fruchterman-Reingold algorithm (Fig 7B) revealed that the salivary flow of only active individuals was highly dissimilar to the other biomarkers and that this flow negatively correlated with systolic blood pressure in the athletes. In contrast, in the physically active group, the negative correlation was greater with heart rate.





Figure 6. Correlation between salivary biomarkers and cardiovascular parameters.

Salivary nitric oxide showed a similar increase in both groups at the time of exhaustion, but five minutes later (T5); while EA showed a reduction, PAE had a 94% increase, with a slower decay for the latter at T15.

Salivary flow was another biomarker where behaviour differed between the groups. The negative percentage variation was almost double in PAE. There was a rapid recovery at T5 for EA, and levels were negative at all times for PAE.



288 **Figure 7.** Dendrogram with Euclidian distance of Salivary Biomarkers. Nearest
289 neighbor clustering plot: single linkage (Fig 7A), network plot with Fruchterman-
290 Reingold algorithm (Fig 7B).

292 Salivary immunoglobulin A showed an increase of 37% in EA and only 7% in PAE
293 (TE). They reached 41% in EA and 15% in PAE for T5 and 26% in EA and 14% in
294 PAE for T15.

295 Salivary amylase showed a higher peak in the EA group (TE). It was less acute in PAE
296 (T5) but decreased to baseline values for both groups at T15.

297 Finally, the concentration of total salivary protein increased by 126% in EA and 438%
298 in PAE, already showing a return to baseline levels at T5 for EA, but remaining with an
299 increase of 213% in PAE, reaching negative levels in the T15 for EA and levels still
300 high in PAE.

301 **4. Discussion**

302 Saliva plays a fundamental role in immunometabolism, food digestion, immune
303 response, drug absorption, and taste. With age, salivary flow can decrease, resulting in
304 malnutrition, recurrent respiratory and digestive tract infections, and a diminution in
305 taste, with implications for the quality of life.⁹

306 Given that the continuous use of different pharmacological agents in this age group is
307 common, for example in the form of polypharmacy (where three or more medications
308 are taken), hyposalivation (a side-effect of many drugs) can be an issue.¹⁰

309 Age, nutritional status, hygiene, medication use, and previous illnesses can all affect
310 salivary flow, but the present study has shown that the level of physical training and
311 regular exercise can be added to the list. The decrease in salivary flow was almost
312 double in PAE relative to AE; recovery was rapid in the second group but remained
313 below baseline levels in the first. These findings corroborate those of a previous study
314 on rats, where regular exercise suppressed all age-induced changes in salivary flow.¹¹
315 We have also indicated that not only regular practice but the level of training also has an
316 influence.

317 To confirm and give robustness to the findings, correlating salivary flow with physical
318 conditioning, we indicate that cardiovascular parameters were inversely correlated with
319 salivary flow; both groups showed a strong correlation with the double product but with

320 a coefficient of almost double in PAE. It was also possible to observe that the change in
321 this parameter was caused by systolic blood pressure in the AS group. In PAE, heart
322 rate was the most closely correlated, with almost twice the correlation coefficient.

323 We also show that not only salivary flow presents itself differently according to the
324 level of physical conditioning. The concentration of salivary nitric oxide, an essential
325 mediator of intra- and extracellular processes mediating immunometabolism, and which
326 even influences body composition,¹² differed between the groups. An elevation of this
327 biomarker in EA of 22% was observed at the time of exhaustion, with a reduction at
328 later times. In PAE, the elevation in the TE was 34%. It continued to increase until it
329 reached 94% of elevation at T5, followed by a reduction to the point where it was 27%
330 at T15 relative to baseline values.

331 Salivary immunoglobulin A, an essential agent in the immunity of the mouth and
332 respiratory and digestive tracts, and which is responsible for oral microbiome
333 homeostasis,¹³ showed an increase of 37% in the TE for EA and only 7% in PAE. At
334 T5, it continued upward in EA, reaching 41% at baseline levels, while PAE saw a 15%
335 increase. Both saw a reduction in the final study time.

336 The concentration and activity of salivary amylase, an enzyme involved in the initiation
337 of polysaccharide digestion and the modulation of immunity, can be altered by mental
338 health,¹⁴ body composition, diet,¹⁵ and age.¹⁶ The present study has shown that it can
339 also be affected by physical conditioning, even in elderly individuals: EA showed an
340 increase of 130% at the time of exhaustion, with a subsequent reduction thereafter. By
341 contrast, PAE showed a 68% increase in the time to exhaustion with a later peak (T5),
342 reaching an elevation of 100%, and a subsequent sudden reduction. The results suggest
343 that EA were more immunometabolically efficient.

344 Total salivary proteins, which have marked antifungal, antiviral, and antibacterial, pH
345 homeostasis, digestive, mineralization, and tissue protection functions, are strongly
346 influenced by age, dietary habits, and environmental factors.¹⁷ The present study has
347 shown that they are also influenced by the level of training and physical conditioning.
348 While EA showed a 126% increase in TE with a reduction to baseline values five
349 minutes later, PAE showed an increase of 438% at the time of exhaustion, remaining
350 high at T5 (213%) and T15 (47%).

351 Finally, antagonistic behaviors were observed between the groups regarding SNO with
352 SBP and DBP, SA with DBP, and synergistically, but with notably different intensities
353 for IgA-s with HR and DP, SA with SBP, and STP with HR. The impact of training
354 levels for the age group under study has been revealed herein for the first time.

355 The unsupervised machine learning model was presented as a crucial exploratory tool to
356 search for correlations between variables and the formation of homogeneous groups
357 among themselves and heterogeneous groups. The present study used the Euclidean
358 measure of dissimilarity to assess the distance between variables and multivariate
359 cluster analysis with Classical Clustering (Agglomerative Hierarchical Method) and
360 Nearest neighbor (single linkage), and sorting by Principal Component Analysis (PCA)
361 and Correspondence Analysis (CA).

362 **5. Conclusions**

363 The level of physical conditioning in the 60+ age group has been shown to impact
364 cardiovascular parameters, salivary biomarkers, and their correlation. Even when the two
365 sub-groups in question regularly practiced the same type of exercise and competed with
366 the same regularity, their level of training affected the behavior of the study variables.

367 Therefore, geriatricians, gerontologists, physical trainers, nutritionists, and other
368 professionals who work with this age group must pay attention not only to relevant
369 reference values but also training levels. They should also have respect for the longevity
370 and quality of life of individuals at such a noble age.

371 *Data Availability*

372 All data generated or analyzed during the study are included in the published article.
373 Raw data can be requested from the corresponding author.

374 *Ethical Approval*

375 The present study was approved by the Human Research Ethics Committee of the
376 Federal University of Mato Grosso, opinion number 5.716.414.

377 *Consent*

378 After an initial lecture on the procedures, objectives, risks, and benefits given by those
379 involved in the study, the participants completed the Free and Informed Consent Form

380 (ICF). They were told that they could withdraw at any time and that they would be
381 withdrawn if they did not follow the protocols.

382 *Conflicts of Interests*

383 The authors declare no conflicts of interest.

384 *Practical applications*

385 Gerontologists, geriatricians, nutritionists, physical education professionals, public
386 managers, and other professionals who work with older people must take into account
387 the age-related traditional reference values for biomarkers and cardiovascular
388 parameters and recognize that the level of physical training in this group affects the
389 data. Studies such as the present one can be used as a parameter for different decision-
390 making.

391 *Author's Contributions*

392 ABJ, FSE and AMMN conceived and designed the study. PJVC, ABJ, LCOG, RJSB,
393 MJSB, FSE and AMMN collected the data. PJVC, ABJ, LCOG, RJSB, MJSB, FSE and
394 AMMN analyzed the data and wrote the manuscript. All authors read and provided
395 critical feedback on the manuscript before approval.

396 *Acknowledgements*

397 We thank the participants who agreed to participate in the study and the health
398 professionals who directly or indirectly helped to make it all happen.

399 **References**

- 400 1. Yuenyongchaiwat K, Boonsinsukh R. Sarcopenia and Its Relationships with
401 Depression, Cognition, and Physical Activity in Thai Community-Dwelling
402 Older Adults. *Current Gerontology and Geriatrics Research*. 2020;
403 2020(8041489):1-6. <https://doi.org/10.1155/2020/8041489>
- 404 2. Vale FA, Voos MC, Brumini C, Suda EY, Silva RL, Caromano FA. Balance as
405 an Additional Effect of Strength and Flexibility Aquatic Training in Sedentary
406 Lifestyle Elderly Women. *Current Gerontology and Geriatrics Research*. 2020;
407 2020(1895473):1-6. <https://doi.org/10.1155/2020/1895473>
- 408 3. Gonçalves LCO, Bessa A, Freitas-Dias R, Luzes R, Werneck-de-Castro JP,
409 Bassini A, Cameron LC. A sportomics strategy to analyze the ability of arginine

- 410 to modulate both ammonia and lymphocyte levels in blood after high-intensity
411 exercise. *Journal of The International Society of Sports Nutrition*. 2012; 9(1):30.
412 <https://doi.org/10.1186/1550-2783-9-30>
- 413 4. Lopes JSS, Neto AMM, Gonçalves LCO, Alves PRL, Almeida AC, Andrade
414 CMB. Kinetics of Muscle Damage Biomarkers at Moments Subsequent to a
415 Fight in Brazilian Jiu-Jitsu Practice by Disabled Athletes. *Frontiers in*
416 *Physiology*. 2019; 10(1055):1-9. <https://doi.org/10.3389/fphys.2019.01055>
- 417 5. Gonçalves LCO, Magalhães-Neto AM, Bassini A, Prado ES, Muniz-Santos R,
418 Verli MVA, Jurisica L, Lopes, JSS, Jurisica I, Andrade, CMB, Cameron LC.
419 Sportomics suggests that albuminuria is a sensitive biomarker of hydration in
420 cross combat. *Scientific Reports*. 2022; 12(1):8159.
421 <https://doi.org/10.1038/s41598-022-12079-7>
- 422 6. Bruce R. Exercise testing of patients with coronary artery disease. *Ann Clin Res*.
423 1971; 3:323-332.
- 424 7. Kawamura T. Avaliação da capacidade física e teste ergométrico. *Rev. Soc.*
425 *Cardiol. Estado de São Paulo*. 2001; 659-672.
- 426 8. Cohen J. Quantitative methods in psychology. A power primer. *Psychological*
427 *Bulletin*. 1992; 112(1):155-159. <https://doi.org/10.1037//0033-2909.112.1.155>
- 428 9. Xu F, Laguna L, Sarkar A. Aging-related changes in quantity and quality of
429 saliva: Where do we stand in our understanding? *Journal of Texture Studies*.
430 2019; 50(1):27-35. <https://doi.org/10.1111/jtxs.12356>
- 431 10. Scelza MFZ, Silva DF, Ahiadzro NK, Silva LE, Scelza P. The influence of
432 medication on salivary flow of the elderly: preliminary study. *Gerodontology*.
433 2010; 27(4):278-282. <https://doi.org/10.1111/j.1741-2358.2009.00326.x>
- 434 11. Jung WK, Park SB, Kim HR, Ryu HY, Kim YH, Kim J. Advanced Glycation
435 End Products Increase Salivary Gland Hypofunction in d-Galactose-Induced
436 Aging Rats and Its Prevention by Physical Exercise. *Current Issues in*
437 *Molecular Biology*. 2021; 43(3):2059-2067.
438 <https://doi.org/10.3390/cimb43030142>
- 439 12. Vitale E, Jirilo E, Magrone T. Correlations between the Youth Healthy Eating
440 Index, body mass index and the salivary nitric oxide concentration in
441 overweight/obese children. *Endocrine, Metabolic & Immune Disorders Drug*
442 *Target*. 2014; 14(2):93-101.
443 <https://doi.org/10.2174/1871530314666140307095630>

- 444 13. Pedersen AML, Belstrom D. The role of natural salivary defences in maintaining
445 a healthy oral microbiota. *Journal of Dentistry*. 2019; 80(1):S3-S12.
446 <https://doi.org/10.1016/j.jdent.2018.08.010>
- 447 14. Jezova D, Trebaticka J, Buzgoova K, Durackova Z, Hlavacova N. Lower
448 activity of salivary alpha-amylase in youths with depression. *Stress*. 2020;
449 23(6):688-693. <https://doi.org/10.1080/10253890.2020.1777975>
- 450 15. Polito R, Valenzano A, Scarinci A, Vilano I, Monda M, Messina A, Cibelli G,
451 Porro C, La Torre E, Pisanelli D, Moscatelli F, Messina G, Monda V. Very
452 Low-Calorie Ketogenic Diet Modulates the Autonomic Nervous System
453 Activity through Salivary Amylase in Obese Population Subjects. *International*
454 *Journal of environmental Research and Public health*. 2021; 18(16):8475.
455 <https://doi.org/10.3390/ijerph18168475>
- 456 16. Wang CH, Woolfolk CA. Salivary amylase activity of the aged. *Gerontology*.
457 1990; 36(4):193-198. <https://doi.org/10.1159/000213199>
- 458 17. Bhuptani D, Kumar S, Vats M, Sagav R. Age and gender related changes of
459 salivary total protein levels for forensic application. *The Journal of Forensic*
460 *Odonto-Stomatology*. 2018; 36(1):26-33.