

Gut microbiota and fecal short chain fatty acids differ with adiposity and country of origin: The METS-Microbiome Study

Gertrude Ecklu-Mensah^{1*}, Candice Choo-Kang^{2*}, Maria Gjerstad Maseng^{3,4,5}, Sonya Donato¹, Pascal Bovet⁶, Kweku Bedu-Addo⁷, Jacob Plange-Rhule⁷, Terrence E. Forrester⁸, Estelle V. Lambert⁹, Dale Rae⁹, Amy Luke², Brian T. Layden^{10,11}, Stephen O'Keefe¹², Jack A. Gilbert^{1**}, Lara R. Dugas^{2, 13**}

*Authors contributed equally.

¹Department of Pediatrics, University of California San Diego, La Jolla, CA, USA

² Public Health Sciences, Parkinson School of Health Sciences and Public Health, Loyola University Chicago, Maywood, IL, USA

³ Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway.

⁴ Dep. of Gastroenterology, Oslo University Hospital, Oslo, Norway.

⁵ Bio-Me, Oslo, Norway.

⁶ University Center for Primary Care and Public Health (Unisanté), Lausanne, Switzerland & Ministry of Health, Republic of Seychelles Department of Physiology, SMS,

⁷ Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

⁸ Solutions for Developing Countries, University of the West Indies, Mona, Kingston, Jamaica

⁹ Research Unit for Exercise Science and Sports Medicine, University of Cape Town, Cape Town, South Africa

¹⁰ Department of Medicine, University of Illinois at Chicago, Chicago, Illinois, USA

¹¹ Jesse Brown Veterans Affairs Medical Center, Chicago, Illinois, USA.

¹² Department of Medicine, University of Pittsburgh, PA, USA

¹³ Division of Epidemiology and Biostatistics, School of Public Health, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa.

Funding: This work is supported by the National Institutes of Health grant R01-DK111848

**Co-Corresponding authors: Jack A Gilbert PhD, Email: jagilbert@health.ucsd.edu; Lara Dugas, PhD, MPH, Email: ldugas@luc.edu

Abstract

The relationship between the gut microbiota, short chain fatty acid (SCFA) metabolism, and obesity remains unclear due to conflicting reports from studies with limited statistical power. Additionally, this association has rarely been explored in large scale diverse populations. Here, we investigated associations between fecal microbial composition, predicted metabolic potential, SCFA concentrations, and obesity in a large ($N = 1,934$) adult cohort of African-origin spanning the epidemiologic transition, from Ghana, South Africa, Jamaica, Seychelles, and the United States (US). The greatest gut microbiota diversity and total fecal SCFA concentration was found in the Ghanaian population, while the lowest levels were found in the US population, respectively representing the lowest and the highest end of the epidemiologic transition spectrum. Country-specific bacterial taxa and predicted-functional pathways were observed, including an increased prevalence of *Prevotella*, *Butyrivibrio*, *Weissella* and *Romboutsia* in Ghana and South Africa, while *Bacteroides* and *Parabacteroides* were enriched in Jamaican and the US populations. Importantly, 'VANISH' taxa, including *Butyricicoccus* and *Succinivibrio*, were significantly enriched in the Ghanaian cohort, reflecting the participants' traditional lifestyles. Obesity was significantly associated with lower SCFA concentrations, a decrease in microbial richness, and dissimilarities in community composition, and reduction in the proportion of SCFA synthesizing bacteria including *Oscillospira*, *Christensenella*, *Eubacterium*, *Alistipes*, *Clostridium* and *Odoribacter*. Further, the predicted proportions of genes in the lipopolysaccharide (LPS) synthesis pathway were enriched in obese individuals, while genes associated with butyrate synthesis via the dominant pyruvate pathway were significantly reduced in obese individuals. Using machine learning, we identified features predictive of metabolic state and country of origin. Country of origin could accurately be predicted by the fecal microbiota (AUC = 0.97), whereas obesity could not be predicted as accurately (AUC = 0.65). Participant sex (AUC = 0.75), diabetes status (AUC = 0.63), hypertensive status (AUC = 0.65), and glucose status (AUC = 0.66) could all be predicted with different success. Interestingly, within country, the predictive accuracy of the microbiota for obesity was inversely correlated to the epidemiological transition, being greatest in Ghana (AUC = 0.57). Collectively, our findings reveal profound variation in the gut microbiota, inferred functional pathways, and SCFA synthesis as a function of country of origin. While obesity could be predicted accurately from the microbiota, the variation in accuracy in parallel with the epidemiological transition suggests that differences in the microbiota between obesity and non-obesity may be larger in low-to-middle countries compared to high-income countries. Further examination of independent study populations using multi-omic approaches will be necessary to determine the factors that drive this association.

Introduction

Obesity, which affects more than 600 million adults worldwide (“Obesity and Overweight” n.d.), over a third of Americans (Hales et al. 2020), and accounts for over 60% of deaths related to high body mass index (BMI) (Tseng and Wu 2019), remains an ongoing global health epidemic that continues to worsen at an alarming rate. A major driver of obesity is the adoption of a western lifestyle, which is characterized by excessive consumption of ultra-processed foods. Obesity is a major risk factor for type 2 diabetes, and according to the most recent National Diabetes Statistics Report almost 13% of the adult US population now have diabetes. Not only do 49.6% of adult African Americans present with obesity but over 17% of them now have diabetes, and are 1.5 times as likely to present with type 2 diabetes compared to whites (“National Diabetes Statistics Report” 2022). Populations of African-origin outside of the US are experiencing similar fates, as the prevalence of obesity among adults living in Sub-Saharan Africa is greater than 13%, and higher than the global obesity prevalence for adults (Agyemang et al. 2016). This has been accompanied by dramatic increases in the prevalence of non-communicable diseases such as type two diabetes and hypertension among people of African-origin (Roth et al. 2020; Gouda et al. 2019). Therefore, disrupting the rapidly expanding obesity epidemic, particularly among African-origin populations is critical to controlling the cardiometabolic disorder epidemic (Geng et al. 2022). However, successfully managing and treating obesity and its comorbidities, and specifically maintaining weight loss long-term, is particularly challenging due to an incomplete understanding of the heterogeneous and complex etiopathology, as well as additional challenges facing populations experiencing rapid urbanization (Nordmo, Danielsen, and Nordmo 2020; Geng et al. 2022; Barone et al. 2022). The epidemiologic transition is a model able to capture these shifts in dietary and rural to urban movements and is characterized by diets that are high in ultra-processed foods with a significant loss in fiber, as evidenced in the US, where less than 50% of the population meet dietary fiber recommendations (Dahl and Stewart 2015).

Gut microbial ecology and metabolism play pivotal roles in the onset and progression of obesity and its related metabolic disorders (Ley 2010). Obese and lean individuals have reported differences in the composition and functional potential of the gut microbiome, with an overall reduction in species diversity in the obese gut (Dugas, Bernabé, et al. 2018; Greenblum, Turnbaugh, and Borenstein 2012; Jumpertz et al. 2011; Ley et al. 2006; Turnbaugh et al. 2009; Le Chatelier et al. 2013), additionally, fecal microbiota transfer from obese donors to mouse models can recapitulate the obese phenotype (Turnbaugh et al. 2006, 2008; Ridaura et al. 2013). Further, fecal microbiota transplant from healthy donors into patients with obese and metabolic syndrome has been shown to improve markers of metabolic health in the recipients (Vrieze et al. 2012). While these studies suggest that modification of microbial ecology may offer new options for the treatment and prevention of obesity, the mechanism that drives the microbiota-obesity relationship is not fully understood. The microbiota may facilitate greater energy exploitation from food, and storage capacity by the host (Turnbaugh et al. 2006; DiBaise et al. 2008), influencing adipose tissue composition and fat mass gain, as well as providing chronic low-grade inflammation and insulin resistance (Cani and Delzenne 2009; J. L. Sonnenburg and Bäckhed 2016).

Among the numerous microbial metabolites modulating obesity, there is an ever-growing interest in the role of short-chain fatty acids (SCFAs), which includes butyrate, acetate, and propionate as

potential biomarkers for metabolic health as well as therapeutic targets. SCFAs derive primarily from microbial fermentation of non-digestible dietary fiber in the colon. They have many effects on host metabolism including serving as an energy source for host colonocytes, used as precursors for the biosynthesis of cholesterol, lipids, proteins and regulating gut barrier activities (Dalile et al. 2019; Koh et al. 2016; van der Hee and Wells 2021). Human and animal studies demonstrate a protective role of SCFAs in obesity and metabolic disease. In experimental animal models, SCFA supplementation reduces body weight, improves insulin sensitivity, and reduces obesity-associated inflammation (Vinolo et al. 2011; Gao et al. 2009; Henagan et al. 2015; Lu et al. 2016; Bonomo et al. 2020). In humans, increased gut production of butyrate correlates with improved insulin response after an oral glucose-tolerance test (Sanna et al. 2019). Although increased SCFA levels are generally observed as positive for health (Valdes et al. 2018), other studies have suggested that overproduction may promote obesity, possibly resulting from greater energy accumulation (Schwiertz et al. 2010; Rahat-Rozenbloom et al. 2014; Teixeira et al. 2013). Indeed, a previous study observed greater fecal SCFA concentrations to be linked with obesity, increased gut permeability, metabolic dysregulation, and hypertension in a human cohort (de la Cuesta-Zuluaga, Mueller, et al. 2018).

The conflicting obesity role of SCFAs identified by existing studies may result from the variation in the gut microbiota, which is shaped by lifestyle and diet. Adequately powered studies in well-characterized populations may permit more rigorous assessments of individual differences. Prior comparative epidemiological studies have broadly focused on either contrasting the gut microbiota of extremely different populations, such as the traditional hunter-gatherers and urban-westernized countries, or ethnically homogenous populations (Pasolli et al. 2019; He et al. 2018; Peters et al. 2018; Zhernakova et al. 2016). Demographic factors represent one of the largest contributors to the individualized nature of the gut microbiome (Falony et al. 2016; Zhernakova et al. 2016; Yatsunen et al. 2012). Thus, it is unclear to what extent the associations between gut microbiome, SCFA and obesity generalize across different geographies and this, additionally limits our understanding and interpretation, especially when considering the substantial geographic disparities in obesity.

The five diverse, well-defined cohorts from the Modeling the Epidemiologic Transition Study (METS) offers a unique opportunity to examine the issues since they are more representative of most of the world's population. METS has longitudinally followed an international cohort of approximately 2,500 African-origin adults spanning the epidemiologic transition from Ghana, South Africa, Jamaica, Seychelles, and the US since 2010 to investigate differences in health outcomes utilizing the framework of the epidemiologic transition. Pioneering microbiome studies from the METS cohorts reveal that cardiometabolic risk factors including obesity is significantly associated with reduced microbial diversity, and the enrichment of specific taxa and predicted functional traits in a geographic-specific manner (Dugas, Bernabé, et al. 2018; Fei et al. 2019). While yielding valuable descriptions of the connections between the gut microbiota ecology and disease, particularly obesity, as well as pioneering the efforts of microbiome studies of populations of African-origin on different stages of the ongoing nutritional epidemiologic transitions, these studies, however, have applied small sample size (N=100 to N=655), and also did not utilize all the countries in the METS cohort. Thus, uncertainties remain as to the precise interpretation of

the microbiome-obesity associations, which hampers further progress towards diagnostic and clinical applications.

Our new study METS-Microbiome investigated associations between the gut microbiota composition and functional patterns, concentrations of fecal SCFAs and obesity in a large ($N = 1,934$) adult population cohort of African-origin, comprised of Ghana, South Africa, Jamaica, Seychelles, and the US spanning the epidemiologic transition (Dugas, Lie, et al. 2018; Luke et al. 2011). The central hypothesis is that shifts towards the highest end of the epidemiologic transition spectrum is associated with alterations in microbiota diversity and community composition, reductions in levels of fecal SCFAs and obesity.

Materials and Methods

Study Cohort. Since 2010, METS, and the currently funded METS-Microbiome study has longitudinally followed an international cohort of African-origin adults spanning the epidemiologic transition from Ghana, South Africa, Jamaica, Seychelles, and US (Dugas, Lie, et al. 2018; Luke et al. 2011). METS utilizes the framework of the epidemiologic transition to investigate differences in health outcomes based on country of origin. The epidemiologic transition is defined using the United Nations Human Development Index (HDI) as an approximation of the epidemiologic transition. Ghana represents a lower-middle income country, South Africa represents a middle-income country, Jamaica and Seychelles represent high income countries and the US represents a very high-income country. This framework has allowed us to investigate aspects of increased Westernization throughout the world (ex. increased consumption of ultra-processed foods) are related to increased prevalence of obesity, diabetes and cardiometabolic diseases. Our data from the original METS cohort demonstrate that the epidemiologic transition has altered habitual diets in the international METS sites, and that reduced fiber intake is associated with higher metabolic risk, inflammation, and obesity across the epidemiologic transition (Mehta et al. 2021). Originally, 2,506 African-origin adults (25–45 yrs), were enrolled in METS between January 2010 and December 2011 and followed on a yearly basis. In 2018, METS participants were recontacted and invited to participate in METS-Microbiome. Participants were excluded from participating in the original METS study if they self-reported an infectious disease, including HIV-positive individuals, pregnancy, breast-feeding or any condition which prevented the individual from participating in normal physical activities. METS-Microbiome was approved by the Institutional Review Board of Loyola University Chicago, IL, US; the Committee on Human Research Publication and Ethics of Kwame Nkrumah University of Science and Technology, Kumasi, Ghana; the Research Ethics Committee of the University of Cape Town, South Africa; the Board for Ethics and Clinical Research of the University of Lausanne, Switzerland; and the Ethics Committee of the University of the West Indies, Kingston, Jamaica. All study procedures were explained to participants in their native languages, and participants provided written informed consent after being given the opportunity to ask any questions.

Participant anthropometry, sociodemographic and biochemical measurements. Participants completed the research visits at the established METS research clinics located in the respective communities (Luke et al. 2011). Briefly, they presented themselves at the site-specific research clinic early in the morning, following an overnight fast. The weight of the participant was measured

without shoes and dressed in light clothing to the nearest 0.1 kg using a standard digital scale (Seca, SC, USA). Height was measured using a stadiometer without shoes and head held in the Frankfort plane to the nearest 0.1 cm. Waist circumference was measured to the nearest 0.1 cm at the umbilicus, while hip circumference was measured to the nearest 0.1 cm at the point of maximum extension of the buttocks. Adiposity (% body fat) was assessed using BIA (Quantum, RJL Systems, Clinton Township, MI), and study specific equations (Luke et al. 2011). Blood pressure was measured using the standard METS protocol using the Omron Automatic Digital Blood Pressure Monitor (model HEM-7471c, Omron Healthcare, Bannockburn, IL, USA), with the antecubital fossa at heart level. Participants were asked to provide a fecal sample using a standard collection kit (EasySampler stool collection kit, Alpcos, NH). Fecal samples were placed within a -80° freezer immediately upon receipt at all the sites. Participants were requested to fast from 8 pm in the evening prior to the clinic examination, during which fasting capillary glucose concentrations were determined using finger stick (Accu-check Aviva, Roche).

Fecal Short Chain Fatty Acid quantification. As in our previous studies (Nooromid et al. 2020; Lewandowski et al. 2021; Reiman, Layden, and Dai 2021; Barengolts et al. 2019; Navarro et al. 2018; Dugas, Bernabé, et al. 2018), fecal SCFAs were measured using LC-MC/MS at the University of Illinois-Chicago Mass Spectrometry Core using previously published methods (Moreau et al. 2003; Richardson et al. 1989). The LC-MC/MS analysis was completed on an AB Sciex Qtrap 5500 coupled to Agilent UPLC/HPLC system. All samples were analyzed by Agilent poroshell 120 EC-C18 Column, 100Å, 2.7 µm, 2.1 mm X 100 mm coupled to an Agilent UPLC system, which was operated at a flow rate of 400 µl/min. A gradient of buffer A (H₂O, 0.1% Formic acid) and buffer B (Acetonitrile, 0.1% Formic acid) were applied as: 0 min, 30% of buffer B; increase buffer B to 100% in 4 min; maintain B at 100% for 5 min. The column was then equilibrated for 3 min at 30% B between the injections with the MS detection is in negative mode. The MRM transitions of all targeted compounds include the precursor ions and the signature production ion. Unit resolution is used for both analyzers Q1 and Q3. The MS parameters such as declustering potential, collision energy and collision cell exit potential are optimized in order to achieve the optimal sensitivity. SCFAs are presented as individual SCFAs (µg/g), including: butyric acid, propionic acid, acetic acid and valeric acid, as well as total SCFAs (sum of 4).

METS data showed Ghanaians consumed the greatest amount of both soluble and insoluble fiber and had the lowest percentage energy from fat (42.5% of the Ghanaian cohort, dietary fiber intake: 24.9 g ± 9.7g/day). The US has the highest proportion of energy from fat and the lowest fiber intake of the five sites (3.2% of the US cohort, dietary fiber intake: 14.2 g ± 7.1 g/day).

DNA extraction, Amplicon Sequencing. Fecal samples were shipped on dry ice to the microbiome core sequencing facility, University of California, San Diego for 16S rRNA gene processing. Fecal samples were randomly sorted, transferred to 96-well extraction plates and DNA was extracted using MagAttract Power Microbiome kit. Blank controls and mock controls (ZymoBiomix) were included per extraction plate, which were carried through all downstream processing steps. Extracted DNA was used for amplification of the V4 region of the 16S rRNA gene with 515F-806R region-specific primers according to the Earth Microbiome Project (Thompson et al. 2017; Walters et al. 2016). Purified amplicon libraries were sequenced on the Illumina NovaSeq platform to produce 150 bp forward and reverse reads through the IGM Genomics Center, University of California San Diego. Full DNA extraction, amplification, quantification, and sequencing protocols

and standards are available at <http://www.earthmicrobiome.org/protocols-and-standards>; (Thompson et al. 2017).

Bioinformatic analysis. The generated raw sequence data were uploaded and processed in Qiita (Gonzalez et al. 2018) (Qiita ID 13512) an open-source, web-enabled microbiome analysis platform. Sequences were demultiplexed, quality filtered, trimmed, erroneous sequences were removed, and amplicon sequence variants (ASVs) were defined using Deblur (Amir et al. 2017). The deblur ASV table was exported to Qiime2 (Bolyen et al. 2019; Bokulich et al. 2018) and representative sequences of the ASVs were inserted into the Greengenes 13.8 99% identity tree with SATé-enabled phylogenetic placement (SEPP) using q2-fragment-insertion (Bolyen et al. 2019; Mirarab, Nguyen, and Warnow 2012) to generate an insertion tree for diversity computation. Additionally, the deblur ASV table was assigned taxonomic classification using the Qiime2 feature-classifier, with Naive Bayes classifiers trained on the SILVA database (version 138; (McLaren 2020)). A total of 463,258,036 reads, 154,952 ASVs and 1902 samples were obtained from the deblur table. The resulting ASV count table, taxonomy data, insertion tree, and sample metadata were exported and merged into a phyloseq (McMurdie and Holmes 2013) object in R (R Foundation for Statistical Computing, Vienna, Austria) for downstream analysis. Features with less than ten reads in the entire dataset and samples with fewer than 6,000 reads were removed from the phyloseq object. In addition, mitochondrial and chloroplast-derived sequences, non-bacterial sequences, as well as ASVs that were unassigned at phylum level were filtered prior to analyses. There were 433,364,873 reads and 13254 ASVs in the remaining 1873 fecal samples in the phyloseq object. The remaining samples after filtering were rarefied to a depth of 6,000 reads to avoid sequencing bias, before generating alpha diversity measures, leaving 9917 ASVs across 1873 samples.

Diversity and differential proportional analyses: Alpha diversity measures based on Observed Amplicon Sequence Variants (ASVs), Faith's Phylogenetic Diversity, and Shannon Index were conducted on rarified samples using phyloseq (McMurdie and Holmes 2013) and picante (Kembel et al. 2010) libraries. Beta diversity was determined using both weighted and unweighted UniFrac distance matrices (Lozupone and Knight 2005), generated in phyloseq. For differential abundance analysis, samples were processed to remove exceptionally rare taxa. First, the non-rarefied reads were filtered to remove samples with < 10,000 reads. Next, ASVs with fewer than 50 reads in total across all samples and/or were present in less than 2% of samples were excluded. This retained 2061 ASVs across 1694 samples. The retained ASVs were binned at genus level, and subsequently used in the analysis of compositions of microbiomes with bias correction (ANCOMBC; (H. Lin and Peddada 2020) to determine specific taxa differentially abundant across sites or obese phenotype. ANCOM-BC is a statistical approach that accounts for sampling fraction, normalizes the read counts by a process identical to log-ratio transformations while controlling for false discovery rates and increasing power. This method applies a library-specific offset term estimated from the observed abundance, which is incorporated into a linear regression model, providing the bias correction. Site, age, sex, BMI were added as covariates in the ANCOM-BC formula to reduce the effect of confounders. The *Bacteroides* *Prevotella* ratio was calculated by dividing the abundance of the genera *Bacteroides* by *Prevotella*. Participants were classified into *Bacteroides* enterotype (B-type) if the ratio was greater than 1, otherwise *Prevotella* enterotype (P-type).

Random forest classifier: Random Forest supervised learning models implemented in Qiime2 were used to estimate the predictive power of microbial community profiles for site and obese phenotype. The classifications were done with 500 trees based on 10-fold cross-validation using the QIIME “sample-classifier classify-samples” plugin (Bokulich et al. 2018). A randomly drawn 80% of samples were used for model training, whereas the remaining 20% were used for validation. Further, the 30 most important ASVs for differentiating between site or obese phenotype were predicted and annotated.

Predicted metabolic gene pathway analysis: The functional potential of microbial communities was inferred using the Phylogenetic Investigation of Communities by Reconstruction of Unobserved States 2 (PICRUSt2) v2.5.1 with the ASV table processed to remove exceptionally rare taxa and the representative sequences as input files (Douglas et al. 2020). The metabolic pathway from the PICRUSt2 pipeline was annotated using the MetaCyc database (Caspi et al. 2016). The predicted MetaCyc abundances (unstratified pathway abundances) were analyzed with ANCOM-BC to determine differentially abundant pathway associations across sites and obese status. Site, age, sex, BMI were added as covariates in the ANCOM-BC formula to reduce the effect of confounders.

Statistical Analysis: All statistical analyses and graphs were done with R software. Kruskal-Wallis test and Permutational Analysis of Variance (PERMANOVA) test with 999 permutations using the Adonis function in the vegan package (Oksanen et al. 2013) were performed to compare alpha and beta diversity measures respectively with multiple groups comparison correction. PERMANOVA models were adjusted for BMI, age, sex for country whereas age, sex and country were accounted for in obese groups. Variables that showed significant differences in the PERMANOVA analyses, PERMDISP test was performed to assess differences in dispersion or centroids. For differential abundance analysis, the false-discovery rate (FDR) method incorporated in the ANCOM-BC library was used to correct P values for multiple testing. A cut-off of $P_{adj} < 0.05$ was used to assess significance. Spearman correlations were performed between concentrations of short chain fatty acids, Shannon diversity or concentrations of short chain fatty acids and differentially abundant taxa that were identified either among study sites or in obese and non-obese individuals. The resulting p-values were adjusted for multiple testing using the false-discovery rate (FDR). P value < 0.05 was considered statistically significant. A mixed model was built using lme4 package to assess whether total SCFAs could be predicted by Shannon diversity, obesity, and country, setting obesity and Shannon diversity as fixed effects and random intercept by country.

Data availability: All 16S rRNA gene sequence data are publicly available via the QIITA platform (<https://qiita.ucsd.edu>) under the study identifier (ID=13512) and will soon be available on the European Bioinformatics Institute (EBI) site.

Results

Obesity differs significantly across the epidemiological transition. From 2018-2019, the METS-Microbiome study recruited 2,085 participants (~60% women) ages 35-55 years old from five different sites (Ghana, South Africa, Jamaica, Seychelles, and US). Of these participants, 1,249 have been followed on a yearly basis since 2010 under the parent METS study. Data from 1,867 participants with complete data sets were used in this analysis. Overall mean age was 42.5 ± 8.0 years (**Table 1**). Mean fasted blood glucose was 105.2 ± 39.4 mg/dL, mean systolic blood pressure was 123.4 ± 18.1 mm Hg and mean diastolic blood pressure was 77.2 ± 13.1 (**Table 1**). When compared to the high-income countries (Jamaica, Seychelles, and US), both women and men from the lower- and middle-income countries (Ghana and South Africa) had significantly lower BMI, fasted blood glucose and blood pressure (systolic and diastolic). Mean BMI was lowest in the South African men ($22.3 \text{ kg/m}^2 \pm 4.1$) and highest in US women ($36.3 \text{ kg/m}^2 \pm 8.8$). When compared to the US, all sites had significantly lower prevalence of obesity ($p < 0.001$ for all sites except for Seychelles: $p = 0.02$). Prevalence of hypertension was lowest in Ghanaian men (33.1%) and highest in US men (72.7%). Prevalence of diabetes was lowest in South African women and men (3.5% for women and men) and highest for Seychellois men (22.8%). When compared to the US, prevalence of hypertension and diabetes was significantly lower in countries at the lower end of the spectrum of HDI (i.e., Ghana and South Africa) when compared to the US ($p < 0.001$).

Microbial community composition and predicted metabolic potential differs significantly between countries and correlates with obesity. Following the removal of control samples and those that had fewer than 6,000 reads and features less than ten reads in the entire dataset, a total of 433,364,873 16S rRNA gene sequences were generated from the 1,873 fecal samples which were clustered into 13,254 ASVs. Country of origin describes most of the variation in microbial diversity and composition, with significant differences in both alpha and beta diversity. Although there were major variations in alpha diversity between countries and large degree of inter-individual variation within countries, Ghana showed significantly greater diversity for all the alpha diversity metrics (Observed ASVs, Shannon Diversity and Faith's phylogenetic diversity) when compared to all other countries. The Seychelles and US had the lowest alpha diversity (**Figure 1, Table 2**). The stool microbiota alpha diversity of non-obese individuals was significantly greater when compared with that of obese individuals (**Figure 1**). Beta diversity was also significantly different between countries (**Figure 1, Table 3 & Supplementary Table 2**; principal coordinate analysis, weighted UniFrac distance; F-statistic = 58.67; $p < 0.001$; unweighted UniFrac distance; $F = 39.87$; $p < 0.001$) and obese group (weighted UniFrac distance; F-statistic = 2.39; $p = 0.031$; unweighted UniFrac distance; $F = 6.06$; $p < 0.001$).

Next, we compared fecal microbiota diversity between obese individuals with their non-obese counterparts within each country independently (**Supplementary Table 1**). Greater alpha diversity was detected in non-obese subjects in the Ghanaian (Observed ASVs, Faith PD; $p < 0.05$) and South African cohorts (Observed ASVs; $p < 0.05$) only. Similarly, significant differences in beta diversity between obese and non-obese microbiota were observed in Ghana (Unweighted UniFrac; $p < 0.05$), South Africa (Unweighted UniFrac; $p < 0.05$) and US (Weighted UniFrac; $p < 0.05$) data sets (**Table 3 & Supplementary Table 2**). These results suggest that the beta diversity differences observed in the Ghanaian and South African participants may partly be due to the presence of more abundant fecal microbiota taxa in the fecal samples whereas among the

US participants, the differences may be related to the abundance of rare taxa. Collectively, these observations suggest that country is a major driver of the variance in gut microbiota diversity and composition among participants with or without obesity with marked contributions from Ghana and South Africa and modest contribution from the US in the overall cohort.

We also examined whether country of origin or obesity relates to the presence of specific microbial genera frequently used to stratify humans into enterotypes (Arumugam et al. 2011). As expected, large differences in enterotype between the countries were observed. The *Prevotella* enterotype (P-type) was enriched on the African continent, with 81% and 62% in Ghanaians and South Africans respectively while *Bacteroides* enterotype (B-type) was dominant in the US (75%), Jamaican cohorts (68%), and comparable proportions of both enterotypes among individuals from Seychelles. Further, obese individuals displayed a greater abundance of B-type whereas a higher proportion of the P-type associated with the non-obese group (**Supplementary Table 3**). Consistent with this observation, the abundance of B-type correlated with higher BMI ($p=0.004$) than P-type. Significantly greater diversity and increased levels of total SCFA were observed in participants in the P-type (**Supplementary Table 3**). The relative abundance of shared and unique features between the different countries illustrated by the Venn diagram showed that Ghana carries the largest proportion of unique taxa than the other countries, and US the lowest (**Figure 1**).

Microbial taxa differ significantly between countries and between lean and obese individuals. In comparison with the US, South African fecal microbiota had a significantly greater proportion of *Clostridium*, *Olsenella*, Bacilli and *Mogibacterium*; Jamaican samples had a significantly greater proportion of Bacilli, *Bacteroides*, Clostridia, *Dialister*, Enterobacteriaceae, and Oscillospiraceae; Seychelles samples had a significantly greater proportion of *Clostridium*, *Olsenella* and *Haemophilus*; and Ghanaian samples had a significantly greater proportion of *Clostridium*, *Prevotella*, *Weissella*, Enterobacteriaceae and Butyricocccaceae. The US samples had a significantly greater proportion of *Aldercruzia*, *Anaerostipes*, *Clostridium*, *Eggerthella*, *Eisenbergiella*, Ruminococcaceae and *Sellimonas* compared to the 4 countries (**Supplementary Figure 1**).

When adjusted for country, age, and sex ($p < 0.05$; false discovery rate (fdr)-corrected), 38 Amplicon Sequence Variants (ASVs) were significantly different between obese and non-obese groups. The obese group was characterized by an increased proportion of *Allisonella*, *Dialister*, *Oribacterium*, *Mitsuokella*, and *Lachnospira*, whereas non-obese microbiota had a significantly greater proportion of *Alistipes*, *Bacteroides*, *Clostridium*, *Parabacteroides*, *Christensenella*, *Oscillospira*, Ruminococcaceae (UBA1819), and Oscillospiraceae (UCG010) (**Supplementary Figure 1**).

Microbial taxonomic features predict obesity overall and within each country. Using supervised Random Forest machine learning, the predictive capacity of the gut microbiota features in stratifying individuals to country of origin, sex, or with metabolic phenotypes were assessed. The predictive performance of the model was calculated by area under the receiver operating characteristic curve (AUC) analysis, which showed a high accuracy for country of origin (AUC = 0.97), and a comparatively lower level of predictive accuracy for obese state (AUC = 0.65) (**Figure 2**). Sex was predicted with AUC = 0.75, the diabetes status with AUC = 0.63, hypertensive status with AUC = 0.65 and glucose status with AUC = 0.66. Random Forest analysis was also

used to identify the top 30 microbial taxonomic features that differentiate between countries and obese states. Similar to the ANCOMBC results, *Prevotella* and *Streptococcus* were at a greater proportion in the microbiota of Ghanaian and non-obese individuals, whereas *Mogibacterium* was at a greater proportion in the South African cohort. A greater proportion of *Megasphaera* was associated with the Jamaican cohort, while a greater proportion of Ruminococcaceae was observed in the American microbiota. *Weissella*, which was identified as having a significantly greater proportion in the Ghanaian cohort using ANCOMBC, was observed to be a discriminatory feature for Seychelles microbiota using Random Forest (**Supplementary Figure 2**).

Similarly, the predictive capacity of the gut microbiota features in stratifying individuals by obese state was assessed at each of the five study sites. The predictive performance of the model was calculated by AUC analysis, which showed a moderate accuracy for obese state for all sites, namely, Ghana (AUC = 0.57), South Africa (AUC = 0.52), Jamaica (AUC = 0.48), Seychelles (AUC = 0.43) and US (AUC = 0.52) (**Supplementary Figure 3**).

Predicted genetic metabolic potential differs by country and obesity status. The predicted potential microbial functional traits resulting from the compositional differences in microbial taxa between countries and obese state were assessed. PICRUST2 predicted a total of 372 MetaCyc functional pathways. ANCOM-BC analysis adjusted for sex, age and BMI identified 67 pathways ($p < 0.05$; false discovery rate (fdr)-corrected, $LFC > 1.4$) that accounted for discriminative features between the 4 different countries with the US (**Supplementary Figure 4**). In comparison with US, MetaCyc pathways differentially increased in Ghana and Jamaica include methylgallate degradation, norspermidine biosynthesis (PWY-6562), gallate degradation I pathway, gallate degradation II pathway, histamine degradation (PWY-6185), and toluene degradation III (via p-cresol) (PWY-5181). South African samples had a greater proportion of L-glutamate degradation VIII (to propanoate) (PWY-5088), isopropanol biosynthesis (PWY-6876), creatinine degradation (PWY-4722), adenosyl cobalamin biosynthesis (anaerobic) (PWY-5507), respiration I (cytochrome c) (PWY-3781). MetaCyc pathways linked to norspermidine biosynthesis (PWY-6562), mycothiol biosynthesis (PWY1G-0), were at a greater proportion in the Seychelles samples, whereas reductive acetyl coenzyme A (CODH-PWY), and chorismate biosynthesis II (PWY-6165) were depleted in the US samples. ANCOM-BC analysis adjusted for site, sex and age identified 24 predicted pathways that differentiated between obese and non-obese individuals (**Supplementary Figure 4**). Notably, the microbiota of non-obese individuals had a greater proportion of predicted pathways including the TCA cycle, amino acid metabolism (P162-PWY, PWY-5154, PWY-5345), ubiquinol biosynthesis-related pathways (PWY-5855, PWY-5856, PWY-5857, PWY-6708, UBISYN-PWY), cell structure biosynthesis and nucleic acid processing (PWY0845, PYRIDOXSYN-PWY).

Next, KEGG orthology (KO) involved in pathways related to butanoate (butyrate) metabolism and LPS biosynthesis were investigated. Predicted genes involved in butyrate biosynthesis pathways showed that enoyl-CoA hydratase enzymes (K01825, K01782, K01692), lysine, glutarate/succinate enzymes (K07250, K00135, K00247), glutarate/Acetyl CoA enzymes (K00175, K00174, K00242, K00241 K01040, K01039) were differentially abundant in participants from Ghana, South Africa, Jamaica, and Seychelles in comparison to the US cohort. The relative abundance of succinic semialdehyde reductase (K18122) was significantly increased only in South Africa, Jamaica, and Seychelles population. Further, predicted genes proportionally

abundant only in specific countries were observed. For instance, succinate semialdehyde dehydrogenase (K18119) was enriched only in the Ghanaian cohort, 4-hydroxybutyrate CoA-transferase (K18122) enriched among South African participants and lysine/glutarate/succinate enzyme (K14268) differentially abundant within the Seychelles population. The relative abundance of predicted genes encoded for enzymes such as maleate isomerase (K10799), 3-oxoacid CoA-transferase (K01027) and pyruvate/acetyl CoA (K00171, K00172, K00169) were greater in the US participants compared with participants from the 4 countries (**Supplementary Figure 5**). The non-obese exhibited a significantly greater abundance of genes that catalyze the production of butyrate via the fermentation of pyruvate or branched amino-acids such as enoyl-CoA hydratase enzyme (K0182), Leucine/Acetyl CoA enzyme (K01640) and pyruvate/acetyl CoA enzyme (K00171, K00172, K00169, K1907) by contrast obese individuals were differentially enriched for succinyl-CoA:acetate CoA-transferase (K18118) (**Supplementary Figure 5**). All analyses were adjusted for country, sex, BMI and age (fdr-corrected $p < 0.05$).

Several gut microbial predicted genes involved in LPS biosynthesis differentially enriched among the countries ($p < 0.05$; false discovery rate (fdr)-corrected) were identified. In particular, the relative abundance of specific LPS genes (K02560, K12973, K02849, K12979, K12975, K12974) were significantly enriched in Ghana, South Africa, Jamaica, and Seychelles when compared with US. Higher proportions of LPS genes including K12981, K12976, K09953, K03280 were significantly increased in Seychelles samples in comparison with US samples and also significantly increased in the US cohorts in comparison with participants from Ghana, South Africa, and Seychelles. US samples had a greater proportion of the following genes (K15669, K09778, K07264, K03273, K03271) in comparison with the other 4 countries (**Supplementary Figure 6**). Non-obese individuals had a greater abundance of predicted genes encoding LPS biosynthesis (K02841, K02843, K03271, K03273, K19353, K02850) whereas only 1 LPS gene (K02841) differentially elevated in the non-obese group (**Supplementary Figure 6**). All analyses were adjusted for country, sex, BMI and age (fdr-corrected $p < 0.05$).

Microbial community composition and predicted metabolic potential correlates with observed fecal SCFA concentrations. All countries had significantly higher weight-adjusted fecal total SCFA levels when compared to the US participants ($p < 0.001$), with Ghanaians having the highest weight-adjusted fecal total SCFA levels (**Supplementary Table 4**). When compared to their obese counterparts, non-obese participants had significantly higher weight-adjusted fecal total and individual SCFA levels (**Supplementary Table 5**). Total SCFA levels displayed weak, but significantly positive correlation with Shannon diversity ($r = 0.074$). A similar trend was observed in the different individual SCFAs, namely valerate ($r = 0.19$), butyrate ($r = 0.12$), propionate ($r = 0.073$) and acetate ($r = 0.058$) (**Figure 3**). Observed ASVs were not significantly correlated with total SCFAs ($p > 0.05$). Levels of acetate, butyrate and propionate exhibited strong significant correlations with total SCFA, whereas valerate levels significantly correlated negatively ($r = -0.09$) with total SCFAs. Next, we assessed if levels of total SCFAs could be predicted by a mixed model. Country explained 45.7% of the variation in SCFAs. No significant effect was explained either by obesity or Shannon diversity.

Microbial taxonomy correlates with SCFA concentration and obesity status. To explore the connection between SCFAs with gut microbiota, Spearman correlations between taxa that were proportionally significantly different between countries and concentrations of SCFAs were

determined. Valerate negatively correlated with the proportion of *Clostridium*, *Prevotella*, *Faecalibacterium*, *Roseburia* and *Streptococcus*, which were all positively correlated with acetate, propionate, and butyrate. Similarly, the proportions of Christensenellaceae, *Eubacterium*, and UCG 002 (Ruminococcaceae) were significantly positively associated with valerate, and negatively correlated with acetate, propionate, and butyrate. In addition, only a single ASV annotated to *Ruminococcus* was observed to be positively associated with all 4 SCFAs (**Figure 4**). Similarly, Spearman's rank correlation coefficients were calculated between the differentially abundant ASVs identified between obese and non-obese group with concentrations of SCFAs. Broadly, the proportions of most ASVs were significantly positively associated with acetate in comparison with the other 3 SCFAs. Consistent with the correlations mentioned above, valerate negatively correlated with most ASVs that were found to be positively correlated with the three major SCFAs, acetate, propionate, and butyrate and vice versa. The relative proportions of ASVs belonging to *Allisonella*, Erysipelotrichaceae and *Libanicoccus* positively correlated with acetate, propionate, and butyrate, whereas significantly negative relationships were observed between *Parabacteroides* and *Bacteroides* abundances with the aforementioned SCFAs. Valerate showed significantly positive associations with Oscillospirales and Ruminococcaceae abundances and significantly negative correlations with *Lachnospira* and *Eggerthella* abundances (**Figure 4**).

Discussion

By leveraging a well characterized large population-based cohort of African origin residing in geographically distinct regions of Ghana, South Africa, Jamaica, Seychelles, and the US, we examined the relationships between gut microbiota, SCFAs and adiposity. Our data revealed profound variations in gut microbiota, which are reflected in the significant changes in community composition, structure, and predicted functional pathways as a function of population obesity and geography, despite their shared ancestral background. Our data further revealed an inverse relation between fecal SCFA concentrations, microbial diversity, and obesity; importantly, the utility of the microbiota in predicting whether an individual was lean or obese was inversely correlated with the income-level of the country of origin. Overall, our findings are important for understanding the complex relationships between the gut microbiota, population lifestyle and the development of obesity, which may set the stage for defining the mechanisms through which the microbiome may shape health outcomes in populations of African-origin.

It has previously been reported that geographic origin can modulate the composition of the gut microbiota (Yatsunenko et al. 2012; De Filippo et al. 2010, 2017). Accordingly, taxonomic profiling revealed significant differences in gut microbiota richness and diversity among the different countries in a continuum manner. Notably, we detected greater microbiota diversity in Ghana, while depleted microbiota diversity was associated with the US, representing the lowest and the highest end along the epidemiologic transition spectrum respectively, while South Africans, Jamaicans and Seychellois ranked in between. Our findings are consistent with our previous METS studies (Fei et al. 2019; Dugas, Bernabé, et al. 2018) and other large scale continental cohort studies (De Filippo et al. 2010, 2017; Yatsunenko et al. 2012; Schnorr et al. 2014; Clemente et al. 2015; Rampelli et al. 2015; Gomez et al. 2016; Mancabelli et al. 2017), that report a higher bacterial diversity and composition/microbial richness in traditionally non-western groups that distinguish them from urban-industrialized individuals whose diets are low in fiber and high in

saturated fats (E. D. Sonnenburg and Sonnenburg 2019; Kolodziejczyk, Zheng, and Elinav 2019). Although we observe enrichment in the relative abundance of several taxa associated with country of origin in our cohorts, we also detect a pattern where the gut microbiota of Ghanaian and South African cohort tends to share many features, while the gut microbiota of the Jamaican cohort shared many features with all 4 countries, possibly reflecting the ongoing epidemiological transitional nature of their communities represented by the overlap with western and traditionally non-western populations. Notably, traditionally non-western associated taxa including *Prevotella*, *Butyrivibrio*, *Weissella* and *Romboutsia* were enriched in participants from Ghana and South Africa, as suggested previously (Mancabelli et al. 2017). Western-associated taxa such as *Bacteroides* and *Parabacteroides* were enriched in individuals from Jamaica and the US (Mancabelli et al. 2017; Kao et al. 2015), while an ASV annotated as *Olsenella* was proportionally abundant in Seychelles microbiota. *Bifidobacterium* and *Aldercreutzia* were enriched in the US cohort. *Clostridium sensu stricto* 1 was over-represented in all 4 countries in comparison with the US. We also found greater enrichment of VANISH taxa including *Butyricicoccus* and *Succinivibrio* in the Ghanaian cohort, in line with individuals practicing traditional lifestyles (Pasolli et al. 2019). *Prevotella* is usually associated with plant-based diets rich in dietary fibers, while *Bacteroides* abundance broadly correlates with diets high in fat, animal protein, and sugars (Gupta, Paul, and Dutta 2017; Wu et al. 2011), which is in agreement with our enterotype analysis where a *Prevotella*-rich microbiota dominates the Ghanaian and South African gut, while a *Bacteroides*-rich microbiota dominated in the high-income countries. *Prevotella* is known to produce high amounts of SCFAs (T. Chen et al. 2017), so its depletion may be associated with the observed concomitant reduction in SCFA concentrations. Increased SCFA synthesis is associated with a reduction in obesity, which is supported by our observations, whereby elevated concentrations of total SCFA and a concomitant reduction in obesity is associated with the *Prevotella* dominated gut of the Ghanaian cohort. Our results support a potential role for geography in reinforcing variations in the gut microbiota in our study cohort despite shared origin. Geography may reflect subtle shifts in lifestyle and/or environmental exposures including heterogeneity of dietary sources, exposure to medications, socioeconomic factors, medical history, and biogeographical patterns in microbial dispersion (Asnicar et al. 2021; Pasolli et al. 2019; Costello et al. 2012; He et al. 2018).

We also inferred the metabolic capacity of the gut microbiota associated with the different countries. Several metabolic pathways linked to carrier, cofactor and vitamin biosynthesis, biosynthesis/degradation of amines, amino acids, aromatic xenobiotics, and tricarboxylic acid (TCA) cycle were differentially enriched between the different countries compared with the US. These pathways are involved in biochemical reactions that regulate several processes including energy metabolism, inflammation, epigenetic processes, and oxidative stress. Several of these observed pathways have been reported in different populations (Yu et al. 2021; Karlsson et al. 2013; N. Qin et al. 2014) indicating that the gut microbiota can directly influence host metabolism, although a majority of these molecules can also be synthesized by the host or supplied through diet. In our cohort, functional shifts observed in participants from Ghana and Jamaica included the enrichment of the metabolic pathway for degradation of gallate. Metabolites generated from the gallate pathway include phenolic catechin metabolites which are thought to alleviate obesity-related pathologies and also promote a healthy and beneficial human gut microbiota composition (Marchesi et al. 2016; Liu et al. 2021). We found pathways related to glutamate degradation which

can be fermented to butyrate and propionate enhanced among South Africans and Ghanaians in comparison with the US. In Seychelles, a pathway involved in mycothiol biosynthesis was upregulated. Mycothiol is a protective antioxidant produced by the members of the Actinobacteria phylum and is involved in the removal of toxic compounds from cells (Newton, Buchmeier, and Fahey 2008). The predicted abundance of mycothiol biosynthesis pathway was identified as underrepresented in the microbiome of individuals with depressive symptoms in a South Korean population (S.-Y. Kim et al. 2022).

We further identified increased abundances in pathways related to the generation of SCFAs such as acetyl coenzyme A pathway, threonine biosynthesis and leucine degradation pathway in the microbiomes of all 4 countries in comparison with the US. Threonine can be metabolized to SCFAs acetate and propionate (Davila et al. 2013) and indeed, genes linked with threonine metabolism have been identified in the human gut microbiome (Abubucker et al. 2012). Taken together, our results suggest that the observed country-specific microbial differences and abundances accompany variance in the distribution of functional pathways abundances, although we are unable to ascertain what the sources are that may explain these differences in the predicted functional enrichment due the inherent limitation in functional resolution of 16S rRNA sequence data in PICRUST2 analysis. Further studies are required to evaluate the potential causal relations of these gut microbial functions with health outcomes using shotgun metagenomic sequencing which offers robust inferences of functional pathways.

Preclinical germ-free mouse models provide early causal links between gut microbial ecology and obesity (Ley et al. 2005; Bäckhed et al. 2007). Thereafter, follow up studies in human cohorts have sought to identify a consistent microbiota signature across populations that can be used to predict obesity. However, identifying obesity-specific microbiome features have proven difficult because the results are often not in agreement (Finucane et al. 2014). Therefore, we sought to examine the fecal levels of individual SCFA types and linking to variations in gut microbiota in obese and non-obese individuals in our large African cohort. The bulk of evidence from prior studies show that obesity is associated with a less diverse bacterial community (Turnbaugh et al. 2009; Dugas, Bernabé, et al. 2018; Peters et al. 2018). Accordingly, we observed that our obese group harbor a significantly lower microbiota diversity and differences in community composition.

Although the mechanism by which the gut microbiota influences obesity are not fully understood, several mechanisms have been proposed. For instance, the regulation of host energy metabolism and body mass concept demonstrate that a perturbed gut bacteria community contributes to the development of obesity by providing excess energy to the host via the fermentation of indigestible carbohydrates into SCFAs. Thus, the altered microbiota explains the ability of the host to extract energy from the diet and further stored in the adipose tissue (Turnbaugh et al. 2006; Jumpertz et al. 2011). In support of this notion, we identified several SCFA producing bacteria significantly under-represented or depleted in obese individuals, indicating that SCFAs beneficially regulate host energy metabolism. For example, the relative abundance of some members of *Oscillospira* have been reported to be markedly greater in healthy individuals and associates with human leanness (Beaumont et al. 2016; Konikoff and Gophna 2016; Gophna, Konikoff, and Nielsen 2017). *Oscillospira* utilizes host glycans to produce SCFAs (Konikoff and Gophna 2016; Gophna, Konikoff, and Nielsen 2017) including butyrate, with beneficial effects on insulin sensitivity, body weight control and inflammation (M.-H. Kim et al. 2020). One of the strongest links that has been

corroborated across several populations between a gut microbial taxa and BMI involves members of the *Christensenella* genus. They are known to produce SCFAs, acetate and butyrate (Morotomi, Nagai, and Watanabe 2012) and associate negatively with markers of obesity, much in agreement with our findings indicating that *Christensenella* may be important for promoting leanness. We also detected several butyrate producing ASVs including *Eubacterium*, *Alistipes*, *Clostridium* and *Odoribacter* to be proportionally enriched in individuals who were non-obese.

Although we observe more SCFA producing taxa in the non-obese group, we also identify taxa that are SCFA producers in the obese group. Notably we observe that obese individuals presented a greater abundance of *Lachnospira*, a finding consistent with our prior study in the same population (Dugas, Bernabé, et al. 2018), and others (Lippert et al. 2017; Meehan and Beiko 2014; de la Cuesta-Zuluaga, Corrales-Agudelo, et al. 2018). Contrary to our results, other studies have shown that a reduction in the abundance of *Lachnospira* positively associates with obesity (Comanys et al. 2021; Stanislowski et al. 2017). It is well known that there are many SCFA producing gut bacteria, raising questions about whether the observed features can be precisely attributed to this mechanism or pathway. However, our predicted functional analysis revealed that genes in the KEGG pathway related to SCFA butyrate synthesis (butanoate metabolism) were significantly depleted or underrepresented in the obese group compared to the non-obese counterparts, which further supports the concept that SCFAs are beneficial. Further, we identified several predicted genes involved in butyrate synthesis via the more-dominant pyruvate pathway in the non-obese group. Altogether, these results suggest that butyrate-producing bacteria may offer protection against obesity (X. Chen and Devaraj 2018). Indeed, butyrate exhibits immunomodulatory effects, improves colon mucosal barrier function, and lowers inflammation.

The SCFA producing microbes dominant in the non-obese group coincided with elevated fecal SCFA levels in these individuals compared with the obese group, which is in line with previous results from other studies that have explored the relation between concentrations of fecal SCFAs and obesity (Yin et al. 2022; Dugas, Bernabé, et al. 2018). Indeed, SCFA supplementation has been documented to protect against a high-fat diet-induced obesity in mice (H. V. Lin et al. 2012; Lu et al. 2016) as well as weight gain in humans (Chambers et al. 2015). Conversely, other studies, mostly from western populations have reported results contrary to our study (Schwartz et al. 2010; Fernandes et al. 2014; Riva et al. 2017; de la Cuesta-Zuluaga, Mueller, et al. 2018). For instance, de la Cuesta-Zuluaga et al observed associations between elevated fecal SCFA levels, central obesity, gut permeability, and hypertension in a Colombian cohort. The specific mechanisms that explain the higher fecal SCFA levels among obese individuals remain a matter of debate and one hypothesis is that disruptions in the obese gut microbiota may lead to less efficient SCFA absorption, hence the observed increased SCFA excretion (de la Cuesta-Zuluaga, Mueller, et al. 2018). Along the same line of notion, our findings of a negative association between obesity and SCFAs could be related to the consumption of diets enriched in fibers and other dietary precursors of SCFAs resulting in elevated SCFA production compared with SCFA absorption, thereby reducing energy harvesting and its associated storage as fat. Indeed, diets high in fiber and Mediterranean diets correlate positively with weight loss (Hu et al. 2013; Esposito et al. 2011) and increased levels of fecal SCFAs (De Filippis et al. 2016) in human studies. Other possible explanations for the observed divergences between our studies and others might be

attributed to differences in population, medication usage, sample size, microbial production capacity and intestinal absorption, underscoring the complex relationships between gut microbiota with SCFA production and host adiposity. Nevertheless, our results demonstrate that the negative associations between obesity and fecal SCFA levels in our study cohort are consistent with the positive associations found between decreased obesity and SCFA synthesizing microbes, although we are aware that fecal SCFA concentrations are not a direct measure of intestinal SCFA production but rather reflect a net result of the difference between SCFA production and absorption (Canfora, Jocken, and Blaak 2015). A measurement of the dynamics of SCFA production and availability with stable isotopes could be determined in future studies. Altogether, the observed differences in SCFA concentrations between obese and non-obese individuals and the several SCFA-producing microbes further reinforce the theory that gut microbiota and its associated SCFA metabolites may have a role in body weight regulation.

Another mechanism by which gut microbiota may contribute to obesity is via the metabolic endotoxemia pathway. Perturbations in the gut microbiota community composition lead to increased production of plasma lipopolysaccharide (LPS) derived from the cell wall of Gram-negative bacteria, provoking low-grade inflammation and increased intestinal permeability which drives adiposity (Zhao 2013; Cani et al. 2008). An increased relative abundance of one ASV assigned to the genus *Dialister* in the gut community of obese individuals was identified from this study. Zhang and colleagues reported proportional increases in *Dialister* in obese persons and suggested that could serve as a potential predictive marker for obesity (Zhang et al. 2021). Additionally, in our recent study (Fei et al. 2021) we observed an increased relative abundance of *Dialister* in subjects with short sleep duration, a condition associated with a chronic inflammatory state. Indeed, *Dialister* has been demonstrated to trigger or aggravate host inflammatory response and insulin resistance by releasing more lipopolysaccharides (Yang et al. 2022). To strengthen these findings, we further observed that several genes in the LPS biosynthesis pathway were differentially enriched within the obese group from our predicted functional analysis. Similar findings have previously been reported where the obese microbiota is enriched by LPS metabolism, initiating inflammation-dependent processes associated with the onset of obesity and insulin resistance (Boulangé et al. 2016) and other related metabolic diseases (Yan et al. 2021; Karlsson et al. 2012; Fei and Zhao 2013; Cani et al. 2007; Fei et al. 2021). Collectively, our results demonstrate that obese individuals harbor a marked inflammatory state favoring the development of obesity, and this is in concordance with the associated metabolic endotoxemia pathway linking gut bacteria to obesity.

This study additionally detected marked depletion in pathways involved in cell structure biosynthesis, vitamin B6 biosynthesis, NAD biosynthesis, amino acid metabolism and SCFA synthesis in our predicted metagenome analysis. Thus, our results further suggest that metabolic pathways important for growth, energy homeostasis and the maintenance of normal gut function are disrupted in individuals with obesity. Conversely, in the obese group, we noted an enrichment of formaldehyde assimilation I (serine pathway) pathway. Ubiquitous formaldehyde can be derived from food, the environment and generated endogenously as a result of human and microbial cellular metabolism of many methylated compounds. Endogenous formaldehyde produced at sufficient levels has carcinogenic properties and detrimental effects on genome stability. To counteract this reactive molecule, organisms have evolved a detoxification system

that converts formaldehyde to formate, a less reactive molecule that can be used for nucleotide biosynthesis (Reingruber and Pontel 2018; N. H. Chen et al. 2016). Thus, we may infer that the pattern of increased formaldehyde assimilation pathway in our data might result from a defect or diminished capacity of formaldehyde detoxification system pathway, an assumption which requires further verification. A study reported increases in the abundance of formaldehyde assimilation pathway in a depressed group when compared with non-depressed controls (S.-Y. Kim et al. 2022). We are the first to show that the gut of obese participants is enriched in the formaldehyde assimilation pathway. Although we do not understand the mechanistic details, it is known that toxic formaldehyde is generated along with reactive oxygen species during inflammatory processes (N. H. Chen et al. 2016). Thus, an increased capacity for formaldehyde pathway may indicate a microbiome-induced increase in reactive oxygen species in the gut of obese individuals. Indeed, prior work has identified induction of oxygen stress by microbial perturbations as one of the mechanisms by which the microbiome can promote weight gain and insulin resistance (J. Qin et al. 2012). The specific alterations of the gut microbiota and the associated predicted functionality may constitute a potential avenue for the development of microbiome-based therapeutics to treat obesity and/or to promote and sustain weight loss.

Study strengths and limitations. While our study has several strengths including a large sample size, diverse population along an epidemiological transition gradient with a comprehensive dataset that allowed the exclusion of the potential effects of origin as well as control of potential interpersonal covariates, and use of validated and standard tools for data collection, we acknowledge some limitations as well. First, the cross-sectional nature of our study design is unable to establish temporality or identify mechanisms by which the gut microbiome may causally influence the observed associations. In that regard, we expect that prospective data from the METS cohort study will provide the basis to assess the longitudinal association between gut microbiota composition, metabolites, and obesity, and we have an ongoing study exploring the potential correlations longitudinally. The use of 16S rRNA sequencing in our analysis for inferences on microbial functional ecology inherently has its limitations for drawing conclusions on species and strain level functionality due to its low resolution. Nevertheless, our results provide insight into the relationship between obesity, gut microbiota, and metabolic pathways in individuals of African-origin across different geographies, stimulating further examination of large-scale studies using multi-omic approaches with deeper taxonomic and functional resolution and animal transplantation studies to investigate potentially novel microbial strains and to explore the clinical relevance of the observed metabolic differences.

Conclusion

This study examined the relationship between the gut microbiota composition and functional patterns, concentrations of fecal short chain fatty acids (SCFAs) and obesity in a large population cohort of African origin, from Ghana, South Africa, Jamaica, Seychelles, and the United States of America, spanning the epidemiologic transition. The Ghanaian cohort exhibited the greatest gut microbiota diversity and the American cohort the least, with corresponding enrichment or depletion in taxa and predicted functional traits. Ghanaian participants were enriched in VANISH taxa reflecting their traditional lifestyle. Significant differences in gut microbiota composition and function were identified in obese individuals compared to the non-obese counterparts. Non-obese

individuals were enriched in SCFA-producing microbes which coincided with increased concentration of total SCFA in feces, extending the evidence that SCFAs mediate body weight regulation. The predictive accuracy of the microbiota for obesity status was greatest in low-income countries, and was reduced in high income countries, suggesting that lifestyle traits in high income countries may result in elevated obesity risk even for lean individuals. The specific alterations of the gut microbiota and the associated predicted metabolic function may constitute a potential avenue to guide the development of microbiome-based solutions to treat obesity and/or to promote and sustain weight loss. Thus, further examination of large-scale studies using multi-omic approaches with deeper taxonomic and functional resolution and animal transplantation studies are warranted to confirm the identified taxonomic and metabolic signatures.

Acknowledgements

We thank the METS participants who continue their ongoing participation in the METS studies, as well as the site-specific clinic staff in Ghana, South Africa, Jamaica, Seychelles and the US. The UCSD Microbiome Core performed sample extractions and library preparation utilizing protocols and primers published on the Earth Microbiome Project website (<https://earthmicrobiome.org/>). This publication includes data generated at the UC San Diego IGM Genomics Center utilizing an Illumina NovaSeq 6000 that was purchased with funding from a National Institutes of Health SIG grant (#S10 OD026929).

Contributions

LRD and BTL conceived the study. LRD, CC-K, PB, KB-A, JP-R, TEF, EVL, DR and AL collected human samples and metadata. GEM, CC-K, DR and AL curated metadata. SD performed sequencing of samples. GE-M, CC-K and MGM conducted formal analysis and visualization. JAG and LRD supervised and provided feedback on formal analysis and visualization. GEM, CC-K, MGM, LRD and JAG wrote the original manuscript. LRD secured the funding. All authors edited and approved the final manuscript.

References

- Abubucker, Sahar, Nicola Segata, Johannes Goll, Alyxandria M. Schubert, Jacques Izard, Brandi L. Cantarel, Beltran Rodriguez-Mueller, et al. 2012. "Metabolic Reconstruction for Metagenomic Data and Its Application to the Human Microbiome." *PLoS Computational Biology* 8 (6): e1002358.
- Agyemang, Charles, Sandra Boatemaa, Grace Agyemang Frempong, and Ama de-Graft Aikins. 2016. "Obesity in Sub-Saharan Africa." In *Metabolic Syndrome: A Comprehensive Textbook*, edited by Rexford S. Ahima, 41–53. Cham: Springer International Publishing.
- Amir, Amnon, Daniel McDonald, Jose A. Navas-Molina, Evguenia Kopylova, James T. Morton, Zhenjiang Zech Xu, Eric P. Kightley, et al. 2017. "Deblur Rapidly Resolves Single-Nucleotide Community Sequence Patterns." *MSystems* 2 (2). <https://doi.org/10.1128/mSystems.00191-16>.
- Arumugam, Manimozhiyan, Jeroen Raes, Eric Pelletier, Denis Le Paslier, Takuji Yamada, Daniel R. Mende, Gabriel R. Fernandes, et al. 2011. "Enterotypes of the Human Gut Microbiome." *Nature* 473 (7346): 174–80.
- Asnicar, Francesco, Sarah E. Berry, Ana M. Valdes, Long H. Nguyen, Gianmarco Piccinno, David A. Drew, Emily Leeming, et al. 2021. "Microbiome Connections with Host Metabolism and Habitual Diet from 1,098 Deeply Phenotyped Individuals." *Nature Medicine* 27 (2): 321–32.
- Bäckhed, Fredrik, Jill K. Manchester, Clay F. Semenkovich, and Jeffrey I. Gordon. 2007. "Mechanisms Underlying the Resistance to Diet-Induced Obesity in Germ-Free Mice." *Proceedings of the National Academy of Sciences of the United States of America* 104 (3): 979–84.
- Barengolts, Elena, Stefan J. Green, George E. Chlipala, Brian T. Layden, Yuval Eisenberg, Medha Priyadarshini, and Lara R. Dugas. 2019. "Predictors of Obesity among Gut Microbiota Biomarkers in African American Men with and without Diabetes." *Microorganisms* 7 (9). <https://doi.org/10.3390/microorganisms7090320>.
- Barone, Monica, Silvia Garelli, Simone Rampelli, Alessandro Agostini, Silke Matysik, Federica D'Amico, Sabrina Krautbauer, et al. 2022. "Multi-Omics Gut Microbiome Signatures in Obese Women: Role of Diet and Uncontrolled Eating Behavior." *BMC Medicine* 20 (1): 500.
- Beaumont, Michelle, Julia K. Goodrich, Matthew A. Jackson, Idil Yet, Emily R. Davenport, Sara Vieira-Silva, Justine Debelius, et al. 2016. "Heritable Components of the Human Fecal Microbiome Are Associated with Visceral Fat." *Genome Biology* 17 (1): 189.
- Bokulich, Nicholas A., Benjamin D. Kaehler, Jai Ram Rideout, Matthew Dillon, Evan Bolyen, Rob Knight, Gavin A. Huttley, and J. Gregory Caporaso. 2018. "Optimizing Taxonomic Classification of Marker-Gene Amplicon Sequences with QIIME 2's Q2-Feature-Classifer Plugin." *Microbiome* 6 (1): 90.
- Bolyen, Evan, Jai Ram Rideout, Matthew R. Dillon, Nicholas A. Bokulich, Christian C. Abnet, Gabriel A. Al-Ghalith, Harriet Alexander, et al. 2019. "Reproducible, Interactive, Scalable and Extensible Microbiome Data Science Using QIIME 2." *Nature Biotechnology* 37 (8): 852–57.
- Bonomo, Raiza R., Tyler M. Cook, Chaitanya K. Gavini, Chelsea R. White, Jacob R. Jones, Elisa Bovo, Aleksey V. Zima, et al. 2020. "Fecal Transplantation and Butyrate Improve Neuropathic Pain, Modify Immune Cell Profile, and Gene Expression in the PNS of Obese

833 Mice." *Proceedings of the National Academy of Sciences of the United States of America*
834 117 (42): 26482–93.

835 Boulangé, Claire L., Ana Luisa Neves, Julien Chilloux, Jeremy K. Nicholson, and Marc-Emmanuel
836 Dumas. 2016. "Impact of the Gut Microbiota on Inflammation, Obesity, and Metabolic
837 Disease." *Genome Medicine* 8 (1): 42.

838 Canfora, Emanuel E., Johan W. Jocken, and Ellen E. Blaak. 2015. "Short-Chain Fatty Acids in
839 Control of Body Weight and Insulin Sensitivity." *Nature Reviews. Endocrinology* 11 (10):
840 577–91.

841 Cani, Patrice D., Jacques Amar, Miguel Angel Iglesias, Marjorie Poggi, Claude Knauf, Delphine
842 Bastelica, Audrey M. Neyrinck, et al. 2007. "Metabolic Endotoxemia Initiates Obesity and
843 Insulin Resistance." *Diabetes* 56 (7): 1761–72.

844 Cani, Patrice D., Rodrigo Bibiloni, Claude Knauf, Aurélie Waget, Audrey M. Neyrinck, Nathalie M.
845 Delzenne, and Rémy Burcelin. 2008. "Changes in Gut Microbiota Control Metabolic
846 Endotoxemia-Induced Inflammation in High-Fat Diet-Induced Obesity and Diabetes in
847 Mice." *Diabetes* 57 (6): 1470–81.

848 Cani, Patrice D., and Nathalie M. Delzenne. 2009. "Interplay between Obesity and Associated
849 Metabolic Disorders: New Insights into the Gut Microbiota." *Current Opinion in*
850 *Pharmacology* 9 (6): 737–43.

851 Caspi, Ron, Richard Billington, Luciana Ferrer, Hartmut Foerster, Carol A. Fulcher, Ingrid M.
852 Keseler, Anamika Kothari, et al. 2016. "The MetaCyc Database of Metabolic Pathways
853 and Enzymes and the BioCyc Collection of Pathway/Genome Databases." *Nucleic Acids*
854 *Research* 44 (D1): D471-80.

855 Chambers, Edward S., Alexander Viardot, Arianna Psichas, Douglas J. Morrison, Kevin G.
856 Murphy, Sagen E. K. Zac-Varghese, Kenneth MacDougall, et al. 2015. "Effects of
857 Targeted Delivery of Propionate to the Human Colon on Appetite Regulation, Body Weight
858 Maintenance and Adiposity in Overweight Adults." *Gut* 64 (11): 1744–54.

859 Chen, Nathan H., Karrera Y. Djoko, Frédéric J. Veyrier, and Alastair G. McEwan. 2016.
860 "Formaldehyde Stress Responses in Bacterial Pathogens." *Frontiers in Microbiology* 7
861 (March): 257.

862 Chen, Tingting, Wenmin Long, Chenhong Zhang, Shuang Liu, Liping Zhao, and Bruce R.
863 Hamaker. 2017. "Fiber-Utilizing Capacity Varies in Prevotella- versus Bacteroides-
864 Dominated Gut Microbiota." *Scientific Reports* 7 (1): 2594.

865 Chen, Xinpu, and Sridevi Devaraj. 2018. "Gut Microbiome in Obesity, Metabolic Syndrome, and
866 Diabetes." *Current Diabetes Reports* 18 (12): 129.

867 Clemente, Jose C., Erica C. Pehrsson, Martin J. Blaser, Kuldip Sandhu, Zhan Gao, Bin Wang,
868 Magda Magris, et al. 2015. "The Microbiome of Uncontacted Amerindians." *Science*
869 *Advances* 1 (3). <https://doi.org/10.1126/sciadv.1500183>.

870 Companys, Judit, Maria José Gosalbes, Laura Pla-Pagà, Lorena Calderón-Pérez, Elisabet
871 Llauradó, Anna Pedret, Rosa Maria Valls, et al. 2021. "Gut Microbiota Profile and Its
872 Association with Clinical Variables and Dietary Intake in Overweight/Obese and Lean
873 Subjects: A Cross-Sectional Study." *Nutrients* 13 (6). <https://doi.org/10.3390/nu13062032>.

874 Costello, Elizabeth K., Keaton Stagaman, Les Dethlefsen, Brendan J. M. Bohannan, and David
875 A. Relman. 2012. "The Application of Ecological Theory toward an Understanding of the
876 Human Microbiome." *Science* 336 (6086): 1255–62.

- Cuesta-Zuluaga, Jacobo de la, Vanessa Corrales-Agudelo, Eliana P. Velásquez-Mejía, Jenny A. Carmona, José M. Abad, and Juan S. Escobar. 2018. "Gut Microbiota Is Associated with Obesity and Cardiometabolic Disease in a Population in the Midst of Westernization." *Scientific Reports* 8 (1): 11356.
- Cuesta-Zuluaga, Jacobo de la, Noel T. Mueller, Rafael Álvarez-Quintero, Eliana P. Velásquez-Mejía, Jelver A. Sierra, Vanessa Corrales-Agudelo, Jenny A. Carmona, José M. Abad, and Juan S. Escobar. 2018. "Higher Fecal Short-Chain Fatty Acid Levels Are Associated with Gut Microbiome Dysbiosis, Obesity, Hypertension and Cardiometabolic Disease Risk Factors." *Nutrients* 11 (1). <https://doi.org/10.3390/nu11010051>.
- Dahl, Wendy J., and Maria L. Stewart. 2015. "Position of the Academy of Nutrition and Dietetics: Health Implications of Dietary Fiber." *Journal of the Academy of Nutrition and Dietetics* 115 (11): 1861–70.
- Dalile, Boushra, Lukas Van Oudenhove, Bram Vervliet, and Kristin Verbeke. 2019. "The Role of Short-Chain Fatty Acids in Microbiota-Gut-Brain Communication." *Nature Reviews. Gastroenterology & Hepatology* 16 (8): 461–78.
- Davila, Anne-Marie, François Blachier, Martin Gotteland, Mireille Andriamihaja, Pierre-Henri Benetti, Yolanda Sanz, and Daniel Tomé. 2013. "Intestinal Luminal Nitrogen Metabolism: Role of the Gut Microbiota and Consequences for the Host." *Pharmacological Research: The Official Journal of the Italian Pharmacological Society* 68 (1): 95–107.
- De Filippis, Francesca, Nicoletta Pellegrini, Lucia Vannini, Ian B. Jeffery, Antonietta La Stora, Luca Laghi, Diana I. Serrazanetti, et al. 2016. "High-Level Adherence to a Mediterranean Diet Beneficially Impacts the Gut Microbiota and Associated Metabolome." *Gut* 65 (11): 1812–21.
- De Filippo, Carlotta, Duccio Cavalieri, Monica Di Paola, Matteo Ramazzotti, Jean Baptiste Poullet, Sebastien Massart, Silvia Collini, Giuseppe Pieraccini, and Paolo Lionetti. 2010. "Impact of Diet in Shaping Gut Microbiota Revealed by a Comparative Study in Children from Europe and Rural Africa." *Proceedings of the National Academy of Sciences of the United States of America* 107 (33): 14691–96.
- De Filippo, Carlotta, Monica Di Paola, Matteo Ramazzotti, Davide Albanese, Giuseppe Pieraccini, Elena Banci, Franco Miglietta, Duccio Cavalieri, and Paolo Lionetti. 2017. "Diet, Environments, and Gut Microbiota. A Preliminary Investigation in Children Living in Rural and Urban Burkina Faso and Italy." *Frontiers in Microbiology* 8 (October): 1979.
- DiBaise, John K., Husen Zhang, Michael D. Crowell, Rosa Krajmalnik-Brown, G. Anton Decker, and Bruce E. Rittmann. 2008. "Gut Microbiota and Its Possible Relationship with Obesity." *Mayo Clinic Proceedings. Mayo Clinic* 83 (4): 460–69.
- Douglas, Gavin M., Vincent J. Maffei, Jesse R. Zaneveld, Svetlana N. Yurgel, James R. Brown, Christopher M. Taylor, Curtis Huttenhower, and Morgan G. I. Langille. 2020. "PICRUSt2 for Prediction of Metagenome Functions." *Nature Biotechnology* 38 (6): 685–88.
- Dugas, Lara R., Beatriz Peñalver Bernabé, Medha Priyadarshini, Na Fei, Seo Jin Park, Laquita Brown, Jacob Plange-Rhule, et al. 2018. "Decreased Microbial Co-Occurrence Network Stability and SCFA Receptor Level Correlates with Obesity in African-Origin Women." *Scientific Reports* 8 (1): 17135.
- Dugas, Lara R., Louise Lie, Jacob Plange-Rhule, Kweku Bedu-Addo, Pascal Bovet, Estelle V. Lambert, Terrence E. Forrester, Amy Luke, Jack A. Gilbert, and Brian T. Layden. 2018. "Gut Microbiota, Short Chain Fatty Acids, and Obesity across the Epidemiologic Transition: The METS-Microbiome Study Protocol." *BMC Public Health* 18 (1): 978.

Esposito, Katherine, Christina-Maria Kastorini, Demosthenes B. Panagiotakos, and Dario Giugliano. 2011. "Mediterranean Diet and Weight Loss: Meta-Analysis of Randomized Controlled Trials." *Metabolic Syndrome and Related Disorders* 9 (1): 1–12.

Falony, Gwen, Marie Joossens, Sara Vieira-Silva, Jun Wang, Youssef Darzi, Karoline Faust, Alexander Kurilshikov, et al. 2016. "Population-Level Analysis of Gut Microbiome Variation." *Science* 352 (6285): 560–64.

Fei, Na, Beatriz Peñalver Bernabé, Louise Lie, Danny Baghdan, Kweku Bedu-Addo, Jacob Plange-Rhule, Terrence E. Forrester, et al. 2019. "The Human Microbiota Is Associated with Cardiometabolic Risk across the Epidemiologic Transition." *PloS One* 14 (7): e0215262.

Fei, Na, Candice Choo-Kang, Sirimon Reutrakul, Stephanie J. Crowley, Dale Rae, Kweku Bedu-Addo, Jacob Plange-Rhule, et al. 2021. "Gut Microbiota Alterations in Response to Sleep Length among African-Origin Adults." *PloS One* 16 (9): e0255323.

Fei, Na, and Liping Zhao. 2013. "An Opportunistic Pathogen Isolated from the Gut of an Obese Human Causes Obesity in Germfree Mice." *The ISME Journal* 7 (4): 880–84.

Fernandes, J., W. Su, S. Rahat-Rozenbloom, T. M. S. Wolever, and E. M. Comelli. 2014. "Adiposity, Gut Microbiota and Faecal Short Chain Fatty Acids Are Linked in Adult Humans." *Nutrition & Diabetes* 4 (6): e121.

Finucane, Mariel M., Thomas J. Sharpton, Timothy J. Laurent, and Katherine S. Pollard. 2014. "A Taxonomic Signature of Obesity in the Microbiome? Getting to the Guts of the Matter." *PloS One* 9 (1): e84689.

Gao, Zhanguo, Jun Yin, Jin Zhang, Robert E. Ward, Roy J. Martin, Michael Lefevre, William T. Cefalu, and Jianping Ye. 2009. "Butyrate Improves Insulin Sensitivity and Increases Energy Expenditure in Mice." *Diabetes* 58 (7): 1509–17.

Geng, Jiafeng, Qingqiang Ni, Wei Sun, Liangge Li, and Xiuqing Feng. 2022. "The Links between Gut Microbiota and Obesity and Obesity Related Diseases." *Biomedicine & Pharmacotherapy = Biomedecine & Pharmacotherapie* 147 (March): 112678.

Gomez, Andres, Klara J. Petrzalkova, Michael B. Burns, Carl J. Yeoman, Katherine R. Amato, Klara Vlckova, David Modry, et al. 2016. "Gut Microbiome of Coexisting BaAka Pygmies and Bantu Reflects Gradients of Traditional Subsistence Patterns." *Cell Reports* 14 (9): 2142–53.

Gonzalez, Antonio, Jose A. Navas-Molina, Tomasz Kosciolk, Daniel McDonald, Yoshiki Vázquez-Baeza, Gail Ackermann, Jeff DeReus, et al. 2018. "Qiita: Rapid, Web-Enabled Microbiome Meta-Analysis." *Nature Methods* 15 (10): 796–98.

Gophna, Uri, Tom Konikoff, and Henrik Bjørn Nielsen. 2017. "Oscillospira and Related Bacteria - From Metagenomic Species to Metabolic Features." *Environmental Microbiology* 19 (3): 835–41.

Gouda, Hebe N., Fiona Charlson, Katherine Sorsdahl, Sanam Ahmadzada, Alize J. Ferrari, Holly Erskine, Janni Leung, et al. 2019. "Burden of Non-Communicable Diseases in Sub-Saharan Africa, 1990-2017: Results from the Global Burden of Disease Study 2017." *The Lancet. Global Health* 7 (10): e1375–87.

Greenblum, Sharon, Peter J. Turnbaugh, and Elhanan Borenstein. 2012. "Metagenomic Systems Biology of the Human Gut Microbiome Reveals Topological Shifts Associated with Obesity and Inflammatory Bowel Disease." *Proceedings of the National Academy of Sciences of the United States of America* 109 (2): 594–99.

- Gupta, Vinod K., Sandip Paul, and Chitra Dutta. 2017. "Geography, Ethnicity or Subsistence-Specific Variations in Human Microbiome Composition and Diversity." *Frontiers in Microbiology* 8 (June): 1162.
- Hales, Craig M., Margaret D. Carroll, Cheryl D. Fryar, and Cynthia L. Ogden. 2020. "Prevalence of Obesity and Severe Obesity Among Adults: United States, 2017-2018." *NCHS Data Brief*, no. 360 (February): 1–8.
- He, Yan, Wei Wu, Hui-Min Zheng, Pan Li, Daniel McDonald, Hua-Fang Sheng, Mu-Xuan Chen, et al. 2018. "Regional Variation Limits Applications of Healthy Gut Microbiome Reference Ranges and Disease Models." *Nature Medicine* 24 (10): 1532–35.
- Hee, Bart van der, and Jerry M. Wells. 2021. "Microbial Regulation of Host Physiology by Short-Chain Fatty Acids." *Trends in Microbiology* 29 (8): 700–712.
- Henagan, Tara M., Barbara Stefanska, Zhide Fang, Alexandra M. Navard, Jianping Ye, Natalie R. Lenard, and Prasad P. Devarshi. 2015. "Sodium Butyrate Epigenetically Modulates High-Fat Diet-Induced Skeletal Muscle Mitochondrial Adaptation, Obesity and Insulin Resistance through Nucleosome Positioning." *British Journal of Pharmacology* 172 (11): 2782–98.
- Hu, Xiaojie, Jinlong Gao, Qianyan Zhang, Yuanqing Fu, Kelei Li, Shankuan Zhu, and Duo Li. 2013. "Soy Fiber Improves Weight Loss and Lipid Profile in Overweight and Obese Adults: A Randomized Controlled Trial." *Molecular Nutrition & Food Research* 57 (12): 2147–54.
- Jumpertz, Reiner, Duc Son Le, Peter J. Turnbaugh, Cathy Trinidad, Clifton Bogardus, Jeffrey I. Gordon, and Jonathan Krakoff. 2011. "Energy-Balance Studies Reveal Associations between Gut Microbes, Caloric Load, and Nutrient Absorption in Humans." *The American Journal of Clinical Nutrition* 94 (1): 58–65.
- Kao, Christina C., Julia L. Cope, Jean W. Hsu, Pratibha Dwarkanath, Jeffrey M. Karnes, Ruth A. Luna, Emily B. Hollister, Minerva M. Thame, Anura V. Kurpad, and Farook Jahoor. 2015. "The Microbiome, Intestinal Function, and Arginine Metabolism of Healthy Indian Women Are Different from Those of American and Jamaican Women." *The Journal of Nutrition* 146 (4): 706–13.
- Karlsson, Fredrik H., Frida Fåk, Intawat Nookaew, Valentina Tremaroli, Björn Fagerberg, Dina Petranovic, Fredrik Bäckhed, and Jens Nielsen. 2012. "Symptomatic Atherosclerosis Is Associated with an Altered Gut Metagenome." *Nature Communications* 3: 1245.
- Karlsson, Fredrik H., Valentina Tremaroli, Intawat Nookaew, Göran Bergström, Carl Johan Behre, Björn Fagerberg, Jens Nielsen, and Fredrik Bäckhed. 2013. "Gut Metagenome in European Women with Normal, Impaired and Diabetic Glucose Control." *Nature* 498 (7452): 99–103.
- Kembel, Steven W., Peter D. Cowan, Matthew R. Helmus, William K. Cornwell, Helene Morlon, David D. Ackerly, Simon P. Blomberg, and Campbell O. Webb. 2010. "Picante: R Tools for Integrating Phylogenies and Ecology." *Bioinformatics* 26 (11): 1463–64.
- Kim, Mi-Hyun, Kyung Eun Yun, Jimin Kim, Eunkyo Park, Yoosoo Chang, Seungho Ryu, Hyung-Lae Kim, and Han-Na Kim. 2020. "Gut Microbiota and Metabolic Health among Overweight and Obese Individuals." *Scientific Reports* 10 (1): 19417.
- Kim, Sun-Young, Eunkyo Park, Weon-Jeong Lim, Soo In Kim, Sang Won Jeon, Yoosoo Chang, Seungho Ryu, Hyung-Lae Kim, and Han-Na Kim. 2022. "Association Between Gut Microbiota and Depressive Symptoms: A Cross-Sectional Population-Based Study in South Korea." *Psychosomatic Medicine* 84 (7): 757–65.

- Koh, Ara, Filipe De Vadder, Petia Kovatcheva-Datchary, and Fredrik Bäckhed. 2016. "From Dietary Fiber to Host Physiology: Short-Chain Fatty Acids as Key Bacterial Metabolites." *Cell* 165 (6): 1332–45.
- Kolodziejczyk, Aleksandra A., Danping Zheng, and Eran Elinav. 2019. "Diet-Microbiota Interactions and Personalized Nutrition." *Nature Reviews. Microbiology* 17 (12): 742–53.
- Konikoff, Tom, and Uri Gophna. 2016. "Oscillospira: A Central, Enigmatic Component of the Human Gut Microbiota." *Trends in Microbiology* 24 (7): 523–24.
- Le Chatelier, Emmanuelle, Trine Nielsen, Junjie Qin, Edi Prifti, Falk Hildebrand, Gwen Falony, Mathieu Almeida, et al. 2013. "Richness of Human Gut Microbiome Correlates with Metabolic Markers." *Nature* 500 (7464): 541–46.
- Lewandowski, Cutler T., Md Wasim Khan, Manel BenAissa, Oleksii Dubrovskiy, Martha Ackerman-Berrier, Mary Jo LaDu, Brian T. Layden, and Gregory R. J. Thatcher. 2021. "Metabolomic Analysis of a Selective ABCA1 Inducer in Obesogenic Challenge Provides a Rationale for Therapeutic Development." *EBioMedicine* 66 (April): 103287.
- Ley, Ruth E. 2010. "Obesity and the Human Microbiome." *Current Opinion in Gastroenterology* 26 (1): 5–11.
- Ley, Ruth E., Fredrik Bäckhed, Peter Turnbaugh, Catherine A. Lozupone, Robin D. Knight, and Jeffrey I. Gordon. 2005. "Obesity Alters Gut Microbial Ecology." *Proceedings of the National Academy of Sciences of the United States of America* 102 (31): 11070–75.
- Ley, Ruth E., Peter J. Turnbaugh, Samuel Klein, and Jeffrey I. Gordon. 2006. "Microbial Ecology: Human Gut Microbes Associated with Obesity." *Nature* 444 (7122): 1022–23.
- Lin, Hua V., Andrea Frassetto, Edward J. Kowalik Jr, Andrea R. Nawrocki, Mofei M. Lu, Jennifer R. Kosinski, James A. Hubert, et al. 2012. "Butyrate and Propionate Protect against Diet-Induced Obesity and Regulate Gut Hormones via Free Fatty Acid Receptor 3-Independent Mechanisms." *PloS One* 7 (4): e35240.
- Lin, Huang, and Shyamal Das Peddada. 2020. "Analysis of Compositions of Microbiomes with Bias Correction." *Nature Communications* 11 (1): 3514.
- Lippert, K., L. Kedenko, L. Antonielli, I. Kedenko, C. Gemeier, M. Leitner, A. Kautzky-Willer, B. Paulweber, and E. Hackl. 2017. "Gut Microbiota Dysbiosis Associated with Glucose Metabolism Disorders and the Metabolic Syndrome in Older Adults." *Beneficial Microbes* 8 (4): 545–56.
- Liu, Xiaoxia, Ke Zhao, Nana Jing, Qingjun Kong, and Xingbin Yang. 2021. "Epigallocatechin Gallate (EGCG) Promotes the Immune Function of Ileum in High Fat Diet Fed Mice by Regulating Gut Microbiome Profiling and Immunoglobulin Production." *Frontiers in Nutrition* 8 (September): 720439.
- Lozupone, Catherine, and Rob Knight. 2005. "UniFrac: A New Phylogenetic Method for Comparing Microbial Communities." *Applied and Environmental Microbiology* 71 (12): 8228–35.
- Lu, Yuanyuan, Chaonan Fan, Ping Li, Yanfei Lu, Xuelian Chang, and Kemin Qi. 2016. "Short Chain Fatty Acids Prevent High-Fat-Diet-Induced Obesity in Mice by Regulating G Protein-Coupled Receptors and Gut Microbiota." *Scientific Reports* 6 (November): 37589.
- Luke, Amy, Pascal Bovet, Terrence E. Forrester, Estelle V. Lambert, Jacob Plange-Rhule, Dale A. Schoeller, Lara R. Dugas, et al. 2011. "Protocol for the Modeling the Epidemiologic Transition Study: A Longitudinal Observational Study of Energy Balance and Change in

- 1057 Body Weight, Diabetes and Cardiovascular Disease Risk.” *BMC Public Health* 11
1058 (December): 927.
- 1059 Mancabelli, Leonardo, Christian Milani, Gabriele Andrea Lugli, Francesca Turrone, Chiara
1060 Ferrario, Douwe van Sinderen, and Marco Ventura. 2017. “Meta-Analysis of the Human
1061 Gut Microbiome from Urbanized and Pre-Agricultural Populations.” *Environmental*
1062 *Microbiology* 19 (4): 1379–90.
- 1063 Marchesi, Julian R., David H. Adams, Francesca Fava, Gerben D. A. Hermes, Gideon M.
1064 Hirschfield, Georgina Hold, Mohammed Nabil Quraishi, et al. 2016. “The Gut Microbiota
1065 and Host Health: A New Clinical Frontier.” *Gut* 65 (2): 330–39.
- 1066 McLaren, Michael R. 2020. *Silva SSU Taxonomic Training Data Formatted for DADA2 (Silva*
1067 *Version 138)*. <https://doi.org/10.5281/zenodo.3731176>.
- 1068 McMurdie, Paul J., and Susan Holmes. 2013. “Phyloseq: An R Package for Reproducible
1069 Interactive Analysis and Graphics of Microbiome Census Data.” *PloS One* 8 (4): e61217.
- 1070 Meehan, Conor J., and Robert G. Beiko. 2014. “A Phylogenomic View of Ecological Specialization
1071 in the Lachnospiraceae, a Family of Digestive Tract-Associated Bacteria.” *Genome*
1072 *Biology and Evolution* 6 (3): 703–13.
- 1073 Mehta, Supal, Lara Ruth Dugas, Candice Choo-Kang, Pascal Bovet, Terrence Forrester, Kweku
1074 Bedu-Addo, Estelle Vicki Lambert, et al. 2021. “Consumption of Monounsaturated Fatty
1075 Acids Is Associated with Improved Cardiometabolic Outcomes in Four African-Origin
1076 Populations Spanning the Epidemiologic Transition.” *Nutrients* 13 (7).
1077 <https://doi.org/10.3390/nu13072442>.
- 1078 Mirarab, S., N. Nguyen, and T. Warnow. 2012. “SEPP: SATé-Enabled Phylogenetic Placement.”
1079 *Pacific Symposium on Biocomputing. Pacific Symposium on Biocomputing*, 247–58.
- 1080 Moreau, N. M., S. M. Goupy, J. P. Antignac, F. J. Monteau, B. J. Le Bizec, M. M. Champ, L. J.
1081 Martin, and H. J. Dumon. 2003. “Simultaneous Measurement of Plasma Concentrations
1082 and ¹³C-Enrichment of Short-Chain Fatty Acids, Lactic Acid and Ketone Bodies by Gas
1083 Chromatography Coupled to Mass Spectrometry.” *Journal of Chromatography. B,*
1084 *Analytical Technologies in the Biomedical and Life Sciences* 784 (2): 395–403.
- 1085 Morotomi, Masami, Fumiko Nagai, and Yohei Watanabe. 2012. “Description of Christensenella
1086 Minuta Gen. Nov., Sp. Nov., Isolated from Human Faeces, Which Forms a Distinct Branch
1087 in the Order Clostridiales, and Proposal of Christensenellaceae Fam. Nov.” *International*
1088 *Journal of Systematic and Evolutionary Microbiology* 62 (Pt 1): 144–49.
- 1089 “National Diabetes Statistics Report.” 2022. June 29, 2022.
1090 <https://www.cdc.gov/diabetes/data/statistics-report/index.html>.
- 1091 Navarro, Guadalupe, Anukriti Sharma, Lara R. Dugas, Terrence Forrester, Jack A. Gilbert, and
1092 Brian T. Layden. 2018. “Gut Microbial Features Can Predict Host Phenotype Response to
1093 Protein Deficiency.” *Physiological Reports* 6 (23): e13932.
- 1094 Newton, Gerald L., Nancy Buchmeier, and Robert C. Fahey. 2008. “Biosynthesis and Functions
1095 of Mycothiol, the Unique Protective Thiol of Actinobacteria.” *Microbiology and Molecular*
1096 *Biology Reviews: MMBR* 72 (3): 471–94.
- 1097 Nooromid, Michael, Edmund B. Chen, Liqun Xiong, Katherine Shapiro, Qun Jiang, Faleen Demas,
1098 Maeve Eskandari, et al. 2020. “Microbe-Derived Butyrate and Its Receptor, Free Fatty
1099 Acid Receptor 3, But Not Free Fatty Acid Receptor 2, Mitigate Neointimal Hyperplasia
1100 Susceptibility After Arterial Injury.” *Journal of the American Heart Association* 9 (13):
1101 e016235.

1102 Nordmo, Morten, Yngvild Sørenbø Danielsen, and Magnus Nordmo. 2020. "The Challenge of
1103 Keeping It off, a Descriptive Systematic Review of High-Quality, Follow-up Studies of
1104 Obesity Treatments." *Obesity Reviews: An Official Journal of the International Association
1105 for the Study of Obesity* 21 (1): e12949.

1106 "Obesity and Overweight." n.d. Accessed February 25, 2023. [https://www.who.int/news-](https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight)
1107 [room/fact-sheets/detail/obesity-and-overweight](https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight).

1108 Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, and H. Wagner. 2013. "Vegan: Community
1109 Ecology Package. R Package Version. 2.0-10," January. <http://dx.doi.org/>.

1110 Pasolli, Edoardo, Francesco Asnicar, Serena Manara, Moreno Zolfo, Nicolai Karcher, Federica
1111 Armanini, Francesco Beghini, et al. 2019. "Extensive Unexplored Human Microbiome
1112 Diversity Revealed by Over 150,000 Genomes from Metagenomes Spanning Age,
1113 Geography, and Lifestyle." *Cell* 176 (3): 649-662.e20.

1114 Peters, Brandilyn A., Jean A. Shapiro, Timothy R. Church, George Miller, Chau Trinh-Shevrin,
1115 Elizabeth Yuen, Charles Friedlander, Richard B. Hayes, and Jiyoung Ahn. 2018. "A
1116 Taxonomic Signature of Obesity in a Large Study of American Adults." *Scientific Reports*
1117 8 (1): 9749.

1118 Qin, Junjie, Yingrui Li, Zhiming Cai, Shenghui Li, Jianfeng Zhu, Fan Zhang, Suisha Liang, et al.
1119 2012. "A Metagenome-Wide Association Study of Gut Microbiota in Type 2 Diabetes."
1120 *Nature* 490 (7418): 55–60.

1121 Qin, Nan, Fengling Yang, Ang Li, Edi Prifti, Yanfei Chen, Li Shao, Jing Guo, et al. 2014.
1122 "Alterations of the Human Gut Microbiome in Liver Cirrhosis." *Nature* 513 (7516): 59–64.

1123 Rahat-Rozenbloom, S., J. Fernandes, G. B. Gloor, and T. M. S. Wolever. 2014. "Evidence for
1124 Greater Production of Colonic Short-Chain Fatty Acids in Overweight than Lean Humans."
1125 *International Journal of Obesity* 38 (12): 1525–31.

1126 Rampelli, Simone, Stephanie L. Schnorr, Clarissa Consolandi, Silvia Turrone, Marco Severgnini,
1127 Clelia Peano, Patrizia Brigidi, Alyssa N. Crittenden, Amanda G. Henry, and Marco
1128 Candela. 2015. "Metagenome Sequencing of the Hadza Hunter-Gatherer Gut Microbiota."
1129 *Current Biology: CB* 25 (13): 1682–93.

1130 Reiman, Derek, Brian T. Layden, and Yang Dai. 2021. "MiMeNet: Exploring Microbiome-
1131 Metabolome Relationships Using Neural Networks." *PLoS Computational Biology* 17 (5):
1132 e1009021.

1133 Reingruber, Hernán, and Lucas Blas Pontel. 2018. "Formaldehyde Metabolism and Its Impact on
1134 Human Health." *Current Opinion in Toxicology* 9 (June): 28–34.

1135 Richardson, A. J., A. G. Calder, C. S. Stewart, and A. Smith. 1989. "Simultaneous Determination
1136 of Volatile and Non-volatile Acidic Fermentation Products of Anaerobes by Capillary Gas
1137 Chromatography." *Letters in Applied Microbiology* 9 (1): 5–8.

1138 Ridaura, Vanessa K., Jeremiah J. Faith, Federico E. Rey, Jiye Cheng, Alexis E. Duncan, Andrew
1139 L. Kau, Nicholas W. Griffin, et al. 2013. "Gut Microbiota from Twins Discordant for Obesity
1140 Modulate Metabolism in Mice." *Science* 341 (6150): 1241214.

1141 Riva, Alessandra, Francesca Borgo, Carlotta Lassandro, Elvira Verduci, Giulia Morace, Elisa
1142 Borghi, and David Berry. 2017. "Pediatric Obesity Is Associated with an Altered Gut
1143 Microbiota and Discordant Shifts in Firmicutes Populations." *Environmental Microbiology*
1144 19 (1): 95–105.

1145 Roth, Gregory A., George A. Mensah, Catherine O. Johnson, Giovanni Addolorato, Enrico
1146 Ammirati, Larry M. Baddour, Noël C. Barengo, et al. 2020. "Global Burden of
1147 Cardiovascular Diseases and Risk Factors, 1990-2019: Update From the GBD 2019
1148 Study." *Journal of the American College of Cardiology* 76 (25): 2982–3021.

1149 Sanna, Serena, Natalie R. van Zuydam, Anubha Mahajan, Alexander Kurilshikov, Arnau Vich
1150 Vila, Urmo Vösa, Zlatan Mujagic, et al. 2019. "Causal Relationships among the Gut
1151 Microbiome, Short-Chain Fatty Acids and Metabolic Diseases." *Nature Genetics* 51 (4):
1152 600–605.

1153 Schnorr, Stephanie L., Marco Candela, Simone Rampelli, Manuela Centanni, Clarissa
1154 Consolandi, Giulia Basaglia, Silvia Turroni, et al. 2014. "Gut Microbiome of the Hadza
1155 Hunter-Gatherers." *Nature Communications* 5 (April): 3654.

1156 Schwartz, Andreas, David Taras, Klaus Schäfer, Silvia Beijer, Nicolaas A. Bos, Christiane Donus,
1157 and Philip D. Hardt. 2010. "Microbiota and SCFA in Lean and Overweight Healthy
1158 Subjects." *Obesity* 18 (1): 190–95.

1159 Sonnenburg, Erica D., and Justin L. Sonnenburg. 2019. "The Ancestral and Industrialized Gut
1160 Microbiota and Implications for Human Health." *Nature Reviews. Microbiology* 17 (6): 383–
1161 90.

1162 Sonnenburg, Justin L., and Fredrik Bäckhed. 2016. "Diet-Microbiota Interactions as Moderators
1163 of Human Metabolism." *Nature* 535 (7610): 56–64.

1164 Stanislawski, Maggie A., Dana Dabelea, Brandie D. Wagner, Marci K. Sontag, Catherine A.
1165 Lozupone, and Merete Eggesbø. 2017. "Pre-Pregnancy Weight, Gestational Weight Gain,
1166 and the Gut Microbiota of Mothers and Their Infants." *Microbiome* 5 (1): 113.

1167 Teixeira, Tatiana F. S., Łukasz Grześkowiak, Sylvia C. C. Franceschini, Josefina Bressan, Célio
1168 L. L. F. Ferreira, and Maria C. G. Peluzio. 2013. "Higher Level of Faecal SCFA in Women
1169 Correlates with Metabolic Syndrome Risk Factors." *The British Journal of Nutrition* 109
1170 (5): 914–19.

1171 Thompson, L. R., J. G. Sanders, D. McDonald, and A. Amir. 2017. "A Communal Catalogue
1172 Reveals Earth's Multiscale Microbial Diversity." *Nature*.
1173 <https://www.nature.com/articles/nature24621?report=reader>.

1174 Tseng, Ching-Hung, and Chun-Ying Wu. 2019. "The Gut Microbiome in Obesity." *Journal of the*
1175 *Formosan Medical Association = Taiwan Yi Zhi* 118 Suppl 1 (March): S3–9.

1176 Turnbaugh, Peter J., Fredrik Bäckhed, Lucinda Fulton, and Jeffrey I. Gordon. 2008. "Diet-Induced
1177 Obesity Is Linked to Marked but Reversible Alterations in the Mouse Distal Gut
1178 Microbiome." *Cell Host & Microbe* 3 (4): 213–23.

1179 Turnbaugh, Peter J., Micah Hamady, Tanya Yatsunenko, Brandi L. Cantarel, Alexis Duncan, Ruth
1180 E. Ley, Mitchell L. Sogin, et al. 2009. "A Core Gut Microbiome in Obese and Lean Twins." *Nature* 457 (7228): 480–84.

1182 Turnbaugh, Peter J., Ruth E. Ley, Michael A. Mahowald, Vincent Magrini, Elaine R. Mardis, and
1183 Jeffrey I. Gordon. 2006. "An Obesity-Associated Gut Microbiome with Increased Capacity
1184 for Energy Harvest." *Nature* 444 (7122): 1027–31.

1185 Valdes, Ana M., Jens Walter, Eran Segal, and Tim D. Spector. 2018. "Role of the Gut Microbiota
1186 in Nutrition and Health." *BMJ* 361 (June): k2179.

1187 Vinolo, Marco A. R., Hosana G. Rodrigues, Renato T. Nachbar, and Rui Curi. 2011. "Regulation
1188 of Inflammation by Short Chain Fatty Acids." *Nutrients* 3 (10): 858–76.

- Vrieze, Anne, Els Van Nood, Frits Holleman, Jarkko Salojärvi, Ruud S. Kootte, Joep F. W. M. Bartelsman, Geesje M. Dallinga-Thie, et al. 2012. "Transfer of Intestinal Microbiota from Lean Donors Increases Insulin Sensitivity in Individuals with Metabolic Syndrome." *Gastroenterology* 143 (4): 913-6.e7.
- Walters, William, Embriette R. Hyde, Donna Berg-Lyons, Gail Ackermann, Greg Humphrey, Alma Parada, Jack A. Gilbert, et al. 2016. "Improved Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers for Microbial Community Surveys." *mSystems* 11 (1). <https://doi.org/10.1128/mSystems.00009-15>.
- Wu, Gary D., Jun Chen, Christian Hoffmann, Kyle Bittinger, Ying-Yu Chen, Sue A. Keilbaugh, Meenakshi Bewtra, et al. 2011. "Linking Long-Term Dietary Patterns with Gut Microbial Enterotypes." *Science* 334 (6052): 105–8.
- Yan, Hang, Qian Qin, Jengfeng Chen, Su Yan, Tiantian Li, Xinxin Gao, Yang Yang, Ang Li, and Suying Ding. 2021. "Gut Microbiome Alterations in Patients With Visceral Obesity Based on Quantitative Computed Tomography." *Frontiers in Cellular and Infection Microbiology* 11: 823262.
- Yang, Teng, Leho Tedersoo, Pamela S. Soltis, Douglas E. Soltis, Miao Sun, Yuying Ma, Yingying Ni, et al. 2022. "Plant and Fungal Species Interactions Differ between Aboveground and Belowground Habitats in Mountain Forests of Eastern China." *Science China. Life Sciences*, December. <https://doi.org/10.1007/s11427-022-2174-3>.
- Yatsunencko, Tanya, Federico E. Rey, Mark J. Manary, Indi Trehan, Maria Gloria Dominguez-Bello, Monica Contreras, Magda Magris, et al. 2012. "Human Gut Microbiome Viewed across Age and Geography." *Nature* 486 (7402): 222–27.
- Yin, Xing-Qi, Ya-Xin An, Cai-Guo Yu, Jing Ke, Dong Zhao, and Ke Yu. 2022. "The Association Between Fecal Short-Chain Fatty Acids, Gut Microbiota, and Visceral Fat in Monozygotic Twin Pairs." *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy* 15 (February): 359–68.
- Yu, Danxia, Yaohua Yang, Jirong Long, Wanghong Xu, Qiuyin Cai, Jie Wu, Hui Cai, Wei Zheng, and Xiao-Ou Shu. 2021. "Long-Term Diet Quality and Gut Microbiome Functionality: A Prospective, Shotgun Metagenomic Study among Urban Chinese Adults." *Current Developments in Nutrition* 5 (4): nzab026.
- Zhang, Qiang, Rong Zou, Min Guo, Mengmeng Duan, Quan Li, and Huajun Zheng. 2021. "Comparison of Gut Microbiota between Adults with Autism Spectrum Disorder and Obese Adults." *PeerJ* 9 (March): e10946.
- Zhao, Liping. 2013. "The Gut Microbiota and Obesity: From Correlation to Causality." *Nature Reviews. Microbiology* 11 (9): 639–47.
- Zhernakova, Alexandra, Alexander Kurilshikov, Marc Jan Bonder, Ettje F. Tigchelaar, Melanie Schirmer, Tommi Vatanen, Zlatan Mujagic, et al. 2016. "Population-Based Metagenomics Analysis Reveals Markers for Gut Microbiome Composition and Diversity." *Science* 352 (6285): 565–69.

Table 1. METS-Microbiome participant characteristics from Ghana, South Africa, Jamaica, Seychelles and US

<i>Women</i>					
	Ghana	South Africa	Jamaica	Seychelles	US
	n=254	n=228	n=263	n=196	n=213
<i>Age (years)</i>	40.74 ± 8.1	35.56 ± 7.8	45.16 ± 7.5	43.84 ± 6.1	45.44 ± 6.4
<i>BMI (kg/m²)</i>	28.30 ± 5.9	33.42 ± 8.6	32.12 ± 7.3	30.32 ± 7.2	36.34 ± 8.8
<i>Obese (%)</i>	45,0%	61,0%	60,4%	49,5%	74,7%
<i>SBP (mm Hg)</i>	117.1 ± 18.5	115.20 ± 17.1	126.08 ± 19.0	123.28 ± 17.8	124.19 ± 18.4
<i>DBP (mm Hg)</i>	70.53 ± 12.2	75.20 ± 12.1	79.41 ± 12.6	79.37 ± 14.4	81.52 ± 12.1
<i>Hypertensive (%)</i>	37,5%	37,3%	57,4%	55,5%	65,4%
<i>Glucose (mg/dL)</i>	110.45 ± 62.7	89.17 ± 20.0	107.46 ± 39.1	111.35 ± 27.2	107.07 ± 44.0
<i>Diabetic (%)</i>	10,0%	3,5%	12,9%	13,9%	19,9%
<i>Men</i>					
	Ghana	South Africa	Jamaica	Seychelles	US
	n=117	n=171	n=133	n=164	n=107
<i>Age (years)</i>	43.92 ± 8.7	36.53 ± 7.2	44.42 ± 7.5	44.57 ± 5.1	47.12 ± 5.5
<i>BMI (kg/m²)</i>	23.7 ± 4.4	22.26 ± 4.1	24.8 ± 5.3	28.46 ± 5.5	30.37 ± 8.2
<i>Obese (%)</i>	13,4%	5,3%	15,7%	39,2%	44,4%
<i>SBP (mm Hg)</i>	121.28 ± 15.4	122.71 ± 15.5	129.23 ± 17.1	130.43 ± 16.2	130.67 ± 16.0
<i>DBP (mm Hg)</i>	68.02 ± 13.0	75.32 ± 11.1	78.07 ± 11.5	81.64 ± 12.1	82.37 ± 12.2
<i>Hypertensive (%)</i>	33,1%	45,0%	50,3%	65,9%	72,7%
<i>Glucose (mg/dL)</i>	100.52 ± 19.4	94 ± 23.4	99.04 ± 33.1	124.26 ± 44.2	107 ± 36.2
<i>Diabetic (%)</i>	4,6%	3,5%	4,8%	22,8%	17,5%

Table 2. Alpha diversity estimated by Shannon, Observed ASVs and Faith's PD (Phylogenetic Diversity) between countries and obesity status. q-value are FDR-corrected p values representing statistical significance ($p < 0.05$) of alpha diversity metrics between the countries. Data are presented by median (interquartile range). FDR = False Discovery Rate

		N	Faith's PD	Shannon	Observed
Ghana	Obese	243	19.2(16.2-21.8)	3.73(3.41,4.09)	228(184,267)
	Non-Obese	89	17.9(14.6,21.6)	3.69(3.30,4.05)	217(155,252)
South Africa	Obese	208	17.2(14.0,19.9)	3.21(2.69,3.52)	174(138,212)
	Non-Obese	179	15.9(13.1,19.9)	3.12(2.65,3.54)	165(126,216)
Jamaica	Obese	217	14.2(11.6,17.2)	3.2(2.72,3.56)	146(110,184)
	Non-Obese	147	13.4(11.5,16.5)	3.15(2.69,3.56)	136(108,173)
Seychelles	Obese	233	18(15.1,20.4)	3.51(3.14,3.79)	204(166,246)
	Non-Obese	141	18.6(15.0,22.4)	3.62(3.21,3.98)	212(165,269)
US	Obese	112	13.6(12.1,16.6)	3.23(2.86,3.50)	144(125,180)
	Non-Obese	195	13.9(12.1,16.5)	3.3(3.00,3.57)	150(122,184)
p-value			<0.001	<0.001	<0.001
q-value			<0.001	<0.001	<0.001

Median (IQR)

p-value: Kruskal-Wallis rank sum test

q-value: False discovery rate correction for multiple testing

Table 3. Adjusted Multivariate Analysis for the entire cohort and by each country.
Statistical significance from permutational multivariate analysis of variance
(PERMANOVA) test, $p < 0.05$. All p -values are generated based on 999 permutations

	Overall		Ghana		South Africa		Jamaica		Seychelles		US	
	R ²	P	R ²	P	R ²	P	R ²	P	R ²	P	R ²	P
Obese	0.003	0.001	0.004	0.032	0.007	0.002	0.002	0.732	0.003	0.279	0.004	0.154
Sex	0.003	0.001	0.005	0.018	0.007	0.002	0.009	0.009	0.01	0.001	0.01	0.001
Age	0.001	0.135	0.113	0.471	0.083	0.708	0.102	0.062	0.063	0.576	0.094	0.252
Country	0.083	0.001										

Figure 1. Variation in gut microbiome diversity and composition. **(A)** Alpha diversity estimated by Shannon, Observed ASVs and Faith's PD (Phylogenetic Diversity) between countries. **(B)** Alpha diversity estimated by Shannon, Observed ASVs and Faith's PD (Phylogenetic Diversity) between obese and non-obese. * $p < 0.05$, **** $p < 0.0001$ Alpha diversity metrics (Faith's PD, Observed ASVs and Shannon) are shown on the y-axis in different panels, while country or obese group are shown on the x-axis. **(C)** Beta diversity principal coordinate analysis based on weighted UniFrac distance between countries. **(D)** Beta diversity principal coordinate analysis based on weighted UniFrac distance between obese and non-obese. **(E)** Beta diversity principal coordinate analysis based on unweighted UniFrac distance between countries. **(F)** Beta diversity principal coordinate analysis based on unweighted UniFrac distance between obese and non-obese. Proportion of variance explained by each principal coordinate axis is denoted in the corresponding axis label. **(G)** Venn diagram of shared and unique ASVs between the five countries. Statistical significance adjusted for multiple comparisons using false discovery rate (FDR) correction are indicated: *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$ across countries and obese groups (Kruskal-Wallis test) for alpha diversity or by permutational multivariate analysis of variance (PERMANOVA) for beta diversity.




Figure 2. Receiver operating characteristic curves showing the classification accuracy of gut microbiota in a Random Forest model. Classification accuracy for estimating (A). All countries; (B) Obesity status, (C). Diabetes status; (D). Glucose status; (E). Hypertensive status; (F). Sex are presented. AUC= area under the curve




Figure 3. Correlations between alpha diversity and concentrations of the different types of fecal short chain fatty acids (SCFAs) among countries. Shannon index correlates positively with (A) total SCFA; (B) Acetate; (C) Butyrate; (D) Propionate; (E) Valerate.




Figure 4. Associations of gut microbiota ASVs with concentrations of short chain fatty acids (SCFAs). **(A)** Heatmap of Spearman's correlation between concentrations of SCFAs and top 30 differentially abundant ASVs (identified by ANCOM-BC) among countries. **(B)** Heatmap of Spearman's correlation between concentrations of SCFAs and differentially abundant ASVs (identified by ANCOM-BC) for obese. Correlations are identified by Spearman's rank correlation coefficient. Brick red squares indicate positive correlation, gray squares represent negative correlation and white squares are insignificant correlation.

Mapping from FDR adjusted p values are denoted as: *, ** and ***, corresponding to $p < 0.05$, <0.01 and <0.001 respectively.

