

Unbiased whole-genome deep sequencing of human and porcine stool samples reveals circulation of multiple groups of rotaviruses and a putative zoonotic infection

Running title: Rotavirus genomes and diversity in human and pigs

Authors: My VT Phan^a, Pham Hong Anh^b, Nguyen Van Cuong^b, Bas B. Oude Munnink^a, Lia van der Hoek^c, Phuc Tran My^b, Tue Ngo Tri^b, Juliet E. Bryant^{b,d}, Stephen Baker^{b,d,e}, Guy Thwaites^{b,d}, Mark Woolhouse^f, Paul Kellam^{g,h,i}, Maia A. Rabaa^{b,d}, and Matthew Cotten^{a,i}, on behalf of the VIZIONS Consortium

Author affiliations:

^aVirus Genomics, Wellcome Trust Sanger Institute, Hinxton, Cambridge, United Kingdom

^bOxford University Clinical Research Unit, Ho Chi Minh City, Vietnam

^cLaboratory of Experimental Virology, Academic Medical Center, University of Amsterdam, Amsterdam, the Netherlands

^dCentre for Tropical Medicine, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

^eLondon School of Tropical Medicine and Hygiene, London, United Kingdom

^fCentre for Immunity, Infection & Evolution, University of Edinburgh, Edinburgh, United Kingdom

^gKymab Inc

^hImperial College, London

ⁱCorresponding authors: Paul Kellam and Matthew Cotten, Virus genomics, Wellcome Trust Sanger Institute, Hinxton Cambridge CB10 1SA, United Kingdom. Email: mc13@sanger.ac.uk

Key words: rotavirus; deep sequencing; whole genomes; virus surveillance; zoonotic infection

Word count: 4,660 words

28

29 **Abstract**

30 Coordinated and synchronous virological surveillance for zoonotic viruses in both human
31 clinical cases and animal reservoirs provides an opportunity to identify interspecies virus
32 movement. Rotavirus is an important cause of viral gastroenteritis in humans and animals. We
33 have documented the rotavirus diversity within co-located humans and animals sampled from the
34 Mekong delta region of Vietnam using a primer-independent, agnostic, deep sequencing approach.
35 A total of 296 stool samples (146 from diarrhoeal human patients and 150 from pigs living in the
36 same geographical region) were directly sequenced, generating the genomic sequences of 60
37 human rotaviruses (all group A) and 31 porcine rotaviruses (13 group A, 7 group B, 6 group C and
38 5 group H). Phylogenetic analyses showed the co-circulation of multiple distinct rotavirus group A
39 (RVA) genotypes/strains, many of which were divergent from the strain components of licensed
40 RVA vaccines, as well as considerable virus diversity in pigs including full genomes of rotaviruses
41 in groups B, C and H, none of which have been previously reported in Vietnam. Furthermore the
42 detection of an atypical RVA genotype constellation (G4-P[6]-I1-R1-C1-M1-A8-N1-T7-E1-H1) in a
43 human patient and a pig from the same region provides some evidence for a zoonotic event.

44

45 **INTRODUCTION**

46 Rotavirus (RV) infections are the leading cause of acute gastroenteritis globally, with a
47 disproportionately greater morbidity and mortality in developing countries of Asia and sub-Saharan
48 Africa (1, 2). RV can infect humans and different animal species and is considered, in part, a
49 zoonotic disease in humans (3). Rotavirus zoonotic infections and transmissions, have been
50 shown with animal strains moving into humans via direct contact with animals or exposure to
51 environmental contamination (4–6), and present a challenge to infection control and management.
52 RV is a non-enveloped double-stranded RNA virus forming the single genus *Rotavirus* in the
53 Reoviridae family, with a 18.5 kb genome of 11 segments encoding six structural (VP1-4, VP6 and
54 VP7) and five or six non-structural proteins (NSP1-NSP5/6) (3, 7). RVs are classified into eight

established groups (A – H) and a new tentative group (I) based on the genetic and antigenic differences of VP6, with viruses of group A, B, C and H known to infect both humans and other animals (8, 9).

Within each rotavirus group, strains are distinguished and classified into G and P genotypes based on VP7 surface glycoprotein and VP4 spike protein, respectively (3). For rotavirus group A (RVA), at least 27 G and 37 P types have been detected in human and animals, with typical combinations of G-types (G1-G4, G9, G12) and P-types (P[4], P[6], P[8]) found in human infections globally, and different combinations of G- and P-types (G3-G5, G9, G11, P[6], P[7], P[13]) commonly found in pigs (6, 10–13). With considerable more sequence data available for RVA as compared to non-RVA strains, a genotype classification system based on 11 genomic segments is recommended for RVA, providing genomic insights into genotype constellations of both common and novel human and animal RVA strains (14). In this RVA genotyping system, Gx-P[x]-Ix-Rx-Cx-Mx-Ax-Nx-Tx-Ex-Hx represents the genotypes of VP7-VP4-VP6-VP1-VP2-VP3-NSP1-NSP2-NSP3-NSP4-NSP5 segments, with the most prevalent human strains belonging to constellations of Wa-like (I1-R1-C1-M1-A1-N1-T1-E1-H1, commonly associated with G1P[8], G3P[8], G4P[8], G9P[8], G12P[8]) and DS-1-like (I2-R2-C2-M2-A2-N2-T2-E2-H2, commonly combined with G2P[4]) (14, 15). Using this system, a common origin has been proposed between the human Wa-like and porcine strains (16). Such an elegant whole genome genotyping system has not yet been established for non-RVA groups.

Determining the sequence of the 18.5kb segmented genome for rotaviruses by standard methods can be biased and cumbersome, requiring an initial PCR step to identify and select primers specific for the RV genogroup and/or strain, with the 11-segment genome providing an additional complication for primer design. Such primer-based sequencing strategies can be further complicated by reassortment possibilities not predicted by the initial PCR typing, leading to sequencing failure of atypical and un-typeable RV strains whose frequency can vary by location, season and environment (10, 17–21). Next-generation sequencing has been recently employed for whole-genome sequencing of RVA with initial 11 PCR amplifications (22–24); however, a single

robust platform for whole-genome deep sequencing of multiple rotavirus genogroups without prior genotype information would be useful. Routine identification of circulating RVA can be performed using commercial enzyme immunoassay kits (based on inner capsid protein VP6) and RT-PCR diagnostic and genotyping assays (based on outer capsid proteins VP4 and VP7) (3, 25). In addition, specific, rapid and cost-effective assays are lacking for the detection of less common rotaviruses such as viruses in group B, C and H (RVB, RVC, RVH), hindering our understanding of molecular epidemiology of these viruses and challenging efforts of genomic sequencing, particularly in resource-limited countries (3, 25).

The rotavirus vaccines (Rotarix and RotaTeq) have been available since 2006 (26, 27), and offer a variable degree of protective immunity against human RVA infections. Reduced RVA vaccine efficacy has been observed in resource-limited countries in comparison to developed countries (28–31). The mechanism responsible for reduced vaccine efficacy in these settings is unclear, but may in part be due to local circulation of genetically and antigenically divergent RVA or zoonotic strains in developing countries (10). Similar to other segmented viruses (32), genetic reassortment has been observed in RVs yielding significant genetic diversity, including a number of cross-species reassortants (3, 4, 25). Hence, assessment of all 11 genome segments through full virus genome sequencing is essential for monitoring the overall RV genomic diversity, complex evolutionary dynamics and the emergence of novel and zoonotic reassortants that may compromise vaccine protection (11, 14, 15).

Vietnam is a low to middle income country located in Southeast Asia and is considered one of the global hot spots of emerging infectious diseases (33). Diarrhoea is the fourth most common cause of mortality in children <5 years of age, accounting for 12% of deaths in this age group in 2013 (34). Among all diarrhoeal pathogens, RVA is responsible for 44% - 67.4% of all childhood diarrhoea cases requiring hospitalisation (35–48). Contrary to the clinical and public health importance, vaccination against RVA is currently not part of the Extended Program on Immunisation for Vietnamese infants. Additionally, diagnosis for rotaviruses in diarrhoeal cases in humans and animals is not routinely performed and systematically genomic surveillance of

circulating human and animal rotaviruses is limited. This leads to relatively little data on the overall rotavirus prevalence and diversity in human and animal populations and their contribution to human infections and their potential to compromise RVA vaccine protection. Given the tropical climate of Vietnam prone to flooding, the frequent close human-animals living proximity and high prevalence of infectious diseases, we hypothesized that rotavirus zoonosis may occur in the region but is under-investigated and under-characterised. There is no report on the overall prevalence of other RVs (non-RVA) in both humans and animals in this region. To address this knowledge gap, we used focused sampling within human healthcare and animal farming populations, combined with high-throughput primer-independent direct genome sequencing from clinical materials (49) to document rotavirus diversity and transmission within and between humans and animals in a region of Vietnam.

MATERIALS AND METHODS

Study setting and design. Human and porcine faecal samples were collected from Dong Thap, a peri-urban province located in the south of Vietnam in the Mekong Delta region (see map, Supplementary Figure S1). The human subjects were diarrhoeal patients (N = 146) admitted to Dong Thap Provincial Hospital in the period from October 2012 to January 2014; a stool specimen was collected from each individual within 24 hours of hospital admission to avoid confounding by nosocomial infections. A total of 150 porcine faecal samples were randomly selected from a collection of porcine stool samples from pig farm baseline surveillance samples collected across the same province from January 2012 to April 2013. For 4 pigs in farms where no faecal specimens were obtained, a boot swab was collected (pig ID 12087_38, 14152_6, 14150_53 and 14250_12). All collected faecal samples were stored in aliquots at -80°C until further processing. Ethical approval for the study was obtained from the Oxford Tropical Research Ethics Committee (OxTREC Approval No. 15-12) (Oxford, United Kingdom), the institutional ethical review board of Dong Thap Provincial Hospital (DTPH) and the Sub-Department of Animal Health Dong Thap province (Dong Thap, Vietnam).

137 **Mapping of the patient residential and pig farm addresses.** The residential district centroid was
 138 recorded for enrolled human patients to maintain participant anonymity, while the exact
 139 geographical location was recorded for the pig farms using an eTrex Legend GPS device (Garmin,
 140 United Kingdom). The decimal degrees of latitude and longitude were entered in a confidential
 141 database and kept separate from patient metadata so that patient identities could not be revealed
 142 based on the residence locations. These addresses were then validated in Google Earth Pro
 143 (<https://www.google.com/earth/>) and finally visualised in QGIS v2.2.0 (<http://www.qgis.org/en/site/>)
 144 overlaid with province-specific geographic data.

145

146 **Sample preparation and nucleic acid extraction.** Total nucleic acid extraction was performed as
 147 previously described (49–51). Briefly, 110 μ L of a 50% stool suspension in PBS was centrifuged for
 148 10 minutes at 10,000 X g. Non-encapsidated DNA in the samples was degraded by addition of 20
 149 U TURBO DNase (Ambion). Virion-protected nucleic acid was subsequently extracted using the
 150 Boom method (52). Reverse transcription was performed using non-ribosomal random hexamers
 151 (53) that avoid transcription of rRNA, and second strand DNA synthesis was performed using 5U
 152 of Klenow fragment 3'-5' exo⁻ (New England Biolabs). Final purification of extracted nucleic acids
 153 was performed with phenol/chloroform and ethanol precipitation.

154

155 **Library preparation and sequencing.** Standard Illumina libraries were prepared for each sample.
 156 In short, nucleic acids in each sample were sheared to 400-500 nt in length, each sample's nucleic
 157 acid was separately indexed and samples were multiplexed at either 7 samples per MiSeq run or
 158 96 samples per HiSeq 2500 run, generating 2-3 million 149 nt (MiSeq) or 250 nt (HiSeq) paired-
 159 end reads per sample.

160

161 **De novo assembly and identification of viral genomes.** Raw sequencing reads were filtered to
 162 remove low quality reads (Phred score >35) and trimmed to remove residual sequencing adapters
 163 using QUASR (54). The reads were assembled into contigs using *de novo* assembly with SPAdes

(55) combined with sSpace (56). RV-encoding contigs and other mammalian virus contigs were identified with a modified SLIM algorithm (49) combined with ublast (57). Coverage was determined for all contigs harvested to filter any process contamination sequences in each run, followed by additional filtering for minimum contig size cutoff (300 nt). Partial but overlapping contigs were joined into full-length sequences using Sequencher (Gene Codes Corporation, USA), and any ambiguities were resolved by consulting the original short reads. Final quality control of genomes included a comparison of the sequences, open reading frames (ORFs) and the encoded proteins with reference sequences retrieved from GenBank.

Genotyping and phylogenetic reconstruction. Assembled RVA sequences were genotyped using the online genotyping tool, RotaC v2.0 (<http://rotac.regatools.be>) (58), according to the guidelines for precise RVA classification using all 11 genomic segments (11). The resulting RVA, RVB, RVC and RVH sequences were combined with additional full-length or nearly full-length sequences from previous Vietnamese studies (if available) and global representatives retrieved from GenBank. The complete genomes from the vaccine components of the monovalent vaccine Rotarix (59) and the pentavalent vaccine RotaTeq (60) were retrieved from GenBank for phylogenetic reconstructions of all 11 RVA segments. Sequences were aligned using MUSCLE v3.8.31 (61) and manually checked in AliView (62); aligned sequences were trimmed to complete ORFs for subsequent analyses. Evolutionary model testing was implemented in IQ-TREE v3.10 (63) using the Akaike Information Criterion (AIC) to determine the best-fit models of nucleotide substitution for all genomic segments. Maximum likelihood (ML) phylogenetic trees were then inferred in IQ-TREE v3.10 with 500 bootstrap replicates under the best-fit model of evolution according to AIC (Supplementary Table 1 summarised the models determined for all segments). Resulting trees were visualised and edited using FigTree v1.4.2 (<http://tree.bio.ed.ac.uk/software/figtree/>). Genetic distances (p-uncorrected) were estimated using Geneious v9.0.4 (Biomatters Ltd).

191 **Bayesian analysis for RVA NSP3 genotype T7.** Available RVA sequences of T7 type (NSP3
192 segment) retrieved from GenBank and new sequences obtained in this study were aligned using
193 MUSCLE v3.8.31 (61), manually checked in AliView (62), and trimmed to complete ORF. A
194 maximum likelihood phylogenetic tree was constructed under the GTR+ Γ_4 model of substitution in
195 IQ-TREE v3.10 (63). The molecular clock model was assessed in Path-O-Gen v1.4
196 (<http://tree.bio.ed.ac.uk/software/pathogen/>), assessing the linear regression between root-to-tip
197 divergence and the date of sampling (year; as data on day and month were not available for
198 GenBank sequences). A Bayesian Markov chain Monte Carlo (MCMC) approach was then
199 performed in BEAST v1.8.0 (64) with three independent chains, using relaxed lognormal molecular
200 clock under HKY85+ Γ_4 substitution model with a Bayesian SkyGid population process for 100
201 million generations chain with sampling performed every 10,000 runs. These triplicate runs were
202 then combined using LogCombiner v1.8.0 (available within the BEAST package) with a removal of
203 10% burn-in, and analysed in Tracer v1.6 (<http://tree.bio.ed.ac.uk/software/tracer/>) to ensure all
204 parameters had converged with effective sample size (ESS) values >200 and to estimate the mean
205 evolutionary rates across branches. Maximum clade credibility trees were annotated using
206 TreeAnnotator v1.8.0 (BEAST) and visualised in FigTree v1.4.2.

207

208 **Bayesian analysis for RVH VP6 gene.** A maximum likelihood phylogenetic tree was inferred for
209 all available RVH VP6 sequences retrieved from GenBank (N=39) and from this study (N=5) under
210 the GTR+ Γ_4 model of substitution in IQ-TREE (63). A time-scaled phylogeny was inferred for
211 sequences within the porcine lineage. Highly similar sequences were removed before running
212 Bayesian analyses (strains BR59, BR60, BR61, BR62, BR63, NC7_64_3, OK5_68_10). The
213 molecular clock model was assessed in Path-O-Gen, and a Bayesian MCMC approach was then
214 performed on the final set of sequences in BEAST (64), employing a relaxed lognormal molecular
215 clock under HKY8+ Γ_4 substitution model with a non-parametric Gaussian Markov Random Fields
216 (GMRF) Bayesian Skyride population with tip dates defined as year, month, day of strain
217 collection. Analyses were run in triplicate for 50 million generations with sampling performed every

5,000 generations. Triplicate runs were combined using LogCombiner with a removal of 10% burn-in, followed by analyses in Tracer, TreeAnnotator and FigTree as outlined in the aforementioned section.

GenBank accession numbers. All sequences generated in this study were deposited into GenBank under accession numbers KX362367 – KX363442. Illumina raw read sets are available at the European Nucleotide Archive under submission ERR471259 - ERR477293, ERR689707 - ERR767572, ERR775471 - ERR780002, ERR780013 - ERR780019, ERR956666, ERR956667, ERR962074, ERR1300950 - ERR1301100.

RESULTS

Overall diversity of rotaviruses in human and pigs

Sequencing of human enteric samples from acute diarrhoeal patients admitted to Dong Thap Provincial Hospital from 2012-2014 yielded 60 *de novo* assembled RVA genome sequences from 146 samples (41.1%). No other RV genogroups were found in these human stool samples (Table 1). The same methods applied to 150 porcine faecal samples collected within the same geographic region (Supplementary Figure 1) identified 31 rotaviruses from 4 different RV groups (A, B, C and H) in a total of 150 samples (20.7%). These *de novo* assembled sequences included 13 RVA (41.9%), 7 RVB (22.6%), 6 RVC (19.4%) and 5 RVH (16.1%) (Table 1). The length of each assembled sequence was determined and expressed as percentage length coverage (length of assembled sequence divided by expected full length of that segment) for the corresponding segment (Figure 1). In samples where 2 distinct contigs were assembled for a segment (e.g. mixed infections), only the longer assembled contig was reported in the heatmap of segment coverage for the purpose of clarity (Figure 1). The overall length coverage in human RVA sequences was higher than porcine RVA, RVB, RVC and RVH, possibly be due to differential viral load or sample quality.

Human and porcine RVA genotype constellations

The genotype constellations were determined for all RVA strains according to established

guidelines from the Rotavirus Classification Working Group (11, 58). Among the human RVA, the two most common constellations were G1-P[8]-I1-R1-C1-M1-A1-N1-T1-E1-H1 (N = 33; Wa-like constellation) and G2-P[4]-I2-R2-C2-M2-A2-N2-T2-E2-H2 (N = 12; DS-1-like constellation) (Figure 2). Reassorted RVA strains between Wa-like and DS-1-like constellations were found in 4 human diarrhoeal patients, including G1-P[8]-I2-R2-C2-M2-A2-N2-T2-E2-H2 (N = 3) and G2-P[8]-I2-R2-C2-M2-A2-N2-T2-E2-H2 (N = 1). Interestingly, the genotype constellation G4-P[6]-I1-R1-C1-M1-A8-N1-T7-E1-H1 was found in 1 human and 1 pig sample. The closely related genotype constellation G4-P[6]-I1-R1-C1-M1-A8-N1-T1-E1-H1, which differs only by the NSP3 segment, predominated among the porcine RVA mono-infections (N = 6) (Figure 2). Among all porcine RVA strains, the internal core gene cassette of R1-C1-M1-A8-N1-T1-E1-H1 (representing genotype of VP1-VP2-VP3-NSP1-NSP2-NSP3-NSP4-NSP5) was relatively conserved, with the exception of co-circulation of T1 and T7 genotypes in the NSP3 segment. The genotypes of capsid proteins (VP7-VP4-VP6) were more diverse in pigs, including G4-P[6]-I1 (6 strains combined with NSP3 T1 and 1 strain with NSP3 T7), G5-P[13]-I5 (N = 2), G9-P[23]-I5 (N = 1) and G11-P[23]-I5 (N = 1).

Mixed infections were identified in 9 samples, 7 in humans and 2 in pigs (Figure 2), with mixed infection being defined as the detection of two assembled but genetically distinct contigs in at least one segment with sufficient contig coverage to exclude potential process contamination among samples in the same run. The two homologous contig segments identified in mixed infections can have different or the same genotype; for example a mixed infection reported in an individual pig (sample ID 12070_4) contained 2 homologous VP7 segments, NSP1, NSP2, NSP3, NSP4 and NSP5 bearing the constellation of G1/G4-P[8]-I1-R1-C1-M1-A1/A8-N1/N1-T1/T1-E1/E1-H1/H1 (Figure 2 and Supplementary Figure S2). Another porcine sample was found with a mixture of G9/G11-P[13]/P[23]-I5/I5-R1/R1-C1-M1-A8/A8-N1/N1-T1/T1-E1/E1-H1 (sample ID 14150_53); however, it is important to note that this particular sample was a boot swab of faecal material in a cage-type pigsty, thus there is the possibility that the sample represents mixed environmental virus from more than one pig. Mixed human RVA infections typically contained genotype 1 and 2 (Wa-like and DS-1 like, respectively) viruses.

Phylogenetic diversity of local human and porcine RVA

Phylogenetic trees were inferred for each RVA segment (Figure 3A-B and Supplementary Figure S3A-B) from assembled sequences in this study along with full-length sequences from previous studies in Vietnam, reference sequences in GenBank and sequences from the RVA vaccine formulations (RotaRix and RotaTeq). The local sequences clustered primarily by genotype as expected (Figure 3A-B and Supplementary Figure S3A-B); for example, VP7 G1 sequences in this study clustered with other G1 sequences from other regions and our G2 sequences clustered with other G2 (Figure 3A). Sequences within the G4 genotype fell into 2 sub-lineages, with the human strain (16020_7) and porcine strains from this study clustering into one common sub-lineage (Figure 3A). The mixed infection in the pig (12070_4) described above comprised 2 distinct contigs for the VP7 segment (belonging to the G1 and G4 genotypes), with the G1 sequence clustering within the human G1 lineage and the G4 sequence falling into a lineage with other G4 porcine sequences from this study (Figure 3A). Similar observations were seen in the phylogenetic tree for VP4 sequences (Figure 3B) and other gene segments (Supplementary Figure S3A-B). It is also noteworthy that sequences from the vaccine strains (Rotarix and RotaTeq) were relatively distinct from the Vietnamese RVA sequences reported here, particularly for genotypes G5, G9, G11, P[6], P[13], P[23] of the two neutralizing antigens, VP7 and VP4 (Figure 3A-B). Comparison of the amino acid sequences of VP7 and VP4 of the local strains to the Rotarix and RotaTeq vaccine sequences indicated a number of amino acid differences observed across the length of the proteins and particularly in the antigenic epitopes of VP7 and VP4 (Supplementary Figure S7). Taken together, multiple RVA genotypes co-circulate in human and pigs in this location; many of these genotypes are genetically dissimilar to currently used vaccine components.

Putative zoonotic infection of human with a porcine-human RVA virus

An atypical RVA genotype constellation of G4-P[6]-I1-R1-C1-M1-A8-N1-T7-E1-H1 was found in both a human patient (16020_7) and a weaning pig (14250_9), whose geographical distance (between the residence and the farm) were approximately 35km apart (Figure 4A-B). The genotype constellation of the core gene cassette (R1-C1-M1-A8-N1-T7-E1-H1) was also identified

in 4 pig samples collected from 2 other farms (Figure 4A); the farms that raised these pigs are also about 35km away from the residential location of the 16020_7 case (Figure 4B). RVA strains with the aforementioned genome constellation have been identified in paediatric diarrhoeal patients in paediatric diarrhoeal patients in Hungary (65), Argentina (66), Paraguay (67), and Nicaragua (68), and were similarly thought to be of zoonotic origin (porcine-like). Given the diversity of geographic locations of reported zoonotic cases over a 15-year period, it is difficult to determine if this strain is sustained as a rare variant in human-to-human infections or has undergone multiple cross-species jumps from a porcine reservoir or an intermediate host.

Phylogenetically, all 11 segments of the human 16020_7 and porcine 14250_9 viruses belonged to lineages comprising porcine and/or porcine-origin human sequences (Figure 3A-B and Supplementary Figure S3A-B). Genetic distance suggested that the strain 16020_7 was most similar to porcine strains: TM-a (for VP1; 96% nt similarity), CMP45 (NSP3; 93%), and porcine-origin human strain 30378 (NSP2; 99%) (Figure 5). The remaining segments were most similar to porcine RVA strains obtained from this study, including the porcine 14150_53 (NSP1; 98.7% and NSP4; 99.1%) and 14225_44 (VP2; 99.2% and NSP5; 99.1%) strains. The capsid proteins of 16020_7 were most similar to the VP7 sequences of porcine samples 12129_48, 12129_49 and 12070_4 (G4 type; 97.2%), to the VP4 of pig 14226_39 (98.7%) and VP6 of pig 14226_42 (99.5%) (Figure 5). The porcine sample 14250_9, despite possessing the same genotype constellation as 16020_7, shared the highest nucleotide homology in only 2 internal genes, VP3 (97.5%) and NSP5 (99.1%), to the corresponding segments of strain 16020_7. Compared to the RVA vaccine Rotarix, 16020_7 was relatively dissimilar, sharing as low as 75.1% and 75.6% nt similarity for the VP4 and VP7 segments, respectively (Figure 5).

In this unusual genotype constellation reported, the NSP3 T7 type is a rare genotype that was first identified in a cow in Great Britain in 1973 (69), then in a bovine-like human strain (70), and later in pigs (71), porcine-bovine human reassortant (72) and porcine-like human strains (65–68, 73) in various geographical locations. The inferred evolutionary rate of RVA NSP3 sequences bearing T7 genotype was 1.3261×10^{-3} substitutions per site per year (95% highest posterior

density (HPD): 8.624×10^{-4} – 1.793×10^{-3}), which is slightly lower than the estimated evolutionary rates for RVA VP7 capsid gene of 1.66×10^{-3} and 1.87×10^{-3} substitutions/site/year for G12 and G9 genotypes, respectively (74). The time-stamped MCC tree also indicate an inter-connection among different host species indicating several host jump events particularly between pig and human hosts, suggesting that viral zoonotic chatter may occur more frequently than hitherto reported (Figure 4C).

Rotavirus group H (RVH), group B (RVB) and C (RVC)

RVH was identified in five Vietnamese pigs (3.33%; 5/150) at several time points and locations with no temporal or geographical associations (Figure 6), suggesting that these infections were sporadic and not linked to a single local outbreak. Furthermore, phylogenetic trees were inferred for all RVH segments to investigate the genetic diversity, comparing the RVH strains identified in this study with RVH sequences retrieved from GenBank (Figure 6 and Supplementary Figure S6). In general, RVH sequences typically clustered according to the host species, i.e. all porcine RVH sequences belonged to a lineage that is separated from human or cow RVH lineages. Within the porcine clade of the VP6 gene (Figure 6), sequences fell into two lineages: one lineage comprising sequences from USA and Japan, and the other lineage of Brazilian and Vietnamese sequences. The evolutionary rate was estimated to be 5.195×10^{-3} substitutions per site per year for sequences in the porcine lineage of RVH VP6 (95% HPD: 1.865×10^{-3} – 8.976×10^{-3}).

RVB was found in 7 pigs (4.67%; 7/150) and RVC was identified in 6 pigs (4%; 6/150). Phylogenetic trees of all segments of RVB and RVC showed that the local porcine sequences belonged to lineages comprising porcine sequences from other geographical locations for RVB (Figure 7 and Supplementary Figure S4) and RVC (Figure 7 and Supplementary Figure S5). In both RVB and RVC groups, the porcine lineages were relatively distant from lineages comprising of human sequences (Figure 7 and Supplementary Figure S4 and S5).

DISCUSSION

357 This study represents the first unbiased genome-wide surveillance, targeting
358 simultaneously multiple groups of rotaviruses infecting humans and animals in the same
359 geographical location. Prior to this study, there were only 3 subgenomic RVC sequences (<300nt)
360 and 9 complete or nearly complete RVA genomes reported from Vietnam in GenBank with no data
361 on RVB and RVH. Data from the current study document genomic sequences from 60 human RVA
362 and 31 porcine RV (group A, B, C and H), providing the largest available collection of genome
363 sequences from human and pigs from a single location in general and from Vietnam in particular.
364 This is also the first report and the first genome characterisation of RVB, RVC and RVH from
365 Vietnamese pigs.

366 Among the RVA, we identified a human (sample 16020_7) and a porcine sample (sample
367 14250_9) with atypical RVA genotype constellation G4-P[6]-I1-R1-C1-M1-A8-N1-T7-E1-H1,
368 detected for the first time in Vietnam and in Asia. This variant may have originated from a direct
369 zoonotic transmission or from reassortment event(s) involving porcine and porcine-origin human
370 strains. The human RVA strain (16020_7) was identified in a sample from a 54 year-old patient,
371 admitted to the hospital due to acute diarrhoea. Rotavirus was the sole enteric pathogen identified
372 from the stool sample and no other common viral and bacterial diarrhoeal pathogens were found
373 by diagnostic testing (39, 75) for norovirus, astrovirus, sapovirus, adenovirus F and aichi virus,
374 *Shigella spp.*, *Salmonella spp.* and *Campylobacter spp.* (data not shown). Although adults can be
375 infected with RVA, such infections in immuno-competent individuals are typically asymptomatic,
376 self-limiting or cause mild disease (76). The rotavirus infection in this particular case required
377 hospitalisation suggesting a moderate-severe end of the clinical spectrum of diarrhoeal disease.
378 Although further studies are required to determine their significance and relevance in human and
379 animal diseases, it is tempting to suggest that this atypical strain may be the cause of the
380 moderate-severe diarrhoeal disease. The close proximity between humans and pigs and common
381 use of river water (Mekong Delta River, Figure 4B) for daily activities and farming might present an
382 enhanced risk of transmitting water-borne infectious pathogens, in this case providing a plausible
383 zoonotic route of atypical rotavirus transmission.

Compared to RVA, human and animal infections with RVB, RVC and RVH are not well understood and the detection rates for these groups of viruses are relatively low (77). This is probably because the majority of rotavirus investigations have been focused on RVA given its clinical and public health relevance and the large genetic distance among these groups of rotaviruses as compared to RVA, which would likely be missed by commonly used diagnostic assays. Recently, there have been increasing numbers of reports on rotavirus groups B, C and H in animals (78–86) and humans (87–89), which possibly reflect improved molecular methods to detect these viruses rather than an actual increase in their prevalence. In Vietnam, the frequencies and relative role in human and animal disease of RVB, RVC and RVH viruses are not yet known. Although no human RVB, RVC and RVH were found in this study, the zoonotic potential of these rotaviruses groups cannot be ignored.

Our documentation of local RV genetic diversity and the potential of RV for zoonosis are highly relevant for the introduction of RVA vaccines into the region. The VP7 surface glycoprotein and VP4 spike protein are considered to be major antigenic targets for the protective host immune responses induced by RVA vaccines (73, 90, 91). Importantly, several amino acid changes were observed in defined antigenic sites of VP7 and VP4 proteins encoded by the local Vietnamese strains compared to available vaccines. Although a genetic comparison alone cannot predict if circulating Vietnamese viruses will be recognized and blocked or if the changes will allow infection in vaccinated subjects, the number of differences in antigenic regions suggests a compromised protective function of the vaccines in Vietnamese children. These data should be considered in light of the clinical observations of reduced vaccine efficacy (28–30). Future vaccine efforts may benefit from this increased knowledge of rotavirus diversity provided by this work.

Our study does have limitations. Firstly, the sample size of 146 human and 150 porcine samples is relatively small. Despite this size, we were able to identify a potential zoonotic infection. This provides a baseline frequency for zoonotic infections and suggest that this may be occurring at a higher rate than previously considered. Secondly, the disease status of the sampled pigs was not well defined and there was no follow-up beyond the sampling time point so the clinical

spectrum of diarrhoeal disease (eg. mild, moderate or severe) in the pigs is unknown. However, the primary objective of this study was characterization of rotavirus pathogenesis or causation of the diarrhoeal disease. The presence of rotavirus material at sufficiently high titres to allow full genome sequencing is consistent with these animals being a common source of the virus for movement to other species. Our findings indicate that porcine faecal material is a source of novel and possibly zoonotic viruses.

It is likely that with the ubiquity and falling costs of sequencing, the unbiased virus sequencing described here will become an important component of infectious disease surveillance and rapid responses to outbreaks (92). The ideal sampling rate, sample numbers and geographical relationship between humans and animals for genetic surveillance are still being defined but the current work provides a good starting point for future efforts. Even within the relatively modest sample set of 296 human and animal enteric samples, a considerable RV genetic diversity was observed including a potential zoonosis. The integration of targeted sampling, sequencing and phylogeography or phylogenetics in different places in the world, perhaps informed by other risk mapping (33, 93) has the ability to inform surveillance and to monitor zoonotic pathogens in human and animals.

Funding

This work was supported by the Wellcome Trust of the UK through the VIZIONS strategic award (WT/093724), and the British Council of the UK through the Researchers Link Travel Award (MVTP, grant application number 127624851).

Competing Interests

The opinions expressed by the authors contributing to this paper do not necessarily reflect the opinions of the Wellcome Trust, the British Council or the institutions with which the authors are affiliated or the grantee bodies. The authors declare no competing interests.

Acknowledgements

We are grateful to all the patients and their families and the farmers in Dong Thap (Vietnam) that have participated in the study. We thank the Illumina C team at the Wellcome Trust Sanger Institute (Hinxton, Cambridge, United Kingdom) for their help in deep sequencing and Simon Watson and Pinky Langat (Wellcome Trust Sanger Institute) for technical assistance on running BEAST analyses. The complete list of the VIZIONS Consortium can be found at (94).

Tables and Figures

Table 1. Number of samples tested and number of samples yielded up to full or nearly full RV genomic sequences.

Host	RVA	RVB	RVC	RVH	Total samples tested
Human	60 (41.1%)	0	0	0	146
Pig	13 (8.7%)	7 (4.7%)	6 (4%)	5 (3.3%)	150

Host indicates human or pig host from which the faecal samples were collected. Percentages of full or nearly full RV genomes were given in brackets. RVA: rotavirus genogroup A; RVB: group B; group C; RVH: group H.

were shown in the first panel, with each column representing a sample and each row showing each RV segment. Similarly, porcine rotavirus samples were shown, each panel representing RVA, RVB, RVC and RVH sequences with the segment names given vertically and sample IDs horizontally.

Figure 2

Genotype constellation												Host count	
	VP7	VP4	VP6	VP1	VP2	VP3	NSP1	NSP2	NSP3	NSP4	NSP5	Human	Pig
Mono-infection	G1	P[8]	I1	R1	C1	M1	A1	N1	T1	E1	H1	33	0
	G2	P[4]	I2	R2	C2	M2	A2	N2	T2	E2	H2	12	0
	G3	P[8]	I1	R1	C1	M1	A1	N1	T1	E1	H1	2	0
	G4	P[6]	I1	R1	C1	M1	A8	N1	T7	E1	H1	1	1
	G4	P[6]	I1	R1	C1	M1	A8	N1	T1	E1	H1	0	6
	G5	P[13]	I5	R1	C1	M1	A8	N1	T7	E1	H1	0	1
	G11	P[13]	I5	R1	C1	M1	A8	N1	T7	E1	H1	0	1
	G9	P[23]	I5	R1	C1	M1	A8	N1	T1	E1	H1	0	1
	G5	P[13]	I5	R1	C1	M1	A8	N1	T7	x	H1	0	1
	G1	P[8]	I2	R2	C2	M2	A2	N2	T2	E2	H2	3	0
	G2	P[8]	I2	R2	C2	M2	A2	N2	T2	E2	H2	1	0
	G1	P[8]	x	R1	C1	M1	A1	N1	T1	E1	H1	1	0
Mixed infection	G1/G2	P[8]/P[4]	I1/I2	R1	C1/C2	M1/M2	A1/A2	N1/N2	T1/T2	E1/E2	H1/H2	1	0
	G1/G2	P[8]/P[4]	I1/I2	R1/R2	C1	M1/M2	A1/A2	N1/N2	T1/T2	E1/E2	H1	1	0
	G1/G2	P[8]/P[4]	I1/I2	R1/R2	C1/C2	M1/M2	A1/A2	N1/N2	T1/T2	E2	H2	1	0
	G1/G2	P[8]/P[4]	I1/I2	R1/R2	C1/C2	M1/M2	A1/A2	N1/N2	T1/T2	E1/E2	H1	1	0
	G1/G2	P[8]/P[4]	I1/I2	R1/R2	C1/C2	M1/M2	A1/A2	N1/N2	T1/T2	E1/E2	H1/H2	1	0
	G1/G2	P[8]/P[4]	I1/I2	R2	C1/C2	M2	A1/A2	N1/N2	T1/T2	E1/E2	H1/H2	1	0
	G1	P[8]	I1	R1	C1	M1	A1	N1/N1	T1	E1	H1	1	0
	G1/G4	P[8]	I1	R1	C1	M1	A1/A8	N1/N1	T1/T1	E1/E1	H1/H1	0	1
	G9/G11	P[13]/P[23]	I5/I5	R1/R1	C1	M1	A8/A8	N1/N1	T1/T7	E1/E1	H1	0	1
	Wa-like genotype (G1/G3/G4/G9-P[8]-I1-R1-C1-M1-A1-N1-T1-E1-H1)									G5			
	DS-1-like genotype (G2/G8-P[4]-I2-R2-C2-M2-A2-N2-T2-E2-H2)									G11			
	G4-P[6]									I5			
	A8-T7									P[13]			
	G9-P[23]									x	Sequence not determined		

Figure 2. Genotype constellations of assembled RVA genomes.

The genotypes of all assembled sequences were determined according to the guidelines of Rotavirus Classification Working Group, representing the genotype of VP7-VP4-VP6-VP1-VP2-VP3-NSP1-NSP2-NSP3-NSP4-NSP5. For mono-infection, each row represents one genotype constellation with the colour block used to illustrate different genotype patterns, such as common human types of Wa-like (orange) and DS-1-like (purple), and other less common genotypes shown in other colour blocks as indicated. The number of each of the genotype constellation identified in samples from human and pigs were given in the column Host count for Human and Pigs, respectively. For mixed infection where 2 distinct contigs were assembled for at least one segment, the genotypes of both contigs were given and indicated the host where strains were identified.

Figure 3




Figure 3. Maximum-likelihood phylogenetic trees inferred from the assembled nucleotide sequences for RVA VP7 and VP4 genes. (A) Maximum-likelihood tree of VP7 gene showed genetic relationships between sequences from this study and additional sequences of corresponding segments from GenBank. Branches and strain names were coloured according to

genotype and host, respectively. Tree is mid-point rooted for the purpose of clarity and bootstrap values of $\geq 75\%$ are shown for major nodes only. All horizontal branch lengths are drawn to the scale of nucleotide substitutions per site. **(B)** Maximum-likelihood tree of VP4 gene showed genetic relationships between sequences from this study and additional sequences of corresponding segments from GenBank. The pattern of tree visualisation is consistent with VP7 tree, see description of Figure 3A for more information.

Figure 4




Figure 4. The analysis of RVA zoonotic strain G4P[6]T7 in the study.

(A) The genotype constellation of the case in investigation 16020_7 and other porcine RVA strains with the NSP3 T7 genotype. Details on dates of collection and age of host are given. The colour coded in the host column is consistent for colour illustration of corresponding case in the map (panel B) in this figure. **(B)** The geographical location of the human case's residency and pig farms that raised the pigs infected with RVA NSP3 T7 strains overlaid on the total sampling area (as shown in Supplementary Figure S1). The colouring of the human case and pig farms is consistent

501 with colour code presented in the column “Host” in panel of this figure. The red star indicates the
502 Dong Thap Provincial Hospital where diarrhoeal patients were admitted. The map scale bar is
503 shown in the units of geometric km. See Supplementary Figure S1 for more information. **(C)** The
504 time-resolved phylogenetic tree of RVA NSP3 T7 genotype sequences comparing local versus
505 global sequences. Reference sequences were retrieved from GenBank (N=69, excluded the
506 duplicate sequence for TM-a strain (JX290174) and BP1901 (KF835960) which is 15 aa shorter
507 than the complete ORF). Strains were coloured according to the host species from which the strain
508 was identified. Sequences identified from this study were highlighted in blue (human case
509 16020_7) or in yellow (porcine sequences). The first T7 sequence identified (in a cow in 1973) was
510 indicated in purple. Posterior probabilities of internal nodes with values ≥ 0.75 are shown and the
511 scale axis indicates time in year of strain identification.

512

513 **Figure 5**

Strain ID	Country	VP7	VP4	VP6	VP1	VP2	VP3	NSP1	NSP2	NSP3	NSP4	NSP5
16020_7	VNM	G4	P[6]	I1	R1	C1	M1	A8	N1	T7	E1	H1
14250_9	VNM	G4 96.9%	P[6] 92.5%	I1 97.3%	R1 85.4%	C1 97.9%	M1 97.5%	A8 81.6%	N1 97.5%	T7 87.5%	E1 96.5%	H1 99.1%
14150_53	VNM	G11/G9 75.7%/72%	P[13]/P[23] 68.2%/72.2%	I5/I5 83.1%/83.5%	R1/R1 87.3%/85.7%	C1 97.2%	M1 86.5%	A8/A8 98.7%/81.9%	N1/N1 86.5%/93.4%	T1/T7 83.1%/91.6%	E1/E1 97.4%/99.1%	H1 94.4%
14150_54	VNM	G11 72.0%	P[13] 68.4%	I5 83.2%	R1 85.6%	C1 97.3%	M1 86.4%	A8 98.1%	N1 86.4%	T7 91.6%	E1 97.5%	H1 95.3%
14225_45	VNM	G5 73.8%	P[13] 67.7%	I5 82.5%	R1 85.6%	C1 98.8%	M1 98.3%	A8 95.7%	N1 97.1%	T7 92.1%	E1 88.7%	H1 98.6%
14225_46	VNM	G5 73.9%	P[13] 67.6%	I5 82.3%	R1 85.5%	C1 98.7%	M1 96.0%	A8 95.7%	N1 94.7%	T7 91.9%	X X	H1 98.4%
14225_44	VNM	G4 96.7%	P[6] 98.3%	I1 99.4%	R1 87.3%	C1 99.2%	M1 86.6%	A8 79.2%	N1 93.8%	T1 83.3%	E1 90.9%	H1 99.1%
14226_42	VNM	G4 96.7%	P[6] 97.9%	I1 99.5%	R1 87.3%	C1 99.1%	M1 86.2%	A8 82.1%	N1 95.5%	T1 83.3%	E1 90.9%	H1 99.1%
14226_39	VNM	G4 97.1%	P[6] 98.7%	I1 83.5%	R1 85.8%	C1 90.0%	M1 87.1%	A8 82.5%	N1 95.5%	T1 82.5%	E1 90.1%	H1 98.1%
14249_23	VNM	G4 97.0%	P[6] 92.4%	I1 98.1%	R1 87.7%	C1 86.7%	M1 86.5%	A8 95.9%	N1 96.3%	T1 82.3%	E1 96.6%	H1 96.0%
12070_4	VNM	G4/G1 97.2%/75.6%	P[8] 75.4%	I1 99.0%	R1 92.5%	C1 87.5%	M1 87.6%	A1/A8 76.9%/80.9%	N1/N1 97.7%/87.6%	T1/T1 84%/82.6%	E1/E1 91.4%/90.3%	H1/H1 98.3%/94%
12129_48	VNM	G4 97.2%	P[6] 91.3%	I1 98.1%	R1 86.8%	C1 86.6%	M1 86.1%	A8 95.9%	N1 96.6%	T1 83.1%	E1 95.6%	H1 96.1%
12129_49	VNM	G4 97.2%	P[6] 92.4%	I1 98.1%	R1 66.6%	C1 82.0%	M1 86.4%	A8 95.9%	N1 96.5%	T1 83.1%	E1 96.3%	H1 95.9%
30378 ^s	VNM	G26 74.8%	P[19] 76.2%	I5 83.4%	R1 87.3%	C1 94.1%	M1 96.9%	A8 82.2%	N1 98.6%	T1 83.8%	E1 96.4%	H1 98.8%
OL ^s	NCA	G4 86.0%	P[6] 86.2%	I1 89.8%	R1 86.1%	C1 87.0%	M1 85.3%	A8 82.4%	N1 92.7%	T7 91.6%	E1 88.0%	H1 95.7%
E931 ^s	CHN	G4 97.1%	P[6] 92.7%	I1 97.7%	R1 94.6%	C1 91.0%	M1 85.1%	A8 82.2%	N1 93.1%	T1 84.1%	E1 96.8%	H1 95.4%
R946 ^s	CHN	G3 73.8%	P[6] 94.5%	I1 96.5%	R1 95.4%	C1 90.8%	M1 84.8%	A1 78.0%	N1 88.9%	T1 84.2%	E1 92.8%	H1 94.7%
CMP45/08	THA	G9 75.5%	P[23] 71.2%	I5 83.1%	R1 86.9%	C1 92.5%	M1 88.1%	A8 92.4%	N1 93.5%	T7 93.1%	E1 89.1%	H1 93.3%
TM-a	CHN	G9 75.5%	P[23] 71.1%	I5 82.9%	R1 95.9%	C1 90.5%	M1 86.6%	A8 80.0%	N1 89.0%	T7 88.2%	E1 91.8%	H1 94.9%
Wa	USA	G1 74.8%	P[8] 75.4%	I1 92.9%	R1 88.8%	C1 88.1%	M1 90.3%	A1 76.5%	N1 88.0%	T1 83.1%	E1 93.9%	H1 93.3%
Rotarix	USA	G1 75.6%	P[8] 75.1%	I1 92.5%	R1 88.6%	C1 88.1%	M1 90.2%	A1 76.2%	N1 88.1%	T1 83.1%	E1 93.3%	H1 92.8%

514

515 **Figure 5. Genotype constellations of RVA 16020_7 compared with representative human**
516 **and animal RVA of known genotypes.**

517 Segments are bold in red and shaded in green to indicate the segments with highest nucleotide
518 sequence similarities to that of strain 16020_7. The strain was coloured according the host from
519 which strain was identified, blue indicates human host, pink for pigs and green for vaccine

component. All porcine samples from this study were shaded in light blue and the human case of interest (16020_7) was shaded in grey. Human strains were previously shown to have porcine origin. Country of isolation abbreviation, VNM: Vietnam; NCA: Nicaragua; CHN: China; THA: Thailand; USA: United States of America.

Figure 6






Figure 6. The analysis of RVH VP6 gene segment.

(A) Maximum-likelihood phylogenetic tree of RVH VP6 sequences. The RVH VP6 sequences in this study (N=5) were compared with available full-length RVH VP6 sequences retrieved from GenBank (N=39). Tree is mid-point rooted for the purpose of clarity and only bootstrap values of $\geq 75\%$ are shown. All horizontal branch lengths are drawn to the scale of nucleotide substitutions per site in the tree. Strains were colour coded according to the host associated with the strain, the

532 country where the strains were identified and the year of strain detection. **(B)** The geographical
533 locations of pig farms that raised the pigs infected with RVH identified in this study. The farms were
534 illustrated as red triangles with strain ID given, overlaid on the overall sampling area as shown in
535 Supplementary Figure S1. The red star indicates the Dong Thap Provincial Hospital. The map
536 scale bar is shown in the units of geometric km. Refer to Supplementary Figure S1 for more
537 information on the background and provincial features colouring. **(C)** Time-resolved phylogenetic
538 tree of porcine RVH VP6 sequences comparing local versus global sequences. Strains were
539 coloured by the country where strains were identified. Porcine sequences identified from this study
540 were highlighted in red, with strain ID given to link with geographical locations on map in panel C of
541 this figure. Strains from Brazil were coloured in dark blue; orange indicates the Japanese porcine
542 strain, and light green refers to the porcine strain from the US. Posterior probabilities of internal
543 nodes with values ≥ 0.75 are shown and the scale axis indicates time in year of strain identification.
544

545 **Figure 7**

Host



546

Figure 7. Maximum-likelihood phylogenetic trees inferred from the assembled nucleotide sequences for VP6 gene of RVB and RVC. Maximum-likelihood trees of RVB and RVC VP6 segments showed genetic relationships between assembled sequences from this study and full-length reference sequences of corresponding segments retrieved from GenBank. Trees are mid-point rooted for the purpose of clarity and only bootstrap values of $\geq 75\%$ are shown. Scale bars are in the unit of nucleotide substitutions per site. Strains were coloured according to the host species that the sequences were identified from.

REFERENCES

1. **Tate JE, Burton AH, Boschi-Pinto C, Steele a D, Duque J, Parashar UD.** 2012. 2008 Estimate of Worldwide Rotavirus-Associated Mortality in Children Younger Than 5 Years Before the Introduction of Universal Rotavirus Vaccination Programmes: a Systematic Review and Meta-Analysis. *Lancet Infect Dis* **12**:136–41.
2. **Tate JE, Burton AH, Boschi-Pinto C, Parashar UD.** 2016. Global, Regional, and National Estimates of Rotavirus Mortality in Children <5 Years of Age, 2000–2013. *Clin Infect Dis* **62**:S96–S105.
3. **Estes MK, Kapikian AZ, Knipe DM, Howley PM.** 2007. Rotaviruses, p. 1917–1974. *In* Knipe, DM, Howley, PM (eds.), *Fields virology* 5th ed. Lippincott Williams & Wilkins, Philadelphia, PA.
4. **Cook N, Bridger J, Kendall K, Gomara MI, El-Attar L, Gray J.** 2004. The zoonotic potential of rotavirus. *J Infect* **48**:289–302.
5. **Martella V, Banyai K, Matthijssens J, Buonavoglia C, Ciarlet M, Bányai K.** 2010. Zoonotic aspects of rotaviruses. *Vet Microbiol* **140**:246–255.
6. **Ghosh S, Kobayashi N.** 2014. Exotic rotaviruses in animals and rotaviruses in exotic animals. *Virus Dis* **25**:158–72.
7. **Ramig RF, Ciarlet M, Mertens PPC, Dermody TS.** 2005. Rotavirus, p. 484–496. *In* Fauquet, CM, Mayo, MA, Maniloff, J, Desselberger, U, Ball, LA (eds.), *Virus taxonomy: eight report of the international committee on taxonomy of viruses*. Elsevier Academic Press, Amsterdam, Holland.
8. **Matthijssens J, Otto PH, Ciarlet M, Desselberger U, Van Ranst M, Johne R.** 2012. VP6-sequence-based cutoff values as a criterion for rotavirus species demarcation. *Arch Virol* **157**:1177–82.
9. **Mihalov-kovács E, Gellért Á, Marton S, Farkas SL, Fehér E, Oldal M, Jakab F, Martella V, Banyai K.** 2015. Candidate New Rotavirus Species in Sheltered Dogs, Hungary. *Emerg Infect Dis* **21**:660–663.
10. **Santos N, Hoshino Y.** 2005. Global distribution of rotavirus serotypes/genotypes and its implication for the development and implementation of an effective rotavirus vaccine. *Rev Med Virol* **15**:29–56.
11. **Matthijssens J, Ciarlet M, McDonald SM, Attoui H, Bányai K, Brister JR, Buesa J, Esona MD, Estes MK, Gentsch JR, Iturriza-Gómara M, Johne R, Kirkwood CD, Martella V, Mertens PPC, Nakagomi O, Parreño V, Rahman M, Ruggeri FM, Saif LJ, Santos N,**

- 588 **Steyer A, Taniguchi K, Patton JT, Desselberger U, Van Ranst M.** 2011. Uniformity of
589 rotavirus strain nomenclature proposed by the Rotavirus Classification Working Group
590 (RCWG). *Arch Virol* **156**:1397–413.
- 591 12. **Trojan E, Sachsenröder J, Twardziok S, Reetz J, Otto PH, Johne R.** 2013. Identification
592 of an avian group A rotavirus containing a novel VP4 gene with a close relationship to those
593 of mammalian rotaviruses. *J Gen Virol* **94**:136–42.
- 594 13. **Dóro R, Farkas SL, Martella V, Bányai K.** 2015. Zoonotic transmission of rotavirus:
595 surveillance and control. *Expert Rev Anti Infect Ther* **13**:1337–1350.
- 596 14. **Matthijnssens J, Ciarlet M, Rahman M, Attoui H, Bányai K, Estes MK, Gentsch JR,
597 Iturriza-Gómara M, Kirkwood CD, Martella V, Mertens PPC, Nakagomi O, Patton JT,
598 Ruggeri FM, Saif LJ, Santos N, Steyer A, Taniguchi K, Desselberger U, Van Ranst M,
599 Bányai K, Iturriza-Gómara M.** 2008. Recommendations for the classification of group A
600 rotaviruses using all 11 genomic RNA segments. *Arch Virol* **153**:1621–1629.
- 601 15. **Matthijnssens J, Van Ranst M.** 2012. Genotype constellation and evolution of group A
602 rotaviruses infecting humans. *Curr Opin Virol* **2**:426–33.
- 603 16. **Matthijnssens J, Ciarlet M, Heiman E, Arijs I, Delbeke T, McDonald SM, Palombo EA,
604 Iturriza-Gómara M, Maes P, Patton JT, Rahman M, Van Ranst M, Iturriza-Gómara M,
605 Van Ranst M.** 2008. Full genome-based classification of rotaviruses reveals a common
606 origin between human Wa-Like and porcine rotavirus strains and human DS-1-like and
607 bovine rotavirus strains. *J Virol* **82**:3204–19.
- 608 17. **Cook SM, Glass RI, Lebaron CW, Ho M.** 1990. Global seasonality of rotavirus infections.
609 *Bull World Health Organ* **66**:171–177.
- 610 18. **Pitzer VE, Viboud C, Simonsen L, Steiner C, Panozzo C a, Alonso WJ, Miller M a,
611 Glass RI, Glasser JW, Parashar UD, Grenfell BT.** 2009. Demographic variability,
612 vaccination, and the spatiotemporal dynamics of rotavirus epidemics. *Science* **325**:290–4.
- 613 19. **Pitzer VE, Viboud C, Lopman B a, Patel MM, Parashar UD, Grenfell BT.** 2011. Influence
614 of birth rates and transmission rates on the global seasonality of rotavirus incidence. *J R*
615 *Soc Interface* **8**:1584–93.
- 616 20. **Atchison C, Iturriza-Gómara M, Tam C, Lopman B.** 2010. Spatiotemporal dynamics of
617 rotavirus disease in Europe: can climate or demographic variability explain the patterns
618 observed. *Pediatr Infect Dis J*, 2010/01/23 ed. **29**:566–568.
- 619 21. **Gentsch JR, Laird AR, Bielfelt B, Griffin DD, Bányai K, Ramachandran M, Jain V,
620 Cunliffe N a, Nakagomi O, Kirkwood CD, Fischer TK, Parashar UD, Bresee JS, Jiang
621 B, Glass RI.** 2005. Serotype diversity and reassortment between human and animal
622 rotavirus strains: implications for rotavirus vaccine programs. *J Infect Dis* **192**:S146–59.
- 623 22. **Magagula NB, Esona MD, Nyaga MM, Stucker KM, Halpin RA, Stockwell TB, Seheri
624 ML, Steele AD, Wentworth DE, Mphahlele MJ.** 2015. Whole Genome Analyses of G1P[8]
625 Rotavirus Strains From Vaccinated and Non-Vaccinated South African Children Presenting
626 With Diarrhea. *J Med Virol* **87**:79–101.
- 627 23. **Jere KC, Mlera L, Neill HGO, Potgieter AC, Page NA, Seheri ML, Dijk AA Van.** 2011.
628 Whole Genome Analyses of African G2 , G8 , G9 , and G12 Rotavirus Strains Using
629 Sequence-Independent Amplification and 454-Pyrosequencing. *J Med Virol* **83**:2018–2042.
- 630 24. **Nyaga MM, Jere KC, Esona MD, Seheri ML, Stucker KM, Halpin R a, Akopov A,
631 Stockwell TB, Peenze I, Diop A, Ndiaye K, Boula A, Maphalala G, Berejena C, Mwenda
632 JM, Steele a D, Wentworth DE, Mphahlele MJ.** 2015. Whole genome detection of
633 rotavirus mixed infections in human, porcine and bovine samples co-infected with various
634 rotavirus strains collected from sub-Saharan Africa. *Infect Genet Evol* **31**:321–334.
- 635 25. **Desselberger U.** 2014. Rotaviruses. *Virus Res* **190C**:75–96.
- 636 26. **Ruiz-Palacios GM, Perez-Schael I, Velazquez FR, Abate H, Breuer T, Clemens SC,**

Cheuvart B, Espinoza F, Gillard P, Innis BL, Cervantes Y, Linhares AC, Lopez P, Macias-Parra M, Ortega-Barria E, Richardson V, Rivera-Medina DM, Rivera L, Salinas B, Pavia-Ruz N, Salmeron J, Ruttimann R, Tinoco JC, Rubio P, Nunez E, Guerrero ML, Yarzabal JP, Damaso S, Tornieporth N, Saez-Llorens X, Vergara RF, Vesikari T, Bouckennooghe A, Clemens R, De Vos B, O’Ryan M. 2006. Safety and efficacy of an attenuated vaccine against severe rotavirus gastroenteritis. *N Engl J Med* **354**:11–22.

27. **Vesikari T, Matson DO, Dennehy P, Van Damme P, Santosham M, Rodriguez Z, Dallas MJ, Heyse JF, Goveia MG, Black SB, Shinefield HR, Christie CDC, Ylitalo S, Itzler RF, Coia ML, Onorato MT, Adeyi B a, Marshall GS, Gothefors L, Campens D, Karvonen A, Watt JP, O’Brien KL, DiNubile MJ, Clark HF, Boslego JW, Offit PA, Heaton PM.** 2006. Safety and efficacy of a pentavalent human-bovine (WC3) reassortant rotavirus vaccine. *N Engl J Med* **354**:23–33.

28. **Breiman RF, Zaman K, Armah G, Sow SO, Duc D, Victor JC, Hille D, Ciarlet M, Neuzil KM, Anh DD.** 2012. Analyses of health outcomes from the 5 sites participating in the Africa and Asia clinical efficacy trials of the oral pentavalent rotavirus vaccine. *Vaccine* **30 Suppl 1**:A24–9.

29. **Zaman K, Dang DA, Victor JC, Shin S, Yunus M, Dallas MJ, Podder G, Vu DT, Le TPM, Luby SP, Le HT, Coia ML, Lewis K, Rivers SB, Sack DA, Schodel F, Steele AD, Neuzil KM, Ciarlet M, Schödel F.** 2010. Efficacy of pentavalent rotavirus vaccine against severe rotavirus gastroenteritis in infants in developing countries in Asia: a randomised, double-blind, placebo-controlled trial. *Lancet* **376**:615–623.

30. **Glass RI, Parashar U, Patel M, Gentsch J, Jiang B.** 2014. Rotavirus vaccines: successes and challenges. *J Infect* **68 Suppl 1**:S9–18.

31. **Armah GE, Sow SO, Breiman RF, Dallas MJ, Tapia MD, Feikin DR, Binka FN, Steele a. D, Laserson KF, Ansah N a., Levine MM, Lewis K, Coia ML, Attah-Poku M, Ojwando J, Rivers SB, Victor JC, Nyambane G, Hodgson A, Schödel F, Ciarlet M, Neuzil KM.** 2010. Efficacy of pentavalent rotavirus vaccine against severe rotavirus gastroenteritis in infants in developing countries in sub-Saharan Africa: A randomised, double-blind, placebo-controlled trial. *Lancet* **376**:606–614.

32. **Nomikou K, Hughes J, Wash R, Kellam P, Breard E, Zientara S, Palmarini M, Biek R, Mertens P.** 2015. Widespread Reassortment Shapes the Evolution and Epidemiology of Bluetongue Virus following European Invasion. *PLOS Pathog* **11**:e1005056.

33. **Hay SI, Battle KE, Pigott DM, Smith DL, Moyes CL, Bhatt S, Brownstein JS, Collier N, Myers MF, George DB, Gething PW.** 2013. Global mapping of infectious disease. *Philos Trans R Soc London Series B, Biol Sci* **368**:20120250.

34. **World Health Organization.** 2015. VietNam: WHO statistical profile.

35. **Nguyen TA, Yagyu F, Okame M, Phan TG, Trinh QD, Yan H, Hoang KT, Cao AT, Le Hoang P, Okitsu S, Ushijima H.** 2007. Diversity of viruses associated with acute gastroenteritis in children hospitalized with diarrhea in Ho Chi Minh City, Vietnam. *J Med Virol* **79**:582–590.

36. **Van Man N, Luan LT, Trach DD, Thanh NTH, Van Tu P, Long NT, Anh DD, Fischer TK, Ivanoff B, Gentsch JR, Glass RI, Luan le T.** 2005. Epidemiological profile and burden of rotavirus diarrhea in Vietnam: 5 years of sentinel hospital surveillance, 1998-2003. *J Infect Dis* **192 Suppl** :S127–32.

37. **Nguyen VMT, Huynh PL, Dang DT, Nguyen TLH, Phan VT, Le TL, Ivanoff B, Gentsch JR, Glass RI.** 2001. The epidemiology and disease burden of rotavirus in Vietnam: sentinel surveillance at 6 hospitals. *J Infect Dis* **183**:1707–1712.

38. **Vu Tra My P, Rabaa M a, Donato C, Cowley D, Vinh Phat V, Thi Ngoc Dung T, Hong Anh P, Vinh H, Bryant JE, Kellam P, Thwaites G, Woolhouse MEJ, Kirkwood CD, Baker S.** 2014. Novel porcine-like human G26P[19] rotavirus identified in hospitalized

pediatric diarrhea patients in Ho Chi Minh City. *J Gen Virol*.

39. **Thompson CN, Phan MVT, Hoang NVM, Minh P Van, Vinh NT, Thuy CT, Nga TTT, Rabaa MA, Duy PT, Dung TTN, Phat VV, Nga TVT, Tu LTP, Tuyen HT, Yoshihara K, Jenkins C, Duong VT, Phuc H Le, Tuyet PTN, Ngoc NM, Vinh H, Chinh NT, Thuong TC, Tuan HM, Hien TT, Campbell JI, Chau NVV, Thwaites G, Baker S.** 2015. A prospective multi-center observational study of children hospitalized with diarrhea in Ho Chi Minh City, Vietnam. *Am J Trop Med Hyg* **92**:1045–52.
40. **Trang N V, Luan LT, Kim-anh LT, Hau VTB, Nhung LTH, Phasuk P, Setrabutr O, Shirley H, Vinje J, Anh DD, Mason CJ.** 2012. Detection and Molecular Characterization of Noroviruses and Sapoviruses in Children Admitted to Hospital With Acute Gastroenteritis in Vietnam **297**:290–297.
41. **Landaeta ME, Dove W, Vinh H, Cunliffe N a, Campbell J, Parry CM, Farrar JJ, Hart CA.** 2003. Characterization of rotaviruses causing diarrhoea in Vietnamese children. *Ann Trop Med Parasitol* **97**:53–59.
42. **Nishio O, Matsui K, Lan DT, Ushijima H, Isomura S.** 2000. Rotavirus infection among infants with diarrhea in Vietnam. *Pediatr Int* **42**:422–424.
43. **Ngo TC, Nguyen BM, Dang DA, Nguyen HT, Nguyen TT, Tran VN, Vu TT, Ogino M, Alam MM, Nakagomi T, Nakagomi O, Yamashiro T.** 2009. Molecular epidemiology of rotavirus diarrhoea among children in Haiphong, Vietnam: the emergence of G3 rotavirus. *Vaccine* **27 Suppl 5**:F75–80.
44. **Doan LTP, Okitsu S, Nishio O, Pham DT, Nguyen DH, Ushijima H.** 2003. Epidemiological features of rotavirus infection among hospitalized children with gastroenteritis in Ho Chi Minh City, Vietnam. *J Med Virol* **69**:588–594.
45. **Anh DD, Thiem VD, Fischer TK, Canh DG, Minh TT, Tho le H, Van Man N, Luan le T, Kilgore P, von Seidlein L, Glass RI.** 2006. The burden of rotavirus diarrhea in Khanh Hoa Province, Vietnam: baseline assessment for a rotavirus vaccine trial. *Pediatr Infect Dis J* **25**:37–40.
46. **Tamura T, Nishikawa M, Anh DD, Suzuki H.** 2010. Molecular epidemiological study of rotavirus and norovirus infections among children with acute gastroenteritis in Nha Trang, Vietnam, December 2005–June 2006. *Jpn J Infect Dis* **63**:405–11.
47. **Nguyen TV, Le Van P, Le Huy C, Weintraub A.** 2004. Diarrhea Caused by Rotavirus in Children Less than 5 Years of Age in Hanoi, Vietnam. *J Clin Microbiol* **42**:5745–5750.
48. **Tra My PV, Rabaa M a, Vinh H, Holmes EC, Hoang NVM, Vinh NT, Phuong LT, Tham NT, Bay PVB, Campbell JI, Farrar J, Baker S.** 2011. The emergence of rotavirus G12 and the prevalence of enteric viruses in hospitalized pediatric diarrheal patients in southern Vietnam. *Am J Trop Med Hyg* **85**:768–75.
49. **Cotten M, Oude Munnink B, Canuti M, Deijs M, Watson SJ, Kellam P, van der Hoek L.** 2014. Full genome virus detection in fecal samples using sensitive nucleic Acid preparation, deep sequencing, and a novel iterative sequence classification algorithm. *PLoS One* **9**:e93269.
50. **de Vries M, Deijs M, Canuti M, van Schaik BDC, Faria NR, van de Garde MDB, Jachimowski LCM, Jebbink MF, Jakobs M, Luyf ACM, Coenjaerts FEJ, Claas ECJ, Molenkamp R, Koekkoek SM, Lammens C, Leus F, Goossens H, Ieven M, Baas F, van der Hoek L.** 2011. A sensitive assay for virus discovery in respiratory clinical samples. *PLoS One* **6**:e16118.
51. **de Vries M, Oude Munnink BB, Deijs M, Canuti M, Koekkoek SM, Molenkamp R, Bakker M, Jurriaans S, van Schaik BDC, Luyf AC, Olabarriaga SD, van Kampen AHC, van der Hoek L.** 2012. Performance of VIDISCA-454 in feces-suspensions and serum. *Viruses* **4**:1328–34.

- 736 52. **Boom R, Sol CJ, Salimans MM, Jansen CL, Wertheim-van Dillen PM, van der Noordaa**
737 **J.** 1990. Rapid and simple method for purification of nucleic acids. *J Clin Microbiol* **28**:495–
738 503.
- 739 53. **Endoh D, Mizutani T, Kirisawa R, Maki Y, Saito H, Kon Y, Morikawa S, Hayashi M.**
740 2005. Species-independent detection of RNA virus by representational difference analysis
741 using non-ribosomal hexanucleotides for reverse transcription. *Nucleic Acids Res* **33**:e65.
- 742 54. **Watson SJ, Welkers MRA, Depledge DP, Coulter E, Breuer JM, Jong D De, Kellam P, B**
743 **PTRS, Jong MD De.** 2013. Viral population analysis and minority-variant detection using
744 short read next-generation sequencing. *Philos Trans R Soc Sci* **368**.
- 745 55. **Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,**
746 **Nikolenko SI, Pham S, Prjibelski AD, Pyshkin A V, Sirotkin A V, Vyahhi N, Tesler G,**
747 **Alekseyev MA, Pevzner PA.** 2012. SPAdes: a new genome assembly algorithm and its
748 applications to single-cell sequencing. *J Comput Biol* **19**:455–77.
- 749 56. **Boetzer M, Henkel C V, Jansen HJ, Butler D, Pirovano W.** 2011. Scaffolding pre-
750 assembled contigs using SSPACE. *Bioinformatics* **27**:578–9.
- 751 57. **Edgar RC.** 2010. Search and clustering orders of magnitude faster than BLAST.
752 *Bioinformatics* **26**:2460–2461.
- 753 58. **Maes P, Matthijnssens J, Rahman M, Van Ranst M.** 2009. RotaC: a web-based tool for
754 the complete genome classification of group A rotaviruses. *BMC Microbiol* **9**:238.
- 755 59. **Gautam R, Esona MD, Mijatovic-Rustempasic S, Ian Tam K, Gentsch JR, Bowen MD.**
756 2013. Real-time RT-PCR assays to differentiate wild-type group A rotavirus strains from
757 Rotarix(®) and RotaTeq(®) vaccine strains in stool samples. *Hum Vaccin Immunother*
758 **10**:767–777.
- 759 60. **Matthijnssens J, Joelsson DB, Warakomski DJ, Zhou T, Mathis PK, van Maanen M-H,**
760 **Ranheim TS, Ciarlet M.** 2010. Molecular and biological characterization of the 5 human-
761 bovine rotavirus (WC3)-based reassortant strains of the pentavalent rotavirus vaccine,
762 RotaTeq. *Virology* **403**:111–27.
- 763 61. **Edgar RC.** 2004. MUSCLE: multiple sequence alignment with high accuracy and high
764 throughput. *Nucleic Acids Res* **32**:1792–7.
- 765 62. **Larsson A.** 2014. AliView: a fast and lightweight alignment viewer and editor for large
766 datasets. *Bioinformatics* **30**:3276–8.
- 767 63. **Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ.** 2015. IQ-TREE: A Fast and
768 Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol Biol*
769 *Evol* **32**:268–274.
- 770 64. **Drummond AJ, Rambaut A.** 2007. BEAST: Bayesian evolutionary analysis by sampling
771 trees. *BMC Evol Biol* **7**:214.
- 772 65. **Papp H, Borzák R, Farkas S, Kisfali P, Lengyel G, Molnár P, Melegh B, Matthijnssens**
773 **J, Jakab F, Martella V, Bányaí K.** 2013. Zoonotic transmission of reassortant porcine
774 G4P[6] rotaviruses in Hungarian pediatric patients identified sporadically over a 15 year
775 period. *Infect Genet Evol* **19**:71–80.
- 776 66. **Degiseppa JI, Beltramino JC, Millán A, Stupka J a., Parra GI.** 2013. Complete genome
777 analyses of G4P[6] rotavirus detected in Argentinean children with diarrhoea provides
778 evidence of interspecies transmission from swine. *Clin Microbiol Infect* **19**:E367–E371.
- 779 67. **Martinez M, Galeano ME, Akopov A, Palacios R, Russomando G, Kirkness EF, Parra**
780 **GI.** 2014. Whole-genome analyses reveals the animal origin of a rotavirus G4P[6] detected
781 in a child with severe diarrhea. *Infect Genet Evol* **27C**:156–162.
- 782 68. **Bucardo F, Rippinger CM, Svensson L, Patton JT.** 2012. Vaccine-derived NSP2 segment
783 in rotaviruses from vaccinated children with gastroenteritis in Nicaragua. *Infect Genet Evol*

12:1282–94.

69. **Ward CW, Elleman TC, Azad AA, Dyal-Smith ML.** 1984. Nucleotide sequence of gene segment 9 encoding a nonstructural protein of UK bovine rotavirus. *Virology* **134**:249–53.
70. **Mukherjee a, Ghosh S, Bagchi P, Dutta D, Chattopadhyay S, Kobayashi N, Chawla-Sarkar M.** 2011. Full genomic analyses of human rotavirus G4P[4], G4P[6], G9P[19] and G10P[6] strains from North-eastern India: evidence for interspecies transmission and complex reassortment events. *Clin Microbiol Infect* **17**:1343–6.
71. **Martel-Paradis O, Laurin M-A, Martella V, Sohal JS, L’Homme Y.** 2013. Full-length genome analysis of G2, G9 and G11 porcine group A rotaviruses. *Vet Microbiol* **162**:94–102.
72. **Wang Y, Kobayashi N, Nagashima S, Zhou X, Ghosh S.** 2010. Full Genomic Analysis of a Porcine – Bovine Reassortant G4P[6] Rotavirus Strain R479 Isolated From an Infant in China. *J Med Virol* **1102**:1094–1102.
73. **Zeller M, Patton JT, Heylen E, De Coster S, Ciarlet M, Van Ranst M, Matthijnsens J.** 2012. Genetic analyses reveal differences in the VP7 and VP4 antigenic epitopes between human rotaviruses circulating in Belgium and rotaviruses in Rotarix and RotaTeq. *J Clin Microbiol* **50**:966–76.
74. **Matthijnsens J, Heylen E, Zeller M, Rahman M, Lemey P, Van Ranst M.** 2010. Phylodynamic analyses of rotavirus genotypes G9 and G12 underscore their potential for swift global spread. *Mol Biol Evol* **27**:2431–2436.
75. **Dung TTN, Phat VV, Nga TTTVT, My PVT, Duy PT, Campbell JI, Thuy CT, Hoang NVM, Van Minh P, Le Phuc H, Tuyet PTN, Vinh H, Kien DTH, Huy HLA, Vinh NT, Hau NTT, Chinh NT, Thuong TC, Tuan HM, Simmons C, Farrar JJ, Baker S, Thi TDNT, Vu TNT, Minh P Van, Phuc H Le, Thi PTN, Thi DKH, Le HHA, Thi NHT.** 2012. The validation and utility of a quantitative one-step multiplex RT real-time PCR targeting Rotavirus A and Norovirus. *J Virol Methods* **187**:1–6.
76. **Anderson EJ, Weber SG.** 2004. Rotavirus infection in adults. *Lancet Infect Dis* **4**:91–99.
77. **Ghosh S, Kobayashi N.** 2011. Whole-genomic ana-lysis of rotavirus strains: current status and future prospects. *Future Microbiol* **6**:1049–1065.
78. **Molinari B, Lorenzetti E, Otonel RAA, Alfieri AF, Alfieri AA.** 2014. Species H rotavirus detected in piglets with diarrhea, Brazil, 2012. *Emerg Infect Dis* **20**:2012–2015.
79. **Marthaler D, Rossow K, Culhane M, Goyal S, Collins J, Matthijnsens J, Nelson M, Ciarlet M.** 2014. Widespread rotavirus H in domesticated pigs, United States. *Emerg Infect Dis* **20**:1195–1198.
80. **Lahon A, Ingle VC, Birade HS, Raut CG, Chitambar SD.** 2014. Molecular characterization of group B rotavirus circulating in pigs from India: Identification of a strain bearing a novel VP7 genotype, G21. *Vet Microbiol* **174**:342–52.
81. **Molinari BLD, Alfieri AF, Alfieri AA.** 2014. Genetic variability of VP6, VP7, VP4, and NSP4 genes of porcine rotavirus group H detected in Brazil. *Virus Res* **197**:48–53.
82. **Marthaler D, Rossow K, Culhane M, Collins J, Goyal S, Ciarlet M, Matthijnsens J.** 2013. Identification, phylogenetic analysis and classification of porcine group C rotavirus VP7 sequences from the United States and Canada. *Virology* **446**:189–98.
83. **Amimo JO, Vlasova a N, Saif LJ.** 2013. Prevalence and genetic heterogeneity of porcine group C rotaviruses in nursing and weaned piglets in Ohio, USA and identification of a potential new VP4 genotype. *Vet Microbiol* **164**:27–38.
84. **Marton S, Mihalov-Kovcs E, Dóro R, Csata T, Feher E, Oldal M, Jakab F, Matthijnsens J, Martella V, Banyai K.** 2015. Canine rotavirus C strain detected in hungary shows marked genotype diversity. *J Gen Virol* **96**:3059–3071.

85. **Marthaler D, Rossow K, Gramer M, Collins J, Goyal S, Tsunemitsu H, Kuga K, Suzuki T, Ciarlet M, Matthijnssens J.** 2012. Detection of substantial porcine group B rotavirus genetic diversity in the United States, resulting in a modified classification proposal for G genotypes. *Virology* **433**:85–96.
86. **Otto PH, Rosenhain S, Elschner MC, Hotzel H, Machnowska P, Trojnar E, Hoffmann K, Johne R.** 2015. Detection of rotavirus species A, B and C in domestic mammalian animals with diarrhoea and genotyping of bovine species A rotavirus strains. *Vet Microbiol* **179**:168–176.
87. **Zhirakovskaia E, Tikunov A, Klemesheva V, Loginovskikh N, Netesov S, Tikunova N.** 2016. First genetic characterization of rotavirus C in Russia. *Infect Genet Evol* **39**:1–8.
88. **Lahon A, Maniya NH, Tambe GU, Chinchole PR, Purwar S, Jacob G, Chitambar SD.** 2013. Group B rotavirus infection in patients with acute gastroenteritis from India: 1994-1995 and 2004-2010. *Epidemiol Infect* **141**:969–75.
89. **Alam MM, Pun SB, Gauchan P, Yokoo M, Doan YH, Tran TNH, Nakagomi T, Nakagomi O, Pandey BD.** 2013. The first identification of rotavirus B from children and adults with acute diarrhoea in kathmandu, Nepal. *Trop Med Health* **41**:129–34.
90. **Dyall-smith ML, Lazdins I, Tregeart GW, Holmes IANH.** 1986. Location of the major antigenic sites involved in rotavirus serotype-specific neutralization. *Proc Natl Acad Sci U S A* **83**:3465–3468.
91. **Nishikawa K, Hoshino Y, Taniguchi K, Green KY, Greenberg HB, Kapikian a Z, Chanock RM, Gorziglia M.** 1989. Rotavirus VP7 neutralization epitopes of serotype 3 strains. *Virology* **171**:503–15.
92. **Woolhouse MEJ, Rambaut A, Kellam P.** 2015. Lessons from Ebola : Improving infectious disease surveillance to inform outbreak management. *Sci Transl Med* **7**:1–9.
93. **Rabaa MA, Tue NT, Phuc TM, Carrique-Mas J, Saylor K, Cotten M, Bryant JE, Nghia HDT, Cuong N Van, Pham HA, Berto A, Phat VV, Dung TTN, Bao LH, Hoa NT, Wertheim H, Nadjm B, Monagin C, van Doorn HR, Rahman M, Tra MPV, Campbell JI, Boni MF, Tam PTT, van der Hoek L, Simmonds P, Rambaut A, Toan TK, Van Vinh Chau N, Hien TT, Wolfe N, Farrar JJ, Thwaites G, Kellam P, Woolhouse MEJ, Baker S.** 2015. The Vietnam Initiative on Zoonotic Infections (VIZIONS): A Strategic Approach to Studying Emerging Zoonotic Infectious Diseases. *Ecohealth* **12**:726–735.
94. **Munnink BBO, Phan MVT, Kellam P, Cotten M, Consortium V.** 2016. Complete Genome Characterization of Two Wild-Type Measles Viruses from Vietnamese Infants during the 2014 Outbreak. *Genome* **4**:1–2.
95. **Chen JZ, Settembre EC, Aoki ST, Zhang X, Bellamy a R, Dormitzer PR, Harrison SC, Grigorieff N.** 2009. Molecular interactions in rotavirus assembly and uncoating seen by high-resolution cryo-EM. *Proc Natl Acad Sci U S A* **106**:10644–8.