

Title: *Mycobacterium ulcerans* low infectious dose and atypical mechanical transmission support insect bites and puncturing injuries in the spread of Buruli ulcer.

John R. Wallace¹, Kirstie M. Mangas², Jessica L. Porter², Renee Marcsisin², Sacha J. Pidot², Brian Howden², Till F. Omansen^{2,3}, Weiguang Zeng², Jason K. Axford⁴, Paul D. R. Johnson⁴ and Timothy P. Stinear^{2,*}

¹Department of Biology, Millersville University, PA, USA

²Department of Microbiology and Immunology, at the Peter Doherty Institute for Infection and Immunity, University of Melbourne, Melbourne, 3000, Australia

³Department of Internal Medicine, University of Groningen, Groningen, 9700 RB, The Netherlands

⁴Pest and Environmental Adaptation Research Group, Bio21 Institute and School of BioSciences, University of Melbourne, Parkville, Vic 3052, Australia.

⁵Department of Infectious Diseases, Austin Health, Heidelberg, Victoria, 3084, Australia

***Corresponding authors:** tstinear@unimelb.edu.au, john.wallace@millersville.edu

Article Summary Line: This study shows that minor puncturing injuries to a skin surface contaminated with *Mycobacterium ulcerans*, caused by accidental trauma or insect bites, can lead to Buruli ulcer.

Short Title: Buruli ulcer transmission

Keywords: Buruli ulcer, *Mycobacterium ulcerans*, neglected tropical diseases, infectious diseases transmission, vector competence.

25 **Abstract**

26 Addressing the transmission enigma of the neglected disease Buruli ulcer (BU) is a World Health
 27 Organization priority. In Australia, we have observed an association between mosquitoes harboring the
 28 causative agent, *Mycobacterium ulcerans*, and BU. Here we tested a contaminated skin model of BU
 29 transmission by dipping the tails from healthy mice in cultures of the causative agent, *Mycobacterium*
 30 *ulcerans*. Tails were exposed to mosquito (*Aedes notoscriptus* and *Aedes aegypti*) blood feeding or punctured
 31 with sterile needles. Two of 11 of mice with *M. ulcerans* contaminated tails exposed to feeding *A.*
 32 *notoscriptus* mosquitoes developed BU. Eighteen of 20 mice subjected to contaminated tail needle puncture
 33 developed BU. Mouse tails coated only in bacteria did not develop disease. We observed a low infectious
 34 dose-50 of four colony-forming units and a median incubation time of 12 weeks, consistent with data from
 35 human infections. We have uncovered a highly efficient and biologically plausible atypical transmission
 36 mode of BU via natural or anthropogenic skin punctures.

37

Author summary

Buruli ulcer is a neglected tropical disease caused by infection with *Mycobacterium ulcerans*. Unfortunately, how people contract this disease is not well understood. Here we show for the first time using experimental infections in mice that a very low dose of *M. ulcerans* delivered beneath the skin by a minor injury caused by a blood-feeding insect (mosquito) or a needle puncture is sufficient to cause Buruli ulcer. This research provides important laboratory evidence to advance our understanding of Buruli ulcer disease transmission.

Introduction

Among the 17 neglected tropical diseases the World Health Organization (WHO) has targeted for control and elimination, only Leprosy and Buruli ulcer (BU) have unknown modes of transmission [1]. The search to understand how humans contract BU spans more than 70 years since the causative agent, *Mycobacterium ulcerans*, was first identified [2]. There are persistent and emerging foci of BU cases across the world, in particular Africa and Australia [3]. BU is characterized by necrotizing skin lesions, caused by localized proliferation of *M. ulcerans* in subcutaneous tissue. BU is rarely fatal, but untreated infections leave patients with significant disfigurement and disability, with damaging personal and economic consequences [4, 5]. Researchers have long been struck by the characteristic epidemiology of BU, with cases occurring in highly geographically circumscribed regions (sometimes less than a few square kilometres) and risk factors for infection that include gardening, insect bites and proximity to (but not necessarily contact with) lacustrine/riverine regions [6-14]. Human-to-human spread is considered unlikely [14]. Disease transmission is thought to occur by contact with an environment contaminated with *Mycobacterium ulcerans* but exactly where the pathogen resides and why it appears so geographically restricted have yet to be determined. [15].

M. ulcerans is very slow growing (doubling time >48 hrs) and this poses a problem for source tracking efforts as it is difficult to isolate the bacteria in pure culture from complex environmental specimens [16]. *M. ulcerans* has only once been isolated from a non-clinical source, an aquatic water bug (Gerridae) from Benin [16]. Quantitative PCR targeting *M. ulcerans*-specific DNA is the most frequently used technique in surveys

of environmental specimens. A comprehensive review of the many field and lab studies that have examined reservoir and transmission of BU has highlighted the range of organisms from aquatic insects, fish, amphibia, and in Australia certain native marsupials that can serve as potential reservoirs for *M. ulcerans* [15, 17]. Since the first observation that biting aquatic insects can harbour *M. ulcerans* [18], studies of BU transmission have largely focused on the potential for insects to biologically vector *M. ulcerans* implying that *M. ulcerans* undergoes a propagative or reproductive mode of development in an insect [19-23]. Several case-control studies, including from both Australia and Africa have suggested insects may play a role in transmission [10, 11]. However, there is no compelling experimental evidence for single-mode biological transmission of *M. ulcerans* via insect vectors.

In southeastern Australia, we noted Buruli lesions on exposed areas likely to attract biting insects, some patients with every brief exposure times to endemic areas [24, 25] and 2004 we began a study that identified *M. ulcerans* DNA associated with mosquitoes captured in endemic areas [19].

Materials and Methods

Bacterial isolates and culture conditions

M. ulcerans strain JKD8049 and bioluminescent *M. ulcerans* JKD8049 (harbouring plasmid pMV306 *hsp:luxG13*) [26, 27] were cultured in 7H9 broth or Middlebrook 7H10 agar, containing 10% oleic-albumin-dextrose-catalase growth supplement (Middlebrook, Becton Dickinson, Sparks, MD, USA) and 0.5% glycerol (v/v) at 30°C. Colony counts from bacterial cultures or tissue specimens were performed using spot plating. Five x 3µl volumes of serial 10-fold dilutions (10^{-1} to 10^{-5}) of a culture or tissue preparation were spotted onto 7H10 agar plates with a 5x5 grid marked. The spots were allowed to dry, the plates loosely wrapped in plastic bags and then incubated as above for 10 weeks before counting colonies. Data analysis was performed using GraphPad Prism (v 6.0). All culture extracts were screened by LC-MS for the presence of mycolactones as previously described to ensure bacteria used in transmission experiments remained fully virulent [28].

90

91 *Experimental animals*

92 The animal ethics committee (AEC) of the University of Melbourne approved all animal experiments under
93 approval number AEC: 1312775.2. □BALB/c mice were purchased from ARC (Canning Vale, Australia)
94 and housed in individual ventilated cages. Upon arrival, animals were acclimatizing for 5 days. Food and
95 water were given *ad libitum*.

96

97 *Aedes notoscriptus* and *Aedes aegypti* rearing. Wild caught mosquitoes were sourced from around Cairns,
98 Queensland, Australia. *A. notoscriptus* and *A. aegypti* colonies were reared in a Physical Containment Level
99 2 (PC2) laboratory environment at 26 °C using previously described methods, with the addition of brown
100 paper used as the oviposition substrate for *A. notoscriptus* [29].

101

102 *Mosquito-mouse transmission experiments*

103 Four-week old female BALB/c mice were anaesthetized and their tails coated in a thin film of *M. ulcerans* by
104 dipping the tails in a Petri dish containing 20mL of bacterial culture (concentration ~10⁶ CFU/mL). The tail
105 only was then exposed to a 200mm x 200mm x 200mm cage of 20 adult female mosquitoes for a period of
106 15 minutes. The number of insects biting each mouse was recorded over the exposure period by continuous
107 observation. Mice were then observed weekly for up to six months for signs of tail lesions. Sterile needle
108 stick (25G or 30G needle) and no-trauma were used as controls. An additional control consisted of tails
109 dipped in sterile culture broth only and subjected to mosquito biting or sterile needle stick.

110

111 *Real time quantitative PCR*. For each mosquito that blood-fed DNA was individually extracted from the
112 dissected head, abdomen and legs of each insect using the Mo Bio Powersoil DNA extraction kit following
113 manufacturer's instructions (Mo Bio Laboratories Inc., Carlsbad CA USA). DNA was similarly extracted
114 from mouse tissue. Procedural extraction control blanks (sterile water) were included at a frequency of 10%
115 to monitor potential PCR contamination, in addition to no-template negative controls. IS2404 quantitative

PCR (qPCR) was performed as described [30]. IS2404 cycle threshold (Ct) values were converted to genome equivalents (GE) to estimate bacterial load within a sample by reference to a standard curve ($r^2=0.9312$, $y=[-3.000\ln(x)+39.33]*Z$, where $y=Ct$ and x =amount of DNA [fg] and Z =the dilution factor]), calculated using dilutions of genomic DNA from *M. ulcerans* strain JKD8049, quantified using fluorimetry (Qubit, Invitrogen) [30].

Preparation of mouse tissue for analysis. At the end of the experimental period or when a clinical end-point was reached mice were humanely killed. The region of a mouse-tail spanning a likely lesion was cut into three equal sections for histology, qPCR and CFU counts. Individual tail pieces for CFU counts were weighed and placed into sterile 2ml screw capped tube containing 0.5g of large glass beads and 600µl of sterile 1x PBS. Tissues were homogenized using four rounds of 2x 30second pulses in a high-speed tissue-disruptor at 6500 rpm, with tubes placed on ice for 5 minutes between each round. A 300µl volume of this homogenate was decontaminated with 300µl of 2% NaOH (v/v) and incubated at room temperature for 15 minutes. The preparation was neutralized drop-wise with a 10% solution of orthophosphoric acid (v/v) with added bromophenol blue until the solution changed from blue to clear. The mixtures were diluted in PBS and CFUs determined by spot plating as described above.

Histology. Sections of mouse-tails were fixed in 10% (w/v) neutral-buffered-formalin and imbedded in paraffin. Each mouse-tail was sectioned transversely (four micron thickness) and subjected to Ziehl-Neelson and hematoxylin/eosin staining. The fixed and stained tissue sections were examined by light microscopy.

Infectious Dose. To estimate the infectious dose we measured the surface area of five dissected mouse-tails to obtain an average surface area ($493.3 \pm 41.1 \text{ mm}^2$). Using ten mouse-tails, we then calculated the average volume of *M. ulcerans* 7H9 Middlebrook culture adhering to the tail surface ($32.4 \pm 4.2 \text{ mL}$), the concentration of bacteria in the cultures used, and the surface area of the tips of 25G and 30G needles used to deliver the puncture wounds (0.207 mm^2 and 0.056 mm^2


respectively). These parameters were then used to calculate the infectious dose, assuming the bacteria were evenly distributed over the tail surface (Fig. 1C). A standard curve was interpolated using non-linear regression and an ID₅₀ estimated using (GraphPad Prism v 7.0a).

Results

M. ulcerans is efficiently transmitted to a mammalian host by an atypical mechanical means.

We established a murine model of *M. ulcerans* transmission (model-1) that represented a skin surface contaminated with the bacteria and then subjected to a minor penetrating trauma, via either a mosquito bite or needle stick puncture. In a first experiment two of six mice with their tails coated with *M. ulcerans* and then bitten by mosquitoes developed lesions (Table 1, Fig. 1C, Fig. 2A). Histology of these lesions confirmed a subcutaneous focus of AFB, within a zone of necrotic tissue. There was also characteristic epithelial hyperplasia adjacent to the site of infection (Fig. 2B,C). Material extracted from the lesions was IS2404 qPCR-positive and culture positive for *M. ulcerans* (Supplementary Table S1). Mice bitten by mosquitoes but with tails coated only with sterile culture media did not develop lesions (Table 1). In the same experiment, we also subjected five mice to a single needle stick puncture. Each mouse had their tail coated with *M. ulcerans* as for the mosquito biting. Four of these five mice developed *M. ulcerans* positive lesions (Table 1, Fig. 2D), with subcutaneous foci of infection and viable bacteria (Fig. 2F). Six mice with their tails coated with *M. ulcerans* but not subjected to a puncturing injury did not develop lesions and remained healthy until the completion of the experiment at six months. This experiment suggested that minor penetrating skin trauma (defined here as a puncture <0.5mm diameter and <2mm deep) to a skin surface contaminated with *M. ulcerans* is sufficient to cause infection. It also revealed a means by which mosquitoes could act as atypical mechanical vectors of *M. ulcerans*.

166



167

168 **Fig.1: Schematic representations of the two BU transmission models tested in this study.** (A) Model-1
 169 tests transmission of *M. ulcerans* present on a skin surface following a puncturing injury created by mosquito
 170 blood-feeding or needle stick. (B) Visualization of bioluminescent *M. ulcerans* JKD8049 (harbouring
 171 plasmid pMV306 *hsp:luxG13*) [26, 27] on the mouse-tail in model-1, showing the distribution of bacteria
 172 immediately after coating for two mice, versus an uncoated animal. *M. ulcerans* concentration was 10^6
 173 CFU/mL.

174

175




Fig. 2: Atypical mechanical transmission of *M. ulcerans*. (A) An example of the development of Buruli ulcer following mosquito blood-feeding through a skin surface (mouse-tail) contaminated with *M. ulcerans*. (B) Composite histological cross-section with Ziehl–Neelsen staining through the infected tail showing the focus of AFB bacteria (arrow) within the subcutaneous tissue. (C) Higher magnification view of the focus of infection, with the yield of viable *M. ulcerans* obtained from the infected tissue. Panels (D) – (F) show the same analyses as for the mosquito-bitten mouse #182, but for a mouse developing a lesion following sterile needle-stick puncture through a contaminated skin surface (mouse #201).

M. ulcerans burden on mosquitoes correlates with transmission.

Then, using approximately the same dose of bacteria to coat the mouse-tails, we repeated experiment-1 but with *Aedes aegypti* because of the close association of this mosquito to humans world-wide and their vector competency for viral pathogens. Despite more mosquito bites per mouse than the first experiment, none of the five insect-exposed mice developed lesions (Table 1). In contrast however, four of five mice subjected to single, needle stick puncture developed *M. ulcerans* positive tail lesions (Table 1). We assessed the burden of *M. ulcerans* by individual IS2404 qPCR of the head, abdomen and legs for each mosquito that blood fed (Fig. 3). A summary of these results is shown in Fig. 3A. We noted that the bacterial load (expressed as genome equivalents [GE]) was significantly higher in the heads of mosquitoes associated with mice that developed lesions ($p < 0.05$) (Fig. 3B, Supplementary Table S1). These data point to a threshold, above which mosquitoes can become competent mechanical vectors for *M. ulcerans* transmission.




Fig. 3: Summary of *M. ulcerans* burden on mosquitoes post-feeding under two models of

transmission. (A) Visualization of the mean number of *M. ulcerans* detected per dissected

mosquito segment, as assessed by IS2404 qPCR and expressed as genome equivalents (GE). ‘N’

indicates the total number of mosquitoes tested. Red-shaded mosquitoes transmitted *M. ulcerans*,

leading to mouse tail lesions. Green-shaded mosquitoes blood-fed on mouse-tails but lesions did not

develop. (B, C, D) Plots of the individual qPCR results for each mosquito segment, listed by

experiment. Red dots correspond to qPCR bacterial load for mosquitoes that transmitted *M.*

ulcerans infection. Null hypothesis (no difference in bacterial load) was rejected (p < 0.05)*

(unpaired, two-tailed *t* test). Horizontal bar indicates the mean bacterial load per mosquito. The y-

axis is GE and x-axis is experiment. The qPCR data for individual insects is contained in

Supplementary Table S1.

Estimation of incubation period and infectious dose of transmission model-1

Based on the time until a tail lesion was first observed, we estimated a median incubation period (IP) of 12 weeks (Fig. 4A). This result overlaps with the IP in humans for BU, estimated in different epidemiological studies from 4-10 weeks in Uganda during the 1960s [14] and 4 - 37 weeks in south east Australia [25]. We also estimated the infectious dose₅₀ (ID₅₀). We used six different concentrations of *M. ulcerans* to coat the tails of mice (n=5 mice/dilution) and observed the number of mice for each dilution that developed Buruli ulcer, allowing an ID₅₀ estimate of 4 CFU (Fig. 4B). To our knowledge this is the first estimate of an *M. ulcerans* infectious dose and indicates that a surprisingly small quantity of this slow growing mycobacterium inoculated just below the skin is sufficient to cause disease.




Fig. 4: *M. ulcerans* incubation period and infectious dose₅₀. (A) Incubation period of *M. ulcerans* based on the time between sterile-needle puncture of an *M. ulcerans* contaminated mouse-tail and first observation of a lesion. (B) Estimated *M. ulcerans* ID₅₀ for transmission model-1.

Discussion

Here, we show for the first time a highly efficient atypical mode of mechanical transmission of *Mycobacterium ulcerans* to a mammalian host that implicates both biting insects and puncturing injuries. This research was designed around established frameworks for implicating vectors in disease transmission and provides the necessary causational evidence to help resolve the 80-year mystery on how *M. ulcerans* is spread to people [15, 31]. The efficient establishment of BU we have shown here via minor penetrating trauma through a contaminated skin surface is an atypical form of mechanical transmission *sensu lato* (s.l.) but it nonetheless satisfactorily fulfills one of the Barnett Criteria [31]. In vector ecology, mechanical transmission, *sensu strictu* (s.s.), is defined as a non-circulative process involving accidental transport of the pathogen. That is, the pathogen, in some fashion, nonspecifically associates or contaminates the mouthparts (stylet) of an arthropod vector. Insect mechanical transmission s.s. of BU implies that if *M. ulcerans* were ingested and then egested via regurgitation or salivation, the mechanism would act more like a syringe than a needle [32]. Such a mode of *M. ulcerans* disease transmission is supported by previous laboratory studies in which *Naucoris* and Belostmatid water bugs were contaminated via feeding on maggot prey that had been injected with *M. ulcerans* or fed naturally on dietary contaminated larval mosquito prey [23, 33].

Our demonstration in the current study of mechanical transmission s.l. implies there are potentially multiple or parallel pathways of *M. ulcerans* infection [31]. Examples of bacterial diseases with multiple transmission modes include tularemia, plague and trachoma [34, 35]. Support for our mechanical transmission s.l. model also comes from the many field reports over the decades of *M. ulcerans* infection following trauma to the skin. Case reports have noted BU following a suite of penetrating injuries ranging from insect bites (ants, scorpions), snake bite, human bite, splinters, gunshot, hypodermic injections of medication and vaccinations [36-38]. Epidemiologists in Uganda during the 1960s and 70s suggested sharp-edged grasses might introduce the bacteria [39]. However, a recent laboratory study established that abrasions of the skin in Guinea pig models and subsequent application of *M. ulcerans* was not enough to cause an ulcer, however, this same study established that a subcutaneous injection would cause an ulcer [40]. As a sequel to this study in Guinea pigs, we raised the question of how likely it was that human skin could be

sufficiently coated in *M. ulcerans* that an injury from natural or anthropogenic sources could lead to infection. Other explanations for the transmission of *M. ulcerans* include linkages with human behavior that increase direct contact with human skin and contaminated water [15]. A recent study from Cameroon recorded the persistence of *M. ulcerans* over a 24-month period in a waterhole used by villagers (including BU patients) for bathing [41]. A scenario could be envisaged where a villager's skin surface becomes contaminated after bathing in such a water body and is primed for infection if (i) the concentration of bacteria is sufficiently high, and (ii) an inoculating event occurs. Whereas, in Australia, earlier studies have shown that *M. ulcerans* contamination of possum feces in and around the gardens of BU patients might present a similar skin surface contamination model in this region [17, 42]. Future experiments will address the possibility that insect vectors may be able to move *M. ulcerans* from one source and inject it into an animal or human.

Our focus on mechanical mosquito transmission s.s. arose from previous surveys in southeastern Australia where a strong association between *M. ulcerans* positive mosquitoes and human cases of BU has shown that *M. ulcerans* has not only been found on adult mosquitoes from both lab and field studies but also a biological gradient, where maximum likelihood estimates (MLE) of the proportion of *M. ulcerans*-positive mosquitoes increased as the number of cases of BU increased [19, 33, 43-46]. However, a recent study in Benin, West Africa found no evidence of *M. ulcerans* in association with adult mosquitoes [47]. The authors concluded that the mode of transmission might differ between southeastern Australia and Africa. Although, laboratory and fieldwork in West Africa suggest that aquatic insects, including mosquito larvae, play a role as reservoirs in nature for *M. ulcerans* that may be indirectly tied to transmission by serving as dispersal mechanisms [18, 23, 48]. Epidemiological studies have shown that direct contact with water is not a universal risk factor for BU [8, 11]. Prior exposure to insect bites and gardening are also independent risk factors for developing BU, while use of insect repellent is protective [11, 49].

Laboratory support to show mosquitoes can be competent vectors to spread BU is important additional

evidence required to satisfy accepted vector ecology criteria for implicating insects in disease transmission [15, 31]. We found that infection was established following very minor penetrating trauma. *Aedes notoscriptus* mosquitoes feed by insertion of a stylet, sheathed within the proboscis, beneath the skin of a host. The stylet has a diameter around 10 μ M tapering to 1 μ M at its tip and extending 1-2 mm below the skin surface. We estimated the density of *M. ulcerans* on the mouse-tails surface was 100-200 CFU/mm². Thus, the number of bacteria potentially injected during mosquito feeding through this contaminated surface is likely to be low, but this is consistent with our infectious dose estimates from needle-stick punctures, indicating an ID₅₀ of only 4 CFU (Fig. 4B). There are strong parallels here with *Mycobacterium leprae*, the agent of leprosy. Like BU, the mode of transmission of the leprosy bacillus is unclear, but the infective dose is known to be very low (10 bacteria) and epidemiological evidence suggests multiple transmission pathways, including entry of the bacteria after skin trauma [50, 51]. Our infective dose estimate for *M. ulcerans* is consistent with observations that pathogens producing locally acting molecules to cause disease (e.g. the polyketide toxin mycolactone of *M. ulcerans*) have lower infective doses [52].

In summary, we have uncovered a highly efficient and biologically plausible atypical transmission mode of *M. ulcerans* infection via natural or anthropogenic skin punctures. Reduction of exposure to insect bites, access to clean water for bathing, and prompt treatment of existing BU are concrete measures likely to interrupt BU transmission.

References

1. Anon. Second WHO report on neglected tropical diseases: sustaining the drive to overcome the global impact of neglected tropical diseases. Geneva, Switzerland: World Health Organization; 2013. p. 11-7.
2. P. M, Tolhurst JC, Buckle G, Sissons HA. A new mycobacterial infection in man. J Pathol Bacteriol. 1948;60(1):93-122. PubMed PMID: 18876541.
3. Johnson PD, Stinear T, Small PL, Pluschke G, Merritt RW, Portaels F, et al. Buruli ulcer (M. ulcerans infection): new insights, new hope for disease control. PLoS Med. 2005;2(4):e108. doi: 10.1371/journal.pmed.0020108. PubMed PMID: 15839744; PubMed Central PMCID: PMC1087202.
4. Amofah GK, Sagoe-Moses C, Adjei-Acquah C, Frimpong EH. Epidemiology of Buruli ulcer in Amansie West district, Ghana. Trans R Soc Trop Med Hyg. 1993;87(6):644-5. PubMed PMID: 8296362.
5. Asiedu K, Etuaful S. Socioeconomic implications of Buruli ulcer in Ghana: a three-year review. Am J Trop Med Hyg. 1998;59(6):1015-22. PubMed PMID: 9886216.
6. Debacker M, Portaels F, Aguiar J, Steunou C, Zinsou C, Meyers W, et al. Risk factors for Buruli ulcer, Benin. Emerg Infect Dis. 2006;12(9):1325-31. doi: 10.3201/eid1209.050598. PubMed PMID: 17073079; PubMed Central PMCID: PMC3294727.
7. Jacobsen KH, Padgett JJ. Risk factors for *Mycobacterium ulcerans* infection. Int J Infect Dis. 2010;14(8):e677-81. doi: 10.1016/j.ijid.2009.11.013. PubMed PMID: 20185351.
8. Landier J, Boisier P, Fotso Piam F, Noumen-Djeunga B, Sime J, Wantong FG, et al. Adequate wound care and use of bed nets as protective factors against Buruli Ulcer: results from a case control study in Cameroon. PLoS Negl Trop Dis. 2011;5(11):e1392. doi: 10.1371/journal.pntd.0001392. PubMed PMID: 22087346; PubMed Central PMCID: PMC3210760.

- 320 9. Nackers F, Johnson RC, Glynn JR, Zinsou C, Tonglet R, Portaels F. Environmental and
321 health-related risk factors for *Mycobacterium ulcerans* disease (Buruli ulcer) in Benin. Am J Trop
322 Med Hyg. 2007;77(5):834-6. PubMed PMID: 17984337.
- 323 10. Pouillot R, Matias G, Wondje CM, Portaels F, Valin N, Ngos F, et al. Risk factors for buruli
324 ulcer: a case control study in Cameroon. PLoS Negl Trop Dis. 2007;1(3):e101. doi:
325 10.1371/journal.pntd.0000101. PubMed PMID: 18160977; PubMed Central PMCID:
326 PMC2154388.
- 327 11. Quek TY, Athan E, Henry MJ, Pasco JA, Redden-Hoare J, Hughes A, et al. Risk factors for
328 *Mycobacterium ulcerans* infection, southeastern Australia. Emerg Infect Dis. 2007;13(11):1661-6.
329 doi: 10.3201/eid1311.061206. PubMed PMID: 18217548; PubMed Central PMCID: PMC3375781.
- 330 12. Raghunathan PL, Whitney EA, Asamoah K, Stienstra Y, Taylor TH, Jr., Amofah GK, et al.
331 Risk factors for Buruli ulcer disease (*Mycobacterium ulcerans* Infection): results from a case-
332 control study in Ghana. Clin Infect Dis. 2005;40(10):1445-53. doi: 10.1086/429623. PubMed
333 PMID: 15844067.
- 334 13. Sopoh GE, Barogui YT, Johnson RC, Dossou AD, Makoutode M, Anagonou SY, et al.
335 Family relationship, water contact and occurrence of Buruli ulcer in Benin. PLoS Negl Trop Dis.
336 2010;4(7):e746. doi: 10.1371/journal.pntd.0000746. PubMed PMID: 20644620; PubMed Central
337 PMCID: PMC2903473.
- 338 14. Uganda-Buruli-Group. Epidemiology of *Mycobacterium ulcerans* infection (Buruli ulcer) at
339 Kinyara, Uganda. Trans R Soc Trop Med Hyg. 1971;65(6):763-75. PubMed PMID: 5157438.
- 340 15. Merritt RW, Walker ED, Small PL, Wallace JR, Johnson PD, Benbow ME, et al. Ecology
341 and transmission of Buruli ulcer disease: a systematic review. PLoS Negl Trop Dis.
342 2010;4(12):e911. doi: 10.1371/journal.pntd.0000911. PubMed PMID: 21179505; PubMed Central
343 PMCID: PMC3001905.
- 344 16. Portaels F, Meyers WM, Ablordey A, Castro AG, Chemlal K, de Rijk P, et al. First
345 cultivation and characterization of *Mycobacterium ulcerans* from the environment. PLoS Negl Trop

- 346 Dis. 2008;2(3):e178. doi: 10.1371/journal.pntd.0000178. PubMed PMID: 18365032; PubMed
347 Central PMCID: PMC2268003.
- 348 17. Fyfe JA, Lavender CJ, Handasyde KA, Legione AR, O'Brien CR, Stinear TP, et al. A major
349 role for mammals in the ecology of *Mycobacterium ulcerans*. PLoS Negl Trop Dis. 2010;4(8):e791.
350 doi: 10.1371/journal.pntd.0000791. PubMed PMID: 20706592; PubMed Central PMCID:
351 PMC2919402.
- 352 18. Portaels F, Elsen P, Guimaraes-Peres A, Fonteyne PA, Meyers WM. Insects in the
353 transmission of *Mycobacterium ulcerans* infection. Lancet. 1999;353(9157):986. doi:
354 10.1016/S0140-6736(98)05177-0. PubMed PMID: 10459918.
- 355 19. Johnson PD, Azuolas J, Lavender CJ, Wishart E, Stinear TP, Hayman JA, et al.
356 *Mycobacterium ulcerans* in mosquitoes captured during outbreak of Buruli ulcer, southeastern
357 Australia. Emerg Infect Dis. 2007;13(11):1653-60. doi: 10.3201/eid1311.061369. PubMed PMID:
358 18217547; PubMed Central PMCID: PMC3375796.
- 359 20. Lavender CJ, Fyfe JA, Azuolas J, Brown K, Evans RN, Ray LR, et al. Risk of Buruli ulcer
360 and detection of *Mycobacterium ulcerans* in mosquitoes in southeastern Australia. PLoS Negl Trop
361 Dis. 2011;5(9):e1305. doi: 10.1371/journal.pntd.0001305. PubMed PMID: 21949891; PubMed
362 Central PMCID: PMC3176747.
- 363 21. Marsollier L, Andre JP, Frigui W, Reysset G, Milon G, Carbonnelle B, et al. Early
364 trafficking events of *Mycobacterium ulcerans* within *Naucoris cimicoides*. Cell Microbiol.
365 2007;9(2):347-55. doi: 10.1111/j.1462-5822.2006.00790.x. PubMed PMID: 16939536.
- 366 22. Marsollier L, Aubry J, Coutanceau E, Andre JP, Small PL, Milon G, et al. Colonization of
367 the salivary glands of *Naucoris cimicoides* by *Mycobacterium ulcerans* requires host plasmatocytes
368 and a macrolide toxin, mycolactone. Cell Microbiol. 2005;7(7):935-43. doi: 10.1111/j.1462-
369 5822.2005.00521.x. PubMed PMID: 15953026.

- 370 23. Marsollier L, Robert R, Aubry J, Saint Andre JP, Kouakou H, Legras P, et al. Aquatic
371 insects as a vector for *Mycobacterium ulcerans*. Appl Environ Microbiol. 2002;68(9):4623-8.
372 PubMed PMID: 12200321; PubMed Central PMCID: PMC124085.
- 373 24. Gordon CL, Buntine JA, Hayman JA, Lavender CJ, Fyfe JA, Hosking P, et al. All-oral
374 antibiotic treatment for buruli ulcer: a report of four patients. PLoS Negl Trop Dis.
375 2010;4(11):e770. doi: 10.1371/journal.pntd.0000770. PubMed PMID: 21152060; PubMed Central
376 PMCID: PMC2994921.
- 377 25. Trubiano JA, Lavender CJ, Fyfe JA, Bittmann S, Johnson PD. The incubation period of
378 Buruli ulcer (*Mycobacterium ulcerans* infection). PLoS Negl Trop Dis. 2013;7(10):e2463. doi:
379 10.1371/journal.pntd.0002463. PubMed PMID: 24098820; PubMed Central PMCID:
380 PMC3789762.
- 381 26. Andreu N, Zelmer A, Fletcher T, Elkington PT, Ward TH, Ripoll J, et al. Optimisation of
382 bioluminescent reporters for use with mycobacteria. PLoS One. 2010;5(5):e10777. doi:
383 10.1371/journal.pone.0010777. PubMed PMID: 20520722; PubMed Central PMCID:
384 PMCPMC2875389.
- 385 27. Omansen TF, Porter JL, Johnson PD, van der Werf TS, Stienstra Y, Stinear TP. In-vitro
386 activity of avermectins against *Mycobacterium ulcerans*. PLoS Negl Trop Dis. 2015;9(3):e0003549.
387 doi: 10.1371/journal.pntd.0003549. PubMed PMID: 25742173; PubMed Central PMCID:
388 PMCPMC4351077.
- 389 28. Hong H, Gates PJ, Staunton J, Stinear T, Cole ST, Leadlay PF, et al. Identification using
390 LC-MSn of co-metabolites in the biosynthesis of the polyketide toxin mycolactone by a clinical
391 isolate of *Mycobacterium ulcerans*. Chem Commun (Camb). 2003;(22):2822-3. PubMed PMID:
392 14651122.
- 393 29. Axford JK, Ross PA, Yeap HL, Callahan AG, Hoffmann AA. Fitness of wAlbB Wolbachia
394 Infection in *Aedes aegypti*: Parameter Estimates in an Outcrossed Background and Potential for

395 Population Invasion. Am J Trop Med Hyg. 2015. doi: 10.4269/ajtmh.15-0608. PubMed PMID:
396 26711515.

397 30. Fyfe JA, Lavender CJ, Johnson PD, Globan M, Sievers A, Azuolas J, et al. Development
398 and application of two multiplex real-time PCR assays for the detection of *Mycobacterium ulcerans*
399 in clinical and environmental samples. Appl Environ Microbiol. 2007;73(15):4733-40. doi:
400 10.1128/AEM.02971-06. PubMed PMID: 17526786; PubMed Central PMCID: PMC1951036.

401 31. Barnett HC, editor The incrimination of arthropods as vectors of disease. . Proceedings of
402 the 11th International Congress of Entomology; 1962.

403 32. Harris KF. An ingestion-egestion hypothesis of noncirculative virus transmission. In:
404 Mandahar L, editor. Plant Viruses. II. Boca Raton Inc., Fla: CRC Press; 1990. p. 177-204.

405 33. Wallace JR, Gordon MC, Hartsell L, Mosi L, Benbow ME, Merritt RW, et al. Interaction of
406 *Mycobacterium ulcerans* with mosquito species: implications for transmission and trophic
407 relationships. Appl Environ Microbiol. 2010;76(18):6215-22. doi: 10.1128/AEM.00340-10.
408 PubMed PMID: 20675453; PubMed Central PMCID: PMC2937476.

409 34. Hugh-Jones M, Blackburn J. The ecology of *Bacillus anthracis*. Mol Aspects Med.
410 2009;30(6):356-67. doi: 10.1016/j.mam.2009.08.003. PubMed PMID: 19720074.

411 35. Petersen JM, Mead PS, Schrieffer ME. *Francisella tularensis*: an arthropod-borne pathogen.
412 Vet Res. 2009;40(2):7. doi: 10.1051/vetres:2008045. PubMed PMID: 18950590; PubMed Central
413 PMCID: PMCPMC2695023.

414 36. Debacker M, Zinsou C, Aguiar J, Meyers WM, Portaels F. First case of *Mycobacterium*
415 *ulcerans* disease (Buruli ulcer) following a human bite. Clin Infect Dis. 2003;36(5):e67-8. doi:
416 10.1086/367660. PubMed PMID: 12594656.

417 37. Hofer M, Hirschel B, Kirschner P, Beghetti M, Kaelin A, Siegrist CA, et al. Brief report:
418 disseminated osteomyelitis from *Mycobacterium ulcerans* after a snakebite. N Engl J Med.
419 1993;328(14):1007-9. doi: 10.1056/NEJM199304083281405. PubMed PMID: 8450852.

- 420 38. Meyers WM, Shelly WM, Connor DH, Meyers EK. Human *Mycobacterium ulcerans*
421 infections developing at sites of trauma to skin. Am J Trop Med Hyg. 1974;23(5):919-23. PubMed
422 PMID: 4451232.
- 423 39. Barker DJ. Epidemiology of *Mycobacterium ulcerans* infection. Trans R Soc Trop Med
424 Hyg. 1973;67(1):43-50. PubMed PMID: 4798218.
- 425 40. Williamson HR, Mosi L, Donnell R, Aqqad M, Merritt RW, Small PL. *Mycobacterium*
426 *ulcerans* fails to infect through skin abrasions in a guinea pig infection model: implications for
427 transmission. PLoS Negl Trop Dis. 2014;8(4):e2770. doi: 10.1371/journal.pntd.0002770. PubMed
428 PMID: 24722416; PubMed Central PMCID: PMC3983084.
- 429 41. Bratschi MW, Ruf MT, Andreoli A, Minyem JC, Kerber S, Wantong FG, et al.
430 *Mycobacterium ulcerans* persistence at a village water source of Buruli ulcer patients. PLoS Negl
431 Trop Dis. 2014;8(3):e2756. doi: 10.1371/journal.pntd.0002756. PubMed PMID: 24675964;
432 PubMed Central PMCID: PMC3967953.
- 433 42. Carson C, Lavender CJ, Handasyde KA, O'Brien CR, Hewitt N, Johnson PD, et al. Potential
434 wildlife sentinels for monitoring the endemic spread of human buruli ulcer in South-East australia.
435 PLoS Negl Trop Dis. 2014;8(1):e2668. doi: 10.1371/journal.pntd.0002668. PubMed PMID:
436 24498452; PubMed Central PMCID: PMC3907424.
- 437 43. Johnson PD, Lavender CJ. Correlation between Buruli ulcer and vector-borne notifiable
438 diseases, Victoria, Australia. Emerg Infect Dis. 2009;15(4):614-5. doi: 10.3201/eid1504.081162.
439 PubMed PMID: 19331750; PubMed Central PMCID: PMC2671447.
- 440 44. Lavender CJ, Fyfe JA. Direct detection of *Mycobacterium ulcerans* in clinical specimens
441 and environmental samples. Methods Mol Biol. 2013;943:201-16. doi: 10.1007/978-1-60327-353-
442 4_13. PubMed PMID: 23104291.
- 443 45. Lavender CJ, Stinear TP, Johnson PD, Azuolas J, Benbow ME, Wallace JR, et al.
444 Evaluation of VNTR typing for the identification of *Mycobacterium ulcerans* in environmental

445 samples from Victoria, Australia. FEMS Microbiol Lett. 2008;287(2):250-5. doi: 10.1111/j.1574-
446 6968.2008.01328.x. PubMed PMID: 18754785.

447 46. Tobias NJ, Seemann T, Pidot SJ, Porter JL, Marsollier L, Marion E, et al. Mycolactone gene
448 expression is controlled by strong SigA-like promoters with utility in studies of *Mycobacterium*
449 *ulcerans* and buruli ulcer. PLoS Negl Trop Dis. 2009;3(11):e553. doi:
450 10.1371/journal.pntd.0000553. PubMed PMID: 19936295; PubMed Central PMCID:
451 PMC2775157.

452 47. Zogo B, Djenontin A, Carolan K, Babonneau J, Guegan JF, Eyangoh S, et al. A Field Study
453 in Benin to Investigate the Role of Mosquitoes and Other Flying Insects in the Ecology of
454 *Mycobacterium ulcerans*. PLoS Negl Trop Dis. 2015;9(7):e0003941. doi:
455 10.1371/journal.pntd.0003941. PubMed PMID: 26196901; PubMed Central PMCID:
456 PMCPMC4510061.

457 48. Roche B, Benbow ME, Merritt R, Kimbirauskas R, McIntosh M, Small PL, et al.
458 Identifying the Achilles' heel of multi-host pathogens: The concept of keystone "host" species
459 illustrated by transmission. Environ Res Lett. 2013;8(4):045009. doi: 10.1088/1748-
460 9326/8/4/045009. PubMed PMID: 24554969; PubMed Central PMCID: PMC3925833.

461 49. Quek TY, Henry MJ, Pasco JA, O'Brien DP, Johnson PD, Hughes A, et al. *Mycobacterium*
462 *ulcerans* infection: factors influencing diagnostic delay. Med J Aust. 2007;187(10):561-3. PubMed
463 PMID: 18021043.

464 50. Bratschi MW, Steinmann P, Wickenden A, Gillis TP. Current knowledge on *Mycobacterium*
465 *leprae* transmission: a systematic literature review. Lepr Rev. 2015;86(2):142-55. PubMed PMID:
466 26502685.

467 51. Shepard CC, McRae DH. *Mycobacterium leprae* in Mice: Minimal Infectious Dose,
468 Relationship between Staining Quality and Infectivity, and Effect of Cortisone. J Bacteriol.
469 1965;89:365-72. PubMed PMID: 14255702; PubMed Central PMCID: PMCPMC305516.

470 52. Leggett HC, Cornwallis CK, West SA. Mechanisms of pathogenesis, infective dose and
 471 virulence in human parasites. PLoS Pathog. 2012;8(2):e1002512. doi:
 472 10.1371/journal.ppat.1002512. PubMed PMID: 22359500; PubMed Central PMCID:
 473 PMC3280976.
 474
 475
 476
 477
 478
 479

Table 1: Summary of transmission experiments

Trauma source	Number of mice	Mouse tail coated	Number of mice bitten	Number of mice developing BU	Estimated dose (CFU)
Experiment 1: (Atypical mechanical transmission model, contaminated skin surface)					
<i>Aedes notoscriptus</i>	12 (4/mosquito cage)*	<i>M. ulcerans</i> (4.1x 10 ⁶ CFU/mL)	6	2 (#182, #191)	21
<i>Aedes notoscriptus</i>	4 (1/mosquito cage)	Media-only	2	0	-
Sterile needle (25G)	5	<i>M. ulcerans</i> (4.1x 10 ⁶ CFU/mL)	-	4 (#186, #200, #201, #202)	55
None	6	<i>M. ulcerans</i> (4.1x 10 ⁶ CFU/mL)	-	0	-
Experiment 2: (Atypical mechanical transmission model, contaminated skin surface)					
<i>Aedes aegypti</i>	5 (1/mosquito cage)	<i>M. ulcerans</i> (1.83 x 10 ⁶ CFU/mL)	5 [#]	0	9
<i>Aedes aegypti</i>	3 (1/mosquito cage)	Media-only	2	0	-
Sterile needle (25G)	5	<i>M. ulcerans</i> (1.83 x 10 ⁶ CFU/mL)	-	4 (#216, #217, #218, #219)	40
None	5	<i>M. ulcerans</i> (1.83 x 10 ⁶ CFU/mL)	-	0	-

Notes: *20 adult female mosquitoes per cage; [#]Multiple bites per mouse with 2 mice receiving 3 bites and 1 mouse receiving 2 bites; [§] “GE” is *M. ulcerans* genome equivalents as estimate based on IS2404 qPCR.

Supporting information Captions

Supplementary Table S1: IS2404 mosquito qPCR results