

1

2

3

4

5

6

7

Non-contact measurement of emotional and physiological changes in heart rate from a webcam

8

9

10 Christopher R. Madan^{1,3,†}, Tyler Harrison¹, & Kyle E. Mathewson^{1,2}

11

12

13 ¹ Department of Psychology, University of Alberta, Edmonton, AB, Canada

14 ² Neuroscience and Mental Health Institute, University of Alberta, Edmonton,
15 AB, Canada

16 ³ Department of Psychology, Boston College, Chestnut Hill, MA, USA

17

18

19

20

21

22

23 [†] Corresponding author.

24 Email address: madanc@bc.edu

25 Boston College, Department of Psychology,

26 McGuinn 300, 140 Commonwealth Ave.,

27 Chestnut Hill, MA, USA 02467

Abstract

Heart rate, measured in beats per minute (BPM), can be used as an index of an individual's physiological state. Each time the heart beats, blood is expelled and travels through the body. This blood flow can be detected in the face using a standard webcam that is able to pick up subtle changes in color that cannot be seen by the naked eye. Due to the light absorption spectrum of blood, we are able to detect differences in the amount of light absorbed by the blood traveling just below the skin (i.e., photoplethysmography). By modulating emotional and physiological stress—i.e., viewing arousing images and sitting vs. standing, respectively—to elicit changes in heart rate, we explored the feasibility of using a webcam as a psychophysiological measurement of autonomic activity. We found a high level of agreement between established physiological measures, electrocardiogram (ECG), and blood pulse oximetry, and heart rate estimates obtained from the webcam. We thus suggest webcams can be used as a non-invasive and readily available method for measuring psychophysiological changes, easily integrated into existing stimulus presentation software and hardware setups.

Keywords: heart rate; webcam; autonomic activity; emotion; arousal

Introduction

Heart rate (HR) is a readily measurable index of an individual's psychophysiological state, specifically autonomic arousal, used in addition to skin conductance response and pupil dilation (Bradley et al., 2008; Kahneman et al., 1969; Robinson et al., 1966). Indeed, the association between the heart and emotional/psychological states dates back to ancient Egypt (Damasio, 1994; Krantz & Falconer, 1997; Schacter & Singer, 1962), as well as permeating into culture throughout the ages (Loe & Edwards, 2004a, 2004b). HR is most often measured using an electrocardiogram (ECG), where changes in voltage generated by innervation of cardiac muscles producing a heartbeat are measured through electrode contacts that are affixed to an individual. However, ECG equipment can be costly, connections can deteriorate over time, and with some participant groups and situations it may be too invasive to apply electrodes. Other less invasive techniques to measure heart rate are therefore needed.

HR can be measured through methods alternative to ECG, such as photoplethysmography (PPG): the detection of variations in transmitted or reflected light (Ackles et al., 1985; Allen, 2007; Jennings et al., 1980; Lu et al., 2009; Schäfer & Vagedes, 2013). Briefly, changes in the light absorbed/reflected by blood can be used to measure the flow of blood. The absorption spectra of blood, and the measurement of the reflectance of skin color in relation to blood, has been studied for many decades within the field of medicine (e.g., Anderson & Parrish, 1981; Angelopoulou, 2001; Brunsting & Sheard, 1929a, 1929b; Edwards & Duntly, 1939; Horecker, 1943; Jakovels et al., 2010, 2011, 2012; Kim & Kim, 2006; Sheard & Brown, 1926; Brunsting & Sheard, 1929; Tsumura et al., 1999, 2000, 2003). A common example of transmission PPG is a pulse

oximeter (PulseOx) measurement in hospital settings in which red light is passed through the finger, wrist, or foot and fluctuations in transmitted light are detected.

More recently, a number of studies performed in biomedical engineering laboratories have demonstrated the feasibility of non-contact measuring of HR with a webcam (i.e., a digital video camera that streams its images to a computer). Poh et al. (2010) demonstrated the validity of HR measurements from a webcam by comparing them with measurements obtained at the same time from (but not time synchronized with) a blood pulse oximetry sensor (also see Kwon et al., 2012; Poh et al., 2011a). Subsequent studies have used webcams to study changes in HR due to exercise (Sun et al., 2011, 2012) and the development of devices designed to aid with health monitoring (Poh et al., 2011b; Verkrusye et al., 2008). There have been additional technical advances in how HR is estimated from the webcam recording (e.g., Lewandowska et al., 2011; Pursche et al., 2012; Sun et al., 2012). While these studies have been beneficial in demonstrating the robustness of this approach to measuring HR, the webcam HR estimates were not compared against time-synchronized standard HR measures, and did not evaluate changes in HR as a psychophysiological measure, i.e., the effect of task-related changes on autonomic arousal. As prior studies have indicated lower limits to the sampling rate required to assess ECG signal (Hejje & Roth, 2004; Pizzuti et al., 1985), it is not clear if the low sampling rate of the webcam will be suitable for measuring heart rate within the context of psychophysiology research.

To test if these techniques could be applied to experimental psychology situations as a method of psychophysiological monitoring, we used a standard webcam to record the light reflected from a participant's face. Acquisition of HR data from the webcam was

marked with respect to events in the stimulus presentation program, which are also marked in concurrently recorded ECG and PulseOx data. While averaging across the face area during recording of the webcam data, to provide anonymity, we measured task-related changes in a participant's HR. Specifically, we modulated emotional and physiological stress (i.e., viewing arousing images and sitting vs. standing, respectively) to elicit changes in HR to demonstrate the use of a webcam as a psychophysiological measurement of autonomic activity.

As a first test of event-related physiological changes in HR, we measured HR in a blocked sitting vs. standing task where we expected to observe large within-subject, task-related differences in HR. HR was measured concurrently from participants using the webcam along with ECG and pulse oximetry, for comparison. Briefly, when standing, the heart has to work harder to pump blood to the extremities to ensure sufficient force to overcome the effects of gravity (Caro et al., 1978; Herman, 2007; Rushmer, 1976). Empirically, the difference in HR for sitting vs. standing is approximately 8-10 BPM in young adults (Guy, 1837; MacWilliam, 1933; Schneider & Truesdell, 1922; also see Stein et al., 1966).

As a test of the feasibility of webcam HR in a task-related context, we next measured changes in HR time locked to emotional pictures, again concurrently with all three measures. Within the literature on emotional processing (e.g., Bradley et al., 2001a, 2008; Buchanan et al., 2006; Critchley et al., 2013; Garfinkel & Critchley, 2016; Lang et al., 1993; Levenson, 2003), it is well known that viewing emotionally arousing stimuli increases autonomic arousal, across a variety of psychophysiological measures. Presentation of unpleasant (i.e., negative valence) pictures elicits a deceleration in HR,

referred to as fear bradycardia, and that this deceleration is primarily mediated by the autonomic/parasympathetic nervous system (Bradley et al., 2001a, 2001b; Campbell et al., 1997). Hare (1973) suggested that this HR deceleration could be due to an orienting response, rather than a defensive response, to viewing the picture (also see Graham & Clifton, 1966; Sokolov, 1963). Empirically, this deceleration is a change of approximately 1-3 beats per minute (BPM), with a time course of approximately 6 seconds (Abercrombie et al., 2008; Bradley et al., 2008; Buchanan et al., 2006; Hare, 1973). Here we tested if our webcam HR technique would provide sufficient sensitivity to measure the subtle changes associated with a typical psychophysiological experiment, with the ECG and pulse oximetry data also acquired for comparison.

Method

Participants

A total of 24 volunteers participated in the experiment (age: $M=21.7$, range=18-25; 14 female) and were recruited from the University of Alberta community using advertisements around campus. Sample size was determined based on pilot studies of the sitting vs. standing task. All participants gave informed, written consent and were compensated at a rate of \$10/hr for their time. The experimental procedures were approved by an internal research ethics board of the University of Alberta.

Equipment

Video was recorded using a Logitech HD Pro Webcam C920 (Logitech International S.A., Newark, CA). The webcam video was recorded in color at a resolution of 640×480, at a mean sampling rate of 12 Hz (0.083 ± 0.016 s [$M \pm SD$] between video frames). Stimuli

were presented on a Dell UltraSharp 24" monitor with a resolution of 1920×1200, using a Windows 7 PC running MATLAB R2012b (The MathWorks Inc., Natick, MA) with the Psychophysics Toolbox v. 3 (Brainard, 1997). Webcam data was simultaneously recorded using in-house code in the same MATLAB script as the stimulus presentation.

ECG signals were collected from bilateral wrists of participants using Ag/AgCl snap-type disposable hydrogel monitoring electrodes (ElectroTrace ET101, Jason Inc., Huntington Beach, CA) in a bi-polar arrangement over the distal extent of the flexor digitorum superficialis muscle, with a ground over the distal extent of the left flexor carpi radialis. Prior to applying the electrodes, the participant's skin was cleaned using alcohol wipes. Blood pulse oximetry data was collected using a finger pulse sensor attached to the index finger of the participant's right hand and enclosed in a black light blocking sheath (Becker Meditec, Karlsruhe, Germany). Both sensors were connected to the AUX ports of a BrainVision V-Amp 16-channel amplifier (Brain Products GmbH, Gilching, Germany) using BIP2AUX converters. Physiological data was recorded at 500 Hz at 1.19 μ V/bit using BrainVision Recorder software (Brain Products GmbH) with a band-pass online filter between 0.628 and 30 Hz.

For the ECG and pulse oximetry data, data was collected for the entire duration of each task (sit-stand, emotion). In order to mark the time of stimulus onset in the ECG and pulse oximetry data, an 8-bit TTL pulse was sent via parallel port by the stimulus presentation software coincident with the onset of important stimuli, marking their time and identity (i.e., onset/offset of the fixation and pictures). The webcam data was recorded in epochs for each block (in the sit-stand task) or trial (in the emotion task) by the stimulus presentation software yoked to the stimulus display. The task presentation

and the data collection through all three measures were done by the same computer, allowing for all signals to be easily synchronized.

Stimuli

The pictures selected for the emotion task comprised four categories, each with 15 pictures/category. The pictures were selected from the International Affective Picture System (IAPS; Lang et al., 2008) database based on normative ratings for valence and arousal and were supplemented with pictures used in prior studies of emotional processing (Singhal et al., 2012; Wang et al., 2005, 2008). Mean IAPS valence/arousal scores (9-point scale, as described below) of the four categories were as follows: Neutral (Neut; 5.8/1.6), Low Arousal (Low; 3.6/3.3), Medium Arousal (Med; 2.3/5.8), and High Arousal (High; 2.3/6.1). A repeated-measures ANOVA showed that valence ratings for each category were significantly different from each adjacent category except for Med and High (i.e., $\text{Neut} > \text{Low} > \text{Med} = \text{High}$, $[F(3,72) = 132.97, p < .001]$). A repeated-measures ANOVA of arousal ratings showed that each category was significantly different from each adjacent category such that, $\text{Neut} < \text{Low} < \text{Med} < \text{High}$ $[F(3,72) = 150.59, p < .001]$. Pair-wise comparisons were Holm-Bonferroni-corrected.

Procedure

The experiment was conducted in a room of an experimental lab with normal lighting conditions. The experiment consisted of two tasks: blocks of sitting and standing (sit-stand task), and passive viewing of emotional and neutral pictures (emotion task). Task order was pseudorandomized across participants. In both cases, participants were seated in front of a webcam, which was placed either on a tripod (sit-stand task) or on top of the computer monitor (emotion task).

Sitting vs. standing task. The sit-stand task contained 10 blocks, of 30 s each. In half of the blocks, participants were instructed to be seated, in the other half they were to stand. The order of the blocks was pseudorandomized such that no more than two blocks from the same condition (e.g., sitting) occurred sequentially.

Before each block, the tripod was adjusted to suit the participant's height. The participant was then instructed to be as still as possible during the 30 s of data collection.

Emotional and neutral picture-viewing task. The emotion task was comprised of three blocks, each consisting of 20 trials. On each trial, participants were first shown a scrambled picture with a fixation cross (“+”) overlaid, followed by an emotional or neutral picture, then followed by the scrambled picture again. Pictures were presented for 2000 ms; scrambled stimuli were presented before and after each picture for 500 and 3000 ms, respectively. The scrambled stimuli were scrambled versions of the emotional or neutral picture, converted to grayscale and kept isoluminant with the picture. The order that the pictures were presented was pseudorandomized such that no more than two stimuli from the same category (e.g., high arousal) were shown sequentially. Trials were separated by jittered inter-trial intervals, ranging from 5000 to 6500 ms.

Prior to each block, the webcam recording was calibrated such that the participant aligned their head with a template indicating the area-of-interest (AOI) using live video feedback. Once the AOI was sufficiently aligned with the participant's face, they were instructed to place their hands on the table in front of them and to remain as still as possible while the stimuli were presented and data was recorded.

Data Analysis

The processing workflow for the webcam analyses is outlined in Figure 1. Based on the calibration, a rectangular AOI positioned over the participant's face constrains the collection of the webcam data. To ensure the collected data preserved participant anonymity, color values for each frame were averaged across this AOI during data collection, rather than maintaining the raw webcam frame. As a result, we only retained three intensity values per webcam frame, corresponding to red, green, and blue (RGB) channels. Data for each block (sit-stand task) or trial (emotion task) were then saved for offline analyses.




Figure 1. Illustration of the analysis pipeline.

Three pre-processing steps were used specifically on the continuous webcam data from entire blocks. First, to maximize the temporal resolution of the webcam data, we had sampled frames from the webcam as quickly as the hardware would allow (using the `videoinput` function in MATLAB), which lead to a non-uniform sampling rate. As minor fluctuations in the interval between successive frames would influence our estimated heart rate, we re-sampled the webcam data with a uniform interpolation of 12

Hz using the `interp` function in MATLAB. As a note to other researchers, if your hardware is able to sample from the webcam at a higher rate reliably, it would be simpler to instead have a uniform sampling rate and not necessitate re-sampling via interpolation. Second, it has been demonstrated that the green RGB color channel is the most sensitive to changes in light reflectance associated with oxygenated vs. deoxygenated blood, though the red and blue channels do still contain plethysmographic information (Lee et al., 2013; Poh et al., 2010; Sun et al., 2011, 2012; Verkrusse et al., 2008). To maximize info from all channels, we submitted the three color-channel time-series data (for the entire block) into a principal component analysis (PCA), allowing us to extract the variability in signal that was common across the three channels. We used the coefficients from the second principal component as our time-series data, as this was the component that corresponded to HR-related changes in all cases (also see Lewandowska et al., 2011; Poh et al., 2010, 2011a, 2011b; Pursche et al., 2012; Tsumura et al., 2000). Third, an additional offline Butterworth band-pass filter was applied to the data (high=0.8 Hz, low=3.0 Hz; see Gribok et al., 2011). This provided a 12-Hz signal from the webcam continuous throughout each block, along with the 500 Hz signals from the ECG and PulseOx.

Finally, for each each measure (webcam, ECG, PulseOx), the continuous data at submitted to a continuous wavelet (Morlet) transform implemented in the BOSC library (“Better OSCillation detection”; Hughes et al., 2012; Whitten et al., 2011). The transform was used to obtain the power spectra for the frequencies corresponding to a range of plausible heart rates, 50-140 BPM, in 1 BPM increments, and a wavelet number of 6. At

each time point of the resulting spectrogram, heart rate was calculated as the frequency with the highest power.

Blocked design. For the sitting vs. standing task, heart rate was estimated as a single value for each trial. Heart rate for each trial, for each measure, was estimated as the median heart rate for the 30-s block.

Event-related design. For the emotional and neutral picture-viewing task, heart rate was measured as a time-varying change, in relation to the onset of the image. To compute the event-related variations in HR, changes in HR were estimated using a sliding time-window. For each trial, epochs spanning from 5 s before to 5 s after the onset of the picture, were segmented from the continuous data.

Preliminary analyses indicated that the webcam data was confounded by stimulus luminance, where the luminance of the presented picture would interact with photoplethysmography signal intended to be recorded. This occurred despite pictures being preceded by an isoluminant scrambled picture; this likely occurred because trial-wise differences in the light emitted by the monitor when presenting the pictures influenced the light reflected by the participants' face and detected by the webcam. To address this confound, luminance for the pictures was regressed out of the individual trial timecourses. Luminance here was quantified by converting the pictures to CIE Lab 1976 color space, and summarized as a single value for each picture by averaging across the L* channel. For future research, we recommend matching the stimulus luminance across pictures if possible, making this regression step unnecessary. The presentation of the scrambled picture is critical, however, to prevent changes in screen luminance that correspond to the onset and offset of the picture-of-interest. We also recommend the

scrambled picture be presented in grayscale as color properties of the original pictures may not be matched across conditions (e.g., high arousing pictures were more red than neutral pictures).

For each trial and measure, the average heart rate in the 2000 ms prior to the picture onset was then subtracted from the entire trial period to align the picture onset across trials, i.e. a baseline correction. Then, for each HR recording type, separate averages are created for each subject in each of the emotional picture conditions. For statistical tests, the peak deceleration between 1500 and 3000 ms was used (based on prior findings; e.g., Abercrombie et al., 2008; Bradley et al., 2008; Buchanan et al., 2006), measured for each participant and emotion condition. See Figure 2 for a demonstration of the analysis pipeline for an event-related design.

Data quality. To ensure that the heart rate estimates obtained from the ECG and PulseOx data were sufficiently reliable, we excluded participants where the power at the peak frequency was less than twice the mean power in the sitting vs. standing task ($N=1$). ANOVA results are reported with Greenhouse-Geisser correction for non-sphericity where appropriate.




Figure 2. Demonstration of the analysis pipeline for an event-related design.

Results

Sitting vs. standing task

We first compared heart rate measurements for sitting vs. standing with each measurement method using a 2 [Posture: Sit, Stand] $\times$ 3 [Measure: ECG, Pulse Oximetry (PulseOx), Webcam] repeated-measures ANOVA, averaging across block. As shown in Figure 3A, we observed a main effect of Posture [$F(1,22)=85.29, p<0.001, \eta_p^2=0.80$], where standing was associated with a 10.4 BPM increase in heart rate relative to sitting. Neither the main effect of Measure [$F(1,28)=2.29, p=0.14, \eta_p^2=0.09$] nor the interaction [$F(2,42)=0.15, p=0.85, \eta_p^2=0.007$] were significant. Planned contrasts showed that the

effect of posture was observable using each measure individually [ECG: $t(22)=8.92$, $p<0.001$, Cohen's $d=0.82$, $M_{diff}=10.5$ BPM; PulseOx: $t(22)=7.84$, $p<0.001$, $d=0.82$, $M_{diff}=10.6$ BPM; Webcam: $t(22)=9.41$, $p<0.001$, $d=0.90$, $M_{diff}=10.2$ BPM].

To evaluate the agreement between the measurements more precisely, we additionally compared the heart-rate estimates from each block, i.e., 10 measurements per participant, between the three measures using correlations and Bland-Altman analyses. All three pairwise correlations were high and of similar magnitude [ECG–PulseOx: $r(458)=0.950$; ECG–Webcam: $r(458)=0.913$; PulseOx–Webcam: $r(458)=0.944$], as were the concordance correlation coefficients (Lin, 1989) [ECG–PulseOx: $r(458)=0.949$; ECG–Webcam: $r(458)=0.907$; PulseOx–Webcam: $r(458)=0.935$]. In all three cases, 2 SD of the difference between the compared measurements was approximately 10 BPM, as shown in Figures 3B-D [ECG–PulseOx: 9.19 BPM; ECG–Webcam: 11.91 BPM; PulseOx–Webcam: 9.67 BPM]. We did, however, observe a greater degree of bias when using the webcam, relative to the other measurements [ECG–PulseOx: -0.56 BPM; ECG–Webcam: 0.63 BPM; PulseOx–Webcam: 1.19 BPM]. This bias suggests that the webcam tends to slightly underestimate heart-rate estimates, perhaps due to the increased noise or slower sampling rate of the webcam measurement. Moreover, considering that certain participants are overrepresented in the outliers it is likely the case that some artifactual noise was impairing the ability to reliably determine the heart rate using some of the measures for these individuals. For instance, hair or clothes, as well as makeup, could interfere with the webcam measurement leading to unreliable estimates of HR on those blocks.

Task-related heart rate estimates from a webcam 16




Figure 3. Results from the sitting vs. standing task. (A) Mean heart rate for sitting and standing from each measure. Error bars represent SEM, corrected for inter-individual differences (within-subject SEM; Loftus & Masson, 1999). Bland-Altman plots for pairs of measures: (B) ECG-PulseOx, (C) ECG-Webcam, and (D) PulseOx-Webcam. Markers represent each block of the task from each participant. Markers in distinct colors represent individual participants; measurements from sitting blocks are shown as circles, standing blocks are shown as triangles.

Emotional and neutral picture-viewing task

As shown in Figure 4A-C the heart-rate decelerations for several of the conditions did not differ. Using the same stimuli in an fMRI study, Hrybouski et al. (2016) found that medium and high arousal stimuli were not distinct in behavioural ratings of emotional arousal or amygdala fMRI (BOLD) activity, and thus collapsed them together in their

reported analyses. Similarly, to maximally index the effect of the emotional pictures on heart rate, here we examined the mean response to the high and medium arousal picture conditions, compared to both the pre-stimulus baseline or viewing of the neutral pictures (Figure 3D). Thus, we pooled high and medium arousal images together and dropping the low arousal condition, as done in Hrybouski et al. (2016), as shown in Figure 4D-F.




Figure 4. Results from the emotional and neutral picture-viewing task. Event-related changes in heart rate in response to viewing each of the picture types, as measured by the (A) ECG, (B) PulseOx, and (C) Webcam. Shaded error bars represent within-subject SEM. The shaded time window (1500-3000 ms) depicts the data used in the statistical analyses. (D-F) Re-plots panels A-C, collapsing the High and Medium arousal conditions and removing the Low arousal condition. (G) Mean heart rate deceleration related to stimulus presentation, relative to the pre-stimulus baseline. Error bars represent SEM, corrected for inter-individual differences (within-subject SEM; Loftus & Masson, 1999).

We examine the heart-rate deceleration effects using a 2 [*Emotion*: High+Medium, Neutral] $\times$ 3 [*Measure*: ECG, Pulse Oximetry (PulseOx), Webcam] repeated-measures ANOVA, based on the mean heart rate during the analyzed window between 1500 and 3000 ms, relative to the pre-stimulus baseline (see Figure 4G). We observed a main effect of Emotion [$F(1,22)=7.94, p=0.010, \eta_p^2=0.23$], where the High+Medium pictures were associated with a 1.01 BPM decrease in heart rate relative to Neutral pictures. Neither the main effect of Measure [$F(1,23)=2.58, p=0.12, \eta_p^2=0.11$] nor the interaction [$F(1,24)=1.56, p=0.22, \eta_p^2=0.068$] were significant.

Despite the non-significant interaction, as planned contrasts we nonetheless report the HR effects for each measure. With the ECG data we observed a significant heart-rate deceleration of 1.71 BPM relative to the pre-stimulus baseline [$t(22)=4.40, p<0.001, d=0.96$], as well as a nominal deceleration of 0.44 BPM relative to viewing neutral pictures in the same window [$t(22)=0.83, p=0.42, d=0.28$]. The pulse oximetry data presented similar effects of viewing the emotional stimuli [relative to baseline: $t(22)=4.81, p<0.001, d=1.04$, 1.62 BPM deceleration; relative to neutral pictures: $t(22)=2.08, p=0.049, d=0.52$, 0.66 BPM deceleration]. With the webcam we observed a significant heart-rate deceleration of 3.33 BPM relative to the pre-stimulus baseline [$t(22)=4.37, p<0.001, d=0.95$], as well as a deceleration of 1.94 BPM relative to viewing neutral pictures in the same window [$t(22)=2.14, p=0.044, d=0.57$]. Thus, we observed significant heart-rate decelerations for emotional pictures with the pulse oximetry and webcam measures, but not with ECG. While the ECG and pulse oximetry obtained similar decelerations due to the arousing pictures, the ECG measure had slightly more variance in the effect (see Figures 4D and E).

It is not clear why the webcam is yielding pronounced, and narrower, heart-rate deceleration effects, particularly since it has less temporal resolution than the other two measures. It is possible that the webcam is measuring autonomic changes in addition to those related to photoplethysmography, such as effects of temperature (influencing skin vasculature) or face-specific responses such as emotion-related changes in facial expressions or blushing. Vasoconstrictive or vasodilative changes associated with sympathetic activity may have also contributed. Future research is needed to better understand how these other factors can influence HR estimates obtained from face recordings. These additional factors may also be responsible for the slight acceleration detected just prior to the deceleration (i.e., the peak at approximately 0.75s in Figure 4F).

Discussion

Heart rate can change in relation to psychological processes, in addition to physiological states. Here we demonstrated that a standard webcam can readily be used as a heart rate measurement device. Despite limitations in sampling rate, we were able to measure small heart-rate decelerations commonly associated with processing emotional pictures, in addition to the much larger changes in heart rate that are known to be associated with physiological state changes.

Our results showed very close agreement with conventional techniques measured simultaneously in both blocked and event-related designs. Differences in the webcam in the block design could largely be attributed to two outlier subjects for whom the webcam reliably underestimated their heart rate (HR). Therefore some individuals seem to be better conceal from the camera their on-going HR. We cannot investigate in the current data set further to determine what characteristics physically or behaviourally were

associated with these imprecisions (e.g., we only saved the webcam data for the face AOI, not the full webcam frame; did not collect inter-individual difference measures), but future work should better understand such individual differences in the measurement success.

Measuring non-contact physiological changes in HR over long periods of time as we showed in our sit-stand results provides an important tool by which one could, in real time, or on recorded footage, identify the ongoing HR of individuals under various levels of physical activity, or in various situations. The live video itself can even be modified to accentuate or visualize the pulse and heart rate on the body (Poh et al., 2011a).

The work here was intended to serve as a proof-of-principle that measurement of HR via webcam is sensitive enough for psychological studies. HR decelerations have been shown to index subsequent memory (Abercrombie et al., 2008; Buchanan et al., 2006; Cunningham et al., 2014; Fiacconi et al., 2016; Garfinkel et al., 2013; Jennings & Hall, 1980), task difficulty (Kahneman et al., 1969), interoceptive awareness (Garfinkel et al., 2013), and state anxiety (Garfinkel et al., 2014; Schachter & Singer, 1962). Heart rate is also known to be coupled to other physiological measures such as pupil dilation, skin conductance, and microsaccades (Bradley et al., 2008; Kahneman et al., 1969; Ohl et al., 2016). Consideration is needed to determine the applicability of this webcam approach, however, as it may not be suitable sensor of heart rate in all cases. For instance, heart-rate variability (HRV) has been associated with physiological well-being, and is related to a variety of factors including autonomic regulation and reactivity to acute stressors (e.g., Francis et al., 2015; Hallman et al., 2011; Shaffer et al., 2014). However, the current sampling rate of 12 Hz is insufficient, where HRV usually requires a sampling rate of 250

Hz or higher (Heijel & Roth, 2004; Pizzuti et al., 1985; Schäfer & Vagedes, 2013). Higher-end webcams or other video cameras, i.e., high-speed cameras, may be able to acquire data at a suitable sampling rate for HRV analyses, though testing will be necessary to determine other limiting factors, such as the rate of MATLAB's video I/O protocol. Further research is also necessary to establish the boundary conditions or other hardware limitations associated with future applications of this webcam approach to measuring HR, such as an index of vasculature function.

From a technical standpoint, measuring heart rate using a webcam can afford several benefits relative to the standard approaches such as ECG and pulse oximetry. While these other measures are non-invasive, a webcam is additionally non-contact. Thus, a webcam can be used equally well with participants that may have sensitive or delicate skin, such as older adults or patient populations, where contact measurements may be problematic. Furthermore, the impedance of the connection between the ECG electrode and the skin may increase over time leading to increased noise in ECG HR estimates. Pulse oximetry can similarly become dislodged over time due to its placement on the finger, and is cumbersome and interferes with normal typing and movements. Webcam equipment is also much more available and affordable than ECG and pulse oximetry, potentially making heart rate analyses more cost effective for pilot studies or researchers with limited funding.

A webcam may also be used to covertly measure heart rate with the participant being unaware that this data is even being collected, as long as proper consent and IRB protocols are followed. For instance, covert heart-rate recording could be beneficial along with a Concealed Information Test (see Matsuda et al., 2012, for a review). In this case, it

is additionally useful to point out that the webcam need not be calibrated towards the participants' face, but merely needs to record video data from exposed skin, e.g., an arm, in the presence of sufficient ambient lighting. Others have previously demonstrated that a single webcam can be used to measure heart rate for several individuals simultaneously (Poh et al., 2010). Additionally, the use of webcams to measure heart rate could be beneficial to medical care, such as when using video communication in patient care (see Armfield et al., 2012). Although animals may seem like unlikely candidates for such measurement, the exposed skin on the face and ears of mammals can also provide a non-invasive window into single or multiple animal HR monitoring.

One could argue that the usefulness of this technique is limited by the requirement of the subject to be still in the camera focus. Others have circumvented by using face detection algorithms (Poh et al., 2010, 2011b) or could take advantage of signal filters designed for detecting skin pigments (Anderson & Parrish, 1981; Changizi et al., 2006; Edwards & Duntly, 1939; Tsumura et al., 1999, 2003). If desired, multiple cameras and 3D motion trackers could be used to improve face/skin localization. Furthermore, movement artifacts are a similar problem for both ECG and PulseOx measurement. For experiment implementation, here we used the Psychophysics Toolbox and MATLAB. Functions within the Psychophysics Toolbox were used to present the stimuli while base MATLAB functions were used to interface with the webcam hardware. This allowed us to yolk webcam data recording to the stimulus presentation, but future studies could further integrate presentation and webcam recording for use with biofeedback (also see Lakens, 2013). In sum, here we demonstrated that the webcam is sufficiently sensitive for

psychologically relevant changes in heart rate, opening many potential lines of future research.

Acknowledgements

This work was supported by a NSERC discovery grant and startup funds from the Faculty of Science to KEM. CRM was supported by a fellowship from the Canadian Institutes of Health Research (FRN-146793).

References

- Abercrombie, H. C., Chambers, A. S., Greischar, L., & Monticelli, R. M. (2008). Orienting, emotion, and memory: Phasic and tonic variation in heart rate predicts memory for emotional pictures in men. *Neurobiology of Learning and Memory*, 90, 644–650. doi: 10.1016/j.nlm.2008.08.001
- Ackles, P. K., Jennings, J. R., & Coles, M. G. H. (1985). *Advances in Psychophysiology* (Vol. 1). Greenwich, CT: JAI.
- Allen, J. (2007). Photoplethysmography and its application in clinical physiological measurement. *Physiological Measurement*, 28, R1-R39.
- Anderson, R. R., & Parrish, J. A. (1981). The optics of human skin. *Journal of Investigative Dermatology*, 77, 13–19. doi:10.1111/1523-1747.ep12479191
- Angelopoulou, E. (2001). Understanding the color of human skin. *SPIE Proceedings*, 4299, 243. doi:10.1117/12.429495
- Armfield, N. R., Gray, L. C., & Smith, A. C. (2012). Clinical use of Skype: a review of the evidence base. *Journal of Telemedicine and Telecare*, 18, 125–127. doi:10.1258/jtt.2012.sft101
- Bradley, M. M., Codispoti, M., Cuthbert, B. N., & Lang, P. J. (2001). Emotion and motivation I: Defensive and appetitive reactions in picture processing. *Emotion*, 1, 276–298. doi: 10.1037/1528-3542.1.3.276
- Bradley, M. M., Codispoti, M., Sabatinelli, D., & Lang, P. J. (2001). Emotion and motivation II: Sex differences in picture processing. *Emotion*, 1, 300–319. doi:10.1037/1528-3542.1.3.300
- Bradley, M. M., Miccoli, L., Escrig, M. A., & Lang, P. J. (2008). The pupil as a measure of emotional arousal and autonomic activation. *Psychophysiology*, 45, 602–607. doi: 10.1111/j.1469-8986.2008.00654.x
- Brainard, D. H. (1997). The psychophysics toolbox. *Spatial Vision*, 10, 433–436. doi: 10.1163/156856897x00357
- Brunsting, L. A., & Sheard, C. (1929a). The color of the skin as analyzed by spectrophotometric methods: III. The rôle of superficial blood. *Journal of Clinical Investigation*, 7, 593–613. doi: 10.1172/jci100245
- Brunsting, L. A., & Sheard, C. (1929b). The color of the skin as analyzed by spectrophotometric methods: II. The rôle of pigmentation. *Journal of Clinical Investigation*, 7, 575–592. doi: 10.1172/jci100244
- Buchanan, T. W., Etzel, J. A., Adolphs, R., & Tranel, D. (2006). The influence of autonomic arousal and semantic relatedness on memory for emotional words. *International Journal of Psychophysiology*, 61, 26–33. doi:10.1016/j.ijpsycho.2005.10.022
- Campbell, B. A., Wood, G., & McBride, T. (1997). Origins of orienting and defensive responses: An evolutionary perspective. In P. J. Lang, R. F. Simons, & M. Balaban (Eds.), *Attention and orienting: Sensory and motivational processes* (pp. 41–67). New York: Lawrence Erlbaum.
- Caro, C., Pedley, T., & Schroter, R. (1978). *The mechanics of the circulation*. Oxford: Oxford University Press.
- Changizi, M. A., Zhang, Q., & Shimojo, S. (2006). Bare skin, blood and the evolution of primate colour vision. *Biology Letters*, 2, 217–221. doi:10.1098/rsbl.2006.0440
- Critchley, H. D., Eccles, J., & Garfinkel, S. N. (2013). Interaction between cognition, emotion, and the autonomic nervous system. *Handbook of Clinical Neurology*, 117, 59–77. doi: 10.1016/b978-0-444-53491-0.00006-7
- Cunningham, T. J., Crowell, C. R., Alger, S. E., Kensinger, E. A., Villano, M. A., Mattingly, S. M., & Payne, J. D. (2004). Psychophysiological arousal at encoding leads to reduced reactivity but enhanced emotional memory following sleep. *Neurobiology of Learning and Memory*, 114, 155–164. doi: 10.1016/j.nlm.2014.06.002
- Damásio, A. (1994). *Descartes' error: Emotion, reason, and the human brain*. New York: Putnam.
- Edwards, E. A., & Duntley, S. Q. (1939). The pigments and color of living human skin. *American Journal of Anatomy*, 65, 1–33. doi:10.1002/aja.1000650102

- Fiacconi, C. M., Peter, E. L., Owais, S., & Köhler, S. (2016). Knowing by heart: Visceral feedback shapes recognition memory judgments. *Journal of Experimental Psychology: General*, 145, 559–572. doi:10.1037/xge0000164
- Francis, H. M., Penglis, K. M., & McDonald, S. (2015). Manipulation of heart rate variability can modify response to anger-inducing stimuli. *Social Neuroscience*, 11, 545–552. doi: 10.1080/17470919.2015.1115777
- Garfinkel, S. N., Barrett, A. B., Minati, L., Dolan, R. J., Seth, A. K., & Critchley, H. D. (2013). What the heart forgets: Cardiac timing influences memory for words and is modulated by metacognition and interoceptive sensitivity. *Psychophysiology*, 50, 505–512. doi: 10.1111/psyp.12039
- Garfinkel, S. N., & Critchley, H. D. (2016). Threat and the body: How the heart supports fear processing. *Trends in Cognitive Sciences*, 20, 34–46. doi:10.1016/j.tics.2015.10.005
- Garfinkel, S. N., Minati, L., Gray, M. A., Seth, A. K., Dolan, R. J., & Critchley, H. D. (2014). Fear from the heart: Sensitivity to fear stimuli depends on individual heartbeats. *Journal of Neuroscience*, 34, 6573–6582. doi:10.1523/jneurosci.3507-13.2014
- Graham, F. K., & Clifton, R. K. (1966). Heart-rate change as a component of the orienting response. *Psychological Bulletin*, 65, 305–320. doi:10.1037/h0023258
- Gribok, A. V., Chen, X., & Reifman, J. (2011). A robust method to estimate instantaneous heart rate from noisy electrocardiogram waveforms. *Annals of Biomedical Engineering*, 39, 824–834. doi:10.1007/s10439-010-0204-2
- Guy, W. A. (1837). The effect produce upon the pulse by change of posture. *Guy's Hospital Reports*, 3, 92–110.
- Hallman, D. M., Olsson, E. M. G., von Schéele, B., Melin, L., & Lyskov, E. (2011). Effects of heart rate variability biofeedback in subjects with stress-related chronic neck pain: A pilot study. *Applied Psychophysiology and Biofeedback*, 36, 71–80. doi:10.1007/s10484-011-9147-0
- Hare, R. D. (1973). Orienting and defensive responses to visual stimuli. *Psychophysiology*, 10, 453–464.
- Heijel, L., & Roth, E. (2004). What is the adequate sampling interval of the ECG signal for heart rate variability analysis in the time domain? *Physiological Measurement*, 25, 1405–1411. doi:10.1088/0967-3334/25/6/006
- Herman, I. P. (2016). *Physics of the human body*. New York: Springer. doi: 10.1007/978-3-319-23932-3
- Horecker, B. L. (1943). The absorption spectra of hemoglobin and its derivatives in the visible and near infra-red regions. *Journal of Biological Chemistry*, 148, 173–183.
- Hrybowski, S., Aghamohammadi-Sereshki, A., Madan, C. R., Shafer, A. T., Baron, C. A., Seres, P., . . . Malykhin, N. V. (2016). Amygdala subnuclei response and connectivity during emotional processing. *NeuroImage*, 133, 98–110. doi:10.1016/j.neuroimage.2016.02.056
- Hughes, A. M., Whitten, T. A., Caplan, J. B., & Dickson, C. T. (2012). BOSC: A better oscillation detection method, extracts both sustained and transient rhythms from rat hippocampal recordings. *Hippocampus*, 22, 1417–1428. doi:10.1002/hipo.20979
- Jakovels, D., Kuzmina, I., Berzina, A., & Spigulis, J. (2012). RGB imaging system for monitoring of skin vascular malformation's laser therapy. *SPIE Proceedings*, 8427, 37. doi: 10.1117/12.922432
- Jakovels, D., Spigulis, J., & Rogule, L. (2011). RGB mapping of hemoglobin distribution in skin. *SPIE Proceedings*, 8087, 2B. doi:10.1117/12.889665
- Jakovels, D., Spigulis, J., & Saknite, I. (2010). Multi-spectral mapping of in vivo skin hemoglobin and melanin. *SPIE Proceedings*, 7715, 2Z. doi:10.1117/12.853928
- Jennings, J. R., & Hall, S. W. (1980). Recall, recognition, and rate: Memory and the heart. *Psychophysiology*, 17, 37–46. doi:10.1111/j.1469-8986.1980.tb02457.x
- Jennings, J.R., Tahmoush, A.J., & Redmond, D.P. (1980). Non-invasive measurement of peripheral vascular activity. In I. Martin and P.H. Venables (Eds.), *Techniques in Psychophysiology* (pp. 69-137). Wiley.

- Kahneman, D., Tursky, B., Shapiro, D., & Crider, A. (1969). Pupillary, heart rate, and skin resistance changes during a mental task. *Journal of Experimental Psychology*, 79, 164–167. doi:10.1037/h0026952
- Kim, D. H., & Kim, M.-J. (2006). Skin color analysis in HSV color space and rendering with fine scale skin structure. *Advances in Computer Graphics*, 254–264. doi:10.1007/11784203_22
- Krantz, D. S., & Falconer, J. J. (1997). Measurement of cardiovascular responses. In S. Cohen, R. C. Kessler, & L. U. Gordon (Eds.), *Measuring stress: A guide for health and social scientists* (pp. 193–212). New York: Oxford University Press.
- Kwon, S., Kim, H., & Park, K. S. (2012). Validation of heart rate extraction using video imaging on a built-in camera system of a smartphone. *Proceedings of the Annual International Conference of the IEEE Engineering in Medicine and Biology Society, 2012*, 2174–2177. doi: 10.1109/embc.2012.6346392
- Lakens, D. (2013). Using a smartphone to measure heart rate changes during relived happiness and anger. *IEEE Transactions on Affective Computing*, 4, 238–241. doi:10.1109/t-affc.2013.3
- Lang, P. J., Bradley, M. M., & Cuthbert, B. N. (2008). *International affective picture system (IAPS): Affective ratings of pictures and instruction manual* (Tech. Rep.). Gainesville, FL: University of Florida.
- Lang, P. J., Greenwald, M. K., Bradley, M. M., & Hamm, A. O. (1993). Looking at pictures: Affective, facial, visceral, and behavioral reactions. *Psychophysiology*, 30, 261–273. doi: 10.1111/j.1469-8986.1993.tb03352.x
- Lee, J., Matsumura, K., Ichi Yamakoshi, K., Rolfe, P., Tanaka, S., & Yamakoshi, T. (2013). Comparison between red, green and blue light reflection photoplethysmography for heart rate monitoring during motion. *Proceedings of the IEEE Annual Conference on Engineering in Medicine and Biology Society (EMBC), 2013*, 1724–1727. doi:10.1109/embc.2013.6609852
- Levenson, R. W. (2003). Autonomic specificity and emotion. In R. J. Davidson, K. R. Sherer, & H. H. Goldsmith (Eds.), *Handbook of affective sciences* (pp. 212–224). New York: Oxford University Press.
- Lewandowska, M., Rumiński, J., Kocejko, T., & Nowak, J. (2011). Measuring pulse rate with a webcam—a non-contact method for evaluating cardiac activity. *Proceedings of the Federated Conference on Computer Science and Information Systems (FedCSIS), 2011*, 405–410.
- Lin, L. I. (1989). A concordance correlation coefficient to evaluate reproducibility. *Biometrics*, 45, 255–268. doi: 10.2307/2532051
- Loe, M. J., & Edwards, W. D. (2004a). A light-hearted look at a lion-hearted organ (or, a perspective from three standard deviations beyond the norm) part 1 (of 2 parts). *Cardiovascular Pathology*, 13, 282–292. doi:10.1016/j.carpath.2004.05.001
- Loe, M. J., & Edwards, W. D. (2004b). A light-hearted look at a lion-hearted organ (or, a perspective from three standard deviations beyond the norm) part 2 (of 2 parts). *Cardiovascular Pathology*, 13, 334–340. doi:10.1016/j.carpath.2004.05.002
- Lu, G., Yang, F., Taylor, J. A., & Stein, J. F. (2009). A comparison of photoplethysmography and ECG recording to analyse heart rate variability in healthy subjects. *Journal of Medical Engineering & Technology*, 33, 634–641. doi: 10.3109/03091900903150998
- MacWilliam, J. A. (1933). Postural effects on heart-rate and blood-pressure. *Quarterly Journal of Experimental Physiology*, 23, 1–33. doi:10.1113/expphysiol.1933.sp000588
- Matsuda, I., Nittono, H., & Allen, J. J. B. (2012). The current and future status of the concealed information test for field use. *Frontiers in Psychology*, 3, 532. doi:10.3389/fpsyg.2012.00532
- Ohl, S., Wohltat, C., Kliegl, R., Pollatos, O., & Engbert, R. (2016). Microsaccades are coupled to heartbeat. *Journal of Neuroscience*, 36, 1237–1241. doi:10.1523/jneurosci.2211-15.2016
- Pizzuti, G., Cifaldi, S., & Nolfi, G. (1985). Digital sampling rate and ECG analysis. *Journal of Biomedical Engineering*, 7, 247–250. doi:10.1016/0141-5425(85)90027-5
- 10.1145/2048259.2048261

- Poh, M.-Z., McDuff, D. J., & Picard, R. W. (2010). Non-contact, automated cardiac pulse measurements using video imaging and blind source separation. *Optics Express*, 18, 10762. doi:10.1364/oe.18.010762
- Poh, M.-Z., McDuff, D. J., & Picard, R. W. (2011a). Advancements in noncontact, multiparameter physiological measurements using a webcam. *IEEE Transactions on Biomedical Engineering*, 58, 7–11. doi:10.1109/tbme.2010.2086456
- Poh, M.-Z., McDuff, D., & Picard, R. (2011b). A medical mirror for non-contact health monitoring. *ACM SIGGRAPH Emerging Technologies*, 2011, 2. doi:
- Pursche, T., Krajewski, J., & Moeller, R. (2012). Video-based heart rate measurement from human faces. *IEEE International Conference on Consumer Electronics (ICCE)*, 2012, 548–549. doi:10.1109/icce.2012.6161965
- Robinson, B. F., Epstein, S. E., Beiser, G. D., & Braunwald, E. (1966). Control of heart rate by the autonomic nervous system: Studies in man on the interrelation between baroreceptor mechanisms and exercise. *Circulation Research*, 19, 400–411. doi:10.1161/01.res.19.2.400
- Rushmer, R. F. (1976). *Cardiovascular dynamics* (4th ed.). Philadelphia: Saunders.
- Schachter, S., & Singer, J. (1962). Cognitive, social, and physiological determinants of emotional state. *Psychological Review*, 69, 379–399. doi:10.1037/h0046234
- Schneider, E. C., & Truesdell, D. (1922). A statistical study of the pulse rate and the arterial blood pressures in recumbency, standing, and after a standard exercise. *American Journal of Physiology*, 61, 429–474.
- Schäfer, A., & Vagedes, J. (2013). How accurate is pulse rate variability as an estimate of heart rate variability? *International Journal of Cardiology*, 166, 15–29. doi: 10.1016/j.ijcard.2012.03.119
- Shaffer, F., McCraty, R., & Zerr, C. L. (2014). A healthy heart is not a metronome: an integrative review of the heart's anatomy and heart rate variability. *Frontiers in Psychology*, 5. doi: 10.3389/fpsyg.2014.01040
- Sheard, C., & Brown, G. E. (1926). The spectrophotometric analysis of the color of the skin. *Archives of Internal Medicine*, 38, 816–831. doi:10.1001/archinte.1926.00120300133011
- Sheard, C., & Brunsting, L. A. (1929). The color of the skin as analyzed by spectrophotometric methods: I. Apparatus and procedures. *Journal of Clinical Investigation*, 7, 559–574. doi: 10.1172/jci100243
- Singhal, A., Shafer, A. T., Russell, M., Gibson, B., Wang, L., Vohra, S., & Dolcos, F. (2012). Electrophysiological correlates of fearful and sad distraction on target processing in adolescents with attention deficit-hyperactivity symptoms and affective disorders. *Frontiers in Integrative Neuroscience*, 6, 119. doi:10.3389/fnint.2012.00119
- Sokolov, E. N. (1963). Higher nervous functions: The orienting reflex. *Annual Review of Physiology*, 25, 545–580. doi:10.1146/annurev.ph.25.030163.002553
- Stein, E., Damato, A. N., Kosowsky, B. D., Lau, S. H., & Lister, J. W. (1966). The relation of heart rate to cardiovascular dynamics: Pacing by atrial electrodes. *Circulation*, 33, 925–932. doi:10.1161/01.cir.33.6.925
- Sun, Y., Papin, C., Azorin-Peris, V., Kalawsky, R., Greenwald, S., & Hu, S. (2011). Comparison of scientific CMOS camera and webcam for monitoring cardiac pulse after exercise. *SPIE Proceedings*, 8135, 6. doi:10.1117/12.893362
- Sun, Y., Papin, C., Azorin-Peris, V., Kalawsky, R., Greenwald, S., & Hu, S. (2012). Use of ambient light in remote photoplethysmographic systems: comparison between a high-performance camera and a low-cost webcam. *Journal of Biomedical Optics*, 17, 037005. doi:10.1117/1.jbo.17.3.037005
- Tsumura, N., Haneishi, H., & Miyake, Y. (1999). Independent-component analysis of skin color image. *Journal of the Optical Society of America A*, 16, 2169–2176. doi: 10.1364/josaa.16.002169
- Tsumura, N., Haneishi, H., & Miyake, Y. (2000). Independent component analysis of spectral absorbance image in human skin. *Optical Review*, 7, 479–482. doi:

674 10.1007/s10043-000-0479-x
675 Tsumura, N., Ojima, N., Sato, K., Shiraishi, M., Shimizu, H., Nabeshima, H., ... Miyake, Y. (2003).
676 Image-based skin color and texture analysis/synthesis by extracting hemoglobin and melanin
677 information in the skin. *ACM SIGGRAPH*, 22, 770–779. doi:
678 10.1145/1201775.882344
679 Verkrusse, W., Svaasand, L. O., & Nelson, J. S. (2008). Remote plethysmographic imaging using
680 ambient light. *Optics Express*, 16, 21434–21445. doi:10.1364/oe.16.021434
681 Wang, L., LaBar, K. S., Smoski, M., Rosenthal, M. Z., Dolcos, F., Lynch, T. R., ... McCarthy, G.
682 (2008). Prefrontal mechanisms for executive control over emotional distraction are altered in
683 major depression. *Psychiatry Research: Neuroimaging*, 163, 143–155. doi:
684 10.1016/j.psychresns.2007.10.004
685 Wang, L., McCarthy, G., Song, A. W., & LaBar, K. S. (2005). Amygdala activation to sad
686 pictures during high-field (4 Tesla) functional magnetic resonance imaging. *Emotion*, 5,
687 12–22. doi:10.1037/1528-3542.5.1.12
688 Whitten, T. A., Hughes, A. M., Dickson, C. T., & Caplan, J. B. (2011). A better oscillation
689 detection method robustly extracts EEG rhythms across brain state changes: The human
690 alpha rhythm as a test case. *NeuroImage*, 54, 860–874. doi:
691 10.1016/j.neuroimage.2010.08.064