

Title: Mapping Cortical Brain Asymmetry in 17,141 Healthy Individuals Worldwide via the ENIGMA Consortium

Abbreviated title: Mapping Cortical Brain Asymmetry

Authors: Xiang-Zhen Kong¹, Samuel Mathias², Tulio Guadalupe¹, Christoph Abé³, Ingrid Agartz^{4,5,6}, Theophilus N. Akudjedu⁷, Andre Aleman⁸, Saud Alhusaini^{9,10}, Nicholas B. Allen^{11,12}, David Ames^{13,14}, Ole A. Andreassen¹⁵, Nicola J. Armstrong¹⁶, Felipe Bergo¹⁷, Mark E. Bastin^{18,19}, Albert Batalla²⁰, Jochen Bauer²¹, Bernhard Baune²², Ramona Baur²³, Joseph Biederman^{24,25}, Sara K. Blaine²⁶, Premika Boedhoe^{27,28,29}, Erlend Bøen⁶, Anushree Bose³⁰, Janita Bralten^{31,32}, Daniel Brandeis^{33,34,35,36}, Silvia Brem^{33,34}, Henry Brodaty^{37,38}, Henrieke Bröhl³⁹, Jan Buitelaar^{33,40,41}, Christian Bürger⁴², Robin Bülow⁴³, Vince Calhoun^{44,45}, Anna Calvo^{46,47}, Erick Jorge Canales-Rodríguez^{48,49}, Jose M. Canive⁵⁰, Dara M. Cannon⁷, Elisabeth C. Caparelli⁵¹, Francisco X. Castellanos^{52,53}, Gianpiero L. Cavalleri⁹, Fernando Cendes¹⁷, Tiffany Moukbel Chaim-Avancini^{54,55}, Kaylita Chantiluke⁵⁶, Qun-lin Chen^{57,58}, Xiayu Chen⁵⁹, Yuqi Cheng⁶⁰, Anastasia Christakou⁶¹, Vincent P. Clark^{44,62}, David Coghill^{63,64}, Colm G. Connolly⁶⁵, Annette Conzelmann⁶⁶, Aldo Córdova-Palomera¹⁵, Janna Cousijn⁶⁷, Tim Crow⁶⁸, Ana Cubillo⁵⁶, Anders Dale^{69,70}, Udo Dannlowski⁴², Sara Ambrosino de Bruttopilo⁷¹, Patrick de Zeeuw⁷¹, Ian J. Deary⁷², Norman Delanty⁹, Damion V. Demeter⁷³, Adriana Di Martino⁵², Bruno Dietsche³⁹, N. Trung Doan¹⁵, Colin P. Doherty⁷⁴, Alysa Doyle^{75,76}, Margaretha Dramsdahl⁷⁷, Sarah Durston⁷¹, Eric Earl⁷³, Stefan Ehrlich⁷⁸, Carl Johan Ekman³, Torbjørn Elvsåshagen^{79,80}, Jeffery N. Epstein^{81,82}, Damien A. Fair^{83,84,85}, Stephen Faraone^{86,87}, Helena Fatouros-Bergman⁸⁸, Geraldo Busatto Filho^{54,55}, Lena Flyckt⁸⁸, Katharina Förster^{42,89}, John Foxe⁹⁰, Thomas Frodl^{91,92}, Paola Fuentes-Claramonte^{48,49}, Janice Fullerton^{93,94}, Hugh Garavan⁹⁵, Danielle do Santos Garcia¹⁷, Ian H. Gotlib⁹⁶, Anna E. Goudriaan^{97,98}, Hans Jörgen Grabe^{99,100}, Nynke A. Groenewold¹⁰¹, Dominik Grotegerd⁴², Oliver Gruber¹⁰², Tiril Gurholt⁴, Jan Haavik^{86,103}, Brandon Hager^{104,105}, Tim Hahn⁴², Narelle K. Hansell¹⁰⁶, Mathew A. Harris^{18,107}, Catharina Hartman¹⁰⁸, Maria del Carmen Valdés Hernández^{18,19}, Dirk Heslenfeld^{109,110}, Rob Hester¹¹¹, Beng-Choon Ho¹¹², Tiffany C. Ho^{65,96}, Pieter Hoekstra¹⁰⁸, Ruth J. van Holst^{113,114}, Martine Hoogman^{31,32}, Marie F. Høvik¹¹⁵, Fleur M. Howells¹¹⁶, Kenneth Hugdahl^{103,117}, Chaim Huyser^{118,119}, Martin Ingvar³, Lourdes Irwin¹²⁰, Akari Ishikawa¹⁷, Anthony James¹²¹, Terry Jernigan^{122,123}, Erik G E G Jönsson^{124,5}, Claas Kähler⁴², Vasily Kaleda¹²⁵, Clare Kelly^{126, 52, 127,128}, Michael Kerich¹²⁹, Anisha Keshavan¹³⁰, Sabin Khadka¹³¹, Tilo Kircher³⁹, Gregor Kohls¹³², Kerstin Konrad¹³², Ozlem Korucuoglu¹³³, Bernd Krämer¹⁰², Axel Krug³⁹, Jun Soo Kwon^{134,135}, Nanda Lambregts-Rommelse^{136,40}, Mikael Landén^{3, 137}, Luisa Lázaro^{138, 139,140,49}, Irina Lebedeva¹²⁵, Rhoshel Lenroot^{141,93,142}, Klaus-Peter Lesch^{143,144,145}, Qinquin Li¹⁴⁶, Kelvin

O. Lim¹⁴⁷, Jia Liu¹⁴⁶, Christine Lochner¹⁴⁸, Edythe D. London¹⁴⁹, Vera Lonning⁴, Valentina Lorenzetti¹⁵⁰, Michelle Luciano⁷², Maartje Luijten¹⁵¹, Astri J. Lundervold^{86,117}, Scott Mackey⁹⁵, Frank P. MacMaster^{152,153,154,155,156}, Sophie Maingault¹⁵⁷, Charles B. Malpas¹⁵⁸, Ulrik F. Malt^{159,160}, David Mataix-Cols⁸⁸, Andrew R. Mayer⁴⁴, Hazel McCarthy⁹², Michael P. Milham^{161,162}, Philip B. Mitchell^{141,163,164}, Bryon A. Mueller¹⁴⁷, Susana Muñoz Maniega^{18,19}, Bernard Mazoyer¹⁵⁷, Colm McDonald⁷, Quinn McLellan^{165,156}, Katie L. McMahon¹⁶⁶, Genevieve McPhilemy⁷, Reza Momenan¹²⁹, Angelica M. Morales¹⁴⁹, Janardhanan C. Narayanaswamy³⁰, José Carlos Vasques Moreira¹⁷, Stener Nerland⁶, Liam Nestor¹⁶⁷, Erik Newman¹²³, Joel T. Nigg¹²⁰, Jan-Egil Nordvik¹⁶⁸, Stephanie Novotny¹³¹, Eileen Oberwelland¹³², Ruth L. O’Gorman^{169,35}, Jaap Oosterlaan^{170,171,172}, Bob Oranje⁷¹, Catherine Orr⁹⁵, Bronwyn Owers⁹³, Paul Pauli²³, Martin Paulus^{173,174}, Kerstin Plessen^{86,175}, Georg G. von Polier¹³², Edith Pomarol-Clotet^{48,49}, Jiang Qiu^{57,58}, Joaquim Radua^{48,49,176,177}, Josep Antoni Ramos-Quiroga^{178,179}, Y.C. Janardhan Reddy³⁰, Andreas Reif¹⁸⁰, Gloria Roberts^{141,163}, Pedro Rosa^{54,55}, Natalie Royle¹⁹, Katya Rubia⁵⁶, Matthew D. Sacchet¹⁸¹, Perminder S. Sachdev^{37,182}, Raymond Salvador^{48,49}, Rocio Santos-Martin¹⁸³, Lianne Schmaal^{11,184,27}, Lisanne Schweren¹⁰⁸, Larry Seidman^{24,185}, Jochen Seitz¹⁸⁶, Mauricio Henriques Serpa^{54,55}, Philip Shaw^{187,188}, Timothy J. Silk^{158,189}, Alan N. Simmons^{190,191}, Egle Simulionyte¹⁰², Rajita Sinha²⁶, Zsuzsika Sjoerds^{192,193}, Runar Elle Smelror⁴, Joan Carlos Soliva¹⁷⁸, Nadia Solowij¹⁹⁴, Scott R. Sponheim¹⁹⁵, Dan J. Stein^{116,148}, Elliot A. Stein⁵¹, Michael Stevens^{196,197,198}, Lachlan T. Strike¹⁰⁶, Gustavo Sudre¹⁸⁷, Jing Sui^{44,199}, Leanne Tamm²⁰⁰, Hendrik S. Temmingh¹¹⁶, Robert J. Thoma^{201,202}, Alexander Tomyshhev¹²⁵, Giulia Tronchin⁷, Jessica Turner²⁰³, Anne Uhlmann^{148,116}, Theo G.M. van Erp²⁰⁴, Odile van den Heuvel^{27,28,29}, Dennis van der Meer⁴, Liza van Eijk²⁰⁵, Alasdair Vance²⁰⁶, Ilya M. Veer²⁰⁷, Dick J. Veltman²⁷, Ganesan Venkatasubramanian³⁰, Oscar Vilarroya^{178,208}, Yolanda Vives-Gilabert²⁰⁹, Henry Völzke^{210,211,212}, Susanne Walitza^{33,34,35}, Henrik Walter²⁰⁷, Esther Walton²¹³, Joanna M. Wardlaw^{19,214,215}, Wei Wen³⁷, Lars T. Westlye^{15,216}, Christopher D. Whelan⁹, Tonya White^{217,218}, Reinout W. Wiers²¹⁹, Margaret J. Wright^{106,166}, Katharina Wittfeld¹⁰⁰, Tony T. Yang⁶⁵, Clarissa L. Yasuda¹⁷, Yuliya Yoncheva⁵², Murat Yücel²²⁰, Je-Yeon Yun^{221,222}, Marcus Vinicius Zanetti^{54,55}, Zonglei Zhen⁵⁹, Xing-xing Zhu^{57,58}, Georg C. Ziegler¹⁴³, Kathrin Zierhut²³, Greig I. de Zubicaray²²³, David C. Glahn^{224,225}, Barbara Franke^{226,20}, Fabrice Crivello¹⁵⁷, Nathalie Tzourio-Mazoyer¹⁵⁷, Simon E. Fisher^{1,227}, Paul M. Thompson²²⁸, Clyde Francks^{1,227}

Affiliations:

1. Language and Genetics Department, Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands
2. Department of Psychiatry, Yale School of Medicine, New Haven, CT, USA

3. Department of Clinical Neuroscience, Osher Centre, Karolinska Institutet, Stockholm, Sweden
4. Norwegian Centre for Mental Disorders Research (NORMENT), K. G. Jebsen Centre for Psychosis Research, Institute of Clinical Medicine, University of Oslo, Oslo, Norway
5. Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden
6. Department of Psychiatric Research, Diakonhjemmet Hospital, Oslo, Norway
7. The Centre for Neuroimaging & Cognitive Genomics (NICOG), Clinical Neuroimaging Lab, NCBES Galway Neuroscience Centre, College of Medicine, Nursing, and Health Sciences, National University of Ireland Galway, H91 TK33 Galway Ireland, Republic of Ireland
8. BCN Neuroimaging Center, Department of Neuroscience, University Medical Center Groningen, University of Groningen, The Netherlands
9. Department of Molecular and Cellular Therapeutics, Royal College of Surgeons in Ireland, Dublin, Ireland
10. Neurology and Neurosurgery Department, Montreal Neurological Hospital and Institute, McGill University, Montreal, Canada
11. Orygen, The National Centre of Excellence in Youth Mental Health, Parkville, Australia
12. Department of Psychology, University of Oregon, Eugene OR, USA
13. National Ageing Research Institute, Melbourne, Australia
14. Academic Unit for Psychiatry of Old Age, University of Melbourne, Melbourne, Australia
15. Norwegian Centre for Mental Disorders Research (NORMENT), K.G. Jebsen Centre for Psychosis Research, Division of Mental Health and Addiction, Oslo University Hospital & Institute of Clinical Medicine, University of Oslo, Oslo, Norway
16. Mathematics and Statistics, Murdoch University, Perth, Australia
17. Laboratory of Neuroimaging, Department of Neurology, University of Campinas, Campinas, Brazil
18. Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK
19. Brain Research Imaging Centre, University of Edinburgh, Edinburgh, UK
20. Department of Psychiatry, Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Center, Nijmegen, The Netherlands
21. Department of Clinical Radiology, School of Medicine, University of Münster, Germany
22. Discipline of Psychiatry, University of Adelaide, Australia
23. Department of Psychology, University of Würzburg, Würzburg, Germany
24. Department of Psychiatry, Harvard Medical School, Boston, Mass, USA
25. Clinical and Research Programs in Pediatric Psychopharmacology and Adult ADHD, Massachusetts General Hospital, Boston, MA, USA
26. Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA
27. Department of Psychiatry, VU University Medical Center, Amsterdam, The Netherlands
28. Department of Anatomy & Neurosciences, VU University Medical Center, Amsterdam, The Netherlands
29. Amsterdam Neuroscience, Amsterdam, The Netherlands
30. Department of Psychiatry, National Institute of Mental Health and Neurosciences, Bengaluru, India

31. Department of Human Genetics, Radboud university medical center, Nijmegen, The Netherlands
32. Donders Institute for Brain, Cognition and Behaviour, Nijmegen, The Netherlands
33. University Clinics for Child and Adolescent Psychiatry (UCCAP), University of Zurich, Zurich, Switzerland
34. Neuroscience Center Zurich, University of Zurich and ETH Zurich, Switzerland
35. Zurich Center for Integrative Human Physiology, University of Zurich, Zurich, Switzerland
36. Department of Child and Adolescent Psychiatry and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim/Heidelberg University, J5, 68159 Mannheim, Germany
37. Centre for Healthy Brain Ageing, School of Psychiatry, UNSW Australia
38. Dementia Collaborative Research Centre ØC Assessment and Better Care, University of New South Wales, Sydney, Australia
39. Department of Psychiatry and Psychotherapy, Philipps-University Marburg, Germany
40. Karakter Child and Adolescent Psychiatry, Nijmegen, The Netherlands
41. Department of Cognitive Neuroscience, Radboud University, Nijmegen, The Netherlands
42. Department of Psychiatry, University of Münster, Germany
43. Department of Diagnostic Radiology and Neuroradiology, University Medicine Greifswald, Greifswald, Germany
44. The Mind Research Network, Albuquerque, NM, USA
45. Department of Electrical and Computer Engineering, University of New Mexico, Albuquerque, NM, USA
46. Medical Image Core Facility, August Pi I Sunyer Biomedical Research Institute (IDIBAPS), Barcelona, Spain
47. CIBERBBN
48. FIDMAG Germanes Hospitalaries Research Foundation, Barcelona, Spain
49. CIBERSAM, Centro de Investigación Biomédica en Red de Salud Mental, Spain
50. Departments of Psychiatry and Neurosciences, University of New Mexico, Albuquerque, NM, USA
51. Neuroimaging Research Branch, National Institute on Drug Abuse, National Institutes of Health, Baltimore, Maryland, USA
52. The Child Study Center at NYU Langone Medical Center, New York, USA
53. Division of Child and Adolescent Psychiatric Research, Nathan Kline Institute for Psychiatric Research, Orangeburg, NY, USA
54. Department of Psychiatry, Faculty of Medicine, University of São Paulo, São Paulo, Brazil
55. Center for Interdisciplinary Research on Applied Neurosciences (NAPNA), University of São Paulo, São Paulo, Brazil
56. King's College London, Institute of Psychiatry, Psychology and Neuroscience, Department of Child and Adolescent Psychiatry, London, UK
57. School of Psychology, Southwest University, Chongqing, China
58. Key Laboratory of Cognition and Personality, Ministry of Education, Chongqing, China

59. State Key Laboratory of Cognitive Neuroscience and Learning & IDG/McGovern Institute for Brain Research, Faculty of Psychology, Beijing Normal University, Beijing, China
60. Department of Psychiatry, First Affiliated Hospital of Kunming Medical University, Kunming, China
61. Department of Child Psychiatry, Institute of Psychiatry, King's College London, London WC2R 2LS, UK
62. Department of Psychology, University of New Mexico, Albuquerque, NM, USA
63. Departments of Paediatrics and Psychiatry, University of Melbourne, Victoria, Australia
64. Division of Neuroscience, Ninewells Hospital and Medical School, University of Dundee, Dundee, UK
65. Department of Psychiatry, Division of Child and Adolescent Psychiatry, and Weill Institute for Neurosciences, University of California, San Francisco, San Francisco, CA, USA
66. Department of Child and Adolescent Psychiatry and Psychotherapy, University of Tübingen, Tübingen, Germany
67. Department of Developmental Psychology, University of Amsterdam, Amsterdam, the Netherlands
68. SANE POWIC, University Department of Psychiatry, Warneford Hospital, Oxford, UK
69. Departments of Neurosciences and Radiology, University of California, San Diego, San Diego, CA, USA
70. UCSD Center for Translational Imaging and Precision Medicine, San Diego, CA, USA
71. NICHE-lab, Brain Center Rudolf Magnus, Department of Psychiatry, University Medical Center Utrecht, Utrecht, The Netherlands
72. Centre for Cognitive Ageing and Cognitive Epidemiology, Psychology, University of Edinburgh, Edinburgh, UK
73. Department of Behavioral Neuroscience at Oregon Health & Science University, Portland, OR, USA
74. Neurology Department, St. James's Hospital, Dublin, Ireland
75. Department of Psychiatry & Center for Human Genetics Research, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA
76. Stanley Center for Psychiatric Research at the Broad Institute, Cambridge, MA, USA
77. Region Zealand, Psychiatry, Roskilde, Denmark
78. Division of Psychological and Social Medicine and Developmental Neurosciences, Faculty of Medicine, Technische Universität Dresden, Dresden, Germany
79. Norwegian Centre for Mental Disorder Research (NORMENT), Institute of Clinical Medicine, University of Oslo, Oslo, Norway
80. Department of Neurology, Oslo University Hospital, Oslo, Norway
81. University of Cincinnati College of Medicine, Cincinnati, OH, USA
82. Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA
83. Department of Behavioral Neuroscience, Oregon Health & Science University, USA
84. Department of Psychiatry, Oregon Health & Science University, USA
85. Advanced Imaging Research Center, Oregon Health & Science University, USA
86. K.G. Jebsen Centre for Neuropsychiatric Disorders, Department of Biomedicine, University of Bergen, Bergen, Norway

87. Department of Psychiatry, SUNY Upstate Medical University, Syracuse, NY, USA
88. Department of Clinical Neuroscience, Centre for Psychiatry Research, Karolinska Institutet, Stockholm, Sweden
89. Institute of Psychiatric Phenomics and Genomics (IPPG), Ludwig-Maximilians-University, Munich, Germany
90. The Ernest J. Del Monte Institute for Neuroscience, Department of Neuroscience, University of Rochester School of Medicine and Dentistry, Rochester, NY, USA
91. Department of Psychiatry and Psychotherapy, University Hospital, Otto-von-Guericke-University, Magdeburg, Germany
92. Department of Psychiatry, Trinity College Dublin, Dublin, Ireland
93. Neuroscience Research Australia, Sydney, NSW, Australia
94. School of Medical Sciences, University of New South Wales, Sydney, New South Wales, Australia
95. Department of Psychiatry, University of Vermont, Burlington, VT, USA
96. Department of Psychology, Stanford University, USA
97. Department of Psychiatry, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands
98. Arkin Mental Health & Amsterdam Institute for Addiction Research, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands
99. Department of Psychiatry and Psychotherapy, University Medicine Greifswald, Germany
100. German Center for Neurodegenerative Diseases (DZNE), Rostock/Greifswald, Germany
101. Interdisciplinary Center Psychopathology and Emotion Regulation, University Medical Center Groningen, University of Groningen, The Netherlands
102. Section for Experimental Psychopathology and Neuroimaging, Department of General Psychiatry, Heidelberg University, Heidelberg, Germany
103. Department of Psychiatry, Haukeland University Hospital, Bergen, Norway
104. Massachusetts Mental Health Center, Boston, MA, USA
105. Division of Public Psychiatry, Beth Israel Deaconess Medical Center/Department of Psychiatry, Harvard Medical School, Boston, MA, USA
106. Queensland Brain Institute, University of Queensland, Brisbane, Australia
107. Division of Psychiatry, University of Edinburgh, Edinburgh, UK
108. University of Groningen, University Medical Center Groningen, Department of Psychiatry, Groningen, The Netherlands
109. Department of Cognitive Psychology, VU University Amsterdam, Amsterdam, The Netherlands
110. Department of Clinical Neuropsychology, VU University Amsterdam, Amsterdam, The Netherlands
111. School of Psychological Sciences, University of Melbourne, Melbourne, Australia
112. Department of Psychiatry, University of Iowa College of Medicine, Iowa City, Iowa, USA
113. Amsterdam Institute for Addiction Research, Academic Medical Center, University of Amsterdam,

Amsterdam, The Netherlands

114. Department of Psychiatry, Amsterdam, The Netherlands
115. Department of Clinical Medicine, University of Bergen, Bergen, Norway
116. Department of Psychiatry and Mental Health, University of Cape Town, South Africa
117. Department of Biological and Medical Psychology, University of Bergen, Bergen, Norway
118. De Bascule, Academic Center for Child and Adolescent Psychiatry, Amsterdam, the Netherlands
119. AMC, Department of Child and Adolescent Psychiatry, Amsterdam, the Netherlands
120. Department of Psychiatry, Oregon Health & Science University, Portland, OR, USA
121. University Department of Psychiatry, Warneford Hospital, Oxford, UK
122. Departments of Cognitive Science, Psychiatry, Radiology, University of California, San Diego, CA, USA
123. Center for Human Development, University of California, San Diego, CA, USA
124. Norwegian Centre for Mental Disorders Research (NORMENT), K.G. Jebsen Centre for Psychosis Research, Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway
125. Mental Health Research Center, Moscow, Russia
126. Department of Psychiatry, University of Dublin, Trinity College Dublin, Dublin, Ireland
127. School of Psychology, Trinity College, Dublin, Ireland
128. Trinity College Institute of Neuroscience, Dublin, Ireland
129. Clinical NeuroImaging Research Core, National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Baltimore, MD, USA
130. McGovern Institute for Brain Research and Poitras Center for Affective Disorders Research, Massachusetts Institute of Technology, Cambridge, Boston, USA
131. Olin Neuropsychiatry Research Center, Hartford CT, USA
132. Child Neuropsychology Section, Department of Child and Adolescent Psychiatry, University Hospital Aachen, Aachen, Germany
133. Department of Psychiatry, Washington University School of Medicine, St Louis, MO, USA
134. Department of Psychiatry, Seoul National University College of Medicine, Seoul, Republic of Korea
135. Department of Brain and Cognitive Sciences, Seoul National University College of Natural Sciences, Seoul, Republic of Korea
136. Department of Psychiatry, Radboud University Medical Center, Nijmegen, The Netherlands
137. Institute of Neuroscience and Physiology, The Sahlgrenska Academy at the Gothenburg University, Goteborg, Sweden
138. Department of Child and Adolescent Psychiatry and Psychology, Hospital Clínic, Barcelona, Spain
139. August Pi I Sunyer Biomedical Research Institute (IDIBAPS), Barcelona, Spain
140. Department of Medicine, University of Barcelona, Barcelona, Spain
141. School of Psychiatry, University of New South Wales, Sydney, NSW, Australia

142. University of New Mexico, Albuquerque, NM, USA
143. Division of Molecular Psychiatry, Center of Mental Health, University of Würzburg, Würzburg, Germany
144. Laboratory of Psychiatric Neurobiology, Institute of Molecular Medicine, I.M. Sechenov First Moscow State Medical University, Moscow, Russia
145. Department of Translational Neuroscience, School for Mental Health and Neuroscience (MHeNS), Maastricht University, Maastricht, The Netherlands
146. Beijing Key Laboratory of Applied Experimental Psychology, National Demonstration Center for Experimental Psychology Education (Beijing Normal University), Faculty of Psychology, Beijing Normal University, Beijing, China
147. Department of Psychiatry, University of Minnesota, Minneapolis, MN, USA
148. SU/UCT MRC Unit on Risk and Resilience in Mental Disorders, Department of Psychiatry, Stellenbosch University, South Africa
149. Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, CA, USA
150. Institute of Psychology Health and Society, University of Liverpool, Liverpool, UK
151. Behavioural Science Institute, Radboud University, Nijmegen, The Netherlands
152. Departments of Psychiatry and Pediatrics, University of Calgary, Calgary AB, Canada
153. Child and Adolescent Imaging Research Program, Alberta Children's Hospital, Calgary AB, Canada
154. Mathison Centre for Mental Health Research & Education, Hotchkiss Brain Institute, University of Calgary, Calgary AB, Canada
155. Strategic Clinical Network for Addictions and Mental Health, Alberta Health Services, Calgary AB, Canada
156. Alberta Children's Hospital Research Institute, Calgary, AB, Canada
157. Institut des Maladies Neurodégénératives, UMR 5293. Groupe d'Imagerie Neurofonctionnelle, CEA - CNRS - Université de Bordeaux, Bordeaux, France
158. Developmental Imaging, Murdoch Childrens Research Institute, Royal Children's Hospital, Melbourne, Australia
159. Department of Research and Education, Oslo University Hospital, Oslo, Norway
160. Institute of Clinical Medicine, University of Oslo, Oslo, Norway
161. Center for the Developing Brain, Child Mind Institute, New York, USA
162. Center for Biomedical Imaging and Neuromodulation, Nathan Kline Institute for Psychiatric Research, Orangeburg, NY, USA
163. Black Dog Institute, Prince of Wales Hospital, Randwick, NSW, Australia
164. Prince of Wales Hospital, Sydney, NSW, Australia
165. Department of Neuroscience, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada
166. Centre for Advanced Imaging, University of Queensland, Brisbane, Australia
167. Division of Brain Sciences, Imperial College London, London, UK
168. Sunnaas Rehabilitation Hospital HT, Nesodden, Norway
169. Center for MR-Research, University Children's Hospital, Zurich, Switzerland

170. Clinical Neuropsychology section, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
171. Emma Children's Hospital Academic Medical Center, Amsterdam, The Netherlands
172. Department of Pediatrics, VU Medical Center, Amsterdam, The Netherlands
173. Laureate Institute for Brain Research, Tulsa, Oklahoma, USA
174. Department of Psychiatry, University of California San Diego, La Jolla, California, USA
175. Child and Adolescent Mental Health Center, Capital Region, Denmark
176. Centre for Psychiatric Research and Education, Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden
177. Department of Psychosis Studies, Institute of Psychiatry, Psychology, and Neuroscience, King's College London, UK
178. Department of Psychiatry and Legal Medicine, Universitat Autònoma de Barcelona, Barcelona, Spain
179. Department of Psychiatry, Hospital Universitari Vall d'Hebron, CIBERSAM, Barcelona, Spain
180. Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, University Hospital Frankfurt, Frankfurt, Germany
181. Department of Psychiatry and Behavioral Sciences, Stanford University, USA
182. Neuropsychiatric Institute, Prince of Wales Hospital, Randwick, Australia
183. Department of Psychiatry and Psychology, University of Barcelona, Barcelona, Spain
184. Centre for Youth Mental Health, The University of Melbourne, Melbourne, Australia
185. Beth Israel Deaconess Medical Center, Boston, MA, USA
186. Department of Child and Adolescent Psychiatry, University Hospital Aachen, Aachen, Germany
187. Neurobehavioral Clinical Research Section, National Human Genome Research Institute, Bethesda, USA
188. National Institute of Mental Health, Bethesda, MD, USA
189. Department of Paediatrics, University of Melbourne, Melbourne, Australia
190. Department of Psychiatry, University of California, San Diego, La Jolla, CA, USA
191. Veterans Affairs San Diego Health Care System, La Jolla, CA, USA
192. Max Planck Institute for Human Cognitive and Brain Sciences, Department of Neurology, Leipzig, Germany
193. Leiden University, Institute of Psychology, Cognitive Psychology Unit & Leiden Institute for Brain and Cognition, Leiden, The Netherlands
194. School of Psychology and Illawarra Health and Medical Research Institute, University of Wollongong, Wollongong, Australia
195. Minneapolis VA Health Care System & University of Minnesota, Minneapolis, MN, USA
196. Clinical Neuroscience and Development Laboratory, Olin Neuropsychiatry Research Center, Hartford CT, USA
197. Child & Adolescent Research, Hartford Hospital/The Institute of Living, Hartford CT, USA
198. Department of Psychiatry, Yale University School of Medicine, Hartford CT, USA

199. National laboratory of Pattern Recognition, Institute of Automation, Chinese Academy of Sciences, Beijing, China

200. Department of Pediatrics, Division of Behavioral Medicine and Clinical Psychology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA

201. Department of Psychiatry and Behavioral Sciences, University of New Mexico, Albuquerque, NM, USA

202. Department of Psychiatry, University of New Mexico, Albuquerque, NM, USA

203. Department of Psychology and Neuroscience Institute, Georgia State University, Atlanta GA, USA

204. Department of Psychiatry and Human Behavior, University of California, Irvine, Irvine, USA

205. School of Psychology, University of Queensland, Brisbane, Australia

206. Academic Child Psychiatry Unit, Royal Children's Hospital, University of Melbourne, Melbourne, Victoria, Australia

207. Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Department of Psychiatry and Psychotherapy, Campus Mitte, Berlin, Germany

208. Fundació IMIM, Barcelona, Spain

209. INNDACYT, Barcelona, Spain

210. Institute for Community Medicine, University Medicine Greifswald, Germany

211. DZHK (German Centre for Cardiovascular Research), partner site Greifswald, Germany

212. German Centre for Diabetes Research (DZD), Site Greifswald, Germany

213. Department of Psychology, Georgia State University, Atlanta GA, USA

214. Scottish Imaging Network, A Platform for Scientific Excellence (SINAPSE) Collaboration, Edinburgh, UK

215. Centre for Clinical Brain Sciences, Centre for Cognitive Ageing and Cognitive Epidemiology, and UK Dementia Research Institute at The University of Edinburgh, Edinburgh, UK

216. Department of Psychology, University of Oslo, Oslo, Norway

217. Department of Child and Adolescent Psychiatry, Erasmus University Medical Centre, Rotterdam, Netherlands

218. Department of Radiology, Erasmus University Medical Centre, Rotterdam, Netherlands

219. Department of Psychology, University of Amsterdam, Amsterdam, the Netherlands

220. Monash Institute of Cognitive and Clinical Neurosciences and School of Psychological Sciences, Monash University, Melbourne, Australia

221. Seoul National University Hospital, Seoul, Republic of Korea

222. Yeongeon Student Support Center, Seoul National University College of Medicine, Seoul, Republic of Korea

223. Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Australia

224. Department of Psychiatry, Yale University, New Haven, CT, USA

225. Olin Neuropsychiatric Research Center, Hartford, CT, USA

226. Department of Human Genetics, Donders Institute for Brain, Cognition and Behaviour, Radboud University

Medical Center, Nijmegen, The Netherlands

227. Donders Institute for Brain, Cognition and Behavior, Radboud University, Nijmegen, The Netherlands

228. Imaging Genetics Center, Institute for Neuroimaging & Informatics, Keck School of Medicine of the University of Southern California, Los Angeles, CA, USA

Correspondence Authors:

Clyde Francks, Ph.D. (clyde.francks@mpi.nl)

Xiang-Zhen Kong, Ph.D. (xiangzhen.kong@outlook.com)

Number of Pages: 35

Number of Figures/Tables: 6/1

Number of words for Abstract: 195; **for Introduction:** 1506; **for Discussion:** 3186.

Total number of words: 11237

Abstract

Hemispheric asymmetry is a cardinal feature of human brain organization. Abnormal brain asymmetry is also linked to numerous cognitive and neuropsychiatric disorders. Here the ENIGMA consortium presents the largest ever analysis of cerebral cortical asymmetry and its variability across individuals. Cortical thickness and surface area were assessed in MRI scans of 17,141 healthy individuals from 99 datasets worldwide. Results revealed widespread asymmetries at both hemispheric and regional levels, with a generally thicker cortex but smaller surface area in the left hemisphere relative to the right. Regionally, asymmetries of cortical thickness and/or surface area were found in the inferior frontal gyrus, transverse temporal gyrus, parahippocampal gyrus, and entorhinal cortex. These regions are involved in lateralized functions, including language and visuospatial processing. In addition to population-level asymmetries, variability in the brain asymmetry was found related to sex, age, and brain size. Surprisingly, we did not find significant associations between asymmetries and handedness. Finally, with two independent pedigree datasets, we found several asymmetries showing considerable heritability. The structural asymmetries identified, and their variabilities and heritability provide a reference resource for future studies on the genetic basis of brain asymmetry and altered laterality in cognitive, neurological, and psychiatric disorders.

Keywords: brain asymmetry; lateralization; cortical thickness; surface area; meta-analysis

Introduction

Understanding the functional specialization of the cerebral hemispheres is a long-standing and central issue in human neuroscience research. At the population-level, hemispheric asymmetry, or lateralization, is involved in various perceptual and cognitive functions, including language (Vigneau et al., 2006; Vigneau et al., 2011), face processing (Badzakova-Trajkov et al., 2010; Willems et al., 2010; Zhen et al., 2015), visuospatial processing (Badzakova-Trajkov et al., 2010; Zago et al., 2017; Zhen et al., 2017), and reasoning (Fornito et al., 2004; Turner et al., 2015), as well as handedness (Steinmetz et al., 1991). For example, language lateralization involves leftward dominance for various processes involved in speech perception and production in most people (Vigneau et al., 2006; Vigneau et al., 2011). Moreover, altered hemispheric lateralization has been associated with numerous cognitive and neuropsychiatric disorders, including dyslexia (Altarelli et al., 2014), Alzheimer's disease (Thompson et al., 1998), attention-deficit/hyperactivity disorder (ADHD) (Shaw et al., 2009), psychotic disorders (Crow, 1990; Schobel et al., 2009; Yucel et al., 2002; Yucel et al., 2003), autism (Eyler et al., 2012) and mood disorders (Liu et al., 2016; Yucel et al., 2009). Various aspects of brain asymmetry, including anatomical asymmetries of perisylvian language-related cortical regions, appear *in utero* during the second trimester of gestation (Chi et al., 1977; Hering-Hanit et al., 2001). Thus, brain laterality is likely to be under the control of genetic-developmental programs which are inherently lateralized, such as those that have been described for the left-right visceral axis (affecting the placement of the heart, lungs etc.) (Concha et al., 2012; Levin, 2005). Together, these observations indicate that asymmetry is a core element of the brain's usual organization, which is required for optimal functioning and influenced by genetic factors.

Although structural and functional asymmetries are likely to be interrelated in the typically lateralized human brain, the nature of structure-function relations are far from clear. For example, it is still not understood whether anatomical asymmetries around the Sylvian fissure are an important aspect of left-hemisphere language dominance (Marie et al., 2015; Tzourio-Mazoyer and Mazoyer, 2017). Furthermore, variations in structural and functional asymmetry have been reported to correlate poorly (Bishop, 2013; Dorsaint-Pierre et al., 2006; Greve et al., 2013; Guadalupe et al., 2014), which

further complicates assessment of the structure-function relations and dependencies. The literature has, however, been hampered by generally small sample sizes and heterogeneous methods for assessing asymmetries and their variabilities (Guadalupe et al., 2016), leading to confusion about which structures are actually anatomically asymmetrical at the population level, and to what degrees (see below). This has also been the case for asymmetry-disorder studies. In this context, and as motivation for the present study, it is important to characterize anatomical asymmetries in a large sample of healthy individuals, in order to provide a definitive and normative reference for future studies of hemispheric specialization in both healthy and clinical populations.

One aspect of structural asymmetry in the human brain is “Yakovlevian torque”, an overall hemispheric twist giving rise to the frontal and occipital petalia, which describes protrusions of the right frontal and left occipital regions over the midline (Narr et al., 2007; Watkins et al., 2001; Yakovlev, 1972). At a regional level, later studies that applied computational methods to MRI data mainly focused on volumetric measures of cortical structures and revealed both replicable and inconsistent findings of asymmetries. For example, Goldberg et al. (2013) summarized in their study that regions implicated in visual processing show rightward volumetric asymmetries, while, in contrast, somatosensory, auditory, and parts of the premotor cortices show leftward volumetric asymmetries (Goldberg et al., 2013). One recent study replicated this distribution of regional asymmetries, especially in the lateral view (Esteves et al., 2017), but several studies have shown quite different asymmetry results (Good et al., 2001; Herve et al., 2006; Watkins et al., 2001). For example, Goldberg et al. (2013) and Esteves et al. (2017) found a greater superior frontal volume in the left hemisphere, while Watkins et al. (2001) found greater superior frontal volume in the right.

Cortical volume is, by definition, a product of two distinct aspects of the brain, i.e. cortical thickness and surface area (Meyer et al., 2014; Raznahan et al., 2011); researchers have also attempted to assess the asymmetries of cortical thickness and surface area separately, using neuroimaging surface-based approaches (Fischl et al., 2002; Van Essen, 2004). Regarding cortical thickness, a number of studies have found mixed results for asymmetry patterns. For example, Luders et al. (2006) found greater left-sided thickness in parts of the cingulate, precentral gyrus, orbital frontal gyrus, and

temporal and parietal lobes, and greater right-sided thickness in the inferior frontal gyrus (Luders et al., 2006). However, other studies (Koelkebeck et al., 2014; Maingault et al., 2016; Plessen et al., 2014; Zhou et al., 2013) revealed somewhat inconsistent patterns of thickness asymmetry. For instance, Zhou et al. (2013), studying individuals of an age range similar to that in Luders et al. (2006), did not find leftward asymmetry in the precentral gyrus, but revealed a strong rightward asymmetry in the lateral parietal and occipital regions. For an overview of mixed results of asymmetry patterns observed in previous studies, please refer to Figure S1 in Supplemental Information. Regarding regional surface area asymmetries, some repeatable findings have been found for the supramarginal gyrus (leftward) (Koelkebeck et al., 2014; Lyttelton et al., 2009; Maingault et al., 2016), the middle temporal gyrus (rightward) (Koelkebeck et al., 2014; Maingault et al., 2016), and the anterior cingulate gyrus (rostral: leftward; caudal: rightward) (Koelkebeck et al., 2014; Maingault et al., 2016). However, there are also many inconsistent results across studies, such as for the lateral occipital cortex, which showed a strong rightward asymmetry in Chiarello et al. (2016) (Chiarello et al., 2016), but leftward asymmetry in Koelkebeck et al. (2016) (see a summary in (Kang et al., 2015)). These mixed results of brain structural asymmetry may reflect differences in many factors, including statistical power and confidence intervals related to sample sizes as well as differences in scanning, brain segmentation, and parcellation methods. Thus, a large-scale survey using harmonized approaches is needed to give a clearer picture of the lateralization in the human brain.

Another potential source for the mixed results in the literature is variability across individuals and in relation to factors like age and sex (Hogstrom et al., 2013; Im et al., 2006; Salat et al., 2004). For example, a recent study has observed that males show, on average, more pronounced gray matter volume asymmetries in superior temporal language regions than females (Guadalupe et al., 2015). Changes in structural asymmetries with age have also been reported (Raz et al., 2004; Shaw et al., 2009), but not consistently (Zhou et al., 2013). Another potential factor linked to brain lateralization is handedness, although the associations are very weak as reported (Guadalupe et al., 2014; Maingault et al., 2016; Willems et al., 2014). For example, with more than 100 left-handed participants and roughly 2000 right-handed participants, Guadalupe et al. (2014) suggested an association of handedness with

the surface area of the left precentral sulcus, but this was not significant after multiple testing adjustment. In addition, greater cortical asymmetry has been observed in participants with larger overall brain size (Kang et al., 2015). Thus, the existing literature on variability in brain structural asymmetries suggests influences of individual differences in age, sex, handedness, and brain size, but again a large-scale study is needed to clarify the nature of any such relations. The largest previous studies of brain asymmetries were conducted by Plessen et al. (2014) and Zhou et al. (2013) in relation to sex and age in sample sizes of 215 and 274 participants, and Maingault et al. (2016) in a sample size of 250 (120 left-handers) in relation to handedness. Each of these studies used different methodological approaches. Thus, a large-scale study of thousands of participants would be a major step forward in achieving a more accurate description of the typical asymmetries of the human brain, as well as variation in these asymmetries and some key biological factors which affect them.

The ENIGMA consortium provides the opportunity for large-scale meta-analysis studies of brain anatomy based on tens of thousands of participants with structural MRI data (Thompson et al., 2014). In this study, we present the largest analysis of structural asymmetries in the human cerebral cortex, with MRI scans of 17,141 healthy individuals from 99 datasets worldwide, in a harmonized multi-site study using meta-analytic methods. Our aim was to identify cortical regions that consistently show asymmetry with regard to either cortical thickness or surface area, to provide a clear picture of population-level asymmetries in the human brain. We also assessed potential influences of age, sex, handedness, and brain size on the variability in asymmetries, as well as of the methodological factor of MRI scanner field strength. Furthermore, as a first step towards elucidating the genetic basis of variability in brain asymmetry, we further analyzed two independent pedigree datasets, i.e. the Genetics of Brain Structure (GOBS; N = 1,443) and Human Connectome Project (HCP; N = 1,113) datasets, to estimate heritability of the asymmetry measures.

Methods

Datasets

The primary datasets used in this study for large-scale meta-analysis were from members of the

Lateralization Working Group within the ENIGMA Consortium (Thompson et al., 2014). There were 99 independent samples, including 17,141 healthy participants from diverse ethnic backgrounds. Samples were drawn from the general population or were healthy controls from clinical studies. Figure 1 summarizes the age ranges and sample sizes of each dataset (for more details on each sample, see Supplemental Information S1). All local institutional review boards permitted the use of extracted measures of the completely anonymized data.




Figure 1. The age ranges and sizes of each sample. Each line covers the age range of an individual sample, with different colors indicating the sample sizes (see color key in figure). The position of the gray/black dot on each line indicates the median age of that sample. Black dots indicate samples with handedness information available. For more details, see Supplemental Information S1.

Handedness was known for a subset of the participants. The method of assessment varied across samples (see Supplemental Information S2). An ambidextrous category was not included, resulting in 827 left-handed and 11,237 right-handed participants in total.

Two additional datasets were used to estimate heritability of asymmetry measures, i.e. the GOBS

dataset and the HCP dataset. GOBS is a family study comprising 1,443 individuals with MRI data (836 females), aged between 18 and 85 years at the time of scanning (McKay et al., 2014). All GOBS subjects are Mexican Americans and belong to pedigrees of varying sizes (the largest pedigree has 143 members). The HCP is a large-scale project comprising 1,113 individuals with MRI data (606 females, age range 22-37 years at the time of scanning) of varying ethnicities (<http://humanconnectome.org/>). The HCP contains 143 monozygotic twin pairs and 85 dizygotic twin pairs, as well as unrelated individuals.

Image Acquisition and Processing

Structural T1-weighted MRI scans were acquired and analyzed locally. Images were acquired at different field strengths (1.5 T and 3 T). The images were analyzed using the fully automated and validated segmentation software FreeSurfer (version 5.3 for 91 of 99 samples). Segmentations of 68 (34 left and 34 right) cortical regions based on the Desikan-Killiany atlas (Desikan et al., 2006) were statistically evaluated and in some cases visually inspected for outliers. Regional measures of cortical thickness and surface area were extracted for each participant. In addition, two hemisphere-level measures (average cortical thickness and total surface area), as well as the intracranial volume (ICV) were obtained. Quality control and data analysis were performed following standardized ENIGMA protocols (see <http://enigma.ini.usc.edu/protocols/imaging-protocols/>). In addition, following Guadalupe et al. (2016), we performed several checks to assess potential errors in the left-right orientation of the data. Further details for the datasets and orientation checking can be found in Supplemental Information S3.

Within-dataset Analyses

For each dataset, descriptive and statistical analyses of the asymmetries in both cortical thickness and surface area were performed at each participating site using a single script in the R language, based on unified table-formatted data. For each global hemispheric or regional measure, an asymmetry index (AI) was defined as $(L-R)/((L+R)/2)$, where L and R are the corresponding thickness or area

measures on the left and right hemisphere, respectively. Thus, positive and negative AI values indicate leftward and rightward asymmetry, respectively. To exclude possible outliers in measures of cortical thickness, surface area, or AIs, we followed Guadalupe et al. (2016) and used an adaptive threshold (SD_{thre}) depending on each dataset's sample size: $N < 150$, $SD_{thre} = 2.5$; $150 \leq N \leq 1000$, $SD_{thre} = 3$; $N > 1000$, $SD_{thre} = 3.5$.

Statistical tests were run for each hemisphere-level or regional measure separately. Paired t-tests were used to assess inter-hemispheric differences. Cohen's d was calculated based on each paired t-test result to estimate the effect size of population-level asymmetry. All differences between sexes (-1=females, 1=males) were assessed with linear regression models adjusted for age, age², and brain size (as indexed by ICV). Cohen's d was calculated to estimate the effect size for each comparison. Furthermore, we examined the age effects on the AIs in cortical thickness and surface area, adjusting for sex and ICV. Similarly, we examined associations between ICV and the AIs in cortical thickness and surface area, adjusting for age, age², and sex. In addition, if handedness information was available, AI differences between handedness groups (-1=left; 1=right) were assessed with linear regression models adjusted for all the other covariates. For each analysis above, additional covariates of scanners were included when more than one scanner was used at one site.

Meta-analyses

All regression models and effect size estimates were fitted at each participating site separately. We then combined the output statistics from each dataset using inverse variance-weighted random-effect meta-analyses (Borenstein et al., 2010) with the R package *metafor*, version 1.9-9. This method tests one overall effect, while weighting each dataset's contribution by the inverse of its corresponding sampling variance. Thus, unlike fixed-effect meta-analysis, this method takes into account variability across different studies.

A Cohen's d effect size estimate of population-level asymmetry (hemispheric difference) was obtained using a random-effect meta-analysis model for each region and each cortical measure (cortical thickness and surface area). Note that including results based on too few participants may

reduce reliability, and therefore we only included datasets with a sample size larger than 15. In the meta-analysis, heterogeneity of each effect was assessed via the I^2 value, which describes the percentage of total variation across studies that is due to heterogeneity rather than chance. I^2 values of 25%, 50%, and 75% indicate a low, moderate, and high heterogeneity, respectively.

Similarly, Cohen's d estimates of sex and handedness effects on asymmetry were obtained by meta-analysis for each cortical region. Again we only included samples with at least 15 participants per group. For meta-analyses of the correlations of asymmetries with age or ICV, predictors were treated as continuous variables, so that effect sizes were expressed as partial-correlation r . For meta-analyses of age effects, we only included samples with a minimum 5-year range in the initial analysis, and in a subsequent analysis we further restricted to samples with age ranges larger than 20 years, to better capture the age effects in our data.

In this study, we report uncorrected p values with a significance threshold determined by Bonferroni correction for multiple comparisons within each separate meta-analysis (i.e. correction separately for analysis of 34 regional surface area asymmetries and 34 regional thickness asymmetries, $p = 0.05/34 = 0.00147$). No correction was done for global hemispheric measures of asymmetry.

Moderator analyses with meta-regression

Meta-regressions were performed to evaluate the potential moderating effects on meta-analysis effect sizes. We tested whether moderating factors, including median age, median ICV, sex ratio, handedness ratio, and MRI scanner field strength (3T, $N = 63$ datasets versus 1.5T, $N = 28$ datasets), influenced the effect size estimates across datasets in the meta-analyses. Each moderator variable was separately included as a fixed effect predictor in the meta-regression model. All statistical analyses were conducted using the R software package *metafor*, and Bonferroni correction was applied for multiple comparisons ($p < 0.00147$; see above).

Heritability estimation

Furthermore, we estimated the heritability of asymmetries in cortical thickness and surface area,

firstly using the GOBS dataset ($N = 1,443$; see ‘Datasets’ above). For more details about this cohort, see McKay et al. (2014). Specifically, we estimated the narrow-sense heritability (i.e., the proportion of overall variance explained by additive genetic effects) of each AI using variance-components analysis (Almasy and Blangero, 1998). Briefly, each AI was entered as a dependent variable into a linear mixed-effects model, which included fixed effects of age, sex, and ICV, and a random effect of genetic similarity, whose covariance structure was determined by the pedigrees. We refer to these as univariate polygenic models. Second, we estimated the genetic correlations between left and right thickness/area measures by extending the univariate polygenic models to incorporate two traits at once; these bivariate polygenic models simultaneously estimate the heritability of left and right paired measures, along with their genetic correlation (which indicates the extent to which their variation is influenced by the same genetic factors).

In principle, family-based analysis such as that using GOBS can conflate shared environmental effects with heritable effects, whereas a twin design is robust to this. Therefore, we also performed heritability analysis in the HCP cohort ($N = 1,113$; see ‘Datasets’ above). HCP is a large-scale project which includes monozygotic and dizygotic twin pairs, as well as unrelated individuals. Precisely the same analyses were conducted in this second cohort.

Results

Meta-analysis of population-level asymmetry

Meta-analysis of population-level asymmetry revealed widely distributed asymmetries in both cortical thickness and surface area. Specifically, we found global differences between the two hemispheres, with generally thicker cortex in the left hemisphere ($b = 0.13$, $Z = 3.64$, $p = 0.00040$; [Figure 2](#)), but larger surface area in the right hemisphere ($b = -0.33$, $Z = -11.30$, $p = 1.36e-29$; [Figure 3](#)).




Figure 2. Forest plots of asymmetry score per dataset, for the overall asymmetry in cortical thickness. Asymmetry score indicates the effect size of the inter-hemispheric difference. The size of a square is proportional to the weights assigned in meta-analysis. The confidence intervals are shown, as well as a dashed vertical line to indicate the point of an asymmetry score of zero.




Figure 3. Forest plots of asymmetry score per dataset, for the overall asymmetry in surface area. Asymmetry score indicates the effect size of the inter-hemispheric difference. The size of a square is proportional to the weights assigned in meta-analysis. The confidence intervals are shown, as well as a dashed vertical line to indicate the point of an asymmetry score of zero.

Substantial, regionally specific differences between the two hemispheres were also observed for

both cortical thickness and surface area. In terms of cortical thickness, 76.5% (26/34) of the regions showed significant asymmetry, after correcting for multiple comparisons ($p < 0.05$, Bonferroni corrected). Specifically, regions showing significant leftward asymmetry (i.e., left > right) of cortical thickness were identified in the anterior cortex, including the lateral, dorsal and medial frontal cortex, the primary sensory, superior parietal, cingulate, and medial temporal cortices (Fig. 4; see [Supplemental Information S4](#)). In contrast, rightward asymmetry (i.e., right > left) was prominent in the posterior regions, including lateral and medial parts of the temporal, parietal, and occipital cortices. This fronto-occipital asymmetry pattern in cortical thickness is striking (Fig. 4) and may also relate to the petalia and Yakovlevian torque effects described above (also see Discussion below). In addition, three temporal regions (especially the inferior temporal and fusiform gyri) showed a trend of rightward asymmetry as defined by uncorrected $P < 0.05$ (inferior temporal: $b = -0.11$, $Z = -2.92$, uncorrected $p = 0.0035$; fusiform: $b = -0.09$, $Z = -2.64$, uncorrected $p = 0.0082$; middle temporal: $b = -0.10$, $Z = -2.19$, uncorrected $p = 0.029$).




Figure 4. Average regional asymmetries in cortical thickness reveal a fronto-occipital pattern. Positive asymmetry (left side in A; red in B) indicates leftward asymmetry, while negative asymmetry (right side in A; blue in B) indicates rightward asymmetry. Asymmetry score indicates the effect size of the inter-hemispheric difference. Error bars indicate standard error of the mean. L, left; R, right.

Similarly, 91.1% (31/34) of the regions showed significant asymmetries of their surface areas after correcting for multiple comparisons ($p < 0.05$, Bonferroni corrected). However, unlike for thicknesses, the surface area asymmetries showed no obvious leftward or rightward patterns involving multiple neighboring areas, or generally along the fronto-occipital axis (Fig. 5; [see Supplemental Information S4](#)). Two language-related regions showed the largest leftward asymmetries of surface area, which were the opercular part of the inferior frontal gyrus (posterior part of the Broca's area) and the transverse temporal gyri (Heschl's gyri). In contrast, however, another two language-related regions, i.e. the triangular part of the inferior frontal gyrus (anterior part of the Broca's area) and the inferior parietal gyrus, showed strong rightward asymmetries of surface area. These findings suggest that opposite asymmetries in morphology of regions within a given network (i.e., language network), or within one functional area (the Broca's area), might be linked to different roles of each constituent part (see Discussion below).




Figure 5. Average asymmetry pattern in surface area. Positive asymmetry (left side in A; red in B) indicates leftward asymmetry, while negative asymmetry (right side in A; blue in B) indicates rightward asymmetry. Asymmetry score indicates the effect size of the inter-hemispheric difference. Error bars indicate standard error of the mean. L, left; R, right.

Effect sizes of cortical thickness and surface area were found to be independent, as illustrated by the absence of a significant correlation between thickness and surface area asymmetries across all cortical regions ($r = -0.14, p = 0.416$).

Moderator analyses using meta-regression

As shown above (e.g., Fig. 2, 3 and **Supplemental Information S4**), we observed moderate to substantial heterogeneity in the asymmetry distributions across datasets (I^2 ranges from 36% to 98%).

To further address the heterogeneity across the samples included in the meta-analyses, we investigated several moderating variables, including sex ratio, median age, handedness ratio, and median ICV. Moderator analyses revealed an influence of the median age of samples on the global hemispheric difference in surface area ($Z = 2.09, p = 0.036$), suggesting a reduced rightward asymmetry with increasing age. No other potential moderators showed significant effects on global cortical thickness or surface area asymmetry ($p > 0.10$). Moderator analyses for each specific region suggested an influence of the median age of samples on the asymmetry of the surface area of the paracentral gyrus ($Z = -4.35, p = 1.38e-5$), and an influence of median ICV on the asymmetry of the surface area in the insula ($Z = -3.18, p = 0.0014$). Given that both the paracentral gyrus and insula showed significant rightward asymmetry in surface area, these findings indicate a decreasing rightward asymmetry with increasing age and with brain size, respectively. In addition, we found a significant effect of scanner field strength on the surface area asymmetry in the insula ($Z = 4.12, p = 3.82e-5$). However, this could be largely reduced by including the other moderating variables (i.e., sex ratio, median age, handedness ratio, and median ICV) in the analysis ($Z = 2.35, p = 0.019$).

Meta-analysis of sex effects on cortical asymmetries

No significant sex effect on the global asymmetry of cortical thickness was found ($p > 0.10$), but notable regionally specific effects on thickness asymmetries were observed in the medial temporal regions (Figure 6), including the parahippocampal gyrus ($Z = 3.57, p = 0.00036$) and the entorhinal cortex ($Z = 3.61, p = 0.00030$), after correcting for multiple comparisons. Together with the

population-level asymmetry observed above, these results indicate that males show more leftward and less rightward asymmetry in cortical thickness of the parahippocampal gyrus and the entorhinal cortex, respectively.

We found a significant sex difference in global asymmetry of surface area ($Z = -2.62, p = 0.0088$), indicating that males have more rightward overall asymmetry in surface area, compared with females. In addition, meta-regression analysis showed that this effect changed with the median ages of samples: we found larger effects of sex (females > males) in the younger samples, compared to the older samples ($Z = 2.80, p = 0.0052$). Regionally specific effects of sex on surface area asymmetry were also revealed (Figure 6; Bonferroni corrected), located in the frontal (superior frontal gyrus, the pars orbitalis region of the left inferior frontal gyrus), temporal (superior temporal gyrus, temporal pole, parahippocampal gyrus and fusiform gyrus), parietal (inferior parietal gyrus and supramarginal gyrus), and anterior cingulate cortices. In addition, various other regions showed nominally significant sex effects (uncorrected $p < 0.05$) without surviving correction for multiple comparisons. More information can be seen in [Supplemental Information S5](#).

We did not observe a significant correlation across regions of the sex effects on the asymmetries of cortical thickness and surface area ($r = 0.14, p = 0.434$), further supporting the independent nature of these two cortical features.




Figure 6. Meta-analysis results for effects of sex, age, brain size, and handedness on regional asymmetry indexes in

cortical thickness and surface area. Red-yellow indicates an increased asymmetry index (AI) in males/with age and brain size; blue-lightblue indicates a decreased AI in males/with age and brain size. A Z threshold of 3.18 ($p = 0.05$, Bonferroni corrected) was used. For more details, please see [Supplemental Information S9](#).

Meta-analysis of age effects on cortical asymmetries

An initial analysis of samples with an age range of larger than five years showed no significant effects of age on global asymmetries of either cortical thickness or surface area ($ps > 0.10$). Several regionally specific, nominally significant effects were found: the superior temporal gyrus (cortical thickness: $Z = 2.38$, $p = 0.017$), the banks of superior temporal sulcus (surface area: $Z = -1.97$, $p = 0.049$), and the entorhinal cortex (surface area: $Z = 2.84$, $p = 0.0045$). However, when restricting the analysis to only those datasets with wider age ranges (at least 20 years range), we observed significant age effects. Specifically, increasing age was associated with more pronounced leftward overall asymmetry in cortical thickness ($Z = 2.65$, $p = 0.0081$), which partly reflects a similar age effect on the thickness asymmetry of the superior temporal gyrus ($Z = 3.17$, $p = 0.0015$; Figure 6). In addition, a similar effect on regional surface area asymmetry was observed in the entorhinal cortex ($Z = 3.21$, $p = 0.0013$). An age effect on surface area asymmetry of the banks of the superior temporal sulcus was nominally significant ($Z = -1.96$, $p = 0.050$). More information can be found in [Supplemental Information S6](#). No significant correlation was found between the age effects across regions on cortical thickness asymmetry and surface area asymmetry ($ps > 0.05$ for both age-range thresholds).

Meta-analysis of group differences by handedness on cortical asymmetries

We did not find significant associations of handedness with cortical asymmetries, even with this unprecedented sample size (from 555 to 608 left-handers versus 6,222 to 7,243 right-handers, depending on the specific regional asymmetry measure). Some temporal regional surface area asymmetries showed nominally significant associations with handedness, including the fusiform gyrus: $Z = 2.00$, $p = 0.046$; the parahippocampal gyrus: $Z = -2.33$, $p = 0.020$; and the superior temporal gyrus: $Z = -2.04$, $p = 0.042$. More information can be found in [Supplemental Information S7](#). No

significant correlation was found between the handedness effects across regions for cortical thickness asymmetry and surface area asymmetry ($r = -0.15, p = 0.403$).

Meta-analysis of brain size effects on cortical asymmetry

Brain size, as indexed with ICV, showed a significant positive effect (i.e., increased leftward asymmetry) on the overall asymmetry in cortical thickness ($Z = 2.14, p = 0.032$). Similar regionally specific effects on cortical thickness asymmetry were found for the inferior parietal gyrus ($Z = 4.51, p = 6.53e-6$), and insula ($Z = 3.71, p = 0.00021$). A negative effect of greater ICV (i.e., decreased leftward asymmetry) was seen in the rostral anterior cingulate gyrus ($Z = -5.23, p = 1.68e-7$) (Figure 6). No significant effect of brain size was found for the overall asymmetry in surface area ($p > 0.10$), but a number of regionally specific effects were revealed (in different directions). Positive effects of greater ICV (i.e., increased leftward/decreased rightward asymmetry) were observed in the medial orbitofrontal gyrus ($Z = 4.17, p = 3.10e-5$), two anterior cingulate gyri (caudal: $Z = 5.71, p = 1.10e-8$; rostral: $Z = 5.67, p = 1.45e-8$), and the isthmus cingulate gyrus ($Z = 4.32, p = 1.59e-5$) (Figure 6). In addition, negative effects of greater ICV (i.e., increased rightward/decreased leftward asymmetry) were seen for the superior frontal gyrus ($Z = -6.58, p = 4.82e-11$), the caudal middle frontal gyrus ($Z = -3.65, p = 0.00026$), the paracentral gyrus ($Z = -5.19, p = 2.11e-7$), the insula ($Z = -5.92, p = 3.13e-9$), the posterior cingulate gyrus ($Z = -3.24, p = 0.0012$), and the cuneus ($Z = -4.49, p = 7.12e-6$) (Figure 6).

More information can be seen in [Supplemental Information S8](#). Similar to the other factors investigated, no significant correlation was found between the effects of brain size across regions on cortical thickness and surface area asymmetry ($r = -0.14, p = 0.417$).

Heritability of cerebral cortical anatomical asymmetries

In the GOBS dataset, the overall hemispheric asymmetries of both cortical thickness and surface area showed low but statistically significant heritabilities (cortical thickness asymmetry: $h^2 = 0.10, p = 0.005$; surface area asymmetry: $h^2 = 0.17, p = 0.00024$). Consistent with a previous study on the genetics of brain structure (Eyler et al., 2014; McKay et al., 2014), regional surface area asymmetries

seemed to be generally more heritable than thickness asymmetries. The most heritable asymmetries in regional cortical thickness were found in the isthmus ($h^2 = 0.17$) and caudal anterior cingulate gyrus ($h^2 = 0.13$), the superior ($h^2 = 0.13$) and rostral middle frontal gyrus ($h^2 = 0.18$), the parahippocampal gyrus ($h^2 = 0.15$), and the lateral occipital gyrus ($h^2 = 0.16$) ($p < 0.05$, Bonferroni corrected; Table 1). The most heritable asymmetries in regional surface area were found in the entorhinal cortex ($h^2 = 0.24$), the superior temporal gyrus ($h^2 = 0.19$), the inferior parietal gyrus ($h^2 = 0.19$), and the isthmus cingulate gyrus ($h^2 = 0.17$) ($p < 0.05$, Bonferroni corrected; Table 1). For each of these regions, we also estimated the genetic correlation between the measures of the left and right structures. While these correlations were high (indicating high pleiotropy), all were significantly different from 1 (see Table 1). These results indicate that most genetic effects on structural variation are shared bilaterally, but some independent genetic effects exist on each hemisphere, which constitute the heritable contributions to structural asymmetry. Finally, we found that the heritability of most of these regions was validated in the HCP dataset. For more details, please see [Supplemental Information S9](#).

Metric	Structure	AI heritability		Genetic correlation between Left and Right		Phenotypic (rho-phen) and environmental (rho-env) correlation between Left and Right		
		h^2	p-value	rho	$p(rho=0)$	$p(rho=1)$	rho-phen	rho-env
Cortical Thickness	Hemisphere	0.102	0.0051	0.993	4.37E-22	0.024	0.950	0.899
	caudalanteriorcingulate	0.129	0.00037	0.718	0.0000016	0.0023	0.285	0.113
	isthmuscingulate	0.171	0.00026	0.823	4.64E-19	0.000045	0.546	0.270
	lateraloccipital	0.163	0.00062	0.914	7.68E-19	0.0012	0.738	0.585
	parahippocampal	0.153	0.00075	0.897	5.1E-28	0.0038	0.656	0.466
	rostralmiddlefrontal	0.181	0.000028	0.881	1.74E-15	0.000053	0.706	0.568
	superiorfrontal	0.128	0.0012	0.950	1.17E-14	0.0051	0.825	0.745
Surface Area	Hemisphere	0.166	0.00025	0.992	3.73E-24	0.0036	0.963	0.925
	entorhinal	0.243	0.000047	0.737	6.15E-09	0.0039	0.437	0.289
	inferiorparietal	0.189	0.000085	0.761	2.43E-11	0.000055	0.504	0.325
	isthmuscingulate	0.167	0.00028	0.729	3.5E-10	0.00015	0.397	0.176
	superiortemporal	0.193	0.000014	0.844	2.64E-18	0.000069	0.591	0.361

Table 1. Significant heritabilities for asymmetry measures based on the GOBS family dataset. In the left part of the table are the heritability and p-values; in the middle part are the genetic correlations between the left and right structural measures, and p-values for whether the genetic correlations differ significantly from 0 or 1. In the right part of the table are the environmental and phenotypic correlation estimates between the left and right regions.

Discussion

In the largest ever analysis of asymmetry of cerebral cortical structure, we applied a meta-analytic approach to brain MRI data from 17,141 healthy individuals from datasets across the world. The findings revealed substantial inter-hemispheric differences in both regional cortical thickness and surface area, and linked some of these asymmetries to sex, age, and brain size. Handedness was not significantly associated with cortical asymmetries. While previous findings are based low hundreds of participants and different methodological approaches, this study of more than 17,000 participants is a major step forward in achieving a more accurate description of the typical asymmetries of the human brain, as well as variation in these asymmetries and some key individual differences factors which affect them. Moreover, with two independent pedigree datasets (i.e., GOBS and HCP), we revealed that several regions showed significant heritability of asymmetry measures.

Cortical thickness

Regions with significant leftward asymmetry in thickness (i.e., left > right) were identified mainly in the frontal cortex, as well as the primary sensory, superior parietal, and medial temporal cortices, while rightward asymmetry was prominent in the posterior cortex, including lateral and medial parts of the temporal, parietal and occipital cortices. This striking asymmetry pattern along the fronto-occipital axis (see Figure 4) is similar to that reported by Plessen et al. (2014) and may be related to the “Yakovlevian torque”, i.e. the frontal/occipital bending in the human brain (Yakovlev, 1972). Specifically, the torque refers to the phenomenon of crossing of the interhemispheric fissure by one hemisphere into the domain of the other. The frontal and occipital bending are the main twisting effects of the torque in opposite directions, with right frontal bending to the left, and left occipital bending to the right (Toga and Thompson, 2003). At the population level, we found that the frontal regions showed leftward asymmetry in cortical thickness, while the occipital regions showed rightward asymmetry.

There were some inconsistencies when comparing our results with previous studies. For example, in 215 healthy participants, Plessen et al. (2014) observed a leftward asymmetry in the inferior frontal cortex, which includes Broca’s area in the inferior frontal gyrus. The authors suggested that this might correspond anatomically with the functional asymmetry for expressive language in these regions, as

has been reported on the basis of brain lesion studies and functional neuroimaging studies (Hagoort, 2005; Horwitz et al., 2003; Rutten et al., 1999). However, this interpretation should be considered with caution in light of a recent study on cortical thickness asymmetries with 250 adults showing an opposite direction of asymmetry (rightward) in this region (Koelkebeck et al., 2014). In the present study, with a much larger sample size, we failed to detect any cortical thickness asymmetry in this region (i.e., the pars opercularis and pars triangularis of the inferior frontal gyrus, uncorrected $p > 0.45$). Another difference with previous findings concerns the supramarginal gyrus, which showed a strong leftward asymmetry in Plessen et al. (2014), but no asymmetry in two other studies (Koelkebeck et al., 2014; Luders et al., 2006), and also not in the present study. This indicates an absence of population-level lateralization in cortical thickness in the supramarginal gyrus, and again underlines the value of the present study in achieving a more accurate characterization of the average anatomical brain laterality.

There are several issues that may contribute to discrepancies of our present results with these previous studies, including the large sample size that we used, as well as the worldwide population. Varying demographic factors, such as sex and age, across the various previous studies might also have played an important role. In the current study, we identified several regions showing significant effects of these factors on the asymmetry of cortical thickness. For sex, notable effects were observed in the medial temporal regions, including the parahippocampal gyrus (more leftward in males) and the entorhinal cortex (more rightward in females), while mixed results have been obtained in previous studies (Kang et al., 2015; Plessen et al., 2014). Considering the critical roles of these two regions in visuospatial processing and spatial navigation (e.g., (Epstein et al., 2003; Solstad et al., 2008)), these sex differences may be related to the tendency for slight male advantage on spatial tasks (Alexander and Wilcox, 2012; Clements et al., 2006; Hines, 2010). In contrast to Plessen et al. (2014), we found no sex differences in cortical thickness asymmetry of core regions of the language network, including the pars opercularis and pars triangularis of the inferior frontal gyrus (the Broca's area), the transverse temporal gyrus (the Heschl's gyrus), and the supramarginal gyrus (uncorrected $p > 0.05$). These results are consistent with two other studies (Kang et al., 2015; Luders et al., 2006), and indicate that subtle

sex differences in the performance on language tasks and language lateralization (Clements et al., 2006) cannot be linked to sex differences in cortical thickness asymmetry of these regions.

In terms of age effects, when limiting our analysis to only the datasets with an age range greater than 20 years, we found a significant correlation between age and the overall hemispheric asymmetry in cortical thickness (i.e., increasing age correlated with more pronounced leftward asymmetry), which was mainly contributed by the superior temporal gyrus. This finding is consistent with previous studies (Plessen et al., 2014; Zhou et al., 2013), though we did not detect age effects in other regions reported by Plessen et al. (2014). Brain size is another factor that can affect functional organization (Razoumnikova, 2000; Yan et al., 2011). In the present study, we found a significant effect of brain size on the overall asymmetry in cortical thickness, such that the leftward asymmetry in cortical thickness increases in larger brains. This effect was the most pronounced in the inferior parietal gyrus and the insula. Our findings on brain size are in accord with the hypothesis that asymmetries increase in larger brains, which might relate to the increased inter-hemispheric distance and transfer time in larger brains (Herve et al., 2013).

No detectable association of handedness with cortical thickness asymmetries was found, even in this unprecedented sample size. This negative result is consistent with one recent study (Maingault et al., 2016), while in other studies handedness has not been included as a variable of interest (Kang et al., 2015; Luders et al., 2006; Plessen et al., 2014; Zhou et al., 2013). It remains possible that handedness is associated with asymmetry measures of other structural metrics and/or in more narrowly defined regions. However, it is clear from the present results that left-handedness does not involve any broad or substantial alterations of cortical thickness asymmetry.

Surface area

Regarding surface area, population-level asymmetry was generally more prominent compared to that of cortical thickness. A large majority of regions (91.1%) showed significant asymmetry in surface area, although with no obvious directional pattern (leftward or rightward) affecting multiple neighboring regions, or along the anterior-posterior axis, as we observed for thicknesses. The present

study detected some similar asymmetry patterns of surface area to those of two previous studies (Chiarello et al., 2016; Koelkebeck et al., 2014). Specifically, consistent results included leftward asymmetry of the superior frontal gyrus, the postcentral gyrus, supramarginal gyrus, and the entorhinal cortex, and rightward asymmetry in the caudal anterior cingulate cortex and the middle temporal gyrus (Chiarello et al., 2016; Kang et al., 2015; Koelkebeck et al., 2014; Maingault et al., 2016). The leftward asymmetry of surface area in the supramarginal gyrus is consistent with the widely-observed volume asymmetry in the Perisylvian regions, which is related to an asymmetrical shift caused by the brain torque (Glasel et al., 2011; Good et al., 2001; Herve et al., 2006; Maingault et al., 2016; Watkins et al., 2001).

We identified several additional regions that are asymmetric in terms of surface area, not previously described. Among these regions, two language-related regions, including the opercular part of the inferior frontal gyrus (the posterior part of Broca's area) and the transverse temporal gyrus (Heschl's gyrus) showed the largest leftward asymmetries. Based on these findings, the asymmetry of surface area (rather than cortical thickness as suggested in Plessen et al., 2014) may correspond anatomically with language lateralization in these regions, although further study is needed investigating both structure and function. Moreover, we found two other language-related regions showing strong asymmetry in the opposite direction (rightward), including the triangular part of the inferior frontal gyrus (the anterior part of Broca's area) and the inferior parietal gyrus. Taking these observations together, it appears that the structural basis of functional language lateralization is more complex than previously thought. For example, as mentioned above, for Broca's area, one of the most well-established areas for language function and language lateralization, while we did not detect asymmetry in terms of cortical thickness, we indeed observed strong asymmetry in surface area within this region. Moreover, the asymmetry was in different directions in two sub-regions of this area: leftward for the posterior part and rightward for the anterior part. These findings may be closely related to distinct roles of these two sub-areas in language functions: these two sub-regions are involved in, respectively, phonology and syntax, related to their distinct connections with areas in inferior parietal and temporal cortex (Hagoort, 2014; Xiang et al., 2010). Thus, these findings

suggested that the opposite directions of structural asymmetry affecting regions within one network or within one functional area might reflect different functional involvements of each component region.

Future studies with both structural and functional data in same participants may help link the structural asymmetries to functional asymmetries in the human brain.

The effects of biological factors on surface area asymmetries were more prominent than on thickness asymmetries. Very few previous studies have reported sex effects. Kang et al. (2015) found no sex differences in asymmetries for surface areas in 138 young adults, while Koelkebeck et al. (2014) only reported a male > female effect for the asymmetry of surface area at the overall hemispheric level in 101 healthy individuals. We also found that males, on average, showed more rightward asymmetry in overall surface area, compared with females, which is consistent with Koelkebeck et al. (2014). We additionally observed a number of regionally specific effects, among which surface area asymmetry in the superior frontal gyrus showed the strongest relation to sex (i.e., males showed more leftward asymmetry in surface area in this region compared to females).

In terms of age, when including only those datasets with an age range of more than 20 years, we found a weak positive correlation between age and the asymmetry of surface area of the entorhinal cortex, that is, the leftward asymmetry of this region was slightly greater in older participants. As far as we are aware, no previous studies have reported possible age effects on the asymmetries of surface area, except one that showed no significant results in 101 participants (Koelkebeck et al., 2014). Note that, in our analyses for either sex or age effects, brain size was included as a covariate to obtain sex- or age-specific effects. In terms of brain size effects themselves (correcting for sex and age), no significant effect was found on the overall asymmetry of surface area, but a number of regionally specific effects were revealed. Specifically, positive effects (increased leftward/decreased rightward asymmetries with brain size) were observed mainly in medial regions such as the anterior cingulate gyri, while negative effects (decreased leftward or increased rightward asymmetries with brain size) were seen in spatially diverse locations, including the posterior cingulate gyrus, the insula, and the caudal middle frontal gyrus. It has been suggested that increased brain size might lead to the development of additional sulci (Kang et al., 2015), which could impact on regional asymmetries as

assessed with the FreeSurfer atlas-based approach.

No significant associations of handedness with cortical surface area asymmetries were found.

This negative result is consistent with two recent studies (Guadalupe et al., 2014; Maingault et al., 2016), while in other studies handedness has not been included as a variable of interest (Chiarello et al., 2016; Kang et al., 2015; Koelkebeck et al., 2014). Note that our results did include several nominally significant handedness associations in several temporal regions, including the fusiform, parahippocampal, and superior temporal gyri. However, these effects did not survive correction of multiple comparisons, and further studies are needed to confirm or refute these associations.

General discussion

Our findings bear on the relationship between asymmetry of cortical thickness and surface area. Previous studies have suggested that thickness and surface area are evolutionarily, genetically, and developmentally distinct (Panizzon et al., 2009; Raznahan et al., 2011), and that therefore separate consideration of these aspects of cortical anatomy is important (e.g., (Winkler et al., 2010)). With a large MRI twin sample, Panizzon et al. (2009) showed that, although average cortical thickness and total surface area are both highly heritable (>0.80), they are essentially unrelated genetically (genetic correlation = 0.08). This genetic independence of cortical thickness and surface area was also found in a large extended family study (Winkler et al., 2010). These results suggest relative independence of the two surface-based measures, and potentially therefore their asymmetry patterns. Data from two recent studies has indeed indicated that the asymmetry measures of cortical thickness and surface area are relatively independent at the overall hemispheric level (Koelkebeck et al., 2014; Maingault et al., 2016). With our larger sample size in the present study (including the BIL&GIN dataset used in Maingault et al., 2016), we confirmed a lack of correlation across regions between the asymmetries of thickness and surface areas, which further supports their independent natures. Moreover, by including data on participants' sex, age, handedness, and brain size, our findings further elaborated the largely independent nature of regional area versus thickness variability. That is, no overall correlations were found between the effects of these factors across regions, on cortical thickness and surface area. Note

that, when zooming in on some individual regions, there may be identifiable relations between thickness and surface area asymmetries, such as reported for the fusiform gyrus and the cingulate cortex (Chiarello et al., 2016; Koelkebeck et al., 2014; Maingault et al., 2016), although further investigation is needed. In future studies of cortical asymmetry, the simultaneous investigation of both cortical thickness and surface area will be important. For example, this may be necessary in order to approach the genetics of brain asymmetry (Guadalupe et al., 2014) and its links with functional lateralization (e.g., language lateralization) (Tzourio-Mazoyer et al., 2015).

With the pedigree datasets from the GOBS and HCP, we revealed that several regions showed significant heritability of their asymmetry measures, while the general distributions of heritability estimates across the cortical regions were quite different for thickness and surface asymmetries (Supplemental Information S9). These data on heritability will be useful in targeting future studies of brain laterality with, for example, genome-wide association scanning aimed at identifying genes involved. Interestingly, cortical asymmetry of the human brain may also be associated with inter-hemispheric differences in gene expression (Francks, 2015; Karlebach and Francks, 2015).

Our data revealed extensive variability in cortical asymmetry across participants and samples. Besides sex, age, brain size, handedness, and heritable effects, further studies on individual variability are needed, from the perspective of cognitive and neuropsychiatric disorders. Some disorders, such as dyslexia (Altarelli et al., 2014), Alzheimer's disease (Thompson et al., 1998), ADHD (Shaw et al., 2009), psychotic disorders (Crow, 1990; Schobel et al., 2009; Yucel et al., 2002; Yucel et al., 2003), autism (Eyler et al., 2012), and mood disorders (Liu et al., 2016; Yucel et al., 2009), may be associated with abnormal cortical asymmetries, though these complex links have not been fully explored.

Asymmetry measures may even be more accurate than unilateral cortical measures to distinguish healthy controls from patients in some contexts (Haller et al., 2009), suggesting the potential for cortical asymmetry to be used as an important biomarker. In this respect, the findings in this paper provide a reference for cortical asymmetry in healthy populations, which may help for further understanding the nature of these disorders in future studies. In fact, studies of cortical asymmetry in several disorders are currently underway within the ENIGMA consortium, using the same

methodology as used here.

Regarding handedness, it is interesting to note that paleoneurologists have attempted to use skull endocasts to assess cerebral asymmetries and to infer the evolution of handedness in hominins (Holloway, 2015). Since we found no significant association between brain anatomical asymmetries and handedness, our analysis does not support the use of indirect measures of brain anatomy to infer the handedness of individuals.

Limitations and future directions

This study has several limitations which could be overcome in future studies. First, for age effects, the cross-sectional study design limits the interpretation of results. Longitudinal studies should ideally be performed to support the findings and provide the normal course of development and aging of brain asymmetry.

Second, when combining already collected data across worldwide samples, data collection protocols are not prospectively harmonized. Imaging acquisition protocols and handedness assessments therefore differed across studies, which resulted in possible sources of heterogeneity. On the other hand, this heterogeneity can be taken as an advantage of our approach, in the sense that our findings are representative of the real-world diversity of MRI acquisition currently in use in the field, and not limited to a single lab-specific protocol.

Third, we note that variability of asymmetry in surface area across samples was relatively lower than that of asymmetry in cortical thickness, at both the global hemispheric and regional levels (see [Supplemental Information S6](#)). The relatively consistent asymmetry in surface area across datasets might be, to an extent, driven by the same parcellation scheme having been used across all samples. The potential impact of this issue is an important direction for future studies. In addition, the region-based approach is necessarily limited in terms of spatial resolution, related to the number of cortical regions defined. A vertex-wise approach combined with cross-hemispheric registration methods could be used in future cortical asymmetry studies (Kang et al., 2015; Maingault et al., 2016; Van Essen et al., 2012).

Finally, besides the directions of the asymmetries, the present study provided the exact effect size distributions for each region with a very large sample size. The results can act as a guide and provide a reference normative resource for future studies of cortical asymmetry. For example, with the population-level effect sizes, researchers can estimate sample sizes required to detect specific effects of interest. Researchers can query the meta-analysis summary statistics with the query tool (<http://conxz.github.io/neurohemi/>).

Summary

In summary, we showed that diverse regions of the human cerebral cortex are asymmetrical in their structural features (i.e., cortical thickness and surface area) with different effect sizes, and that the asymmetry patterns are different between cortical thickness and surface area. Moreover, we showed widespread effects of several biological factors (e.g., sex, age and brain size) on the cortical asymmetries, but found no significant handedness effects. Finally, we revealed that the human brain is composed of regions with significant heritability of the asymmetry characteristics. This study not only contributes to the understanding of human brain asymmetry in the healthy population, but also provides informative data for future studies of the genetics of brain asymmetry, and potentially abnormal brain asymmetry in cognitive and neuropsychiatric disorders.

Acknowledgements

Funding information for each site is available in a separate Acknowledgements document. All sites within the ENIGMA-Lateralization working group are very grateful to the participants for their generosity of time and willingness to participate in each of the collaborating studies.

Compliance with ethical standards

All participating studies were approved by the local ethical committee. Informed consent was obtained from all participants involved. The authors declare no conflicts of interest except for the authors listed in Conflicts of interest.

References

Alexander, G.M., Wilcox, T., 2012. Sex Differences in Early Infancy. *Child Development Perspectives* 6, 400-406.

Almasy, L., Blangero, J., 1998. Multipoint quantitative-trait linkage analysis in general pedigrees. *Am J Hum Genet* 62, 1198-1211.

Altarelli, I., Leroy, F., Monzalvo, K., Fluss, J., Billard, C., Dehaene-Lambertz, G., Galaburda, A.M., Ramus, F., 2014. Planum temporale asymmetry in developmental dyslexia: Revisiting an old question. *Hum Brain Mapp* 35, 5717-5735.

Badzakova-Trajkov, G., Haberling, I.S., Roberts, R.P., Corballis, M.C., 2010. Cerebral asymmetries: complementary and independent processes. *PLoS One* 5, e9682.

Bishop, D.V., 2013. Cerebral asymmetry and language development: cause, correlate, or consequence? *Science* 340, 1230531.

Borenstein, M., Hedges, L.V., Higgins, J.P., Rothstein, H.R., 2010. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods* 1, 97-111.

Chi, J.G., Dooling, E.C., Gilles, F.H., 1977. Left-right asymmetries of the temporal speech areas of the human fetus. *Arch Neurol* 34, 346-348.

Chiarello, C., Vazquez, D., Felton, A., McDowell, A., 2016. Structural asymmetry of the human cerebral cortex: Regional and between-subject variability of surface area, cortical thickness, and local gyration. *Neuropsychologia* 93, 365-379.

Clements, A.M., Rimrodt, S.L., Abel, J.R., Blankner, J.G., Mostofsky, S.H., Pekar, J.J., Denckla, M.B., Cutting, L.E., 2006. Sex differences in cerebral laterality of language and visuospatial processing. *Brain Lang* 98, 150-158.

Concha, M.L., Bianco, I.H., Wilson, S.W., 2012. Encoding asymmetry within neural circuits. *Nature Reviews Neuroscience* 13, 832-843.

Crow, T.J., 1990. Temporal-Lobe Asymmetries as the Key to the Etiology of Schizophrenia. *Schizophrenia Bulletin* 16, 433-&.

Desikan, R.S., Segonne, F., Fischl, B., Quinn, B.T., Dickerson, B.C., Blacker, D., Buckner, R.L., Dale, A.M., Maguire, R.P., Hyman, B.T., Albert, M.S., Killiany, R.J., 2006. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* 31, 968-980.

Dorsaint-Pierre, R., Penhune, V.B., Watkins, K.E., Neelin, P., Lerch, J.P., Bouffard, M., Zatorre, R.J., 2006. Asymmetries of the planum temporale and Heschl's gyrus: relationship to language lateralization. *Brain* 129, 1164-1176.

Epstein, R., Graham, K.S., Downing, P.E., 2003. Viewpoint-specific scene representations in human parahippocampal cortex. *Neuron* 37, 865-876.

Esteves, M., Marques, P., Magalhaes, R., Castanho, T.C., Soares, J.M., Almeida, A., Santos, N.C., Sousa, N., Leite-Almeida, H., 2017. Structural laterality is associated with cognitive and mood outcomes: An assessment of 105 healthy aged volunteers. *Neuroimage* 153, 86-96.

Eyler, L.T., Pierce, K., Courchesne, E., 2012. A failure of left temporal cortex to specialize for language is an early emerging and fundamental property of autism. *Brain* 135, 949-960.

Eyler, L.T., Vuoksimaa, E., Panizzon, M.S., Fennema-Notestine, C., Neale, M.C., Chen, C.H., Jak, A., Franz, C.E., Lyons, M.J., Thompson, W.K., Spoon, K.M., Fischl, B., Dale, A.M., Kremen, W.S., 2014. Conceptual and data-based investigation of genetic influences and brain asymmetry: a twin study of multiple structural phenotypes. *J Cogn Neurosci* 26, 1100-1117.

Fischl, B., Salat, D.H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., van der Kouwe, A., Killiany, R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N., Rosen, B., Dale, A.M., 2002. Whole brain segmentation: Automated labeling of neuroanatomical structures in the human brain. *Neuron* 33, 341-355.

Fornito, A., Yucel, M., Wood, S., Stuart, G.W., Buchanan, J.A., Proffitt, T., Anderson, V., Velakoulis, D., Pantelis, C., 2004. Individual differences in anterior cingulate/paracingulate morphology are related to executive functions in healthy males. *Cereb Cortex* 14, 424-431.

Francks, C., 2015. Exploring human brain lateralization with molecular genetics and genomics. *Ann N*

Y Acad Sci 1359, 1-13.

Glasel, H., Leroy, F., Dubois, J., Hertz-Pannier, L., Mangin, J.F., Dehaene-Lambertz, G., 2011. A robust cerebral asymmetry in the infant brain: the rightward superior temporal sulcus. Neuroimage 58, 716-723.

Goldberg, E., Roediger, D., Kucukboyaci, N.E., Carlson, C., Devinsky, O., Kuzniecky, R., Halgren, E., Thesen, T., 2013. Hemispheric asymmetries of cortical volume in the human brain. Cortex 49, 200-210.

Good, C.D., Johnsrude, I., Ashburner, J., Henson, R.N., Friston, K.J., Frackowiak, R.S., 2001. Cerebral asymmetry and the effects of sex and handedness on brain structure: a voxel-based morphometric analysis of 465 normal adult human brains. Neuroimage 14, 685-700.

Greve, D.N., Van der Haegen, L., Cai, Q., Stufflebeam, S., Sabuncu, M.R., Fischl, B., Brysbaert, M., 2013. A Surface-based Analysis of Language Lateralization and Cortical Asymmetry. Journal of Cognitive Neuroscience 25, 1477-1492.

Guadalupe, T., Mathias, S.R., vanErp, T.G., Whelan, C.D., Zwiers, M.P., Abe, Y., Abramovic, L., Agartz, I., Andreassen, O.A., Arias-Vasquez, A., Aribisala, B.S., Armstrong, N.J., Arolt, V., Artiges, E., Ayesa-Arriola, R., Baboyan, V.G., Banaschewski, T., Barker, G., Bastin, M.E., Baune, B.T., Blangero, J., Bokde, A.L., Boedhoe, P.S., Bose, A., Brem, S., Brodaty, H., Bromberg, U., Brooks, S., Buchel, C., Buitelaar, J., Calhoun, V.D., Cannon, D.M., Cattrell, A., Cheng, Y., Conrod, P.J., Conzelmann, A., Corvin, A., Crespo-Facorro, B., Crivello, F., Dannlowski, U., de Zubicaray, G.I., de Zwart, S.M., Deary, I.J., Desrivieres, S., Doan, N.T., Donohoe, G., Dorum, E.S., Ehrlich, S., Espeseth, T., Fernandez, G., Flor, H., Fouche, J.P., Frouin, V., Fukunaga, M., Gallinat, J., Garavan, H., Gill, M., Suarez, A.G., Gowland, P., Grabe, H.J., Grotegerd, D., Gruber, O., Hagenaars, S., Hashimoto, R., Hauser, T.U., Heinz, A., Hibar, D.P., Hoekstra, P.J., Hoogman, M., Howells, F.M., Hu, H., Hulshoff Pol, H.E., Huyser, C., Ittermann, B., Jahanshad, N., Jonsson, E.G., Jurk, S., Kahn, R.S., Kelly, S., Kraemer, B., Kugel, H., Kwon, J.S., Lemaitre, H., Lesch, K.P., Lochner, C., Luciano, M., Marquand, A.F., Martin, N.G., Martinez-Zalacain, I., Martinot, J.L., Mataix-Cols, D., Mather, K., McDonald, C., McMahon, K.L., Medland, S.E., Menchon, J.M., Morris, D.W., Mothersill, O., Maniega, S.M., Mwangi, B., Nakamae, T., Nakao, T., Narayanaswamy, J.C., Nees, F., Nordvik, J.E., Onnink, A.M., Opel, N., Ophoff, R., Paillere Martinot, M.L., Papadopoulos Orfanos, D., Pauli, P., Paus, T., Poustka, L., Reddy, J.Y., Renteria, M.E., Roiz-Santianez, R., Roos, A., Royle, N.A., Sachdev, P., Sanchez-Juan, P., Schmaal, L., Schumann, G., Shumskaya, E., Smolka, M.N., Soares, J.C., Soriano-Mas, C., Stein, D.J., Strike, L.T., Toro, R., Turner, J.A., Tzourio-Mazoyer, N., Uhlmann, A., Hernandez, M.V., van den Heuvel, O.A., van der Meer, D., van Haren, N.E., Veltman, D.J., Venkatasubramanian, G., Vetter, N.C., Vuletic, D., Walitza, S., Walter, H., Walton, E., Wang, Z., Wardlaw, J., Wen, W., Westlye, L.T., Whelan, R., Wittfeld, K., Wolfers, T., Wright, M.J., Xu, J., Xu, X., Yun, J.Y., Zhao, J., Franke, B., Thompson, P.M., Glahn, D.C., Mazoyer, B., Fisher, S.E., Francks, C., 2016. Human subcortical brain asymmetries in 15,847 people worldwide reveal effects of age and sex. Brain Imaging Behav.

Guadalupe, T., Willems, R.M., Zwiers, M.P., Vasquez, A.A., Hoogman, M., Hagoort, P., Fernandez, G., Buitelaar, J., Franke, B., Fisher, S.E., Francks, C., 2014. Differences in cerebral cortical anatomy of left- and right-handers. Frontiers in Psychology 5.

Guadalupe, T., Zwiers, M.P., Wittfeld, K., Teumer, A., Vasquez, A.A., Hoogman, M., Hagoort, P., Fernandez, G., Buitelaar, J., van Bokhoven, H., Hegenscheid, K., Volzke, H., Franke, B., Fisher, S.E., Grabe, H.J., Francks, C., 2015. Asymmetry within and around the human planum temporale is sexually dimorphic and influenced by genes involved in steroid hormone receptor activity. Cortex 62, 41-55.

Hagoort, P., 2005. On Broca, brain, and binding: a new framework. Trends in Cognitive Sciences 9, 416-423.

Hagoort, P., 2014. Nodes and networks in the neural architecture for language: Broca's region and beyond. Curr Opin Neurobiol 28, 136-141.

Haller, S., Borgwardt, S.J., Schindler, C., Aston, J., Radue, E.W., Riecher-Rossler, A., 2009. Can Cortical Thickness Asymmetry Analysis Contribute to Detection of At-Risk Mental State and First-Episode Psychosis?: A Pilot Study. Radiology 250, 212-221.

Hering-Hanit, R., Achiron, R., Lipitz, S., Achiron, A., 2001. Asymmetry of fetal cerebral hemispheres: in utero ultrasound study. *Arch Dis Child Fetal Neonatal Ed* 85, F194-196.

Herve, P.Y., Crivello, F., Perchey, G., Mazoyer, B., Tzourio-Mazoyer, N., 2006. Handedness and cerebral anatomical asymmetries in young adult males. *Neuroimage* 29, 1066-1079.

Herve, P.Y., Zago, L., Petit, L., Mazoyer, B., Tzourio-Mazoyer, N., 2013. Revisiting human hemispheric specialization with neuroimaging. *Trends in Cognitive Sciences* 17, 69-80.

Hines, M., 2010. Sex-related variation in human behavior and the brain. *Trends in Cognitive Sciences* 14, 448-456.

Hogstrom, L.J., Westlye, L.T., Walhovd, K.B., Fjell, A.M., 2013. The structure of the cerebral cortex across adult life: age-related patterns of surface area, thickness, and gyration. *Cereb Cortex* 23, 2521-2530.

Holloway, R.L., 2015. The evolution of the hominid brain. *Handbook of paleoanthropology*, 1961-1987.

Horwitz, B., Amunts, K., Bhattacharyya, R., Patkin, D., Jeffries, K., Zilles, K., Braun, A.R., 2003. Activation of Broca's area during the production of spoken and signed language: a combined cytoarchitectonic mapping and PET analysis. *Neuropsychologia* 41, 1868-1876.

Im, K., Lee, J.M., Lee, J., Shin, Y.W., Kim, I.Y., Kwon, J.S., Kim, S.I., 2006. Gender difference analysis of cortical thickness in healthy young adults with surface-based methods. *Neuroimage* 31, 31-38.

Kang, X., Herron, T.J., Ettlinger, M., Woods, D.L., 2015. Hemispheric asymmetries in cortical and subcortical anatomy. *Laterality* 20, 658-684.

Karlebach, G., Francks, C., 2015. Lateralization of gene expression in human language cortex. *Cortex* 67, 30-36.

Koelkebeck, K., Miyata, J., Kubota, M., Kohl, W., Son, S., Fukuyama, H., Sawamoto, N., Takahashi, H., Murai, T., 2014. The contribution of cortical thickness and surface area to gray matter asymmetries in the healthy human brain. *Hum Brain Mapp* 35, 6011-6022.

Levin, M., 2005. Left-right asymmetry in embryonic development: a comprehensive review. *Mech Dev* 122, 3-25.

Liu, W., Mao, Y., Wei, D.T., Yang, J.Y., Du, X., Xie, P., Qiu, J., 2016. Structural Asymmetry of Dorsolateral Prefrontal Cortex Correlates with Depressive Symptoms: Evidence from Healthy Individuals and Patients with Major Depressive Disorder. *Neuroscience Bulletin* 32, 217-226.

Luders, E., Narr, K.L., Thompson, P.M., Rex, D.E., Jancke, L., Toga, A.W., 2006. Hemispheric asymmetries in cortical thickness. *Cereb Cortex* 16, 1232-1238.

Lyttelton, O.C., Karama, S., Ad-Dab'bagh, Y., Zatorre, R.J., Carbonell, F., Worsley, K., Evans, A.C., 2009. Positional and surface area asymmetry of the human cerebral cortex. *Neuroimage* 46, 895-903.

Maingault, S., Tzourio-Mazoyer, N., Mazoyer, B., Crivello, F., 2016. Regional correlations between cortical thickness and surface area asymmetries: A surface-based morphometry study of 250 adults. *Neuropsychologia* 93, 350-364.

Marie, D., Jobard, G., Crivello, F., Perchey, G., Petit, L., Mellet, E., Joliot, M., Zago, L., Mazoyer, B., Tzourio-Mazoyer, N., 2015. Descriptive anatomy of Heschl's gyri in 430 healthy volunteers, including 198 left-handers. *Brain Struct Funct* 220, 729-743.

McKay, D.R., Knowles, E.E., Winkler, A.A., Sprooten, E., Kochunov, P., Olvera, R.L., Curran, J.E., Kent, J.W., Jr., Carless, M.A., Goring, H.H., Dyer, T.D., Duggirala, R., Almasy, L., Fox, P.T., Blangero, J., Glahn, D.C., 2014. Influence of age, sex and genetic factors on the human brain. *Brain Imaging Behav* 8, 143-152.

Meyer, M., Liem, F., Hirsiger, S., Jancke, L., Hanggi, J., 2014. Cortical Surface Area and Cortical Thickness Demonstrate Differential Structural Asymmetry in Auditory-Related Areas of the Human Cortex. *Cerebral Cortex* 24, 2541-2552.

Narr, K.L., Bilder, R.M., Luders, E., Thompson, P.M., Woods, R.P., Robinson, D., Szeszko, P.R., Dintcheva, T., Gurbani, M., Toga, A.W., 2007. Asymmetries of cortical shape: Effects of handedness, sex and schizophrenia. *Neuroimage* 34, 939-948.

Panizzon, M.S., Fennema-Notestine, C., Eyler, L.T., Jernigan, T.L., Prom-Wormley, E., Neale, M., Jacobson, K., Lyons, M.J., Grant, M.D., Franz, C.E., Xian, H., Tsuang, M., Fischl, B., Seidman, L.,

Dale, A., Kremen, W.S., 2009. Distinct genetic influences on cortical surface area and cortical thickness. *Cereb Cortex* 19, 2728-2735.

Plessen, K.J., Hugdahl, K., Bansal, R., Hao, X., Peterson, B.S., 2014. Sex, age, and cognitive correlates of asymmetries in thickness of the cortical mantle across the life span. *J Neurosci* 34, 6294-6302.

Raz, N., Gunning-Dixon, F., Head, D., Rodriguez, K.M., Williamson, A., Acker, J.D., 2004. Aging, sexual dimorphism, and hemispheric asymmetry of the cerebral cortex: replicability of regional differences in volume. *Neurobiol Aging* 25, 377-396.

Raznahan, A., Shaw, P., Lalonde, F., Stockman, M., Wallace, G.L., Greenstein, D., Clasen, L., Gogtay, N., Giedd, J.N., 2011. How Does Your Cortex Grow? *Journal of Neuroscience* 31, 7174-7177.

Razoumnikova, O.M., 2000. Functional organization of different brain areas during convergent and divergent thinking: an EEG investigation. *Brain Res Cogn Brain Res* 10, 11-18.

Rutten, G.J., van Rijen, P.C., van Veelen, C.W., Ramsey, N.F., 1999. Language area localization with three-dimensional functional magnetic resonance imaging matches intrasulcal electrostimulation in Broca's area. *Ann Neurol* 46, 405-408.

Salat, D.H., Buckner, R.L., Snyder, A.Z., Greve, D.N., Desikan, R.S., Busa, E., Morris, J.C., Dale, A.M., Fischl, B., 2004. Thinning of the cerebral cortex in aging. *Cereb Cortex* 14, 721-730.

Schobel, S.A., Kelly, M.A., Corcoran, C.M., Van Heertum, K., Seckinger, R., Goetz, R., Harkavy-Friedman, J., Malaspina, D., 2009. Anterior hippocampal and orbitofrontal cortical structural brain abnormalities in association with cognitive deficits in schizophrenia. *Schizophr Res* 114, 110-118.

Shaw, P., Lalonde, F., Lepage, C., Rabin, C., Eckstrand, K., Sharp, W., Greenstein, D., Evans, A., Giedd, J.N., Rapoport, J., 2009. Development of cortical asymmetry in typically developing children and its disruption in attention-deficit/hyperactivity disorder. *Arch Gen Psychiatry* 66, 888-896.

Solstad, T., Boccara, C.N., Kropff, E., Moser, M.B., Moser, E.I., 2008. Representation of geometric borders in the entorhinal cortex. *Science* 322, 1865-1868.

Steinmetz, H., Volkmann, J., Jancke, L., Freund, H.J., 1991. Anatomical left-right asymmetry of language-related temporal cortex is different in left- and right-handers. *Ann Neurol* 29, 315-319.

Thompson, P.M., Moussai, J., Zohoori, S., Goldkorn, A., Khan, A.A., Mega, M.S., Small, G.W., Cummings, J.L., Toga, A.W., 1998. Cortical variability and asymmetry in normal aging and Alzheimer's disease. *Cereb Cortex* 8, 492-509.

Thompson, P.M., Stein, J.L., Medland, S.E., Hibar, D.P., Vasquez, A.A., Renteria, M.E., Toro, R., Jahanshad, N., Schumann, G., Franke, B., Wright, M.J., Martin, N.G., Agartz, I., Alda, M., Alhusaini, S., Almasy, L., Almeida, J., Alpert, K., Andreasen, N.C., Andreassen, O.A., Apostolova, L.G., Appel, K., Armstrong, N.J., Aribisala, B., Bastin, M.E., Bauer, M., Bearden, C.E., Bergmann, O., Binder, E.B., Blangero, J., Bockholt, H.J., Boen, E., Bois, C., Boomsma, D.I., Booth, T., Bowman, I.J., Bralten, J., Brouwer, R.M., Brunner, H.G., Brohawn, D.G., Buckner, R.L., Buitelaar, J., Bulayeva, K., Bustillo, J.R., Calhoun, V.D., Cannon, D.M., Cantor, R.M., Carless, M.A., Caseras, X., Cavalleri, G.L., Chakravarty, M.M., Chang, K.D., Ching, C.R., Christoforou, A., Cichon, S., Clark, V.P., Conrod, P., Coppola, G., Crespo-Facorro, B., Curran, J.E., Czisch, M., Deary, I.J., de Geus, E.J., den Braber, A., Delvecchio, G., Depondt, C., de Haan, L., de Zubicaray, G.I., Dima, D., Dimitrova, R., Djurovic, S., Dong, H., Donohoe, G., Duggirala, R., Dyer, T.D., Ehrlich, S., Ekman, C.J., Elvsashagen, T., Emsell, L., Erk, S., Espeseth, T., Fagerness, J., Fears, S., Fedko, I., Fernandez, G., Fisher, S.E., Foroud, T., Fox, P.T., Francks, C., Frangou, S., Frey, E.M., Frodl, T., Frouin, V., Garavan, H., Giddaluru, S., Glahn, D.C., Godlewska, B., Goldstein, R.Z., Gollub, R.L., Grabe, H.J., Grimm, O., Gruber, O., Guadalupe, T., Gur, R.E., Gur, R.C., Goring, H.H., Hagenaars, S., Hajek, T., Hall, G.B., Hall, J., Hardy, J., Hartman, C.A., Hass, J., Hatton, S.N., Haukvik, U.K., Hegenscheid, K., Heinz, A., Hickie, I.B., Ho, B.C., Hoehn, D., Hoekstra, P.J., Hollinshead, M., Holmes, A.J., Homuth, G., Hoogman, M., Hong, L.E., Hosten, N., Hottenga, J.J., Hulshoff Pol, H.E., Hwang, K.S., Jack, C.R., Jr., Jenkinson, M., Johnston, C., Jonsson, E.G., Kahn, R.S., Kasperaviciute, D., Kelly, S., Kim, S., Kochunov, P., Koenders, L., Kramer, B., Kwok, J.B., Lagopoulos, J., Laje, G., Landen, M., Landman, B.A., Lauriello, J., Lawrie, S.M., Lee, P.H., Le Hellard, S., Lemaitre, H., Leonardo, C.D., Li, C.S., Liberg, B., Liewald, D.C., Liu, X., Lopez, L.M., Loth, E., Lourdusamy, A., Luciano, M.,

Macciardi, F., Machielsen, M.W., Macqueen, G.M., Malt, U.F., Mandl, R., Manoach, D.S., Martinot, J.L., Matarin, M., Mather, K.A., Mattheisen, M., Mattingsdal, M., Meyer-Lindenberg, A., McDonald, C., McIntosh, A.M., McMahon, F.J., McMahon, K.L., Meisenzahl, E., Melle, I., Milaneschi, Y., Mohnke, S., Montgomery, G.W., Morris, D.W., Moses, E.K., Mueller, B.A., Munoz Maniega, S., Muhleisen, T.W., Muller-Myhsok, B., Mwangi, B., Nauck, M., Nho, K., Nichols, T.E., Nilsson, L.G., Nugent, A.C., Nyberg, L., Olvera, R.L., Oosterlaan, J., Ophoff, R.A., Pandolfo, M., Papalampropoulou-Tsiridou, M., Papmeyer, M., Paus, T., Pausova, Z., Pearson, G.D., Penninx, B.W., Peterson, C.P., Pfennig, A., Phillips, M., Pike, G.B., Poline, J.B., Potkin, S.G., Putz, B., Ramasamy, A., Rasmussen, J., Rietschel, M., Rijpkema, M., Risacher, S.L., Roffman, J.L., Roiz-Santianez, R., Romanczuk-Seiferth, N., Rose, E.J., Royle, N.A., Rujescu, D., Ryten, M., Sachdev, P.S., Salami, A., Satterthwaite, T.D., Savitz, J., Saykin, A.J., Scanlon, C., Schmaal, L., Schnack, H.G., Schork, A.J., Schulz, S.C., Schur, R., Seidman, L., Shen, L., Shoemaker, J.M., Simmons, A., Sisodiya, S.M., Smith, C., Smoller, J.W., Soares, J.C., Sponheim, S.R., Sprooten, E., Starr, J.M., Steen, V.M., Strakowski, S., Strike, L., Sussmann, J., Samann, P.G., Teumer, A., Toga, A.W., Tordesillas-Gutierrez, D., Trabzuni, D., Trost, S., Turner, J., Van den Heuvel, M., van der Wee, N.J., van Eijk, K., van Erp, T.G., van Haren, N.E., van 't Ent, D., van Tol, M.J., Valdes Hernandez, M.C., Veltman, D.J., Versace, A., Volzke, H., Walker, R., Walter, H., Wang, L., Wardlaw, J.M., Weale, M.E., Weiner, M.W., Wen, W., Westlye, L.T., Whalley, H.C., Whelan, C.D., White, T., Winkler, A.M., Wittfeld, K., Woldehawariat, G., Wolf, C., Zilles, D., Zwiers, M.P., Thalamuthu, A., Schofield, P.R., Freimer, N.B., Lawrence, N.S., Drevets, W., Alzheimer's Disease Neuroimaging Initiative, E.C.I.C.S.Y.S.G., 2014. The ENIGMA Consortium: large-scale collaborative analyses of neuroimaging and genetic data. *Brain Imaging Behav* 8, 153-182.

Toga, A.W., Thompson, P.M., 2003. Mapping brain asymmetry. *Nat Rev Neurosci* 4, 37-48.

Turner, B.O., Marinsek, N., Ryhal, E., Miller, M.B., 2015. Hemispheric lateralization in reasoning. *Ann N Y Acad Sci* 1359, 47-64.

Tzourio-Mazoyer, N., Mazoyer, B., 2017. Variations of planum temporale asymmetries with Heschl's Gyri duplications and association with cognitive abilities: MRI investigation of 428 healthy volunteers. *Brain Struct Funct* 222, 2711-2726.

Tzourio-Mazoyer, N., Petit, L., Zago, L., Crivello, F., Vinuesa, N., Joliot, M., Jobard, G., Mellet, E., Mazoyer, B., 2015. Between-hand difference in ipsilateral deactivation is associated with hand lateralization: fMRI mapping of 284 volunteers balanced for handedness. *Frontiers in Human Neuroscience* 9.

Van Essen, D.C., 2004. Surface-based approaches to spatial localization and registration in primate cerebral cortex. *Neuroimage* 23, S97-S107.

Van Essen, D.C., Glasser, M.F., Dierker, D.L., Harwell, J., Coalson, T., 2012. Parcellations and hemispheric asymmetries of human cerebral cortex analyzed on surface-based atlases. *Cereb Cortex* 22, 2241-2262.

Vigneau, M., Beaucousin, V., Herve, P.Y., Duffau, H., Crivello, F., Houde, O., Mazoyer, B., Tzourio-Mazoyer, N., 2006. Meta-analyzing left hemisphere language areas: Phonology, semantics, and sentence processing. *Neuroimage* 30, 1414-1432.

Vigneau, M., Beaucousin, V., Herve, P.Y., Jobard, G., Petit, L., Crivello, F., Mellet, E., Zago, L., Mazoyer, B., Tzourio-Mazoyer, N., 2011. What is right-hemisphere contribution to phonological, lexico-semantic, and sentence processing? Insights from a meta-analysis. *Neuroimage* 54, 577-593.

Watkins, K.E., Paus, T., Lerch, J.P., Zijdenbos, A., Collins, D.L., Neelin, P., Taylor, J., Worsley, K.J., Evans, A.C., 2001. Structural asymmetries in the human brain: a voxel-based statistical analysis of 142 MRI scans. *Cereb Cortex* 11, 868-877.

Willems, R.M., Peelen, M.V., Hagoort, P., 2010. Cerebral lateralization of face-selective and body-selective visual areas depends on handedness. *Cereb Cortex* 20, 1719-1725.

Willems, R.M., Van der Haegen, L., Fisher, S.E., Francks, C., 2014. On the other hand: including left-handers in cognitive neuroscience and neurogenetics. *Nat Rev Neurosci* 15, 193-201.

Winkler, A.M., Kochunov, P., Blangero, J., Almasy, L., Zilles, K., Fox, P.T., Duggirala, R., Glahn, D.C., 2010. Cortical thickness or grey matter volume? The importance of selecting the phenotype for imaging genetics studies. *Neuroimage* 53, 1135-1146.

Xiang, H.D., Fonteijn, H.M., Norris, D.G., Hagoort, P., 2010. Topographical Functional Connectivity Pattern in the Perisylvian Language Networks. *Cerebral Cortex* 20, 549-560.

Yakovlev, P., 1972. A proposed definition of the limbic system. In: Hockman, C. (Ed.), *Limbic System Mechanisms and Autonomic Function*. Charles C. Thomas, Springfield, IL, pp. 241-283.

Yan, C., Gong, G., Wang, J., Wang, D., Liu, D., Zhu, C., Chen, Z.J., Evans, A., Zang, Y., He, Y., 2011. Sex- and Brain Size-Related Small-World Structural Cortical Networks in Young Adults: A DTI Tractography Study. *Cereb Cortex* 21, 449-458.

Yucel, K., McKinnon, M., Chahal, R., Taylor, V., Macdonald, K., Joffe, R., MacQueen, G., 2009. Increased subgenual prefrontal cortex size in remitted patients with major depressive disorder. *Psychiatry Research-Neuroimaging* 173, 71-76.

Yucel, M., Stuart, G.W., Maruff, P., Wood, S.J., Savage, G.R., Smith, D.J., Crowe, S.F., Copolov, D.L., Velakoulis, D., Pantelis, C., 2002. Paracingulate morphologic differences in males with established schizophrenia: a magnetic resonance imaging morphometric study. *Biol Psychiatry* 52, 15-23.

Yucel, M., Wood, S.J., Phillips, L.J., Stuart, G.W., Smith, D.J., Yung, A., Velakoulis, D., McGorry, P.D., Pantelis, C., 2003. Morphology of the anterior cingulate cortex in young men at ultra-high risk of developing a psychotic illness. *British Journal of Psychiatry* 182, 518-524.

Zago, L., Petit, L., Jobard, G., Hay, J., Mazoyer, B., Tzourio-Mazoyer, N., Karnath, H.O., Mellet, E., 2017. Pseudoneglect in line bisection judgement is associated with a modulation of right hemispheric spatial attention dominance in right-handers. *Neuropsychologia* 94, 75-83.

Zhen, Z., Kong, X.Z., Huang, L., Yang, Z., Wang, X., Hao, X., Huang, T., Song, Y., Liu, J., 2017. Quantifying the variability of scene-selective regions: Interindividual, interhemispheric, and sex differences. *Hum Brain Mapp* 38, 2260-2275.

Zhen, Z., Yang, Z., Huang, L., Kong, X.Z., Wang, X., Dang, X., Huang, Y., Song, Y., Liu, J., 2015. Quantifying interindividual variability and asymmetry of face-selective regions: a probabilistic functional atlas. *Neuroimage* 113, 13-25.

Zhou, D., Lebel, C., Evans, A., Beaulieu, C., 2013. Cortical thickness asymmetry from childhood to older adulthood. *Neuroimage* 83, 66-74.

Figure S1. Mixed results of cortical thickness asymmetry in previous publications.

