

Structures of the stator complex that drives rotation of the bacterial flagellum

Justin C. Deme^{1,2}, Steven Johnson¹, Owen Vickery^{3,4}, Amy Muellbauer¹, Holly Monkhouse¹, Thomas Griffiths¹, Rory Hennell James¹, Ben C. Berks³, James W. Coulton^{5,6}, Phillip J. Stansfeld^{3,4} & Susan M. Lea^{1,2,*}

1. Sir William Dunn School of Pathology, University of Oxford, South Parks Road, Oxford OX1 3RE, UK
2. Central Oxford Structural Molecular Imaging Centre, University of Oxford, South Parks Road, Oxford, OX1 3RE, UK
3. Department of Biochemistry, University of Oxford, South Parks Road, Oxford OX1 3QU
4. School of Life Sciences & Department of Chemistry, University of Warwick, Coventry, CV4 7AL, UK.
5. Department of Microbiology and Immunology, McGill University, 3775 University Street, Montreal, H3A 2B4, Canada
6. Département de biochimie et médecine moléculaire, Université de Montréal, 2900 boulevard Édouard-Montpetit, Montréal, H3T 1J4, Canada

*For Correspondence: susan.lea@path.ox.ac.uk

Summary

The bacterial flagellum is the proto-typical protein nanomachine and comprises a rotating helical propeller attached to a membrane-embedded motor complex¹. The motor consists of a central rotor surround by stator units that couple ion flow across the cytoplasmic membrane to torque generation. Here we present the structures of stator complexes from multiple bacterial species, allowing interpretation of the extensive body of data on stator mechanism. The structures reveal an unexpected asymmetric A₅B₂ subunit assembly in which the five A subunits enclose the two B subunits. Comparison to novel structures of other ion-driven motors indicates that this A₅B₂ architecture is fundamental to bacterial systems that

couple energy from ion-flow to generate mechanical work at a distance, and suggests that such events involve rotation in the motor structures.

Main

A motor is a machine that supplies motive power for a device with moving parts. Biological systems use both linear and rotary motors to generate a variety of outputs. One of the most fascinating and complex biological rotary motors is the flagellar apparatus used by bacteria to propel themselves through fluid environments. Although bacterial swimming was first observed in the 17th century², a mechanistic understanding of how the bacterial flagellum generates rotation is still lacking. The core of the flagellum is a highly conserved motor (Fig. 1a) consisting of a cytoplasmic-membrane embedded rotor complex surrounded by varying numbers of stator complexes (hereafter termed simply stators) that generate torque³. While high resolution information has recently been obtained for the rotor component⁴, structural detail of the stators has thus far been limited to modelling studies⁵.

Stators harvest energy from either H⁺ or Na⁺ ion flow across the cytoplasmic membrane, generating torque in the cytoplasmic portion (C-ring) of the rotor complex⁶⁻⁹. Chimeras between H⁺- and Na⁺-dependent stators are functional, implying that the mechanism converting ion flow into work is the same for the two coupling ions¹⁰. Stators are built from two cytoplasmic membrane proteins, which for simplicity are generically referred to here as MotA and MotB. MotA is predicted to contain four transmembrane helices (TMH) with a large cytoplasmic insertion between TMH2 and TMH3. MotB is predicted to contain a short cytoplasmic sequence, a single TMH, and a C-terminal peptidoglycan binding (PGB) domain. Early biochemical work defined the stator stoichiometry as MotA₄B₂¹¹, and this subunit composition has informed attempts to derive mechanism for conversion of ion flow

into rotation (reviewed in ¹²). Extensive experimental studies have led to a model of stator function in which docking of the MotA cytoplasmic loop to the rotor C-ring simultaneously induces ion permeation through the stator and release of the MotB-PGB domain to bind to the peptidoglycan (PG) surrounding the flagellar basal body^{13,14}. Ion flow is proposed to lead to conformational changes in the cytoplasmic domain of MotA that generate torque in the rotor¹⁵⁻¹⁷. In the absence of a stator structure various mechanistic hypotheses have been proposed to explain the coupling of ion flow to conformational change, most of which explicitly use the predicted 2-fold symmetry of a MotA₄B₂ complex¹⁷⁻¹⁹.

Flagellar stators are MotA₅B₂ complexes

We used cryo-electron microscopy (cryo-EM) to study stator complexes from a range of bacterial species with different ion specificities (Extended Data Fig. 1). Two-dimensional class averages of the complexes from three species (*Vibrio mimicus*, *Clostridium sporogenes* and *Bacillus subtilis*) clearly showed a distorted, pentagonal, structure (Fig. 1b). 3D reconstructions of these complexes yielded volumes that could only be interpreted as MotA₅B₂ assemblies (Fig. 1c; Extended Data Fig. 2), with five copies of MotA fully enclosing the TMHs of two copies of MotB (Fig. 2a). Although we do not observe the PGB-domains of MotB in the resolved structures, these domains must be present in the imaged complexes because the stators were purified using an affinity tag located after the PGB domain. Thus, the PGB-domain of MotB has no fixed location with respect to the core complex in the context of the isolated protein. The stator structures are compatible with sequence conservation data, with inter- and intra-molecular co-evolution data, and with published cysteine crosslinking^{20,21} and tryptophan scanning mutagenesis^{22,23} (Extended data Fig. 3).

The four TMHs of MotA are arranged in two layers. TMH3 and TMH4 line the central pore, while TMH1 and TMH2 form a surrounding outer layer of helices (Fig. 2a,b). TMH1 and TMH2 are not in contact with each other within a single subunit but instead interact between adjacent subunits, thereby stabilising the MotA assembly. Immediately following TMH2 there is an amphipathic helix (AMPH) running perpendicular to the TMHs at the cytoplasmic membrane surface, with the five copies of this helix forming a belt around the outside of the structure. TMH3 and TMH4 extend 30 Å outside the membrane to form the core of the MotA cytoplasmic domain, with the rest of the domain built from helices inserted in the loop between the AMPH and TMH3. Both within and outside the membrane domain the pentameric arrangement of MotA is distorted (Fig. 2c). Charged residues shown to be essential for the interaction of the stator with the rotor C-ring²⁴ are located towards the base of this domain, forming a ring that decorates the surface of the pentamer (Extended data Fig. 4).

The TMHs of the two copies of MotB are located in the central pore of the distorted MotA pentamer, with their hydrophobic sidechains completely buried within the MotA ring. From the N-terminal ends of the MotB TMHs clear densities extend down to contact the inner surfaces of TMH3 and TMH4 in the cytoplasmic domains of MotA (Fig. 3a,b; Extended Data Fig. 5a,b). Although the densities are too weak for the sequence to be traced, they are of sufficient length to account for most of the MotB N-terminus, including a cluster of positive charges essential for motor function²⁵. At the non-cytoplasmic face of the complex the MotB TMHs emerge vertically from the MotA pentamer and are followed by another short helical section that packs down between the TMH3-TMH4 loops of the MotA chains. The connectivity of these densities defines them as the plug helices previously implicated by mutagenesis as critical to sealing the complexes in an off state²⁶.

Prior mutagenesis studies have established that a series of conserved residues in the TMHs of MotA and MotB are important for flagellar motion and/or ion-flow through the stator (reviewed in ³). Invariant MotB_{D32} (using the *Escherichia coli* numbering system) is the key protonatable residue and both copies are seen to lie within a ring formed by the five copies of another invariant polar residue, MotA_{T209} (Fig. 3c). A second Thr residue (at a position corresponding to residue A180 in *E. coli* MotA) that is conserved in the Na⁺-dependent stators also contributes to this ring, and forms part of a track of Na⁺/H⁺ specificity determining residues that line the inner surface of the MotA pore (Extended data Fig. 1b). Two conserved Pro residues in MotA have been shown to be important for torque generation²⁷. One of these, MotA_{P222} can now be seen to be required for contacts between neighbouring MotA monomers. The other, MotA_{P173}, forms a second ring of conserved residues with invariant MotA_{Y217}, two helical turns down from the Thr ring. This hydrophobic ring contacts MotB at the completely conserved MotB_{W26}. A MotB_{W26A} substitution completely abolished motility confirming the importance of this contact (Fig. 3d).

Asymmetry and the implications for activation of ion flow

The 5:2 stoichiometry of the stator complex leads to multiple levels of asymmetry in the structure (Fig. 2c; Extended Data Fig. 5c). The pentagon formed by the MotA subunits within the membrane is distorted to accommodate and seal around the two MotB TMHs. The asymmetry of this part of the complex is also driven by the two MotB plug helices sitting between the MotA loops, which divide the MotA chains into two groups separated in the extracytoplasmic region. Removal of the MotB plug has been shown to lead to uncontrolled ion flow through the MotAB channel²⁶. However, our structures show that the plug helices are not the sole block to ion permeation since there are no detectable channels across the

cytoplasmic membrane compartment (Fig. 3e; Extended Data Fig. 5d). Embedding plug-free structures in full lipid bilayer models and running extended simulations demonstrated the observed structures are stable, low energy, states (Extended Data Fig. 6). No ion permeation across the bilayer was seen in any simulation, supporting the idea that the complexes currently seen will require rearrangement for activity.

Activation of ion flow is proposed to be triggered by docking of the inactive stator onto the flagellar C-ring via the MotA cytoplasmic domains resulting in signal propagation from the cytoplasm to the plug region and plug release^{13,14}. Our structures reveal two potential routes for such a signal. The first involves the cytoplasmic N-termini of the MotB subunits which contain functionally essential residues²⁵ that interact with the inside of the MotA pentamer through highly evolutionarily coupled contacts (Extended data Fig. 3d). C-ring-induced movement of MotA would be communicated to MotB at this site leading to alterations at the opposite end of the MotB TMHs. The second possible route of signal propagation is directly through the MotA subunits, with hinging of the long TMH3 and TMH4 helices altering the conformation of the plug helix binding loops to allow plug release. Our structures provide insight into the conformational changes that the MotA cytoplasmic domains can undergo. The structures show differing degrees of hinging of the MotA cytoplasmic domains relative to the membrane embedded helices (Fig. 3f; Extended Data Fig. 5e) suggesting changes in the degree of asymmetry may link to functional state. Our structures also reveal that the two MotA residues known to be essential for interaction with the C-ring protein Flg²⁴ are located on opposite sides of the MotA cytoplasmic domain, with MotA_{R90} from one copy facing MotA_{E98} from the neighbouring copy (Extended data Fig. 4). Therefore docking of the C-terminal domain of Flg between two MotA subunits could trigger conformational change in the stator.

Coupling of ion-flow to flagellar rotation

The most striking feature of the asymmetry of the 5:2 subunit stoichiometry is that it places the TMHs of the two copies of MotB, including the critical MotB_{D32} residue, in different environments within the distorted MotA pentagon. The system is therefore primed for differential binding of H⁺ or Na⁺ at the critical MotB_{D32} residue to induce changes in the relative positioning of the MotB and MotA helices, most likely by rotation of one relative to the other. Because MotB becomes tethered to the PG upon stator activation¹³ (Fig. 4a) it follows that the MotA ring rotates around the MotB dimer. The 5:2 subunit stoichiometry implies a model whereby a single channel opens to allow ion binding to MotB_{D32} on one MotB triggering rotation of the MotA ring by ~36° (Fig. 4b). This motion would bring the second MotB chain into the same position relative to the surrounding MotA subunits as the starting arrangement of the first MotB chain, closing the first channel and opening the second. Each subsequent ion binding event would trigger a further ratchet motion of 36°, with each turn of the MotA cytoplasmic domains providing a “power stroke” to the rotor. An alternative model in which the second ion binding event leads to a reset to the original position and full rotation does not occur, would also be compatible with the structure. Such a mechanism would still be capable of driving full rotation of the rotor component, acting like an energised escapement mechanism.

Any mechanism for coupling ion flow to flagellar rotation must also explain how the direction of rotation of the flagellum can reverse in response to chemotactic stimuli. All experimental evidence (reviewed in ²⁸) shows that the chemotaxis machinery acts on the FlhG subunit of the C-ring rather than the stator. Our unidirectional rotation model for the stator mechanism can account for flagellar reversal if the chemotaxis-linked conformational

changes induced in the C-ring lead to an alteration in the side of the stator that is driving the rotation (Fig. 4c). Consistent with this proposal large conformational changes in the stator-interacting Flg component of the C-ring have been observed in crystal structures of Flg fragments, where 180° rotations of the C-terminal domain relative to the middle domain have been observed²⁹. The model also predicts that any reversal of the ion flow through the stator would have the potential to reverse the direction of flagellar rotation even in the absence of switching by the chemotaxis machinery, and this phenomenon has been observed in *Streptococcus* species assayed under high pH conditions^{30,31}.

Common architecture across multiple bacterial ion-driven machines

The MotAB system is related at the sequence level to the ExbBD complex found in Gram-negative bacteria that uses ion-flow across the cytoplasmic membrane to power transport processes at the outer membrane via the trans-periplasmic TonB protein³². We determined cryo-EM structures of ExbBD complexes from *E. coli* and *Pseudomonas savastanoi* (Extended Data Fig. 7). Both displayed a 5:2 ExbB:ExbD stoichiometry that differs from the subunit composition of earlier structures^{33,34}, but agrees with the subunit stoichiometry of a novel structure of the *E. coli* ExbBD reported whilst this manuscript was in preparation³⁵. Comparison of these new ExbB₅D₂ structures to the stator complexes reveals a high level of structural conservation, particularly within the membrane domain (Extended Data Fig. 8a,b). Both the flattened pentagon geometry and the alignment of mechanistically important residues, such as the conserved Asp within a ring of Thr residues, suggest that the two systems use the same molecular mechanism. We therefore predict that the ExbB will rotate relative to the ExbD helices in response to proton flow. Outside the core TMH region there are structural differences between the systems that presumably reflect their very different

biologies. ExbB is very differently elaborated relative to MotA, with only one TMH packing across the pair of helices that form the core inner ring and no bracing helices strengthening packing between subunits (Extended Data Fig. 8c). The ExbB cytoplasmic domains are only superficially related to the corresponding MotA domain and lack the short pair of C-terminal helices found in MotA (Extended Data Fig. 8d).

P. savastanoi ExbB and ExbD were purified as a complex with TonB when all three proteins were co-expressed (Extended data Fig. 9a). However, no extra density was observed in the cryo-EM maps of this complex relative to the ExbBD complex alone, suggesting that TonB is located on the outside of the ExbBD complex and dissociates upon sample freezing. A peripheral location for TonB is consistent with both co-evolution and mutagenesis/suppressor data³⁶, which suggest that the TonB binding site is on the outside of the ExbB transmembrane domain (Extended data Fig. 9b). TonB consists of a single pass TMH, followed by an extended periplasmic region that interacts with the periplasmic domain of ExbD³⁷, and terminates in a folded domain that links with outer-membrane receptor proteins³⁸. We speculate that the TonB TMH packs against the exterior of the ExbBD complex so that conformational change in TonB is driven by rotation of the ExbB component relative to ExbD. By extension, the homologous TolQRA system will also share this architecture and be mechanistically related³⁹.

Bacteria from the *Bacteroidetes* phylum possess a motor complex that harvests energy from ion-flow to drive protein secretion and to power bacterial motility via a non-flagellar mechanism termed gliding motility⁴⁰. The structure of this complex is described in a companion paper (Hennell-James et al, companion paper). Although the constituent GldL and GldM subunits of this motor have no sequence similarity to the subunits of the MotAB or ExbBD complexes, the *Bacteroidetes* motor complex exhibits the same 5:2 subunit

stoichiometry as these complexes (Fig. 5a). All three complexes have an intramembrane core consisting of a central subunit TMH dimer surrounded by a 10 TMH ring. Structural comparisons demonstrate the similarity between the three motors in the arrangement of this intramembrane core and of the height within the membrane at which charged residues critical to function are located (Fig. 5b,c). Such shared underlying architecture between otherwise highly dissimilar motors (Fig. 5d) implies an unexpected commonality in their mechanism.

Acknowledgements


We thank David Blair for providing *E. coli* RP6894. We thank E. Johnson and A. Costin of the Central Oxford Structural Molecular Imaging Centre (COSMIC) for assistance with data collection, and H. Elmlund (Monash) for access to SIMPLE code ahead of release. We acknowledge the use of the Central Oxford Structural Microscopy and Imaging Centre (COSMIC). The Central Oxford Structural Microscopy and Imaging Centre is supported by the Wellcome Trust (grant no. 201536), The EPA Cephalosporin Trust, The Wolfson Foundation and a Royal Society/Wolfson Foundation Laboratory Refurbishment Grant (no. WL160052). Research in S.M.L.'s laboratory is supported by a Wellcome Trust Investigator Award (grant no. 100298), a Collaborative award (no. 209194) and an Medical Research Council (London) Programme Grant (no. MR/M011984/1). Research in B.C.B.'s laboratory is supported by a Wellcome Trust Investigator Award (grant no. 107929/Z/15/Z). Research in J.W.C.'s laboratory is supported by the Canadian Institutes of Health Research (grant 178048-BMA-CFAA-11449). Research in P.J.S.'s lab is funded by Wellcome (208361/Z/17/Z), the MRC (MR/S009213/1) and BBSRC (BB/P01948X/1, BB/R002517/1 and BB/S003339/1). This project made use of time on ARCHER and JADE granted via the UK High-End Computing Consortium

for Biomolecular Simulation, HECBioSim (<http://hecbiosim.ac.uk>), supported by EPSRC (grant no. EP/R029407/1), and Athena at HPC Midlands+, which was funded by the EPSRC on grant EP/P020232/1, and used the University of Warwick Scientific Computing Research Technology Platform for computational access.

Author contributions

J.C.D carried out all biochemical work except as credited otherwise, prepared cryo-EM grids, collected and processed EM data and determined the structures. J.C.D., S.J., and S.M.L designed the project, interpreted the data, built models, and wrote the first draft of the paper. S.J. also performed MALS experiments. O.V. and P.J.S. performed molecular dynamics simulations. A.M., H.M., and T.G. carried out biochemical work on *Pseudomonas* TonB-ExbB-ExbD. R.H.J. and B.C.B. contributed the GldLM structure. J.W.C. initiated and provided materials for the ExbBD project. All authors commented on drafts of the manuscript.

262 **Figures**



263
264

265 **Figure 1. Stators from multiple organisms have a MotA₅MotB₂ stoichiometry.** **a**, Composite
266 cartoon showing the general organisation of bacterial flagellar complexes in Gram-negative
267 (left side) and Gram-positive (right side) bacteria with major components labelled. Stators are
268 orange and rotor components, MS- and C-ring, are grey. OM, outer membrane; CM,
269 cytoplasmic membrane; PG, peptidoglycan. **b**, 2D class averages of cryo-EM particles of
270 stators from the bacterial species indicated. Upper panels are representative ‘top’ views of
271 the 5:2 complexes. Lower panels are ‘side’ views from data collected in the presence of
272 fluorinated octyl maltoside. **c**, Cryo-EM volumes of stators from the three bacterial species.
273 The MotA subunits are coloured pink, blue, green, teal, and yellow, and the centrally-located
274 MotB subunits white and dark grey. Bound detergent is shown as transparent density at the

275 periphery. Upper panels show views from the cytoplasm, lower panels show side views with
 276 the likely membrane location (assigned from the position of the detergent micelle and from
 277 simulations; Extended Data Fig. 6) indicated by the grey bar.
 278




Figure 2. Structures of stators from *C. sporogenes* and *B. subtilis*. **a**, The *C. sporogenes* (left) and *B. subtilis* (right) stators are shown as cartoon representations and coloured as in Fig. 1c. Upper panel, side view with membrane indicated in grey. Lower panel, view from the cytoplasm. **b**, Two views of a single MotA subunit (*C. sporogenes*) coloured from blue at the N-terminus to red at the C-terminus. **c**, Slabs (viewed from cytoplasm) through the *C. sporogenes* complex at the indicated positions on the inset structure (arrow indicates the cytoplasmic side of the complex). Distortion of the MotA subunits from a regular pentagon arrangement becomes more extreme in the cytoplasmic regions.




Figure 3. Functionally critical regions of the stator complex. **a**, Structure of the *C. sporogenes* stator shown as cartoon representations coloured as in Figs. 1 and 2 with the unmodeled density for the MotB N-terminal extensions (Left) and plug helices (Right) shown. **b**, (i) Isolated MotB dimer extracted from the *C. sporogenes* stator and (ii) superposition of the TMHs of the two MotB chains showing the relative rotation of the N-terminal extensions and plug helices. **c**, The environment around MotB_{D32} within the membrane. (Left) Only the core MotA helices within the transmembrane region are shown and the two copies of MotA at the front of the view are removed. (Right) Slabs through the stator core at the indicated heights. Residue numbering is that of the *E. coli* MotAB stator but displayed on the *C. sporogenes* stator structure. **d**, Motility in soft agar of *E. coli* RP6894 (Δ *motAB*) complemented with plasmids expressing *motAB* with the indicated mutations or the vector control (vector_{ctrl}). **e**, Surface representation of the model shows close packing. (Top) Side view with front of complex removed. (Bottom) Top-down view of the slab indicated by dashed lines. **f**, Overlay of the five copies of the *C. sporogenes* MotA chain reveals they fall into two conformational classes which differ in the degree of flexing at the highlighted prolines.




Figure 4. Mechanistic model for the generation of bi-directional flagellar torque. **a**, Activation of the stator complex from the structurally resolved state (here termed ‘relaxed’) to form a ‘tensioned’ state permissive to ion-flow. This conformational change is likely driven by interactions between the C-terminal peptidoglycan binding domains (black and white ovals) of MotB and the peptidoglycan layer, as well as interactions between the stator and the C-ring complex. **b**, Cartoon showing top views of the intra-membrane core of the stator complex with five MotA chains surrounding two MotB subunits. Bound ions are shown as red spheres. Ion flow leads to rotation of the MotA ring through alternating formation of MotB-ion-MotA interactions by the two MotB chains that processes around the surrounding MotA subunits. **c**, Model describing how a stator that rotates in one direction can drive either clockwise (CW) or counterclockwise (CCW) rotation of the flagellum depending on the conformational state of the C-ring.




Figure 5. Conservation of core architecture between diverse families of ion-driven motors.

a, Representatives of three ion-driven motor families that share a common structural core. Complexes are shown as grey cartoons with helices equivalent to those in the MotA inner ring displayed in coloured surface representation. **b**, Overlay of the three complexes (common core in grey cartoons, other structure semi-transparent cartoons). Mechanistically essential charged residues within the common core (space filling side chains; C, grey; O, red) occur at the same height with respect to the membrane irrespective of whether they occur on the MotA- or MotB-equivalent chain. **c**, Overlay of the common core of the three complexes. The distortion from pentamer symmetry within the membrane is shared between all three families **d**, Cartoon summarising the updated view of how the three families of ion-driven motors are coupled to their different biological effects. Note that ion movement drives rotation of the central subunits in GldLM but of the peripheral subunits in MotAB/ExbBD. OM, outer membrane; CM, cytoplasmic membrane; PG, peptidoglycan

Table 1. Cryo-EM data collection, refinement and validation statistics

	<i>C. sporogenes</i> MotAB (EMDB-10895) (PDB ID 6YSF)	<i>B. subtilis</i> MotAB (EMDB-10899) (PDB ID 6YSL)	<i>V. mimicus</i> PomAB (EMDB-10901)	<i>E. coli</i> ExbBD (EMDB-10902)	<i>P. savastanoi</i> ExbBD (EMDB-10897)
Data collection and processing					
Magnification	165,000	165,000	165,000	165,000	165,000
Voltage (kV)	300	300	300	300	300
Electron exposure (e-/Å ²)	48	48	48	48	48
Defocus range (µm)	1.0-3.0	1.0-3.0	1.0-3.0	1.0-2.5	1.0-3.0
Pixel size (Å)	0.822	0.822	0.822	0.822	0.822
Symmetry imposed	C1	C1	C1	C1	C1
Initial particle images (no.)	1,998,900	1,532,430	2,383,022	2,045,350	1,342,937
Final particle images (no.)	314,230	122,615	244,654	227,700	65,617
Map resolution (Å)	3.4	3.5	4.2	4.6	3.8
FSC threshold	0.143	0.143	0.143	0.143	0.143
Map resolution range (Å)	3.2-5.2	3.3-6.0	3.9-6.5	4.0-6.5	3.6-6.1
Refinement					
Initial model used (PDB code)	None	None			
Model resolution (Å)	3.4	3.5			
FSC threshold	0.143	0.143			
Model resolution range (Å)	3.2-5.2	3.3-6.0			
Map sharpening B factor (Å ²)	-117	-104			
Model composition					
Non-hydrogen atoms	10220	10128			
Protein residues	1327	1324			
Ligands	0	0			
B factors (Å²)					
Protein	53	99			
Ligand	NA	NA			
R.m.s. deviations					
Bond lengths (Å)	0.004	0.007			
Bond angles (°)	0.757	1.425			
Validation					
MolProbity score	2.15	1.85			
Clashscore	13.66	5.93			
Poor rotamers (%)	0.90	0.47			
Ramachandran plot					
Favored (%)	91.47	90.92			
Allowed (%)	8.53	9.01			
Disallowed (%)	0.00	0.08			

Methods

Bacterial strains and plasmids

Bacterial strains and plasmids used in this study are listed in Supplementary Table 1. The pT12 backbone used for all protein expression was derived from Kuhlen *et al*⁴¹. Plasmids were generated by Gibson assembly of PCR fragments using the NEBuilder HiFi Master Mix (NEB). Fragments were created by PCR with the relevant primers (listed in Supplementary Table 2) using Q5 polymerase (NEB) and genomic DNA templates obtained from the Leibniz Institute [dsmz.de]: *Vibrio mimicus* (DSM 19130), *Bacillus subtilis* 168 (DSM 402), *Clostridium sporogenes* 388 (DSM 795), *Escherichia coli* W (DSM 1116), *Pseudomonas savastanoi*, pv. phaseolicola 1448A (DSM 21482). Gibson assembly and PCR were carried out following the manufacturer's recommendations. *E. coli* RP6894 (Δ motAB) for motility assays was generated by J. S. Parkinson and gifted by D.F. Blair.

Purification of MotAB/PomAB and ExbBD complexes

V. mimicus PomAB, its derivative PomAB_{Δ61-120} lacking unstructured periplasmic residues of PomB, *B. subtilis* MotAB, *C. sporogenes* MotAB, *E. coli* ExbBD, and *P. savastanoi* TonB-ExbBD complexes were expressed in *E. coli* MT56 as a single operon from a pT12 vector encoding a C-terminal twin-strep tag. Purification steps were similar across all constructs and carried out at 4 °C. Briefly, cells were grown at 37 °C for 16 h in TB media containing kanamycin (50 µg/mL) and rhamnose monohydrate (0.1% w/v) then collected by centrifugation at 4,000g. Cell pellets were resuspended in TBS (100 mM Tris, 150 mM NaCl, 1 mM EDTA pH 8.0) plus 30 µg/mL DNase I and 400 µg/mL lysozyme for 30 mins before passage through an EmulsiFlex C5 homogenizer (Avestin) at 15,000 psi. Unbroken cells were

removed by centrifugation at 24,000g for 20 min. The supernatant was recovered and total membranes were collected by centrifugation at 200,000g for 1.5 h. Membranes were resuspended in TBS and solubilized by incubation with 1% (w/v) lauryl maltose neopentyl glycol (LMNG; Anatrace) for 2 h. Insoluble material was removed by centrifugation at 100,000g for 30 min. Solubilized membranes were then applied to a Streptactin XT column (IBA Lifesciences). The resin was washed with 10 column volumes (CV) of TBS containing 0.02% (w/v) LMNG and proteins were eluted in 5 CV of TBS supplemented with 0.01% (w/v) LMNG and 50 mM D-biotin (IBA Lifesciences). Eluates were concentrated using a 100-kDa molecular weight cutoff (MWCO) Vivaspin 6 (GE Healthcare) centrifugal filter unit and injected onto a Superose 6 Increase 10/300 GL size exclusion column (GE Healthcare) pre-equilibrated in TBS plus 0.01% (w/v) LMNG. Peak fractions were collected and concentrated using a 100-kDa MWCO Vivaspin 500 (GE Healthcare) centrifugal filter unit.

For *P. savastanoi* TonB-ExbBD and ExbBD complexes, SEC-MALS analysis was carried out by injecting 100 μ L ($A_{280\text{nm}} = 1.0$) of either sample onto a Superose 6 increase 10/300 GL (GE Healthcare) equilibrated in TBS containing 0.02% (w/v) LMNG. Light scattering and refractive index changes were measured using a Dawn Heleos-II light-scattering detector and an Optilab-TrEX refractive index monitor. Analysis was carried out using ASTRA 6.1.1.17 software using a theoretical extinction coefficient of 1.02 ($\text{Abs}_{0.1\%}$) and a protein dn/dc value of 0.186 mL/g and a detergent dn/dc value of 0.143 mL/g.

Cryo-EM sample preparation and imaging

Purified complexes (4 μ L each) of *V. mimicus* PomAB ($A_{280\text{nm}} = 0.5$), PomAB $_{\Delta 61-120}$ ($A_{280\text{nm}} = 0.55$), *B. subtilis* MotAB ($A_{280\text{nm}} = 1.0$), *C. sporogenes* MotAB ($A_{280\text{nm}} = 0.8$), *E. coli* ExbBD ($A_{280\text{nm}} = 3.2$), or *P. savastanoi* TonB-ExbBD ($A_{280\text{nm}} = 2.0$) were adsorbed to glow-

discharged holey carbon-coated grids (Quantifoil 300 mesh, Au R1.2/1.3) for 10 s. Grids were then blotted for 2 s at 100% humidity at 8°C and frozen in liquid ethane using a Vitrobot Mark IV (FEI). Alternatively, specimens were prepared by supplementing *V. mimicus* PomAB ($A_{280\text{nm}} = 2.3$), PomAB $_{\Delta 61-120}$ ($A_{280\text{nm}} = 3.7$), *B. subtilis* MotAB ($A_{280\text{nm}} = 7.2$), *C. sporogenes* MotAB ($A_{280\text{nm}} = 8.6$), *E. coli* ExbBD ($A_{280\text{nm}} = 4.2$) with 0.7 mM fluorinated octyl maltoside (fluo OM; Anatrace) prior to grid preparation.

Data were collected in counting mode on a Titan Krios G3 (FEI) operating at 300 kV with a GIF energy filter (Gatan) and K2 Summit detector (Gatan) using a pixel size of 0.822 Å and a total dose of 48 e⁻/Å² spread across 20 or 32 fractions. Except for *P. savastanoi* TonB-ExbBD, all datasets included movies from grids prepared with and without the presence of fluo OM to improve distribution of particle orientations.

Cryo-EM data processing

Motion correction and dose weighting were performed using MotionCorr implemented in Relion 3.0⁴². Contrast transfer functions were calculated using CTFFIND4⁴³. Particles were picked in Simple⁴⁴ and processed in Relion 3.0⁴². Gold standard Fourier shell correlations using the 0.143 criterion and local resolution estimations were calculated within Relion⁴² (Extended Data Fig. 2).

V. mimicus PomAB particles (1,172,445) underwent one round of reference-free 2D classification, from which 253,681 particles were selected and used to generate an *ab initio* initial model. This model was low-pass filtered to 30 Å and used as reference for 3D classification, generating a class that refined to 6.8 Å from 155,280 particles.

For the deletion construct PomAB $_{\Delta 61-120}$ that improved particle orientations and data quality, particles (2,383,062) were extracted from 13,980 movies. Following one round of

reference-free 2D classification, 800,844 particles were classified in 3D (4 classes) against a 40 Å low-pass filtered map of PomAB. A class containing 244,654 particles was further subjected to masked refinement yielding a 4.8 Å map. Refinement after Bayesian particle polishing and per-particle defocus with beamtilt estimation further improved map quality to 4.2 Å.

B. subtilis MotAB particles (1,532,430) were extracted over 11,588 movies. After 2D classification, selected particles (397,584) underwent two rounds of 3D classification (3 classes each) using a 40 Å low-pass filtered map generated from a subset of particles refined against a 60 Å low-pass filtered map of PomAB_{Δ61-120}. A class made up of 122,615 particles was refined to 3.9 Å. Bayesian particle polishing further improved map resolution by 0.2 Å, and subsequent CTF refinement using per-particle defocus with beamtilt estimation generated a 3.5 Å map. To improve MotB N-terminal and plug densities, a subset of fluorinated particles (43,375) was selected and refined against the 3.5 Å reconstruction, generating a 5.0 Å map that was used to depict these regions in Extended Data Fig. 5.

C. sporogenes MotAB particles (1,998,900) were extracted from 9,148 movies and subjected to a round of reference-free 2D classification. Initial 3D classification performed against a 60 Å low-pass filtered map of *B. subtilis* MotAB revealed two prominent classes which represented a monomeric MotAB complex and a non-physiological end-to-end dimer of MotAB. These classes were used as references in a supervised multi-reference 3D classification against the full 1,137,357 particle set to exclude dimeric particles. Unsupervised 3D classification (4 classes) performed against 865,446 monomeric particles and further refinement yielded 3.8 Å from 314,230 particles. Bayesian particle polishing followed by per-particle defocus with beamtilt estimation further improved map quality to 3.4 Å.

Movies (6,902) were collected for *E. coli* ExbBD, resulting in the extraction of 2,045,350 particles. Following one round of reference-free 2D classification, an initial model of ExbBD was generated by 3D classification and refinement of a particle subset against a 40 Å low-pass filtered 5:1 ExbBD complex³⁴ (EMD-6928). The resulting map was used as initial model for multiple rounds of 3D classification against the full 2D-classified particle set (755,677). After refinement of 227,700 particles, this protocol generated a 5.8 Å map, improving to 4.6 Å following Bayesian particle polishing and per-particle defocus plus beamtilt estimation.

Movies (4,232) were collected for *P. savastanoi* TonB-ExbBD and 1,342,900 particles were extracted. Particles were subjected to two rounds of 2D classification with centered re-extraction between classifications. The cleaned 499,697 particles were subjected to C5-symmetric 3D classification against a 40 Å low-pass filtered 5:2 ExbBD map from *E. coli*. The resultant 202,356 particles were refined with C5 symmetry to generate a 3.5 Å map that lacked density for the transmembrane helices (TMHs) of ExbD. Particles were polished and subjected to an additional round of 2D classification followed by 3D classification with C1 symmetry, resulting in 3.9 Å map from 110,164 particles after refinement in C1. An additional round of Bayesian polishing and refinement (C1) yielded a 3.8 Å map. Per-particle defocus and beamtilt estimation followed by alignment-free 3D classification and subsequent local refinement (C1) yielded a 3.8 Å map with improved density for the TMHs of ExbD from 65,617 particles.

Model building and refinement

Atomic models were built using Coot⁴⁵. Multiple rounds of rebuilding (in both the globally sharpened and local-resolution filtered maps) and real-space refinement in Phenix⁴⁶

using secondary structure, rotamer and Ramachandran restraints yielded the final models described in Table 1. All models were validated using Molprobity⁴⁷. Conservation analysis was carried out using the Consurf server⁴⁸. A homology model of *E. coli* MotAB was generated by sequence threading against the *Clostridium* model using Phyre2⁴⁹. Figures were prepared using UCSF ChimeraX⁵⁰ and Pymol (The PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC). All models depicted in figures are based on the highest resolution *Clostridium* model, unless otherwise specified. Residue numbering adopts the reference *E. coli* sequence and model; a residue conversion table is provided (Supplementary Table 3).

Evolutionary covariance analysis

Coevolutionary contacts for *E. coli* W MotA were determined by the Gremlin web-server⁵. Searches used the Jackhmmer algorithm for multiple sequence alignment, an E-value threshold of 10^{-10} and a minimum coverage of 75%. Intra- and intermolecular contacts were mapped to the *E. coli* MotA structure using Gremlin beta⁵. Intermolecular contacts between MotA and MotB (residues 1-120) were determined using an E-value threshold of 10^{-20} and 10^{-2} , respectively. Intermolecular contacts between TonB and ExbB were determined using an E-value threshold of 10^{-20} . Contacts with a probability score greater than 0.9 were regarded as significant and listed in Supplementary Table 4.

Simulation setup

All simulations were run using GROMACS 2018⁵¹. The systems were initially setup using the Martini 2.2 coarse-grain (CG) force field and solvated with water and 0.15 M NaCl to neutralise the system⁵². The membranes were constructed using INSANE with a 4:1 ratio of POPE:POPG lipids⁵³. An elastic network of $1000 \text{ kJ mol}^{-1} \text{ nm}^{-2}$ was applied between all

backbone beads between 0.5 and 1 nm. Electrostatics were described using the reaction field method, with a cut-off of 1.1 nm using the potential shift modifier and the van der Waals interactions were shifted between 0.9-1.1 nm. The systems were first energy minimised by steepest descent algorithm to 1000 kJ mol⁻¹ nm⁻¹ and then simulated for a total of 1 µs. The temperature and pressure were kept constant throughout the simulation at 310 K and 1 bar respectively, with protein, lipids and water/ions coupled individually to a temperature bath by the V-rescale method⁵⁴ and a semi-isotropic Parrinello-Rahman barostat⁵⁵. The final snapshots from the CG simulations were then converted back to an atomistic description using CG2AT⁵⁶.

Atomistic simulations

The charged N- and C- termini of the converted protein were capped using acetyl and methyl moieties, respectively. All ionisable groups were simulated with default protonation states, unless otherwise mentioned. The virtual site model for hydrogen atoms⁵⁷, adapted for the charmm36 forcefield⁵⁸ was employed, allowing the use of a 4 fs timestep during the simulations. Electrostatics were described using PME, with a cut-off of 1.2 nm and the van der Waals interactions were shifted between 1-1.2 nm. The tip3p water model was used, the water bond angles and distances were constrained by SETTLE⁵⁹. All other bonds were constrained using the LINCS algorithm⁶⁰. The systems were then equilibrated for a further 1 ns using a 4 fs timestep with positional restraints of 1000 kJ mol⁻¹ nm⁻² on the heavy atoms, in a NPT ensemble with temperature V-rescale coupling at 310 K⁵⁴ and semi-isotropic Parrinello-Rahman barostat at 1 bar with protein, lipids and water/ions coupled individually⁵⁵. The Production simulations were performed without position restraints for a total of 200 ns and were run in triplicate.

508

509 **Motility assay**

510 *E. coli* RP6894 (Δ *motAB*) was transformed with pT12-derived plasmids encoding C-
511 terminal twin-strep tagged MotAB containing point mutations or appropriate controls.
512 Saturated overnight cultures (2 μ L) were injected into soft agar plates (0.3% w/v agar in
513 tryptone broth) containing kanamycin (30 μ g/mL) plus rhamnose monohydrate (0.5% w/v)
514 and incubated in a humidified chamber for 23 h at 25 °C.

515

516 **Pulldowns**

517 *E. coli* MT56 was transformed with pT12-derived plasmids encoding C-terminal twin-
518 strep tagged MotAB harbouring point mutations or appropriate controls. Small scale (5 mL)
519 cultures were grown at 37 °C for 16 h in TB media containing kanamycin (50 μ g/mL) and
520 rhamnose monohydrate (0.1% w/v). OD₆₀₀-normalized cell counts were centrifuged at 4,000g
521 for 5 min and pellets frozen at -80 °C. After thawing, cells were lysed by resuspension in 200
522 mM Tris pH 8.0, 300 mM NaCl, 2 mM EDTA plus 30 μ g/mL DNase I and 400 μ g/mL lysozyme
523 for 30 min, then solubilized in 1.5 % w/v LMNG for 1 h. Insoluble material was removed by
524 centrifugation at 18,000g for 30 min. LMNG-solubilized lysates were added to 5 μ g of TBS-
525 prewashed MagStrep XT magnetic beads (IBA Lifesciences) for 1 h with mild shaking. Beads
526 were isolated and washed twice with TBS plus 0.025% w/v LMNG followed by elution with
527 TBS plus 0.025% w/v LMNG and 50 mM D-biotin. Eluates were diluted in SDS-PAGE sample
528 buffer and run on a 4–20% polyacrylamide gel (NuSep). The gel was stained with InstantBlue
529 (Expedeon) to determine the presence of MotA and MotB.

530

531 **Reporting summary**

532

533 Data availability

534 The data that support the findings of this study are available from the corresponding author
535 upon reasonable request. Cryo-EM volumes and atomic models have been deposited to the
536 EMDB (accession codes EMD-10895, EMD-10899, EMD-10901, EMD-10902, EMD-10897) and
537 PDB (accession codes 6YSF and 6YSL), respectively. Source data for gel shown in Extended
538 Data Fig. 10 are provided with the paper.

539

540

541 References

- 542 1 Berg, H. C. The rotary motor of bacterial flagella. *Annu Rev Biochem* **72**, 19-54,
543 doi:10.1146/annurev.biochem.72.121801.161737 (2003).
- 544 2 Leeuwenhoek, A. Observation, communicated to the publisher by Mr Anthony van
545 Leewenhoek, in a Dutch letter of the 9 Octob. 1676 here English'd: concerning little
546 animals by him observed in rain-well-sea and snow water; as also in water wherein
547 pepper had lain infused. *Phil. Trans.* **12**, 821-831, doi:doi:10.1098/rstl.1677.0003
548 (1677).
- 549 3 Nakamura, S. & Minamino, T. Flagella-Driven Motility of Bacteria. *Biomolecules* **9**,
550 doi:10.3390/biom9070279 (2019).
- 551 4 Johnson, S. *et al.* Symmetry mismatch in the MS-ring of the bacterial flagellar rotor
552 explains the structural coordination of secretion and rotation. *Nat Microbiol*,
553 doi:10.1038/s41564-020-0703-3 (2020).
- 554 5 Ovchinnikov, S. *et al.* Large-scale determination of previously unsolved protein
555 structures using evolutionary information. *Elife* **4**, e09248, doi:10.7554/eLife.09248
556 (2015).
- 557 6 Kojima, S. Dynamism and regulation of the stator, the energy conversion complex of
558 the bacterial flagellar motor. *Curr Opin Microbiol* **28**, 66-71,
559 doi:10.1016/j.mib.2015.07.015 (2015).
- 560 7 Blair, D. F. & Berg, H. C. Restoration of torque in defective flagellar motors. *Science*
561 **242**, 1678-1681, doi:10.1126/science.2849208 (1988).
- 562 8 Blair, D. F. & Berg, H. C. The MotA protein of *E. coli* is a proton-conducting component
563 of the flagellar motor. *Cell* **60**, 439-449, doi:10.1016/0092-8674(90)90595-6 (1990).
- 564 9 Larsen, S. H., Adler, J., Gargus, J. J. & Hogg, R. W. Chemomechanical coupling without
565 ATP: the source of energy for motility and chemotaxis in bacteria. *Proc Natl Acad Sci*
566 *U S A* **71**, 1239-1243, doi:10.1073/pnas.71.4.1239 (1974).

- 567 10 Asai, Y., Yakushi, T., Kawagishi, I. & Homma, M. Ion-coupling determinants of Na⁺-
568 driven and H⁺-driven flagellar motors. *J Mol Biol* **327**, 453-463, doi:10.1016/s0022-
569 2836(03)00096-2 (2003).
- 570 11 Kojima, S. & Blair, D. F. Solubilization and purification of the MotA/MotB complex of
571 Escherichia coli. *Biochemistry* **43**, 26-34, doi:10.1021/bi035405l (2004).
- 572 12 Nirody, J. A., Sun, Y.-R. & Lo, C.-J. The biophysicist's guide to the bacterial flagellar
573 motor. *Advances in Physics: X* **2**, 324-343, doi:10.1080/23746149.2017.1289120
574 (2017).
- 575 13 Kojima, S. *et al.* The Helix Rearrangement in the Periplasmic Domain of the Flagellar
576 Stator B Subunit Activates Peptidoglycan Binding and Ion Influx. *Structure* **26**, 590-598
577 e595, doi:10.1016/j.str.2018.02.016 (2018).
- 578 14 Zhu, S. *et al.* Conformational change in the periplasmic region of the flagellar stator
579 coupled with the assembly around the rotor. *Proc Natl Acad Sci U S A* **111**, 13523-
580 13528, doi:10.1073/pnas.1324201111 (2014).
- 581 15 Kim, E. A., Price-Carter, M., Carlquist, W. C. & Blair, D. F. Membrane segment
582 organization in the stator complex of the flagellar motor: implications for proton flow
583 and proton-induced conformational change. *Biochemistry* **47**, 11332-11339,
584 doi:10.1021/bi801347a (2008).
- 585 16 Kojima, S. & Blair, D. F. Conformational change in the stator of the bacterial flagellar
586 motor. *Biochemistry* **40**, 13041-13050, doi:10.1021/bi011263o (2001).
- 587 17 Mandadapu, K. K., Nirody, J. A., Berry, R. M. & Oster, G. Mechanics of torque
588 generation in the bacterial flagellar motor. *Proc Natl Acad Sci U S A* **112**, E4381-4389,
589 doi:10.1073/pnas.1501734112 (2015).
- 590 18 Boschert, R., Adler, F. R. & Blair, D. F. Loose coupling in the bacterial flagellar motor.
591 *Proc Natl Acad Sci U S A* **112**, 4755-4760, doi:10.1073/pnas.1419955112 (2015).
- 592 19 Nishihara, Y. & Kitao, A. Gate-controlled proton diffusion and protonation-induced
593 ratchet motion in the stator of the bacterial flagellar motor. *Proc Natl Acad Sci U S A*
594 **112**, 7737-7742, doi:10.1073/pnas.1502991112 (2015).
- 595 20 Braun, T. F., Al-Mawsawi, L. Q., Kojima, S. & Blair, D. F. Arrangement of core
596 membrane segments in the MotA/MotB proton-channel complex of Escherichia coli.
597 *Biochemistry* **43**, 35-45, doi:10.1021/bi035406d (2004).
- 598 21 Braun, T. F. & Blair, D. F. Targeted disulfide cross-linking of the MotB protein of
599 Escherichia coli: evidence for two H⁺ channels in the stator Complex. *Biochemistry*
600 **40**, 13051-13059, doi:10.1021/bi011264g (2001).
- 601 22 Sharp, L. L., Zhou, J. & Blair, D. F. Features of MotA proton channel structure revealed
602 by tryptophan-scanning mutagenesis. *Proc Natl Acad Sci U S A* **92**, 7946-7950,
603 doi:10.1073/pnas.92.17.7946 (1995).
- 604 23 Sharp, L. L., Zhou, J. & Blair, D. F. Tryptophan-scanning mutagenesis of MotB, an
605 integral membrane protein essential for flagellar rotation in Escherichia coli.
606 *Biochemistry* **34**, 9166-9171, doi:10.1021/bi00028a028 (1995).
- 607 24 Yakushi, T., Yang, J., Fukuoka, H., Homma, M. & Blair, D. F. Roles of charged residues
608 of rotor and stator in flagellar rotation: comparative study using H⁺-driven and Na⁺-
609 driven motors in Escherichia coli. *J Bacteriol* **188**, 1466-1472,
610 doi:10.1128/JB.188.4.1466-1472.2006 (2006).
- 611 25 Hosking, E. R. & Manson, M. D. Clusters of charged residues at the C terminus of MotA
612 and N terminus of MotB are important for function of the Escherichia coli flagellar
613 motor. *J Bacteriol* **190**, 5517-5521, doi:10.1128/JB.00407-08 (2008).

614 26 Hosking, E. R., Vogt, C., Bakker, E. P. & Manson, M. D. The Escherichia coli MotAB
615 proton channel unplugged. *J Mol Biol* **364**, 921-937, doi:10.1016/j.jmb.2006.09.035
616 (2006).

617 27 Braun, T. F. *et al.* Function of proline residues of MotA in torque generation by the
618 flagellar motor of Escherichia coli. *J Bacteriol* **181**, 3542-3551 (1999).

619 28 Minamino, T., Kinoshita, M. & Namba, K. Directional Switching Mechanism of the
620 Bacterial Flagellar Motor. *Comput Struct Biotechnol J* **17**, 1075-1081,
621 doi:10.1016/j.csbj.2019.07.020 (2019).

622 29 Lam, K. H. *et al.* Multiple conformations of the FliG C-terminal domain provide insight
623 into flagellar motor switching. *Structure* **20**, 315-325, doi:10.1016/j.str.2011.11.020
624 (2012).

625 30 Khan, S., Dapice, M. & Humayun, I. Energy transduction in the bacterial flagellar
626 motor. Effects of load and pH. *Biophys J* **57**, 779-796, doi:10.1016/S0006-
627 3495(90)82598-4 (1990).

628 31 Manson, M. D., Tedesco, P. M. & Berg, H. C. Energetics of flagellar rotation in bacteria.
629 *J Mol Biol* **138**, 541-561, doi:10.1016/s0022-2836(80)80017-9 (1980).

630 32 Marmon, L. Elucidating the origin of the ExbBD components of the TonB system
631 through Bayesian inference and maximum-likelihood phylogenies. *Mol Phylogenet*
632 *Evol* **69**, 674-686, doi:10.1016/j.ympev.2013.07.010 (2013).

633 33 Celia, H. *et al.* Structural insight into the role of the Ton complex in energy
634 transduction. *Nature* **538**, 60-65, doi:10.1038/nature19757 (2016).

635 34 Maki-Yonekura, S. *et al.* Hexameric and pentameric complexes of the ExbBD energizer
636 in the Ton system. *Elife* **7**, doi:10.7554/eLife.35419 (2018).

637 35 Celia, H. *et al.* Cryo-EM structure of the bacterial Ton motor subcomplex ExbB-ExbD
638 provides information on structure and stoichiometry. *Commun Biol* **2**, 358,
639 doi:10.1038/s42003-019-0604-2 (2019).

640 36 Swayne, C. & Postle, K. Taking the Escherichia coli TonB transmembrane domain
641 "offline"? Nonprotonatable Asn substitutes fully for TonB His20. *J Bacteriol* **193**, 3693-
642 3701, doi:10.1128/JB.05219-11 (2011).

643 37 Ollis, A. A., Kumar, A. & Postle, K. The ExbD periplasmic domain contains distinct
644 functional regions for two stages in TonB energization. *J Bacteriol* **194**, 3069-3077,
645 doi:10.1128/JB.00015-12 (2012).

646 38 Pawelek, P. D. *et al.* Structure of TonB in complex with FhuA, E. coli outer membrane
647 receptor. *Science* **312**, 1399-1402, doi:10.1126/science.1128057 (2006).

648 39 Cascales, E., Lloubes, R. & Sturgis, J. N. The TolQ-TolR proteins energize TolA and share
649 homologies with the flagellar motor proteins MotA-MotB. *Mol Microbiol* **42**, 795-807,
650 doi:10.1046/j.1365-2958.2001.02673.x (2001).

651 40 McBride, M. J. & Zhu, Y. Gliding motility and Por secretion system genes are
652 widespread among members of the phylum bacteroidetes. *J Bacteriol* **195**, 270-278,
653 doi:10.1128/JB.01962-12 (2013).

654 41 Kuhlen, L. *et al.* Structure of the core of the type III secretion system export apparatus.
655 *Nat Struct Mol Biol* **25**, 583-590, doi:10.1038/s41594-018-0086-9 (2018).

656 42 Zivanov, J., Nakane, T. & Scheres, S. H. W. A Bayesian approach to beam-induced
657 motion correction in cryo-EM single-particle analysis. *IUCrJ* **6**, 5-17,
658 doi:10.1107/S205225251801463X (2019).

659 43 Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from
660 electron micrographs. *J Struct Biol* **192**, 216-221, doi:10.1016/j.jsb.2015.08.008
661 (2015).

662 44 Reboul, C. F., Eager, M., Elmlund, D. & Elmlund, H. Single-particle cryo-EM-Improved
663 ab initio 3D reconstruction with SIMPLE/PRIME. *Protein Sci* **27**, 51-61,
664 doi:10.1002/pro.3266 (2018).

665 45 Brown, A. *et al.* Tools for macromolecular model building and refinement into electron
666 cryo-microscopy reconstructions. *Acta Crystallogr D Biol Crystallogr* **71**, 136-153,
667 doi:10.1107/S1399004714021683 (2015).

668 46 Afonine, P. V. *et al.* Real-space refinement in PHENIX for cryo-EM and crystallography.
669 *Acta Crystallogr D Struct Biol* **74**, 531-544, doi:10.1107/S2059798318006551 (2018).

670 47 Williams, C. J. *et al.* MolProbity: More and better reference data for improved all-atom
671 structure validation. *Protein Sci* **27**, 293-315, doi:10.1002/pro.3330 (2018).

672 48 Ashkenazy, H. *et al.* ConSurf 2016: an improved methodology to estimate and visualize
673 evolutionary conservation in macromolecules. *Nucleic Acids Res* **44**, W344-350,
674 doi:10.1093/nar/gkw408 (2016).

675 49 Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. The Phyre2 web
676 portal for protein modeling, prediction and analysis. *Nat Protoc* **10**, 845-858,
677 doi:10.1038/nprot.2015.053 (2015).

678 50 Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and
679 analysis. *Protein Sci* **27**, 14-25, doi:10.1002/pro.3235 (2018).

680 51 Abraham, M. J. *et al.* GROMACS: High performance molecular simulations through
681 multi-level parallelism from laptops to supercomputers. *SoftwareX* **1-2**, 19-25,
682 doi:10.1016/j.softx.2015.06.001 (2015).

683 52 de Jong, D. H. *et al.* Improved Parameters for the Martini Coarse-Grained Protein
684 Force Field. *Journal of Chemical Theory and Computation* **9**, 687-697,
685 doi:10.1021/ct300646g (2012).

686 53 Wassenaar, T. A., Ingólfsson, H. I., Böckmann, R. A., Tieleman, D. P. & Marrink, S. J.
687 Computational Lipidomics with insane: A Versatile Tool for Generating Custom
688 Membranes for Molecular Simulations. *Journal of Chemical Theory and Computation*
689 **11**, 2144-2155, doi:10.1021/acs.jctc.5b00209 (2015).

690 54 Bussi, G., Donadio, D. & Parrinello, M. Canonical sampling through velocity rescaling.
691 *The Journal of Chemical Physics* **126**, doi:10.1063/1.2408420 (2007).

692 55 Parrinello, M. & Rahman, A. Polymorphic transitions in single crystals: A new
693 molecular dynamics method. *Journal of Applied Physics* **52**, 7182-7190,
694 doi:10.1063/1.328693 (1981).


695 56 Stansfeld, P. J. & Sansom, M. S. P. From Coarse Grained to Atomistic: A Serial
696 Multiscale Approach to Membrane Protein Simulations. *Journal of Chemical Theory
697 and Computation* **7**, 1157-1166, doi:10.1021/ct100569y (2011).

698 57 Feenstra, K. A., Hess, B. & Berendsen, H. J. C. Improving efficiency of large time-scale
699 molecular dynamics simulations of hydrogen-rich systems. *Journal of Computational
700 Chemistry* **20**, 786-798, doi:10.1002/(sici)1096-987x(199906)20:8<786::Aid-
701 jcc5>3.0.Co;2-b (1999).

702 58 Olesen, K., Awasthi, N., Bruhn, D. S., Pezeshkian, W. & Khandelia, H. Faster Simulations
703 with a 5 fs Time Step for Lipids in the CHARMM Force Field. *Journal of Chemical Theory
704 and Computation* **14**, 3342-3350, doi:10.1021/acs.jctc.8b00267 (2018).

705 59 Miyamoto, S. & Kollman, P. A. Settle: An analytical version of the SHAKE and RATTLE
706 algorithm for rigid water models. *Journal of Computational Chemistry* **13**, 952-962,
707 doi:10.1002/jcc.540130805 (1992).
708 60 Hess, B., Bekker, H., Berendsen, H. J. C. & Fraaije, J. G. E. M. LINCS: A linear constraint
709 solver for molecular simulations. *Journal of Computational Chemistry* **18**, 1463-1472,
710 doi:10.1002/(sici)1096-987x(199709)18:12<1463::Aid-jcc4>3.0.Co;2-h (1997).
711
712
713
714
715
716
717
718
719
720
721
722
723
724

725 Extended data figures and tables




726


727 **Extended Data Fig. 1. Ion specificity of stator complexes. a-c**, Sequence alignment of the (a)
728 transmembrane and plug helices of MotB and (b) transmembrane helices (TMHs) 3 and 4 of
729 MotA from *E. coli* and other relevant bacterial species. Species are classified based on ion
730 specificity (red, H⁺-driven; blue, Na⁺-driven). Residues that are conserved just within either
731 the H⁺ or Na⁺-driven classes of stator and map to the MotA-MotB interface (c) are indicated
732 with orange asterisks in the alignment and same residue colouring across H⁺-driven (Left) and
733 Na⁺-driven (Right) models (using *E. coli* MotAB numbering equivalents). The threonine ring is
734 highlighted with light blue asterisks (a,b) or residues (c).

735

736




Extended Data Fig. 2. Cryo-EM map quality and resolution estimates of stators. **a**, Gold-standard Fourier shell correlation (FSC) curves of RELION-postprocessed stator volumes. Resolution at the gold-standard cutoff (FSC = 0.143) is indicated. Curves: red, phase-randomized; green, unmasked; blue, masked; black, MTF-corrected. **b**, Local resolution estimates (in Å) of the sharpened volumes. **c**, Representative modelled densities.




Extended Data Fig. 3. Conservation, covariance, and prior mutagenesis data mapped onto the stator structure. **a**, Surface conservation as determined by Consurf⁴⁸ (maroon high conservation, cyan low conservation). (Left) View from the cytoplasm showing conservation

at the cytoplasmic MotA domains, including residues previously identified to be important for torque generation²⁴ (R90 and E98). (Centre) Side view showing poor conservation within membrane-interfacing residues of MotA. (Right) Cutaway displaying high level of conservation at the MotA-MotB interface; MotA shown as surface representation, MotB as ribbon. **b-d**, Evolutionary co-variation of residues (b) within MotA, (c) between MotA subunits, with boxes highlighting regions of strong covariance that are illustrated in more detail in the adjacent fragments, or (d) between MotA and MotB with the left hand side showing contacts for modelled MotB regions and the right hand side showing contacts in the unmodelled N-terminal MotB density represented here as a poly-alanine backbone. Predictions were carried out in Gremlin⁵ and contacts with a probability score of > 0.9 are shown. Contacts are coloured by Cα-Cα distance (≤10 Å in green, ≤15 Å in yellow). **e**, Mapping previous tryptophan scanning mutagenesis performed on MotA TMHs²² (i) or MotB²³ (ii) to the stator structure. MotA and MotB are coloured as in Fig. 2,3 with targeted residues coloured according to toleration to mutagenesis; yellow corresponds to tolerated mutants (relative swarm rates > 0.5), red are poorly tolerated mutants (relative swarm rates of ≤ 0.5). In (i), TMH4 (green) and TMH2 (red) of neighbouring MotA subunits and MotB (white) are shown as transparent silhouettes. Poorly tolerated mutants cluster at subunit interfaces. In (ii) only TMH3-TMH4 of three MotA subunits are shown for clarity. **f**, Mapping previously determined cysteine crosslinks between (i) MotA-MotA²⁰, (ii) MotB-MotB²¹, or (iii) MotA-MotB²⁰ to our structure. For displaying crosslinks, a yield of ≥ 30% disulfide-linked adduct under iodine oxidizing conditions was used as threshold, except for MotA_{TMH4}-MotB crosslinks which used a ≥ 10% threshold. All analyses in this figure were performed using an *E. coli* MotAB structure generated by homology threading onto *C. sporogenes* MotAB.




Extended Data Fig. 4. Residues that interact with the flagellar C-ring form a charged ring on the cytoplasmic face of MotA. **a**, An isolated MotA subunit with the essential torque-generating charged residues R90 and E98²⁴ displayed in yellow spheres representation. **b**, The full *C. sporogenes* stator complex viewed from the side (Left) or from the cytoplasm (Right), coloured as in Figs. 2-3, and with the torque-generating charged residues represented as in (a) using *E. coli* MotAB numbering scheme.


10




Extended Data Fig. 5. Structural elements of the *B. subtilis* stator. **a**, Structure of the *B. subtilis* stator depicting unmodeled density for the MotB N-terminal extensions (Left) and plug helices (Right). **b**, (i) Isolated *B. subtilis* MotB dimer represented as in (a), and (ii) superposition of the TMHs of the two MotB chains showing the relative rotation of the plug helices. **c**, Slabs at the heights indicated through *B. subtilis* MotAB show distortion from a regular pentagon (arrow indicates the cytoplasmic side of the complex) as viewed from the cytoplasm). **d**, Surface representation of *B. subtilis* MotAB showing tight packing. (Top) Side view with the front of the complex removed. (Bottom) Top-down view of the slab indicated by dashed lines. **e**, Structural alignment of the five *B. subtilis* MotA chains reveal they fall into two conformational classes which differ in the degree of flexing at the highlighted prolines.



Extended Data Fig. 6. Molecular dynamics simulations of stator structures in lipid bilayers.
a, Side (*top*) and top-down (*bottom*) views of (i) *C. sporogenes* and (ii) *B. subtilis* MotAB inserted within a lipid bilayer after extended simulations (coarsegrain for 1 μ s then atomistic for a further 200 ns). **b**, Cartoon representation of (i) *Clostridium* MotAB and (ii) *Bacillus* MotAB coloured (blue to red) by the average rmsf of the 3 replica simulations performed.



Extended Data Fig. 7. Cryo-EM map quality and resolution estimates of ExbBD complexes a, Gold-standard Fourier shell correlation (FSC) curves of RELION-postprocessed ExbBD maps. Resolution at the gold-standard cutoff (FSC = 0.143) is indicated. Curves: red, phase-randomized; green, unmasked; blue, masked; black, MTF-corrected. **b,** Local resolution estimates (in Å) of the sharpened maps. **c,** Structure of the 5:2 ExbBD complex from *E. coli*³⁵ (PDB 6tyi) fit into the *E. coli W* (top) and *P. savastanoi* ExbBD (bottom) maps. ExbB is coloured dark grey and ExbD is coloured blue. Top right panels have three ExbB subunits removed to demonstrate density for the two TMHs of ExbD. Bottom right panels reveal the 5:2 ExbB:ExbD arrangement as viewed from the cytoplasm.



818

819 **Extended Data Fig. 8. Structural alignment of MotAB and ExbBD.**

820 **a**, side views of the cores of the *C. sporogenes* MotAB (orange) and *E. coli* ExbBD³⁵ (yellow)

821 complexes with two front MotA/ExbB subunits removed showing the conserved Asp residues

822 lying at the same height with respect to a ring of Thr on the MotA/ExbD components **b**,

823 Overlaying the cores of *C. sporogenes* MotAB (orange) and *E. coli* ExbBD (yellow) by alignment

824 of the MotA/ExbB helices demonstrates alignment of the critical polar residues between the

825 two systems. Regions outside the core shown as transparent grey; MotB/ExbD shown in grey.


826 **c**, Aligning a single subunit of MotA (rainbow colouring) with ExbB (white) via the pore lining

827 helices reveals the different elaborations of this core unit in each protein. **d**, Aligning a single

828 subunit of MotA with ExbB via the cytoplasmic extensions of the two core helices reveals the

829 different folding of the rest of the cytoplasmic regions. Proteins coloured as in (c).

830



Extended Data Fig. 9. TonB recruitment to the ExbBD complex. **a**, SEC-MALS profile of purified *Pseudomonas* TonB-ExbBD (black absorption curve) and ExbBD (grey absorption curve) with SDS-PAGE analysis of each sample inlayed. Total protein-detergent complex molar mass (green) and deconvoluted protein (blue) and detergent (orange) molar masses are shown. A ~30 kDa difference in molar mass is observed between the complexes consistent with the the TonB-ExbBD complex containing one TonB subunit. **b**, Evolutionary co-variation of residues between TonB and ExbB displayed on the *E. coli* ExbBD structure³⁵ (PDB 6tyi). For clarity, a topological model of the TMH of TonB is shown in orange with TonB-ExbB contacts indicated as dashed lines. Covarying residues decorating the periphery of ExbB are displayed in green. ExbB (grey) and ExbD (blue) are displayed as ribbon cartoons. Contacts shown were generated by Gremlin⁵ and have a probability score of > 0.9.

846 **Supplementary Data**

847 **Supplementary Table 1.** Bacterial strains and plasmids used in this study

848 **Supplementary Table 2.** Oligonucleotides used in this study.

849 **Supplementary table 3.** Species conversion of residues specified in text

850 **Supplementary Table 4.** List of evolutionary contacts predicted by Gremlin

851