

Automated phenotyping of mosquito larvae enables high-throughput screening for novel larvicides and smartphone-based detection of larval insecticide resistance

Steven D. Buckingham^{1†}, Frederick A. Partridge^{1†}, Beth C. Poulton^{1,2†}, Ben Miller^{3†},
Rachel A. McKendry^{3,4} Gareth J Lycett² and David B. Sattelle^{1*}

¹UCL Centre for Respiratory Biology, UCL Respiratory, Division of Medicine, Rayne Building, 5 University Street, London WC1E 6JF, UK.

²Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, Merseyside, L3 5QA, UK.

³London Centre for Nanotechnology, Faculty of Maths & Physical Sciences, University College London, 17-19, Gordon Street London, WC1H 0AH, UK.

⁴Division of Medicine, UCL, Rayne Building, 5 University Street, London WC1E 6JF, UK

†These authors contributed equally; * To whom correspondence should be directed

Abstract

Pyrethroid-impregnated nets have contributed significantly to halving the burden of malaria but resistance threatens their future efficacy and the pipeline of new insecticides is short. Here we report that an invertebrate automated phenotyping platform (INVAPP), combined with the algorithm Paragon, provides a robust system for measuring larval motility in *Anopheles gambiae* (and *An. coluzzi*) as well as *Aedes aegypti* with the capacity for high-throughput screening for new larvicides. By this means, we reliably quantified both time- and concentration-dependent actions of chemical insecticides faster than using the WHO standard larval assay. We illustrate the effectiveness of the system using an established larvicide (temephos) and demonstrate its capacity for library-scale chemical screening using the Medicines for Malaria (MFP) Pathogen-Box library. As a proof-of-principle, this library screen identified a compound, subsequently confirmed to be tolfenpyrad, as an effective larvicide. We have also used the INVAPP / Paragon system to enable detection of resistance to deltamethrin. We show how this approach to monitoring larval susceptibility to insecticides can be adapted for use with a smartphone camera application and therefore has potential for further development as a simple portable field-assay for insecticide resistance with associated real-time, geo-located information to identify hotspots.

Author summary

We have developed an automated platform for recording the motility of mosquito larvae and applied it to larvae of a mosquito vector of malaria and a mosquito vector of dengue, Zika, yellow fever and other human diseases. The platform facilitates high-throughput, chemical screening for new compounds to control mosquito larvae and also allows detection of larval resistance to the pyrethroid insecticide deltamethrin. Pyrethroid-impregnated bednets have helped to halve the deaths from malaria in recent years but pyrethroid resistance is an important threat to this progress. Our approach assays insecticide actions faster than the current WHO standard test and we show that it can be adapted for use with a smartphone, which offers the prospect of a future field assay for resistance with the added benefit of precise satellite-based location.

Introduction

Mosquito-borne diseases kill more than one million people every year, while severely debilitating diseases including malaria, dengue, Zika, West Nile virus, chikungunya, yellow fever and Japanese encephalitis, transmitted by these vectors cause suffering for hundreds of millions. Successful attempts to control the incidence of such diseases have mainly targeted mosquito population size and hence infective biting frequency [1]. As a result, the burden of malaria has been halved in the period 2000-2015, mostly due to the use of insecticide-treated nets (ITNs) containing pyrethroids, which target the adult disease-transmitting *Anopheles* mosquitoes that tend to feed nocturnally in homes [2]. However, the future efficacy of ITNs to control malaria is threatened by growing mosquito resistance to the available pool of insecticides [3] and by an increase in the prevalence of *Anopheles* mosquitoes that blood-feed outdoors [4].

ITNs are of limited use against viral diseases transmitted by *Aedes aegypti* mosquitoes, such as dengue, Zika, yellow fever and chikungunya, since the adult vectors favour feeding at dawn and dusk. The ineffectiveness of ITNs against Aedes and the absence of drugs or vaccines for most vector-borne viruses means that a mainstay of viral disease control has been larval source management (LSM) [5]. This integrated approach targets aquatic egg-laying sites with the aim of reducing the number of immature forms, leading in turn to fewer biting adults. LSM can incorporate community environmental measures to limit temporary aquatic breeding sites in and around homes, as well as the application of larvicidal agents, such as temephos, to important, permanent water sources. These traditional LSM methods are effective where the breeding sites that produce the majority of adults can be readily identified and treated accordingly [6]. Alternative methods, attempting to overcome this limitation, involve the dusting of resting female mosquitoes with highly potent larval growth inhibitors. As a result, growth inhibitors are delivered to oviposition sites at doses proportional to the frequency of visits. By this means, the most important breeding sites are targeted [7].

For malaria control, there is growing awareness that LSM may have a more prominent role to play under certain situations, complementing adult-targeted insecticide interventions. Examples include: targeting zones where the prominent vectors rest and feed outdoors; tackling transmission foci, where saturating ITN usage is having limited effect on disease transmission; deployment in areas where adults are resistant to all known insecticides; treatment of urban areas in which water sources that suit Anopheline development may be more readily identified [8]. Technologies, such as drones and mapping have been combined to improve identification and prioritisation of habitats in rural areas [9,10]. There may be additional opportunities to control multiple larval mosquito species simultaneously that will affect both malaria and viral disease transmission [6].

Larvicides in current use include surface oils and films that suffocate immature forms, bacteria (including *Bacillus thuringiensis* that produces toxic protein crystals that target the larval midgut), insect growth regulators (including pyriproxyfen that inhibits adult emergence) and synthetic chemicals (such as chlorpyrifos, pirimiphos-methyl and temephos) that target the insect nervous system [6] .

Standardized methods to examine relative larval susceptibility to compounds are necessary to maintain the effective deployment of current larvicides by identifying the emergence of resistance in mosquito populations [11]. Moreover, a high-throughput method to screen larvicides would offer the possibility to examine large chemical libraries for new classes of compounds. Currently the World Health Organisation issues guidelines for larvicidal toxicity assays [12], which consist of exposing larvae in small cups to increasing doses of chemical and manual observation of mosquito killing. In such assays, larval death is defined by the moribund appearance of larvae and their failure to respond to tapping of the cup. This is labour-intensive and the end-point can be difficult to assign unambiguously.

We have developed an automated phenotyping assay based on larval motility, which is simple to deploy, provides a fast readout and, being multi-well plate based, offers much needed high-throughput screening capabilities. Our system uses an INVertebrate Automated Phenotyping Platform (INVAPP) previously described for monitoring nematode motility in conjunction with the Paragon algorithm to estimate motility from moving images of larvae [13]. We illustrate the capability of our system by (a) quantifying the actions of the larvicide temephos on larval motility, (b) screening a

chemical library, which showed larvicidal actions of tolfenpyrad and (c) detecting pyrethroid (deltamethrin) resistance in *Anopheles coluzzii* and *Aedes aegypti* mosquito larvae. We also demonstrate the potential for its further development as a smartphone-based field assay, for rapid detection of pyrethroid resistance, using the inbuilt camera to track larval motility.

Methods

Mosquito rearing and strains

The mosquito strains used are established colonies which have been characterised by WHO protocols as either deltamethrin sensitive or resistant in female adults. They have been in lab culture for 2 - 20 years and were not infected with any known pathogen. Resistant (Cayman) and sensitive (New Orleans) *Ae. aegypti* strains and resistant *An. coluzzii* (Tiassale) and sensitive *An. gambiae* (G3) strains were studied. It should be noted that there exists a considerable degree of genetic variation in Anopheline mosquitoes especially in the species complex *Anopheles gambiae sensu lato*, which is widespread in sub-Saharan Africa. *Anopheles gambiae* and *Anopheles coluzzii* are morphologically indistinguishable species and were for many years considered as distinct molecular forms (S and M respectively) of the same species. The G3 strain we here deploy has always been called *Anopheles gambiae* but is in fact a hybrid of molecular forms. Pure *An. coluzzii* strains can be found in Tiassale, Cote D'Ivoire, but the lab strain Tiasalle 13 deployed here is also a hybrid.

Larval hatching and maintenance

Anopheles

Anopheles eggs laid day 1 evening or day 2 morning were shipped securely from Liverpool School of Tropical Medicine (LSTM) to University College London (UCL) on filter paper ensuring that they remain moist throughout the journey. At UCL on day 3, eggs were washed into a shallow dish containing deoxygenated (pre-boiled and cooled) water containing 0.001% pond guardian tonic salt (Blagdon) to a depth of about 2.5 cm. The eggs and newly-hatched larvae were maintained at 25°C. Larvae were fed 1/3 pellet of cat food per dish and tested on days 5-6.

Aedes aegypti

Ae. aegypti egg papers were shipped securely to UCL from LSTM and stored at room temperature for up to 1 month. They were hatched over 1-2 days by placing the egg papers in a tray of deoxygenated (pre-boiled and cooled) tap water to which a crushed yeast tablet (Holland and Barrett) had been added.

Harvesting larvae and transfer to 96-well plates

Experiments were performed on first instar larvae. The water in which larvae were swimming was passed through a 100 µm Nylon mesh cell-strainer (Fisher Scientific) to concentrate the larvae. This concentrated suspension of larvae was diluted until 100 µL of suspension contained 5-10 larvae. A 100 µL aliquot of this suspension was added using a standard Gilson pipette to each well of a 96-well plate, with the tip of the pipette being cut back to reduce damage to the larvae. Subsequently, 100 µL of the compound under test was added (dissolved in water from a 10⁻²M DMSO stock to yield the required final concentration). Wells containing DMSO alone diluted to appropriate concentrations served as controls.

Filming larvae using INVAPP and analysis of motility using Paragon

Larvae were filmed using INVAPP, a device for monitoring invertebrate motility that is well suited to high-throughput chemical screening and has been deployed for the study of nematode motility (11). INVAPP consists of an Andor Neo “sCMOS” camera (2560 x 2160 resolution, maximum 100 frames per second frame rate) fitted with a Pentax YF3528 line-scan lens. The camera is mounted beneath a microplate supporting platform illuminated from above by an LED array panel. An acrylic diffuser ensures that the field of view of the camera is evenly lit (Figure 1).

Images (30) were acquired every 10 ms. This was repeated at approximately 5 s intervals until 5 or 10 series of image sequences were obtained. Acquiring images at this rate detected the slow, drifting movements associated with filter feeding as well as rapid “jumping” that larvae undergo at sporadic intervals. For every plate, in addition to filming at selected time points, readings were collected immediately prior to addition of chemicals. Image sequences were stored offline for later analysis.

Paragon algorithm

The analysis uses an implementation of the Paragon algorithm [13] and is described in outline below:

1. Read frames 1, 11 and 21 of the image sequence into a $n \times m \times 3$ array where n , m are the height (approximately 700 pixels) and width (approximately 950 pixels) of the image sequences.
2. Calculate the variance in the time dimension
3. Identify the pixels for which the variance exceeds the mean by 3 standard deviations:

$$\begin{cases} 1 & \text{if } x_i > \text{mean}(x) + 3 \times \text{std}(x); \\ 0 & \text{if } x_i \leq \text{mean}(x) + 3 \times \text{std}(x) \end{cases}$$

4. For each well, count the number of pixels of value 1 to produce a *Movement Index*
5. Normalize the movement index for all time points by dividing it by the value obtained for movies acquired immediately before adding compounds.

Image processing and inferencing was performed with Python scripts utilizing the following publicly available software libraries: Numpy, Pandas, Scikit-Image and Statsmodels. Matplotlib and Seaborn software libraries were employed for plotting.

Filming a plate requires 5 (in some cases 10) image sequences to be made each with 30 frames at 10 ms intervals, requiring a total of about 1 s for each image sequence. Analyzing one movie takes approximately 3 s on a desktop computer. The speed of recording and analysis permits library-scale screening of re-profiled and novel, candidate insecticide chemistry.

Chemicals

Temephos, an organophosphate anti-cholinesterase and established mosquito larvicide, was obtained from Sigma Aldrich (Item 31526-250MG). Deltamethrin (DM) is a pyrethroid insecticide used in bednets for which acts on insect sodium channels [14–16] and was obtained from Chem Service (Item N-11579). This particular pyrethroid was selected for study as the strains used had previously been characterised for deltamethrin resistance in the adult stage, and we were keen to examine if resistance could also be detected during larval stages using the INVAPP / Paragon system. Temephos and deltamethrin were prepared as stock solutions (1×10^{-2} M in DMSO) then diluted to the required final concentrations in water.

The Medicines for Malaria Pathogen box library contains 400 drug-like molecules targeting a wide range of neglected tropical diseases. We used this to explore the potential of library-scale drug screening using INVAPP / Paragon.

Smartphone-based detection of resistance

The smartphone application software was written using Android Studio with the OpenCV 3.3.0 library to implement a camera, handling the RAW pixel values in real time. The smartphone used was an LG G4 (16 MP, f/1.8, 28mm). After adding the larvae to wells of a 96 well plate and leaving to settle, recording is initiated with the phone camera pointing at the well plate. The application then automatically initialises two 1D arrays for the real-time mean and variances of each pixel. On each

camera frame, the raw pixel values are extracted as a 1D byte array and converted to a float array. The mean and variance matrices are updated with the following formulae:

$$M_n = M_{n-1} + \frac{P - M_{n-1}}{n}$$

where, M is the mean matrix, P is the pixel value matrix and n is the number of frames.

$$V_n = \frac{(n-1)V_n + (P - M_{n-1})(P - M_n)}{n}$$

where V is the variance matrix.

After 60 frames, the mean and variance matrices are exported as a text file. This measurement is repeated every 10 s. The frame-rate, and therefore sensitivity, depends on the resolution of the image. Here, 320×240 images are used, giving around 30 frames per s, and therefore 2 s measurements every 10 s.

At $t=200$ s, deltamethrin or control (water) was added to each well, and the variance tracked over time as described. The resulting time-variance plots are fitted to exponential a decay function (Mathematica 11.3):

$$V_t = A + B \exp^{-k(t-T)}$$

where V is the variance, A is the baseline variance, B is the maximum variance due to the larvae, k is the variance decay rate, and T is the time when the variance begins to decrease.

The fitted values for k are used to distinguish between susceptible and resistant larvae.

WHO Larval Assay

The WHO larval assay was published in 2005 with the aim of standardizing testing of compounds for their larvicidal actions on mosquitoes. It provides a detailed protocol for quantifying these toxic

actions of insecticides on mosquito larvae in the field and in the laboratory and is available from

https://apps.who.int/iris/bitstream/handle/10665/69101/WHO_CDS_WHOPES_GCDPP_2005.13.pdf.

A quarter yeast tablet (Holland and Barrett) was added to 200 ml of dH₂O containing 0.001% pond salt (Blagdon), prior to addition of 25 third-instar *Ae. aegypti* (New Orleans or Cayman) larvae per cup. The appropriate volume of deltamethrin stock (dissolved in acetone) was then added to achieve the desired concentrations (5×10^{-10} – 1×10^{-6} M respectively) Mortality was assessed visually after 24 h continuous exposure and performed in triplicate over three biological replicates.

Results

Using the INVAPP system to study the actions of the larvicide temephos on larvae of Aedes aegypti

Mosquito larvae swim [17,18] , jump [19] and rest [18]. Filming at 30 frames per second ensures that all types of larval movements are captured. Temephos, a widely used larvicide targeting *Ae. aegypti*, was tested on susceptible larvae of this species. A concentration-dependent reduction in motility was detected after 240 min exposure to this larvicide (Fig. 1, bottom right panel). Thus, the INVAPP / Paragon system enables robust, automated detection of larval motility and the severe motility impairment resulting from exposure to the larvicide temephos.

Screening a chemical library for compounds with larvicidal activity

Given the system's ability to provide a monitor of impaired motility in the presence of an established larvicide, we tested the system using a small chemical library, the Medicines for Malaria Venture (MMV) Pathogen Box. This library was supplied as 1.0×10^{-2} M solutions in DMSO and was further diluted in DMSO to generate a 1.0×10^{-3} M stock of each compound. The library was then screened against *An. gambiae* using the INVAPP / Paragon system; all chemicals from the collection were initially tested at 1.0×10^{-5} M, $n = 3$, 1% v/v final DMSO. The actions of each chemical on the movement index was recorded and ranked by their median (see supplementary data Table S1), and the top 5 compounds (MMV007803, MMV637229, MMV687775, MMV687776, MMV688934) were used in further analysis. All 5 were re-sourced in powder form from Sigma Aldrich, prepared as a fresh DMSO stock, and tested for concentration-dependent actions over the range 3×10^{-7} - 1×10^{-4} M. Of these compounds, MMV688934 (tolfenpyrad) showed a marked concentration-dependent immobilization of larvae (Fig. 2). Tolfenpyrad is an inhibitor of complex I of the respiratory electron transport chain in the mitochondria. It has been successfully trialled in toxic sugar baits to kill

mosquitoes [20]. A weaker action was observed for MMV637229 (Clemastine, a first-generation antihistamine for the treatment of hayfever) which has also shown to be effective in killing *Trypanosoma cruzi* and *Plasmodium falciparum* [21,22].

Detecting deltamethrin resistance in *An. coluzzii* and *Ae. aegypti*

We tested whether our INVAPP / Paragon system was able to detect and measure resistance to insecticides in mosquito larvae. Larvae from deltamethrin susceptible and resistance strains were examined for each species (G3 [susceptible] and Tiassale [resistant] for *Anopheles*, and New Orleans [susceptible] and Cayman [resistant] for *Ae. aegypti*). Image sequences were filmed immediately before addition of Deltamethrin and at intervals (2, 5, 10, 20, 45 and 60 min) afterwards.

There was a concentration-dependent action of deltamethrin on motility for the 4 different strains (Fig. 3). The concentration dependence was detected after 5 min of exposure. At briefer exposures larvae were not completely immobilized, whereas at exposures longer than 60 min even the lowest concentrations tested had immobilized all larvae. The curves for the 4 strains after 60 min exposure to deltamethrin were fitted to a sigmoid function (see Methods), from which $PI_{C_{50}}$ s were estimated.

The $PI_{C_{50}}$ for Tiassale was estimated to be 7.0 ± 1.0 (100 nM), while that for G3 was estimated to be 8.0 ± 1.0 (10 nM). Similarly, the $PI_{C_{50}}$ for Cayman was estimated to be 8.0 ± 1.0 (10 nM), while that for New Orleans was estimated to be 9.0 ± 0.0 (1 nM). The differences between susceptible and resistant strains were statistically significant (2-way ANOVA, $F(1.0,24.0) = 7.0$, $P = 0.016$) but no significant difference was detected between genus (2-way ANOVA, $F(1.0,24.0) = 3.0$, $P = 0.091$) and there was no statistically significant interaction between genus and susceptibility (2-way ANOVA, $F(1.0,24.0) = 0.0$, $P = 0.906$). Hence, the algorithm detected resistance to deltamethrin in both *Anopheles* and *Aedes* larvae.

Comparison of INVAPP / Paragon data with the standard WHO assay to detect relative resistance

To determine how the Paragon algorithm compared to the WHO larval assay, we tested the *Ae. aegypti* New Orleans and Cayman strains with traditional larval cup bioassays (Fig. 4). The WHO assay yielded LC₅₀s of 8.60 ± 0.04 (2.5 nM) and 7.29 ± 0.14 (51.3 nM) for New Orleans and Cayman respectively (2-tailed, independent t-test, T(4)=-3.9, P=0.0). The inhibition and resistance ratios estimated from the two methods were thus 13.5 and 11.8 respectively. Hence, the resistance ratio estimated from WHO assays is similar to the inhibition ratio obtained from Paragon. This suggests that the inhibition of movement at the 4hr time point in the Paragon assay correlates well with the 24hr killing observed in WHO assays.

Smartphone detection of pyrethroid resistance

Larvae were dispensed into a petri dish or 96 well plate and allowed to settle. Films of 2000-3000 s duration were made at the smartphone camera default settings (LG G4: 16 MP, f/1.8, 28 mm). The INVAPP/Paragon algorithm was then applied with no changes other than the threshold for allocating a pixel as “having movement” or “not having movement”. In the absence of deltamethrin both resistant and susceptible larvae yielded a roughly constant movement index over the duration of the recording (Fig. 5A, B, dotted lines). In the presence of 1×10^{-6} M deltamethrin, the movement index began to fall after about 500-1000 s to a steady-state, low level. The rate of this decline was slower for resistant strains ($\tau = 0.18 \pm 0.08 \times 10^3 \text{ s}^{-1}$ for Cayman and $3.5 \pm 0.16 \times 10^3 \text{ s}^{-1}$ for Tiassale) than it was for susceptible strains ($\tau = 2.7 \pm 0.2 \times 10^3 \text{ s}^{-1}$ for New Orleans and $12 \pm 2.3 \times 10^3 \text{ s}^{-1}$ for G3 – *Anopheles*: one-tailed unpaired t-test T(4) = 7.0, P = 0.003 ; *Ae. aegypti*: one-tailed unpaired t-test T (4) = 20.0, P < 0.001). Thus, INVAPP/Paragon detected differential susceptibility between strains for two significant

human disease vectors following short term exposure to 1 μ M deltamethrin in real-time on a readily available smartphone.

Discussion

Here we evaluate the utility of an automated screening platform / algorithm combination in quantifying the actions of chemicals tested against mosquito larvae for their capacity to impair larval motility. Previously, the “INVertebrate Automated Phenotyping platform (INVAPP), with the Paragon algorithm [13] had been shown to enable high-throughput chemical screening on the nematode genetic model organism *C. elegans* and parasitic nematodes such as *Haemonchus contortus*, *Teladorsagia circumcincta*, and *Trichuris muris* worms in the search for novel anthelmintic candidates [13]. The system allows 96 well plate-based chemical screening for anthelmintic activity. By this means we detected compounds affecting the motility and development of worms. We deployed known anthelmintics to validate the utility of the INVAPP/Paragon system and screened the MMV Pathogen Box chemical library identifying compounds already known to have anthelmintic or anti-parasitic activity and other compounds previously not known to have anthelmintic activity [13]. In separate studies we showed how this approach could help identify novel classes of chemistry with anthelmintic activity, the dihydrobenz[e] [1,4] oxazepin-2 (3*H*)-ones (21) and the 2,4-diaminothienol [3,2-*d*] pyrimidines [23].

Here we demonstrate the use of INVAPP paired with the analysis algorithm Paragon as a rapid, convenient and robust method for estimating a biologically-relevant parameter (motility) of the behaviour of mosquito larvae from two key human disease vectors. It was first used to detect the actions on motility of *Ae. aegypti* larvae of a known larvicide, temephos. Following this validation of the ability of INVAPP / Paragon to detect larval motility impairment by an established larvicide, we

explored its potential use in library-scale chemical screening. The discovery of new and re-profiled insecticidally-active chemicals remains important. As a proof-of- principle exercise, we screened (blind) a 400-compound library – The Medicines for Malaria Venture (MMV) Pathogen Box. We identified tolfeprad, an established insecticide, as a toxic agent for mosquito larvae. Tolfeprad is a pyrazole insecticide developed by the Mitsubishi Chemical Corporation in 1991. It is active against whiteflies [24] and aphids [25] as well as *H. contortus* [26] . Since tolfeprad is the only known insecticide in the MMV Pathogen Box, this is indicative of the specificity of the screen.

As well as readily identifying the potent insecticide, tolfeprad, our screen of the MMV Pathogen Box also showed that clemastine has a paralysing action on mosquito larvae, suggesting that the algorithm can be useful in high throughput chemical re-profiling screens. Clemastine is a sedating anti-histaminergic drug used in the treatment of allergic reactions, including seasonal allergic rhinitis (“hay-fever”) [27]. In vertebrates it blocks primarily the H1-subtype of histamine receptor, which is a member of the 7-transmembrane, G-protein-coupled receptor family [28]. Insects are known to use histamine as a neurotransmitter and to possess a histamine-gated chloride channel with some pharmacological similarities to the vertebrate H1 class [28,29]. Perhaps of equal interest is the fact that histamine-gated chloride channels are not present in vertebrates [30] and insects do not possess a metabotropic histamine receptor [31]. Cys-loop ligand-gated anion channels are known targets of insecticides and anthelmintics [32,33]. It has also been reported that mutations in the two *Drosophila* histamine-gated chloride channels (HCIA and HCIB) have opposite effects on ivermectin sensitivity: mutation of HCIA confers enhanced sensitivity whereas knockout of HCIB diminished sensitivity to avermectin neurotoxins [34,35], suggesting that these channels as well as GluCl_s may play a role in the action of macrocyclic lactone derivatives. Furthermore, the finding that two histamine-gated

chloride channels cloned from the housefly, *Musca domestica*, are insensitive to other insecticides known to target GABA and glutamate-gated ion channels [36] suggests that histamine-gated chloride channels may have a very distinct pharmacological profile offering a quite new candidate insecticidal target. However, it should be noted that neither HClA [37] nor HClB [35] null mutants are lethal, suggesting a possible limitation on the usefulness of these receptors as targets of insecticides.

Resistance to drugs and chemicals used to combat malaria seriously threatens the success of control measures and the investment therein over the past 15 years. The widespread emergence of pyrethroid resistance threatens the hitherto successful deployment of ITNs [3,38]. Resistance in the adult mosquito vectors are established but much less is known concerning resistance in mosquito larvae for which targeting by larvicides is a valid control approach especially for *Ae. aegypti*. Here using behavioural phenotyping, we have shown that larvae of *An. coluzzii* and *Ae. aegypti* also show resistance to a pyrethroid (deltamethrin) in strains where the adults are known to be resistant. These findings are robust in our laboratory studies. In future, it will be of interest to explore whether dual larvicidal treatments can help combat resistance and whether the INVAPP / Paragon system can be used to assist in identifying new areas of chemistry and / or new targets not already compromised by known resistance mechanisms that will be needed for such approaches.

INVAPP combined with the Paragon algorithm allows for faster measurement than can be accomplished using the currently deployed manual WHO assay for larvicidal activity. It may not be as sensitive as the manual assay, but it allows the effectiveness of compounds to be measured more rapidly, and provides more detail in the time-dependence of the actions of compounds to be identified. It also requires a smaller bench footprint than the WHO assay and is less labour intensive.

Also, manual assays such as the current WHO assay are subject to investigator fatigue and the end-point assessment is not always straightforward. INVAPP / Paragon, which is automated, reduces errors due to these factors.

Increasingly, sensitive smartphone applications and cameras are being used for the diagnosis and study of infectious diseases including malaria. For example, a smartphone polarised microscope has been used for malaria diagnosis [39]. Automated detection of malaria parasites in blood smears has been achieved [40,41]. A rapid and robust field detection of resistance that utilises a simple smartphone assay could also yield valuable information. Geo-located data on larval resistance could be readily transmitted to central databases that provide regional information of the temporal and spatial development of insecticide resistance which can be mapped alongside the emergence of adult resistance and malaria prevalence data [42,43]. We have shown that relative larval resistance to a pyrethroid insecticide can be detected using a smartphone camera. This paves the way for the development of a smartphone application perhaps to be used in conjunction with a simple, slide-on test kit that will facilitate such measurements. These assays are a promising first step in developing such a simple-to-operate field assay.

References

1. Curtis CF, Jana-Kara B, Maxwell CA. Insecticide treated nets: impact on vector populations and relevance of initial intensity of transmission and pyrethroid resistance. *J Vector Borne Dis.* 2003;40: 1–8.
2. Bhatt S, Weiss DJ, Mappin B, Dalrymple U, Cameron E, Bisanzio D, et al. Coverage and system efficiencies of insecticide-treated nets in Africa from 2000 to 2017. Kyobutungi C, editor. *eLife.* 2015;4: e09672. doi:10.7554/eLife.09672
3. Hemingway J, Ranson H, Magill A, Kolaczinski J, Fornadel C, Gimnig J, et al. Averting a malaria disaster: will insecticide resistance derail malaria control? *The Lancet.* 2016;387: 1785–1788. doi:10.1016/S0140-6736(15)00417-1

4. Russell TL, Beebe NW, Bugoro H, Apairamo A, Chow WK, Cooper RD, et al. Frequent blood feeding enables insecticide-treated nets to reduce transmission by mosquitoes that bite predominately outdoors. *Malar J.* 2016;15: 156. doi:10.1186/s12936-016-1195-8
5. Medical Entomology for Students. 5 edition. Cambridge: Cambridge University Press; 2012.
6. Larval source management: a supplementary measure for malaria vector control: an operational manual. Geneva: World Health Organization; 2013.
7. Devine GJ, Perea EZ, Killeen GF, Stancil JD, Clark SJ, Morrison AC. Using adult mosquitoes to transfer insecticides to *Aedes aegypti* larval habitats. *Proc Natl Acad Sci.* 2009;106: 11530–11534. doi:10.1073/pnas.0901369106
8. Tusting LS, Thwing J, Sinclair D, Fillinger U, Gimnig J, Bonner KE, et al. Mosquito larval source management for controlling malaria. *Cochrane Database Syst Rev.* 2013 [cited 16 Apr 2020]. doi:10.1002/14651858.CD008923.pub2
9. Carrasco-Escobar G, Manrique E, Ruiz-Cabrejos J, Saavedra M, Alava F, Bickersmith S, et al. High-accuracy detection of malaria vector larval habitats using drone-based multispectral imagery. *PLoS Negl Trop Dis.* 2019;13: e0007105. doi:10.1371/journal.pntd.0007105
10. Majambere S, Fillinger U, Sayer DR, Green C, Lindsay SW. Spatial Distribution of Mosquito Larvae and the Potential for Targeted Larval Control in The Gambia. *Am J Trop Med Hyg.* 2008;79: 19–27. doi:10.4269/ajtmh.2008.79.19
11. Grisales N, Poupardin R, Gomez S, Fonseca-Gonzalez I, Ranson H, Lenhart A. Temephos resistance in *Aedes aegypti* in Colombia compromises dengue vector control. *PLoS Negl Trop Dis.* 2013;7: e2438. doi:10.1371/journal.pntd.0002438
12. Organization WH. Guidelines for laboratory and field testing of mosquito larvicides. 2005 [cited 16 Apr 2020]. Available: <https://apps.who.int/iris/handle/10665/69101>
13. Partridge FA, Brown AE, Buckingham SD, Willis NJ, Wynne GM, Forman R, et al. An automated high-throughput system for phenotypic screening of chemical libraries on *C. elegans* and parasitic nematodes. *Int J Parasitol Drugs Drug Resist.* 2018;8: 8–21. doi:10.1016/j.ijpddr.2017.11.004
14. Laufer J, Roche M, Pelhate M, Elliott M, Janes NF, Sattelles DB. Pyrethroid insecticides: Actions of deltamethrin and related compounds on insect axonal sodium channels. *J Insect Physiol.* 1984;30: 341–349. doi:10.1016/0022-1910(84)90089-1
15. Laufer J, Pelhate M, Sattelle DB. Actions of pyrethroid insecticides on insect axonal sodium channels. *Pestic Sci.* 1985;16: 651–661. doi:10.1002/ps.2780160611
16. Sattelle DB, Yamamoto D. Molecular targets of pyrethroid insecticides. In: Evans PD, Wigglesworth VB, editors. *Advances in Insect Physiology*. Academic Press; 1988. pp. 147–213. doi:10.1016/S0065-2806(08)60025-9
17. Christophers SR. *Aedes Aegypti* (L.), the Yellow Fever mosquito: its life history, bionomics, and structure. University Press; 1960.

18. Brackenbury J. Locomotion through use of the mouth brushes in the larva of *Culex pipiens* (Diptera: Culicidae). *Proc R Soc Lond B Biol Sci.* 2001;268: 101–106. doi:10.1098/rspb.2000.1336
19. Brackenbury J. Regulation of swimming in the *Culex pipiens* (Diptera, Culicidae) pupa: kinematics and locomotory trajectories. *J Exp Biol.* 1999;202: 2521–2529.
20. Stewart ZP, Oxborough RM, Tungu PK, Kirby MJ, Rowland MW, Irish SR. Indoor application of attractive toxic sugar bait (ATSB) in combination with mosquito nets for control of pyrethroid-resistant mosquitoes. *PLOS ONE.* 2013;8: e84168. doi:10.1371/journal.pone.0084168
21. Planer JD, Hulverson MA, Arif JA, Ranade RM, Don R, Buckner FS. Synergy testing of FDA-approved drugs identifies potent drug combinations against *Trypanosoma cruzi*. *PLoS Negl Trop Dis.* 2014;8: e2977. doi:10.1371/journal.pntd.0002977
22. Duffy S, Sykes ML, Jones AJ, Shelper TB, Simpson M, Lang R, et al. Screening the Medicines for Malaria Venture Pathogen Box across multiple pathogens reclassifies starting points for open-source drug discovery. *Antimicrob Agents Chemother.* 2017;61. doi:10.1128/AAC.00379-17
23. Partridge FA, Murphy EA, Willis NJ, Bataille CJR, Forman R, Heyer-Chauhan N, et al. Dihydrobenz[e][1,4]oxazepin-2(3H)-ones, a new anthelmintic chemotype immobilising whipworm and reducing infectivity *in vivo*. *PLoS Negl Trop Dis.* 2017;11: e0005359. doi:10.1371/journal.pntd.0005359
24. Sugiyama K, Katayama H, Saito T. Effect of insecticides on the mortalities of three whitefly parasitoid species, *Eretmocerus mundus*, *Eretmocerus eremicus* and *Encarsia formosa* (Hymenoptera: Aphelinidae). *Appl Entomol Zool.* 2011;46: 311–317. doi:10.1007/s13355-011-0044-z
25. Ohta I, Takeda M. Acute toxicities of 42 pesticides used for green peppers to an aphid parasitoid, *Aphidius gifuensis* (Hymenoptera: Braconidae), in adult and mummy stages. *Appl Entomol Zool.* 2015;50: 207–212. doi:10.1007/s13355-015-0323-1
26. Jiao Y, Preston S, Song H, Jabbar A, Liu Y, Baell J, et al. Assessing the anthelmintic activity of pyrazole-5-carboxamide derivatives against *Haemonchus contortus*. *Parasit Vectors.* 2017;10: 272. doi:10.1186/s13071-017-2191-8
27. Kaiser HB. HI-receptor antagonist treatment of seasonal allergic rhinitis. : 4.
28. Yamashita M, Fukui H, Sugama K, Horio Y, Ito S, Mizuguchi H, et al. Expression cloning of a cDNA encoding the bovine histamine H1 receptor. *Proc Natl Acad Sci.* 1991;88: 11515–11519. doi:10.1073/pnas.88.24.11515
29. Hardie RC. Is histamine a neurotransmitter in insect photoreceptors? *J Comp Physiol [A].* 1987;161: 201–213. doi:10.1007/bf00615241
30. Zheng Y, Hirschberg B, Yuan J, Wang AP, Hunt DC, Ludmerer SW, et al. Identification of two novel *Drosophila melanogaster* histamine-gated chloride channel subunits expressed in the eye. *J Biol Chem.* 2002;277: 2000–2005. doi:10.1074/jbc.M107635200

31. Roeder T. Metabotropic histamine receptors—nothing for invertebrates? *Eur J Pharmacol.* 2003;466: 85–90. doi:10.1016/S0014-2999(03)01553-X
32. Raymond V, Sattelle DB. Novel animal-health drug targets from ligand-gated chloride channels. *Nat Rev Drug Discov.* 2002;1: 427–436. doi:10.1038/nrd821
33. Bloomquist JR. Chloride channels as tools for developing selective insecticides. *Arch Insect Biochem Physiol.* 2003;54: 145–156. doi:10.1002/arch.10112
34. Iovchev M, Kodrov P, Wolstenholme AJ, Pak WL, Semenov EP. Altered drug resistance and recovery from paralysis in *Drosophila melanogaster* with a deficient histamine-gated chloride channel. *J Neurogenet.* 2002;16: 249–261. doi:10.1080/01677060216293
35. Yusein S, Velikova N, Kuppenova P, Hardie R, Wolstenholme A, Semenov E. Altered ivermectin pharmacology and defective visual system in *Drosophila* mutants for histamine receptor HCLB. *Invert Neurosci.* 2008;8: 211–222. doi:10.1007/s10158-008-0078-2
36. Kita T, Irie T, Nomura K, Ozoe F, Ozoe Y. Pharmacological characterization of histamine-gated chloride channels from the housefly *Musca domestica*. *NeuroToxicology.* 2017;60: 245–253. doi:10.1016/j.neuro.2016.04.019
37. Pantazis A, Segaran A, Liu C-H, Nikolaev A, Rister J, Thum AS, et al. Distinct Roles for Two Histamine Receptors (hclA and hclB) at the *Drosophila* Photoreceptor Synapse. *J Neurosci.* 2008;28: 7250–7259. doi:10.1523/JNEUROSCI.1654-08.2008
38. Hemingway J. The way forward for vector control. *Science.* 2017;358: 998–999. doi:10.1126/science.aaj1644
39. Pirnstill CW, Coté GL. Malaria diagnosis using a mobile phone polarized microscope. *Sci Rep.* 2015;5: 1–13. doi:10.1038/srep13368
40. Rosado L, da Costa JMC, Elias D, Cardoso JS. Mobile-based analysis of malaria-infected thin blood smears: automated species and life cycle stage determination. *Sensors.* 2017;17. doi:10.3390/s17102167
41. Das DK, Mukherjee R, Chakraborty C. Computational microscopic imaging for malaria parasite detection: a systematic review. *J Microsc.* 2015;260: 1–19. doi:10.1111/jmi.12270
42. Moyes CL, Wiebe A, Gleave K, Trett A, Hancock PA, Padonou GG, et al. Analysis-ready datasets for insecticide resistance phenotype and genotype frequency in African malaria vectors. *Sci Data.* 2019;6: 121. doi:10.1038/s41597-019-0134-2
43. Hancock PA, Hendriks CJM, Tangena J-A, Gibson H, Hemingway J, Coleman M, et al. Mapping trends in insecticide resistance phenotypes in African malaria vectors. *PLOS Biol.* 2020;18: e3000633. doi:10.1371/journal.pbio.3000633

Figure Legends

Figure 1. The procedure for automating the analysis of mosquito larval swimming. In each trial, 30 images are acquired at 10 ms intervals and stored for later offline analysis. An index of the amount of movement is obtained by measuring the variance for each pixel over time. Pixels for which the variance more than 3 standard deviations from the mean variance are scored as 1, the remainder as 0. The movement index for each well is taken as the sum of these scores for that well. The output of the algorithm for quantifying movement plotted against the concentration of temephos (bottom right), a larvicide commonly used in the control of mosquitoes, is shown at the end of the pipeline. Alongside this similar data for larvae of *An. gambiae* are shown. A concentration-dependent inhibition of movement is seen in studies on larvae of *An. gambiae* and *Ae. aegypti*.

Figure 2. Screening a chemical library on *An. gambiae* G3 larvae using the INVAPP / Paragon system. Concentration response curves for the five compounds in the Medicines for Malaria Venture (MMV) Pathogen Box library identified as active in the primary screen. The movement index for each well was measured before adding the compounds, and then again 240 min later. The second reading is divided by the first to normalize for background variations between the wells caused, for example, by differences in the number of larvae dispensed in each well. The compounds were applied over the range 3.0×10^{-9} M - 1.0×10^{-4} M, although only concentrations above 7.8×10^{-7} M are shown for clarity.

Figure 3. Motility measurement following deltamethrin exposure in concentration- and time-dependent manner for *An. Gambiae* susceptible (G3) and resistant (Tiassale) strains as well as for *Ae. Aegypti* susceptible (New Orleans) and resistant (Cayman) strains. The normalized movement index, which is the movement index (see Methods) divided by the index recorded before the application of

insecticide, is plotted against the concentration after 60 min applications (A) and the time following exposure (B) to 10^{-6} M deltamethrin. Increased susceptibility is apparent in the adult susceptible strains (blue: G3 (G), New Orleans (N)) as compared to the adult resistant strains (red: Tiassale (T), Cayman (C)). The dotted line indicates the value before insecticide application. Error bars indicate ± 1 s.e.m.

Figure 4. The application of the standard WHO larval assay to detect resistance in *Ae. aegypti*.


Deltamethrin was tested over a range between 5.0×10^{-10} M to 1.0×10^{-6} M. (A) The concentration-response curves are given for the wild-type New Orleans strain (blue) and the Deltamethrin-resistant Cayman strain (red). (B) The fitted P/C_{50} s estimated from fitting the curves in A to a Sigmoid curve. Each point in (B) represents the mean of 3 (WHO) separate experiments and error bars indicate the standard error of the mean.

Figure 5. The output of a smartphone application that determines deltamethrin resistance by mosquito larvae motion, quantified by the average pixel variance over 2 s. (A) The motility of *Aedes aegypti* New Orleans and Cayman larvae, presented as pixel variance over time for the sample with 10^{-5} M deltamethrin in water and DMSO/water only controls. The spike at t 200s is the addition of the deltamethrin or control (water). The solid lines show the mean and the shaded area, the standard deviation (n = 3). (B) The motility of *Anopheles gambiae* G3 and *An. coluzzii* Tiassale larvae presented as pixel variance over time for the sample with 10^{-5} M deltamethrin in water and DMSO/water controls. Fitting the variance over time of all the DM samples to exponential decay functions allows the extraction of the decay rate in each case. These are plotted on the bar chart with error bars showing the standard deviation of the fitted decay rate (n = 3). A higher decay rate is observed for the


susceptible strains compared to their respective resistant strains. *Aedes aegypti* $p = 1.9 \times 10^{-5}$ (one-tailed t-test, $T(2)=19.8$); *Anopheles* $p = 1.1 \times 10^{-2}$ (one-tailed t-test, $T(2)=6.6$).

Supplementary Table S1. The top 20 hits identified in a screen of the Malaria for Medicines Venture box. The rows are sorted by the values of the movement index. The phenotype and location in the box are shown. In the “Phenotype” column, G stands for G3 (susceptible) and T for Tiassale (resistant).


Supplementary Table S2. The raw data from the Malaria for Medicines Venture box are shown, with details provided with the MMV box. In the “Phenotype” column, G stands for G3 (susceptible) and T for Tiassale (resistant).



Acquire images at 67 s^{-1}



Save as 30-image stack



Calculate intensity variance over time dimension




Mark pixels where variance is more than 3 std above mean


Count these pixels in each well

8 x 12 matrix of movement indices


mean of 10




DMSO




MMV687775




MMV007803




MMV687776









MMV637229




MMV688934







A



B

