

1 **Antioxidant nanozyme counteracts HIV-1 by modulating intracellular redox** 2 **potential**

3
4 Shalini Singh^{1, 2}, Sourav Ghosh³, Virender Kumar Pal^{1, 2}, MohamedHusen Munshi²,
5 Raju S Rajmani², Govindasamy Mugesh^{3*}, and Amit Singh^{1, 2*}
6

7 ¹Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore-
8 560012, India

9 ²Centre for Infectious Disease Research (CIDR), Indian Institute of Science,
10 Bangalore 560012, India

11 ³Department of Inorganic and Physical Chemistry, Indian Institute of Science,
12 Bangalore 560012, India.

13

14

15

16 *** Correspondence:**

17 Amit Singh (asingh@iisc.ac.in) and Govindasamy Mugesh (mugesh@iisc.ac.in)

18

19 **Classification**

20 Major – Biological sciences; Minor – Applied biological sciences

21 **Keywords**

22 Nanozymes, HIV, glutathione peroxidase, glutathione, latency

23

24

25

26

27

Abstract

Reactive oxygen species (ROS) regulates the proliferation of human immunodeficiency virus (HIV-1) and *Mycobacterium tuberculosis* (*Mtb*) inside the infected immune cells. However, the application of this knowledge to develop therapeutic strategies remained unsuccessful due to unfavorable consequences of manipulating cellular antioxidant systems that respond to ROS. Here, we show that vanadium pentoxide (V_2O_5) nanosheets functionally mimic the activity of natural glutathione peroxidase (GPX) to mitigate ROS associated with HIV-1 infection without triggering detrimental changes in cellular physiology. Using genetic reporters of glutathione (GSH) redox potential (E_{GSH} ; Grx1-roGFP2) and H_2O_2 (Orp1-roGFP2), we showed that V_2O_5 -nanosheets catalyze GSH-dependent neutralization of ROS in HIV-1 infected cells. Notably, V_2O_5 -nanosheets uniformly blocked HIV-1 reactivation, multiplication, and impaired survival of drug-resistant *Mtb* during HIV-TB co-infection. Mechanistically, V_2O_5 -nanosheets suppressed HIV-1 by affecting the expression of pathways coordinating redox balance, virus transactivation (e.g., NF- κ B and FOS), inflammation, and apoptosis. Importantly, a combination of V_2O_5 -nanosheets with a pharmacological inhibitor of NF- κ B (BAY11-7082) abrogated activation of HIV-1 from latency. Lastly, V_2O_5 -nanosheets counteracted ROS, disease pathophysiology, and virus expression in HIV-1 transgenic mice. Our data successfully revealed the usefulness of V_2O_5 -nanosheets against human pathogens and suggest nanozymes as future platforms to develop interventions against infectious diseases.

53 **Significance Statement**

54 Redox stress, such as those caused by the deregulation of the antioxidant
 55 glutathione, promotes the multiplication of human immunodeficiency virus-1 (HIV-1)
 56 and *Mycobacterium tuberculosis* (*Mtb*). Here we present a vanadium pentoxide
 57 (V₂O₅)-based antioxidant nanozyme that targets cells infected with HIV-1. The
 58 nanozyme, by mimicking the activity of glutathione peroxidase, reprograms redox
 59 signaling to subvert HIV-1 from monocytes, lymphocytes, and HIV-1 transgenic mice.
 60 Treatment with nanozyme bolsters the antiviral potential of immune cells by reducing
 61 the expression of genes involved in virus activation, inflammation, and apoptosis.
 62 The nanozyme also inhibited the proliferation of *Mtb*, which is a major cause of
 63 lethality in HIV patients. These V₂O₅-based nanozymes may be applied to numerous
 64 human pathogens where redox signaling contributes to disease progression.

65

66

67

68

69

70 Introduction

71 Nanomaterials with intrinsic enzyme-mimetic properties (nanozymes) have
 72 been explored as low-cost alternatives to natural enzymes (1). The application of
 73 these nanomaterials was largely restricted to industries for chemical synthesis,
 74 detection of biomolecules, and bioremediation (1, 2). While largely ignored by the
 75 biomedical community, recent studies provide evidence for the clinical importance of
 76 artificial nanozymes *in vivo* (3). For example, ceria-based nanoparticles (NPs) mimic
 77 superoxide dismutase (SOD) activity and exhibit neuroprotection and reduced
 78 inflammation (4, 5). Similarly, iron oxide-based nanoparticles mimic peroxidase-like
 79 activity and protect from bacterial biofilms associated with oral infection (6).
 80 Moreover, ferumoxytol, an FDA approved iron-oxide nanoparticle has been shown to
 81 inhibit tumor growth in mice (7). Recently, we reported that vanadium pentoxide
 82 (V_2O_5) nanomaterial could protect mammalian cells from oxidative damage and
 83 apoptosis by glutathione peroxidase (GPX)-like activity in the presence of cellular
 84 antioxidant glutathione (GSH) (8). These findings indicate the importance of
 85 nanozymes mimicking the activity of natural antioxidant enzymes for biomedical
 86 applications.

87

88 Our particular interest is to apply antioxidant nanozymes in oxidative stress
 89 linked human infections where therapeutic options are inadequate. In this context,
 90 oxidative stress is central to the infection caused by one of the major global
 91 pathogen human immuno-deficiency virus (HIV-1; the causative agent of the
 92 acquired immuno-deficiency syndrome [AIDS]) (9). A major barrier to curing HIV-1
 93 infection is latency, wherein the infected cells harbor the intact viral genome that is
 94 replication-competent but transcriptionally silent. Interestingly, oxidative stress is

known to reactivate HIV-1 from latent reservoirs via NF- κ B directed transcriptional activation of the viral long terminal repeat (LTR) (10, 11). Further studies linking oxidative stress with HIV-1 infection demonstrate variations in GSH levels in infected cells and tissues (12-14). Using a non-invasive biosensor of GSH redox potential (E_{GSH} ; Grx1-roGFP2), we discovered that reductive E_{GSH} sustains viral latency, whereas a marginal oxidative shift in E_{GSH} promotes HIV-1 reactivation (15).

101

In addition to GSH, decreased expression levels of host GPXs are associated with HIV-1 reactivation and replication (15, 16). The HIV-1 genome also encodes a fully functional GPX (HIV-1 vGPX) module (17), which protects cells from reactive oxygen species (ROS) induced apoptosis and possibly helps HIV-1 to maintain latency (17). Consistent with this, majority of the viral isolates from individuals naturally competent to maintain HIV-1 in a latent state (long-term non-progressors [LTNPs]) possess a functional vGPX gene (18). In contrast, HIV-1 strains from the patients with actively replicating virus contain a non-functional vGPX (18). Since LTNPs serve as a natural model for slowing progression of HIV in humans, the association of functional vGPX with these individuals reinforces the physiological importance of the GSH/GPX axis in HIV-1 latency. Despite these observations, efforts to mitigate oxidative stress for subverting HIV-1 reactivation by either supplementation of GSH precursor (N-acetylcysteine [NAC]) or activation/over-expression of GPXs yielded inconsistent results (19-22). While counterintuitive, these findings are in agreement with several studies showing the adverse influence of uncontrolled overproduction of natural antioxidant systems (including GPXs) on redox metabolism and disease outcome (22). Further, over-expression of GPXs might not be sufficient as cellular GPX activity is enhanced by oxidative stress

responsive post-translational modifications (PTMs) such as phosphorylation, carbonylation and O-GlcNAcylation (23-25). Besides this, the activity of GPXs is also dependent on the availability of selenium (Se), an essential micronutrient that has been reported to be low in HIV patients (16). In this context, artificial nanozymes mimicking GPX-like activity (e.g., V_2O_5 NPs) with high sensitivity and specificity under physiological conditions found in the human body (i.e. mild temperature, pH 4-8, and aqueous buffer) can provide a suitable alternate to natural GPXs. We envisage that antioxidant nanozymes can be exploited to generate new knowledge on redox signaling mechanisms underlying HIV-1 latency and reactivation, which could aid the development of fresh therapeutic approaches for targeting HIV.

In this work, we exploited multiple interdisciplinary approaches to describe the utility of V_2O_5 thin-nanosheet (Vs) displaying functional GPX activity in dissecting redox signaling underlying HIV-1 latency. By exploiting genetic reporters, cellular models of HIV-1 latency, and HIV transgenic mice (Tg26), we discovered that Vs-mediated remediation of ROS efficiently subverted reactivation of HIV-1 and survival of another human pathogen (*Mycobacterium tuberculosis* [Mtb]) that frequently infects HIV patients. Our study provides an elegant example of how antioxidant nanozymes can be explored to understand the redox basis of human infections.

Results

Synthesis and biophysical characterization of catalytically efficient V_2O_5 thin-nanosheets (Vs)

We have recently shown that the V_2O_5 nanomaterials exhibit excellent isoform-specific GPX activity, which is dependent on the surface exposed crystal

facets (26). To make V_2O_5 nanomaterials biologically useful, we focused on two distinct morphologies of V_2O_5 nanomaterials i.e. wires (VNw) and sheets (VSh) (Fig. S1, A and B). Both the morphologies displayed similar surface-exposed crystal facet {001}. However, VSh exhibited a higher GPX reaction rate as compared to VNw as determined by measuring the decrease in NADPH absorbance at 340 nm using glutathione reductase (GR)-coupled assay (see *Materials and Methods*) (Fig. S1C). Smaller nanoparticles often display higher catalytic activity (27-31). Therefore, we made subtle modifications in VSh preparation to generate ultrathin-nanosheets (Vs) (Fig. 1A). We confirmed the morphology of Vs by scanning electron microscopy (SEM) (Fig. 1B). Examination of the rate of GPX activity revealed that Vs possesses a 1.6- to 2-fold higher capacity to reduce H_2O_2 in the GR-coupled assay (Fig. 1C) as compared to VNw and VSh (Fig. 1D). A comparison of activities with three different peroxides - H_2O_2 , tertiary-butyl hydroperoxide (*t*-BuOOH) and cumene hydroperoxide (Cum-OOH), indicates that Vs is very selective towards H_2O_2 (Fig. 1E). Based on this, we carried out the extensive biophysical and biochemical characterization of Vs *in vitro*.

First, we recorded the crystalline nature of the lyophilized, thin nanosheets, Vs, by the powder X-ray diffraction pattern (PXRD) (Fig. S2A). The PXRD pattern was indexed to the standard V_2O_5 orthorhombic phase ($a = 11.5160 \text{ \AA}$, $b = 3.5656 \text{ \AA}$, $c = 4.3727 \text{ \AA}$, JCPDS = 41-1426, Space group Pmmn). Second, we examined the crystal facets exposed in the Vs material using high-resolution transmission electron microscopy (HRTEM) and selected area electron diffraction (SAED) pattern analysis. The observed lattice fringes for Vs are (200) and (110) with d-spacing of 0.58 nm and 0.34 nm, respectively, with the interfacial angle of 72.8° . The interfacial angle

between two planes (110) and (-110) with equidistant d-spacing is 34.4° (Fig. S2B and 1F). These three planes in the HRTEM fall in the common zone axis [001]. In both the figures, SAED patterns (inset) were well indexed along [001] zone axis, which confirms that the surface exposed facets are indeed [001]. These results agree with the most intense peak due to (001) plane observed in the PXRD pattern of Vs.

Third, we performed FT-Raman spectroscopy to determine the nature of bonding formed between the metal and oxygen atoms in the orthorhombic V_2O_5 crystals. The FT-Raman spectra showed a peak around 995 cm^{-1} , which corresponds to the terminal (V=O) resulting from the unshared oxygen atom of the V_2O_5 crystal (Fig. 1G). The peaks detected at lower vibration frequencies are consistent with the lattice vibrations of layered material (32-34). We examined the purity of Vs by confirming the detection signal for vanadium (V) and oxygen (O_2) using point energy dispersive X-ray spectroscopy (EDS) (Fig. 1H). A small signal detected at 2 KeV is due to the gold (Au) sputtering of the sample during spectrum recording. Fourth, we confirmed the elemental composition and purity of the Vs material by selected area bright field (SABF) images and X-ray mapping images (Fig. 1I). Both these techniques confirmed that Vs material has a homogenous distribution of vanadium (V) and oxygen (O_2) (Fig. 1, J and K). Finally, we determined the oxidation state of the Vs material using X-ray photoelectron spectroscopy (XPS). The analysis revealed binding energies (BE) and full width at half maxima (FWHM) for the $V2p_{3/2}$ and $V2p_{1/2}$ peaks as well as the difference in the BE between $O1s$ and $V2p_{3/2}$ orbitals (12.8 eV). All of this confirms that vanadium exists in the +5-oxidation state in Vs (Fig. S3A – S3C).

195

196 **Vs displays efficient H₂O₂ linked GPX activity**

197 Having established the biophysical characteristics of Vs, we examined the
 198 GPX-mimetic activity of Vs using GR-coupled assay as described earlier (Fig. 1C).
 199 Various control experiments, such as reactions lacking GSH/GR/H₂O₂, were
 200 performed to rule out the possibility of nonspecific reactions. We observed that in the
 201 absence of any one of the required constituents, Vs does not show GPX-like activity
 202 i.e., no reduction of H₂O₂ takes place (Fig. 2A, and S4A). Varying the concentration
 203 of Vs from 0 to 20 ng/μL led to a proportional dependence of the initial rate for the
 204 reduction of H₂O₂ with first-order reaction kinetics (Fig. 2B). Since both H₂O₂ and
 205 GSH are important for the GPX activity, we performed the activity assay by varying
 206 concentrations of H₂O₂ (0-400 μM) and GSH (0-7 mM) under steady state condition.
 207 Typical enzymatic Michaelis-Menten kinetics was observed for both H₂O₂ and GSH
 208 (Fig. 2, C and D). The corresponding Lineweaver-Burk plots are depicted in fig. 2E
 209 and 2F. For comparison, we simultaneously performed kinetics of Vs, VSh, and
 210 VNw. Interestingly, the V_{max} values for Vs and VSh correlate with their surface area
 211 with an exception of VNw (Fig. 2, G and H). This is consistent with the differences in
 212 the {001} exposed facets among three morphologies. For H₂O₂, the K_M values
 213 obtained for VNw, VSh, and Vs were 44.4 ± 1.7, 57.3 ± 3.8, and 112.2 ± 3.8 μM,
 214 respectively (Fig. 2H). This indicates that the surface of the nanowires and
 215 nanosheets (VNw & VSh) are saturated at lower concentrations of H₂O₂ (26),
 216 whereas relatively higher concentrations of H₂O₂ are required for the saturation of
 217 the surface of thin nanosheets (Vs).

218

The stability of the nanomaterials for the reduction of H₂O₂ was examined by performing multiple assay cycles, which demonstrate only a marginal loss of catalytic activity (Fig. S5A). TEM measurements of nanomaterial surface before and after multiple rounds of catalysis indicate no alterations (Fig. S5B and S5C), confirming that Vs performs H₂O₂ reduction with unprecedented stability, specificity, and sensitivity *in vitro*.

Vs mimics GPX activity inside the HIV-1 infected cells

To test the Vs-related GPX activity inside mammalian cells and to understand its influence on HIV-1, we selected the U1 cell line model of HIV-1 latency and reactivation. The U1 cell line is derived from the parent promonocytic cell line U937, wherein two copies of the HIV-1 genome is latently integrated (35). The viral replication can be induced by treatment of U1 cells with various pro-inflammatory agents such as phorbol myristate acetate (PMA), tumor necrosis factor-alpha (TNF- α), and granulocyte-macrophage colony-stimulating factor GM-CSF (35,36). We first examined the uptake of Vs nanomaterial by U1 through inductively coupled plasma mass spectrometry (ICP-MS). As shown in fig. 3A, 15 min of Vs exposure led to its buildup inside U1, which was gradually decreased over time such that only a fraction of Vs was retained intracellularly (Fig. 3A). We also found that survival of U1 was not adversely affected by a range of Vs concentration (Fig. S6A).

To examine the role of Vs in the intracellular reduction of H₂O₂ through GPX activity, we exploited Orp1-roGFP2 and Grx1-roGFP2 biosensors that allow non-invasive imaging of the intracellular H₂O₂ and E_{GSH} , respectively (37-39). The roGFP2 moiety has two surface exposed cysteines, which undergo disulfide bond

formation upon oxidation resulting in an increase in fluorescence excitation intensity at 405 nm along with a relative decrease at 488 nm excitation at a fixed emission of 510 nm (37). An inverse relation in 405/488-biosensor ratio was detected upon the reduction of disulfides. The specific equilibration of the roGFP2 dithiol-disulfide redox pair (roGFP2_{red}/roGFP2_{oxi}) either with H₂O₂ or with glutathione redox pair (GSH/GSSG) is efficiently catalyzed by the covalently fused peroxidase Orp1 or glutaredoxin Grx1, respectively (38, 39). The fusion of Orp1 with roGFP2 creates a redox relay in which Orp1 mediates near-stoichiometric oxidation of roGFP2 by H₂O₂ (38) (Fig. 3B). Likewise, Grx1 mediates oxidation of roGFP2 in response to a nanomolar increase in intracellular GSSG upon H₂O₂ stress (37). The redox relay created by Orp1-roGFP2 or Grx1-roGFP2 demonstrates dynamic behavior as the biosensor ratio returns to basal during recovery from oxidative stress due to normalization of H₂O₂ and GSSG levels (37, 39).

We created stable transfected U1 cells that express either Orp1-roGFP2 (U1-Orp1-roGFP2) or Grx1-roGFP2 (U1-Grx1-roGFP2) in the cytosol (Fig. S7, A and B). Exposure of U1-Orp1-roGFP2 to H₂O₂ for 2 min showed a concentration-dependent increase in the biosensor ratio, consistent with the Orp1-mediated oxidation of roGFP2 by H₂O₂ (Fig. 3C). In contrast, pretreatment of U1-Orp1-roGFP2 with 25 and 50 ng/μL of Vs for 15 min diminished biosensor oxidation by H₂O₂, consistent with the Vs-catalyzed reduction of H₂O₂ (Fig. 3C). Because H₂O₂ exposure also leads to oxidation of reduced GSH to GSSG (9), we monitored this transformation using Grx1-roGFP2 biosensor (Fig. 3D). The U1-Grx1-roGFP2 cells were challenged with various concentrations of H₂O₂ for 2 min, and the sensor response was quantified. We found that the biosensor responds to increasing concentrations of H₂O₂ and

treatment with 100 μ M of H_2O_2 for 2 min results in 90% oxidation of Grx1-roGFP2 (Fig. 3E). The corresponding E_{GSH} was -240 mV, which is higher than the basal E_{GSH} for U1 cells, -320 mV. Pretreatment of U1-Grx1-roGFP2 with Vs for 15 min effectively reduced H_2O_2 mediated oxidation of biosensor in a concentration dependent manner (Fig. 3E). Next, we measured the time kinetics of Grx1-roGFP2 oxidation to a low concentration of H_2O_2 (50 μ M). An increase in the biosensor ratio was observed within 2 min of H_2O_2 exposure followed by a gradual decrease to the baseline levels in 30 min, indicating efficient mobilization of cellular antioxidant machinery (37) (Fig. 3F). In contrast, the addition of 50 ng/ μ L of Vs at post H_2O_2 treatment decreased the biosensor oxidation to baseline levels within 10 min (Fig. 3F). Importantly, a single dose of Vs completely prevented subsequent oxidation of biosensor by H_2O_2 (Fig. 3F). This data is fully consistent with earlier results demonstrating multiple cycles of H_2O_2 reduction by a single dose of Vs *in vitro*.

Since GPX function is dependent on GSH as an electron donor (40), we tested the requirement of GSH in Vs-mediated H_2O_2 reduction. We treated U1-Grx1-roGFP2 cells with 0.5 mM buthionine sulfoximine (BSO), which lowers cellular GSH content by inhibiting γ -glutamyl cysteine synthetase (GCS) activity (41). Following this, cells were treated with Vs for 15 min and exposed to 50 and 100 μ M of H_2O_2 for 2 min. As shown earlier, both the concentrations of H_2O_2 achieved nearly complete oxidation of Grx1-roGFP2, which was effectively blocked by Vs pretreatment (Fig. 3G). In contrast, pretreatment with BSO attenuated Vs ability to prevent biosensor oxidation by H_2O_2 (Fig. 3G). Supplementation of exogenous GSH (15 mM) restored Vs activity as shown by a significant decrease in the biosensor oxidation upon challenge with a saturating concentration of H_2O_2 (Fig. 3H).

Vs subverts HIV-1 reactivation

Studies have shown that H₂O₂ treatment reactivates HIV-1 from latency (15, 42). Increased oxidative stress was shown to activate the HIV-1 LTR through redox-sensitive transcription factors, such as NF-κB (11). On this basis, we reasoned that Vs displaying efficient antioxidant activity could affect redox-dependent reactivation of HIV-1. We first induced HIV-1 expression using low concentrations of PMA (5 ng/mL) and prostatin (1.25 μM), two well-established activators of HIV-1 (43, 44). The expression of the HIV-1 *gag* transcript was monitored as a marker of HIV-1 activation by qRT-PCR at various time points post-treatment with PMA/prostatin. Both activators induced HIV-1 transcription with a significant increase observed at 24 h post-treatment (Fig. 4A). Pre-exposure of U1 with Vs or N-acetyl cysteine (NAC- a well-established antioxidant) effectively blocked PMA/prostatin-mediated viral reactivation (Fig. 4, A and B). Using U1-Orp1-roGFP2, we confirmed an increase in the intracellular levels of H₂O₂ at 6 and 12 h post-PMA treatment, which was significantly reduced upon Vs pretreatment (Fig. 4C). This indicates that oxidative stress precedes PMA-stimulated virus reactivation and GPX activity associated with Vs counteracted redox-dependent HIV-1 reactivation. The capacity of Vs in lessening HIV-1 activation was also confirmed in a lymphocytic model of HIV-1 latency (J1.1) (Fig. 4D), corroborating that the effect of Vs is not restricted to a cell-type.

HIV infected individuals suffer from selenium (Se) deficiency that adversely affects the activity of Se-dependent GPX enzyme leading to oxidative stress, HIV reactivation, and exacerbation of disease pathology (16, 45). Therefore, Se limitation

is a physiologically relevant stimulus that induces oxidative stress and HIV-1 reactivation (46). We envisage that Se-independent GPX activity of Vs could replenish the impaired activity of cellular GPX under Se deficient conditions to subvert HIV-1 reactivation. To examine this, we starved U1-Grx1-roGFP2 of fetal bovine serum (FBS; the source of Se) and monitored the change in its antioxidant response over time. We observed an increase in biosensor ratio within 30 min of FBS removal, indicating oxidative stress (Fig. S8A). Supplementation of Vs or Se in the culture medium of Se-deficient U1-Grx1-roGFP2 decreased biosensor ratio, signifying alleviation of oxidative stress by Vs (Fig. 4E). As expected, Se-deficiency triggered HIV-1 reactivation in U1, and addition of Vs or Se had an opposite effect (Fig. 4F).

Vs adversely affects intracellular replication of HIV-1 and *Mtb*

Along with reactivation, oxidative stress has been shown to promote HIV-1 replication (9). Therefore, we next examined the influence of Vs activity on HIV-1 replication. First, we used a stable CD4⁺ T cell clone expressing EGFP (CEM-GFP) under HIV-1 LTR. Infection with HIV-1 significantly induces the expression of CEM-GFP (47). The infection of CEM-GFP cells with T cell-tropic HIV-1 provirus (pNL4.3) progressively increased GFP fluorescence over 5 days (Fig. 5A). Addition of 50 ng/μL of Vs for 15 min every 24 h completely blocked GFP expression in the infected CEM-GFP cells (Fig. 5A). We also infected Jurkat CD4⁺ T cells with pNL4.3 virus and measured *gag* transcript, and p24 HIV capsid protein in the whole cell lysate and in the supernatant. Each technique showed a time-dependent increase in HIV-1 replication, while treatment with Vs resulted in a significant inhibition (Fig. 5, B to D). Finally, we infected U937 promonocytic cells with macrophage (M)-tropic HIV pro-

virus, pNLAD8, and HIV-1 replication was measured by measuring *gag* transcript at 24 h post-treatment. As shown in fig. 5E, HIV-1 infected U937 showed a 15-fold increase in *gag* transcript, which was reduced to 3-fold in case of Vs pre-treatment. We also examined if antioxidant potential of Vs confers anti-viral response in primary human CD4 T lymphocytes. We pretreated primary CD4 T cells isolated from human peripheral blood mononuclear cells (PBMCs) with Vs (25 ng/μl), infected with HIV-1 NL4.3, and measured HIV-1 replication at 3- and 5-days post-infection. The p24 ELISA confirmed 100 to 200-fold increase in virus replication at 3 to 5 days post-infection, respectively (Fig. 5F). In contrast, pretreatment of Vs restricted virus load to 20- to 80- fold at these time-points (Fig. 5F).

We have previously shown that oxidative stress promotes synergy between *Mtb* and HIV (15, 48). Antioxidant GSH has been shown to reduce *Mtb* load in HIV-infected individuals, while NAC lowers the survival of *Mtb* in mice and guinea pigs (49, 50). On this basis, we assessed if the antioxidant potential of Vs affects the survival of *Mtb* in macrophages infected with HIV-1. PMA-differentiated U1 were infected with a laboratory strain of *Mtb* (H37Rv) and a multidrug-resistant patient isolate (JAL 2287) (51). Bacterial survival was monitored at 4 h and 24 h post-infection. *Mtb* H37Rv and Jal 2287 displayed 1.78- and 2.74-fold increase in survival at 24 h post-infection, respectively (Fig. 5, G and H). Notably, pretreatment with Vs completely abrogated the ability of *Mtb* strains to multiply inside U1 (Fig. 5, G and H). In sum, Vs catalytic activity efficiently counteracts replication of HIV-1 and *Mtb*.

Vs dampens the expression of host genes involved in HIV-1 reactivation

Having shown the utility of Vs in counteracting oxidative stress and HIV-1 reactivation, we next examined the underlying mechanisms. We performed expression analysis using the NanoString nCounter system, which permits absolute quantification of multiple RNA transcripts without any requirements for reverse transcription (52). We focused on 185 host genes that are known to respond to HIV infection and oxidative stress (Table S1A-S1C). We performed expression analysis on RNA isolated from U1, PMA treated U1, Vs treated U1, and Vs plus PMA treated U1. The fold change (> 1.5-fold, P-value < 0.05) was calculated by normalizing the raw mRNA counts to the geometric mean of the internal control β_2 microglobulin (B2M).

A total of 123 genes showed differential expression under the conditions tested (Fig. 6A). Overlap analysis confirmed 55 of 123 genes to be common in each category (Fig. 6B). Treatment with PMA induced the expression of genes associated with ROS and RNS (reactive nitrogen species) generation (e.g., NADPH oxidase subunits [NCF1, NCF2] and nitric oxide synthase [NOS2]). Genes involved in antioxidant response, including catalase (CAT) and superoxide dismutase 2 (SOD2), were downregulated upon PMA treatment (Fig. 6C). Overall, these changes are consistent with increased oxidative stress in response to PMA triggered HIV-1 reactivation (15). Up-regulation of genes involved in reducing free iron pool (ferritin heavy chain1; FTH1), maintaining GSH balance (cysteine/glutamate transporter; SLC7A11), (Fig. 6C), indicate a compensatory mechanism to protect from oxidative conditions induced by PMA (53,54). The transcription factor Nrf2 is the major activator of antioxidant systems (55). Surprisingly, a majority of Nrf2 dependent antioxidant systems such as GSH biosynthesis/recycling (e.g., GSS, GPX1, GPX4, GSTP1), thioredoxins (e.g., TXNRD2), and peroxiredoxins (PRDX6) were down

regulated upon treatment with Vs alone or Vs plus PMA (Fig. 6C), indicating an adverse influence of Vs on U1 cells. One likely possibility is that the natural antioxidant defense mechanisms are attenuated by a feedback-like mechanism because of the potent antioxidant properties of Vs. Also, the expression of superoxide producing system (NCF1 and NCF2) was repressed in Vs alone and Vs plus PMA treated U1 (Fig. 6C), which can further reduce intracellular ROS levels.

Genes known to be associated with HIV-1 activation such as transcription factors (*e.g.*, FOS and CEBPB) (56, 57), inflammatory cytokines/receptors (TGF β 1, TNFRSF1B, and IL16) (58-60), and chemokines (CCL3 and CCL4) (61) were induced upon PMA treatment and repressed by Vs plus PMA (Fig. 6C). Several genes encoding proteins associated with HIV-1 replication, packaging, budding, and fitness (*e.g.*, APOBEC3G, CD44, XPO1, VPS4A, DHCR24) were down-regulated upon Vs plus PMA treatment as compared to PMA alone (Fig. 6D). It is known that cells latently infected with HIV-1 are refractory to apoptosis, whereas increased apoptosis promotes HIV-1 reactivation (62). Consistent with this, a majority of genes encoding pro-apoptotic proteins (*e.g.*, BAD, BAX, CASP3, and CASP8) were substantially repressed upon Vs plus PMA treatment as compared to PMA alone (Fig. 6C). In addition, a cellular inhibitor of transcription factor NF- κ B (*i.e.* NFKBIA) was highly induced upon Vs or Vs plus PMA treatment (Fig. 6D). Since NF- κ B is known to activate HIV-1 in response to oxidative stress (10), the induction of its inhibitor (NFKBIA) by Vs is indicative of reduced reactivation of HIV-1 through NF- κ B. Based on this, we hypothesize that a well-established pharmacological inhibitor of NF- κ B (*E*)-3-[(4-methylphenyl)sulfonyl]-2-propenenitrile (BAY11-7082) (63) would synergies with Vs to efficiently subvert HIV-1 reactivation. To examine this, we

exposed U1 cells pretreated with Vs to BAY11-7082 and HIV-1 reactivation in response to PMA was monitored by measuring the levels of *gag* transcript and p24 capsid protein. Consistent with our hypothesis, exposure of U1 to both Vs and BAY11-7082 suppresses reactivation of HIV-1 which supersedes that produced by either Vs or BAY11-7082 alone (Fig 6E and 6F). Overall, Vs not only affected the expression of redox pathways but as a consequence also modulates the expression of pathways coordinating the inflammatory response, viral fitness, transcription, and apoptosis to subvert HIV-1 reactivation.

Vs mitigates oxidative stress and HIV-1 associated pathologies *in vivo*

Given our findings that Vs diminishes oxidative stress to reduce HIV-1 reactivation in cell lines, we sought to determine the impact of Vs on oxidative stress and HIV-1 associated pathologies *in vivo*. We used an HIV-1 transgenic mouse (Tg26), which harbors a replication-incompetent, Δgag variant of pNL4.3/HIV (64). Expression of HIV-1 proteins in Tg26 promotes oxidative stress and recapitulates clinical lesions observed in HIV infected patients (*e.g.*, lung damage and nephropathy) (64-66).

We treated Tg26 C57BL/6 mice with Vs (1 mg/kg body weight) intra-peritoneally (*i.p.*). Untreated or Vs-treated animals were euthanized 3 h post-treatment and oxidative stress was measured in the lungs (Fig. 7A). The lung homogenates were stained with redox-active fluorescent dye 2',7' – dichlorofluorescein diacetate (DCFDA), and subjected to flow cytometry. As shown in figure 7B, total cellular populations within lung homogenates derived from Vs treated Tg26 displayed diminished fluorescence as compared untreated Tg26. Based on the

forward and side scatter gating, we gated different populations representing mostly granulocytes (P2), monocytes (P3) and lymphocytes (P4). The level of fluorescence within multiple cell types was consistently lower in the case of Vs treated Tg26 than untreated Tg26 animals (Fig. 7B). Using antibodies against CD64 and MERTK markers, we specifically gated lung macrophages and confirmed lower ROS in cells derived from Vs treated Tg26 animals than untreated Tg26 (Fig. 7C). The lung cells derived from Wt C57BL/6 showed ROS levels comparable to Vs treated Tg26 animals (Fig. 7, B and C). Consistent with a strong antioxidant activity of Vs, expression of other cellular antioxidant genes were either down-regulated (e.g, GPX1) or unchanged (GPX4, SOD1, and SOD2) in the lungs of Vs treated Tg26 as compared to untreated animals (Fig. 7D and Fig. S9A). One of the clinical signs of HIV infection is inflammation (67). The Tg26 mice display edema in various organs, which is a marker for inflammation and fluid accumulation (65). Treatment with Vs significantly reduced lung edema in Tg26 animals (Fig. 7E). Lastly, we confirmed that Vs treatment reduced the expression of HIV-1 transcript (*tat-rev* region) by ~65% in Tg26 mice (Fig. 7F). Altogether, our findings position Vs as a potential alternative to natural GPX enzyme(s) for regulating redox signaling, inflammation, and virus expression *in vivo*.

Discussion

Studies exploring the application of antioxidant nanozymes in targeting human pathogens are limited. In-depth cellular studies using laboratory models that mimic the physiological environment during infection could help predict the clinical potential of nanozymes and will encourage new designs that are efficacious in humans. Using several biophysical, biochemical, cellular, genetic biosensors, animal model, and

expression technologies, we demonstrated the biomedical application of V₂O₅ based nanozymes in counteracting redox stress and reactivation of HIV-1. We showcase the remarkable efficacy of Vs against various physiological conditions, including Se starvation and infection with secondary pathogens such as *Mtb*.

V₂O₅ nanoparticles possess intrinsic peroxidase-mimicking activity (8). By modifying V₂O₅ nanomaterial into various morphologies, we reported efficient GSH-dependent GPX activity of V₂O₅ nanozymes (26). In the present study, we further fine-tuned the crystal facets of V₂O₅ nanomaterial to generate a thinner nanosheet (Vs), which displayed high catalytic activity in the presence of GSH. Other properties, such as the formation of stable V-peroxo species that ensure GSH specificity, multiple recycling without loss of activity, and stability in a range of organic solvents (698) motivated us to exploit V₂O₅ nanomaterial in understanding GSH/GPX based redox signaling during HIV-1 infection. Until now, studies examining the antioxidant function of nanozymes relied on *in vitro* enzymatic assays or chemical analyses of redox metabolites (e.g., GSH/GSSG) in whole cells or tissues. These invasive methods introduce oxidation artifacts and preclude observation of real-time changes in redox physiology upon nanozyme treatment. We circumvented these issues by applying non-invasive genetically encoded biosensors of H₂O₂ (Orp1-roGFP2) and *E*_{GSH} (Grx1-roGFP2) to dynamically assess the activity of Vs in reducing intracellular H₂O₂ and maintaining GSH homeostasis.

Numerous studies have indicated a link between GPX activity and HIV-1 *in vitro* and *in vivo* (16, 21). Remediation of H₂O₂ by GPXs potentially reversed NF-κB-mediated HIV-1 transcription (21). We have demonstrated that latently infected cells

efficiently metabolized H₂O₂ likely via endogenous GPXs (15). Expression of GPXs was elevated in monocytes and lymphocytes harboring latent HIV-1, whereas expression was diminished in cells and in PBMCs of patients during active HIV-1 replication (15). Since the activity of cellular GPXs is selenium (Se)-dependent, studies have found a high correlation between Se-deficiency, GPX activation, and HIV-related mortality (45). Importantly, HIV-1 infected T-cells showed the general downregulation of cellular Se-proteins (69). We found that Se depletion induced the oxidative shift in *E_{GSH}* and promotes HIV reactivation. Importantly, Vs supplementation under Se depleted conditions was sufficient to diminish oxidative stress and HIV-1 reactivation. Mechanistically, our Nanostring data confirm that GPX-like activity of Vs efficiently suppresses the expression of redox-dependent transcription factors, pro-inflammatory cytokines/chemokines, and pro-apoptotic molecules required for HIV-1 reactivation. In line with our findings, transcriptomics of whole blood RNA from elite controllers (ECs) to that of individuals with high viral load showed that cytokine signaling, apoptosis, anti-viral response, and immune activation are mainly affected (70-72). Our data align well with the meta-analyses where pathways promoting oxidative stress and viral transcription such as NCF2, ACTB, FOS, CEBPB are downregulated in ECs and antioxidants (GPX1 and GPX4) were induced as compared to viremic progressors (71,72).

Individuals co-infected with HIV and *Mtb* pose major problems in disease management. We showed that basic mechanisms such as intracellular redox potential and bioenergetics contribute to synergy between HIV and *Mtb* (15,48,73). Diminished levels of host GSH, GPX4, and increased ROS contribute to the survival of *Mtb in vivo* (74). Similar to these observations, supplementation of GPX activity by

Vs hampered multiplication of *Mtb* and drug-resistant clinical isolate in U1 cells pre-exposed to Vs. Finally, the biological importance of our findings comes from the data showing reduced oxidative stress, inflammation, and expression of HIV-1 transcripts in Tg26 mice treated with Vs. While our findings provide a clear relationship between catalytic activity of Vs with HIV-1 reactivation and redox signaling, future experiments are needed to exploit Vs for the development of sustainable, affordable, and safe therapeutics against HIV-1 infection. The importance of nanomaterials in infectious diseases is emerging as silver and gold-based nanomaterials have been shown as promising candidates for therapeutic applications against bacterial and viral pathogens (75-78).

In conclusion, we have presented compelling experimental evidence that strongly suggests the application of Vs in modulating redox signaling associated with HIV-1 latency and reactivation. Antiretroviral therapy (ART), while effective in reducing viral load, remains unsuccessful in eradicating HIV from latent reservoirs and induces systemic oxidative stress (79). Recently, a concept called “block and lock” showed that blocking host (e.g., heat shock protein 90 (HSP90), NF-κB, protein kinase C (PKC) or viral factors (e.g., Tat) involved in reactivation significantly delayed viral rebound upon interruption of ART (80). Our findings on Vs showed promise in preventing reactivation from latency, raising the possibility of locking HIV in an extended latency by blocking redox-mediated HIV-1 transcription, and also improving the health of individuals on ART by minimizing oxidative stress.

Materials and methods

Preparation of thin vanadia (V₂O₅) nanosheets (Vs)

Thin V_2O_5 nanosheets (Vs) were synthesized from the crude V_2O_5 nanosheets (VSh). Briefly, 2 mM of V_2O_5 powder was dispersed in 15 ml ultrapure water for 20 min. Then 15 mL H_2O_2 (30% w/v) was added dropwise. During the addition of H_2O_2 the colour of the solution changed from yellow to orange and then to red. The red solution turns dark brown after stirring for 2.5 h at room temperature (RT). This reaction was strictly performed in a fume hood due to its exothermic nature. After continuous stirring for 2.5 h, 10 mL of ultrapure water was added into the mixture and heated to 60°C overnight to form a brownish gel ($V_2O_5 \cdot nH_2O$). This gel was dried at 100°C for 12 h and subsequently calcined at 400°C for 2 h to get crude VSh. Following this, VSh was probe sonicated in ultrapure water for 2 h to get a dense dispersion of nanosheets. The dispersion was then centrifuged at 3000 rpm for 5 min and the bright yellow colour supernatant was lyophilised to obtain the powdered form of thin V_2O_5 nanosheets (Vs).

Characterization of Vs

Powder X-ray diffraction (PXRD) was recorded by Phillips PANalytical diffractometer using a $CuK\alpha$ ($\lambda = 1.5406 \text{ \AA}$) radiation. The emission current and accelerating voltage used in the diffractometer were 30 mA and 40 kV respectively. For morphological and elemental characterization, EDS and scanning electron microscopy (SEM) were performed on FEI Sirion UHR SEM and ESEM-Quanta respectively. Transmission electron microscopy (TEM), High resolution transmission electron microscopy (HRTEM), and X-ray mapping images were recorded on JEOL transmission electron microscope operated at 200 kV after casting a drop of nanoparticle dispersion in isopropyl alcohol, over a Cu grid. FT-Raman spectra were recorded using a Renishaw in-Via Raman Microscope (Renishaw Inc, UK), with

excitation wavelength 514 nm. To perform all the enzyme mimetic activity assay, SHIMADAZU UV-2600 spectrophotometer was used. X-ray photoelectron spectroscopy (XPS) was performed using AXIS Ultra, KRATOS ANALYTICAL, SHIMADAZU. The surface area measurement was performed by Brunauer-Emmett-Teller (BET) method on the micromeritics surface area analyzer model ASAP 2020.

GPX – mimicking activity of V₂O₅ nanoparticles (NPs)

The GPX-like activity of V₂O₅ NPs was assessed spectrophotometrically by using the standard GR-coupled GPX assay (8). The components and the concentration used in this assay mixture were GSH (2.0 mM), NADPH (0.2 mM), GR 1.7 U, catalyst 20 ng/μL, and H₂O₂ (0.2 mM) in sodium phosphate buffer pH 7.4 at 25°C. The rate of the reaction was quantified by following the decrease in the absorbance of NADPH ($\epsilon = 6220 \text{ M}^{-1} \text{ cm}^{-1}$ at 340 nm) to form NADP⁺ which is equal to the rate of conversion of H₂O₂ to H₂O.

Dynamic response of U1-Grx1-roGFP2 cells towards oxidative stress

Oxidation-reduction kinetics of the Grx1-roGFP2 biosensor were measured by flow cytometry, as demonstrated earlier (15). Briefly, the basal redox state of 1×10^6 U1-Grx1-roGFP2 cells was measured, following which 50 μM H₂O₂ was added after 2 min. Biosensor oxidation and the kinetics of its subsequent recovery were monitored. Parallely, Vs was added to a set of H₂O₂ treated cells at the point of maximum oxidation, and recovery of the biosensor in the presence of Vs was noted. After complete recovery of the cells from oxidative insult, both untreated and Vs treated cells were challenged with another bolus of 50 μM H₂O₂, and the biosensor dynamics were monitored by flow cytometry. Percentage oxidation of the Grx1-

roGFP2 biosensor was determined by equating maximal oxidation by 10 mM H₂O₂ as 100%.

Determination of the specificity of Vs towards GSH

The specificity of Vs towards GSH as cofactor was determined by modulating cellular GSH levels. U1-Grx1-roGFP2 cells were treated with 0.5 mM of BSO, an inhibitor of GSH biosynthesis, or supplemented with 7.5 or 15 mM GSH for 16 h. Following this, the cells were treated with 50 ng/μL of Vs and challenged with various H₂O₂ concentrations for 2 min. The biosensor response in U1 cells was measured by flow cytometry.

Assessing the effect of Vs on active HIV replication.

CD4⁺ T cell line – CEM-GFP and Jurkat- and monocytic cell line U937 were infected with laboratory adapted HIV-1 strains pNL4.3 (T-tropic) and pNLAD8 (M-tropic), respectively. 0.5×10^6 untreated or Vs treated cells of each type were suspended in 500 μL of Opti MEM media and infected at multiplicity of infection (moi) 0.1. The cells were incubated at 37°C for 4 h and mixed intermittently during the infection period. After 4 h, the cells were washed to remove unbound virus and supplemented with complete RPMI media with 10% FBS. Vs treatment was repeated every 24 h for Jurkat and CEM-GFP cells. CEM-GFP cells were grown till 5 days and LTR activation was assessed from day 2 to day 5 post-infection by flow cytometry. Viral replication in Jurkat and U937 cells was monitored by qRT-PCR at indicated time points.

Nanostring gene expression analysis

Expression levels of 185 genes responsive to oxidative stress and HIV infection were analysed in untreated U1 cells, U1 cells treated with PMA or Vs alone, and a combination of Vs plus PMA. The Nanostring nCounter analysis system was utilized for this purpose. Briefly, the assay was performed with 100 ng of total RNA, isolated from untreated or treated cells using the Qiagen RNeasy kit. The purity of the RNA was confirmed spectrophotometrically using Nanodrop Lite Spectrophotometer (Thermo Scientific). The nCounter probes are barcoded DNA oligonucleotides complementary to the target mRNA. Hybridization and counting were performed according to the manufacturer's protocol (52) using a customized panel of 185 genes. 6 housekeeping control genes were included in the panel. Data analysis was done using nSolver 4.0. B2M was used as an internal control due to its minimum % CV.

Selenium starvation and HIV-1 reactivation

U1-Grx1-roGFP2/U1 cells grown in complete RPMI medium were harvested and washed three times with serum-free RPMI to remove traces of Se. Cells were seeded in 24 well plates and incubated for 30 min, 1 h, and 2 h in serum free medium. Parallely, the cells were treated with various concentrations of Vs for 15 min and cultured as mentioned above. 0.5 nM sodium selenite (Se source) was used as a positive control. Cells were harvested at indicated time points and the biosensor response was measured by flow cytometry. Viral reactivation and expression of host antioxidant genes were analysed 6 h post starvation, as mentioned earlier.

Methods included in the supplementary information

Dispersion of V₂O₅ NPs, treatment of cell lines with Vs, internalization of Vs by U1, mammalian and bacterial Cell Culture, preparation of stable cell lines and validation using flow cytometry, assessment of Vs antioxidant activity and redox potential measurement, propidium iodide (PI) staining, HIV reactivation in U1 cells and qRT-PCR analysis, p24 detection by Immunoblotting and ELISA, bacterial Survival assays in U1 cells, isolation and infection of primary CD4⁺ T cells, animal experiments with HIV-Tg mice, pulmonary edema analysis, ROS staining of lung homogenates.

Statistical analysis

All statistical analyses were performed using the GraphPad Prism software (Version 8.1). The data values are indicated as mean ± S.D. Statistical significance between two non-parametric test groups was determined using the Mann Whitney Rank Sum test, unless specified. Analysis of Nanostring data was performed using the nSolver platform. Differences in *P* values <0.05 were considered significant.

References

1. H. Wei, E. Wang, Nanomaterials with enzyme-like characteristics (nanozymes): next-generation artificial enzymes. *Chem. Soc. Rev.***42**, 6060-6093 (2013).
2. R. de la Rica, M. M. Stevens, Plasmonic ELISA for the ultrasensitive detection of disease biomarkers with the naked eye. *Nat. nanotechno.* **7**, 821-824 (2012)

- 668 3. O. Salata, Applications of nanoparticles in biology and medicine. *J.*
669 *Nanobiotechnology* **2**, 3 (2004).
- 670 4. C. Korsvik, S. Patil, S. Seal, W. T. Self, Superoxide dismutase mimetic
671 properties exhibited by vacancy engineered ceria nanoparticles. *Chem.*
672 *Commun. (Camb)*, 1056-1058 (2007).
- 673 5. M. Das, S. Patil, N. Bhargava, J. F. Kang, L. M. Riedel, S. Seal, J. J.
674 Hickman, Auto-catalytic ceria nanoparticles offer neuroprotection to adult rat
675 spinal cord neurons. *Biomaterials* **28**, 1918-1925 (2007).
- 676 6. L. Gao, Y. Liu, D. Kim, Y. Li, G. Hwang, P. C. Naha, D. P. Cormode, H. Koo,
677 Nanocatalysts promote *Streptococcus mutans* biofilm matrix degradation and
678 enhance bacterial killing to suppress dental caries in vivo. *Biomaterials* **101**,
679 272-284 (2016).
- 680 7. S. Zanganeh, G. Hutter, R. Spitler, O. Lenkov, M. Mahmoudi, A. Shaw, J. S.
681 Pajarinen, H. Nejadnik, S. Goodman, M. Moseley, L. M. Coussens, H. E.
682 Daldrup-Link, Iron oxide nanoparticles inhibit tumour growth by inducing pro-
683 inflammatory macrophage polarization in tumour tissues. *Nat. Nanotechnol.*
684 **11**, 986-994 (2016).
- 685 8. A. A. Vernekar, D. Sinha, S. Srivastava, P. U. Paramasivam, P. D'Silva, G.
686 Mugesh, An antioxidant nanozyme that uncovers the cytoprotective potential
687 of vanadia nanowires. *Nat. Commun.* **5**, 5301 (2014).
- 688 9. A. Perl, K. Banki, Genetic and metabolic control of the mitochondrial
689 transmembrane potential and reactive oxygen intermediate production in HIV
690 disease. *Antioxid. Redox Signal.* **2**, 551-573 (2000).

- 691 10. F. J. Staal, M. Roederer, L. A. Herzenberg, L. A. Herzenberg, Intracellular
692 thiols regulate activation of nuclear factor kappa B and transcription of human
693 immunodeficiency virus. *Proc. Natl. Acad. Sci. U.S.A.* **87**, 9943-9947 (1990).
- 694 11. C. W. Pyo, Y. L. Yang, N. K. Yoo, S. Y. Choi, Reactive oxygen species
695 activate HIV long terminal repeat via post-translational control of NF-kappaB.
696 *Biochem. Biophys. Res. Commun.* **376**, 180-185 (2008).
- 697 12. H. P. Eck, H. Gmunder, M. Hartmann, D. Petzoldt, V. Daniel, W. Droge, Low
698 concentrations of acid-soluble thiol (cysteine) in the blood plasma of HIV-1-
699 infected patients. *Biol. Chem. Hoppe Seyler* **370**, 101-108 (1989).
- 700 13. R. Buhl, H. A. Jaffe, K. J. Holroyd, F. B. Wells, A. Mastrangeli, C. Saltini, A. M.
701 Cantin, R. G. Crystal, Systemic glutathione deficiency in symptom-free HIV-
702 seropositive individuals. *Lancet* **2**, 1294-1298 (1989).
- 703 14. L. A. Herzenberg, S. C. De Rosa, J. G. Dubs, M. Roederer, M. T. Anderson,
704 S. W. Ela, S. C. Deresinski, L. A. Herzenberg, Glutathione deficiency is
705 associated with impaired survival in HIV disease. *Proc. Natl. Acad. Sci. U.S.A.*
706 **94**, 1967-1972 (1997).
- 707 15. A. Bhaskar, M. Munshi, S. Z. Khan, S. Fatima, R. Arya, S. Jameel, A. Singh,
708 Measuring glutathione redox potential of HIV-1-infected macrophages. *J. Biol.*
709 *Chem.* **290**, 1020-1038 (2015).
- 710 16. M. P. Look, J. K. Rockstroh, G. S. Rao, K. A. Kreuzer, S. Barton, H. Lemoch,
711 T. Sudhop, J. Hoch, K. Stockinger, U. Spengler, T. Sauerbruch, Serum
712 selenium, plasma glutathione (GSH) and erythrocyte glutathione peroxidase
713 (GSH-Px)-levels in asymptomatic versus symptomatic human
714 immunodeficiency virus-1 (HIV-1)-infection. *Eur. J. Clin. Nutr.* **51**, 266-272
715 (1997).

- 716 17. L. Zhao, A. G. Cox, J. A. Ruzicka, A. A. Bhat, W. Zhang, E. W. Taylor,
717 Molecular modeling and in vitro activity of an HIV-1-encoded glutathione
718 peroxidase. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 6356-6361 (2000).
- 719 18. I. Cohen, P. Boya, L. Zhao, D. Metivier, K. Andreau, J. L. Perfettini, J. G.
720 Weaver, A. Badley, E. W. Taylor, G. Kroemer, Anti-apoptotic activity of the
721 glutathione peroxidase homologue encoded by HIV-1. *Apoptosis* **9**, 181-192
722 (2004); .
- 723 19. S. C. De Rosa, M. D. Zaretsky, J. G. Dubs, M. Roederer, M. Anderson, A.
724 Green, D. Mitra, N. Watanabe, H. Nakamura, I. Tjioe, S. C. Deresinski, W. A.
725 Moore, S. W. Ela, D. Parks, L. A. Herzenberg, L. A. Herzenberg, N-
726 acetylcysteine replenishes glutathione in HIV infection. *Eur. J. Clin. Invest.* **30**,
727 915-929 (2000).
- 728 20. A. Witschi, E. Junker, C. Schranz, R. F. Speck, B. H. Lauterburg,
729 Supplementation of N-acetylcysteine fails to increase glutathione in
730 lymphocytes and plasma of patients with AIDS. *AIDS Res. Hum. Retroviruses*
731 **11**, 141-143 (1995).
- 732 21. C. Sappey, S. Legrand-Poels, M. Best-Belpomme, A. Favier, B. Rentier, J.
733 Piette, Stimulation of glutathione peroxidase activity decreases HIV type 1
734 activation after oxidative stress. *AIDS Res. Hum. Retroviruses* **10**, 1451-1461
735 (1994).
- 736 22. P. A. Sandstrom, J. Murray, T. M. Folks, A. M. Diamond, Antioxidant defenses
737 influence HIV-1 replication and associated cytopathic effects. *Free Radic.*
738 *Biol. Med.* **24**, 1485-1491 (1998).

- 739 23. C. Cao, Y. Leng, W. Huang, X. Liu, D. Kufe, Glutathione peroxidase 1 is
740 regulated by the c-Abl and Arg tyrosine kinases. *J. Biol. Chem.* **278**, 39609-
741 39614 (2003).
- 742 24. T. Wiedenmann, N. Dietrich, T. Fleming, S. Altamura, L. E. Deelman, R. H.
743 Henning, M. U. Muckenthaler, P. P. Nawroth, H. P. Hammes, A. H. Wagner,
744 M. Hecker, Modulation of glutathione peroxidase activity by age-dependent
745 carbonylation in glomeruli of diabetic mice. *J. Diabetes Complications* **32**,
746 130-138 (2018).
- 747 25. W. H. Yang, S. Y. Park, S. Ji, J. G. Kang, J. E. Kim, H. Song, I. Mook-Jung, K.
748 M. Choe, J. W. Cho, O-GlcNAcylation regulates hyperglycemia-induced GPX1
749 activation. *Biochem. Biophys. Res. Commun.* **391**, 756-761 (2010).
- 750 26. S. Ghosh, P. Roy, N. Karmodak, E. D. Jemmis, G. Mugesh, Nanoisozymes:
751 Crystal-Facet-Dependent Enzyme-Mimetic Activity of V₂O₅ Nanomaterials.
752 *Angewandte Chemie* **57**, 4510-4515 (2018).
- 753 27. J. R. Morones, J. L. Elechiguerra, A. Camacho, K. Holt, J. B. Kouri, J. T.
754 Ramirez, M. J. Yacaman, The bactericidal effect of silver nanoparticles.
755 *Nanotechnology* **16**, 2346-2353 (2005).
- 756 28. W. Jiang, B. Y. Kim, J. T. Rutka, W. C. Chan, Nanoparticle-mediated cellular
757 response is size-dependent. *Nat. Nanotechnol.* **3**, 145-150 (2008).
- 758 29. Y. Pan, S. Neuss, A. Leifert, M. Fischler, F. Wen, U. Simon, G. Schmid, W.
759 Brandau, W. Jahnke-Dechent, Size-dependent cytotoxicity of gold
760 nanoparticles. *Small* **3**, 1941-1949 (2007).
- 761 30. M. Auffan, J. Rose, J. Y. Bottero, G. V. Lowry, J. P. Jolivet, M. R. Wiesner,
762 Towards a definition of inorganic nanoparticles from an environmental, health
763 and safety perspective. *Nat. Nanotechnol.* **4**, 634-641 (2009).

- 764 31. X. Zhou, W. Xu, G. Liu, D. Panda, P. Chen, Size-dependent catalytic activity
765 and dynamics of gold nanoparticles at the single-molecule level. *J. Am.*
766 *Chem. Soc.* **132**, 138-146 (2010).
- 767 32. B. Zhou, D. He, Raman spectrum of vanadium pentoxide from
768 density-functional perturbation theory. *J. Raman Spectrosc.* **39**, 1475-1481
769 (2008).
- 770 33. W. Avansi, J. L. Q. Maia, C. Ribeiro, E. R. Leite, V. R. Mastelaro, Local
771 structure study of vanadium pentoxide 1D-nanostructures. *J. Nanopart. Res.*
772 **13**, (2011).
- 773 34. C. Sanchez, J. Livage, G. Lucazeau, Infrared and Raman study of amorphous
774 V₂O₅. *J. Raman Spectrosc.* **12**, 68-72 (1982).
- 775 35. T. M. Folks, J. Justement, A. Kinter, C. A. Dinarello, A. S. Fauci, Cytokine-
776 induced expression of HIV-1 in a chronically infected promonocyte cell line.
777 *Science* **238**, 800-802 (1987).
- 778 36. G. Poli, A. Kinter, J. S. Justement, J. H. Kehrl, P. Bressler, S. Stanley, A. S.
779 Fauci, Tumor necrosis factor alpha functions in an autocrine manner in the
780 induction of human immunodeficiency virus expression. *Proc. Natl. Acad. Sci.*
781 *U.S.A.* **87**, 782-785 (1990).
- 782 37. M. Gutscher, A. L. Pauleau, L. Marty, T. Brach, G. H. Wabnitz, Y. Samstag, A.
783 J. Meyer, T. P. Dick, Real-time imaging of the intracellular glutathione redox
784 potential. *Nat. Methods* **5**, 553-559 (2008).
- 785 38. B. Morgan, M. C. Sobotta, T. P. Dick, Measuring E(GSH) and H₂O₂ with
786 roGFP2-based redox probes. *Free Radic. Biol. Med.* **51**, 1943-1951 (2011).

- 787 39. M. Gutscher, M. C. Sobotta, G. H. Wabnitz, S. Ballikaya, A. J. Meyer, Y.
788 Samstag, T. P. Dick, Proximity-based protein thiol oxidation by H₂O₂-
789 scavenging peroxidases. *J. Biol. Chem.* **284**, 31532-31540 (2009).
- 790 40. R. Brigelius-Flohe, M. Maiorino, Glutathione peroxidases. *Biochim. Biophys.*
791 *Acta* **1830**, 3289-3303 (2013).
- 792 41. R. Drew, J. O. Miners, The effects of buthionine sulfoximine (BSO) on
793 glutathione depletion and xenobiotic biotransformation. *Biochemical*
794 *Pharmacology* **33**, 2989-2994 (1984).
- 795 42. S. Legrand-Poels, D. Vaira, J. Pincemail, A. van de Vorst, J. Piette, Activation
796 of human immunodeficiency virus type 1 by oxidative stress. *AIDS Res. Hum.*
797 *Retroviruses* **6**, 1389-1397 (1990).
- 798 43. C. H. Kim, S. Gollapudi, A. Kim, T. Lee, S. Gupta, Role of protein kinase C-
799 beta isozyme in activation of latent human immunodeficiency virus type 1 in
800 promonocytic U1 cells by phorbol-12-myristate acetate. *AIDS Res. Hum.*
801 *Retroviruses* **12**, 1361-1366 (1996).
- 802 44. R. J. Gulakowski, J. B. McMahon, R. W. Buckheit, Jr., K. R. Gustafson, M. R.
803 Boyd, Antireplicative and anticytopathic activities of prostratin, a non-tumor-
804 promoting phorbol ester, against human immunodeficiency virus (HIV).
805 *Antiviral Res.* **33**, 87-97 (1997).
- 806 45. A. Campa, G. Shor-Posner, F. Indacochea, G. Zhang, H. Lai, D. Asthana, G.
807 B. Scott, M. K. Baum, Mortality risk in selenium-deficient HIV-positive children.
808 *J. Acquir. Immune Defic. Syndr. Hum. Retrovirol.* **20**, 508-513 (1999).
- 809 46. M. P. Look, J. K. Rockstroh, G. S. Rao, K. A. Kreuzer, U. Spengler, T.
810 Sauerbruch, Serum selenium versus lymphocyte subsets and markers of

811 disease progression and inflammatory response in human immunodeficiency
812 virus-1 infection. *Biol. Trace Elem. Res.* **56**, 31-41 (1997).

813 47. A. Gervais, D. West, L. M. Leoni, D. D. Richman, F. Wong-Staal, J. Corbeil, A
814 new reporter cell line to monitor HIV infection and drug susceptibility in vitro.
815 *Proc. Natl. Acad. Sci. U.S.A.* **94**, 4653-4658 (1997).

816 48. P. Tyagi, V. K. Pal, R. Agrawal, S. Singh, S. Srinivasan, A. Singh,
817 *Mycobacterium tuberculosis* Reactivates HIV-1 via Exosome-Mediated
818 Resetting of Cellular Redox Potential and Bioenergetics. *mBio* **11**, (2020).

819 49. E. P. Amaral, E. L. Conceicao, D. L. Costa, M. S. Rocha, J. M. Marinho, M.
820 Cordeiro-Santos, M. R. D'Imperio-Lima, T. Barbosa, A. Sher, B. B. Andrade,
821 N-acetyl-cysteine exhibits potent anti-mycobacterial activity in addition to its
822 known anti-oxidative functions. *BMC Microbiol.* **16**, 251 (2016).

823 50. G. Teskey, R. Cao, H. Islamoglu, A. Medina, C. Prasad, R. Prasad, A.
824 Sathananthan, M. Fraix, S. Subbian, L. Zhong, V. Venketaraman, The
825 Synergistic Effects of the Glutathione Precursor, NAC and First-Line
826 Antibiotics in the Granulomatous Response Against *Mycobacterium*
827 *tuberculosis*. *Front. Immunol.* **9**, 2069 (2018).

828 51. D. Kumar, L. Nath, M. A. Kamal, A. Varshney, A. Jain, S. Singh, K. V. Rao,
829 Genome-wide analysis of the host intracellular network that regulates survival
830 of *Mycobacterium tuberculosis*. *Cell* **140**, 731-743 (2010).

831 52. M. M. Kulkarni, Digital multiplexed gene expression analysis using the
832 NanoString nCounter system. *Curr. Protoc. Mol. Biol.* **Chapter 25**, Unit25B 10
833 (2011).

834 53. R. Eid, E. Boucher, N. Gharib, C. Khoury, N. T. Arab, A. Murray, P. G. Young,
835 C. A. Mandato, M. T. Greenwood, Identification of human ferritin, heavy

- 836 polypeptide 1 (FTH1) and yeast RGI1 (YER067W) as pro-survival sequences
837 that counteract the effects of Bax and copper in *Saccharomyces cerevisiae*.
838 *Exp. Cell. Res.* **342**, 52-61 (2016).
- 839 54. H. Sato, A. Shiiya, M. Kimata, K. Maebara, M. Tamba, Y. Sakakura, N.
840 Makino, F. Sugiyama, K. Yagami, T. Moriguchi, S. Takahashi, S. Bannai,
841 Redox imbalance in cystine/glutamate transporter-deficient mice. *J. Biol.*
842 *Chem.* **280**, 37423-37429 (2005).
- 843 55. C. Espinosa-Diez, V. Miguel, D. Mennerich, T. Kietzmann, P. Sanchez-Perez,
844 S. Cadenas, S. Lamas, Antioxidant responses and cellular adjustments to
845 oxidative stress. *Redox biology* **6**, 183-197 (2015).
- 846 56. K. A. Roebuck, D. S. Gu, M. F. Kagnoff, Activating protein-1 cooperates with
847 phorbol ester activation signals to increase HIV-1 expression. *AIDS* **10**, 819-
848 826 (1996).
- 849 57. A. J. Henderson, K. L. Calame, CCAAT/enhancer binding protein (C/EBP)
850 sites are required for HIV-1 replication in primary macrophages but not
851 CD4(+) T cells. *Proc. Natl. Acad. Sci. U.S.A.* **94**, 8714-8719 (1997).
- 852 58. R. Hu, N. Oyaizu, S. Than, V. S. Kalyanaraman, X. P. Wang, S. Pahwa, HIV-1
853 gp160 induces transforming growth factor-beta production in human PBMC.
854 *Clin. Immunol. Immunopathol.* **80**, 283-289 (1996).
- 855 59. G. Herbein, U. Mählknecht, F. Batliwalla, P. Gregersen, T. Pappas, J. Butler,
856 W. A. O'Brien, E. Verdin, Apoptosis of CD8+ T cells is mediated by
857 macrophages through interaction of HIV gp120 with chemokine receptor
858 CXCR4. *Nature* **395**, 189-194 (1998).
- 859 60. C. Amiel, E. Darcissac, M. J. Truong, J. Dewulf, M. Loyens, Y. Mouton, A.
860 Capron, G. M. Bahr, Interleukin-16 (IL-16) inhibits human immunodeficiency

861 virus replication in cells from infected subjects, and serum IL-16 levels drop
862 with disease progression. *J. Infect. Dis.* **179**, 83-91 (1999)

863 61. W. Choe, D. J. Volsky, M. J. Potash, Induction of rapid and extensive beta-
864 chemokine synthesis in macrophages by human immunodeficiency virus type
865 1 and gp120, independently of their coreceptor phenotype. *J. Virol.* **75**, 10738-
866 10745 (2001)

867 62. S. Z. Khan, N. Hand, S. L. Zeichner, Apoptosis-induced activation of HIV-1 in
868 latently infected cell lines. *Retrovirology* **12**, 42 (2015).

869 63. K. Devadas, N. J. Hardegen, L. M. Wahl, I. K. Hewlett, K. A. Clouse, K. M.
870 Yamada, S. Dhawan, Mechanisms for Macrophage-Mediated HIV-1
871 Induction. *J. Immunol.* **173** (11), 6735-6744 (2004).

872 64. P. Dickie, J. Felser, M. Eckhaus, J. Bryant, J. Silver, N. Marinos, A. L.
873 Notkins, HIV-associated nephropathy in transgenic mice expressing HIV-1
874 genes. *Virology* **185**, 109-119 (1991).

875 65. B. A. Jacob, K. M. Porter, S. C. Elms, P. Y. Cheng, D. P. Jones, R. L. Sutliff,
876 HIV-1-induced pulmonary oxidative and nitrosative stress: exacerbated
877 response to endotoxin administration in HIV-1 transgenic mouse model. *Am.*
878 *J. Physiol. Lung. Cell. Mol. Physiol.* **291**, L811-819 (2006).

879 66. J. B. Kopp, M. E. Klotman, S. H. Adler, L. A. Bruggeman, P. Dickie, N. J.
880 Marinos, M. Eckhaus, J. L. Bryant, A. L. Notkins, P. E. Klotman, Progressive
881 glomerulosclerosis and enhanced renal accumulation of basement membrane
882 components in mice transgenic for human immunodeficiency virus type 1
883 genes. *Proc. Natl. Acad. Sci. U.S.A.* **89**, 1577-1581 (1992).

884 67. S. G. Deeks, R. Tracy, D. C. Douek, Systemic effects of inflammation on
885 health during chronic HIV infection. *Immunity* **39**, 633-645 (2013).

- 886 68. S. Ghosh, Prasad, S. , Mugesh, G., Understanding the role of oxo and
887 peroxido species in the glutathione peroxidase (GPx)-like activity of metal
888 based nanozymes. *Inorganica Chimica Acta* **484**, 283-290 (2019).
- 889 69. V. N. Gladyshev, T. C. Stadtman, D. L. Hatfield, K. T. Jeang, Levels of major
890 selenoproteins in T cells decrease during HIV infection and low molecular
891 mass selenium compounds increase. *Proc. Natl. Acad. Sci. U.S.A.* **96**, 835-
892 839 (1999).
- 893 70. J. Q. Wu, T. R. Sasse, G. Wolkenstein, V. Conceicao, M. M. Saksena, M.
894 Soedjono, S. S. Perera, B. Wang, D. E. Dwyer, N. K. Saksena, Transcriptome
895 analysis of primary monocytes shows global down-regulation of genetic
896 networks in HIV viremic patients versus long-term non-progressors. *Virology*
897 **435**, 308-319 (2013)
- 898 71. S. Y. Lee, Y. K. Park, C. H. Yoon, K. Kim, K. C. Kim, Meta-analysis of gene
899 expression profiles in long-term non-progressors infected with HIV-1. *BMC*
900 *Med. Genomics* **12**, 3 (2019)
- 901 72. L. L. Zhang, Z. N. Zhang, X. Wu, Y. J. Jiang, Y. J. Fu, H. Shang,
902 Transcriptomic meta-analysis identifies gene expression characteristics in
903 various samples of HIV-infected patients with nonprogressive disease. *J.*
904 *Transl. Med.* **15**, 191 (2017).
- 905 73. R. Mishra, S. Kohli, N. Malhotra, P. Bandyopadhyay, M. Mehta, M. Munshi, V.
906 Adiga, V. K. Ahuja, R. K. Shandil, R. S. Rajmani, A. S. N. Seshasayee, A.
907 Singh, Targeting redox heterogeneity to counteract drug tolerance in
908 replicating *Mycobacterium tuberculosis*. *Sci. Transl. Med.* **11**, (2019).
- 909 74. E. P. Amaral, D. L. Costa, S. Namasivayam, N. Riteau, O. Kamenyeva, L.
910 Mittereder, K. D. Mayer-Barber, B. B. Andrade, A. Sher, A major role for

911 ferroptosis in *Mycobacterium tuberculosis*-induced cell death and tissue
 912 necrosis. *J. Exp. Med.* **216**, 556-570 (2019).

913 75. L. Xu, Y. Liu, Z. Chen, W. Li, Y. Liu, L. Wang, Y. Liu, X. Wu, Y. Ji, Y. Zhao, L.
 914 Ma, Y. Shao, C. Chen, Surface-engineered gold nanorods: promising DNA
 915 vaccine adjuvant for HIV-1 treatment. *Nano letters* **12**, 2003-2012 (2012).

916 76. M. C. Bowman, T. E. Ballard, C. J. Ackerson, D. L. Feldheim, D. M. Margolis,
 917 C. Melander, Inhibition of HIV fusion with multivalent gold nanoparticles. *J*
 918 *Am. Chem. Soc.* **130**, 6896-6897 (2008).

919 77. H. H. Lara, N. V. Ayala-Nunez, L. Ixtepan-Turrent, C. Rodriguez-Padilla,
 920 Mode of antiviral action of silver nanoparticles against HIV-1. *J.*
 921 *nanobiotechnology* **8**, 1 (2010).

922 78. H. H. Lara, N. V Ayala-Núñez, L. C. I. Turrent, C. R. Padilla, Bactericidal
 923 effect of silver nanoparticles against multidrug-resistant bacteria. *World*
 924 *Journal of Microbiology and Biotechnology* **26**, 615-621 (2010).

925

926 79. Mandas A, Iorio EL, Congiu MG, Balestrieri C, Mereu A, Cau D, Dessì S,
 927 Curreli N. Oxidative imbalance in HIV-1 infected patients treated with
 928 antiretroviral therapy. *J Biomed Biotechnol.* (2009).

929 80. M. M. Elsheikh, Y. Tang, D. Li, G. Jiang, Deep latency: A new insight into a
 930 functional HIV cure. *EBioMedicine* **45**, 624-629 (2019).

931 81. J. Mendiàdua, R. Casanova, Y. Barbaux, XPS studies of V₂O₅, V₆O₁₃, VO₂
 932 and V₂O₃. *J. Electron Spectrosc. Relat. Phenom.* **71**, 249-261 (1995).

933 82. E. Hryha, E. Rutqvist, L. Nyborg, Stoichiometric vanadium oxides studied by
 934 XPS. *Surf. Interface Anal.* **44** 1022-1025 (2012).

935

Acknowledgements

This work was supported by Wellcome Trust-Department of Biotechnology (DBT) India Alliance grant IA/S/16/2/502700 (A.S.) and in part by DBT grants BT/PR11911/BRB/10/1327/2014, BT/PR13522/COE/34/27/2015 and BT/HRD/NBA/39/07/2018-19 (A.S.), DBT-IISc Partnership Program grant 22-0905-0006-05-987 436, and the Infosys Foundation. A.S. is a senior fellow of Wellcome Trust-DBT India Alliance. G. M. acknowledges the DST Nano Mission (SR/NM/NS-1380/2014) and SERB (SB/S2/JCB-067/2015), DST, New Delhi for funding. SS and SG acknowledge fellowships from the University Grants Commission (UGC) and Indian Institute of Science (IISc) respectively. We thank Dr. Amit A. Vernekar for helpful discussions. We gratefully acknowledge the NanoString services provided by TheraCUES Innovations Pvt Ltd, Bangalore. We also thank the AFMM Facility, CeNSE, IISc for the microscopic and spectroscopic facilities.

Figures

Figure 1




Figure 1. Synthesis and characterization of vanadia (V_2O_5) nanoparticles. (A) Schematic representation of the synthesis of thin V_2O_5 nanosheets (Vs) from VSh. **(B)** SEM image of thin-nanosheets (Vs), scale – 20 μm . **(C)** Schematic representing

the glutathione reductase (GR)-coupled assay to measure the GPX-like activity of Vs. **(D)** Comparison of initial activity rate among all three forms of V_2O_5 nanomaterials having a common exposed facet [001]. *** $P < 0.001$, **** $P < 0.0001$ by Student's t test. Data is representative of three independent experiment (mean $\pm$ SD). **(E)** Bar diagram of the initial rate of Vs with 3 different peroxides, t -BuOOH – tertiary-butyl hydroperoxide, Cum-OOH – cumene hydroperoxide and H_2O_2 . **(F)** High-resolution TEM (HRTEM) and fast Fourier transform (FFT) (inset) of Vs showing the lattice fringes and the exposed plane. **(G)** FT-Raman spectroscopy of Vs showing the peaks corresponding to the orthorhombic phase of the material. **(H)** Energy dispersive spectroscopy (EDS) of Vs. The small peak at 2.0 KeV is due to Au spurting while recording the spectra (Scale – 50 μ m). The peak of C is coming from atmospheric carbon. X-Ray mapping images of Vs (Scale – 300 nm). **(I)** Left column: Selective area bright field (SABF) image, **(J)** middle column: distribution of vanadium (V) atoms in red, **(K)** right column: distribution of oxygen (O_2) atoms in green.

Figure 2




Figure 2. Enzymatic parameters and surface area measurements of Vs. (A) Bar diagram shows the initial rate of GPX-like activity of Vs under different assay conditions. **(B)** Dependence of the initial reaction rate for H_2O_2 reduction on varying concentration of the catalyst, Vs. **(C)** Michaelis–Menten plot with the variation of H_2O_2 (0–400 μM) in the presence of Vs (20 $\text{ng}/\mu\text{L}$), GSH (2 mM), NADPH (0.2 mM), GR (1.7 units) in phosphate buffer (100 mM, pH 7.4) at 25°C. **(D)** Michaelis–Menten plot with the variation of GSH (0–7 mM) in the presence of Vs (20 $\text{ng}/\mu\text{L}$), H_2O_2 (200 μM), NADPH (0.2 mM), GR (1.7 units) in phosphate buffer (100 mM, pH 7.4) at 25°C. **(E)** and **(F)** Lineweaver-Burk plot with varying concentration of H_2O_2 and GSH in presence of Vs nanozyme respectively. The concentration of NADPH was

constant (0.2 mM) in all the assay conditions. **(G)** Surface area measurement by N₂ adsorption, desorption isotherm and distribution of pore size (Inset). The measured surface area of Vs was 13.3 m²/g. **(H)** Enzyme kinetic parameters and BET surface area values of different forms of V₂O₅ nanomaterials [#] reported from our previous literature (26).

Figure 3




Figure 3. Vs acts as a mimic of GPX in U1 cells. (A) U1 cells were treated with 50 ng/μL of Vs for 15 min (grey area), washed, and kinetics of Vs retention (yellow area) was quantified by measuring cellular and extra-cellular Vs content using ICP-MS. (B) Schematic representation showing working principle of Orp1-roGFP2 biosensor. (C) U1-Orp1-roGFP2 cells were treated with Vs for 15 min, followed by exposure to H₂O₂ for 2 min, and ratiometric response was measured. (D) Schematic representation showing working principle of Grx1-roGFP2 biosensor. (E) U1-Grx1-roGFP2 cells were pre-treated with Vs for 15 min, exposed to H₂O₂ for 2 min, and ratiometric response was measured. (F) U1-Grx1-roGFP2 cells were treated with two

1005 doses of 50 μ M H₂O₂ (indicated by the arrows) and the ratiometric response was
1006 measured (blue line). Parallely, U1 cells treated similarly with H₂O₂ were exposed to
1007 Vs at the indicated time point and the ratiometric response was measured (green
1008 line). ****P<0.0001, by Wilcoxon matched-pairs signed rank test. **(G)** and **(H)** U1
1009 Grx1-roGFP2 cells were supplemented with BSO or GSH for 16 h to deplete or
1010 replenish GSH, respectively. Following this, cells were treated with Vs for 15 min,
1011 exposed to H₂O₂, and the ratiometric response was measured. **** P<0.0001, ***
1012 P<0.001, ** P<0.01, * P<0.05, by Mann Whitney Test. Asterisks (*) compare Vs
1013 treated cells with control cells (C). Data are representative of results from three
1014 independent experiments performed in triplicate (mean $\pm$ SEM).
1015
1016

Figure 4





Figure 4. Vs subverts HIV reactivation. (A) Vs-treated U1 cells were challenged with 5 ng/mL PMA or 1.25 μM prostratin (Pros) for 24 h and HIV-1 induction was monitored by *gag* RT-PCR. 10 mM NAC, an antioxidant known to subvert PMA mediated viral reactivation, was used as a positive control. (B) Vs-treated U1 cells were exposed to PMA and viral activation was measured as a function of time, by *gag* RT-PCR. U1 cells were also treated with Vs or PMA alone. (C) Untreated or Vs-treated U1-Orp1-roGFP2 cells were exposed to PMA and the biosensor response was measured at the indicated time points. The biosensor response was also measured for untreated or PMA treated cells. (D) J1.1 cells were treated twice with Vs for 15 min at 0 and 24 h time point. HIV-1 induction was measured by *gag* RT-PCR at 24 h and 48 h post-treatment. An untreated control was used for normalization. (E) U1-Grx1-roGFP2 cells were serum starved for 30 min in the

1033 presence or absence of Vs and sodium selenite (0.5 nM), and the biosensor
 1034 response was measured. Data were compared to serum starved control cells (C). **(F)**
 1035 U1 cells were either serum-starved or supplemented with Se (0.5 nM) or Vs (0.62
 1036 ng/μL) and HIV reactivation was measured at 6 h post starvation by *gag* RT-PCR.
 1037 **P<0.01, \$/##*P<0.05, by Mann Whitney Test. Data are representative of results
 1038 from three independent experiments performed in triplicate (mean ± SD).

1039

1040

1041 **Figure 5**



1042

1043 **Figure 5. Vs reduces replication of HIV-1 and *Mtb*.** (A) CEM-GFP cells were pre-

1044 treated with 50 ng/ μ L of Vs for 15 min and infected with 0.1 moi of T-tropic HIV-1

1045 provirus (pNL4.3) and GFP fluorescence was measured by at 488 nm as an indicator

1046 of HIV LTR activity. Vs treatment was repeated every 24 h for the experiment. A


1047 similar assay was performed using Jurkat (CD4⁺ T cell line) and viral replication was

1048 assessed by (B) *gag* RT PCR, (C) p24 ELISA in the culture supernatant, and (D)

1049 immunoblotting for p24 (viral capsid protein) in the whole cell lysate. (E) U937

(promonocytes) were pre-treated with 50 ng/μL of Vs for 15 min followed by infection with 1 moi of M-tropic HIV pro-virus (pNLAD8) and viral activation was measured by *gag* qRT-PCR at 24 h post-infection (hpi). **(F)** Primary CD4⁺ T cells purified from human PBMCs were activated, pre-treated with 25 ng/μL Vs for 15 min, and infected with 0.1 moi of T-tropic HIV-1 provirus (pNL4.3). Virus released in supernatant was quantified by p24 ELISA. Vs treatment was repeated every 48 h. Data analysed by Student's *t* test (mean ± SD). **(G and H)** U1 macrophages pre-treated with 50 ng/μL Vs were infected with drug-sensitive *Mtb* H37Rv or multidrug-resistant JAL2287 strains and bacterial survival was evaluated by enumerating colony forming units (CFUs). Percent survival was calculated by normalizing with CFUs derived from *Mtb* infected Vs-untreated U1 at 4 hpi. All figures except **(A)** and **(C)** were analysed by Mann Whitney test. **(A)** and **(C)** were analysed by 2-way ANOVA with Tukeys multiple comparison test. **** P<0.0001, *** P<0.001, * P<0.05. Data are representative of results from three independent experiments performed in triplicate (mean ± SD).

1075 **Figure 6**




1076

1077

1078 **Figure 6. Expression analysis of host pathways associated with HIV-1**
1079 **activation.** U1 cells were treated with Vs (50 ng/μL) for 15 min, followed by PMA
1080 treatment for 12 h. Total RNA was isolated from untreated (UT), PMA-treated, Vs-
1081 treated and Vs+PMA treated U1 cells and expression analysis of 185 genes specific
1082 to oxidative stress and HIV host response was measured using the NanoString
1083 technology. **(A)** Number of differentially expressed genes (DEGs) under the three

comparison conditions i. PMA/UT, ii. Vs/UT and iii. Vs+PMA/UT. **(B)** The venn diagram showing overlap of DEGs significantly perturbed under different comparison conditions. **(C and D)** Heat map showing multiple categories of DEGs under PMA/UT, Vs/UT and PMA+Vs/UT comparisons. mRNA counts were normalised using the internal control β_2 microglobulin (B2M), and fold change (FC) was calculated using the nSolver 4.0 software. Genes showing an absolute FC >1.5, and P <0.05 were considered as significantly altered. **(E and F)** Vs-treated or untreated U1 cells were exposed to Bay11-7082 (7.5 μ M). PMA mediated HIV reactivation at 12 h was monitored by *gag* RT PCR and p24 ELISA. **P<0.01, *P<0.05, by Mann Whitney Test. Data are representative of results from three independent experiments performed in triplicate (mean $\pm$ SEM). Asterisks (*) compare different treatment conditions with PMA-treated cells.

1101 **Figure 7**



1102

Figure 7. Vs reduces oxidative stress, edema, and HIV expression *in vivo*. (A)

Strategy to investigate the antioxidant function of Vs in HIV Transgenic mice, Tg26.

HIV Tg26 (n=8), were left untreated or treated with 1 mg/kg Vs for 3 h. Untreated WT

C57BL/6 was used as a control. **(B)** Lung homogenates were stained with DCFDA

and analysed by flow cytometry for ROS detection. Singlets, followed by different cell

clusters comprising mainly of granulocytes (P2), monocytes (P3) and lymphocytes

(P4) were gated from the whole lung homogenate and ROS levels were plotted as

median fluorescence intensity (MFI). **(C)** Pan-macrophage surface markers, CD64

and MERTK, were used to stain total lung macrophage population, ROS levels were

quantified by DCFDA staining and plotted as MFI.**(D)** RNA was isolated from the

right lung lobe and expression of GPX1 in Tg UT and Tg Vs animals was checked by

qRT-PCR. GPX1 expression values were normalised with Tg UT. **(E)** Lung edema

plotted as the ratio of wet versus the dry weight of left lobe. **(F)** Expression of *tat-rev*

region as a marker for HIV-1 transcription was evaluated by qRT-PCR using RNA

from lungs of Tg UT and Tg Vs animals. Data has been plotted as median $\pm$

interquartile range. *** P<0.001, ** P<0.01, * P<0.05, by Mann Whitney Test.

1128 **Supplementary Material**

1129

1130 **List of Supplementary Materials**

1131 Fig. S1. SEM and initial reaction rate for Vanadia Nanowires and crude nanosheets.

1132 Fig. S2. XRD pattern and TEM of Vs

1133 Fig. S3. X-Ray Photoelectron Spectroscopy (XPS) of Vs.

1134 Fig. S4. GR- coupled assay for Vs under varied conditions.

1135 Fig. S5. *In vitro* recycling ability of Vs.

1136 Fig. S6. Survival assay for Vs-treated U1 cells.

1137 Fig. S7. Preparation of Stable cell lines.

1138 Fig. S8. Time dependent induction of Selenium (Se) starvation-mediated oxidative
1139 stress.

1140 Fig. S9. Antioxidant gene expression in lungs of Vs treated Tg26.

1141 Table S1. List of differentially expressed genes from U1 cells either left Untreated
1142 (UT), or treated with PMA, Vs or Vs+PMA.

1143 Table S2. List of primers used in the study.

1144 Supplementary Experimental procedures

1145

1146

1147

1148

1149

Supplementary Figures

Figure S1




Figure. S1 SEM and initial reaction rate for Vanadia Nanowires and crude nanosheets. Scanning electron microscopy (SEM) images of **(A)** nanowires (VNw; scale – 5 μm) and **(B)** crude nanosheets (VSh; scale – 200nm). **(C)** Bar graph shows the initial rate of activity for VNw and VSh. * P<0.05 by Student's *t* test. Data is representative of three independent experiment (mean ± SD).

Figure S2




Figure S2. XRD pattern and TEM of Vs **(A)** Powder XRD of Vs where all the diffraction peaks were indexed to standard V₂O₅ orthorhombic phase (JCPDS = 41-

1426, space group Pmmn). **(B)** Transmission electron microscopy (TEM) image of Vs thin nanosheets and selected area electron diffraction (SAED) (inset) shows the crystalline nature of the material.

Figure S3




Figure S3. X-Ray Photoelectron Spectroscopy (XPS) of Vs. **(A)** Left: wide spectra and **(B)** right: deconvoluted spectra of oxygen and vanadium peaks. The spectra were calibrated by taking C1s (284.6 eV) as a standard. As described by *Mendialdua et al* (81, 82) the difference in binding energies between the O1s core level and the V2p3/2 level ($\Delta = \text{BE (O1s)} - \text{BE (V2p3/2)}$) was used to determine the oxidation state of V₂O₅ nanozymes which confirmed +5 oxidation states of vanadium in Vs. **(C)** Full width half maxima (FWHM) & binding energies of deconvoluted oxygen and vanadium XPS peaks.

Figure S4




Figure S4. GR- coupled assay for Vs under varied conditions. (A) The decrease in absorbance of NADPH (0.2 mM) under different assay conditions were monitored during GR coupled assay.

Figure S5




Figure S5. *In vitro* recycling ability of Vs. (A) Recycling activity of Vs during multiple rounds of catalysis was analysed by addition of fresh substrates in the

reaction mixture up to six cycles. The curves observed due to reduction in NADPH absorbance, for every cycle were parallel to each other, indicating no change in their initial rate and persistence of the activity. This clearly depicts robustness of the catalyst. Conditions used for the assay was sodium phosphate buffer (100 mM, pH 7.4), GSH (2 mM), NADPH (0.2 mM), catalyst (20 ng/ μ L), GR (~1.7 U) and H₂O₂ (20 μ M) at 25°C. High Resolution TEM (HRTEM) image and FFT patterns of Vs before **(B)** and after **(C)** catalysis. **(D)** SAED pattern of Vs after catalysis. The pattern was indexed in a common zone axis [001] which indicates the surface exposed facets are retained after multiple rounds of catalysis.

Figure S6




Figure S6. Survival assay for Vs-treated U1 cells. (A) U1 cells were treated with increasing concentrations of Vs – 10 to 100 ng/ μ L for 15 min and cell survival was analysed flow cytometrically after 24 h by propidium iodide (PI) staining. Experiment is representative of two independent experiments done in triplicate (mean $\pm$ SD). *P<0.05, by Mann Whitney test.

Figure S7




Figure S7. Preparation of Stable cell lines. Dot plot validating the stable expression of the biosensors in (A) U1 cyto-Orp1-roGFP2 and (B) U1 cyto-Grx1-roGFP2 cell lines.

Figure S8




Figure S8. Time dependent induction of Selenium (Se) starvation-mediated oxidative stress. (A) U1-Grx1-roGFP2 cells were cultured in serum free medium (to deplete Se) for various time periods and biosensor response was measured by flow cytometry. N=2. Data is representative of two experiments done in duplicates (mean $\pm$ SEM). * P<0.05, by Mann Whitney test.

Figure S9




Figure S9. Antioxidant gene expression in lungs of Vs treated Tg26. (A) Tg26 mice were left untreated or treated with 1 mg/kg of Vs for 3 h. RNA was isolated from the right lung lobe, reverse transcribed and expression levels of GPX4, SOD1 and SOD2 in Tg UT and Tg Vs animals were analysed. Values were normalised with Tg UT. n=6, ns=non-significant.

Supplementary Table

Table S2

Primers for qRT-PCR			
Organism	Gene		Primer Sequence
HIV-1	p24	Forward	5'-ATAATCCACCTATCCCAGTAGGAGAAAT-3'
		Reverse	5'-TTGGTTCCCTTGCTTATGTCCAGAATGC-3'
	tat-rev	Forward	5'- TGAAGCATCCAGGAAGTCAGCC-3'
		Reverse	5'-TTCTTCTTCTATTCTTCCTTCGGGCC-3'
<i>Homo sapiens</i>	β -Actin	Forward	5'-ATGTGGCCGAGGACTTTGATT-3'
		Reverse	5'-AGTGGGGTGGCTTTTAGGATG-3'
<i>Mus musculus</i>	GPX1	Forward	5'-GGTTCGAGCCCAATTTTACA-3'
		Reverse	5'-CCCACCAGGAACCTTCTCAAA-3'
	GPX4	Forward	5'-CTCCATGCACGAATTCTCAG-3'
		Reverse	5'-ACGTCAGTTTTGCCTCATTG-3'
	SOD1	Forward	5'-CAGAAGGCAAGCGGTGAA-3'
		Reverse	5'-CAGCCTTGTGTATTGTCCCCATA-3'
	SOD2	Forward	5'-GACCTGCCTTACGACTATGG-3'
		Reverse	5'-GACCTTGCTCCTTATTGAAGC-3'
	β -ACTIN	Forward	5'-GCAAGCAGGAGTACGATGAG-3'
		Reverse	5'-CCATGCCAATGTTGTCTCTT-3'
Primers for Genotyping			
	Tq26 F18296	Forward	5'-TCCAGTTTGGAAGGACCAG-3'
	Tq26 F18297	Reverse	5'-TTGCCACACAATCATCACCT-3'
	Int Ct F21157	Forward	5'-CTCCCAACCCCAGAGGTAGT-3'
	Int Ct F21225	Reverse	5'-AGACCCCAGATCCAGAAAGG-3'

Table S2 List of primers used in the study.

Supplementary Materials and Methods

Dispersion of V₂O₅ NPs

V₂O₅ NPs were dispersed in sterile water at a concentration of 2 mg/mL. The dispersion was carried out by sonication using probe sonicator under the following conditions: time – 5 min, amplitude – 5 sec ON, 5 sec OFF. After dispersion, the vanadia NPs form a yellowish colloidal solution.

Treatment of cell lines with Vs

Indicated cell lines were treated with different concentrations of Vs at a cell density of 0.2×10^6 cells/mL for 15 min at 37°C in a CO₂ incubator. Following Vs treatment, the cells were washed and re-suspended in complete medium for further culturing or in FACS buffer (1× PBS + 3% FBS) for measuring the antioxidant response by flow cytometry, as required. For experiments with J1.1, Jurkat, or CEM-GFP cell lines, which required long term culturing, Vs treatment was repeated every 24 h, as mentioned above.

Internalization of Vs by U1

U1 cells were treated with 50 ng/μL as mentioned above. The cell pellet was re-suspended in 10 mL of dilute nitric acid (HNO₃; 3 N) and incubated for 16 h to remove cellular organic matter. The samples are analysed by ICP-AES (inductively coupled plasma atomic emission spectroscopy) and their vanadium (V) content was estimated in parts per billion (ppb). The ppb content of vanadium sulphate treated parallelly with HNO₃ was used as a standard to estimate the cellular V content.

Mammalian and bacterial Cell Culture

The human monocytic cell line U937, CD4⁺ T lymphocytic cell line Jurkat (ATCC, Manassas, VA), the chronically infected U1 and J1.1, and CD4⁺ reporter T cell line, CEM-GFP (AIDS Research and Reference Reagent program, NIH, USA) were grown in RPMI 1640 (Cell Clone), with 10% FBS (MP Biomedicals) and 2 mM L-glutamine (MP Biomedicals) supplementation. Vs treatment, transfection, and HIV-1 infection was carried out in Opti MEM media (Hyclone). Differentiation and HIV activation in U1 cells were carried out by treatment with 5 ng/mL of phorbol ester PMA (Sigma) or 1.25 μM prostratin (Sigma). Laboratory strain of *Mtb* H37Rv and

clinical isolate JAL2287 (MDR) were cultured in 7H9 (liquid) or 7H11 (solid) synthetic medium supplemented with 0.4% glycerol, 0.1% tween-80 and 10% albumin-dextrose-sodium chloride (ADS) or 0.4% glycerol and 10% oleic acid-albumin-dextrose-catalase (OADC), respectively.

Preparation of Stable Cell lines and validation using flow cytometry

Various cell lines stably expressing the cytosolic biosensor Grx1-roGFP2 or Orp1-roGFP2 were prepared by electroporation of 10×10^6 U1 cells with 5 μ g of the pMSCVpuro-Grx1-roGFP2 or pMSCVpuro-Orp1-roGFP2 constructs, followed by selection on 350 ng/mL puromycin. The ratiometric responses of the biosensors were measured by excitation at 405 and 488 nm, and recording emission at 510/10 nm, using BD FACSVerse flow cytometer (BD Biosciences). The data were analysed using FACSuite software (BD Biosciences).

Assessment of Vs antioxidant activity and redox potential measurement

0.1 $\times 10^6$ untreated and Vs treated U1-Orp1-roGFP2 or U1-Grx1-roGFP2 cells were exposed to increasing concentrations of H_2O_2 - 50 and 100 μ M or 50, 100 and 200 μ M, respectively- and incubated at RT for 2-3 min. These cells were analysed by flow cytometry at excitation of 405 nm (V500) and 488 nm (FITC), while the emission was fixed at 510 nm. Ratio of fluorescence intensities at 405/488 was calculated and normalized using a cell permeable oxidant H_2O_2 or the reductant DTT to calculate the responsiveness of both the biosensors.

Intracellular redox potential was measured for cells expressing the Grx1-roGFP2 biosensor, as mentioned earlier (15). Briefly, for each experiment, 100% biosensor oxidation or reduction corresponding to maximal and minimal fluorescence

intensity ratios was determined by treatment with 10 mM H₂O₂ and 10 mM DTT, respectively. The observed ratios were used to determine the degree of biosensor oxidation and ultimately equated in a modified form of the Nernst equation to obtain the intracellular glutathione redox potential (E_{GSH}).

Propidium iodide (PI) staining

U1 cells were treated with increasing concentrations of Vs for 15 min and cultured in complete RPMI medium for 24 h. After 24 h, cells were washed, suspended in 1× PBS, and stained with 3 μM PI for 15 min in the dark. After washing twice with 1× PBS, cells were analyzed on a flow cytometer using the phycoerythrin (PE) detector (575/26 nm) by excitation at 488 nm. emission.

HIV reactivation in U1 cells and qRT-PCR analysis

U1 cells were treated with either 5 ng/mL PMA or 1.25 μM prostratin and incubated at 37°C in a CO₂ incubator. Samples were harvested at 6 h, 12 h, and 24 h post-activation, and RNA was isolated using the Qiagen RNAeasy kit (manufacturer's protocol). cDNA was synthesized using 400 ng RNA by the Biorad iScript cDNA synthesis kit. qRT-PCR was performed using primers against *gag* transcript (a marker for HIV reactivation). Actin was used as an internal control. To inhibit NF-κB pathway, untreated or Vs-treated cells were exposed to 7.5 μM Bay11-7082 (TCI chemicals) for 12 h.

p24 detection by Immunoblotting and ELISA

Untreated or Vs treated HIV-infected Jurkat cells were harvested at 36, 48, and 72 h post infection. Cells were lysed in 300 µL of passive lysis buffer (Promega) supplemented with 1×protease inhibitor cocktail (Roche). Protein was quantified using Pierce™ BCA Protein Assay Kit (Thermo Fisher Scientific). 50 µg of whole cell lysate (WCL) was mixed with Laemmli buffer, heated at 95°C for 5 min, and was separated on a 12% SDS PAGE gel. Immunoblotting was performed using primary antibodies against HIV-p24 (Abcam; ab9071) and GAPDH (CST; D4C6R) as an internal control. Horse anti-mouse IgG (CST; 7076) was used as the secondary antibody. For ELISA, we collected supernatant from Jurkat cells infected with HIV at 24, 48, and 72 hpi. p24 levels were determined by sandwich ELISA using the J.J. Mitra's kit as per the manufacturers' instructions. Standard curve was prepared with known amount p24 and utilised for calculating the viral p24 concentration in the medium.

Bacterial Survival assays in U1 cells

For survival experiments of *Mtb* H37Rv and JAL2287 strains in U1 cell lines, the cells were differentiated with 5 ng/mL of PMA for 18 h. Differentiated U1 macrophages were either left untreated or treated with 50 ng/µL of Vs for 15 min, washed, and allowed to rest for 2 h. H37Rv and JAL2287 were opsonized in 50% horse serum for 1 h. Untreated or Vs treated U1 cells were infected at moi 2 for 4 h. After 4 h of incubation, extracellular bacilli were killed by treatment with 0.2 mg/mL amikacin for 1 h. The infected cells were washed three times with 1× PBS and then cultured in complete RPMI media for 24 h. The infected cells were lysed in 0.06% sodium dodecyl sulphate (SDS) and plated on 7H11 plates supplemented with 10% OADC. Bacterial survival was assessed 24 h post infection by enumerating colony forming units (CFUs) on agar media. The initial bacterial burden was determined by

plating at 0 h post infection and the percentage increase in bacterial survival was plotted.

Isolation and infection of primary CD4⁺ T cells

Peripheral blood mononuclear cells (PBMCs) were isolated using Ficoll-Paque-based density gradient centrifugation from blood samples of healthy donors, donated after informed consent, approved by the IISc Ethical Committee. Primary CD4⁺ T cells were purified from PBMCs using an EasySep human CD4⁺ T cell isolation kit (Stem Cell Technologies, Canada) and cultured for 3 days after isolation at 37°C in 5% CO₂ in complete media containing RPMI 1640 supplemented with 10% FBS, 100 U/mL interleukin-2 (IL-2) (Peprotech, London, United Kingdom) (specific activity, 10 U/ng), and 1 µg/mL phytohemagglutinin (PHA) (Thermo Fisher Scientific). Subsequently, 250,000 activated primary CD4⁺ T cells, were pre-treated with 25 ng/µL Vs for 15 min and infected with 0.1 moi pNL4.3 virus by spinoculation at 1000 g for 90 min at 32°C. Cells were then washed and replenished with complete media containing 100 U/mL IL-2. Vs treatment was repeated every 48 h. To quantify the virion release, supernatant was harvested from infected cells and centrifuged at 400 g for 10 min and virus concentration was estimated by HIV-1 p24 ELISA.

Animal experiments

Heterozygous HIV transgenic mouse Tg26 containing HIV proviral DNA with mutated 3 kb region of *gag* and *pol* genes was used (64). In order to create a more stable line, the HIV Tg26 (FVB/N) was backcrossed with C57BL/6 for eight generations to create the current Tg26 by Dr. Roy L. Sutliff (Veterans Affairs Medical Center, Emory University, Atlanta, GA). HIV Tg26 breeding colony was maintained in the BSL3 animal facility at CIDR, IISc, according to the guidelines set by Institutional

Animal Ethical Committee. Genotyping was done at 4 weeks of age by tail vein PCR. 6-9 weeks old sex-matched transgenic mice were used for the experiment. Wild type (Wt) littermates were used as controls. 1 mg/kg body weight Vs was administered by intra-peritoneal injection in Tg26 treatment group. Control and treated animals were euthanized at 3 h post Vs treatment.

Pulmonary edema analysis

Left lobe of the lungs from 8 mice in each group- untreated C57BL/6 Tg26 and Vs treated Tg26 were removed aseptically and weighed in 1.5 mL microfuge tubes. The lungs were left for 16-24 h in a dry oven at 55°C for desiccation and the dry weight of the tissue was recorded. The ratio of lung wet weight to dry weight was plotted for each group.

ROS staining of lung homogenates

Untreated WT, untreated Tg26, and Vs treated Tg26 mice were euthanized at 3 h post Vs treatment and a portion of the right lobe of the lung was aseptically removed. Single cell suspensions for lung homogenates were prepared for DCFDA staining and flow cytometry analysis as per an earlier study (73). Briefly, the tissue was minced and digested in serum free RPMI containing 0.2 mg/mL Liberase DL and 0.1 mg/mL DNase I for 60 min at 37°C and agitation at 180 rpm. The minced tissue was mechanically disrupted using GentleMACS tissue dissociator to obtain a finer suspension. Larger cell clumps were removed by passing the suspension through a 40 µm cell strainer (BD Falcon). Red blood cells were removed by incubating the cells in RBC lysis buffer.

qRT-PCR analysis from animal tissues

RNA was isolated from a part of the right lobe of the lungs using Trizol reagent (Ambion). 1 µg RNA was reverse transcribed using the Biorad iScript cDNA synthesis kit. qRT-PCR was performed using primers against GPX1, GPX4, CAT, SOD1, and SOD2 to assess the host antioxidant response under different treatment groups. β-Actin was used as an internal control.