

Chicken cGAS senses fowlpox virus infection and regulates macrophage effector functions

Marisa Oliveira¹, Damaris Ribeiro Rodrigues², Vanaique Guillory³, Emmanuel Kut³, Efstathios S. Giotis^{4,5}, Michael A. Skinner⁴, Rodrigo Guabiraba^{3†}, Clare E Bryant^{2†}, Brian J Ferguson^{1†*}

¹Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge, UK

² Department of Veterinary Medicine, University of Cambridge, Cambridge, UK

³ ISP, INRAE, Université de Tours, 37380, Nouzilly, France.

⁴ Department of Infectious Diseases, Imperial College London, London, United Kingdom

⁵ School of Life Sciences, University of Essex, Colchester, United Kingdom

[†]Co-senior authorship

* Correspondence:

Brian J Ferguson

bf234@cam.ac.uk

Keywords: DNA, Fowlpox, chicken, macrophages, cGAS, STING, PRR

Abstract

The anti-viral immune response is dependent on the ability of infected cells to sense foreign nucleic acids. In multiple species, the pattern recognition receptor (PRR) cyclic GMP-AMP synthase (cGAS) senses viral DNA as an essential component of the innate response. cGAS initiates a range of signalling outputs that are dependent on generation of the second messenger cGAMP that binds to the adaptor protein stimulator of interferon genes (STING). Here we show that in chicken macrophages, the cGAS/STING pathway is essential not only for the production of type-I interferons in response to intracellular DNA stimulation, but also for regulation of macrophage effector functions including the expression of MHC-II and co-stimulatory molecules. In the context of fowlpox, an avian DNA virus infection, the cGAS/STING pathway was found to be responsible for type-I interferon production and MHC-II transcription. The sensing of fowlpox virus DNA is therefore essential for mounting an anti-viral response in chicken cells and for regulation of a specific set of macrophage effector functions.

Introduction

The ability of virally infected cells to mount an effective innate immune response is dependent on the intracellular sensing of nucleic acids by pattern recognition receptors (Mansur et al., 2014). The PRRs that sense and respond to intracellular DNA are well characterised in a number of mammalian and non-mammalian organisms but are less studied in avian species, including chickens (Bryant et al., 2015). The PRR cyclic cAMP-GMP (cGAMP) synthase (cGAS) binds intracellular viral DNA and, via production of the second-messenger 2'3'-cGAMP, triggers a range of signalling outputs including type-I interferon (IFN-I) production, cell death and cellular senescence (Li and Chen, 2018). The absence of cGAS or the adaptor protein, stimulator of interferon genes (STING), which

binds cGAMP, results in the susceptibility to DNA virus infection in knockout mice and impairs IFN-I production by cells infected with DNA viruses or transfected with linear double stranded DNA (Li et al., 2013). Through its ability to sense mislocalised self-DNA, the cGAS/STING signalling axis is also a potent regulator of autoinflammatory and anti-tumour immune responses (Ablasser et al., 2013a; Mullard, 2017). People with activating mutations in STING or loss-of function mutations in the 5'-3' exonuclease TREX, which removes excess cytoplasmic dsDNA, suffer from interferonopathies (Crow and Rehwinkel, 2009).

The ability of cGAS/STING signalling to drive multiple downstream signalling outputs is dependent on the activation of a number of distinct signalling mechanisms, some of which are better defined than others. The production of IFN-I in this context is dependent on STING recruiting and facilitating activation of TANK-binding kinase-1 (TBK1) and the transcription factor interferon regulatory factor-3 (IRF3) (Tanaka and Chen, 2012). IRF3 phosphorylation, dimerisation and translocation to the nucleus results in IFN-I transcription. The mechanism or mechanisms by which STING can promote cell death are less well described, but include inflammasome activation (Gaidt et al., 2017) and apoptosis of various cell types including myeloid and T cells (Gulen et al., 2017; Sze et al., 2013). cGAS can also activate a programme of cellular senescence in fibroblasts by sensing damaged self-DNA (Glück et al., 2017). It is not currently clear in what contexts these disparate signalling outputs are activated by cGAS/STING and to what extent they cross-talk with each other.

Chickens are economically important livestock birds that are infected by numerous viruses including fowlpox virus (FWPV). Fowlpox is a virus from the *poxviridae* family that replicates its double stranded DNA genome in the cytoplasm of infected cells. The infection is characterised by proliferative lesions in the skin that progress to thick scabs (cutaneous form) and by lesions in the upper GI and respiratory tracts (diphtheritic form) (Giotis and Skinner, 2019). Transmitted mechanically by biting insects, it causes significant losses to all forms of poultry production systems (from backyard, through extensive to intensive commercial flocks). It is particularly challenging in tropical climates where control of biting insects is difficult. FWPV is also used as a live recombinant vaccine vector in avian and mammalian species (Lousberg et al., 2010). Like other poxviruses the cytoplasmic replication cycle of FWPV exposes large amounts of foreign DNA to intracellular DNA sensing PRRs, making cGAS a likely candidate for sensing FWPV infection and making FWPV a potentially useful tool for delineating nucleic acid sensing mechanisms in avian systems. The mechanisms by which FWPV is sensed by PRRs during infection have not, however, been described.

In this study we show the existence of a cGAS/STING pathway in chicken macrophages and determine its downstream signalling outputs. Using cGAS and STING CRISPR/Cas9 knockout HD11 cells and pharmacological inhibitors of STING and TBK1 in primary macrophages, we show that the activation of cGAS by intracellular DNA drives a IFN-I response and that this response can be enhanced by priming cells with IFN α . As well as driving IFN-I production, we show that cGAS/STING signalling in macrophages can enhance transcription of specific immune recognition molecules including genes encoding the class II major histocompatibility complex (MHC-II) and co-stimulatory proteins, but without altering phagocytosis. Using FWPV mutants that are deficient in specific immunomodulators we are able to overcome the immunosuppression of wild type FWPV and show that this virus is sensed by cGAS, resulting in IFN-I and MHC-II transcription. These data show that the cGAS/STING/TBK1 pathway senses viral DNA in chicken macrophages and that this pathway regulates not only the antiviral interferon response but also modulates specific components of macrophage effector function machinery.

82 **Materials and Methods**

83 **Reagents**

84 Calf Thymus (CT) DNA (Sigma), Herring Testes (HT) DNA (Sigma), polyinosinic-polycytidylic
85 acid (poly(I:C), Invivogen), 2'3'-cGAMP (Invivogen) and chicken interferon alpha (Yeast-derived
86 Recombinant Protein, Kingfisher Biotech, Inc) were diluted in nuclease-free water (Ambion,
87 ThermoFisher). H-151 and BX795 (Invivogen) were diluted in DMSO, following the manufacturer's
88 protocols.

89 **Cell Culture**

90 HD11 cells, an avian myelocytomatosis virus (MC29)-transformed chicken macrophage-like cell line
91 (Beug et al., 1979), were incubated at 37°C, 5% CO₂. They were grown in RPMI (Sigma-Aldrich,
92 Germany) complemented with 2.5% volume per volume (v/v) heat-inactivated foetal bovine serum
93 (FBS; Sera Laboratories International Ltd), 2.5% volume per volume (v/v) chicken serum (New
94 Zealand origin, Gibco, Thermo Fisher Scientific), 10% Tryptose Phosphate Broth solution (Gibco,
95 Thermo Fisher Scientific), 2 mM L-glutamine (Gibco, Thermo Fisher Scientific), 50 µg/mL of
96 penicillin/streptomycin (P/S; Gibco, Thermo Fisher Scientific).

97 Chicken embryonic fibroblasts (CEFs) (Pirbright Institute, Woking, UK) were incubated at 37°C, 5%
98 CO₂ and were grown in Dulbecco's Modified Eagle Medium (DMEM) -F12 with Glutamax (Gibco),
99 5% v/v FBS, and 50 µg/mL P/S.

100 **Knock-out HD11 cell line generation by CRISPR-Cas9**

101 **CRISPR guide design**

102 According to the *MB21D1* (cGAS) and *TMEM137* (STING) sequences obtained from the Ensembl
103 database (release 94), single guide (sg)RNA sequences (Table 1) were designed targeting the
104 catalytic domain (residues 11-13 and 109) and start of the open reading frame, for cGAS and STING,
105 respectively.

106 **Knock-out cell lines generation using CRISPR-Cas9**

107 Genome editing of HD11 was performed using ribonucleoprotein (RNP) delivery. tracrRNA was
108 mixed with the target specific sgRNA (Table 1), followed by an incubation at 95°C. To form the
109 RNP complex, the tracrRNA/sgRNA mix was incubated with the Cas9 protein (IDT, Leuven,
110 Belgium) and electroporation enhancer at 21°C.

111 To generate knockout cells, 1x10⁶ cells per guide were electroporated with the corresponding RNP
112 complex using Lonza Electroporation Kit V (Lonza). After 48 h, the cells were expanded for future
113 experiments and their DNA were extracted using the PureLink Genomic DNA Kit (Thermo
114 Scientific, Waltham, MA, USA). The knockout efficiency was evaluated by genotyping the
115 polyclonal cell populations using MiSeq (Illumina) according to a published method (Schmidt et al.,
116 2016). The primers used for the sequencing are listed Table 2.

117 The successfully edited populations (using guides cGAS sg3 and STING sg1) were diluted to a
118 concentration of 0.5 cell/well and seeded in 96-well plates. Individual clones were sequenced by
119 MiSeq and the confirmed knockout clones were expanded for experiments.

120 **Primary macrophages**

121 Chicken bone marrow derived macrophages (BMDM) were generated as previously described (Garrido
122 et al., 2018). Briefly, femurs and tibias of 4 week-old immunologically mature White Leghorn (PA12
123 line) outbred chickens were removed, both ends of the bones were cut and the bone marrow was flushed
124 with RPMI supplemented with P/S. Cells were then washed and re-suspended in RPMI, loaded onto
125 an equal volume of Histopaque-1077 (Sigma-Aldrich, Germany), and centrifuged at 400 g for 20 min.
126 Cells at the interface were collected and washed twice in RPMI. Purified cells were seeded at
127 1×10^6 cells/ml in sterile 60 mm bacteriological petri dishes in RPMI supplemented with 10% FBS,
128 25 mM HEPES, 2 mM L-glutamine, P/S and 25 ng/ml recombinant chicken colony stimulating factor
129 1 (CSF-1) (Kingfisher Biotech, Inc) at 41 °C and 5% CO₂. Half of the medium was replaced with fresh
130 medium containing CSF-1 at day 3. At day 6, adherent cells were harvested and cultured in RPMI
131 supplemented with 10% FBS, 25 mM HEPES, 2 mM L-glutamine, and P/S prior to stimulation.

132 **Stimulation Assays**

133 HD11 (WT, cGAS and STING knockouts) were seeded in 12-well plates at a density of 3×10^5
134 cells/well. In the following day, the cells were transfected using TransIT-LT1 (Mirus Bio, USA) with
135 HT-DNA (1, 2 or 5 µg/mL), CT-DNA (1, 2 or 5 µg/mL) or Poly(I:C) (1 µg/mL), and harvested 6 h or
136 16 h post-transfection. In the priming assays, IFNα (200 ng/mL) was added 16 h hours prior to
137 transfection. 2'3' cGAMP was added at a concentration of 2.5 µg/mL and cells were harvested 6 h
138 post-treatment.

139 BMDM were seeded in 6-well plates at 8×10^5 cells/ml. In the following day, cells were transfected
140 using TransIT-LT1 with HT-DNA (2 µg/mL), CT-DNA (2 µg/mL) or Poly(I:C) (1 µg/mL), and
141 harvested 6 h post-transfection. In the priming assays, IFNα (50 ng/ml) was added 16 h prior
142 transfection to the cells supernatants. 2'3' cGAMP was added to cells supernatants at the
143 concentration of 10 µg/mL and the cells were harvested 6 h post-treatment.

144 **Chicken IFN-I bioassay**

145 The presence of IFN-I in supernatants of stimulated BMDM was measured indirectly using a
146 luciferase-based Mx-reporter bioassay (Schwarz et al., 2004). Briefly, cells from the quail fibroblast
147 cell line CEC32 carrying the luciferase gene under the control of chicken Mx promoter (kindly
148 provided by Prof. Peter Stäheli, University of Freiburg, Germany) were seeded at 2.5×10^5 cells/well
149 in 24-well plates and incubated at 41 °C under 5% CO₂. The next day, cells were incubated for 6 h
150 with the diluted supernatants (1/10 of total volume). Medium was removed and cells were washed
151 twice with PBS. Cells were lysed using the Cell Culture Lysis Reagent (Promega, USA), according
152 to the manufacturer's instructions, and luciferase activity was measured using the Luciferase assay
153 reagent (Promega, USA) and a GloMax-Multi Detection System (Promega, USA).

154 **Cell viability**

155 BMDM viability following different stimuli was assessed using the fluorescent DNA intercalator 7-
156 aminoactinomycin D (7-AAD, BD Biosciences, USA). Briefly, following stimulations, supernatants
157 were discarded and the cells were harvested and washed in PBS. Cells were stained according to the
158 manufacturer's protocol and the viability was analyzed by flow cytometry (BD FACS Calibur). Data
159 were expressed as the percentage of 7AAD positive cells over total acquired events (50,000 cells).

160 RNA Extraction

161 Cells were lysed by overlaying with 250 μ L of lysis buffer containing 4 M guanidine thiocyanate, 25
162 mM Tris pH 7, and 143 mM 2-mercaptoethanol. As a second step, 250 μ L of ethanol was added, and
163 the solution was transferred to a silica column (Epoch Life Science, Inc., Sugar Land, TX, USA) and
164 centrifuged; all centrifugation steps were performed for 90 seconds at 16600 g. The bound RNA was
165 washed by centrifugation with 500 μ L of buffer containing 1 M guanidine thiocyanate, 25 mM Tris
166 pH 7, and 10% ethanol, followed by a double washing step with 500 μ L of wash buffer 2 (25 mM
167 Tris pH 7 and 70% (v/v) ethanol). RNA was eluted by centrifugation in 30 μ L of nuclease-free water
168 and the concentration was measured using a NanoDrop 2000 Spectrophotometer (Thermo Scientific,
169 Waltham, MA, USA).

170 cDNA and qPCR

171 Using 500 ng of RNA extracted from HD11 cells, cDNA was produced using SuperScript III reverse
172 transcriptase, following the manufacture's protocol (Thermo Scientific, Waltham, MA, USA).
173 Samples were diluted in nuclease-free water in a 1:2.5 ratio. 1 μ L of the diluted product was used for
174 quantitative PCR (qPCR) in a final volume of 10 μ L. qPCR was performed using SybrGreen Hi-Rox
175 (PCR Biosystems Inc.) using primers described in Table 3. Fold change in mRNA expression was
176 calculated by relative quantification using hypoxanthine phosphoribosyltransferase (HPRT) as
177 endogenous control.

178 Total RNA (up to 1 μ g per reaction) from BMDM was reverse transcribed with iScript cDNA
179 synthesis kit (Bio-Rad, USA). Quantitative PCR was performed using 1 μ L of cDNA, 5 μ L of iQ
180 SYBR Green Supermix (Bio-Rad, USA), 0.25 μ L of each primer pair and 3.5 μ L of nuclease-free
181 water in a total reaction volume of 10 μ L. Fold-increase in gene expression was calculated by relative
182 quantification using HPRT and Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) as
183 endogenous controls.

184 Phagocytosis assay by flow cytometry

185 HD11 WT cells were seeded at a confluence of 3×10^5 cells/ml in 12-well plates. The cells were
186 primed with IFN α for 16h and then with transfected exogenous DNA (HT- and CT-DNA – 2 μ g/mL)
187 or treated with 2'3'cGAMP (5 μ g/mL) for 6 h. After this, the cells were incubated with Zymosan
188 coated beads conjugated with FITC at a ratio of 30 beads to 1 cell for all conditions for 40 min at
189 37 °C. The cells were wash two times in PBS and fixed in suspension using the solution (missing ref;
190 BD Biosciences) with 4% PFA. Cell populations were counted by analysis on a CytoFLEX
191 cytometer.

192 Fowlpox virus growth and titration

193 Fowlpox WT (FP9) and mutants (FPV012 (Laidlaw et al., 2013) and FPV184 (Giotis & Skinner,
194 unpublished)) were propagated in primary chicken embryonic fibroblasts (CEFs) and grown in
195 DMEM-F12 (Thermo Fisher Scientific, Waltham, MA, USA) containing 1% FBS and 5% P/S, and
196 harvested 5 days later. 10-fold dilutions of cell supernatants were prepared in serum-free DMEM-F12
197 and used to inoculate confluent monolayers of CEFs for 1.5 h at 37°C. Cells were then overlaid with
198 2xMEM:CMC (1/1 ratio). The foci were counted 7 days later after staining with Toluidine Blue.

199 Fowlpox virus infection

HD11 cells were seeded in 12-well plates in the day prior infection. Fowlpox viruses were diluted in serum-free DMEM-F12 at a multiplicity of infection (MOI) of 3 and added in the cells (1 ml per well). Infected cells and supernatants were collected from infections at 8h and 24h post-infection.

Statistical Analysis

Prism 7 (GraphPad) was used to generate graphs and perform statistical analysis. Data were analyzed using an unpaired t test with Welch's correction unless stated otherwise. Data with $P < 0.05$ was considered significant and 2-tailed P-value were calculated and presented as: <0.0001 - ****, >0.0001 - ***, >0.001 - **, >0.01 - *. Each experiment has at least two biological replicates unless stated.

Results

Intracellular DNA activates a IFN-I response in chicken macrophages

In order to assess the ability of chicken macrophages to sense and respond to intracellular DNA we used a combination of the monocytic cell line HD11 and primary bone marrow derived macrophages (BMDM). Transfection of CT DNA, increasing doses of HT-DNA or the RNA analogue poly(I:C) into HD11 cells resulted in transcription of chicken interferon- β (IFN β) and the interferon stimulated gene (ISG) ISG12.2, an orthologue of mammalian IFI6 (Figure 1A). A dose-dependent response to DNA was observed. Transfection of DNA into primary BMDMs also resulted in IFN β and ISG12.2 transcription (Figure 1B) and IFN-I secretion as measured by a bioassay (Figure 1C), indicating that this response is present in both primary macrophages and the transformed monocytic HD11 cell line.

Since in mammalian systems STING is recognised as an interferon stimulated gene (ISG) (Ma et al., 2015), we sought to understand the effect of IFN-I priming of macrophages on the response to intracellular DNA. Pre-treatment of HD11 or BMDM with chIFN α resulted in an enhancement of IFN β transcription following DNA stimulation and confirmed ISG12.2 as an ISG (Figure 1D). This signalling enhancement might be explained by increased transcription of STING and/or IRF7 following IFN α treatment (Supplementary Figure 1). Across all HD11 and BMDM DNA stimulations we found that there was little observable or measurable cell death (Figure 1E), indicating that, in chicken macrophages, cell death is not a specific output of STING signalling.

Intracellular DNA stimulates transcription of MHC-II and co-stimulatory molecules

The sensing of both intracellular and extracellular pathogens activates macrophages, causing up-regulation or enhancement of effector functions designed to combat infection. We hypothesised that DNA transfection, mimicking the presence of intracellular infection, might result in direct effects on the molecules that contribute to T cell stimulation. There are two chicken MHC-II beta chain genes, BLB1 and BLB2, both of which were transcriptionally upregulated by DNA stimulation in chicken BMDMs (Figure 2A). In HD11 cells, BLB1 transcription was upregulated by DNA stimulation, while BLB2 transcription was upregulated only by IFN α pre-treatment (not shown), highlighting possible differences between primary and transformed cells in this specific context (Figure 2B). CD86 and CD40 are key co-stimulatory molecules in T cell activation. In BMDM CD86 and CD40 transcription was upregulated in response to DNA stimulation (Figure 2A). There was, however, no measurable impact of DNA stimulation on phagocytosis as measured by bead-uptake assays in HD11 cells (Figure 2C). As such, key molecules involved in T cell activation by macrophages are regulated by DNA stimulation, but not all macrophage effector functions are equally enhanced by this signal.

241 **STING and TBK1 contribute to DNA-driven transcriptional responses in chicken BMDMs**

242 In order to dissect the signalling pathway downstream of intracellular DNA sensing, we first used the
243 ligand 2'3'-cGAMP, the enzymatic product of cGAS that directly binds and activates STING
244 (Ablasser et al., 2013b). Treatment of BMDMs or HD11 cells with 2'3'-cGAMP led to increased
245 transcription levels of IFN β , ISG12.2, BLB1, BLB2, CD86 and CD40 (Figure 3A,B). This response,
246 and the response to DNA stimulation, could be reduced by small molecule inhibitors of STING
247 (H151) and the kinase TBK1 (BX795), indicating the existence of a STING and TBK1-dependent
248 signalling pathway in chicken macrophages and evidencing the cross-species utility of these two
249 pharmacological inhibitors (Figure 3D). As with DNA stimulation, there was no measurable impact
250 of cGAMP treatment on phagocytosis in HD11 cells (Figure 3E).

251 **cGAS is essential for intracellular DNA-dependent IFN-I and MHC-II transcription in HD11** 252 **cells**

253 To address the possibility that cGAS is a principle PRR responsible for sensing intracellular DNA in
254 chicken macrophages, we generated HD11 knockout cell lines using CRISPR/Cas9 genome editing.
255 To do this we analysed the annotated cGAS sequence in the current release of the *Gallus gallus*
256 genome and designed gRNA sequences targeting regions of the gene which exhibited high
257 conservation across multiple orthologues. By sequencing single cell clones we generated multiple
258 cGAS knockout cell lines with two different gRNAs. By sequencing across the gRNA PAM target
259 sites, we characterised indels to confirm the knockout status in these clones (eg Figure 4A).
260 Stimulation of multiple cGAS knockout HD11 clones, each with a different indel, with DNA resulted
261 in an abrogation of IFN-I and ISG transcription indicating that cGAS is a key PRR for sensing
262 intracellular DNA in chicken macrophages (Figure 4B and Supplementary Figure 2). cGAS knockout
263 also abrogated the upregulation of DNA-driven BLB1 stimulation, indicating the cGAS-dependent
264 signalling is responsible for regulation of MHC class II transcription in this context (Figure 4B).
265 These data were independent of IFN α pre-treatment, which enhanced IFN-I and BLB1 transcription
266 in WT DNA-stimulated cells, but did not affect cGAS KO cells (Figure 4C). Consistent with the
267 mammalian cGAS mechanism, stimulation of WT or cGAS KO cells with 2'3'-cGAMP resulted in
268 robust IFN-I transcription, indicating IFN-I production by direct STING ligation was not affected by
269 cGAS KO (Figure 4D). These data confirm the intracellular DNA PRR function of cGAS in chicken
270 macrophages.

271 **STING is essential for intracellular DNA-dependent IFN-I transcription in HD11 cells**

272 In parallel, using the same methodology, we generated multiple STING knockout HD11 cell lines
273 (Figure 5A). Stimulation of these cells with DNA phenocopied the cGAS knockout lines, confirming
274 the function of chicken STING downstream of cGAS in the intracellular DNA sensing pathway
275 (Figure 5B, Supplementary Figure 3). These data are consistent with the presence of a cGAS/STING
276 pathway in HD11 cells and, in concert with the data using H151 in BMDMs, indicate the function of
277 STING as a critical adaptor protein for intracellular DNA sensing in chicken macrophages.

278

279 **Fowlpox triggers a cGAS / STING dependent DNA sensing pathway in HD11 cells**

280 FWPV replication exposes large quantities of DNA to the cytoplasm of infected cells making it a
281 prime target for intracellular DNA sensing PRRs. Despite this, using the wild-type vaccine strain FP9
282 we, and others (Giotis and Skinner, 2019; Laidlaw et al., 2013), observe little or no IFN-I

transcription in infected cells, and indeed a downregulation of IFN and MHC transcription (Figure 6A). The lack of IFN-I response in poxvirus infected cells is likely due to the presence of numerous virally-encoded suppressors of PRR signalling and IFN-I production (Laidlaw et al., 2013; Smith et al., 2013), hence deletion of specific innate immunomodulators from the viral genome can result in a virus that stimulates host IFN-I signalling. We made use of FWPV mutants FPV012 and FPV184 ((Laidlaw et al., 2013, Giotis & Skinner, unpublished), each deficient in single genes that are proposed immunomodulators, and both of which induce IFN-I production from infected cells (Laidlaw et al., 2013), including HD11 cells (Figure 6B). In the absence of cGAS or STING the transcription of IFN-I, ISG12.2, BLB1 and CD40 by FPV184 or FPV012 was significantly lower at 24h post infection (Figure 6B), despite robust infection of HD11 cells by all three virus strains (Figure 6C), indicating that FWPV is sensed in infected cells by the DNA sensing PRR cGAS and that the cGAS/STING pathway is responsible for FWPV-induced IFN-I production and MHC-II transcription.

Discussion

The ability of innate immune cells to detect virus infection is dependent on a set of PRRs that directly bind viral nucleic acids. Macrophages act in this context as tissue-resident sentinel sensors of infection that express a broad repertoire of PRRs and mount a rapid and robust innate immune response to viruses and other pathogens. Indeed intracellular DNA sensing was first described in macrophages (Stetson and Medzhitov, 2006). As well as interferon and cytokine production, activated macrophages use effector functions for pathogen clearance and for activation of adaptive immunity. In mammalian systems the signalling outputs downstream of intracellular DNA detection in macrophages include IRF-dependent IFN and cytokine production and cell death driven by the AIM2 inflammasome. In chicken macrophages, which lack AIM2, we find that intracellular DNA sensing produces IFN but doesn't result in measurable cell death, rather it upregulates a specific set of antigen presentation machinery including the MHC-II gene BLB1 and co-stimulatory molecules, providing a direct link between anti-viral innate sensing and the initiation of adaptive immunity.

During DNA virus infection, the cGAS/STING-dependent signalling pathway is triggered by viral DNA, resulting in type-I interferon production via activation of TBK1 and the IRF family of transcription factors. Although well defined in mammalian systems, the function of chicken cGAS and STING has only more recently been identified (Gao et al., 2018; Vitak et al., 2016). FWPV is an avian poxvirus that causes skin lesions and respiratory infections and can infect multiple cell types including macrophages (Williams et al., 2010). Here we show that the cGAS/STING pathway in chicken macrophages can sense FWPV infection and is responsible for the IFN-I response as well as for upregulation of BLB1.

In order to escape detection and evade host anti-viral responses, poxviruses like FWPV encode a broad range of immunomodulatory proteins that target PRR signalling pathways resulting in these viruses being able to effectively inhibit IFN production from infected cells. These immune evasion mechanisms mask the signalling outputs of PRR signalling during infection with wild type poxviruses. To overcome this issue, we used two mutant FWPVs with deletions in individual genes that block IFN-I production during infection. Infection of cells with FPV184 and FPV012 (Giotis et al., 2016) resulted in interferon and ISG transcription, which was lost in cGAS and STING knockout lines. FWPV DNA is therefore sensed by the cGAS/STING pathway and the downstream signalling response leading to IFN-I production is effectively blocked by the wild type virus.

Birds occupy the same habitats as mammals, have comparable ranges of life span and body mass, and confront similar pathogen challenges, yet birds have a different repertoire of organs, cells, molecules and genes of the immune system compared to mammals (Kaiser, 2010). It is increasingly evident that the immune system of avian species is rather different from those of model mammalian species. Untested extrapolation from mammalian systems cannot provide the quality of knowledge that is required for understanding host-pathogen relationships in birds. Here we find that the signalling downstream of chicken cGAS leading to IFN-I transcription is similar to that found in mammalian systems. The presence of orthologues of STING and TBK1 in the chicken genome and their functional inhibition by small molecule compounds (H151 and BX795) is indicative of mechanistic signalling pathway conservation. The chicken genome also contains an orthologue of IRF3, which is the main transcription factor downstream of STING/TBK1 activation, although chicken IRF7 (as this gene is annotated) is not equivalent to mammalian IRF3 or IRF7 and may be considered more as a hybrid these two genes (Grant et al., 1995). It is likely that chicken IRF7 and TBK1 are recruited by STING following 2'3'-cGAMP ligation and that subsequent phosphorylation, dimerisation and nuclear translocation of IRF7 leads to DNA-induced IFN-I transcription (Cheng et al., 2019; Gao et al., 2018). Recent evidence has implicated chicken cGAS and STING in avian antiviral defence, in particular against Marek's Disease Virus (MDV) and chicken adenovirus 4 (Li et al., 2019; Wang et al., 2020) in fibroblasts. Using CRISPR/Cas9 technology to knockout STING and cGAS in a transformed monocytic cell line (HD11) and complementing these data in primary macrophages with pharmacological inhibitors we have been able to show this cGAS/STING/TBK1 pathway is active in chicken macrophages. The use of primary cells in this context is important as transformation or immortalisation can significantly alter PRR pathways so as to obscure physiological signalling mechanisms.

IFN-I is one of the most effective anti-viral innate immune mediators. Secretion and subsequent ISG transcription induced by autocrine and paracrine IFN receptor signalling sets an anti-viral/inflammatory state in infected and bystander cells. As an example, chicken IFN β was shown to be an autocrine/paracrine pro-inflammatory mediator in chicken macrophages (Garrido et al., 2018), with direct effects in macrophage effector functions. Nucleic acid sensing PRRs therefore provide a rapid and potent innate response helping to combat infection and reduce viral spread in infected tissues. At the same time, innate immune responses can initiate and amplify adaptive immune responses for example, by regulating functions of antigen presenting cells (APCs), promoting cross-priming and stimulating antibody production (Desmet and Ishii, 2012; Loré et al., 2003; Schulz et al., 2005). In both mammals and birds, macrophages are key regulators of adaptive immunity as principle APCs. By processing and presenting antigen to T and B cells, macrophages directly trigger adaptive responses. The discovery that cGAS/STING signalling can directly regulate the transcription of MHC genes in macrophages provides further evidence linking PRR signalling with the activation of adaptive immunity during infection. It remains to be explored exactly how the transcription of BLB1 and BLB2 is regulated by cGAS/STING signalling. In tissues, macrophages survey the local environment for infection and damage. In this context, macrophage effector functions may be modulated by the presence of innate immune mediators in the tissue. The priming effect of IFN α as an enhancer of macrophage DNA sensing, by upregulating STING expression, suggests a possible mechanism of bystander surveillance. Tissue resident macrophages may respond to signals, including IFN-I and cGAMP, secreted from virally infected stromal cells by enhancing specific effector functions appropriate to defend against viral infection in the tissue (Ablasser et al., 2013c; Schadt et al., 2019).

Our data adds to the list of chicken cGAS/STING functions in sensing of avian DNA viruses such as MDV and Adenovirus 4 that replicate in the nucleus or FWPV that replicates in the cytoplasm, and in

the regulation of macrophage effector functions. The ability of this pathway to sense a broad range of DNA viruses that replicate in different compartments in avian innate immune cells indicates that this pathway is a primary DNA sensing mechanism for DNA viruses in chickens.

Figure Legends

Figure 1. Intracellular DNA activates an IFN-I response in chicken macrophages. (A) HD11 cells were transfected with HT-DNA (1, 2 and 5 $\mu\text{g/mL}$), CT-DNA (5 $\mu\text{g/mL}$) or Poly(I:C) and transcription of IFNB and ISG12.2 measured by qRT-PCR 6 h later. (B) Chicken BMDM were transfected with HT-DNA, CT-DNA (2 $\mu\text{g/mL}$) or Poly(I:C) (1 $\mu\text{g/mL}$) and transcription of IFNB and ISG12.2 measured by qRT-PCR 6 h later. (C) Resting BMDMs or BMDMs primed with IFN α for 6 h were transfected with HT-DNA, CT-DNA (2 $\mu\text{g/mL}$) or Poly(I:C) (1 $\mu\text{g/mL}$) and interferon activity in the supernatants was measured after 24 h using a bioassay. (D) HD11 or BMDM were primed with IFN α for 6h, transfected with HT-DNA, CT-DNA, or Poly(I:C) and transcription of IFNB and ISG12.2 measured by qRT-PCR 6 h later. (E) BMDM were primed with IFN α for 6h, transfected with HT-DNA, CT-DNA, or Poly(I:C) and cell viability measured by 7AAD staining 24 h later. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$; ****: $p < 0.0001$; ns: no significant difference.

Figure 2. Intracellular DNA stimulates transcription of MHC-II and co-stimulatory molecules. (A) BMDMs or (B) HD11 cells were transfected with HT-DNA, CT-DNA, or Poly(I:C) and transcription of BLB1, BLB2, CD40 and CD86 measured by qRT-PCR 6 h later. (C) HD11 cells were stimulated with HT-DNA, CT-DNA, or Poly(I:C) and 6 h later phagocytosis was monitored by FITC-conjugated, zymosan coated bead uptake. Histograms of non-treated versus treated cells (left panels) and respective percentages of FITC positive cells for each treatment tested (right panel) are presented. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$; ****: $p < 0.0001$; ns: no significant difference.

Figure 3. STING and TBK1 contribute to DNA-driven transcriptional responses in chicken BMDMs. (A) HD11 and (B) BMDM cells were treated with 2'3'cGAMP (10 $\mu\text{g/mL}$) and qRT-PCR carried out 6 h later for the indicated genes. (C) BMDM were treated with the STING inhibitor H-151 (10 μM) or TBK1 inhibitor BX795 (1 μM) for 1 h before transfection with HT-DNA and CT-DNA. 6 h later RNA was extracted and qRT-PCR carried out for the indicated genes. (D) BMDM were treated with the STING inhibitor H-151 (10 μM) or TBK1 inhibitor BX795 (1 μM) for 1 h before treatment with 2'3'cGAMP (10 $\mu\text{g/mL}$). 6 h later RNA was extracted and qRT-PCR carried out for the indicated genes. (E) HD11 cells were treated with 2'3'cGAMP (2.5 $\mu\text{g/mL}$) 6 h later phagocytosis was monitored by FITC-conjugated, zymosan coated bead uptake. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$; ****: $p < 0.0001$; ns: no significant difference

Figure 4. cGAS is essential for intracellular DNA-dependent IFN-I and MHC-II transcription in HD11 cells. (A) Example of identification of indel in clonally selected HD11 cGAS KO using NGS sequencing. (B, C) WT and cGAS KO HD11 cells were transfected with HT-DNA, CT-DNA (2 $\mu\text{g/mL}$) or Poly(I:C) (1 $\mu\text{g/mL}$) for 6 h and transcription of the indicated genes measured by qRT-PCR. (D) cGAS KO HD11 cells were primed with IFN α for 6h, transfected with HT-DNA, CT-DNA, or Poly(I:C) and transcription of IFNB and ISG12.2 measured by qRT-PCR 6 h later (E) WT or cGAS KO cells were treated with 2'3'cGAMP (10 $\mu\text{g/mL}$) and transcription of IFNB measured by qRT-PCR 6 h later. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$; ****: $p < 0.0001$; ns: no significant difference

Figure 5. STING is essential for intracellular DNA-dependent IFN-I transcription in HD11

cells. (A) Example of identification of indel in clonally selected HD11 STING KO using NGS sequencing. (B) WT and STING KO HD11 cells were transfected with HT-DNA, CT-DNA (2 µg/mL) or Poly(I:C) (1 µg/mL) for 6 h and transcription of the indicated genes measured by qRT-PCR 6 h later. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$; ****: $p < 0.0001$; ns: no significant difference

Figure 6. Fowlpox triggers a cGAS / STING dependent DNA sensing pathway in HD11 cells.

(A) HD11 cells were infected with FWPV strain FP9 at a multiplicity of infection of three. 24 h later RNA was extracted and qRT-PCR carried out for the indicated genes. (B, C) HD11 WT, cGAS or STING KO cells were infected with FP9, FPV012 or FPV184 at a multiplicity of infection of three. 24 h later RNA was extracted and qRT-PCR carried out for the indicated genes. *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$; ****: $p < 0.0001$; ns: no significant difference

Supplementary Figure 1. Effect of IFN α priming on expression levels of STING and IRF7 in

BMDM and HD11. BMDM or HD11 cells were treated with IFN α for 6 h and transcription of STING and IRF7 measured by qRT-PCR 6 h later.

Supplementary Figure 2. cGAS is essential for intracellular DNA-dependent IFN-I

transcription in HD11 cells. WT or three individual cGAS knockout clones with different indels were stimulated with HT-DNA (2 µg/mL) and IFNB transcription measured by qRT-PCR 6 h later.

Supplementary Figure 3. STING is essential for intracellular DNA-dependent IFN-I

transcription in HD11 cells. (A) WT or three individual cGAS knockout clones with different indels were stimulated with HT-DNA (2 µg/mL) and IFNB transcription measured by qRT-PCR 6 h later.

Tables

Table 1. CRISPR/Cas9 guide RNAs

Gene	Target	Guide	Sequence
cGAS	Catalytic Domain	sgRNA1	CTCTTTCTCGCATATCGAGA
		sgRNA2	ACGGCCTCAACATAGAATGC
		sgRNA3	TTTGGTTCAGATATCTGCAA
		sgRNA4	ACTGTGAAAAGGAAAAAGCG
STING	Coding Region	sgRNA1	GTAGCCGATGTAGTAGGAC
		sgRNA2	GTGCAGACGCTGCGGATGA

Table 2. Illumina sequencing primers

Gene	Guide	Forward primer	Reverse primer
cGAS	sgRNA1	CTATTTAAATCTCGTGCTCACCCC	CTCACTCCCTGTTCTAAATAACG
	sgRNA2/ sgRNA4	GTGTTTCTTCTGTTATGGAAAAGG	GCTTGGCCACTAAGTAAATTGG
	sgRNA3	CCACTTGAATGCACATCAGTCTGG	CCAGTGTCGTCACTCTCATCTAGCT
STING	sgRNA1	TCCACAGGGCCACCACT	TGCAGGAGCCGTTTCCATCT
	sgRNA2	CAACCAGGAGCAGCCCTGCT	CTGGAGTGCAGGTGGAAGATCTCC

Table 3. qRT-PCR primer sequences

Gene	Forward primer	Reverse primer
HPRT	TGGTGGGGATGACCTCTCAA	GGCCGATATCCCACACTTCG
IFNB	TCCTGCAACCATCTTCGTCA	CACGTCTTGTTGTGGGCAAG
ISG12-2	TGACCAGAACGTCCACAAAGCCG	ACCTGCTCCTGGACCGATGCTT
BLB1	GTGAGCCGCAAGCTGAATAC	ACCGTGAAGGACTCCACAAC
BLB2	ATGAATGAAGTGGACAGGGTCT	TTCAGGAACCACTTCACCTCG
CD40	AGCCATGCCACTTCTGGAC	ATCGGAAGTGTTTCGTCCCTT
CD86	TATGCACGTGGACAAGGGAC	AACCTCCGCTGGAAGAACAG
STING	AGCTCCCTACCTCCATCAGGA	TCTGGAAAACCCCAGCATCTC
IRF7	TGCCTCAGGCGTCCCCAATG	TGTGTGCCACAGGGTTGGC
FPV094	TATAATGAATGGCGCTGTGT	GTTTTGCTATCTTGGCTGT
FPV168	ACCTCAAACAACCTCATC	GTTAATACTTGTGACTGCTG

476

477

478 **Conflict of Interest**

479 The authors declare that the research was conducted in the absence of any commercial or financial
480 relationships that could be construed as a potential conflict of interest.

481 **Author Contributions**

482 BF and CB provided the funding and supervised the work. MO, DR, RG, VG and EK performed the
483 experiments and statistical analysis. SG and MS generated the mutant fowlpox viruses. BF, CB, RG
484 designed the study and wrote the manuscript.

485 **Funding**

486 This work was funded by BBSRC grants RG94719 (BF and CEB), BB/E009956/1, BB/G018545/1,
487 BB/H005323/1 & BB/K002465/1 (MAS)

488 Acknowledgments

489 This manuscript has been released as a Pre-print at BioRxiv. We thank the experimental facility PFIE
490 (Plateforme d'Infectiologie Expérimentale, Centre INRAE Val de Loire, Nouzilly, France) for
491 providing the animals used for the isolation of primary macrophages.

492 References

- 493 Ablasser, A., Hertrich, C., Wassermann, R., and Hornung, V. (2013a). Nucleic acid driven sterile
494 inflammation. *Clin. Immunol.* *147*, 207–215.
- 495 Ablasser, A., Goldeck, M., Cavlar, T., Deimling, T., Witte, G., Röhl, I., Hopfner, K.-P., Ludwig, J.,
496 and Hornung, V. (2013b). cGAS produces a 2'-5'-linked cyclic dinucleotide second messenger that
497 activates STING. *Nature* *498*, 380–384.
- 498 Ablasser, A., Schmid-Burgk, J.L., Hemmerling, I., Horvath, G.L., Schmidt, T., Latz, E., and
499 Hornung, V. (2013c). Cell intrinsic immunity spreads to bystander cells via the intercellular transfer
500 of cGAMP. *Nature* *503*, 530–534.
- 501 Beug, H., von Kirchbach, A., Döderlein, G., Conscience, J.F., and Graf, T. (1979). Chicken
502 hematopoietic cells transformed by seven strains of defective avian leukemia viruses display three
503 distinct phenotypes of differentiation. *Cell* *18*, 375–390.
- 504 Bryant, C.E., Orr, S., Ferguson, B., Symmons, M.F., Boyle, J.P., and Monie, T.P. (2015).
505 International union of basic and clinical pharmacology. XCVI. Pattern recognition receptors in health
506 and disease. *Pharmacol. Rev.* *67*.
- 507 Cheng, Y., Zhu, W., Ding, C., Niu, Q., Wang, H., Yan, Y., and Sun, J. (2019). IRF7 Is Involved in
508 Both STING and MAVS Mediating IFN- β Signaling in IRF3-Lacking Chickens. *J. Immunol.* *203*,
509 1930–1942.
- 510 Crow, Y.J., and Rehwinkel, J. (2009). Aicardi-Goutieres syndrome and related phenotypes: linking
511 nucleic acid metabolism with autoimmunity. *Hum. Mol. Genet.* *18*, R130-6.
- 512 Desmet, C.J., and Ishii, K.J. (2012). Nucleic acid sensing at the interface between innate and adaptive
513 immunity in vaccination. *Nat. Rev. Immunol.* *12*, 479–491.
- 514 Gaidt, M.M., Ebert, T.S., Chauhan, D., Ramshorn, K., Pinci, F., Zuber, S., O'Duill, F., Schmid-
515 Burgk, J.L., Hoss, F., Buhmann, R., et al. (2017). The DNA Inflammasome in Human Myeloid Cells
516 Is Initiated by a STING-Cell Death Program Upstream of NLRP3. *Cell* *171*, 1110-1124.e18.
- 517 Gao, L., Li, K., Zhang, Y., Liu, Y., Liu, C., Zhang, Y., Gao, Y., Qi, X., Cui, H., Wang, Y., et al.
518 (2018). Inhibition of DNA-Sensing Pathway by Marek's Disease Virus VP23 Protein through
519 Suppression of Interferon Regulatory Factor 7 Activation. *J. Virol.* *93*.
- 520 Garrido, D., Alber, A., Kut, E., Chanteloup, N.K., Lion, A., Trotereau, A., Dupont, J., Tedin, K.,
521 Kaspers, B., Vervelde, L., et al. (2018). The role of type I interferons (IFNs) in the regulation of
522 chicken macrophage inflammatory response to bacterial challenge. *Dev. Comp. Immunol.* *86*, 156–
523 170.

524 Giotis, E.S., and Skinner, M.A. (2019). Spotlight on avian pathology: fowlpox virus. *Avian Pathol.*
525 48, 87–90.

526 Giotis, E.S., Robey, R.C., Skinner, N.G., Tomlinson, C.D., Goodbourn, S., and Skinner, M.A.
527 (2016). Chicken interferome: avian interferon-stimulated genes identified by microarray and RNA-
528 seq of primary chick embryo fibroblasts treated with a chicken type I interferon (IFN- α). *Vet. Res.*
529 47, 75.

530 Glück, S., Guey, B., Gulen, M.F., Wolter, K., Kang, T.W., Schmacke, N.A., Bridgeman, A.,
531 Rehwinkel, J., Zender, L., and Ablasser, A. (2017). Innate immune sensing of cytosolic chromatin
532 fragments through cGAS promotes senescence. *Nat. Cell Biol.* 19, 1061–1070.

533 Grant, C.E., Vasa, M.Z., and Deeley, R.G. (1995). cIRF-3, a new member of the interferon regulatory
534 factor (IRF) family that is rapidly and transiently induced by dsRNA. *Nucleic Acids Res.* 23, 2137–
535 2146.

536 Gulen, M.F., Koch, U., Haag, S.M., Schuler, F., Apetoh, L., Villunger, A., Radtke, F., and Ablasser,
537 A. (2017). Signalling strength determines proapoptotic functions of STING. *Nat. Commun.* 8.

538 Kaiser, P. (2010). Advances in avian immunology-prospects for disease control: A review. *Avian*
539 *Pathol.* 39, 309–324.

540 Laidlaw, S.M., Robey, R., Davies, M., Giotis, E.S., Ross, C., Buttigieg, K., Goodbourn, S., and
541 Skinner, M.A. (2013). Genetic Screen of a Mutant Poxvirus Library Identifies an Ankyrin Repeat
542 Protein Involved in Blocking Induction of Avian Type I Interferon. *J. Virol.* 87, 5041–5052.

543 Li, T., and Chen, Z.J. (2018). The cGAS–cGAMP–STING pathway connects DNA damage to
544 inflammation, senescence, and cancer. *J. Exp. Med.* 215, 1287–1299.

545 Li, K., Liu, Y., Xu, Z., Zhang, Y., Luo, D., Gao, Y., Qian, Y., Bao, C., Liu, C., Zhang, Y., et al.
546 (2019). Avian oncogenic herpesvirus antagonizes the cGAS-STING DNA-sensing pathway to
547 mediate immune evasion. *PLoS Pathog.* 15.

548 Li, X.-D., Wu, J., Gao, D., Wang, H., Sun, L., and Chen, Z.J. (2013). Pivotal roles of cGAS-cGAMP
549 signaling in antiviral defense and immune adjuvant effects. *Science* 341, 1390–1394.

550 Loré, K., Betts, M.R., Brenchley, J.M., Kuruppu, J., Khojasteh, S., Perfetto, S., Roederer, M., Seder,
551 R.A., and Koup, R.A. (2003). Toll-Like Receptor Ligands Modulate Dendritic Cells to Augment
552 Cytomegalovirus- and HIV-1-Specific T Cell Responses. *J. Immunol.* 171, 4320–4328.

553 Lousberg, E.L., Fraser, C.K., Tovey, M.G., Diener, K.R., and Hayball, J.D. (2010). Type I
554 interferons mediate the innate cytokine response to recombinant fowlpox virus but not the induction
555 of plasmacytoid dendritic cell-dependent adaptive immunity. *J. Virol.* 84, 6549–6563.

556 Ma, F., Li, B., Yu, Y., Iyer, S.S., Sun, M., and Cheng, G. (2015). Positive feedback regulation of
557 type I interferon by the interferon-stimulated gene STING. *EMBO Rep.* 16, 202–212.

558 Mansur, D.S., Smith, G.L., and Ferguson, B.J. (2014). Intracellular sensing of viral DNA by the
559 innate immune system. *Microbes Infect.* 16, 1002–1012.

560 Mullard, A. (2017). Can innate immune system targets turn up the heat on “cold” tumours? *Nat. Rev.*
561 *Drug Discov.* 2017 171.

562 Schadt, L., Sparano, C., Schweiger, N.A., Silina, K., Cecconi, V., Lucchiari, G., Yagita, H.,
563 Guggisberg, E., Saba, S., Nascakova, Z., et al. (2019). Cancer-Cell-Intrinsic cGAS Expression
564 Mediates Tumor Immunogenicity. *Cell Rep.* 29, 1236-1248.e7.

565 Schmidt, T., Schmid-Burgk, J.L., Ebert, T.S., Gaidt, M.M., and Hornung, V. (2016). Designer
566 nuclease-mediated generation of knockout THP1 cells. In *Methods in Molecular Biology*, (Humana
567 Press Inc.), pp. 261–272.

568 Schulz, O., Diebold, S.S., Chen, M., Näslund, T.I., Nolte, M.A., Alexopoulou, L., Azuma, Y.-T.,
569 Flavell, R.A., Liljeström, P., and Reis e Sousa, C. (2005). Toll-like receptor 3 promotes cross-
570 priming to virus-infected cells. *Nature* 433, 887–892.

571 Schwarz, H., Harlin, O., Ohnemus, A., Kaspers, B., and Staeheli, P. (2004). Synthesis of IFN-β by
572 Virus-Infected Chicken Embryo Cells Demonstrated with Specific Antisera and a New Bioassay. *J.*
573 *Interf. Cytokine Res.* 24, 179–184.

574 Smith, G.L., Benfield, C.T.O., Maluquer de Motes, C., Mazzon, M., Ember, S.W.J., Ferguson, B.J.,
575 and Sumner, R.P. (2013). Vaccinia virus immune evasion: mechanisms, virulence and
576 immunogenicity. *J. Gen. Virol.* 94, 2367–2392.

577 Stetson, D.B., and Medzhitov, R. (2006). Recognition of cytosolic DNA activates an IRF3-dependent
578 innate immune response. *Immunity* 24, 93–103.

579 Sze, A., Belgnaoui, S.M., Olganier, D., Lin, R., Hiscott, J., and Van Grevenynghe, J. (2013). Host
580 restriction factor SAMHD1 limits human T cell leukemia virus type 1 infection of monocytes via
581 STING-mediated apoptosis. *Cell Host Microbe* 14, 422–434.


582 Tanaka, Y., and Chen, Z.J. (2012). STING specifies IRF3 phosphorylation by TBK1 in the cytosolic
583 DNA signaling pathway. *Sci. Signal.* 5, ra20.

584 Vitak, N., Hume, D.A., Chappell, K.J., Sester, D.P., and Stacey, K.J. (2016). Induction of interferon
585 and cell death in response to cytosolic DNA in chicken macrophages. *Dev. Comp. Immunol.* 59,
586 145–152.

587 Wang, J., Ba, G., Han, Y.Q., Ming, S.L., Wang, M. Di, Fu, P.F., Zhao, Q.Q., Zhang, S., Wu, Y.N.,
588 Yang, G.Y., et al. (2020). Cyclic GMP-AMP synthase is essential for cytosolic double-stranded DNA
589 and fowl adenovirus serotype 4 triggered innate immune responses in chickens. *Int. J. Biol.*
590 *Macromol.* 146, 497–507.


591 Williams, S.M., Smith, J.A., Garcia, M., Brinson, D., Kiupel, M., and Hofacre, C. (2010). Severe
592 histiolymphocytic and heterophilic bronchopneumonia as a reaction to in ovo fowlpox vaccination in
593 broiler chicks. *Vet. Pathol.* 47, 177–180.

594




BMDM

A




B


HD11












C









Beads Intake






A**B****C****D**





IFNB



IFNB

