

Coupling of H3K27me3 recognition with transcriptional repression through the BAH-PHD-CPL2 complex in *Arabidopsis*

Yi-Zhe Zhang^{1,2†}, Jianlong Yuan^{1,2†}, Lingrui Zhang^{3†}, Chunxiang Chen¹, Yuhua Wang¹, Guiping Zhang¹, Li Peng¹, Si-Si Xie^{1,2}, Jing Jiang⁴, Jian-Kang Zhu^{1,3*}, Jiamu Du^{5*} and Cheng-Guo Duan^{1,4*}

¹Shanghai Center for Plant Stress Biology and CAS Center for Excellence in Molecular Plant Sciences, Chinese Academy of Sciences, Shanghai 201602, China.

²University of Chinese Academy of Sciences, Beijing 100049, China.

³Department of Horticulture and Landscape Architecture, Purdue University, West Lafayette, IN 47907, USA

⁴State Key Laboratory of Crop Stress Adaptation and Improvement, School of Life Sciences, Henan University, Kaifeng 475004, China

⁵Key Laboratory of Molecular Design for Plant Cell Factory of Guangdong Higher Education Institutes, Institute of Plant and Food Science, School of Life Sciences, Southern University of Science and Technology, Shenzhen 518055, China.

[†]These authors contributed equally to this work.

*CORRESPONDING AUTHORS:

Cheng-Guo Duan (cgduan@psc.ac.cn)

Jiamu Du (dujm@sustech.edu.cn)

Jian-Kang Zhu (jzkzhu@psc.ac.cn)

SUMMARY

Histone 3 Lys 27 trimethylation (H3K27me₃)-mediated epigenetic silencing plays a critical role in multiple biological processes. However, the H3K27me₃ recognition and transcriptional repression mechanisms are only partially understood. Here, we report a new mechanism for H3K27me₃ recognition and transcriptional repression. Our structural and biochemical data showed that the BAH domain protein AIPP3 and the PHD proteins AIPP2 and PAIPP2 cooperate to read H3K27me₃ and unmodified H3K4 histone marks, respectively, in *Arabidopsis*. The BAH-PHD bivalent histone reader complex silences a substantial subset of H3K27me₃-enriched loci, including a number of development and stress response-related genes such as the RNA silencing effector gene *ARGONAUTE 5 (AGO5)* and We found that the BAH-PHD module associates with CPL2, a plant-specific Pol II carboxyl terminal domain (CTD) phosphatase, to form the BAH-PHD-CPL2 complex (BPC) for transcriptional repression. The BPC complex represses transcription through CPL2-mediated CTD dephosphorylation, thereby causing inhibition of Pol II release from the transcriptional start site. Our work reveals a mechanism coupling H3K27me₃ recognition with transcriptional repression through the alteration of Pol II phosphorylation states, thereby contributing to our understanding of the mechanism of H3K27me₃-dependent silencing.

KEYWORDS

H3K27me₃; CPL2; BAH; PHD; Pol II; H3K4; Flowering;

INTRODUCTION

In eukaryotic cells, the N-terminal histone tails undergo numerous posttranslational modifications (PTMs). Histone modification acts as a mark to specify the chromatin status as well as potential functional indications¹. The deposition, recognition and removal of specific histone PTMs are dynamically regulated by different proteins or protein complexes called ‘writer’, ‘reader’, and ‘eraser’ modules, respectively^{2, 3, 4}. The reader module can specifically recognize certain histone mark in both sequence- and modification-specific manners, and subsequently transmits the signal to downstream effectors. As a repressive epigenetic mark localized in euchromatin, deposition of trimethylation on histone H3 lysine 27 (H3K27me3) has been observed in many important functional genes⁵. The polycomb repressive complexes (PRCs), consisting of a different polycomb group (PcG) of proteins, have been shown to be involved in the deposition and downstream action of H3K27me3 mark⁶. Different PcG proteins associate to form two functionally distinct complexes, PRC1 and PRC2. The PRC1 complex has E3 ligase activity which has been shown to catalyze the monoubiquitination of histone H2A at lysine (H2Aub1), and the PRC2 complex catalyzes H3K27me2 and H3K27me3^{7, 8, 9, 10}. Three major models have been proposed to explain the mechanisms of PRC complex-mediated transcription repression⁸. For those bivalent promoters marked by both H3K27me3 and H3K4me3 marks, PcG complexes are believed to hold the poised Pol II at the transcription start site (TSS), resulting in the inhibition of Pol II release. Alternatively, PcG complexes can alter the chromatin environment by inducing chromatin condensation, thereby blocking the accessibility of chromatin remodeling complexes that is required for transcription activation^{11, 12, 13}. Third, the histone PTMs might directly prevent Pol II processivity during transcription elongation⁸. For example, studies in *Drosophila* have indicated that H3K27me3 could limit Pol II recruitment to gene promoters¹⁴. H2Aub1 has been implicated in restraining Pol II elongation^{15, 16}. However, the detailed mechanisms through which H3K27me3 reading is connected to transcriptional repression are not fully understood.

Histone mark recognition in plants is generally similar to that in animals but sometimes possesses plant-specific mark-reader pairs¹⁷. The PHD and BAH domains are two types of histone binding domains in eukaryotes^{18, 19}. The PHD finger has been reported to recognize methylated/unmethylated H3K4 marks and lysine acetylation marks¹⁹, and

the BAH domain can bind distinct histone marks, including H3K9me²⁰, H4K20me²¹, unmodified H3K4²², nucleosome core particle^{23, 24, 25, 26} and the more recently identified H3K27me^{27, 28, 29, 30}. In plants and animals, a large number of development and environmental response-related processes are subjected to H3K27me3-dependent regulation. Among them, flowering control has been a paradigmatic model for PRC complexes-mediated transcriptional repression in plants. The H3K27me3 dynamics in the flowering repressor gene *FLOWERING LOCUS C (FLC)* and the florigen gene *FLOWERING LOCUS T (FT)* play essential roles in the flowering time control³¹, and the H3K27me3 regulators influence flowering time in different ways^{27, 28, 29, 32, 33, 34, 35, 36, 37, 38, 39, 40}. Here, we demonstrated that the BAH domain-containing protein ASI1-IMMUNOPRECIPITATED PROTEIN 3 (AIPP3) and two PHD domain-containing proteins AIPP2/PARALOG OF AIPP2 (PAIPP2) could form a BAH-PHD module to read H3K27me3 and unmethylated H3K4, respectively, and coordinate in implementing transcriptional repression of hundreds of genes, particularly those development and stress-responsive genes in *Arabidopsis* such as the florigen gene *FT* and the RNA silencing effector gene *AGO5*. Moreover, our structural and biochemical studies further revealed the molecular basis for the specific recognition of these histone marks. We also revealed that the BAH-PHD module represses the release of Pol II from TSS regions by cooperating with CPL2, a known plant-specific Pol II CTD Ser5 phosphatase. Collectively, our findings reveal a coupling of the H3K27me3 recognition and downstream transcriptional repression through the BPC complex. This pathway may represent a mechanism of H3K27me3-mediated gene silencing.

RESULTS

BAH protein AIPP3 associates with PHD proteins and CPL2 to form a protein complex in *Arabidopsis*

Through a mass spectrometry analysis of the chromatin regulator Anti-silencing 1 (ASI1), we previously demonstrated that BAH domain-containing protein AIPP3 and PHD protein AIPP2 and CPL2 are associated with ASI1^{41,42}. AIPP2 is known to interact with AIPP3 and CPL2⁴¹. This association was further confirmed by immunoprecipitation assays coupled to a mass spectrometry analysis (IP-MS) of AIPP3, AIPP2 and CPL2 in which AIPP3, AIPP2 and CPL2 could be mutually co-purified with one another except for ASI1 (Fig. 1a and Supplementary Table 1). Interestingly, another PHD protein encoded by *AT5G16680* (Supplementary Fig. 1), the closest paralog of AIPP2 (hereafter referred to as PAIPP2) in *Arabidopsis*, was co-purified with AIPP3 and CPL2, and AIPP3 and CPL2 were also present in the IP-MS of PAIPP2 (Fig. 1a). The yeast two-hybrid (Y2H) and split luciferase assays indicated that PAIPP2 could also interact with AIPP3 and CPL2 but not with AIPP2 (Fig. 1b and Supplementary Fig. 2). Regarding the domain requirements for protein interactions, the Y2H results indicated that the BAH domain-containing N terminus of AIPP3 and the RBM motif-containing C terminus region of CPL2 are required for their interactions with AIPP2 and PAIPP2, respectively (Supplementary Fig. 3). AIPP2 and PAIPP2 were divided into three parts: the N terminus (N), PHD that is followed by a frequently associated polybasic region (PHD-PBR) and the C terminus (C) (Supplementary Fig. 4). The PHD-PBR part is indispensable for AIPP2/PAIPP2-AIPP3 interactions. Intriguingly, the PHD-PBR interaction with AIPP3 could be strengthened and inhibited by the N and C termini of AIPP2/PAIPP2, respectively (Supplementary Fig. 4), indicating the presence of an intramolecular regulation mode within AIPP2 and PAIPP2. The C termini of AIPP2 and PAIPP2 were fully responsible for the interaction with CPL2. Thus, we reasoned that the PHD proteins AIPP2 and PAIPP2 interact with AIPP3 and CPL2 independently to associate *in vivo* (Fig. 1c). Moreover, the gel filtration assay that was performed using epitope-tagged transgenic lines indicated that these four proteins co-eluted in the same fractions (Fig. 1d). Thus, these data support the notion that the BAH protein AIPP3 associates with two PHD proteins and CPL2 to form a protein complex *in vivo* (which are hereafter referred to as the BAH-PHD-CPL2 complex or the BPC complex).




Fig. 1 The BAH protein AIPP3 associates with PHD proteins and CPL2 to form a protein complex. **a** Mass spectrometry analysis of epitope-tagged *AIPP3*, *AIPP2*, *PAIPP2* and *CPL2* transgenic plants. **b** Y2H results showing the reciprocal interactions within the tested proteins. **c** A diagram showing the interaction network within AIPP3, AIPP2, PAIPP2 and CPL2. **d** Immunoblotting results of gel filtration assays.

The BAH-PHD-CPL2 complex represses flowering by inhibiting *FT* expression

By generating native promoter-driven β -glucuronidase (*GUS*) reporter transgenes, we observed that the BAH-PHD-CPL2 complex genes were ubiquitously expressed in *Arabidopsis* (Supplementary Fig. 5). To explore the biological function of the BAH-PHD-CPL2 complex, the null mutants of *PAIPP2* were generated using CRISPR/Cas9-mediated gene editing (Supplementary Fig. 6), and the morphology and flowering phenotypes of the generated *paipp2-1* mutant as well as the reported *aipp3-1*, *aipp2-1* and *cpl2-2* mutants were investigated⁴¹. The *aipp3-1*, *cpl2-2* and *aipp2-1/paipp2-1* displayed multiple developmental defects, such as a dwarfed size, small leaves and poor fertility (Fig. 2a). Instead, the *aipp2-1* and *paipp2-1* mutants only showed mild developmental defects. Moreover, the *aipp3-1* and *cpl2-2* mutants showed obvious earlier flowering during both long day (LD) and short day (SD) photoperiods in comparison with Col-0 (Fig. 2b, c). Although only mild early flowering was observed in the *aipp2-1* and *paipp2-1* single mutants, *aipp2-1/paipp2-1* showed similar early flowering compared with the *aipp3-1* and *cpl2-2* mutants (Fig. 2b, c), suggesting a redundancy in these two PHD proteins in relation to the flowering time control. To dissect their genetic relationship, we attempted to generate double, triple and quadruple mutants. Unfortun-

nately, we failed to obtain the *aipp3/aipp2/paipp2/cpl2* quadruple mutant due to a severe developmental defect. Compared to Col-0, the *aipp3-1/aipp2-1/paipp2-1*, *aipp3-1/cpl2-2* and *aipp2-1/paipp2-1/cpl2-2* mutants flowered earlier, and the time to flower was similar to the single mutants under the LD condition (Fig. 2d), suggesting that the BAH-PHD-CPL2 complex acts in the same genetic pathway in flowering time control.




Fig. 2 The BAH-PHD-CPL2 complex regulates plant development and flowering time. **a** The phenotypic developmental defects of *bpc* mutants. The morphological phenotypes of whole plants, inflorescence tissues, rosette and cauline leaves were shown. **b-c** The flowering phenotypes (**b**) and the numbers of rosette leaf (RLs) at flowering (**c**) in selected mutants during LD and SD photoperiods. **d** Comparison of the

number of RLs at flowering in selected mutants under the LD condition. **e** The relative mRNA levels of the *FLC*, *FT* and *SOC1* genes in the selected mutants. The mRNA levels were first normalized to *ACT2* and then to Col-0. The data are the means $\pm$ S.D. of three biological repeats. Significance analysis (t-test) was performed and * represent p value <0.01 . **f** The circadian accumulation of *FT* mRNA in selected mutants. The *FT* mRNA levels were normalized to *ACT2*. The data are the means $\pm$ S.D. of three biological repeats. The white and black boxes represent light and dark periods, respectively. **g-h** The numbers of RLs in the BAH-PHD-CPL2 complex mutants and their double mutants with *flc-3* (g) and *ft-10* (h) at flowering during the LD photoperiod.

In *Arabidopsis*, FLC, which is a MADS-box transcription factor that integrates multiple flowering signals, acts as a key floral repressor^{31, 43, 44}. FLC directly represses the expression of florigen gene *FT* and *SUPPRESSOR OF OVEREXPRESSION OF CO 1* (*SOC1*) by binding to the promoter of *SOC1* and the first intron of *FT*⁴⁵. The quantitative RT-PCR (RT-qPCR) indicated that the *FLC* RNA levels were reduced in the *aipp3-1*, *aipp2-1*, *cpl2-2*, *paipp2-1* and *aipp2-1/paipp2-1* mutants compared to Col-0 (Fig. 2e). Instead, the *FT* RNA levels were significantly increased in the *aipp3-1*, *cpl2-2* and *aipp2-1/paipp2-1* mutants but not in the *aipp2-1* and *paipp2-1* single mutants, further supporting the functional redundancy of AIPP2 and PAIPP2. By contrast, the *SOC1* RNA level was not significantly changed in all the tested mutants. It is known that the accumulation of FT protein has a circadian rhythm that peaks before dusk during LD photoperiod^{46, 47, 48}. We noticed that the loss of the BAH-PHD-CPL2 complex did not change the circadian rhythm of *FT* mRNA but led to a constitutive increase (Fig. 2f). To determine the genetic relationship between FLC, FT and the BAH-PHD-CPL2 complex in relation to the flowering time control, *flc-3* and *ft-10* mutants (in Col-0 background) were crossed with the tested mutants. Surprisingly, the *aipp3-1/flc-3*, *cpl2-2/flc-3*, *aipp2-1/paipp2-1/flc-3* mutants displayed similar early flowering compared with the with *aipp3-1*, *cpl2-2* and *aipp2-1/paipp2-1* mutants, but they flowered earlier than the *flc-3* single mutant (Fig. 2g). By contrast, the early flowering phenotypes of *aipp3-1*, *cpl2-2* and *aipp2-1/paipp2-1* mutants were completely rescued by *ft-10* (Fig. 2h), indicating that the BAH-PHD-CPL2 complex represses flowering primarily by repressing the expression of *FT*.

The AIPP3-BAH domain specifically recognizes the H3K27me3 mark

The chromatin-based mechanisms play vital roles in the flowering time control^{1, 49, 50}. The BAH domain is commonly identified as an epigenetic reader module of a particular histone mark¹⁸. To decipher the molecular function of the AIPP3-BAH domain (Fig. 3a), we firstly performed a histone peptide pull-down assay using purified AIPP3-BAH

protein. Among all the tested histone peptides, AIPP3-BAH could only be pulled down by H3K27me1, H3K27me2 and H3K27me3 peptides in a sequentially increasing manner (Fig. 3b). The H3K27me3-binding activity was further confirmed by isothermal titration calorimetry (ITC) binding analysis (Fig. 3c). Among all four tested histone methylation marks, H3K4me3, H3K9me2, H3K27me3 and H3K36me3, the AIPP3-BAH domain showed a significant preference for the H3K27me3 mark. Moreover, consistent with the histone pull-down result, AIPP3-BAH has a preference for the higher methylation level in H3K27. Thus, the evidence above fully demonstrated that AIPP3-BAH is a H3K27me3 reader module.

Structure of the AIPP3-BAH domain in complex with an H3K27me3 peptide

To gain molecular insight into the interaction between the AIPP3-BAH domain and H3K27me3, we successfully determined the crystal structure of the AIPP3-BAH domain in complex with an H3K27me3 peptide at a resolution of 2.4 Å (Fig. 3d and Supplementary Table 2). Overall, the AIPP3-BAH domain has a classic β -barrel structure that is similar to other reported BAH domain structures¹⁸. The H3K27me3 peptide has a good electron density map and a β -strand-like extended conformation (Supplementary Fig. 7a). The peptide is captured by a negatively charged cavity that is formed on the surface of the AIPP3-BAH domain with extensive hydrophobic and hydrophilic interactions (Fig. 3e). In detail, the three aromatic residues of the AIPP3-BAH domain, Tyr149, Trp170 and Try172, form an aromatic cage to accommodate the trimethyl-lysine of the H3K27me3 peptide (Fig. 3f), and it resembles other typical methyl-lysine-reading histone readers⁵¹. At the N-terminus, the H3A25 forms two main chain-main chain hydrogen bonds with the AIPP3 Val140 and Glu142 (Fig. 3f). In the middle, the AIPP3 residues Trp170, Tyr172, His198, and Asp200 form an extensive hydrogen-binding network with H3S28, which further fixes the imidazole ring of His198 in a special rotamer state that is parallel with the prolyl ring of H3P30. This parallel alignment of the two rings enables the CH- π and stacking interactions between




Fig. 3 Structural analysis of the AIPP3-BAH domain and PAIPP2-PHD finger in complex with H3K27me3 and unmodified H3K4 peptide, respectively. **a** The domain architecture of AIPP3 (lower panel) and the BAH domain construct used for structural and biochemical studies (upper panel). **b** The immunoblotting results showing the pull-down results for AIPP3-BAH and mutated proteins using different histone peptides. The 0.5% inputs serve as positive controls. **c** The ITC binding curves showing the AIPP3-BAH domain-binding preference for different histone methylation marks (left panel) and different H3K27 methylation levels (right panel). **d** The overall structure of the AIPP3-BAH domain-H3K27me3 peptide complex shown in ribbon with the AIPP3-BAH domain and the H3K27me3 peptide colored in green and yellow, respectively. **e** An electrostatic surface view of AIPP3-BAH domain with the

H3K27me3 peptide in the space-filling model showing the peptide fit inside a negatively charged surface cleft of the AIPP3-BAH domain. **f** The detailed interaction between AIPP3-BAH domain and the H3K27me3 peptide with the interacting residues highlighted in sticks and the hydrogen bonds highlighted in dashed yellow lines. **g-h** The ITC binding curves show that the mutations of essential residues for the H3P30 recognition (**g**) and aromatic cage residues (**h**) of the AIPP3 BAH domain significantly decrease the binding towards the H3K27me3 peptide. **i** The ITC binding curves showing the specific preference of the PAIPP2-PHD finger for the unmodified H3K4. **j** The overall modeled structure of the PAIPP2-PHD finger in complex with the unmodified H3 peptide with the PHD finger and peptide shown in ribbon and space-filling models. The modeled PAIPP2-PHD finger and the modeling template ATXR5 PHD finger are colored in cyan and silver, respectively, and they were superimposed together. **k** The detailed interaction between the PAIPP2-PHD finger (in cyan) and the unmodified H3K4 peptide with the interacting residues are highlighted in stick model and hydrogen bonds highlighted in dashed yellow lines. The corresponding residues from modeling template ATXR5 (in silver) were overlain and highlighted, showing that almost all the interacting residues are conserved. **l** The ITC binding curves showing that the D306K mutation of PAIPP2-PHD finger, which potentially disrupts H3R2 recognition, totally abolishes the unmodified H3K4 binding by PAIPP2.

AIPP3 His198 and H3P30, which resemble the recognition of H3K27me3 by the EBS and SHL BAH domains with conserved key residues (Supplementary Fig. 7b)^{27, 28}. In the C-terminus, H3G33 interacts with AIPP3 Val240 through a main chain-main chain hydrogen bond. To validate these structural observations, we performed a mutagenesis analysis on the key residues. The mutations D200A and H198A, which are essential for H3P30 recognition, in addition to Y149A, W170A, and Y172A, which represent important aromatic cage residues, showed reduced or no detectable binding to the H3K27me3 peptide (Fig. 3g, h).

PHD fingers of AIPP2 and PAIPP2 recognize the unmodified H3K4 mark

In addition to AIPP3-BAH, the PHD fingers of AIPP2 and PAIPP2 may also be involved in the recognition of histone marks. AIPP2 and PAIPP2 share a conserved PHD finger in their sequences (Supplementary Fig. 8a), and their sequences of PHD fingers are the typical signatures of unmodified H3 recognition PHD fingers¹⁹. We first detected their histone substrate binding properties by ITC method. Although the AIPP2-PHD finger does not behave well *in vitro* and tends to precipitate, we successfully detected the binding between the PAIPP2-PHD finger and the differentially methylated H3K4 peptides (Fig. 3i). The PAIPP2-PHD finger prefers to bind to unmethylated H3K4 and the binding affinities were clearly decreased when the methylation level of the H3K4 peptide was increased. Considering the high sequence

similarity between the two PHD fingers (Supplementary Fig. 8a), we believe that the AIPP2-PHD finger may possess the same binding preference on the unmodified H3K4. Moreover, both AIPP2 and PAIPP2 PHD fingers share approximately 35% sequence identity with the PHD finger of *Glycine max* ATXR5 (PDB ID: 5VAB) (Supplementary Fig. 8a), which recognizes the unmodified H3K4⁵². We modelled the AIPP2 and PAIPP2 PHD fingers using the ATXR5 PHD finger as a template to analyze the interactions with unmodified H3K4 (Fig. 3j and Supplementary Fig. 8b). In the modelled structure, we noticed that almost all the peptide-binding residues are conserved (Supplementary Fig. 8a). For instance, the Pro322 and Gly324 of PAIPP2, which correspond to the ATXR5 Pro60 and Gly62, respectively, are involved in the hydrogen-binding interaction with H3A1 (Fig. 3k). Similarly, the Ile300, Cys301, Ser302, and Asp306 of PAIPP2, which are equivalent to ATXR5 Leu39, Cys40, Asp41 and Asp44, respectively, contribute to the recognition of H3R2 (Fig. 3k). The Val23, Gly31, and Leu37 of PAIPP2, which correspond to the Val285, Gly292, and Leu298 of ATXR5, participate in the recognition of the unmodified H3K4 (Fig. 3k). To validate the modelling results, we performed mutagenesis experiment, too. As most of the peptide recognition is achieved by the hydrogen bonding interactions of the main chain of the PHD finger, we only mutated Asp306 of PAIPP2, which is involved in the recognition of H3R2 by its side chain. As shown in Fig. 3l, the D306K mutation of PAIPP2 PHD finger almost totally disrupts the peptide binding, further supporting our modelling data. The AIPP2 PHD finger possesses similar unmodified H3K4 recognition residues and interactions (Supplementary Fig. 8c).

The BAH-PHD-CPL2 complex represses the expression of the genes marked by H3K27me3 and low-methylated H3K4

H3K27me3 is usually considered a repressive mark that functions in transcriptional repression by recruiting recognition proteins or protein complexes^{4, 8}. To dissect the molecular function of the BAH-PHD-CPL2 complex, mRNA-Seq was performed using their null mutants as well as CLF and LHP1 mutants^{53, 54}, which are one of the known H3K27me3 methyltransferases and a reader protein in *Arabidopsis*, respectively. Using a two-fold change cutoff, we noticed that the numbers of up-regulated differentially expressed genes (up-DEGs) were far greater than the numbers of down-DEGs in the *aipp3-1*, *cpl2-2* and *aipp2-1/paipp2-1* mutants (Fig. 4a). Supporting the functional redundancy of AIPP2 and PAIPP2, very few DEGs were identified in *aipp2-1* and

paipp2-1 single mutants. It is noteworthy that most of the up-DEG genes regulated by the BAH-PHD-CPL2 complex display very low expression levels in the wild-type, which is a similar pattern to that the *clf* and *lhp1* mutants (Fig. 4b). Under a strict criterion, 155 genes were commonly up-regulated in the *bah-phd-cpl2* mutants (Fig. 4c, 4d, Supplementary Table 3), and a substantial subset are stress-responsive genes (Supplementary Fig. 9), such as *SEC31A* (*AT1G18830*), which participates in endoplasmic reticulum stress responses^{55, 56, 57}, and *AGO5* (*AT2G27880*), which is involved in antiviral RNA silencing^{58, 59, 60} and gametophyte development^{61, 62}. In fact, for most of the *aipp2-1/paipp2-1* up-DEGs, higher expression was also observed in the *aipp3-1* and *cpl2-2* mutants (Fig. 4d), indicating that the BAH-PHD-CPL2 complex targets a common subset of genes for repression.

We next explored the chromatin feature of the target genes by plotting the distributions of the histone marks on the up-DEGs using published histone modifications ChIP-seq data⁶³. As expected, the H3K27me3 mark was heavily enriched on the bodies of the common target genes (Fig. 4e). Consistent with the ITC result in which the higher methylation of H3K4 inhibits the binding of PAIPP2-PHD (Fig. 3i), the levels of active H3K4me3 mark were lower in the regions around the TSS. H3K36me3 deposition was also low on the whole gene bodies. The depositions of H3K9me2 and H3K27me1 were markedly lower on the target genes, suggesting that the BAH-PHD-CPL2 complex primarily targets euchromatic genes. The distribution patterns of H3K27me3 and H3K4me3 marks at BPC complex target genes were confirmed by ChIP-qPCR assays at representative target genes *AGO5*, *SUC5* and *FT* (Fig. 4f, g). We next investigated the impacts of BPC complex dysfunction on global histone mark levels. Interestingly, compared to the reduction of global H3K27me2/3 levels in *clf-81* mutant, the absence of the BPC complex did not result in obvious changes in global levels of H3K27me1/2/3 and H3K4me1/2/3 (Supplementary Fig. 10a). Consistent with this pattern, H3K27me3 deposition was not obviously changed between Col-0 and *bpc* mutants at selected target genes *AGO5*, *SUC5* and *FT* (Fig. 4g), whereas H3K4me3 levels are slightly increased (Fig. 4h). To further verify this observation, more target genes were selected. As shown in Supplementary Fig. 10b-d, nuclear run-on assay showed that *AT1G43570*, *AT3G59480* and *AT3G53650*, indicating that these three genes are subjected to BPC complex-mediated transcriptional repression. Interestingly, ChIP-qPCR results indicated that H3K27me3 levels were not obviously changed in *bpc*

mutants at *AT2G43570* and *AT3G59480*, consistent with our observation at *AGO5*, *SUC5* and *FT* (Fig. 4g). While, H3K27me3 deposition showed significant reduction at *AT3G53650*, implying that BPC dysfunction has substantial impacts on H3K27me3 deposition at this gene. Combined with these data, we speculate that, for most target genes, the BPC complex may serve as a surveillance system to prevent reactivation of H3K27me3-marked genes which are already silenced by PRC2, but in some specific genes, BPC is required for H3K27me3 through unknown mechanism.




Fig. 4. BAH-PHD-CPL2 complex represses the expression of H3K27me3-enriched genes. **a** The numbers of up- and down-DEGs identified in the selected mutants. A two-fold cutoff was used for the DEG criterion. **b** The BAH-PHD-CPL2 complex primarily represses low-expression genes. The gene expression levels of up-DEGs were classified into four groups according to their FPKM values in *Col-0* mRNA-seq. All the annotated genes serve as controls. **c** A Venn diagram showing the overlap of up-DEGs between

the selected mutants. **d** A heatmap showing the expression analysis in selected mutants. The genes in the upper and lower panels represent the common 155 up-DEGs and up-DEGs of *aipp2-1paipp2-1*, respectively. **e** The distribution of different histone marks in the respective up-DEGs of selected mutants. TTS, transcription termination site. **f** Snapshots showing the expression of the selected target genes. One representative replicate of mRNA-seq and histone ChIP-seq are shown. The adjacent genes with high H3K4me3/low H3K27me3 were also shown as parallel controls. **g-h** ChIP-qPCR showing the H3K27me3 (**g**) and H3K4me3 (**h**) density at the selected target genes. The ChIP signals were normalized to histone H3. The data are means $\pm$ S.D. of three technical repeats. One representative result of three biological replicates is shown. The lowercase letters represent the ChIP-qPCR examined regions as shown in (**f**).

The BAH-PHD-CPL2 complex reads the H3K27me3 at a genome-wide level

We next investigated the interplay between the BAH-PHD-CPL2 complex and the chromatin of the target genes by performing a ChIP assay in epitope-tagged transgenic plants. The ChIP-qPCR results indicated that AIPP3, AIPP2 and PAIPP3 were enriched in the selected target genes, particularly in the regions close to TSSs (Fig. 5a), but they were low in the adjacent high H3K4me3/low H3K27me3 genes. To confirm this point, an AIPP3 ChIP-seq was performed. As shown in Fig. 5b, AIPP3 specifically binds to the selected target genes. At the genome-wide level, AIPP3 binding peaks were enriched in the DEGs that were up-regulated in the *aipp3-1* mutant (Fig. 5c, d). Moreover, a pattern of high H3K27me3 and low H3K4me3 was clearly observed in AIPP3-bound genes (Fig. 5e). Next, we compared the genome-wide occupancy of AIPP3 with a published global analysis of the H3K27me3-marked regions¹. As shown in Fig. 5f, 455 loci, or approximately half of the AIPP3-enriched loci, significantly overlap with H3K27me3-enriched loci, indicating that a substantial part of the H3K27me3 loci were targeted by the AIPP3 complex. In *Arabidopsis*, EMBRYONIC FLOWER 1 (EMF1) is a plant-specific PRC1 component that is essential for conferring H3K27me3-dependent silencing at thousands of loci²⁹. Recently, EMF1 has been shown to interact with two H3K27me3 readers to form the BAH-EMF1 complex and implement silencing²⁹. We compared the AIPP3 loci with the published EMF1-bound loci and found that approximately 59% of AIPP3-enriched loci were also occupied by EMF1 (Fig. 5f). While, no EMF1 peptides were found in the BPC complex co-purified proteins (Supplementary Table 1). One possible explanation is that both EMF1 and AIPP3 associate with H3K27me3 mark independently, although the possibility of indirect association between these two reader proteins cannot be excluded.




Fig. 5. BAH-PHD-CPL2 complex binds to the H3K27me3-marked genes to repress the transcriptional initiation and elongation of Pol II. **a** ChIP-qPCR showing the fold enrichments of AIPP3, AIPP2 and PAIPP2 on the selected genes. The occupancy was first normalized to the internal control *AtSN1*, and shown are relative fold enrichments compared to Col-0. The data are means $\pm$ S.D. of three biological repeats. Significance analysis (t-test) was performed and * represent p value < 0.01, p value < 0.01. **b** Snapshots of AIPP3-4Myc ChIP-seq showing the distribution of the AIPP3 protein at the selected genes. Note: the ChIP-qPCR and ChIP-seq results of *AIPP3-4Myc* were from independent samples. **c-d** The diagrams showing AIPP3 occupancies on *aipp3-1* up-DEGs (b) and AIPP3-bound genes (c). **e** The diagrams showing the H3K27me3 (top) and H3K4me3 (bottom) density on AIPP3-bound genes. Random regions serve as negative controls. **f** A Venn diagram showing the overlap between AIPP3-bound genes, H3K27me3-enriched genes and EMF1-bound genes.

To decipher whether H3K27me3 binding is indispensable for flowering time control and transcription repression, the wild-type and mutated AIPP3 genomic DNA in which the crucial Tyr149, Trp170, and Try172 residues required for H3K27me3 binding were mutated into alanine was introduced into the *aipp3-1* mutant under the direction of the native promoter to generate *AIPP3*, *W170A* and *Y149A/W170A/Y172A* transgenic *Arabidopsis*. The early flowering of the *aipp3-1* mutation was rescued by the wild-type *AIPP3* transgene, but not by the *W170A* or *Y149A/W170A/Y172A* transgenes (Fig. 6a, 6b). Interestingly, compared to the comparable accumulation levels of AIPP3 and

456 W170A proteins in transgenic plants, Y149A/W170A/Y172A protein level was much
 457 lower (Fig. 6c), indicating that Y149A/W170A/Y172A mutation alters the stability of
 458 AIPP3 protein. Therefore, we used *AIPP3* and *W170A* transgenes in the following
 459 experiment. RT-qPCR results indicated that the W170A mutation could not recover the
 460 repressive state of the selected target genes (Fig. 6d), suggesting that the H3K27me3-
 461 binding activity is essential for the repression of flowering and gene expression. To test
 462 whether W170A mutation has an impact on AIPP3 binding at target genes, ChIP-qPCR
 463 assay was performed in Col-0, *AIPP3* and *W170A* transgenic plants. As shown in Fig.
 464 6e, compared to the significant enrichment at selected target genes in AIPP3 transgene,
 465 AIPP3 binding was disrupted by the W170A mutation. Considering the fact that W170
 466 is essential for H3K27me3 binding activity (Fig. 3h), this result strengthens our
 467 conclusion that H3K27me3 binding activity is indispensable for AIPP3-mediated
 468 flowering time control and transcriptional repression. Interestingly, the mutations did
 469 not affect the AIPP3 interactions with AIPP2 and PAIPP2 (Supplementary Fig. 11),
 470 indicating that the BAH-PHD-CPL2 complex is not dissociated by the disabling of
 471 H3K27me3-binding.




Fig. 6. The H3K27me3-binding activity is essential for AIPP3-mediated repression of flowering and gene expression. **a**, the flowering phenotypes of the *aipp3-1* mutant for its complementary lines when transformed with the wild-type *AIPP3* genomic DNA, mutated *AIPP3* in which the key amino acids for H3K27me3-binding were mutated. For the mutated transgene, two randomly selected transgenes were used for the analysis. The plants were grown under LD. **b**, the column showing the numbers of rosette leaves (LDs) of different *AIPP3* transgene plants upon flowering under the LD condition. **c**, Western blotting result showing the accumulation levels of AIPP3 proteins in different transgenes. Ponceaus staining serves as protein loading controls. **d**, The relative mRNA levels of the selected target genes of the BAH-PHD-CPL2 complex in the wild-type, *aipp3-1* and different *AIPP3* transgene plants. Their levels are presented relative to Col-0. The data are the means $\pm$ S.D. of three biological repeats. Significance analysis (t-test) was performed and * represent p value < 0.01 . **e**, ChIP-qPCR resulting showing AIPP3 occupancy at the selected target genes in *AIPP3* and *W170A* transgenes. The data are the means $\pm$ S.D. of three biological repeats. Significance analysis (t-test) was performed and * represent p value < 0.01 . ns, no significance.

BAH-PHD-CPL2 couples the recognition of H3K27me3 and the repression of Pol II release

It has been well documented that differential phosphorylation on the Ser2 (Ser2P) and

Ser5 (Ser5P) of the Pol II CTD plays essential roles in the switches between distinct transcriptional stages⁶⁴. During transcription, Pol II is first assembled at the promoter region. After initiation, Pol II is phosphorylated at Ser 5 and Ser 2 and is then released from the proximal promoter region to engage in productive elongation^{64, 65}. CPL2 has been shown to dephosphorylate the CTD-Ser5-PO₄ of Pol II⁶⁶. Therefore, it is reasonable to hypothesize that the BAH- and PHD-mediated recognition of H3K27me₃ and unmodified H3K4 directly represses transcription through the CPL2-mediated dephosphorylation of Pol II. To confirm this hypothesis, we first determined the states of different forms of Pol II in the target genes. Recently, Zhu et al. revealed Pol II dynamics with single-nucleotide resolution in *Arabidopsis* using native elongating transcript sequencing (NET-seq)⁶⁷. Surprisingly, only a sharp peak of unphosphorylated Pol II signals was observed at the TSS region of the selected target genes, and both the Ser5P and Ser2P signals were quite low or even undetectable throughout the promoter-proximal regions and gene bodies (Fig. 7a). By contrast, in the H3K4me₃-enriched active expressing genes that were independent from the regulation of the BPC complex (Fig. 4f), the maximum signals for unphosphorylated Pol II were detected at approximately + 200 bp and sharply decreased to a very low level, and Ser5P and Ser2P Pol II peaked in the promoter-proximal regions but decreased to a mild level when entering the gene body regions (Fig. 7a). These results strongly support the idea that transcription initiation and subsequent elongation of target genes were repressed in the wild-type by the BAH-PHD-CPL2 complex. We then checked what happens to the Pol II occupancy when the complex is absent. Compared to the wild-type, higher accumulations of Ser5P-Pol II were observed at the selected target genes in *aipp3-1*, *cpl2-2* and *aipp2-1/paipp2-1* mutants (Fig. 7b), and the Pol II signals maintained high levels towards the 3' end of the selected genes. Similar to Ser5P-Pol II, the occupancy of total (unphosphorylated) and Ser2P-Pol II also displayed higher levels at selected target genes *AGO5*, *SUC5* and *FT* in the *bpc* mutants in comparison with Col-0 (Supplementary Fig. 12). In contrast, the occupancies of all types of Pol II at *AGO5* and *SUC5* downstream non-target genes were not significantly changed (Supplementary Fig. 12), demonstrating that *bpc* mutations led to reactivation of Pol II initiation specifically at BPC complex target genes and the initiated Pol II successfully switched to an elongating state in the *bpc* mutants. These evidence, combined with the known knowledge that CPL2 is a Ser5P-Pol II phosphatase, prompts us to hypothesize that the BPC complex represses gene expression by connecting the BAH-PHD module-

mediated recognition of H3K27me3/unmodified H3K4 and the CPL2-mediated dephosphorylation of Pol II CTD Ser5-PO4.




Fig. 7 BPC complex directly connects H3K27me3 recognition with transcriptional repression. **a** Snapshots showing the occupancies of different Pol II forms on selected target genes using a published Pol II NET-seq database. The red and blue lines indicate the Pol II signals in the plus and minus strands, respectively. The dashed red boxes indicate the occupancies of Pol II on the selected target genes. The black arrows indicate the transcriptional direction. **b** The ChIP-qPCR validation of the occupancy of Ser5P-Pol II on selected genes in different mutants. The lowercase letters the examined regions (the same below). The occupancy was normalized to the *ACT7*. The Data are the means $\pm$ S.D. of three biological repeats. Significance analysis (t-test) was performed and * represent p value < 0.01 (the same below). ns, no significance. **c** Nuclear

Run-On analysis showing the relative Pol II transcription rate at the selected target genes in Col-0 and *bpc* mutants with or without FLA treatment (mock). *AT2G27900* and *AT1G71900* genes serve as control genes. The relative transcription rate was normalized to 18S rRNA. **d** CPL2 ChIP-qPCR results showing the relative occupancy of CPL2 at selected target genes in the presence and absence of AIPP3 and AIPP2/PAIPP2. The occupancy was first normalized to AtSN1 and then normalized to Col-0. The Data are the means $\pm$ S.D. of three biological repeats.

To support the above hypothesis, flavopiridol (FLA) treatment assay was performed which can reduce the phosphorylation of Pol II CTD by inhibiting the activity of CDK kinases⁶⁷, and the nascent RNA levels of the selected target genes were measured by Nuclear Run-On assay. The results indicated that the nascent RNA levels of *AGO5* and *SUC5*, but not that of the downstream non-target genes, were dramatically increased in the *bpc* mutants compared to Col-0 in mock condition. This evidence strongly supports our conclusion of transcriptional repression implemented by the BPC complex. In contrast, the nascent RNA levels of both *AGO5*, *SUC5* and the downstream non-target genes were greatly reduced by FLA treatment, and no significant changes were observed between different genotypes (Fig. 7c), indicating that inhibition of Pol II CTD phosphorylation dramatically repressed the transcription reactivation caused by the malfunctions of the BPC complex. Considering the likelihood of general effects of FLA treatment on transcriptional at a global level, to strengthen our hypothesis, the chromatin binding of CPL2 was compared in the presence or absence of AIPP3 and PHD proteins. To this end, *CPL2-GFP* transgene was crossed into *aipp3-1* and *aipp2-1/paipp2-1* mutants and CPL2 ChIP-qPCR assay was performed in *CPL2-GFP/Col-0*, *CPL2-GFP/aipp3-1* and *CPL2-GFP/aipp2-1/paipp2-1* plants. The result indicated that CPL2 has significant binding at selected target genes in wild-type background, whereas this binding was completely abolished in the *aipp3-1* and *aipp2-1/paipp2-1* mutants (Fig. 7d). This result provides a link between chromatin marks and CPL2-mediated dephosphorylation of Pol II, in which the chromatin localization of CPL2 at BPC complex target genes largely depends on the recognition of H3K27me3/H3K4me0 marks by BAH-PHD proteins. Based on these evidences, we proposed a working model of the transcriptional repression conferred by the BPC complex (Fig. 8). In this model, the BAH-PHD bivalent histone reader recognizes H3K27me3 and unmodified H3K4 marks and recruits CPL2 to dephosphorylate the Ser5P of Pol II CTD, resulting in the inhibition of Pol II release from a transcriptional initiation state to elongation. When the BPC complex is absent, active H3K4me3 mark is deposited at BPC target genes.

The Ser5 and Ser2 residues of Pol II CTD are sequentially phosphorylated, leading to transcriptional reactivation of BPC target genes.




Fig.8 A working model of the BPC complex-mediated transcription repression. When BPC is present, AIPP3-BAH and AIPP2/PAIPP2-PHD motifs recognize H3K27me3, which is deposited by PRC2 complex, and unmodified H3K4 around the TSS, respectively. Then, the BAH-PHD histone reader module recruit CPL2 to dephosphorylate Pol II at the 5th Ser of CTD, thereby repressing the transcriptional initiation and subsequent elongation of Pol II. When BPC is absent, active H3K4me3 mark is deposited. Pol II CTD Ser2 and Ser5 residues can be phosphorylated sequentially, leading to release of Pol II from initiation to elongation state.

DISCUSSION

BAH and PHD proteins form a bivalent histone reader complex for H3K27me3 and H3K4me0 marks

Recently, we structurally described several plant H3K27me3-reading BAH domain proteins, such as EBS, SHL and AIPP3 in this research^{27, 28}, which has encouraged us to characterize the H3K27me3-reading BAH domains. The recognitions of H3K27me3 by all these BAH domains depend on the aromatic cage to recognize the methyl-lysine, and a histidine and an aspartic acid residue to specifically interact with H3P30 for sequence specificity. Using the aromatic cage and the specific His and Asp residues as a criterion, we identified a subfamily of BAH domain proteins that can potentially recognize the H3K27me3 mark, which are widely distributed in plants, fungi, and animals (Supplementary Fig. 13). Interestingly, the predicted H3K27me3-binding BAH domain-containing protein human BAHD1 was reported to be an H3K27me3 reader³⁰,

which further supported our prediction. Therefore, we believe that the aromatic cage and conserved His and Asp residues are the key features of H3K27me3-recognition BAH domains.

Different histone marks do not function in a totally independent manner. Instead, they engage in communications and cooperation with each other. The cooperation between different histone marks can be a combinatorial reading to enhance binding or be mutually exclusive to balance the binding between different marks, which occurs at both the single protein level and the multiple protein complex level⁶⁸. Recently, we reported that the two flowering regulators EBS and SHL could dynamically recognize the antagonist histone marks H3K27me3 and H3K4me3 at the single protein level to regulate floral phase transitions^{27, 28}. Here, AIPP3 and AIPP2/PAIPP2 could recognize H3K27me3 and H3K4me0, respectively, and form a bivalent histone reader complex. This observation is consistent with our functional data showing that the BAH-PHD-CPL2 complex colocalizes with higher H3K27me3 and lower H3K4me3 marked chromatin regions. The BAH domain of AIPP3 is responsible for the colocalization of the complex with H3K27me3. The binding of unmodified H3K4 by the PHD finger of AIPP2/PAIPP2 may have two roles. First, the binding of unmodified H3K4 may prevent the binding towards methylated H3K4 to make sure the complex is targeted to the gene repressive H3K4me3 depletion region. Second, the binding of H3K4me0 together with the H3K27me3 binding by AIPP3 may combine to enhance the overall binding of the complex towards a certain chromatin region. Therefore, the proper targeting of the BAH-PHD-CPL2 complex relies on the crosstalk between the H3K4me0 and H3K27me3 marks.

Coupling H3K27me3 recognition and transcription repression through the CPL2-mediated dephosphorylation of Pol II

RNA Pol II-dependent transcription is a stepwise process involving the formation of the preinitiation complex (PIC), initiation, elongation and polyadenylation/termination stages. Each of the stages is associated with a distinct pattern of CTD phosphorylation^{8, 65}. The transcriptional machinery is first recruited to the promoter regions. Once incorporated into the PIC, the mediator stimulates cyclin-dependent kinase to phosphorylate serine 5 of the CTD heptad repeat, and Ser5P helps in the release of Pol II from the PIC complex, thereby allowing Pol II to escape the promoter and the

subsequent initiation of transcription. Ser5P is retained during the first several hundred nucleotides, in preparation for productive elongation⁶⁹.

Here, on the selected target genes *AGO5* and *SUC5*, unphosphorylated Pol II is restricted in the TSS region, whereas Ser5P and Ser2P were below detectable levels (Fig. 7a), indicating that Ser5P-dependent transcription initiation is inhibited, possibly by CPL2, in the presence of the BPC complex. Consistent with this notion, the density of the Ser5P Pol II was significantly increased, and it peaked within the first several hundred nucleotides in the BPC complex mutants (Fig. 7b), indicating that the Ser5P-dependent transcription initiation was derepressed. In the canonical model of H3K27me3-mediated transcription repression, the recognition of H3K27me3 recruits the PcG proteins in the PRC1 complex to impose the monoubiquitination of H2A, which represses transcription through three possible mechanisms, as mentioned in the introduction⁸. Consistent with this model, a recent report showed that the mutations in H3K27me3 reader proteins LHP1, EBS and SHL in *Arabidopsis* led to a reduction in H2Aub1²⁹. We found that the H2Aub1 levels were reduced in the *lhp1-3* mutant but were not affected in the *aipp3-1/cpl2-2*, *aipp3-1/aipp2-1/paipp2-1* and *aipp2-1/paipp2-1/cpl2-2* mutants (Supplementary Fig. 10a). This observation is consistent with our hypothesis that the BPC complex represses transcription mainly through the inhibition of Ser5P-dependent transcription initiation. While, we cannot rule out the possibility that BPC the complex has direct/indirect interaction with H2Aub1 at specific target genes. This study unveiled a direct connection between H3K27me3 recognition and transcription repression; the BAH-PHD module recognizes H3K27me3/H3K4me0 and recruits CPL2 to dephosphorylate Pol II CTD, resulting in the failure of Pol II to enter into the initiation and elongation form. Although Pol II CTD phosphatases are conserved in eukaryotes, no evidence shows that histone modifiers manipulate CTD phosphatases, including CPL2 analogies in other systems, to affect transcription. Regarding histone modifiers, several studies have reported that histone PTM modifiers can affect transcription via modulating Pol II CTD phosphorylation state^{70, 71}. JMJD3, a H3K27me3 demethylase in human, has been shown to directly interact with CTD-Ser2P to affect gene expression⁷². Knocking down JMJD3 or JHDM1D, a H3K27me1/2 demethylase, reduces the enrichment of Pol II CTD-Ser2P at specific genes in human promyelocytic leukemia cells⁷³. In addition to histone methylation, other histone PTMs including histone ubiquitylation and phosphorylation, are also associated with Pol II

CTD phosphorylation-dependent transcriptional elongation⁷¹. For example, knocking down histone ubiquitylation modifiers has been shown to affect CTD-Ser2 phosphorylation⁷¹. The phosphorylation of histone H3 on S10 and S28 has been reported to be associated with phosphorylated Pol II during transcriptional activation in humans and *Drosophila*^{74, 75}. These studies support a notion that modulation of Pol II CTD phosphorylation represents an important regulatory mechanism adopted by chromatin regulators to regulate gene expression. Our finding that the BPC complex reading histone information and conferring transcriptional repression through CPL2 phosphatase-mediated modulation of Pol II CTD phosphorylation state provides a direct evidence to support this notion. Considering that CPL2 is a plant-specific Pol II phosphatase that bears a unique RBM domain, and that this domain is required for its interaction with PHD proteins (Fig. 1c), the BAH-PHD-CPL2 pathway may represent a newly evolved silencing pathway in plants. Our findings suggest a greater complexity and diversity of H3K27me3-mediated transcriptional regulation. In addition, it is well-known that H3K27me3-mediated silencing mechanisms participate in multiple biological processes in both plant and animals, particularly development and stress responsive genes. The obvious developmental defects (Fig. 2a) and reactivation of many development- and stress response-related genes (Supplementary Fig. 9) in *bpc* mutants imply that the BPC complex may play more important roles in plant development and stress responses in addition to flowering time control.

METHODS

Plant materials and growth conditions

All the plant seeds were sown and grown on 1/2 MS medium containing 1% sucrose. The seedlings were grown under a long- (16 h light/8 h dark) or short-day (8 h light/16 h dark) photoperiod at 23 °C. The T-DNA insertion mutants *aipp2-1*, *aipp3-1* and *cpl2-2* were described in our previous study⁴¹. The *paipp2-1* mutant was generated by CRISPR/Cas9-mediated mutagenesis in a Col-0 background (Supplementary Fig. 4). *clf-81* and *lhpl-3* have been described previously^{53, 54}. For the epitope-tagged transgenic expression of the *AIPP2*, *AIPP3*, *PAIPP2* and *CPL2* genes, the wild-type and mutated genomic DNA driven by their native promoters were cloned into binary vectors with different tags and then transformed into the corresponding mutants using the flowering dip method. T3 generation transgenic plants were used for analysis. FLA treatment assay was performed as previous report⁶⁷. In brief, two-week-old

seedlings were collected and incubated with 200 μ M FLA or mock solution (DMSO) overnight. The treated seedlings were subjected to total RNA extraction.

RT-qPCR and RNA-seq analysis

Total RNA was extracted from 2-week-old seedlings using Trizol reagent (Thermo), and cDNAs were synthesized using HiScript II Reverse Transcriptase (Vazyme). Quantitative PCR was performed using a CFX96 Touch Deep Well Real-Time PCR Detection System (Bio-Rad). Three biological replicates were created. The primers used in this study are listed in Supplementary Table 4. For the RNA-seq analysis, total RNAs were extracted from 12-day-old seedlings grown during long days using a RNeasy Plant Mini kit (Qiagen). Following RNA purification, reverse transcription and library construction, the libraries were quantified by TBS380, and a paired-end RNA sequencing library was performed with Illumina NovaSeq 6000 (2 $\times$ 150 bp read length). The raw paired-end reads were trimmed and subject to quality control with SeqPrep (<https://github.com/jstjohn/SeqPrep>) and Sickle (<https://github.com/najoshi/sickle>) using the default parameters.

IP and mass spectrometry analysis

Immunoprecipitation and mass spectrometry analyses were performed as previously described⁷⁶. In brief, the total proteins were extracted from the inflorescence tissues with IP buffer (50 mM Tris-HCl, pH 7.6, 150 mM NaCl, 5 mM MgCl₂, 10% Glycerol, 0.1% NP-40, 0.5 mM DTT, and protease inhibitor cocktail) and then precipitated with anti-Flag (Sigma-Aldrich) or anti-Myc (Millipore) antibodies for 2 h at 4 °C. The precipitated protein mixtures were subjected to MS analysis.

Protein interaction analysis and gel filtration assays

For the Y2H assays, the full-length and truncated coding sequences of *AIPP2*, *AIPP3*, *CPL2* and *PAIPP2* were cloned into the pGADT7 and pGBKT7 vectors to generate AD and BD constructs. After the transformation, the yeast cultures were spotted onto SD plates lacking Trp and Leu (-LW) or lacking Trp, Leu and His (-LWH) and incubated at 30 °C for 3 d.

Gel filtration assays were performed as described previously⁷⁶. In brief, the total proteins were extracted from 4 g of seedling tissues expressing *AIPP3-FLAG*, *AIPP2-*

4MYC, *PAIPP2-3FLAG* and *CPL2-4MYC* with IP buffer and then loaded on to a Superdex 200 10/300GL column (GE Healthcare). The eluted fractions were collected in 96-well plates and the target proteins were detected by standard western blotting.

Protein expression and purification

The sequence containing the AIPP3-BAH domain (residues 112-279) was constructed into a self-modified pMal-p2X vector to fuse a hexahistidine tag plus a maltose-binding protein (MBP) tag to the N-terminus of the target protein. The plasmids were transformed into *Escherichia coli* strain BL21 (DE3) RIL (Stratagene). Expression was induced at 16 °C overnight with 0.2 mM of IPTG. The recombinant proteins were purified with a pre-packaged HisTrap FF column (GE Healthcare). The His-MBP tags were cleaved by TEV protease overnight and removed by flowing through a HisTrap FF column (GE Healthcare) again. The target protein was further purified using a Heparin column (GE Healthcare) and a Superdex G200 column (GE Healthcare). All the AIPP3-BAH mutations were generated by standard PCR-based mutagenesis procedure. The mutations of AIPP3-BAH and truncated AIPP2/PAIPP2 fragments were purified using the same protocols as those used for the wild-type AIPP3-BAH. For the GST-AIPP3-BAH proteins used in histone peptide pull-down assays, the wild-type and mutated AIPP3-BAH proteins were purified with glutathione-Sepharose (GE Healthcare) and eluted with elution buffer (50 mM Tris-HCl pH 8, and 10 mM reduced glutathione). The peptides were purchased from GL Biochem or EpiCypher.

Histone peptide pull-down

Histone peptide pull-down was performed according to a previous report²⁷. In brief, 1.5 µg of biotinylated histone peptides were incubated with streptavidin beads (NEB) in binding buffer (50 mM Tris-HCl 8.0, 300 mM NaCl, 0.1% NP-40) for 1 h at 4 °C and then washed with binding buffer. A 1.5 µg quantity of AIPP3-BAH proteins was incubated with a peptide-bead mixture in 0.5 ml of binding buffer for 3 h at 4 °C and then washed with binding buffer five times. The protein-bead mixtures were subjected to immunoblotting using anti-GST antibody (Abmart).

Chromatin immunoprecipitation assays and ChIP-Seq analysis

ChIP assays were performed according to a reported procedure⁷⁷. In brief, 3 g of seedlings was harvested and fixed with 1% formaldehyde in the cross-linking buffer (0.4 M

sucrose, 10 mM Tris-HCl pH 8, 1 mM PMSF, and 1 mM EDTA). After nuclei isolation with isolation buffer (0.25 M sucrose, 15 mM PIPES pH 6.8, 5 mM MgCl₂, 60 mM KCl, 15 mM NaCl, 1 mM CaCl₂, 0.9% Triton X-100, and protease inhibitor cocktail) and the following centrifugation, nuclei were resuspended in 500 µl of lysis buffer (50 mM HEPES pH 7.5, 150 mM NaCl, 1 mM EDTA, 1 mM PMSF, 1% SDS, 0.1% Na deoxycholate, 1% Triton X-100, and protease inhibitor cocktail) and sonicated with a Bioruptor (Diagenode). The nuclei lysate was precipitated with anti-Flag (Sigma-Aldrich), anti-Myc (Millipore), anti-H3 (Abcam), anti-H3K27me3 (Millipore), Ser2P-Pol II (Abcam), Ser5P-Pol II (Abcam) and unphosphorylated Pol II (Abcam) antibodies overnight and incubated with Dynabeads (Thermo) for 2 h. The precipitated protein-DNA mixtures were washed and eluted with elution buffer (0.5% SDS and 0.1 M NaHCO₃) at room temperature. The DNA was recovered after reverse cross-linking and proteinase K treatment.

For ChIP-Seq analysis, clean reads were mapped to the *Arabidopsis thaliana* genome (TAIR10) by Bowtie2 (version 2.2.8) with default parameters⁷⁸. Enriched peaks were identified by MACS (version 1.4) with default parameters. We defined the region of a target gene as the range from 1 kb upstream of TSS to TTS. The target genes of each peak were annotated by annotatePeak function in ChIPseeker package. The visualization of the average read coverage over gene body and additional 1kb up- and downstream of the TSS and TES was performed by deepTools (version 2.4.1)⁷⁹.

Crystallization, data collection and structure determination

Crystallization screening was performed using the sitting drop vapor diffusion method at 4 °C. The sample was concentrated and mixed with peptide at a molar ratio of 1:4. All the crystals emerged in a solution of 0.1 M HEPES, pH 7.0, and 2.4 M ammonium sulfate. The crystals were cryo-protected in the reservoir solution supplemented with 20% glycerol and flash-cooled in liquid nitrogen for X-ray diffraction. All the diffraction data were collected at beamline BL19U1 of the National Center for Protein Sciences Shanghai (NCPSS) at the Shanghai Synchrotron Radiation Facility (SSRF). The data were processed with the HKL3000 program⁸⁰. The structure was determined by molecular replacement method as implemented in the Phenix program⁸¹ using the ZMET2 BAH domain (PDB ID: 4FT2) as the searching model²⁰. Model building and

structure refinement were performed using the Coot and Phenix programs, respectively^{81, 82}. The statistics of data collection and structure refinement are listed in Supplementary Table 2.

Isothermal titration calorimetry

All the binding experiments were performed on a Microcal PEAQ-ITC instrument (Malvern) at 25 °C. The proteins were dialyzed against a buffer consisting of 50 mM NaCl and 20 mM Tris-HCl, pH 7.5 overnight at 4 °C. The lyophilized peptides were dissolved into the dialysis buffer. The titration was performed using the standard protocol and the data were processed using the Origin 7.0 program.

Nuclear Run-On assay

Nuclear Run-On assay was performed as previous report with minor changes⁸³. In brief, after nuclei isolation with buffer (0.25 M sucrose, 15 mM PIPES pH 6.8, 5 mM MgCl₂, 60 mM KCl, 15 mM NaCl, 1 mM CaCl₂, 0.9% Triton X-100, and protease inhibitor cocktail), the nuclear pellet was resuspended with nuclei storage buffer (50 mM Tris-HCl pH 8, 0.1 mM EDTA, 5 mM MgCl₂ and 40% glycerol) and mixed with transcription buffer (10 mM Tris-HCl, pH 8, 2.5 mM MgCl₂, 150 mM KCl, 2 mM DTT, 80 units RNase inhibitor, 0.5 mM BrUTP, 1 mM ATP, 1 mM GTP, 1 mM CTP, 0.5 mM UTP and 0.2% sarkosyl). After incubation at 30 °C for 15 min, the run-on reaction was stopped by adding Trizol reagent (Thermo). The nuclear RNAs were extracted and treated with DNase. The DNA-free RNAs were then inoculated with 2 µg anti-BrdU antibody (abcam) at room temperature for 30 min and precipitated by Dynabeads (Thermo). After two times washing, RNAs were extracted using Trizol reagent (Thermo). cDNAs were synthesized using SuperScript IV Reverse Transcriptase (Thermo) and subjected to qPCR analysis.

Reporting summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

Coordinates and structure factors have been deposited in the RCSB Protein Data Bank with the accession code: 7CCE. The ChIP-Seq and mRNA-seq data have been deposited in the GEO with the accession codes GSE147981 and GSE157196, respectively.

Supplementary Table 1 is provided as a Supplementary Data file. All other data are available from the corresponding authors on request.

Author contributions

C.-G.D., J.D. and J.-K.Z. designed this study. Y.Z., J.Y., L.Z., Y.W., G.Z. and Si-Si Xie conducted the experiments. Y.Z., J.Y., L.Z., J.J., J.D., J.K.Z. and C.-G.D. analyzed the data. C.C. and P.L. performed bioinformatic analysis. J.D., J.K.Z. and C.-G.D. wrote the paper.

Competing interests

The authors declare no competing interests.

Acknowledgments

We thank Dr. Yuehui He for the seeds of *clf-81*, *lhp1-3* and *flc-3* mutants. We thank staff at beamline BL19U1 of the National Center for Protein Sciences Shanghai (NCPSS) at the Shanghai Synchrotron Radiation Facility (SSRF) for data collection. Dr. C.G.D. was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB27040203) and by the National Natural Science Foundation of China (31570155). Dr. J.K.Z. was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences(XDB2704). J.D. was supported by the National Natural Science Foundation of China (31770782), Shenzhen Science and Technology Program (KQTD20190929173906742), SUSTech (G02226301) and the Key Laboratory of Molecular Design for Plant Cell Factory of Guangdong Higher Education Institutes (2019KSYS006).

References

1. Kim SY, Lee J, Eshed-Williams L, Zilberman D, Sung ZR. EMF1 and PRC2 cooperate to repress key regulators of Arabidopsis development. *PLoS genetics* **8**, e1002512 (2012).
2. Strahl BD, Allis CD. The language of covalent histone modifications. *Nature* **403**, 41-45 (2000).
3. Campos EI, Reinberg D. Histones: annotating chromatin. *Annu Rev Genet* **43**,

- 875 559-599 (2009).
876
- 877 4. Liu C, Lu F, Cui X, Cao X. Histone methylation in higher plants. *Annual review*
878 *of plant biology* **61**, 395-420 (2010).
879
- 880 5. Chang YN, Zhu C, Jiang J, Zhang H, Zhu JK, Duan CG. Epigenetic regulation
881 in plant abiotic stress responses. *Journal of integrative plant biology*, (2019).
882
- 883 6. Kim DH, Sung S. Polycomb-mediated gene silencing in *Arabidopsis thaliana*.
884 *Molecules and cells* **37**, 841-850 (2014).
885
- 886 7. Simon JA, Kingston RE. Mechanisms of polycomb gene silencing: knowns and
887 unknowns. *Nature reviews Molecular cell biology* **10**, 697-708 (2009).
888
- 889 8. Aranda S, Mas G, Di Croce L. Regulation of gene transcription by Polycomb
890 proteins. *Science advances* **1**, e1500737 (2015).
891
- 892 9. Di Croce L, Helin K. Transcriptional regulation by Polycomb group proteins.
893 *Nature structural & molecular biology* **20**, 1147-1155 (2013).
894
- 895 10. Blackledge NP, Rose NR, Klose RJ. Targeting Polycomb systems to regulate
896 gene expression: modifications to a complex story. *Nature reviews Molecular*
897 *cell biology* **16**, 643-649 (2015).
898
- 899 11. Shao Z, *et al.* Stabilization of chromatin structure by PRC1, a Polycomb
900 complex. *Cell* **98**, 37-46 (1999).
901
- 902 12. Levine SS, Weiss A, Erdjument-Bromage H, Shao Z, Tempst P, Kingston RE.
903 The core of the polycomb repressive complex is compositionally and
904 functionally conserved in flies and humans. *Molecular and cellular biology* **22**,
905 6070-6078 (2002).
906
- 907 13. Francis NJ, Kingston RE, Woodcock CL. Chromatin compaction by a polycomb
908 group protein complex. *Science* **306**, 1574-1577 (2004).
909
- 910 14. Chopra VS, Hendrix DA, Core LJ, Tsui C, Lis JT, Levine M. The polycomb
911 group mutant *esc* leads to augmented levels of paused Pol II in the *Drosophila*
912 embryo. *Molecular cell* **42**, 837-844 (2011).
913
- 914 15. Zhou W, *et al.* Histone H2A monoubiquitination represses transcription by

- 915 inhibiting RNA polymerase II transcriptional elongation. *Molecular cell* **29**, 69-
916 80 (2008).
917
- 918 16. Stock JK, *et al.* Ring1-mediated ubiquitination of H2A restrains poised RNA
919 polymerase II at bivalent genes in mouse ES cells. *Nature cell biology* **9**, 1428-
920 1435 (2007).
921
- 922 17. Liu R, Li X, Chen W, Du J. Structure and mechanism of plant histone mark
923 readers. *Science China Life sciences* **61**, 170-177 (2018).
924
- 925 18. Yang N, Xu RM. Structure and function of the BAH domain in chromatin
926 biology. *Critical reviews in biochemistry and molecular biology* **48**, 211-221
927 (2013).
928
- 929 19. Li Y, Li H. Many keys to push: diversifying the 'readership' of plant
930 homeodomain fingers. *Acta biochimica et biophysica Sinica* **44**, 28-39 (2012).
931
- 932 20. Du J, *et al.* Dual binding of chromomethylase domains to H3K9me2-containing
933 nucleosomes directs DNA methylation in plants. *Cell* **151**, 167-180 (2012).
934
- 935 21. Kuo AJ, *et al.* The BAH domain of ORC1 links H4K20me2 to DNA replication
936 licensing and Meier-Gorlin syndrome. *Nature* **484**, 115-119 (2012).
937
- 938 22. Li S, *et al.* Structural Basis for the Unique Multivalent Readout of Unmodified
939 H3 Tail by Arabidopsis ORC1b BAH-PHD Cassette. *Structure* **24**, 486-494
940 (2016).
941
- 942 23. Armache KJ, Garlick JD, Canzio D, Narlikar GJ, Kingston RE. Structural basis
943 of silencing: Sir3 BAH domain in complex with a nucleosome at 3.0 Å
944 resolution. *Science* **334**, 977-982 (2011).
945
- 946 24. Yang D, *et al.* Nalpha-acetylated Sir3 stabilizes the conformation of a
947 nucleosome-binding loop in the BAH domain. *Nat Struct Mol Biol* **20**, 1116-
948 1118 (2013).
949
- 950 25. Arnaudo N, Fernandez IS, McLaughlin SH, Peak-Chew SY, Rhodes D, Martino
951 F. The N-terminal acetylation of Sir3 stabilizes its binding to the nucleosome
952 core particle. *Nat Struct Mol Biol* **20**, 1119-1121 (2013).
953
- 954 26. Wang F, *et al.* Heterochromatin protein Sir3 induces contacts between the
955 amino terminus of histone H4 and nucleosomal DNA. *Proc Natl Acad Sci U S*

- 956 *A* **110**, 8495-8500 (2013).
957
- 958 27. Yang Z, *et al.* EBS is a bivalent histone reader that regulates floral phase
959 transition in Arabidopsis. *Nature genetics* **50**, 1247-1253 (2018).
960
- 961 28. Qian S, *et al.* Dual recognition of H3K4me3 and H3K27me3 by a plant histone
962 reader SHL. *Nature communications* **9**, 2425 (2018).
963
- 964 29. Li Z, Fu X, Wang Y, Liu R, He Y. Polycomb-mediated gene silencing by the
965 BAH-EMF1 complex in plants. *Nature genetics* **50**, 1254-1261 (2018).
966
- 967 30. Zhao D, *et al.* The BAH domain of BAHD1 is a histone H3K27me3 reader.
968 *Protein Cell* **7**, 222-226 (2016).
969
- 970 31. Luo X, He Y. Experiencing winter for spring flowering - a molecular epigenetic
971 perspective on vernalization. *Journal of integrative plant biology*, (2019).
972
- 973 32. Noh B, *et al.* Divergent roles of a pair of homologous jumonji/zinc-finger-class
974 transcription factor proteins in the regulation of Arabidopsis flowering time. *The*
975 *Plant cell* **16**, 2601-2613 (2004).
976
- 977 33. Goodrich J, Puangsomlee P, Martin M, Long D, Meyerowitz EM, Coupland G.
978 A Polycomb-group gene regulates homeotic gene expression in Arabidopsis.
979 *Nature* **386**, 44-51 (1997).
980
- 981 34. Yan W, *et al.* Dynamic and spatial restriction of Polycomb activity by plant
982 histone demethylases. *Nature plants* **4**, 681-689 (2018).
983
- 984 35. Zheng S, *et al.* The Arabidopsis H3K27me3 demethylase JUMONJI 13 is a
985 temperature and photoperiod dependent flowering repressor. *Nature*
986 *communications* **10**, 1303 (2019).
987
- 988 36. Yang H, Howard M, Dean C. Physical coupling of activation and derepression
989 activities to maintain an active transcriptional state at FLC. *Proceedings of the*
990 *National Academy of Sciences of the United States of America* **113**, 9369-9374
991 (2016).
992
- 993 37. Gaudin V, *et al.* Mutations in LIKE HETEROCHROMATIN PROTEIN 1 affect
994 flowering time and plant architecture in Arabidopsis. *Development* **128**, 4847-
995 4858 (2001).
996

38. Gomez-Mena C, Pineiro M, Franco-Zorrilla JM, Salinas J, Coupland G, Martinez-Zapater JM. early bolting in short days: an Arabidopsis mutation that causes early flowering and partially suppresses the floral phenotype of leafy. *The Plant cell* **13**, 1011-1024 (2001).
39. Sung S, *et al.* Epigenetic maintenance of the vernalized state in Arabidopsis thaliana requires LIKE HETEROCHROMATIN PROTEIN 1. *Nature genetics* **38**, 706-710 (2006).
40. Mylne JS, *et al.* LHP1, the Arabidopsis homologue of HETEROCHROMATIN PROTEIN1, is required for epigenetic silencing of FLC. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 5012-5017 (2006).
41. Duan CG, *et al.* A protein complex regulates RNA processing of intronic heterochromatin-containing genes in Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* **114**, E7377-E7384 (2017).
42. Wang X, *et al.* RNA-binding protein regulates plant DNA methylation by controlling mRNA processing at the intronic heterochromatin-containing gene IBM1. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 15467-15472 (2013).
43. Michaels SD, Amasino RM. FLOWERING LOCUS C encodes a novel MADS domain protein that acts as a repressor of flowering. *The Plant cell* **11**, 949-956 (1999).
44. Sheldon CC, *et al.* The FLF MADS box gene: a repressor of flowering in Arabidopsis regulated by vernalization and methylation. *The Plant cell* **11**, 445-458 (1999).
45. Searle I, *et al.* The transcription factor FLC confers a flowering response to vernalization by repressing meristem competence and systemic signaling in Arabidopsis. *Genes & development* **20**, 898-912 (2006).
46. Suarez-Lopez P, Wheatley K, Robson F, Onouchi H, Valverde F, Coupland G. CONSTANS mediates between the circadian clock and the control of flowering in Arabidopsis. *Nature* **410**, 1116-1120 (2001).
47. Imaizumi T, Kay SA. Photoperiodic control of flowering: not only by coincidence. *Trends in plant science* **11**, 550-558 (2006).

- 1039 48. Turck F, Fornara F, Coupland G. Regulation and identity of florigen:
1040 FLOWERING LOCUS T moves center stage. *Annual review of plant biology*
1041 **59**, 573-594 (2008).
1042
- 1043 49. Dennis ES, Peacock WJ. Epigenetic regulation of flowering. *Current opinion in*
1044 *plant biology* **10**, 520-527 (2007).
1045
- 1046 50. Duan CG, Zhu JK, Cao X. Retrospective and perspective of plant epigenetics
1047 in China. *Journal of genetics and genomics = Yi chuan xue bao* **45**, 621-638
1048 (2018).
1049
- 1050 51. Patel DJ. A Structural Perspective on Readout of Epigenetic Histone and DNA
1051 Methylation Marks. *Cold Spring Harb Perspect Biol* **8**, a018754 (2016).
1052
- 1053 52. Bergamin E, *et al.* Molecular basis for the methylation specificity of ATXR5 for
1054 histone H3. *Nucleic acids research* **45**, 6375-6387 (2017).
1055
- 1056 53. Kim GT, Tsukaya H, Uchimiya H. The CURLY LEAF gene controls both
1057 division and elongation of cells during the expansion of the leaf blade in
1058 *Arabidopsis thaliana*. *Planta* **206**, 175-183 (1998).
1059
- 1060 54. Lindroth AM, *et al.* Dual histone H3 methylation marks at lysines 9 and 27
1061 required for interaction with CHROMOMETHYLASE3. *The EMBO journal*
1062 **23**, 4286-4296 (2004).
1063
- 1064 55. Takagi J, *et al.* MAIGO5 functions in protein export from Golgi-associated
1065 endoplasmic reticulum exit sites in *Arabidopsis*. *The Plant cell* **25**, 4658-4675
1066 (2013).
1067
- 1068 56. Pastor-Cantizano N, Bernat-Silvestre C, Marcote MJ, Aniento F. Loss of
1069 *Arabidopsis* p24 function affects ERD2 trafficking and Golgi structure, and
1070 activates the unfolded protein response. *Journal of cell science* **131**, (2018).
1071
- 1072 57. Gimeno-Ferrer F, *et al.* alpha2-COP is involved in early secretory traffic in
1073 *Arabidopsis* and is required for plant growth. *Journal of experimental botany*
1074 **68**, 391-401 (2017).
1075
- 1076 58. Brosseau C, Moffett P. Functional and Genetic Analysis Identify a Role for
1077 *Arabidopsis* ARGONAUTE5 in Antiviral RNA Silencing. *The Plant cell* **27**,
1078 1742-1754 (2015).
1079

- 1080 59. Garcia-Ruiz H, *et al.* Roles and programming of Arabidopsis ARGONAUTE
1081 proteins during Turnip mosaic virus infection. *PLoS pathogens* **11**, e1004755
1082 (2015).
1083
- 1084 60. Minoia S, *et al.* Specific argonautes selectively bind small RNAs derived from
1085 potato spindle tuber viroid and attenuate viroid accumulation in vivo. *Journal*
1086 *of virology* **88**, 11933-11945 (2014).
1087
- 1088 61. Tucker MR, Okada T, Hu Y, Scholefield A, Taylor JM, Koltunow AM. Somatic
1089 small RNA pathways promote the mitotic events of megagametogenesis during
1090 female reproductive development in Arabidopsis. *Development* **139**, 1399-1404
1091 (2012).
1092
- 1093 62. Borges F, Pereira PA, Slotkin RK, Martienssen RA, Becker JD. MicroRNA
1094 activity in the Arabidopsis male germline. *Journal of experimental botany* **62**,
1095 1611-1620 (2011).
1096
- 1097 63. Luo C, Sidote DJ, Zhang Y, Kerstetter RA, Michael TP, Lam E. Integrative
1098 analysis of chromatin states in Arabidopsis identified potential regulatory
1099 mechanisms for natural antisense transcript production. *The Plant journal : for*
1100 *cell and molecular biology* **73**, 77-90 (2013).
1101
- 1102 64. Harlen KM, Churchman LS. The code and beyond: transcription regulation by
1103 the RNA polymerase II carboxy-terminal domain. *Nature reviews Molecular*
1104 *cell biology* **18**, 263-273 (2017).
1105
- 1106 65. Hajheidari M, Koncz C, Eick D. Emerging roles for RNA polymerase II CTD
1107 in Arabidopsis. *Trends in plant science* **18**, 633-643 (2013).
1108
- 1109 66. Koiwa H, *et al.* Arabidopsis C-terminal domain phosphatase-like 1 and 2 are
1110 essential Ser-5-specific C-terminal domain phosphatases. *Proceedings of the*
1111 *National Academy of Sciences of the United States of America* **101**, 14539-
1112 14544 (2004).
1113
- 1114 67. Zhu J, Liu M, Liu X, Dong Z. RNA polymerase II activity revealed by GRO-
1115 seq and pNET-seq in Arabidopsis. *Nature plants* **4**, 1112-1123 (2018).
1116
- 1117 68. Du J, Patel DJ. Structural biology-based insights into combinatorial readout and
1118 crosstalk among epigenetic marks. *Biochim Biophys Acta* **1839**, 719-727
1119 (2014).
1120

- 1121 69. Ding Y, Avramova Z, Fromm M. Two distinct roles of ARABIDOPSIS
1122 HOMOLOG OF TRITHORAX1 (ATX1) at promoters and within transcribed
1123 regions of ATX1-regulated genes. *The Plant cell* **23**, 350-363 (2011).
1124
- 1125 70. Kooistra SM, Helin K. Molecular mechanisms and potential functions of
1126 histone demethylases. *Nature reviews Molecular cell biology* **13**, 297-311
1127 (2012).
1128
- 1129 71. Srivastava R, Ahn SH. Modifications of RNA polymerase II CTD: Connections
1130 to the histone code and cellular function. *Biotechnology advances* **33**, 856-872
1131 (2015).
1132
- 1133 72. Estaras C, Fueyo R, Akizu N, Beltran S, Martinez-Balbas MA. RNA
1134 polymerase II progression through H3K27me3-enriched gene bodies requires
1135 JMJD3 histone demethylase. *Molecular biology of the cell* **24**, 351-360 (2013).
1136
- 1137 73. Chen S, *et al.* The histone H3 Lys 27 demethylase JMJD3 regulates gene
1138 expression by impacting transcriptional elongation. *Genes & development* **26**,
1139 1364-1375 (2012).
1140
- 1141 74. Ivaldi MS, Karam CS, Corces VG. Phosphorylation of histone H3 at Ser10
1142 facilitates RNA polymerase II release from promoter-proximal pausing in
1143 *Drosophila*. *Genes & development* **21**, 2818-2831 (2007).
1144
- 1145 75. Rossetto D, Avvakumov N, Cote J. Histone phosphorylation: a chromatin
1146 modification involved in diverse nuclear events. *Epigenetics* **7**, 1098-1108
1147 (2012).
1148
- 1149 76. Duan CG, *et al.* A pair of transposon-derived proteins function in a histone
1150 acetyltransferase complex for active DNA demethylation. *Cell research* **27**,
1151 226-240 (2017).
1152
- 1153 77. Saleh A, Alvarez-Venegas R, Avramova Z. An efficient chromatin
1154 immunoprecipitation (ChIP) protocol for studying histone modifications in
1155 Arabidopsis plants. *Nature protocols* **3**, 1018-1025 (2008).
1156
- 1157 78. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nature*
1158 *methods* **9**, 357-359 (2012).
1159
- 1160 79. Ramirez F, *et al.* deepTools2: a next generation web server for deep-sequencing
1161 data analysis. *Nucleic acids research* **44**, W160-165 (2016).

1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199

80. Otwinowski Z, Minor W. Processing of X-ray diffraction data collected in oscillation mode. *Methods in enzymology* **276**, 307-326 (1997).
81. Adams PD, *et al.* PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta crystallographica Section D, Biological crystallography* **66**, 213-221 (2010).
82. Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. *Acta crystallographica Section D, Biological crystallography* **66**, 486-501 (2010).
83. Roberts TC, Hart JR, Kaikkonen MU, Weinberg MS, Vogt PK, Morris KV. Quantification of nascent transcription by bromouridine immunocapture nuclear run-on RT-qPCR. *Nature protocols* **10**, 1198-1211 (2015).
84. Tariq M, Saze H, Probst AV, Lichota J, Habu Y, Paszkowski J. Erasure of CpG methylation in Arabidopsis alters patterns of histone H3 methylation in heterochromatin. *Proceedings of the National Academy of Sciences of the United States of America* **100**, 8823-8827 (2003).

SUPPORTING INFORMATION

- Supplementary Fig. 1 PAIPP2 is the closest paralog of AIPP2 in *Arabidopsis*.
- Supplementary Fig. 2 Protein interactions revealed by split luciferase assay
- Supplementary Fig. 3 Domain requirements for the interactions between BAH-PHD-CPL2 complex proteins
- Supplementary Fig. 4 Domain requirement for protein interactions
- Supplementary Fig. 5 Expression analysis of the *GUS* reporters from BAH-PHD-CPL2 complex genes in seedlings and inflorescence tissues
- Supplementary Fig. 6 CRISPR/Cas9-mediated mutagenesis of *PAIPP2*
- Supplementary Fig. 7 Structural analysis of the AIPP3 BAH-H3K27me3 complex
- Supplementary Fig. 8 Structural analysis of the AIPP2 PHD-H3 modeled complex
- Supplementary Fig. 9 FDR analysis of commonly up-regulated genes in the mutants of the BAH-PHD-CPL2 complex
- Supplementary Fig. 10 The effects of BPC dysfunctions on the deposition of different histone marks
- Supplementary Fig. 11 The 170A and Y149A/W170A/Y172A mutations of AIPP3 did not affect its interaction with AIPP2 and PAIPP2
- Supplementary Fig. 12 Relative occupancy of unphosphorylated and Ser2P-Pol II at selected target genes
- Supplementary Fig. 13 A structure-based sequence alignment of potential H3K27me3 reading BAH domains from different species
- Supplementary Table 1. IP-MS analysis
- Supplementary Table 2. Data collection and refinement statistics
- Supplementary Table 3. Commonly up-regulated genes in BPC complex mutants
- Supplementary Table 4. Primers used in this study