

1 Trends of genetic changes uncovered by Env- and Eigen-GWAS in wheat and barley

2 Rajiv Sharma^a, James Cockram^b, Keith A. Gardner^b, Joanne Russell^c, Luke Ramsay^c, William TB
3 Thomas^c, Donal M. O'Sullivan^d, Wayne Powell^a, Ian J. Mackay^{a1}

4 ^aScotland's Rural College (SRUC), Kings Buildings, West Mains Road, Edinburgh, EH9 3JG,
5 United Kingdom. ^bThe John Bingham Laboratory, NIAB, 93 Lawrence Weaver Road, Cambridge,
6 CB3 0LE, United Kingdom; ^cThe James Hutton Institute, Invergowrie, Dundee, DD2 5DA, United
7 Kingdom; ^dSchool of Agriculture, Policy and Development, University of Reading, Reading, RG6
8 6AR, United Kingdom

9 ¹ To whom correspondence may be addressed. E-mail: Ian Mackay ian.mackay@sruc.ac.uk

10 Abstract

11 The process of crop breeding over the last century has delivered new varieties with increased
12 genetic gains, resulting in higher crop performance and yield. However in many cases, the
13 underlying alleles and genomic regions that have underpinned this success remain unknown. This
14 is due, in part, to the difficulty in generating sufficient phenotypic data on large numbers of
15 historical varieties to allow such analyses to be undertaken. Here we demonstrate the ability to
16 circumvent such bottlenecks by identifying genomic regions selected over 100 years of crop
17 breeding using the age of a variety as a surrogate for yield. Using 'environmental genome-wide
18 association scans' (EnvGWAS) on variety age in two of the world's most important crops, wheat
19 and barley, we found strong signals of selection across the genomes of our target crops.
20 EnvGWAS identified 16 genomic regions in barley and 10 in wheat with contrasting patterns
21 between spring and winter types of the two crops. To further examine changes in genome structure
22 in wheat and barley over the past century, we used the same genotypic data to derive eigenvectors
23 for deployment in EigenGWAS. This resulted in the detection of seven major chromosomal
24 introgressions that contributed to adaptation in wheat. The deployment of both EigenGWAS and
25 EnvGWAS based on variety age avoids costly phenotyping and will facilitate the identification of
26 genomic tracts that have been under selection during plant breeding in underutilized historical
27 cultivar collections. Our results not only demonstrate the potential of using historical cultivar
28 collections coupled with genomic data to identify chromosomal regions that have been under

29 selection but to also guide future plant breeding strategies to maximise the rate of genetic gain and
30 adaptation in crop improvement programs.

31 **Significance Statement**

32 100 years of plant breeding have greatly improved crop adaptation, resilience, and productivity.
33 Generating the trait data required for these studies is prohibitively expensive and can be
34 impossible on large historical traits. This study reports using variety age and eigenvectors of the
35 genomic relationship matrix as surrogate traits in GWAS to locate the genomic regions that have
36 undergone selection during varietal development in wheat and barley. In several cases these were
37 confirmed as associated with yield and other selected traits. The success and the simplicity of the
38 approach means it can easily be extended to other crops with a recent recorded history of plant
39 breeding and available genomic resources.

40 **Introduction**

41 In the last century, significant improvements in yield and quality have been reported in
42 almost all crop species as a result of plant breeding driven by market demand (1). However, the
43 growing demand for food, feed and fibre to meet the expanding global human population requires
44 an acceleration in the pace of crop genetic improvement (2). Identification of the genetic loci
45 responsible for these changes will help accelerate the genetic gains required to meet future food
46 security needs, via their incorporation in marker assisted selection breeding strategies (3). Over
47 the last decade, genome wide association studies (GWAS) has become a prominent method for
48 genetic analysis in plants (4). In crops, GWAS require trait data on large collections of varieties or
49 accessions, which is typically expensive to collect and can therefore result in underpowered
50 studies with relatively low numbers of lines (5, 6). An alternative is to exploit the availability of
51 historical data, such as that collected during varietal development programmes.

52 For almost every major crop, yield is the most important breeding target. Breeding
53 programmes invest large amounts of resources into realising the incremental genetic gains in yield
54 that are required for continual varietal improvement. Accordingly, the process of developing new
55 crop varieties involves rigorous screening in large multi-location and multi-environmental trials over

several years. Large historical phenotypic data sets from such trials have been successfully employed for GWAS in the past (7), and in several cases have identified the functional genes underlying the genetic control of the investigated traits (8–11). However, the availability of seed for variety collections with appropriate trait data is not common for many crops. Alternatively, seed of historical varieties may be available, but the associated trait data may be lost or disjointed. In both cases, the cost of collecting *de novo* trait data can be prohibitive. In many cases however, the release date, subsequently termed here ‘age’, of varieties is known. Given that in most crops, the breeding process has improved the genetic potential of key agronomic traits over time, variety age can be used as a surrogate measure of merit and mapped in GWAS. This approach, in which environmental or non-genetic variables are treated as traits in GWAS to map loci associated with those variables, has been termed EnvGWAS (12). For many crops, the predominant genetic change over time has been to increase yield (e.g.(13)), and the age of a variety may function directly as a surrogate for yield, although loci detected may also be associated with other temporal changes. EnvGWAS on variety age can also be regarded as a simple genome-wide test for genetic loci under directional selection, which may be subsequently associated with traits. This approach may also provide a way of identifying alleles associated with adaptation (14) which otherwise have been difficult to detect. Finally, EnvGWAS can be a cost effective strategy since it can access large pre-existing datasets but is not dependent on historical or *de novo* trait data.

A related approach requiring no trait data is EigenGWAS (15). Using genotypic data alone, the singular value decomposition of the genomic relationship matrix provides loadings (eigenvectors) for each variety on each eigenvalue of the matrix. For the largest eigenvalues, these loadings are then treated as independent traits for GWAS. Significant associations with any particular component highlight genomic regions or markers of greatest importance for that eigenvalue, and therefore the potential major drivers of population structure. Subsequent study of varieties differing in these regions may also be interpretable in terms of drivers of adaptation. EigenGWAS and EnvGWAS have recently been used to study diversity among maize landraces and identify lines and traits suitable for downstream analysis without large scale phenotyping (12).

83 In this study, we demonstrate for the first time the utility of treating variety age as a
 84 surrogate trait for crop productivity when combined with EnvGWAS and EigenGWAS to identify
 85 target regions and quantitative trait loci (QTL) underpinning genetic improvements in crop
 86 performance that have occurred during modern plant breeding. This is a powerful but cost effective
 87 method that does not require extensive trait data or complex software. We demonstrate the utility
 88 of these complementary approaches by: i) using EnvGWAS on variety age to identify loci
 89 responsible for genetic improvement in four complimentary datasets of modern winter and spring
 90 types of wheat (*Triticum aestivum*) and barley (*Hordeum vulgare*) from the United Kingdom (UK)
 91 and Brazil. ii) Validating the results from (i) by GWAS on subsets of these varieties for which
 92 historic yield data were also available. iii) Evaluating the temporal changes of allelic state at the loci
 93 identified. iv) Performing EigenGWAS on the same four datasets. EigenGWAS compliments
 94 EnvGWAS in that it too does not required trait data and may also identify genomic regions that
 95 have undergone selection. However, unlike EnvGWAS, it does not explicitly search for regions
 96 associated with variety age and is more likely to detect features associated with local adaptation,
 97 which may change little in frequency over time. As far as we are aware, no EnvGWAS analysis
 98 has been published in plants for which variety age has been used as a trait. The combination of
 99 EnvGWAS with EigenGWAS used here provides insights into the recent breeding history and
 100 population structure of two of the world's most important crops, and highlights the effectiveness
 101 and simplicity of these approaches to study recent selection history without the requirement for
 102 phenotype data.

103 **Results**

104 **Year of variety release as a surrogate measure for yield**

105 The Pearson correlations between historical yield data and age of variety were calculated for the
 106 subsets of 192 UK wheat and 197 UK barley varieties for which historical yield data were available
 107 (***SI Appendix, Fig. S1***). High correlations between yield and year of release (range 0.896 – 0.974)
 108 were found in both UK data sets. This confirms year of release could be used as a good measure
 109 of genetic progress in UK wheat and barley yield potential. No historical yield data for the Brazilian
 110 wheat panel were available.

111 **EnvGWAS for variety age**

112 *EnvGWAS wheat.* Using variety age for EnvGWAS in the UK winter wheat panel ($n=404$) identified
113 thirteen significant ($-\log_{10}(p) > 4.0$) genomic regions, of which four loci were found to be highly
114 significant ($-\log_{10}(p) > 6.0$), located on chromosomes 1A, 2A, 2D and 6A (**Fig. 1A, Table 1, SI**
115 **Appendix, Table S1**). For Brazilian spring wheat ($n=355$), three significant genetic loci were
116 detected, two on chromosome 2B (251 cM, 318 cM) and one on 5A (710 cM), none of which were
117 identified in the UK winter wheat panel (**Fig. 1B, Table 1, SI Appendix, Table S1**).

118 *EnvGWAS Barley.* We identified three highly significant genetic loci in the winter barley panel
119 ($n=297$), and seven in the spring barley panel ($n=406$) (**Table 1; Fig. 1C-D**); a summary of the
120 associated markers is listed in **SI Appendix, Table S2**. Two significant loci were identified in both
121 barley panels (chromosome 3H, ~68-70 cM; 5H, ~20 cM) (**Fig. 1** and **Table S2**). Subsequently,
122 EnvGWAS was performed on the combined winter and spring panels ($n=704$), identifying the same
123 four significant loci we identified in the spring panel alone (**SI Appendix, Fig. S2A, Table S2** and
124 **Table 1**). We repeated the analysis using seasonal growth habit ('spring' or 'winter' types) as a
125 covariate, without any major changes in results (**SI Appendix, Fig. S2B**). In addition, we
126 performed GWAS on seasonal growth habit itself, identifying three major genetic loci on the long
127 arms of chromosomes 1H, 4H and 5H (**SI Appendix, Fig. S2C**), corresponding to major flowering
128 time and vernalization genes known to be the major determinants of winter and spring seasonal
129 growth type (*PPD-H2* on chromosome 1H, *VRN-H2* on 4H and *VRN-H1* on 5H) (16, 17).
130 EnvGWAS for variety age was then repeated with these QTL as covariates (**SI Appendix, Fig.**
131 **S2D**). The most significant results mainly on chromosome 5H from the analyses with and without
132 covariates changed little. However, the magnitude of other significant peaks differed, such as the
133 locus on chromosome 1H.

134 **Validation of EnvGWAS based on trait analysis and a multi-founder experimental population**

135 To validate the EnvGWAS analyses, we performed GWAS on the subset of 192 UK winter wheat
136 varieties for which historical yield data were available together with EnvGWAS on variety age for
137 direct comparison of the results. In this subset, we found that GWAS for yield identified the same
138 genomic region on chromosome 1A (**Fig. S3A**) as was detected by EnvGWAS for variety age (**SI**

139 **Appendix, Fig. S3, Table S1**). This is the same region that we identified in EnvGWAS for variety
140 age in the complete set of 404 UK wheat varieties. Interestingly, while the chromosome 5A QTL
141 was detected with low-significance ($-\log_{10}(p) = 4.45$) by GWAS on yield, it was not identified using
142 EnvGWAS on variety age. In addition, EnvGWAS analysis of variety life-span also detected a locus
143 on chromosome 1B that was not detected in any other of our analyses.

144 Similarly, EnvGWAS on variety age and GWAS on yield was repeated using the subset of 197
145 winter and spring barley varieties for which historic yield data was available, detecting highly
146 significant hits ($-\log_{10}(p) > 4.0$) on chromosome 5H for variety age, variety life-span and yield,
147 using seasonal growth habit as a covariate (**SI Appendix, Fig. S4, Table S2**). Although not
148 identified in the larger panel of 703 varieties, analysis of our subset of 197 lines consistently
149 identified a highly significant genetic locus on the short arm of chromosome 3H for variety age,
150 variety life-span and yield. An additional peak was detected with EnvGWAS for variety life-span on
151 the long arm of chromosome 2H.

152 To further validate our EnvGWAS findings, we analysed data from a 16 founder wheat multiparent
153 advanced generation inter cross (MAGIC) population consisting of 550 recombinant inbred lines
154 generated by inter-crossing 16 wheat varieties released between 1935 to 2004 (18). We found that
155 the four major genomic regions previously identified by EnvGWAS of variety age on chromosomes
156 1A, 2A, 2D and 6A were also significant in the MAGIC population for several yield and grain related
157 traits, height, and yellow-rust resistance (**Table 2**). Further details of the 213 agronomic and
158 disease resistance traits analysed and the corresponding significance levels are listed in (**SI**
159 **Appendix, Table S3**).

160 **Allele-shift over time**

161 To illustrate the changes in allele frequency present in our variety collections over time, we
162 generated rocket plots (**SI Appendix, Fig. S5–S8**) for the major genomic regions identified by
163 EnvGWAS on variety age (**SI Appendix, Table S4–S7**). Different patterns and intensity of selection
164 were evident across chromosomal regions over time. For wheat, these fell into three broad
165 classes: (1) Late introduction of ‘modern’ alleles followed by a rapid increase in frequency (**SI**
166 **Appendix, Fig. S5A**), (2) retention of both ‘modern’ and ‘old’ alleles at similar frequency across

time (e.g. **SI Appendix S5E**), (3) relatively early introduction of the ‘modern’ allele, followed by its retention at low frequency (e.g. **SI Appendix S5F**). Details of the alleles-shifts examples are provided in the Supplementary Notes. In barley, the rocket plots illustrated both gradual and rapid shifts in allele frequency at the genomic regions identified by EnvGWAS on variety age (**SI Appendix, Fig. S5I-N**). For example, for the UK spring barley genetic locus on chromosome 7H (~8.8 Mbp), only one allele was present until 1992 (**SI Appendix, Fig. S5N and Table S6**), after which the ‘modern’ allele remained at low frequency, even among modern varieties. A genomic region on chromosome 5H which was identified separately in winter and spring barley displays a pattern where the ‘modern’ allele is introduced in 1986, after which both alleles are found at intermediate frequencies among the most recent varieties in winter barleys. However, modern spring barleys were predominantly of ‘modern’ allele type.

178 **EigenGWAS scans**

While EnvGWAS allowed us to use variety age to investigate the genomic regions underlying QTL for yield and adaptation, we hypothesised that the complementary method, EigenGWAS, would allow us to detect changes in larger scale structural variants in our target crop genomes over time. After determining the first ten PCs in each of our UK and Brazilian wheat populations (**SI Appendix, Table S8**), EigenGWAS detected numerous significant hits ($N=11567$ SNPs with $-\log_{10}(p) > 4.0$) (**Fig. 2 & SI Appendix, Table S9**). Seven genetic loci distributed on chromosomes 1A, 1B, 2B, 5B, 6A and 6B were found to be significant with multiple PCs, as well as within the Brazilian (spring) and UK (winter) panels (**Fig. 2**). These loci corresponded to major chromosomal introgressions from related cereal species into wheat (**SI Appendix, Table S9**). For instance, the 1B locus co-locates with the chromosome 1B/1R introgression from rye (*Secale cereale*), which is known to regulate multiple traits including disease resistance and yield (19, 20). We identified an additional seventeen putative introgressions that were supported by a recent introgression survey by (21), along with another 58 novel putative introgressions (**SI Appendix, Table S9**). Among these novel putative introgressions were regions on chromosome 5A, depicted in **Fig. 2** as 5A_2 and 5A_3, which displayed amongst the most significant hits across the UK and Brazilian wheat data sets and multiple PCs. Interestingly, two highly significant genomic regions (1A_2 and 5A_5)

identified by EigenGWAS on PC2 in winter wheat were also detected by GWAS on yield in the validation data set (**SI Appendix, Table S13**). In addition, three genomic regions (5B_2, 6A_1 and 7B_1) identified in the winter wheat EigenGWAS analysis were also detected in EnvGWAS on variety age, suggesting both approaches are not exclusively identifying different genomic regions (**SI Appendix, Table S13**).

In contrast to wheat, EigenGWAS in the winter and spring barley varieties did not detect any major loci with highly significant peaks across multiple PCs (**Fig. 3** and **SI Appendix, Table S8 & results in Table S11**). Although two genomic regions in winter (1H_3 and 4H_3) and three in spring barleys (2H_3, 3H_1 and 7H_1) were identified in at least three PCs. Nevertheless, peaks were also identified close to the locations of known genes controlling flowering time and height (**SI Appendix, Table S11**), e.g. the PC5 hit on chromosome 3H ~632 Mbp (explaining 2.46% of the variation) is near the semi-dwarfing gene *sdw1* in spring barley. Interestingly, one of the most significant hits in the spring barley panel (3H_1, identified using PC1 and explaining 6.91% of the variation) was also detected using EnvGWAS on variety age and by GWAS on yield (**SI Appendix, Table S14**). Given the location of this hit in a highly recombinogenic region of the barley genome, and that it was detected only in the spring barley panel, this may indicate a major locus under selection specific to spring barley breeding. No strong peak in winter barley was found for PC1, with the most significant peak obtained using PC6. As UK elite winter barley is more genetically diverse than UK elite spring barley, these results indicate that UK elite winter barley may be subjected to weaker selection pressures. Interestingly, hits on genomic regions (5H_2 and 7H_1) from the spring barley EigenGWAS analysis were also identified in GWAS analysis of seasonal growth-habit and variety age, highlighting the importance of these loci under selection (**SI Appendix, Table 14**).

Discussion

We demonstrate that use of variety age for EnvGWAS can detect regions of crop genomes under selection during breeding. In addition, we show variety age is a good proxy for yield, with the genetic loci identified for wheat validated in an independent experimental multi-founder population (18). Lastly, we showed that the genetic loci detected by EnvGWAS showed gradual, as well as

sharp, shifts in allele-frequency over time, indicating subtle changes at these loci by breeders, which are less discernible to detection using approaches such as partitioning the populations on age and searching for differences based on F_{st} .

It is perhaps not surprising that selection of loci varies between the UK winter and Brazilian spring wheat, given that the target agricultural environments and growth types are very different. Wheat yields in both Brazil and the UK have improved greatly over the years (13, 22). Our contrasting results in wheat indicate that different sets of genes have been selected over the years, and are likely involved in both yield component and local adaptation traits. Future efforts will shed more lights on the types of genes underpinning these loci, allowing changes in allelic diversity over the years to be investigated.

Our results for UK barley contrast with those for UK wheat. Firstly, more hits were associated with variety age in spring compared to winter barley, and secondly an identical peak on chromosome 5H (at ~19cM, ~7.5 Mbp) was identified in both panels (as well as in the combined spring and winter analysis). This is surprising as breeders rarely cross spring and winter barley, and since the breeding targets in the two pools differ (malting and largely animal feed, respectively). To further investigate this region, we tested the candidate SNPs against phenotypic data available from national trial data (**SI Appendix, Table S10**), finding it to be associated with several malting quality traits, powdery mildew resistance and yield in fungicide untreated trials. These findings suggest the potential importance of this region for breeding for disease resistance and end use quality. Interestingly this region on 5H houses a cluster of terpene synthases that have been implicated in fungal disease resistance in other species (23) and that potentially have been selected alongside direct targets such as *Mla* and *mlo* genes (24).

The detection of significant hits with EnvGWAS provides an opportunity to explore their relationship with yield and other agronomically important traits. Some hits coincide with previously published QTL in wheat and barley, for example the highly significant loci on wheat chromosomes 1A and 6A (25–27). Our EnvGWAS hits on chromosomes 1A and 2A also overlapped with the reduced diversity peaks identified in the recent analysis of the UK wheat pedigree by (28). Specifically, the

250 2A locus may correspond to a stripe rust resistance gene described by (29), as the peak markers
251 overlap. Interestingly a group of R genes *Lr37-Yr17-Sr38* (30) which were important sources of
252 resistance in the past also lie in this region and might be more plausible candidates, rising in
253 frequency before their resistance broke down. Similarly the highly significant genetic locus on the
254 short arm of barley chromosome 3H for variety age and yield found in the subset of 197 barley
255 lines corresponds to the genomic region associated with the a malting quality trait, hot water
256 extract, in UK spring barley that demonstrated a major change in allele frequency over the last
257 thirty years (31). In addition, the region identified on chromosome 3H (~68cM) for variety age in
258 winter barley in the larger dataset has been shown previously to be associated with yield
259 component traits (grain length and grain area) in European winter barley (32).

260 Similarly, in barley, the region identified on chromosome 2H (~65cM, ~621 Mbp) for variety life-
261 span has been shown previously to be associated with yield and yield component traits (32, 33)
262 and may correspond to the *OsBR1/D61* candidate genes reported previously that are associated
263 with yield traits in barley (32, 33).

264 This is interesting as old varieties, despite being less-productive than modern varieties, were under
265 cultivation for longer periods. It may however be noted that with the introduction of modern
266 breeding practices yield increases, but with drastic effects on variety life-span due to the more
267 frequent introduction of new varieties that outperform contemporary varieties. In wheat, EnvGWAS
268 on variety life-span also identified a hit on chromosome 1A that co-located with a hit for variety age.
269 This further indicates a direct relationship between variety age and variety life-span in wheat and
270 barley.

271 Using EigenGWAS, we detected major introgressions in the wheat varietal panels investigated,
272 with several of these found to be in common between the UK winter and Brazilian spring wheat
273 panels, indicating their wide use in breeding. (18), analysing the 16 founder MAGIC population we
274 used in our validation studies here, proposed a major role for multiple introgressions from wild
275 species in UK wheat breeding to date. In contrast, EigenGWAS results in barley provide no
276 evidence of a similar pattern of introgressions in either the winter or spring panels. Wheat and
277 barley breeding differ in their exploitation of genetic resources. In wheat, several alien-

278 introgressions from related species are known to have occurred (34). While wheat is an
279 allohexaploid and can support large tracts of non-recombining alien chromosome, this may not be
280 the case in diploid barley. However, examples of introgressions in barley from landraces and
281 spontaneous mutant lines for agronomically important genes have been reported, such as the
282 semi-dwarfing allele *sdw1d* from the variety Diamant and the disease resistance gene *mlo11* from
283 Ethiopian landraces (24, 35).

284 Interestingly, within the genomic region of 6A_1, detected by EigenGWAS in wheat (a non-
285 recombining peri-centromeric region) lies the gene *TaGW2* (36) which influences grain-weight and
286 protein content traits that further suggest that the present approach is very effective in discovering
287 genomic regions undergoing selection for yield. Another interesting finding is that the semi-
288 dwarfing *Rht2* gene in wheat (chromosome 2D) was not detected despite its importance in the
289 breeding history of the crop. This could be due to population structure control of the analyses. In
290 the case of *Rht2*, it is noteworthy that GWAS on a panel of French, German and UK lines failed to
291 detect an effect on yield or height unless a locus specific marker was used (37, 38), suggesting
292 weak LD and low marker coverage on the 4D chromosome as the cause of failure here too.

293 **Conclusion**

294 Breeding has resulted in considerable and sustained genetic improvement of wheat and barley in
295 recent decades, and our results identify at least some of the major loci that have contributed, and
296 are still contributing, to these improvements. Using EnvGWAS, we demonstrate the utility of
297 analysing variety age as a surrogate for traits selected by breeders to detect the genetic loci under
298 selection over time, and to assess the temporal changes in their respective allele frequencies. For
299 UK cereals, trends over time suggest that these loci are likely QTL for yield or yield components.
300 While the resolution of this study in the non-recombining peri-centromeric region is insufficient to
301 definitively associate known QTLs with the loci we have found, several such QTLs were found.
302 EigenGWAS on the same data proved a simple method of detecting contrasting features of
303 genome organisation in wheat and barley, and in some cases these too could be related to traits.
304 We advocate the use of variety age as a surrogate trait, and the use of EnvGWAS and
305 EigenGWAS to identify the genetic loci under selection that have underpinned the productivity

gains made via breeding. These extensions to GWAS that exploit historical datasets are useful additions to the analysis toolbox of crop quantitative genetics.

Materials and methods

Germplasm, age and trait data.

For both wheat and barley, we selected two panels of varieties representing national list entries and some older varieties from the UK (404 winter wheat; 297 winter and 406 spring barleys) and Brazil (355 spring wheat) (**Table S12**). The Brazilian spring wheat panel included entries released between 1922 to 2013. Year of varietal release and trait data were obtained from (39). The UK wheat panel consists of winter wheat varieties that were either registered or in use from 1916 to 2010. The winter and spring barley panels consisted of varieties grown in the UK from 1960 to 2016. Only two-rowed spike morphology types were included and all hybrid varieties were excluded. Variety age for UK germplasm was determined from the year of entry into national list trials or from the first reported year of trial data and was manually checked across different local data and published sources ((13); <https://ahdb.org.uk/ri> & <https://www.gov.uk/government/publications/plant-varieties-and-seeds-gazette-2020> <https://www.niab.com/services/seed-certification/botanical-descriptions-varieties>) with unresolvable ambiguities removed, reducing the UK wheat panel from 450 to 404 varieties. Following (13), only varieties with either three years trials data or equivalently which were known to be successful in national list trials were included in the dataset. In addition to variety age, we computed life-span of UK varieties as the difference between the last and first year in national trials plus one. This is usually equally to the total number of years each variety remained in trial, though with some rare breaks in the testing sequence over years. Grain yield data for the UK wheat and barley panels were sourced from (13).

Genotyping

Genotypic data were sourced from NIAB (<https://www.niab.com/research/agricultural-crop-research/resources>) and JHI (<http://www.barleyhub.org/projects/impromalt/>) by permission through WAGTAIL and IMPROMALT projects.

333 For wheat, 14654 SNPs derived from genotyping with the 90K Illumina iSelect SNP array (Wang et
334 al. 2014) generated within the Biotechnology and Biological Sciences Research Council grant
335 BB/J002542/1 were sourced with permission from NIAB, and available at
336 <https://www.niab.com/research/agricultural-crop-research/resources>. For barley, 43799 SNPs
337 genotyped using the 50K Illumina iSelect array (40) were sourced from (31). Genetic maps for
338 wheat (41) and barley (31, 40) were previously described. The physical map locations of wheat
339 and barley SNPs were retrieved from (42) and (40), respectively. SNPs with a minor allele
340 frequency <5%, missing values <10% and heterozygosity >10% were removed, leaving 12656
341 wheat SNPs and 25562 barley SNPs for downstream analyses.

342 **EnvGWAS and EigenGWAS analysis**

343 EnvGWAS and EigenGWAS analyses were performed using the R-package GWASpoly (43)
344 implemented in R version 3.5.2 (<http://www.R-project.org/>). To determine the population structure
345 of the panels, principal component analysis (PCA) was performed using the R-package SNPRelate
346 (44). The first ten principal components associated with the largest eigenvalues were used for
347 EigenGWAS. Population structure and kinship effects were controlled by inclusion of a mixed
348 model of a canonical relationship (kinship) matrix (45), generated from a subset of SNPs (741 for
349 wheat and 2500 for barley) pruned based on genetic positions. For ease of comparison across
350 GWAS scans, the threshold for significance was set to $-\log_{10}(p\text{-value}) = 4.0$ which in several GWAS
351 scans was above the threshold obtained using false discovery rate
352 (<http://www.strimmerlab.org/software/fdrtool/index.html>). Manhattan plots and circular plots were
353 generated using R-packages qqman (46) and CMplot (47), respectively.

354 **MAGIC wheat analysis**

355 The significant SNPs from the wheat EnvGWAS were used for validation against agronomic traits
356 identified in the 'NIAB Diverse MAGIC' population (18). Analysis was performed in R using
357 adjustments for the funnel structure of the cross (48). All data used were obtained from the
358 following websites that hosts the genotyping and phenotyping data of the 550 MAGIC-diverse RILs
359 <http://mtweb.cs.ucl.ac.uk/mus/www/MAGICdiverse/index.html>.

360 Data availability

361 Genotypic data sets of the study are available from NIAB, UCL and JHI via following websites:

362 <https://www.niab.com/research/agricultural-crop-research/resources>

363 <http://mtweb.cs.ucl.ac.uk/mus/www/MAGICdiverse/index.html>

364 <http://www.barleyhub.org/projects/impromalt/>.

365 Acknowledgments

366 This project was part funded by projects IMPROMALT BB/K/0070251/1 and WAGTAIL
367 BB/J002542/1, jointly funded by the BBSRC and UK wheat and barley breeders; and by direct
368 funding of Rajiv Sharma from SRUC. We thank Mark Looseley and Hazel Bull (JHI), for sharing the
369 genotypic data set of the barley varieties analysed here, as well as Chin Jian Yang, Ian Dawson
370 and David Marshall (SRUC) for helpful discussion throughout the work.

371 Figure Legends

372 **Fig. 1.** Wheat EnvGWAS for variety age. Manhattan plots of the four panels are shown. On the x-
373 axis genetic positions are based on the consensus map (41) are displayed for (A) UK winter wheat
374 and (B) Brazilian spring wheat panels; a pseudo-genetic map positions that relates to the physical
375 positions (40) of the UK winter (C) and spring (D) barley panels are shown. On the y-axis
376 $-\log_{10}(p)$ -values are displayed. The red line indicates the threshold value of the significance
377 corresponding to $-\log_{10}(p) = 4$.

378 **Fig. 2.** Wheat EigenGWAS for the first ten principal components (PCs). Circular plots of the two
379 wheat panels investigated are shown. Highly significant PCs are in the inner circle and the least
380 significant outer circle are displayed. Genetic positions based on a consensus map (41) are
381 displayed for (A) UK winter and (B) Brazilian spring wheat panels. Chromosomal introgressions
382 significant across multiple PCs are highlighted (See **SI Appendix, Table S9**).

383 **Fig. 3.** Barley EigenGWAS for the first ten principal components (PCs). Circular plots of the four
384 panels are shown. Highly significant PCs are in the inner circle and the least significant outer circle
385 are displayed. Pseudo-genetic map positions that relate to the physical positions (40) are displayed
386 for (A) UK winter and (B) UK spring barley panels. Chromosomal introgressions significant across
387 multiple PCs are highlighted (See **SI Appendix, Table S11**).

388 References

- 389 1. R. A. Fischer, G. O. Edmeades, Breeding and cereal yield progress. *Crop Science* **50**, S-85-
390 S-98 (2010).
- 391 2. R. K. Varshney, V. K. Singh, A. Kumar, W. Powell, M. E. Sorrells, Can genomics deliver
392 climate-change ready crops? *Current Opinion in Plant Biology* **45**, 205–211 (2018).
- 393 3. T. Chiurugwi, S. Kemp, W. Powell, L. T. Hickey, Speed breeding orphan crops. *Theoretical*
394 *and Applied Genetics* **132**, 607–616 (2019).
- 395 4. P. K. Ingvarsson, N. R. Street, Association genetics of complex traits in plants. *New*
396 *Phytologist* **189**, 909–922 (2011).
- 397 5. D. Macarthur, Methods: Face up to false positives. *Nature* **487**, 427–428 (2012)
- 398 6. I. Mackay, H.P. Piepho, A. A. F. Garcia, “Statistical methods for plant breeding” in *Handbook*
399 *of Statistical Genomics*, (John Wiley & Sons, Ltd, 2019), pp. 501–520.
- 400 7. X. Huang, B. Han, Natural variations and genome-wide association studies in crop plants.
401 *Annual Review of Plant Biology* **65**, 531–551 (2014).
- 402 8. J. Cockram, *et al.*, Genome-wide association mapping to candidate polymorphism resolution
403 in the unsequenced barley genome. *Proceedings of the National Academy of Sciences of*
404 *the United States of America* **107**, 21611–21616 (2010).
- 405 9. L. Ramsay, *et al.*, *INTERMEDIUM-C*, a modifier of lateral spikelet fertility in barley, is an
406 ortholog of the maize domestication gene *TEOSINTE BRANCHED 1*. *Nature Genetics* **43**,
407 169–172 (2011).
- 408 10. J. Comadran, *et al.*, Natural variation in a homolog of *Antirrhinum CENTRORADIALIS*
409 contributed to spring growth habit and environmental adaptation in cultivated barley. *Nature*
410 *Genetics* **44**, 1388–1392 (2012).
- 411 11. M. T. Hamblin, E. S. Buckler, J. L. Jannink, Population genetics of genomics-based crop
412 improvement methods. *Trends Genet* **27**, 98–106 (2011).
- 413 12. J. Li, *et al.*, Identifying loci with breeding potential across temperate and tropical adaptation
414 via EigenGWAS and EnvGWAS. *Molecular Ecology* **28**, 3544–3560 (2019).
- 415 13. I. Mackay, *et al.*, Reanalyses of the historical series of UK variety trials to quantify the
416 contributions of genetic and environmental factors to trends and variability in yield over time.
417 *Theoretical and Applied Genetics* **122**, 225–238 (2011).
- 418 14. T. N. Rowan, *et al.*, Powerful detection of polygenic selection and environmental adaptation
419 in US beef cattle populations. Detecting ongoing selection with Generation Proxy Selection
420 Mapping (GPSM). *BioRxiv* (2020) <https://doi.org/10.1101/2020.03.11.988121>.
- 421 15. G. B. Chen, S. H. Lee, Z. X. Zhu, B. Benyamin, M. R. Robinson, EigenGWAS: Finding loci
422 under selection through genome-wide association studies of eigenvectors in structured
423 populations. *Heredity* **117**, 51–61 (2016).
- 424 16. J. Cockram, R. Horsnell, E. hee Soh, C. Norris, D. M. O’Sullivan, Molecular and phenotypic
425 characterization of the alternative seasonal growth habit and flowering time in barley
426 (*Hordeum vulgare* ssp. *vulgare* L.). *Molecular Breeding* **35**, 165 (2015).
- 427 17. J. Cockram, *et al.*, Control of flowering time in temperate cereals: genes, domestication, and
428 sustainable productivity. *Journal of Experimental Botany* **58**, 1231–1244 (2007).
- 429 18. M. F. Scott, *et al.*, Limited haplotype diversity underlies polygenic trait architecture across 70
430 years of wheat breeding. *bioRxiv* (2020). <https://doi.org/10.1101/2020.09.15.296533>
- 431 19. J. S. Heslop-Harrison, A. R. Leitch, T. Schwarzacher, K. Anamthawat-Jónsson, Detection
432 and characterization of 1B/1R translocations in hexaploid wheat. *Heredity* **65**, 385–392

- 433 (1990).
- 434 20. S. Rajaram, C. Mann, G. Qrtiz-Ferrara, A. Mujeeb-kazi, Adaptation, stability and high yield
435 potential of certain 1B/1R CIMMYT wheats. in *In: Sakamoto S. (Ed); "Proc. 6th Int. Wheat.
436 Genet. Symp."*, Kyoto University, Japan., (1983), pp. 613–621.
- 437 21. H. Cheng, *et al.*, Frequent intra- and inter-species introgression shapes the landscape of
438 genetic variation in bread wheat. *Genome Biology* **20**, 1–16 (2019).
- 439 22. O. Rodrigues, J. C. B. Lhamby, A. D. Didonet, J. A. Marchese, Fifty years of wheat breeding
440 in Southern Brazil: Yield improvement and associated changes. *Pesquisa Agropecuaria
441 Brasileira* **42**, 817–825 (2007).
- 442 23. H. Chen, *et al.*, Combinatorial evolution of a terpene synthase gene cluster explains terpene
443 variations in *oryza*. *Plant Physiology* **182**, 480–492 (2020).
- 444 24. I. H. Jørgensen, Discovery, characterization and exploitation of *Mlo* powdery mildew
445 resistance in barley. *Euphytica* **63**, 141–152 (1992).
- 446 25. L. Yang, *et al.*, QTL mapping for grain yield-related traits in bread wheat via SNP-based
447 selective genotyping. *Theoretical and Applied Genetics* **133**, 857–872 (2020).
- 448 26. H. Lehnert, A. Serfling, W. Friedt, F. Ordon, Genome-wide association studies reveal
449 genomic regions associated with the response of wheat (*Triticum aestivum* L.) to
450 mycorrhizae under drought stress conditions. *Frontiers in Plant Science* **871** (2018).
- 451 27. C. D. Zanke, *et al.*, Analysis of main effect QTL for thousand grain weight in European winter
452 wheat (*Triticum aestivum* L.) by genome-wide association mapping. *Frontiers in Plant
453 Science* **6**, 1–14 (2015).
- 454 28. N. Fradgley, *et al.*, A large-scale pedigree resource of wheat reveals evidence for adaptation
455 and selection by breeders. *PLoS Biology* **17**, 1–20 (2019).
- 456 29. U. Beukert, *et al.*, The potential of hybrid breeding to enhance leaf rust and stripe rust
457 resistance in wheat. *Theoretical and Applied Genetics* **133**, 2171–2181 (2020).
- 458 30. M. Helguera, *et al.*, PCR assays for the Lr37-Yr17-Sr38 cluster of rust resistance genes and
459 their use to develop isogenic hard red spring wheat lines *Crop Science* **43**, 1839–1847
460 (2003).
- 461 31. M. E. Looseley, *et al.*, Association mapping of malting quality traits in UK spring and winter
462 barley cultivar collections. *Theoretical and Applied Genetics* **133**, 2567–2582 (2020).
- 463 32. X. Xu, *et al.*, Genome-Wide Association Analysis of Grain Yield-Associated Traits in a Pan-
464 European Barley Cultivar Collection. *The Plant Genome* **11**, 11 (2018).
- 465 33. R. Sharma, *et al.*, Genome-wide association of yield traits in a nested association mapping
466 population of barley reveals new gene diversity for future breeding. *Journal of Experimental
467 Botany* **69**, 3811–3822 (2018).
- 468 34. B. S. Gill, B. R. Friebe, F. F. White, Alien introgressions represent a rich source of genes for
469 crop improvement. *Proceedings of the National Academy of Sciences of the United States of
470 America* **108**, 7657–7658 (2011).
- 471 35. V. Haahr, D. von Wettstein, Studies of an induced high-yielding dwarf-mutant of spring
472 barley . *3rd International Barley Genetics Symposium* , 215–218 Barley Genetics III. (1976).
- 473 36. Y. Zhang, *et al.*, Analysis of the functions of *TaGW2* homoeologs in wheat grain weight and
474 protein content traits. *Plant Journal* **94**, 857–866 (2018).
- 475 37. A. R. Bentley, *et al.*, Applying association mapping and genomic selection to the dissection
476 of key traits in elite European wheat. *Theoretical and Applied Genetics* **127**, 2619–2633
477 (2014).

- 478 38. O. Ladejobi, *et al.*, Reference Genome Anchoring of High-Density Markers for Association
479 Mapping and Genomic Prediction in European Winter Wheat. *Frontiers in Plant Science*
480 **10**, 1278 (2019).
- 481 39. G. Mellers, *et al.*, Genetic Characterization of a Wheat Association Mapping Panel Relevant
482 to Brazilian Breeding Using a High-Density Single Nucleotide Polymorphism Array. *G3*
483 (*Bethesda, Md.*) **10**, 2229–2239 (2020).
- 484 40. M. M. Bayer, *et al.*, Development and Evaluation of a Barley 50k iSelect SNP Array.
485 *Frontiers in Plant Science* **8**, 1–10 (2017).
- 486 41. S. Wang, *et al.*, Characterization of polyploid wheat genomic diversity using a high-density
487 90,000 single nucleotide polymorphism array. *Plant Biotechnology Journal* **12**, 787–796
488 (2014).
- 489 42. C. Sun, *et al.*, The Wheat 660K SNP array demonstrates great potential for marker-assisted
490 selection in polyploid wheat. *Plant Biotechnology Journal* **18**, 1354–1360 (2020).
- 491 43. U. R. Rosyara, W. S. De Jong, D. S. Douches, J. B. Endelman, Software for Genome-Wide
492 Association Studies in Autopolyploids and Its Application to Potato. *The Plant Genome* **9**, 2
493 (2016).
- 494 44. X. Zheng, *et al.*, A high-performance computing toolset for relatedness and principal
495 component analysis of SNP data. *Bioinformatics* **28**, 3326–3328 (2012).
- 496 45. P. M. VanRaden, Efficient Methods to Compute Genomic Predictions. *Journal of Dairy*
497 *Science* **91**, 4414–4423 (2008).
- 498 46. S. D. Turner, qqman: an R package for visualizing GWAS results using Q-Q and manhattan
499 plots. *Journal of Open Source Software* (2018) <https://doi.org/10.21105/joss.00731>.
- 500 47. L. Yin, *et al.*, rMVP: A Memory-efficient, Visualization-enhanced, and Parallel-1 accelerated
501 tool for Genome-Wide Association Study. *bioRxiv* (2020)
502 <https://doi.org/10.1101/2020.08.20.258491>.
- 503 48. I. J. Mackay, *et al.*, An eight-parent multiparent advanced generation inter-cross population
504 for winter-sown wheat: Creation, properties, and validation. *G3: Genes, Genomes, Genetics*
505 (2014) <https://doi.org/10.1534/g3.114.012963>.

506 **Supplementary Figures**

507 **Fig S1.** Linear trend in variety year of release and yield (t/ha). (A) UK wheat; (B) UK barley.

508 **Fig S2.** UK barley Manhattan plots for EnvGWAS from combined winter and spring varieties

509 ($n=704$). (A) Manhattan for the age from the analysed barley varieties; (B) analysis using seasonal
510 growth-habit (winter and spring-type) as a covariate; (C); GWAS analysis of seasonal growth-habit;
511 (D) GWAS analysis of seasonal growth habit with the four peaks identified in C as covariates.

512 **Fig S3.** UK winter wheat Manhattan plots for EnvGWAS validation using a subset of 192 varieties.

513 Manhattan plots are shown for: (A) the first year of variety in National trial; (B) the last year each
514 variety is enlisted on the national list; (C) variety life-span, i.e. how long a variety is on the national
515 list; (D) GWAS on yield.

516 **Fig S4.** UK barley Manhattan plots for EnvGWAS validation using a subset of 197 varieties.
 517 Manhattan plots are shown for: (A) the first year of variety in National trial; (B) the last year each
 518 variety is enlisted in the national list; (C) variety life-span, i.e. how long a variety is on the national
 519 list; (D) GWAS on yield.

520 **Fig S5.** Rocket-plots displaying temporal changes in allele frequencies in wheat (UK-wheat: A-D &
 521 Brazilian-wheat: E-H) and barley panels (winter: I-J & spring: K-N). Also displayed wheat
 522 categories in parenthesis within the panel.

523 **Supplementary Tables**





524

Table 1. Summary of the significant hits detected by EnvGWAS on variety age.
Details in SI Appendix, Table S1 & Table S2.

Pop-name	SNP-name	Chrom	Position (cM)	Ref-allele	Ref-Allele-F	-log(p)	Effects
Winter-Wheat	wsnp_Ex_c572_1A	1A	221.0	A	0.50	6.01	-7.61
	Kukri_c18109_1B	1B	350.0	A	0.92	4.62	11.64
	Excalibur_c153_2A	2A	20.0	A	0.66	6.31	9.50
	RFL_Contig403_2A	2A	62.0	A	0.65	5.24	8.32
	BS00071630_5_2A	2A	87.0	A	0.66	6.18	9.26
	IACX6178_2A	2A	158.0	A	0.66	6.18	9.26
	BS00022799_5_2D	2D	33.0	A	0.66	6.31	9.50
	BobWhite_rep_5B	5B	381.0	A	0.13	4.31	6.94
	BS00021901_5_5D	5D	180.0	T	0.85	5.04	9.58
	BS00022120_5_6A	6A	190.0	T	0.83	8.11	12.87
	Kukri_c16404_6B	6B	322.0	A	0.06	4.06	10.33
	Kukri_c67076_7A	7A	383.0	A	0.14	4.29	8.48
	BobWhite_c429_7B	7B	236.0	A	0.94	4.88	-12.92
Spring-Wheat	Ku_c5725_892_2B	2B	251.0	A	0.49	4.44	-7.35
	RFL_Contig484_2B	2B	318.0	T	0.76	4.20	-9.34
	RAC875_c8642_5A	5A	710.0	A	0.08	4.51	-13.21
Winter-Barley	JHI-Hv50k-201_3H	3H	68.7	A	0.29	4.74	-1.95
	JHI-Hv50k-201_3H	3H	124.5	C	0.64	4.22	1.71
	JHI-Hv50k-201_5H	5H	19.2	A	0.73	5.92	-1.87
Spring-Barley	JHI-Hv50k-201_1H	1H	51.0	A	0.41	4.08	-3.07
	SCRI_RS_148_2H	2H	0.0	A	0.42	5.17	-2.59
	JHI-Hv50k-201_3H	3H	1.7	C	0.22	4.40	3.69
	JHI-Hv50k-201_3H	3H	77.7	C	0.95	4.52	-4.42
	JHI-Hv50k-201_5H	5H	20.5	C	0.12	4.90	3.38
	12_30230_6H	6H	53.1	A	0.88	5.22	4.45
	JHI-Hv50k-201_7H	7H	7.8	A	0.93	5.40	5.37
Spring&Winter-	JHI-Hv50k-201_2H	2H	0.0	C	0.74	4.17	-2.15
	JHI-Hv50k-201_2H	2H	20.3	C	0.92	5.74	-2.86
	JHI-Hv50k-201_3H	3H	45.2	C	0.92	4.15	3.07
	JHI-Hv50k-201_3H	3H	68.7	C	0.14	7.13	-4.41
	JHI-Hv50k-201_3H	3H	126.6	C	0.80	4.29	3.64
	JHI-Hv50k-201_5H	5H	19.2	C	0.82	7.67	-3.43
	JHI-Hv50k-201_5H	5H	105.0	A	0.09	4.51	3.53
	11_20546_5H	5H	160.7	A	0.89	4.70	-2.94
	JHI-Hv50k-201_7H	7H	3.8	C	0.05	5.56	-4.65

Table 2: Collocation of significant loci ($-\log(p) > 3$) in MAGIC with the three major winter wheat GWAS peaks. Collocation used 35K and 90K physical maps.

Trait (MAGIC coding)	Trait decription	Region
GLA_on_31_03_17_Year1	Grain Length and Area	1A
waxiness_leaf_score_Year1	Waxiness	1A
FL_length_x_width_Year1	Flag-leaf length	1A
Pigmentation_score_Year1	Pigmentation-score	1A
Height_FL_to_ear_base_Year1	Height	1A
Spikelets_paired_frequency_in_20_Year1	Spikelets-pair-freq	1A
Spring_type_Year1	Spring-type	1A
Yellow_rust_on_11_05_17_Year1	Yellow-rust	2A
GLA_on_20_04_17_Year1	Grain Length and Area	2A
Chlorosis_score_Year1	Chlorosis-score	2A
GS65_DAS_Year1	Heading	2A
FL_width_Year1	Flag-leaf width	2A
Yield_Year1	Yield	2A
GLA_on_07_03_17_Year1	Grain Length and Area	6A
waxiness_leaf_score_Year1	Waxiness	6A
FL_angle_Year1	Flag-leaf angle	6A
FL_length_Year1	Flag-leaf length	6A
FL_length_x_width_Year1	Flag leaf area	6A
FL_length_width_ratio_Year1	Flag leaf length/width ratio	6A
Height_to_FL_Year1	Height	6A
Height_to_ear_base_Year1	Height	6A
Height_to_ear_tip_Year1	Height	6A
Ear_length_Year1	Ear-length	6A
Height_FL_to_ear_base_Year1	Height	6A

A**C****B****D**

Chromosome

Chromosome

