

LncRNA GAS5 attenuates fibroblast activation through inhibiting Smad3 signaling

Rui Tang^{2,##}, Yung-Chun Wang¹⁺, Xiaohan Mei^{1,2+}, Ning Shi¹, Chengming Sun², Ran Ran², Gui Zhang², Wenjing Li², and Kevin F. Staveley-O'Carroll^{1,4}, Guangfu Li¹, Shi-You Chen^{1,2,3*}

¹Department of Surgery, University of Missouri School of Medicine, Columbia, MO 65212;

²Department of Physiology & Pharmacology, University of Georgia, Athens, GA, 30602;

³Department of Medical Pharmacology & Physiology, University of Missouri School of Medicine, Columbia, MO 65212

⁴ The Research Service, Harry S. Truman Memorial Veterans Hospital, Columbia, MO 65212

Running title: GAS5 in fibroblast activation.

⁺These authors contribute equally to this paper.

[#]Current address: Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305

^{*}To whom correspondence should be addressed: Department of Sugery, University of Missouri School of Medicine, 1 Hospital Drive, Columbia, MO 65212, Tel: (573) 884-0371; Fax: (573) 884-4585; Email: scqvd@missouri.edu

Keywords: LncRNA, GAS5, TGF- β , fibroblast activation, skin fibrosis

Abbreviations:

TGF- β	Transforming Growth Factor β
lncRNA	Long noncoding RNA
GAS5	Growth arrest-specific transcript 5
ECM	Extracellular matrix
Col1A	Collagen 1A
T β R	TGF- β receptor
JNK	c-Jun N-terminal kinase
FISH	Fluorescence in situ hybridization
RIP	RNA immunoprecipitation
Co-IP	Co-immunoprecipitation
Smad3	Mothers against decapentaplegic homolog 3
PPM1A	Protein phosphatase 1A
α -SMA	Smooth muscle α -actin
IHC	Immunohistochemistry
RNP	RNA-Protein complex

Abstract:

Transforming Growth Factor β (TGF- β)-induced fibroblast activation is a key pathological event during tissue fibrosis. Long noncoding RNA (lncRNA) is a class of versatile gene regulators participating in various cellular and molecular processes. However, the function of lncRNA in fibroblast activation is still poorly understood. In this study, we identified growth arrest-specific transcript 5 (GAS5) as a novel regulator for TGF- β -induced fibroblast activation. GAS5 expression was downregulated in cultured fibroblasts by TGF- β and in resident fibroblasts from bleomycin-treated skin tissues. Overexpression of GAS5 suppressed TGF- β -induced fibroblast to myofibroblast differentiation. Mechanistically, GAS5 directly bound Smad3 and promoted Smad3 binding to PPM1A, a Smad3 dephosphatase, and thus accelerated Smad3 dephosphorylation in TGF- β -treated fibroblasts. In addition, GAS5 inhibited fibroblast proliferation. Importantly, local delivery of GAS5 via adenoviral vector suppressed bleomycin-induced skin fibrosis in mice. Collectively, our data revealed that GAS5 suppresses fibroblast activation and fibrogenesis through inhibiting TGF- β /Smad3 signaling, which provides a rationale for an lncRNA-based therapy to treat fibrotic diseases.

69 Introduction

70

71 Tissue fibrosis is a wound healing progress following injury, during which normal
72 parenchymal tissue is gradually replaced by connective tissue, accompanied by deposition of
73 extracellular matrix (ECM) components (7, 51, 53). The pathogenesis of fibrosis is initiated in
74 the residential area where local injury triggers innate immune response and causes recruitment
75 of immune cells (26, 32, 40). Pro-fibrotic cytokines or growth factors such as transforming
76 growth factor β (TGF- β) secreted from circulating monocytes and residential macrophages
77 induce fibroblast activation through fibroblast-myofibroblast transition (6, 22, 37, 38).
78 Activated fibroblasts proliferate and thus increase tissue mass. Meanwhile, myofibroblasts
79 express extracellular matrix proteins such as collagen 1A (CollA), the main component of
80 excessive ECM deposition in human fibrotic diseases such as skin, cardiovascular, pulmonary,
81 and liver fibrosis. (57, 58).

82

83 TGF- β signaling regulates a wide spectrum of cellular processes, including cell differentiation,
84 proliferation, migration, and apoptosis (33, 35). TGF- β induces fibrosis through inducing
85 fibroblast trans-differentiation into collagen-producing myofibroblasts and increasing
86 fibroblast proliferation (5, 29). TGF- β transduces its signal mainly through Smad-dependent
87 pathways. TGF- β binds and activates TGF- β receptors (T β Rs), leading to Smad
88 phosphorylation and translocation into nuclei where it activates the transcription of target genes
89 (14). However, how Smad signaling is precisely modulated during TGF- β -induced fibroblast
90 activation and how TGF- β coordinates fibroblast transdifferentiation and proliferation during
91 fibrogenesis are not completely understood.

92

93 Long non-coding RNAs (lncRNAs) are a class of versatile regulators which regulate gene
94 expression at various levels ranging from chromatin modulation to protein degradation (25).
95 Growth Arrest Specific 5 (GAS5) is a well-known lncRNA that suppresses cell proliferation
96 and induces apoptosis (27, 36). Our previous studies have identified GAS5 as a regulator in
97 both Smad3-dependent cell differentiation and p53-dependent cell survival (47, 49). Since
98 TGF- β signaling regulates both Smad-dependent fibroblast-myofibroblast transition and
99 fibroblast proliferation, we hypothesized that GAS5 may be involved in TGF- β -induced
100 fibroblast activation and fibrogenesis.

101

102 In the current study, we firstly observed that GAS5 was downregulated in TGF- β -treated
103 fibroblast cells and bleomycin-injected skin tissues. By both gain-of- and loss-of-function
104 studies, we found that GAS5 suppressed TGF- β -induced fibroblast-myofibroblast transition
105 through modulating Smad3 signaling. Mechanistically, GAS5 directly bound to Smad3, which
106 enhanced phosphatase PPM1A binding to Smad3, and thus accelerated Smad3
107 dephosphorylation. GAS5 also inhibited fibroblast proliferation by blocking c-Jun N-terminal
108 kinase (JNK) signaling pathway. In vivo, local delivery of adenoviral vector expressing GAS5
109 inhibited bleomycin-induced skin fibrosis in mice.

110

111 Material and Methods:

112

113 **Cells and Reagents:** NIH-3T3 cells were purchased from American Type Culture Collection
114 (ATCC). Cells were maintained at 37 °C in a humidified 5% CO₂ incubator in Dulbecco's
115 Modified Eagle's Medium (GIBCO, CA, USA) containing 10% fetal bovine serum (GIBCO,
116 CA, USA), 100 units/ml penicillin and 100 μ g/ml streptomycin. TGF- β 1 was obtained from
117 R&D Systems (Minneapolis, MN). mGAS5 siRNA (n251731) was purchased from Life
118 Technologies (Gaithersburg, MD). Smad3 inhibitor SIS3 was purchased from Sigma-Aldrich

(St. Louis, MO, USA). PPM1A (PA5–29275) antibody was purchased from ThermoFisher Scientific (Pittsburgh, PA)(15). Akt (4691S), phospho-Akt (9271S), JNK (9252S), phospho-JNK (9251S), p38 (9212S), phospho-p38 (4511S), Smad3 (9523S), phospho-Smad3 (9520S), and Smad4 (38454) antibodies were purchased from Cell Signaling (Danvers, MA, USA)(48, 49). GAPDH (G8795) and α -SMA (A2547) antibodies were purchased from Sigma-Aldrich (St. Louis, MO, USA)(43). PCNA (sc-56) and Type I Collagen (Col1A) (sc-25974) antibodies were purchased from Santa Cruz Biotechnology (Dallas, TX, USA) (43, 47). Smad3 overexpression plasmid was constructed by subcloning human Smad3 cDNA into pcDNA3.0 backbone plasmid (Addgene). GAS5 adenoviral vector was constructed by inserting mouse GAS5 cDNA into pShuttle-IRES-hrGFP-1 vector (Agilent), and adenovirus was packaged as described previously (49). All constructs were confirmed by Sanger sequencing.

Primary mouse skin fibroblast preparation: Skin tissues of C57BL/6J mice (The Jackson Laboratory) were dissected, cut into small pieces, and digested in 5 ml tissue digest media (3.5 ml HBSS-Ca²⁺ free, 0.5 mL Trypsin-EDTA (0.25%), 5 mg Collagenase IV (Worthington), 25 U Dispase (Corning)) in a hybridization chamber with rotation at 37°C for 30 minutes. Digestion was then neutralized by adding 5 ml ice-cold Quench Solution (4.5 ml L15 media, 0.5 mL FBS, 94 μ g DNase). Single cell suspensions were generated by filtering through a 40 μ m cell strainer, spinning down at 500 rcf for 5 minutes followed by washing with PBS twice. Skin fibroblasts were resuspended and cultured in Dulbecco's Modified Eagle's Medium (GIBCO, CA, USA) containing 10% fetal bovine serum (GIBCO, CA, USA), 100 units/ml penicillin and 100 μ g/ml streptomycin.

Animals and skin fibrosis model: All animals were housed under conventional conditions in the animal care facility and received humane care in compliance with the Principles of Laboratory Animal Care formulated by the National Society for Medical Research and the Guide for the Care and Use of Laboratory Animals. Bleomycin-induced dermal sclerosis/fibrosis was generated following previously published protocol (43). 8-10 week old male C57BL/6 mice were injected subcutaneously with bleomycin (0.02U) in PBS every other day for 14 or 28 days. PBS was injected as control. Adenovirus (1×10^8 pfu per mouse) expressing GFP or mGAS5 was injected twice on the first day and 14 days following the first bleomycin injection. After 2 or 4 weeks, the animals were euthanized by CO₂ asphyxiation and cervix dislocation. The skin areas with bleomycin injection were removed and processed for biochemical or histological analysis. All animal surgical procedures were approved by the Institutional Animal Care and Use Committee of the University of Georgia.

Quantitative RT-qPCR (qPCR): Total RNA was extracted from cells or tissues using Trizol reagent (Life Technologies, Gaithersburg, MD) and reverse-transcribed to cDNA using iScript™ cDNA Synthesis Kit (Bio-Rad, Hercules, CA). qPCR was performed using a Stratagene Mx3005 qPCR thermocycler (Agilent Technologies, La Jolla, CA). All reactions including no template controls were run in triplicates. After the reaction, the CT values were determined using fixed threshold settings. LncRNA expression was normalized to Cyclophilin (CYP). Primers used in this study were listed in Supplementary Table 1.

RNA immunoprecipitation (RIP) assay: RIP assay was performed as described (49). Cells at 80-90% confluence in 15 cm² culture dishes were fixed with 1% Paraformaldehyde (PFA) and lysed in FA lysis buffer (50 mM HEPES, 140 mM NaCl, 1 mM EDTA, 1% (v/v) Triton X-100, 0.1% (w/v) sodium deoxycholate, pH7.5) containing 40U/ml RNase inhibitor (Sigma-Aldrich, St. Louis, MO) and 1X Halt™ Protease Inhibitor Cocktail (ThermoFisher Scientific, Grand Island, NY). After 4-6 rounds of 50% power output sonication, 300 μ l of whole cell

extracts (around 500 µg total proteins) were incubated with normal rabbit IgG, Smad3, Smad4, or PPM1A antibodies (1 µg) at 4°C overnight. Next day, the immune complexes were captured with 50 µl protein A/G agarose beads (Santa Cruz Biotechnology, Dallas, TX, USA). After washing with FA lysis buffer, samples were incubated with Proteinase K at 42°C for 1 hour to digest the protein and then immunoprecipitated RNA was isolated. Purified RNA was subjected to qRT-PCR analysis for detecting the presence of GAS5.

Western blot: Cultured cells or tissue samples were lysed in RIPA buffer (50 mM Tris-HCl, pH 7.4; 150 mM NaCl; 1% NP-40; and 0.1% SDS), and incubated with continuous rotation for 10 min at 4 °C, and then centrifuged at 12 000 ×g. The supernatant was collected, and the protein concentration was determined by a BCA assay (Pierce, Rockford, USA). Protein extracts (60-100 µg) were dissolved on 10% sodium dodecyl sulfate-polyacrylamide gels (SDS-PAGE) and transferred to polyvinylidene difluoride (PVDF) membranes. The membranes were blocked with 5% non-fat milk in Tris-buffered saline (TBS) plus Tween-20 (TBST) at room temperature for 1 h followed by incubation with primary antibodies diluted in TBST at 4 °C overnight. After three 10-min washing with TBST, blots were incubated with the appropriate secondary antibody conjugated to HRP at room temperature for 1 h. The protein expression was detected with enhanced chemiluminescent reagent.

Co-immunoprecipitation assay (Co-IP): Cells were transduced with Ad-GFP or Ad-GAS5 for 24 hours, and lysed with ice-cold lysis buffer containing protease inhibitor mix (Sigma). The lysates were incubated with IgG or anti-PPM1A antibody for 1 h followed by incubation with protein A/G-beads at 4 °C for 12 h. The immunoprecipitates were pelleted, washed, and subjected to immunoblotting.

Chromatin immunoprecipitation (ChIP): Fresh tissues were minced into small pieces in ice-cold PBS with a clean razor blade. Formaldehyde was then added to a final concentration of 1% and incubated with shaking at room temperature for 10 min. The cross-linked tissues were collected by centrifugation at 4°C and washed with PBS containing protease inhibitors before final collection. The tissues were resuspended by rotating in 1% SDS lysis buffer at 4°C for 20 min followed by sonication on ice to shear DNA into 500-1000 bp fragments. The lysates were immunoprecipitated with 2 µg of IgG (negative control) or Smad3 antibody in coimmunoprecipitation reagents (17-195, Millipore). Semi-quantitative and quantitative PCR were performed to amplify α-SMA or Col1A promoter regions containing SBE.

Luciferase reporter assay: 3T3 cells cultured in 12-well plates were transduced with AdGFP or AdGAS5 and transfected with 250 ng of firefly luciferase reporter plasmid driven by α-SMA promoter using Lipofectamie LTX (Invitrogen, USA). Cells were treated with vehicle or 5 ng/ml of TGF-β1 for 8 hours, and luciferase activities were measured using a luciferase assay kit (Promega) by following the manufacturer's protocol. The experiments were repeated for three times with triplicates.

Immunohistochemistry (IHC) staining: Tissue sections were rehydrated, permeabilized with 0.01% Triton X-100 in PBS, blocked with 10% goat serum, and incubated with primary antibodies at 4°C overnight followed by incubation with horseradish peroxidase (HRP)-conjugated secondary antibody. The sections were counterstained with hematoxylin.

MTT cell proliferation assay: Cell proliferation was evaluated with 3-(4,5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium (MTT) assay using a TACS MTT Cell Proliferation Assay Kit (Trivegen). The optical density at 570 nm was measured.

Fluorescence in situ hybridization (FISH): GAS5 RNA probes (sense/antisense) were synthesized and labeled using the FISH Tag RNA Multicolor kit (Life Technologies, Gaithersburg, MD). Skin tissue cryosections were digested with 20 µg/ml of proteinase K at 37 °C for 1 hour, and washed with 2 × SSC solution and with water for 5 min each at room temperature. The slides were dehydrated, air-dried and incubated with pre-denatured GAS5 probes in a dark and humid environment for hybridization at 55°C for 24 hours. The slides were then washed in 50% formamide in 2x SSC for 4 times before mounting. Nuclear was counterstained with 5,6-diamidino-2-phenylindole (DAPI).

Statistical analysis: Sample or experiment sizes were determined empirically to achieve sufficient statistical power. For animal studies, the sample size was chosen to minimize the number of sacrificed animals while obtaining sufficient statistical power. In all of the experiments reported in this study, no data point was excluded. No randomization was used in this study. There was no blinding method used to assign individuals to experimental groups. The variance is similar between groups that are being statistically compared.

All in vitro experiments is repeated at least three times in triplicates. At least five mice were used for each treatment group in bleomycin-induced skin fibrosis studies. All values are presented as means ± SEM. Comparisons of parameters between two groups were made by two-tailed Student's t-tests. The differences among several groups will be evaluated by one-way ANOVA with Tukey-Kramer post hoc evaluation. p-values <0.05 will be considered statistically significant. P values less than 0.05 and 0.01 were considered significant (*) or very significant (**), respectively.

Results

GAS5 expression was decreased in TGF-β-activated fibroblasts.

TGF-β activated fibroblasts as shown by the induction of myofibroblast marker α-SMA and CollA in both 3T3 fibroblasts and primary cultured mouse skin fibroblasts (Fig. 1A-1D). To test if GAS5 is involved in the fibroblast activation, we first examined GAS5 expression in TGF-β-treated fibroblasts. 3T3 fibroblasts were treated with vehicle, 1, 2, 5, or 10 ng/ml TGF-β for 12 hours in serum-free DMEM. Total RNA was extracted, and GAS5 expression was detected by RT-qPCR. As shown in Fig. 1E, GAS5 was down-regulated in TGF-β-treated 3T3 cells. Interestingly, 10 ng/ml of TGF-β slightly increased the GAS5 expression as compared to the treatment with 5 ng/ml of TGF-β (Fig 1E). This is probably because high concentration of TGF-β (≥10 ng/ml) can inhibit cell proliferation, which could re-activate GAS5 transcription as a feedback response. TGF-β-induced GAS5 reduction was also validated in primary cultured mouse skin fibroblasts (Fig. 1F). Since fibroblast activation is an essential process leading to tissue fibrosis, and bleomycin-induced skin fibrosis involves TGF-β signaling (13, 19, 43), we assessed if GAS5 expression is altered in bleomycin-treated skin tissues. C57BL/6 mice were injected subcutaneously with 0.02 U bleomycin every other day for 14 days, skin tissues were dissected, tissue RNA was extracted, and GAS5 expression was detected by qPCR. As shown in Fig. 1G, GAS5 was indeed downregulated in fibrotic skin tissues. Moreover, we observed GAS5-expressing cells by fluorescence in situ hybridization (FISH) assay. Comparing to the vehicle-treated skins, the numbers of GAS5-positive cells were significantly reduced in bleomycin-treated skin tissues (Fig. 1H-1I). These results suggested that GAS5 may be involved in TGF-β-induced fibroblast activation and skin fibrosis.

GAS5 blocked TGF-β-induced fibroblast-myofibroblast transition.

Since TGF- β induces fibroblast activation through Smad-dependent pathway, and our previous studies have shown that GAS5 blocks TGF- β /Smad3 signaling in SMC differentiation, we sought to determine if GAS5 affects TGF- β -induced fibroblast to myofibroblast transition. Thus, we detected α -SMA and Col1A protein expression in 3T3 cells transduced with AdGFP or AdGAS5 along with transfection with control or GAS5 siRNA. GAS5 overexpression and knockdown efficacies were detected by RT-qPCR (Supplemental Fig. S1). As shown in Fig. 2A-2B, overexpression of GAS5 suppressed Col1A and α -SMA protein expression both at the basal state and under TGF- β treatment. Conversely, knockdown of GAS5 by its siRNA increased TGF- β -induced Col1A and α -SMA expression (Fig. 2C-2D). Consistent with the protein expression, GAS5 negatively regulated Col1A and α -SMA mRNA expression (Fig. 2E-2F). Interestingly, GAS5 caused 5 times more reduction in Col1A and 7 times more reduction in α -SMA expression in TGF- β -treated cells than the vehicle-treated cells (Fig. 2E), suggesting that GAS5 regulates Col1A and α -SMA expression primarily in association with TGF- β signaling. Moreover, overexpression of GAS5 also suppressed TGF- β -induced α -SMA promoter activity (Fig. 2G). These results indicated that GAS5 inhibits TGF- β -induced fibroblast activation by negatively regulating myofibroblast marker gene transcription.

GAS5 promoted Smad3 dephosphorylation.

Previous studies have shown that TGF- β /Smad3 signaling is continuously activated or phosphorylated during tissue fibrogenesis, which causes sustainable Smad nuclear retention (46). We have reported that Smad3 is phosphorylated and translocated into nuclei during the initial stage of TGF- β stimulation while shuttling back to cytoplasm at the later stage of TGF- β -induced smooth muscle differentiation (55). These observations prompted us to hypothesize that GAS5 may alter Smad phosphorylation status/nuclear retention in order to regulate TGF- β -induced fibroblast activation. Since Smad nuclear localization depends on its phosphorylation status, we tested if GAS5 affects Smad2/3 phosphorylation/dephosphorylation turnover in 3T3 cells because Smad2/3 are the two major Smad proteins downstream of TGF- β signaling. As shown in Fig. 3A-3B, TGF- β induced both Smad2 and Smad3 phosphorylation. However, overexpression of GAS5 decreased Smad3, but not Smad2, phosphorylation. These data suggest that GAS5 regulates myofibroblast transition through promoting Smad3 dephosphorylation in 3T3 cells.

GAS5 bound to Smad3 to increase Smad3 binding to PPM1A.

Smad phosphorylation status in the nuclei is controlled by Smad phosphatase (8, 10, 15). To determine the mechanism by which GAS5 promotes Smad dephosphorylation, we first predicted the interactions between GAS5 and various known Smad phosphatases through IncPro (<http://bioinfo.bjmu.edu.cn/Incpro/>) (31). As shown in supplementary Table 2, PPM1A is the top candidate binding GAS5. Since GAS5 only binds Smad3, but not Smad2 (49), and GAS5 overexpression caused Smad3, but not Smad2, dephosphorylation (Fig 3A-3B), we sought to test how GAS5 affects Smad3 phosphorylation and hypothesized that GAS5 regulates Smad3 turnover through binding both Smad3 and PPM1A. Thus, we experimentally determined if GAS5 interacts with PPM1A and Smad3 by performing RNA immunoprecipitation (RIP) assays. RNA-protein complex from 3T3 cells were pulled down using anti-Smad3 or anti-PPM1A antibody with IgG used as a negative control. The presence of GAS5 in the complex was detected by RT-qPCR. As shown in Fig. 3C-3D, GAS5 directly bound to Smad3 (Fig. 3C) as well as PPM1A (Fig. 3D). To test if Smad3 interacts with PPM1A in 3T3 fibroblasts and determine if GAS5 affects Smad3-PPM1A interaction, co-immunoprecipitation (Co-IP) was performed using anti-PPM1A antibody and cell lysates isolated from 3T3 cells transduced with AdGFP/AdGAS5 or transfected with siCtrl/siGAS5. As shown in Fig. 3E-3F, Smad3 indeed interacted with PPM1A. And importantly,

overexpression of GAS5 enhanced Smad3-PPM1A interaction while knockdown of GAS5 suppressed Smad3-PPM1A interaction. These results indicated that GAS5 causes Smad3 dephosphorylation by promoting PPM1A binding to Smad3.

GAS5 inhibited myofibroblast activation through PPM1A-mediated Smad3 dephosphorylation.

To determine if PPM1A is important for GAS5 function in Smad3 signaling during myofibroblast activation, we knocked down PPM1A in 3T3 cells by its siRNA along with GAS5 overexpression and TGF- β treatment. As shown in Fig. 4A-4B, overexpression of GAS5 down-regulated Col1A and α -SMA expression as well as Smad3 phosphorylation. However, knockdown of PPM1A rescued, at least partially, the inhibitory effect of GAS5 on Smad3 phosphorylation and Col1A and α -SMA expression, indicating that PPM1A mediated GAS5 function in Smad3 phosphorylation during the myofibroblast activation.

To determine if GAS5 regulates myofibroblast gene expression through Smad3, we overexpressed GAS5 in 3T3 cells via adenoviral transduction along with transfection of Smad3 expression plasmid followed by TGF- β induction. As shown in Fig. 4C-4D, overexpression of GAS5 suppressed TGF- β -induced Col1A expression in 3T3 cells. However, Smad3 overexpression restored the Col1A expression. On the other hand, knockdown of GAS5 increased Col1A expression, but blockade of Smad3 activity by its inhibitor SIS3 demolished the effect of silencing GAS5 on Col1A expression (Fig. 4E-4F). These results demonstrated that GAS5 negatively regulated the myofibroblast gene expression by suppressing PPM1A-induced Smad3 dephosphorylation.

GAS5 suppressed TGF- β induced fibroblast proliferation.

Fibroblast proliferation is one of the main sources for the tissue mass increase during fibrosis (12, 18). TGF- β signaling is known to regulate cell proliferation (1, 4, 24). To confirm the function of TGF- β in fibroblast proliferation, 3T3 cells were treated with 0, 1, 5 or 10 ng/ml TGF- β for 48 hours in serum-free DMEM, following by MTT assay to assess cell proliferation. As shown in Supplemental Fig. S2A, TGF- β increased 3T3 cell proliferation in a dose-dependent manner. Consistently, PCNA expression was also dose-dependently increased in response to TGF- β (Fig. S2B-2C). To determine the role of GAS5 in TGF- β -induced 3T3 proliferation, we overexpressed GAS5 by transducing 3T3 cells with control or adenoviral vector expressing GAS5 (AdGAS5) followed by treatment with vehicle or 5 ng/ml of TGF- β for 48 hours for MTT assay or 24 hours for Western blot. As shown in Figure S2D-2F, overexpression of GAS5 suppressed TGF- β -induced 3T3 proliferation (Fig. S2D) as well as PCNA protein expression (Fig. S2E-2F). On the other hand, knockdown of GAS5 by its siRNA increased 3T3 cell proliferation (Fig. S2G) and increased PCNA protein expression (Fig. S2H-2I). These results indicated that GAS5 suppressed TGF- β -induced fibroblast proliferation.

GAS5 suppressed fibroblast proliferation through TGF- β /JNK signaling

It is known that TGF- β /Smad signaling cross-talks with multiple non-Smad signaling pathways, most often PI3K/Akt, JNK and p38 pathways, to regulate cell proliferation (2, 3, 17, 23, 41, 59). To determine if GAS5 inhibits fibroblast proliferation by targeting these signaling pathways, we assessed the phosphorylation of key components in PI3K/Akt, JNK and p38 signaling pathways in TGF- β -treated 3T3 cells. As shown in Supplemental Figure S3A-3B, TGF- β increased Akt and JNK phosphorylation, suggesting these signaling pathways were activated during TGF- β -induced fibroblast proliferation. However, comparing to AdGFP-transduced cells, overexpression of GAS5 (AdGAS5) suppressed JNK phosphorylation while had no effect on p38 phosphorylation and increased the Akt phosphorylation (Fig. S3A-3B). These results suggested that GAS5 suppressed TGF- β -induced fibroblast proliferation only

through JNK pathway. To further validate the function of JNK signaling in GAS5-regulated fibroblast proliferation, we knocked down GAS5 in 3T3 cells via siRNA transfection followed by treatment with vehicle or JNK pathway inhibitor, SP600125. As shown in Fig. S3C, knockdown of GAS5 increased 3T3 cell proliferation, but JNK inhibitor reversed the effect of GAS5. These results demonstrated that GAS5 inhibits fibroblast proliferation through interfering with TGF- β -JNK pathways.

In vivo local delivery of GAS5 via adenoviral vector suppressed skin fibrosis in mice.

Since fibroblast activation is an important mechanism underlying pathological tissue/organ fibrosis, we tested the function of GAS5 in skin fibrosis by using a bleomycin-induced TGF- β -dependent skin sclerosis model (43). GAS5 was overexpressed via adenoviral delivery in bleomycin-treated mouse skin. As shown in Fig. 5A-5D, overexpression of GAS5 attenuated bleomycin-induced skin sclerosis as indicated by the reduction of both skin thickness (Fig. 5A-5B) and skin tissue collagen deposition (Fig. 5C-5D). Immunohistochemistry (IHC) staining showed that Col1A- (Fig. 5E-5F) and α SMA-positive cells (Fig. 5G-5H) were significantly reduced in AdGAS5-transduced skin tissues, suggesting that GAS5 inhibited bleomycin-caused skin fibrosis. Moreover, overexpression of GAS5 decreased Col1A and α -SMA protein expression (Fig. 6A-6B) in bleomycin-treated skins. Importantly, GAS5 reduced the Smad3 binding to the Col1A and α -SMA promoters in a chromatin setting in vivo in the bleomycin-treated skin tissues (Fig. 6C-6F), confirming that GAS5 impedes skin fibrogenesis by inhibiting Smad3-mediated transcription of myofibroblast genes.

Discussion

Fibrosis is a chronic wound healing progress characterized by fibroblast activation involving fibroblast proliferation and fibroblast-myofibroblast transition (20, 42). In this study, we identified lncRNA GAS5 as an essential regulator for TGF- β -induced fibroblast activation and skin fibrosis. Specifically, GAS5 directly bound to both Smad3 and PPM1A, thus promoted Smad3 dephosphorylation and suppressed Smad3-induced myofibroblast activation. In addition, GAS5 inhibited 3T3 cell proliferation via blocking JNK signaling (Fig. 7). Importantly, Local adenoviral delivery of GAS5 effectively suppressed bleomycin-induced skin fibrosis in mice, suggesting that GAS5 may be used as a promising RNA-based therapeutic agent for treating fibrotic diseases.

LncRNA normally functions as a suppressive protein sponge for transcription factors (27, 49). LncRNA can also serve as a scaffold RNA which brings spatial proximity of different components thus facilitates protein complex formation and enhances their functions (45, 52). GAS5 has been shown to regulate different signaling cascades through recruiting protein co-factors to form RNA-Protein complex (RNP). Indeed, we have previously identified GAS5 as a Smad3 sponge RNA in the initial stage of TGF- β signaling transduction. In the current study, we identified a new function of GAS5 at a later stage of TGF- β signaling. I.e., GAS5 can quench the persistent TGF- β /Smad signaling observed in fibrosis by promoting PPM1A-mediated Smad3 dephosphorylation. Although transient TGF- β activity is important for tissue repair/regeneration, persistent TGF- β signaling results in fibrosis and scarring in fibrotic disease (11, 30). Our results suggested that GAS5 may act as a brake for the persistent TGF- β signaling in tissue fibrosis. GAS5 affects myofibroblast marker gene Col1A and α -SMA expression at both basal and TGF- β -treated states, suggesting that GAS5 may be required for maintaining fibroblast homeostasis of the quiescent cells. Interestingly, the effect of silencing GAS5 on α -SMA is much less than Col1A in the basal state, which is probably because α -SMA expression is also regulated by many factors other than GAS5. Of significance, GAS5

has more profound impacts on Col1A and α -SMA expression in TGF- β -treated cells, suggesting that GAS5 may be more important and powerful in blocking the fibroblast activation and fibrosis than maintaining quiescent cell activity. Indeed, GAS5 has also been identified as an anti-fibrotic lncRNA in cardiac and hepatic fibrosis although the mechanism is mainly through functioning as a miRNA sponge (16, 50, 56). Whether GAS5 regulates TGF- β /Smad signaling through sponging miRNAs in skin fibrosis requires further studies.

TGF- β not only initiates fibroblast-myofibroblast transition but also stimulates fibroblast proliferation, usually through cross-talking with Smad independent pathways, such as PI3K/Akt, JNK and p38 (1, 4, 24). Given the fact that GAS5 is a potent anti-proliferation lncRNA, it is expected that GAS5 also suppresses TGF- β -induced fibroblast proliferation. Although previous studies have shown that TGF- β signaling interacts with almost all major cellular signaling pathways (54), our results indicate that JNK is the major downstream signaling pathway mediating GAS5 function in TGF- β -induced fibroblast proliferation. There are reports showing that knockdown of GAS5 suppresses JNK phosphorylation in primary retinal ganglion cells and a neuroblastoma cell line (34, 60). These discrepancies in GAS5 function may be due to cell type-specific effects. How GAS5 interacts with JNK signaling, and whether the interaction between GAS5 and JNK signaling is Smad3-dependent are interesting subjects for future studies.

In the progression of fibrogenesis, intensive interplays occur among different cell types. Epithelial cells, fibroblast, smooth muscle cells and immune cells all have shown to play important roles in this chronic disease (21, 28, 39). M2 macrophages, for example, is known to play important roles not only in the initial wound healing response (anti-fibrotic) but also in the later TGF- β secretion and fibroblast activation stages (pro-fibrotic) (9). Interestingly, GAS5 has been reported to inhibit M2 macrophage polarization (44). Thus, GAS5 may have many unidentified functions in fibrosis. For example, it could coordinate or modulate TGF- β signaling among different cell types during fibrogenesis. The investigation into these topics may shed new light on the RNA-based therapy in treating TGF- β -related fibrotic diseases.

Acknowledgments

This work was supported by grants from National Institutes of Health (HL123302, HL119053, HL135854, and HL147313 to S.-Y.C.) and VA merit Award (I01 BX004065-1 to K. F. S). R. T. is a recipient of Stanford University School of Medicine Dean's Postdoctoral Fellowship and a TRDRP Postdoctoral Fellowship (27FT-0044).

Disclosures: None

Endnotes:

The Supplemental Materials are available in the following link:

<https://doi.org/10.6084/m9.figshare.12140880>

Reference:

1. **Abdullah N, Torres B, Basu M, and Johnson H.** Differential effects of epidermal growth factor, transforming growth factor- α , and vaccinia virus growth factor in the positive regulation of IFN- γ production. *The Journal of Immunology* 143: 113-117, 1989.
2. **Bakin AV, Rinehart C, Tomlinson AK, and Arteaga CL.** p38 mitogen-activated protein kinase is required for TGF β -mediated fibroblastic transdifferentiation and cell migration. *Journal of cell science* 115: 3193-3206, 2002.
3. **Battegay EJ, Raines EW, Seifert RA, Bowen-Pope DF, and Ross R.** TGF- β induces bimodal proliferation of connective tissue cells via complex control of an autocrine PDGF loop. *Cell* 63: 515-524, 1990.
4. **Bettinger DA, Yager DR, Diegelmann RF, and Cohen IK.** The effect of TGF- β on keloid fibroblast proliferation and collagen synthesis. *Plastic and reconstructive surgery* 98: 827-833, 1996.
5. **Biernacka A, Dobaczewski M, and Frangogiannis NG.** TGF- β signaling in fibrosis. *Growth factors* 29: 196-202, 2011.
6. **Bonfield TL, Panuska JR, Konstan MW, Hilliard KA, Hilliard JB, Ghnaim H, and Berger M.** Inflammatory cytokines in cystic fibrosis lungs. *American journal of respiratory and critical care medicine* 152: 2111-2118, 1995.
7. **Border WA, and Noble NA.** Transforming growth factor β in tissue fibrosis. *New England Journal of Medicine* 331: 1286-1292, 1994.
8. **Bourgeois B, Gilquin B, Tellier-Lebègue C, Östlund C, Wu W, Pérez J, El Hage P, Lallemand F, Worman HJ, and Zinn-Justin S.** Inhibition of TGF- β signaling at the nuclear envelope: characterization of interactions between MAN1, Smad2 and Smad3, and PPM1A. *Sci Signal* 6: ra49-ra49, 2013.
9. **Braga TT, Agudelo JSH, and Camara NOS.** Macrophages during the fibrotic process: M2 as friend and foe. *Frontiers in immunology* 6: 602, 2015.
10. **Bruce DL, and Sapkota GP.** Phosphatases in SMAD regulation. *FEBS letters* 586: 1897-1905, 2012.
11. **Cutroneo KR.** TGF- β -induced fibrosis and SMAD signaling: oligo decoys as natural therapeutics for inhibition of tissue fibrosis and scarring. *Wound Repair and Regeneration* 15: S54-S60, 2007.
12. **Daly TJ, and Weston WL.** Retinoid effects on fibroblast proliferation and collagen synthesis in vitro and on fibrotic disease in vivo. *Journal of the American Academy of Dermatology* 15: 900-902, 1986.
13. **Daniels CE, Wilkes MC, Edens M, Kottom TJ, Murphy SJ, Limper AH, and Leof EB.** Imatinib mesylate inhibits the profibrogenic activity of TGF- β and prevents bleomycin-mediated lung fibrosis. *The Journal of clinical investigation* 114: 1308-1316, 2004.
14. **Derynck R, and Zhang YE.** Smad-dependent and Smad-independent pathways in TGF- β family signalling. *Nature* 425: 577-584, 2003.
15. **Dong K, Guo X, Chen W, Hsu AC, Shao Q, Chen J-F, and Chen S-Y.** Mesenchyme homeobox 1 mediates transforming growth factor- β (TGF- β)-induced smooth muscle cell differentiation from mouse mesenchymal progenitors. *Journal of Biological Chemistry* 293: 8712-8719, 2018.
16. **Dong Z, Li S, Wang X, Si L, Ma R, Bao L, and Bo A.** lncRNA GAS5 restrains CCl4-induced hepatic fibrosis by targeting miR-23a through the PTEN/PI3K/Akt signaling pathway. *American Journal of Physiology-Gastrointestinal and Liver Physiology* 316: G539-G550, 2019.

17. **Engel ME, McDonnell MA, Law BK, and Moses HL.** Interdependent SMAD and JNK signaling in transforming growth factor- β -mediated transcription. *Journal of Biological Chemistry* 274: 37413-37420, 1999.
18. **Garbuzenko E, Nagler A, Pickholtz D, Gillery P, Reich R, Maquart FX, and Levi-Schaffer F.** Human mast cells stimulate fibroblast proliferation, collagen synthesis and lattice contraction: a direct role for mast cells in skin fibrosis. *Clinical & Experimental Allergy* 32: 237-246, 2002.
19. **Giri SN, Hyde DM, and Hollinger MA.** Effect of antibody to transforming growth factor beta on bleomycin induced accumulation of lung collagen in mice. *Thorax* 48: 959-966, 1993.
20. **Grotendorst GR, Rahmanie H, and Duncan MR.** Combinatorial signaling pathways determine fibroblast proliferation and myofibroblast differentiation. *The FASEB Journal* 18: 469-479, 2004.
21. **Hays S, Ferrando R, Carter R, Wong H, and Woodruff P.** Structural changes to airway smooth muscle in cystic fibrosis. *Thorax* 60: 226-228, 2005.
22. **Heymann F, Trautwein C, and Tacke F.** Monocytes and macrophages as cellular targets in liver fibrosis. *Inflammation & Allergy-Drug Targets (Formerly Current Drug Targets-Inflammation & Allergy)* 8: 307-318, 2009.
23. **Huang SS, and Huang JS.** TGF- β control of cell proliferation. *Journal of cellular biochemistry* 96: 447-462, 2005.
24. **Huang W-Y, Li Z-G, Rus H, Wang X, Jose PA, and Chen S-Y.** RGC-32 mediates transforming growth factor- β -induced epithelial-mesenchymal transition in human renal proximal tubular cells. *Journal of Biological Chemistry* 284: 9426-9432, 2009.
25. **Huarte M.** The emerging role of lncRNAs in cancer. *Nature medicine* 21: 1253-1261, 2015.
26. **Huax F, Liu T, McGarry B, Ullenbruch M, Xing Z, and Phan SH.** Eosinophils and T lymphocytes possess distinct roles in bleomycin-induced lung injury and fibrosis. *The Journal of Immunology* 171: 5470-5481, 2003.
27. **Kino T, Hurt DE, Ichijo T, Nader N, and Chrousos GP.** Noncoding RNA gas5 is a growth arrest- and starvation-associated repressor of the glucocorticoid receptor. *Sci Signal* 3: ra8-ra8, 2010.
28. **Krieg T, Abraham D, and Lafyatis R.** Fibrosis in connective tissue disease: the role of the myofibroblast and fibroblast-epithelial cell interactions. *Arthritis research & therapy* 9: S4, 2007.
29. **Leask A, and Abraham DJ.** TGF- β signaling and the fibrotic response. *The FASEB Journal* 18: 816-827, 2004.
30. **Leask A, Denton CP, and Abraham DJ.** Insights into the molecular mechanism of chronic fibrosis: the role of connective tissue growth factor in scleroderma. *Journal of Investigative Dermatology* 122: 1-6, 2004.
31. **Lu Q, Ren S, Lu M, Zhang Y, Zhu D, Zhang X, and Li T.** Computational prediction of associations between long non-coding RNAs and proteins. *BMC genomics* 14: 651, 2013.
32. **Luzina IG, Todd NW, Iacono AT, and Atamas SP.** Roles of T lymphocytes in pulmonary fibrosis. *Journal of leukocyte biology* 83: 237-244, 2008.
33. **Massagué J, and Chen Y-G.** Controlling TGF- β signaling. *Genes & development* 14: 627-644, 2000.
34. **Miao X, and Liang A.** Knockdown of long noncoding RNA GAS5 attenuates H₂O₂-induced damage in retinal ganglion cells through upregulating miR-124: Potential role in traumatic brain injury. *Journal of cellular biochemistry* 120: 2313-2322, 2019.

35. **Miyazono K.** TGF- β signaling by Smad proteins. *Cytokine & growth factor reviews* 11: 15-22, 2000.
36. **Mourtada-Maarabouni M, Pickard M, Hedge V, Farzaneh F, and Williams G.** GAS5, a non-protein-coding RNA, controls apoptosis and is downregulated in breast cancer. *Oncogene* 28: 195-208, 2009.
37. **Pedroza M, Le TT, Lewis K, Karmouty-Quintana H, To S, George AT, Blackburn MR, Tweardy DJ, and Agarwal SK.** STAT-3 contributes to pulmonary fibrosis through epithelial injury and fibroblast-myofibroblast differentiation. *The FASEB Journal* 30: 129-140, 2015.
38. **Phan SH.** The myofibroblast in pulmonary fibrosis. *Chest* 122: 286S-289S, 2002.
39. **Sakai N, and Tager AM.** Fibrosis of two: Epithelial cell-fibroblast interactions in pulmonary fibrosis. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* 1832: 911-921, 2013.
40. **Simonian PL, Roark CL, Wehrmann F, Lanham AK, del Valle FD, Born WK, O'Brien RL, and Fontenot AP.** Th17-polarized immune response in a murine model of hypersensitivity pneumonitis and lung fibrosis. *The Journal of Immunology* 182: 657-665, 2009.
41. **Song K, Wang H, Krebs TL, and Danielpour D.** Novel roles of Akt and mTOR in suppressing TGF- β /ALK5-mediated Smad3 activation. *The EMBO journal* 25: 58-69, 2006.
42. **Strutz F, and Zeisberg M.** Renal fibroblasts and myofibroblasts in chronic kidney disease. *Journal of the American Society of Nephrology* 17: 2992-2998, 2006.
43. **Sun C, and Chen S-Y.** RGC32 promotes bleomycin-induced systemic sclerosis in a murine disease model by modulating classically activated macrophage function. *The Journal of Immunology* 200: 2777-2785, 2018.
44. **Sun D, Yu Z, Fang X, Liu M, Pu Y, Shao Q, Wang D, Zhao X, Huang A, and Xiang Z.** LncRNA GAS5 inhibits microglial M2 polarization and exacerbates demyelination. *EMBO reports* 18: 1801-1816, 2017.
45. **Sun M, Nie F, Wang Y, Zhang Z, Hou J, He D, Xie M, Xu L, De W, and Wang Z.** LncRNA HOXA11-AS promotes proliferation and invasion of gastric cancer by scaffolding the chromatin modification factors PRC2, LSD1, and DNMT1. *Cancer research* 76: 6299-6310, 2016.
46. **Takagawa S, Lakos G, Mori Y, Varga J, Yamamoto T, and Nishioka K.** Sustained activation of fibroblast transforming growth factor- β /Smad signaling in a murine model of scleroderma. *Journal of investigative dermatology* 121: 41-50, 2003.
47. **Tang R, Mei X, Wang Y-C, Cui X-B, Zhang G, Li W, and Chen S-Y.** LncRNA GAS5 regulates vascular smooth muscle cell cycle arrest and apoptosis via p53 pathway. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* 1865: 2516-2525, 2019.
48. **Tang R, Zhang G, and Chen S-Y.** Smooth muscle cell proangiogenic phenotype induced by Cyclopentenyl Cytosine promotes endothelial cell proliferation and migration. *Journal of Biological Chemistry* 291: 26913-26921, 2016.
49. **Tang R, Zhang G, Wang Y-C, Mei X, and Chen S-Y.** The long non-coding RNA GAS5 regulates transforming growth factor β (TGF- β)-induced smooth muscle cell differentiation via RNA Smad-binding elements. *Journal of Biological Chemistry* 292: 14270-14278, 2017.
50. **Tao H, Zhang J-G, Qin R-H, Dai C, Shi P, Yang J-J, Deng Z-Y, and Shi K-H.** LncRNA GAS5 controls cardiac fibroblast activation and fibrosis by targeting miR-21 via PTEN/MMP-2 signaling pathway. *Toxicology* 386: 11-18, 2017.
51. **Todd NW, Luzina IG, and Atamas SP.** Molecular and cellular mechanisms of pulmonary fibrosis. *Fibrogenesis & tissue repair* 5: 11, 2012.

52. **Wang KC, and Chang HY.** Molecular mechanisms of long noncoding RNAs. *Molecular cell* 43: 904-914, 2011.
53. **Wynn TA.** Cellular and molecular mechanisms of fibrosis. *The Journal of Pathology: A Journal of the Pathological Society of Great Britain and Ireland* 214: 199-210, 2008.
54. **Xiao L, Du Y, Shen Y, He Y, Zhao H, and Li Z.** TGF-beta 1 induced fibroblast proliferation is mediated by the FGF-2/ERK pathway. *Front Biosci (Landmark Ed)* 17: 2667-2674, 2012.
55. **Xie W-B, Li Z, Miano JM, Long X, and Chen S-Y.** Smad3-mediated Myocardin Silencing A NOVEL MECHANISM GOVERNING THE INITIATION OF SMOOTH MUSCLE DIFFERENTIATION. *Journal of Biological Chemistry* 286: 15050-15057, 2011.
56. **Yu F, Zheng J, Mao Y, Dong P, Lu Z, Li G, Guo C, Liu Z, and Fan X.** Long non-coding RNA growth arrest-specific transcript 5 (GAS5) inhibits liver fibrogenesis through a mechanism of competing endogenous RNA. *Journal of Biological Chemistry* 290: 28286-28298, 2015.
57. **Zeisberg EM, Potenta SE, Sugimoto H, Zeisberg M, and Kalluri R.** Fibroblasts in kidney fibrosis emerge via endothelial-to-mesenchymal transition. *Journal of the American Society of Nephrology* 19: 2282-2287, 2008.
58. **Zeisberg M, Bonner G, Maeshima Y, Colorado P, Müller GA, Strutz F, and Kalluri R.** Renal fibrosis: collagen composition and assembly regulates epithelial-mesenchymal transdifferentiation. *The American journal of pathology* 159: 1313-1321, 2001.
59. **Zhang YE.** Non-Smad pathways in TGF- β signaling. *Cell research* 19: 128-139, 2009.
60. **Zhou X-B, Lai L-F, Xie G-B, Ding C, Xu X, and Wang Y.** LncRNA GAS5 sponges miRNA-221 to promote neurons apoptosis by up-regulated PUMA under hypoxia condition. *Neurological Research* 1-9, 2019.

Figure Legends:

Figure 1: LncRNA GAS5 was downregulated in fibroblasts by TGF- β . A-D), TGF- β (5ng/ml) induced the expression of myofibroblast cell marker CollA and α SMA in 3T3 fibroblasts (A-B) and primary skin fibroblasts (C-D). **B)** Quantification of protein expression in **A** by normalizing to α -Tubulin. **D)** Quantification of protein expression in **C** by normalizing to α -Tubulin. **E)** TGF- β down-regulated GAS5 in 3T3 fibroblasts in a dose-dependent manner. 3T3 fibroblasts were cultured in serum-free DMEM with 0, 1, 2, 5, or 10 ng/ml of TGF- β for 12 hrs. GAS5 was detected by RT-qPCR. **F)** TGF- β (5ng/ml) down-regulated GAS5 in primary mouse skin fibroblasts. **G)** GAS5 was down-regulated in skin tissues with bleomycin treatment, as analyzed by RT-qPCR. C57BL/6 mice were injected subcutaneously with bleomycin (Bleo; 0.02 U) every other day for 14 days. GAS5 RNA expression was calculated by normalizing to cyclophilin mRNA. **H)** GAS5 positive cells in skin tissues were decreased by bleomycin treatment. Skin tissues from (G) was fixed in 4% PFA, and GAS5 positive cells were shown by RNA-FISH staining. Positive and false-positive staining were indicated by white and yellow arrows, respectively. **I)** Quantification of GAS5-expressing cells by counting the positive staining from 10 different fields, shown as the fold change. Bar: 200 μ m. * p<0.05; ** p<0.01; n=3-5. All values are presented as means $\pm$ SEM. T-tests were performed for B, D, F, G, I, and one-way ANOVA test was performed for E.

Figure 2. GAS5 suppressed TGF- β -induced fibroblast-myofibroblast transition. A-B) GAS5 suppressed α SMA and CollA protein expression. 3T3 cells were transduced with AdGFP or AdGAS5 followed by vehicle or 5 ng/ml of TGF- β treatment for 24 hrs. α SMA and CollA expression was assessed by Western blot (A) and quantified by normalizing to α -

Tubulin (B). **C-D**) Knockdown of GAS5 increased α SMA and Col1A protein expression. 3T3 cells were transfected with siCtrl or siGAS5 followed by vehicle or 5 ng/ml of TGF- β treatment for 24 hrs. α SMA and Col1A expression was assessed by Western blot (C) and quantified by normalizing to α -Tubulin (D). **E**) GAS5 suppressed TGF- β -induced α SMA and Col1A mRNA expression. 3T3 cells were transduced with AdGFP or AdGAS5 followed by vehicle (Basal) or 5 ng/ml of TGF- β treatment for 12 hrs. mRNA expression was assessed by RT-qPCR and normalized to cyclophilin. **F**) Knockdown of GAS5 increased α SMA and Col1A mRNA expression. 3T3 cells were transfected with siCtrl or siGAS5 followed by vehicle (Basal) or 5 ng/ml of TGF- β treatment for 12 hrs. mRNA expression was assessed by qPCR and normalized to cyclophilin. **G**) GAS5 suppressed TGF- β -induced α SMA promoter activity. 3T3 cells were transduced with AdGFP or AdGAS5 and transfected with α SMA promoter luciferase reporter for 24 hours prior to the treatment with vehicle (-) or 5 ng/ml of TGF- β for 8 hours. Luciferase assays were performed. NS: not significant; * $p < 0.05$; ** $p < 0.01$; $n = 3$. All values are presented as means $\pm$ SEM. one-way ANOVA tests were performed.

Figure 3. GAS5 promoted Smad3 dephosphorylation through facilitating PPM1A binding to Smad3. **A**) GAS5 accelerated Smad3 dephosphorylation in TGF- β -treated fibroblasts. 3T3 cells were transduced with AdGFP or AdGAS5 followed by vehicle (0 h) or 5 ng/ml of TGF- β treatment for 2 or 4 h. Phospho- and total Smad2 and Smad3 were detected by Western blot. **B**) Phospho-Smad2 and Phospho-Smad3 were quantified by normalizing to their total protein, respectively. **C**) GAS5 bound to Smad3 as shown by RIP assay. Smad3- and Smad4-interacting molecules in 3T3 cells were pulled down by their antibodies, respectively. The presence of GAS5 was detected via qPCR. **D**) GAS5 bound PPM1A in 3T3 fibroblasts. PPM1A-interacting molecules in 3T3 cells was pulled down by its antibody, and the presence of GAS5 was detected by qPCR. **E-F**) Overexpression of GAS5 promoted while Knockdown of GAS5 suppressed Smad3-PPM1A interaction. 3T3 cells were transduced with AdGFP or AdGAS5 (E) or transfected with siCtrl or siGAS5 (F) as indicated for 24 h. Co-immunoprecipitation was performed by using PPM1A antibody, and the presence of Smad3 was detected by Western blot. ** $p < 0.01$; $n = 3$. All values are presented as means $\pm$ SEM. One-way ANOVA tests were performed for B, C, and T-test was performed for D.

Figure 4. GAS5 regulated myofibroblast transition through PPM1A-induced Smad3 dephosphorylation. **A-B**) Knockdown of PPM1A rescued the inhibitory effect of GAS5 on Smad3 phosphorylation and myofibroblast marker gene expression. 3T3 cells were transduced with AdGFP or AdGAS5 along with transfection of siCtrl or siPPM1A followed by vehicle (-) or TGF- β treatment (5 ng/ml) for 24 hours. Protein expression was assessed by Western blot (A) and quantified by normalizing to α -Tubulin (for α SMA and Col1A) or total Smad3 (for pSmad3) (B). **C-D**) Overexpression of Smad3 rescued GAS5-blocked Col1A expression. 3T3 cells were transduced with AdGFP or AdGAS5 and transfected with control or Smad3 expression plasmid followed by vehicle or TGF- β treatment (5 ng/ml) for 24 hours. Col1A expression was assessed by Western blot (C) and quantified by normalizing to α -Tubulin (D). **E-F**) Smad3 inhibitor SIS3 blunted GAS5 knockdown-enhanced Col1A expression. 3T3 cells were transfected with siCtrl or siGAS5 followed by vehicle or SIS3 (10 μ M) treatment for 24 hours. Col1A expression was assessed by Western blot (E) and quantified by normalizing to α -Tubulin (F). ** $p < 0.01$; $n = 3$. All values are presented as means $\pm$ SEM. One-way ANOVA tests were performed.

Figure 5. GAS5 suppressed skin fibrosis in mice. **A-D**) Forced expression of GAS5 via local adenoviral delivery suppressed bleomycin-induced skin fibrosis in mice. Mouse skins were treated with vehicle (PBS) or bleomycin (0.02U/day) every other day for 28 days. The skin

tissues were stained with H&E for structural changes (A), Masson's trichrome (MT) for collagen deposition (C). Bar: 200 μ m. Skin thicknesses shown in A were averaged from 10 different fields (B), and collagen deposition in C was quantified by measuring the staining intensity from 10 different fields (D). **E-H**) Forced expression of GAS5 via local adenoviral delivery suppressed bleomycin-induced expression of Col1A (E-F) and α SMA (G-H). Bar: 100 μ m. Skin tissue sections underwent immunohistochemistry (IHC) staining using Col1A (E) and α SMA (G) antibodies, respectively. Col1A (F) and α SMA (H)-positive cells were averaged from 10 different fields. The large rectangle inserts are enlarged images from the small rectangle boxes in C, E and G, respectively. * $p < 0.05$; ** $p < 0.01$; $n = 5$. All values are presented as means $\pm$ SEM. One-way ANOVA tests were performed.

Figure 6. GAS5 inhibited Col1A and α SMA protein expression in fibrotic skin tissues and attenuated Smad3 binding to their promoters in vivo. **A)** GAS5 suppressed Col1A and α SMA protein expression in skin tissues with bleomycin-induced fibrosis. **B)** Col1a and α SMA protein levels in (A) were quantified by normalizing to GAPDH. ** $p < 0.01$; $n = 5$. **C-F)** GAS5 suppressed Smad3 binding to Col1a (C-D) and α -SMA (E-F) promoters in vivo that were significantly enriched during bleomycin-induced skin fibrosis. Smad3 binding to Col1a (C) and α -SMA (E) promoters in a chromatin setting was measured by in vivo chromatin immunoprecipitation (CHIP) assay. Smad3 binding enrichments were quantified by qPCR relative to the Smad3 binding in vehicle-treated skin tissues (D-F). ** $p < 0.01$ vs AdGFP-treated groups; $n = 5$. All values are presented as means $\pm$ SEM. One-way ANOVA tests were performed.

Figure 7. A diagram of GAS5 function in TGF- β -induced skin fibrosis. During the onset of skin fibrosis, infiltrating immune cells secrete TGF- β , which activates resident fibroblasts through Smad3 signaling pathway. GAS5 suppresses the progression of fibrosis by promoting Smad3 dephosphorylation through facilitating PPM1A binding to Smad3, thus inhibiting Smad3-mediated myofibroblast activation. GAS5 also inhibits fibroblast proliferation through impeding JNK signaling pathway.

Fig 1




Fig 2




Fig 3




Fig 4




Fig 5




Fig 6




Fig 7

