

Evolutionary transcriptomics implicates *HAND2* in the origins of implantation and regulation of gestation length

Mirna Marinić^{1,2}, Katelyn Mika^{1,2}, Sravanthi Chigurupati^{1,3}, and Vincent J. Lynch^{4*}

¹ Department of Human Genetics, University of Chicago, 920 E 58th Street, Chicago, IL, 60637, USA.

² Current address: Department of Organismal Biology and Anatomy, University of Chicago, 1025 E 57th Street, Chicago, IL, 60637, USA.

³ Current address: AbbVie, 1N Waukegan Rd, North Chicago, IL, 60064-6119

⁴ Department of Biological Sciences, University at Buffalo, SUNY, 551 Cooke Hall, Buffalo, NY, 14260, USA.

*Correspondence: vjlynch@buffalo.edu

Abstract

The developmental origins and evolutionary histories of cell types, tissues and organ systems contribute to the ways in which their dysfunction leads to disease. In mammals for example, the nature and extent of maternal-fetal interactions, how those interactions develop, and their evolutionary history likely influence diseases of pregnancy such as infertility and preterm birth. Here we show genes that evolved to be expressed at the maternal-fetal interface in Eutherian ('Placental') mammals play essential roles in the evolution of pregnancy and are associated with immune system disorders and preterm birth. Among these genes is the transcription factor *HAND2*, which suppresses estrogen signaling, an innovation of Eutherians, thereby allowing blastocyst implantation. We found that *HAND2* is dynamically expressed in the decidua throughout the menstrual cycle and pregnancy, gradually decreasing to reach a low at term. *HAND2* regulates a small but distinct set of target genes in endometrial stromal fibroblasts including the cytokine *IL15*, which was also dynamically expressed throughout the menstrual cycle and gestation, and promoted the migration of natural killer cells and extravillous cytotrophoblasts. Remarkably, we found that the *HAND2* promoter loops to a distal enhancer containing SNPs implicated in the regulation of gestation length and birth weight. Collectively, these data connect *HAND2* expression at the maternal-fetal interface with the evolution of implantation and gestation length regulation, and preterm birth.

Introduction

The ontogeny and evolutionary history of cell types, tissues and organ systems, as well as the life histories of organisms biases the ways in which dysfunctions in those systems underlie disease (Varki, 2012). Thus a mechanistic understanding of how cells, tissues and organs evolved their functions, and how organism's life histories influence them, may provide clues to the molecular etiologies of disease. The most common way of utilizing evolutionary information to characterize the genetic architecture of disease is to link genetic variation within a species to phenotypes using quantitative trait loci (QTL) or genome wide association studies (GWAS). An alternative approach is to identify fixed genetic differences between species that are phylogenetically correlated with different disease relevant phenotypes. While the risk of cancer increases with age across mammals, for example, the prevalence of cancer types varies by species (Abegglen *et al.*, 2015), likely because of differences in genetic susceptibility to specific cancers, structure of organ and tissue systems, and life exposures to carcinogens (Varki and Varki, 2015). Similarly, the risk of cardiovascular disease (CVD) increases with age across species, but the pathophysiology of CVD can differ even between closely related taxa such as humans, in which CVD predominantly results from coronary artery atherosclerosis, and the other Great Apes (Hominids), in which CVD is most often associated with interstitial myocardial fibrosis (Varki *et al.*, 2009).

Extant mammals span major stages in the origins and diversification of pregnancy, thus a mechanistic understanding of how pregnancy originated and diverged may provide unique insights into the ontogenetic origins of pregnancy disorders. The platypus and echidna (Monotremes) are oviparous, but the embryo is retained in the uterus for 10–22 days, during which the developing fetus is nourished by maternal secretions delivered through a simple placenta, prior to the laying of a thin, poorly mineralized egg that hatches in ~2 weeks (Hill, 1936). Live birth (viviparity) evolved in the stem-lineage of Therian mammals, but Marsupials and Eutherian ('Placental') mammals have dramatically different reproductive strategies. In Marsupials, pregnancies are generally short (~25 days) and completed within the span of a single estrous cycle (Renfree and Shaw, 2001; Renfree, 2010). Eutherians, in contrast, evolved a suite of traits that support prolonged pregnancies (up to 670 days in African elephant), including an interrupted estrous cycle, which allows for gestation lengths longer than a single reproductive cycle, maternal-fetal communication, maternal recognition of pregnancy, implantation of the blastocyst and placenta into uterine tissue, differentiation (decidualization) of

endometrial stromal fibroblasts (ESFs) in the uterine lining into decidual stromal cells (DSCs), and maternal immunotolerance of the antigenically distinct fetus, i.e. the fetal allograft (Guleria and Pollard, 2000; Moffett and Loke, 2004; Erlebacher, 2013).

Gene expression changes at the maternal-fetal interface underlie evolutionary differences in pregnancy (Hou *et al.*, 2012; Lynch *et al.*, 2015; Armstrong *et al.*, 2017), and thus likely also pathologies of pregnancy such as infertility, recurrent spontaneous abortion (Kosova *et al.*, 2015), preeclampsia (Elliot, 2017; Arthur *et al.*, 2018; Varas Enriquez, McKerracher and Elliot, 2018) and preterm birth (Plunkett *et al.*, 2011; Swaggart, Pavlicev and Muglia, 2015; LaBella *et al.*, 2019). Here, we assembled a collection of gene expression data from the pregnant/gravid maternal-fetal interface of tetrapods and used evolutionary methods to reconstruct gene expression changes during the origins of mammalian pregnancy. We found that genes that evolved to be expressed at the maternal-fetal interface in the Eutherian stem-lineage were enriched for immune functions and diseases, as well as preterm birth. Among the recruited genes was the transcription factor *Heart- and neural crest derivatives-expressed protein 2 (HAND2)*, which plays essential roles in neural crest development (Srivastava *et al.*, 1997), cardiac morphogenesis (Srivastava *et al.*, 1997; Shen *et al.*, 2010; Tamura *et al.*, 2013; Lu *et al.*, 2016; Sun *et al.*, 2016), and suppressing estrogen signaling during the period of uterine receptivity to implantation (Huyen and Bany, 2011; Li *et al.*, 2011; Shindoh *et al.*, 2014; Fukuda *et al.*, 2015; Mestre-Citrinovitz *et al.*, 2015; Murata *et al.*, 2019; Šućurović *et al.*, 2020). We determined that *HAND2* expression at the first trimester maternal-fetal interface was almost entirely restricted to cell types in ESF lineage, and is regulated by multiple transcription factors that control progesterone responsiveness. Moreover, the *HAND2* promoter loops to an enhancer with single nucleotide polymorphisms (SNPs) that have been implicated by GWAS in the regulation of gestation length (Warrington *et al.*, 2019; Sakabe *et al.*, 2020). Furthermore, we showed that *HAND2* regulates interleukin 15 (*IL15*) expression in ESFs, and that ESF-derived *IL15* influences the migration of natural killer and trophoblast cells. These data suggest that *HAND2* and *IL15* signaling played an important role in the evolution of implantation and regulation of gestation length.

Results

Genes that evolved endometrial expression in Eutherian mammals are enriched in immune functions

We previously used comparative transcriptomics to reconstruct the evolution of gene expression at the maternal-fetal interface during the origins of mammalian pregnancy (Lynch *et al.*, 2015). Here, we assembled a collection of new and existing transcriptomes from the pregnant/gravid endometria of 15 Eutherian mammals, three Marsupials, one Monotreme (platypus), two birds, six lizards, and one amphibian (**Supplementary Table 1**). The complete dataset includes expression information for 21,750 genes from 28 species. Next, we transformed continuous transcript abundance estimates values into discrete character states such that genes with Transcripts Per Million (TPM) ≥ 2.0 were coded as expressed (state=1), genes with TPM < 2.0 were coded as not expressed (state=0), and genes without data in specific species were coded as missing (state=?). We then used parsimony to reconstruct ancestral transcriptomes and trace the evolution of gene expression gains ($0 \rightarrow 1$) and losses ($1 \rightarrow 0$) in the endometrium.

We identified 958 genes that potentially evolved endometrial expression in the Eutherian stem-lineage (**Figure 1A**), including 149 that unambiguously evolved endometrial expression (**Supplementary Table 2**). These 149 genes were significantly enriched in pathways related to the immune system (**Supplementary Table 3**), although only two pathways were enriched at False Discovery Rate (FDR) ≤ 0.10 , namely, “Cytokine Signaling in Immune System” (hypergeometric $P=1.97 \times 10^{-5}$, FDR=0.054) and “Signaling by Interleukins” (hypergeometric $P=4.09 \times 10^{-5}$, FDR=0.067). Unambiguously recruited genes were also enriched in numerous human phenotype ontology terms (**Supplementary Table 4**) but only two, “Immune System Diseases” (hypergeometric $P=3.15 \times 10^{-8}$, FDR=2.52x10⁻⁴) and “Preterm Birth” (hypergeometric $P=4.04 \times 10^{-4}$, FDR=8.07x10⁻⁴), at FDR ≤ 0.10 . In contrast, these genes were enriched in numerous biological process gene ontology (GO) terms (**Supplementary Table 5**), nearly all of which were related to regulation of immune system, including “Leukocyte Migration” (hypergeometric $P=1.17 \times 10^{-7}$, FDR=1.29x10⁻³), “Inflammatory Response” (hypergeometric $P=8.07 \times 10^{-7}$, FDR=2.21x10⁻³), and “Cytokine-mediated Signaling Pathway” (hypergeometric $P=2.18 \times 10^{-5}$, FDR=0.013).

Recruitment of *HAND2* and anti-estrogenic signaling in Eutherians

Among the genes that unambiguously evolved endometrial expression in the Eutherian stem-lineage was the basic helix-loop-helix family transcription factor *Heart- and neural crest derivatives-expressed protein 2 (HAND2)*. *HAND2* plays an essential role in mediating the anti-estrogenic action of progesterone and the establishment of uterine receptivity to implantation (**Figure 1B**) (Huyen and Bany, 2011; Li *et al.*, 2011; Fukuda *et al.*, 2015; Mestre-Citrinovitz *et al.*, 2015), suggesting an active role of this hormone during pregnancy in non-Eutherian mammals. To test this, we performed immunohistochemistry (IHC) on endometrial sections from day 12.5 of pregnancy (after the transition from the histotrophic to placental phase) in the short-tailed opossum (*Monodelphis domestica*). We observed strong staining for estrogen receptor alpha (*ESR1*; ER α) phosphorylated at serine 118 (a mark of transcriptionally active ER α) (Kato *et al.*, 1995), phosphorylated ERK1/2, and MUC1 (**Figure 1C**). We also found that *ESR1*, *FGF2*, *FGF9*, *FGF18*, *MUC1*, several *WNT* genes that stimulate proliferation of the luminal epithelia, and *MKI67* (Ki-67), a mark of actively proliferating cells (Scholzen and Gerdes, 2000), were abundantly expressed in RNA-Seq data from day 12.5 pregnant *M. domestica* endometrium (**Figure 1C**). These data are indicative of persistent estrogen signaling during pregnancy in opossum.

***HAND2* is expressed in endometrial stromal fibroblast lineage cells**

To determine which cell types at the human maternal-fetal interface express *HAND2*, we used previously generated single-cell RNA-Seq (scRNA-Seq) data from first trimester decidua (Vento-Tormo *et al.*, 2018). *HAND2* expression was almost entirely restricted to cell populations in the endometrial stromal fibroblast lineage (ESF1, ESF2, and DSC), with particularly high expression in ESF2s and DSCs (**Figure 2A**). *HAND2* protein was localized to nuclei in ESF lineage cells in human pregnant decidua (**Figure 2B**) from Human Protein Atlas IHC data (Uhlén *et al.*, 2015). We also used existing functional genomics data to explore the regulatory status of the *HAND2* locus (see Methods). Consistent with active expression, the *HAND2* locus in human DSCs is marked by histone modifications that typify enhancers (H3K27ac) and promoters (H3K4me3), and is located in a region of open chromatin as assessed by ATAC-, DNaseI- and FAIRE-Seq. Additionally, it is bound by transcription factors that establish endometrial stromal cell-type identity and mediate decidualization, including the progesterone receptor (PR), NR2F2 (COUP-TFII), GATA2, FOSL2, FOXO1, as well as polymerase II (**Figure 2C**). The *HAND2* promoter loops to several distal enhancers, as assessed by H3K27ac ChIP data generated from a normal hTERT-immortalized endometrial cell line (E6E7hTERT),

including a region bound by PR, NR2F2, GATA2, FOSL2 and FOXO1, that also contains SNPs associated with gestation length in recent GWAS (Warrington *et al.*, 2019; Sakabe *et al.*, 2020) (**Figure 2C**). *HAND2* was significantly up-regulated by decidualization of human ESFs into DSCs by cAMP/progesterone treatment ($\text{Log}_2\text{FC}=1.28$, $P=2.62\times 10^{-26}$, $\text{FDR}=1.16\times 10^{-24}$), and significantly down-regulated by siRNAs targeting PR ($\text{Log}_2\text{FC}=-0.90$, $P=7.05\times 10^{-15}$, $\text{FDR}=2.03\times 10^{-13}$) and GATA2 ($\text{Log}_2\text{FC}=-2.73$, $P=0.01$, $\text{FDR}=0.19$) (**Figure 2D**). In contrast, siRNA-mediated knockdown of neither NR2F2 ($\text{Log}_2\text{FC}=-0.91$, $P=0.05$, $\text{FDR}=1.0$) nor FOXO1 ($\text{Log}_2\text{FC}=0.08$, $P=0.49$, $\text{FDR}=0.74$) significantly altered *HAND2* expression (**Figure 2D**).

Differential *HAND2* expression throughout the menstrual cycle and pregnancy

Our observation that *HAND2* is progesterone responsive suggests it may be differentially expressed throughout the menstrual cycle and pregnancy. To explore this possibility, we utilized previously published gene expression datasets generated from the endometrium across the menstrual cycle (Talbi *et al.*, 2006) and from the basal plate from mid-gestation to term (Winn *et al.*, 2007). *HAND2* expression tended to increase from proliferative through the early and middle secretory phases, reaching a peak in the late secretory phase of the menstrual cycle (**Figure 2E**). In stark contrast, *HAND2* expression decreases throughout pregnancy reaching a low at term (**Figure 2F**). For comparison, 9% of genes were down-regulated between weeks 14-19 and 37-40 of pregnancy ($\text{FDR}\leq 0.10$). We also used previously published gene expression datasets (see Methods) to explore if *HAND2* was associated with disorders of pregnancy and found significant *HAND2* dysregulation in the endometria of women with infertility (IF) and recurrent spontaneous abortion (RSA) compared to fertile controls (**Figure 2G**). *HAND2* was not differentially expressed in ESFs or DSCs from women with preeclampsia (PE) compared to controls (**Figure 2G**).

HAND2 regulates a distinct set of target genes

HAND2 expression has previously been shown to play a role in orchestrating the transcriptional response to progesterone during decidualization in human and mouse DSCs (Huyen and Bany, 2011; Li *et al.*, 2011; McConaha *et al.*, 2011; Shindoh *et al.*, 2014). However, whether *HAND2* has functions in other endometrial stromal lineage cells such as ESFs is unknown. Therefore, we used siRNA to knockdown *HAND2* expression in human hTERT-immortalized ESFs (T-HESC) and assayed global gene expression changes by RNA-Seq 48 hours after knockdown. We found that *HAND2* was knocked-down $\sim 78\%$ ($P=7.79\times 10^{-3}$) by

siRNA treatment (**Figure 3A**), which dysregulated the expression of 553 transcripts (489 genes) at FDR ≤ 0.10 (**Figure 3A** and **Supplementary Table 6**). Genes dysregulated by *HAND2* knockdown had very little overlap with genes dysregulated by *PR*, *NR2F2*, *GATA2*, or *FOXO1* knockdown (**Figure 3B**). These data indicate that *HAND2* regulates a distinct set of target genes compared to transcription factors that establish cell-type identity and mediate the decidualization response. Indeed, only 30 genes are co-regulated by cAMP/progesterone, *PR*, *GATA2*, and *HAND2* (**Figure 3B**).

Genes dysregulated by *HAND2* knockdown were enriched in several pathways and human phenotype ontologies relevant to endometrial stromal cells and pregnancy in general (**Figure 3C**, **Supplementary Table 7**, and **Supplementary Table 8**). Enriched pathways play a role in decidualization, such as “Wnt Signaling” (Peng *et al.*, 2008; Hayashi *et al.*, 2009; Sonderegger, Pollheimer and Knöfler, 2010; Franco *et al.*, 2011; Wang *et al.*, 2013), “BMP Signaling” (Ying and Zhao, 2000; Lee *et al.*, 2007; Li *et al.*, 2007; Wetendorf and DeMayo, 2012), “ErbB Signaling” (Lim, Dey and Das, 1997; Klonisch *et al.*, 2001; Large *et al.*, 2014), “TGF-beta Receptor Signaling” (Jones *et al.*, 2006; Li, 2014; Ni and Li, 2017) and “BMP2-WNT4-FOXO1 Pathway in Human Primary Endometrial Stromal Cell Differentiation” (Gellersen and Brosens, 2003; Buzzio *et al.*, 2006; Lee *et al.*, 2007; Li *et al.*, 2007; Brayer, Lynch and Wagner, 2009; Lynch *et al.*, 2009; Kajihara, Brosens and Ishihara, 2013). Other enriched pathways included those involved in placental bed development disorders and preeclampsia (Chekir *et al.*, 2006; Oliver *et al.*, 2011; Guedes-Martins *et al.*, 2013), the induction of pro-inflammatory factors via nuclear factor- κ B (NF κ B) through the “AGE/RAGE pathway” (Lappas, Permezel and Rice, 2007), the “Aryl Hydrocarbon Receptor Pathway”, which mediates maternal immunotolerance of the fetal allograft (Munn *et al.*, 1998; Abbott *et al.*, 1999; Funeshima *et al.*, 2005; Hao *et al.*, 2013), “Circadian Rhythm Related Genes”, which have been associated with implantation (Greenhill, 2014) and parturition (Roizen *et al.*, 2007; Olcese, 2012; Olcese, Lozier and Paradise, 2013; Menon *et al.*, 2016), and the “RAC1/PAK1/p38/MMP2” pathway, which has also been implicated in decidual inflammation, senescence and parturition (Menon *et al.*, 2016). Enriched human phenotype ontology terms were related to complications of pregnancy, including “Premature Rupture of Membranes”, “Premature Birth”, “Toxemia of Pregnancy” (preeclampsia) and “Abnormal Delivery”. We also observed that several genes in the NF κ B pathway, such as *MYD88*, *CHUK*, *IkBKE*, *NFkBIE* and *RTKN2* were differentially expressed; NF κ B signaling has been associated with the molecular etiology of preterm birth (Allport, 2001;

Lindstrom and Bennett, 2005).

HAND2 regulates IL15 expression in endometrial stromal fibroblast lineage cells

Among the genes dysregulated by *HAND2* knockdown in ESFs was *IL15* ($\text{Log}_2\text{FC}=7.98$, $P=7.91\times 10^{-8}$, $\text{FDR}=1.49\times 10^{-5}$), a pleiotropic cytokine previously shown to be expressed in the endometrium and decidua (Figure 3A and Table 1) (Kitaya *et al.*, 2000; Okada, 2000; Dunn, Critchley and Kelly, 2002; Okada *et al.*, 2004; Godbole and Modi, 2010). *IL15* was robustly expressed at the first trimester maternal-fetal interface in stromal fibroblast lineage cells (Figure 4A) and there was a general correlation between *HAND2* and *IL15* expression in single cells (Figure 4A inset) (Vento-Tormo *et al.*, 2018). *IL15* protein localized to cytoplasm in ESF lineage cells in human pregnant decidua (Figure 4B) in Human Protein Atlas data. The *IL15* promoter loops to several distal sites in H3K27ac HiChIP data from E6E7hTERT endometrial cells including to regions bound by PR, NR2F2, GATA2, FOSL2, FOXO1, and SRC2, an intrinsic histone acetyltransferase that is a transcriptional co-factor of ligand-dependent hormone receptors (Figure 4C). The *IL15* promoter also loops to a putative enhancer in its first intron that contains a PGR binding site and SNPs marginally associated with a maternal effect on offspring birth weight (rs190663174, $P=6\times 10^{-4}$) by GWAS (Warrington *et al.*, 2019). *IL15* was significantly up-regulated by *in vitro* decidualization of human ESFs into DSCs by cAMP/progesterone treatment ($\text{Log}_2\text{FC}=2.15$, $P=2.58\times 10^{-33}$, $\text{FDR}=1.59\times 10^{-31}$), and significantly down-regulated by siRNAs targeting PR ($\text{Log}_2\text{FC}=-1.24$, $P=6.23\times 10^{-15}$, $\text{FDR}=1.80\times 10^{-13}$) and GATA2 ($\text{Log}_2\text{FC}=-2.08$, $P=4.16\times 10^{-3}$, $\text{FDR}=0.14$) (Figure 4D), but not NR2F2 ($\text{Log}_2\text{FC}=0.19$, $P=0.38$, $\text{FDR}=0.93$) or FOXO1 ($\text{Log}_2\text{FC}=0.29$, $P=0.04$, $\text{FDR}=0.22$) (Figure 4D). Although *HAND2* binding data is not available for human stromal fibroblast lineage cells, several *HAND2* binding motifs (≥ 0.85 motif match) are located within enhancers that loop to the *IL15* promoter.

Differential IL15 expression throughout the menstrual cycle and pregnancy

Our observations that *HAND2* is progesterone responsive and differentially expressed throughout the menstrual cycle and pregnancy suggests that *IL15*, which is controlled by *HAND2* as well as cAMP/progesterone/PR/GATA2, may be similarly regulated. Indeed, like *HAND2*, we found that *IL15* expression increased as the menstrual cycle progressed, peaking in the middle-late secretory phases (Figure 4E) and decreased throughout pregnancy reaching a low at term (Figure 4F). *IL15* expression was also dysregulated in the endometria of women with infertility but not recurrent spontaneous abortion, compared to fertile controls (Figure 4G).

In women with preeclampsia, *IL15* was not dysregulated in ESFs, but it was expressed significantly higher in DSCs, compared to controls. Thus, like *HAND2*, *IL15* is differentially expressed throughout the menstrual cycle and pregnancy, and in the endometria of women with infertility.

ESF-derived IL15 promotes NK and trophoblast migration

Endometrial stromal cells promote the migration of uterine natural killer (uNK) (Chen *et al.*, 2011) and trophoblast cells (Graham and Lala, 1991; Paiva *et al.*, 2009; Zhu *et al.*, 2009; Godbole *et al.*, 2011). IL15, in particular, stimulates the migration of uNK cells (Allavena *et al.*, 1997; Verma *et al.*, 2000; Ashkar *et al.*, 2003; Barber and Pollard, 2003; Kitaya, Yamaguchi and Honjo, 2005) and the human choriocarcinoma cell line, JEG-3 (Zygmunt *et al.*, 1998). Therefore, we tested whether ESF-derived IL15 influenced the migration of primary human NK cells and immortalized first trimester extravillous trophoblasts (HTR-8/SVneo) in trans-well migration assays (**Figure 5A**). We found that ESF media supplemented with recombinant human IL15 (rhIL15) was sufficient to stimulate the migration of NK and HTR-8/SVneo cells to the lower chamber of trans-wells, compared to non-supplemented media (**Figure 5B,C; Supplementary Tables 9 and 10**). Conditioned media from ESFs with siRNA-mediated *HAND2* knockdown increased migration of both NK and HTR-8/SVneo compared to negative control (i.e., non-targeting siRNA) (**Figure 5B,C**). Conditioned media from ESFs with siRNA-mediated *IL15* knockdown reduced migration of both NK and HTR-8/SVneo cells compared to negative control (**Figure 5B,C**). Similarly, ESF conditioned media supplemented with anti-IL15 antibody reduced cell migration compared to media supplemented with control IgG antibody (**Figure 5B,C**).

Discussion

Eutherian mammals evolved a suite of traits that support pregnancy, including an interrupted estrous cycle allowing for prolonged gestation lengths, maternal-fetal communication, implantation, maternal immunotolerance and recognition of pregnancy, and thus are uniquely afflicted by disorders of these processes. When searching for clues as to how variation in normal physiological functions can lead to dysfunction and disease, deeper understanding of the evolutionary and developmental histories of organ and tissues has the potential to provide novel insights. Here, we used evolutionary transcriptomics to identify genes that evolved to be expressed on the maternal side (endometrium) of the maternal-fetal interface

during the origins of pregnancy in Eutherians, and hence may also contribute to pregnancy complications such as infertility, recurrent spontaneous abortion, and preterm birth.

Among the genes recruited into endometrial expression in Eutherians we identified *HAND2*, a pleiotropic transcription factor that plays an essential role in suppressing estrogen signaling at the time of uterine receptivity to blastocyst embedding, through its down-regulation of pro-estrogenic genes and by directly inhibiting the transcriptional activities of the estrogen receptor (Huyen and Bany, 2011; Li *et al.*, 2011; Shindoh *et al.*, 2014; Fukuda *et al.*, 2015; Mestre-Citrinovitz *et al.*, 2015; Murata *et al.*, 2019). Consistent with these functions, we found evidence of estrogen activity in the endometrium of pregnant opossum. Earlier research detected similar activity in the gravid oviduct of birds and reptiles (Means *et al.*, 1975; Kato *et al.*, 1992; Girling, 2002; González-Morán, 2015). These data indicate that suppression of estrogen signaling during the window of uterine receptivity to implantation is an evolutionary innovation of Eutherian mammals, which involved the recruitment of *HAND2* and its anti-estrogenic functions into endometrial expression.

The roles of *HAND2* in DSCs and implantation are well-understood (Huyen and Bany, 2011; Li *et al.*, 2011; Shindoh *et al.*, 2014; Fukuda *et al.*, 2015; Mestre-Citrinovitz *et al.*, 2015; Murata *et al.*, 2019; Šućurović *et al.*, 2020). In contrast, the function(s) of *HAND2* at other stages of pregnancy and in ESFs remain relatively unexplored. We knocked down *HAND2* in ESFs and found that downstream dysregulated genes were enriched for human phenotype ontologies related to disorders of pregnancy, including “Premature Rupture of Membranes”, “Premature Birth”, and “Abnormal Delivery”, suggesting that *HAND2* has functions throughout pregnancy and in parturition. Indeed, we discovered that SNPs recently implicated in the regulation of gestation length and birth weight by GWAS (Warrington *et al.*, 2019; Sakabe *et al.*, 2020) make long-range interactions to the *HAND2* promoter. Also of note, *HAND2* expression is significantly higher in placental villous samples from idiopathic spontaneous preterm birth (isPTB) compared to term controls (Brockway *et al.*, 2019). However, this difference may be related to gestational age rather than the etiology of PTB (Eidem *et al.*, 2016).

Additionally, analyzing previously published datasets we noticed that *HAND2* expression decreases throughout gestation. Unlike the majority of Eutherians, where parturition closely follows the significant drop in progesterone concentrations in maternal peripheral blood, this is

not the case for humans and other Old-World primates (Ratajczak, Fay and Muglia, 2010), where progesterone levels keep rising throughout gestation, reaching maximum at birth. These observations, combined with the central role of *HAND2* in mediating the anti-estrogenic actions of progesterone, suggest decreased *HAND2* at the end of pregnancy may contribute to the estrogen dominant uterine environment at the onset of labor (Pinto *et al.*, 1966; Pepe and Albrecht, 1995; Mesiano *et al.*, 2002; R. Smith *et al.*, 2009; Ratajczak, Fay and Muglia, 2010; Welsh *et al.*, 2012), despite high systemic progesterone. Low *HAND2* at the end of pregnancy in humans is therefore most likely not directly related to progesterone concentrations, suggesting that an unidentified inhibitory signal reduces endometrial *HAND2* expression. While taken collectively these data indicate a role for *HAND2* in pre/term birth, a direct mechanistic link between *HAND2* expression and the timing of parturition remains to be demonstrated.

One of the genes dysregulated by *HAND2* knockdown was the multifunctional cytokine *IL15*, which plays important roles in innate and adaptive immunity. In the context of pregnancy, it is important for the recruitment of uterine natural killer (uNK) cells to the endometrium (Kitaya *et al.*, 2000; Verma *et al.*, 2000; Ashkar *et al.*, 2003; Barber and Pollard, 2003; Kitaya, Yamaguchi and Honjo, 2005; Laskarin *et al.*, 2006). The roles of uNK cells in the remodeling of uterine spiral arteries and regulating trophoblast invasion are well known (Zygmunt *et al.*, 1998; Hanna *et al.*, 2006; S. D. Smith *et al.*, 2009; Burke *et al.*, 2010; Hazan *et al.*, 2010; Lash, Robson and Bulmer, 2010; Bany, Scott and Eckstrum, 2012; Robson *et al.*, 2012; Zhang, Dunk and Lye, 2013; Lima *et al.*, 2014; Fraser *et al.*, 2015; Felker and Croy, 2016; Renaud *et al.*, 2017). Endometrial stromal cell-derived *IL15* is also necessary for the selective targeting and clearance of senescent endometrial stromal cells from the implantation site by uNK cells, which is essential for endometrial rejuvenation and remodeling at embryo implantation (Brighton *et al.*, 2017). Dysregulation of uNK cell-mediated clearance of these senescent cells has also been implicated in recurrent pregnancy loss (Lucas *et al.*, 2020). We found that, like *HAND2*, *IL15* decreases throughout gestation and both genes increase in expression as the menstrual cycle progresses. Unexpectedly however, while previous studies showed that *HAND2* induces *IL15* in DSCs (Shindoh *et al.*, 2014; Murata *et al.*, 2020), we discovered that *HAND2* inhibited *IL15* expression in ESFs, indicating a switch in regulatory activity sometime during early menstrual cycle. These data suggest that *HAND2* regulates the appropriate timing of endometrial *IL15* expression during the menstrual cycle and pregnancy, and thus the appropriate timing of uNK cell recruitment, trophoblast migration and the clearance of senescent endometrial stromal cells

from the implantation site (**Figure 5D**). Additional work is needed to elucidate the mechanisms that underlie change in *HAND2-IL15* dynamics. An intriguing possibility, however, is that progressive cell-state changes during gestation promote the transition from ESFs to DSCs to senescent DSCs (snDSCs), which have much lower *HAND2* and *IL15* expression than DSCs (Lucas *et al.*, 2020). Thus, our observation of low *HAND2* and *IL15* near term may reflect a reduction of anti-inflammatory DSCs (high *HAND2* and *IL15*) and an accumulation of pro-inflammatory snDSCs (low *HAND2* and *IL15*) at the maternal-fetal interface.

Decreased *HAND2* and *IL15* expression near term and their influence on immune cells at the maternal-fetal interface may also play a role in parturition. While the signals that initiate the onset of parturition in humans and other Old-World monkeys are not known, pre/term labor is known to be associated with elevated inflammation and an influx of immune cells into utero-placental tissues (Thomson *et al.*, 1999; Young *et al.*, 2002; Osman *et al.*, 2003; Gomez-Lopez, Guilbert and Olson, 2010; Rinaldi *et al.*, 2011, 2014; Hamilton *et al.*, 2012; Shynlova *et al.*, 2013; Bartmann *et al.*, 2014; Menon *et al.*, 2016; Peters *et al.*, 2016; Wilson and Mesiano, 2020). uNK cells are abundant throughout gestation (Bulmer *et al.*, 1991; King *et al.*, 1991; Moffett-King, 2002; Williams *et al.*, 2009; Bartmann *et al.*, 2014), but whether they play a role in late pregnancy and parturition is unclear. However, depletion of uNK cells rescues LPS-induced preterm birth in *IL10*-null mice (Murphy *et al.*, 2005), indicating they contribute to infection/inflammation-induced preterm parturition (Murphy *et al.*, 2009). CD16⁺CD56^{dim} (cytotoxic) uNK cells have also been observed in the decidua and the placental villi of women with preterm but not term labor, suggesting an association between dysregulation of uNK cells and preterm birth in humans (Gomaa *et al.*, 2017). uNK cells are associated with other pregnancy complications in humans such as fetal growth restriction, preeclampsia, and recurrent spontaneous abortion (Moffett, Regan and Braude, 2004; Hiby *et al.*, 2010; Wallace *et al.*, 2013, 2014; Kieckbusch *et al.*, 2014). Taken together, these data indicate that uNK cells may act downstream of *HAND2-IL15* signaling in the timing of parturition.

Conclusions

Here we show that *HAND2* evolved to be expressed in endometrial stromal cells in the Eutherian stem-lineage, coincident with the evolution of suppressed estrogen signaling during the window of implantation and an interrupted reproductive cycle during pregnancy, which necessitated a means to regulate the length of gestation. Our data suggest that *HAND2* may

contribute to the regulation of gestation length by promoting an estrogen dominant uterine environment near term and through its effect on IL15 signaling and uNK cell function. To further expand our understanding of *HAND2* functions at the molecular mechanistic level, multiple technical and ethical difficulties associated with studying human pregnancy *in vivo* will need to be overcome. Therefore, recently developed organoid models of the human maternal-fetal interface (Rinehart, Lyn-Cook and Kaufman, 1988; Boretto *et al.*, 2017; Turco *et al.*, 2017, 2018; Marinić and Lynch, 2019), which allow for *in vitro* 3D manipulation, will prove instrumental.

Methods

Endometrial gene expression profiling and ancestral transcriptome reconstruction

Data Collection: We obtained previously generated RNA-Seq data from pregnant endometria of amniotes by searching NCBI BioSample, Short Read Archive (SRA), and Gene Expression Omnibus (GEO) databases for anatomical terms referring to the portion of the female reproductive tract (FRT), including “uterus”, “endometrium”, “decidua”, “oviduct”, and “shell gland”, followed by manual curation to identify those datasets that included the FRT region specialized for maternal-fetal interaction or shell formation. Datasets that did not indicate whether samples were from pregnant or gravid females were excluded, as were those composed of multiple tissue types. Species included in this study and their associated RNA-Seq accession numbers are included in **Supplementary Table 1**.

New RNA-Seq data: Endometrial tissue samples from the pregnant uteri of baboon (x3), mouse (x3), hamster (x3), bat (x2), and squirrel (x2) were dissected and mailed to the University of Chicago in RNA-Later. These samples were further dissected to remove myometrium, luminal epithelium, and extra-embryonic tissues, and then washed 3x in ice cold PBS to remove unattached cell debris and red blood cells. Total RNA was extracted from the remaining tissue using the RNeasy Plus Mini Kit (74134, QIAGEN) per manufacturer’s instructions. RNA concentrations were determined by Nanodrop 2000 (Thermo Scientific). A total amount of 2.5 μ g of total RNA per sample was submitted to the University of Chicago Genomics Facility for Illumina Next Gen RNA sequencing. Quality was assessed with the Bioanalyzer 2100 (Agilent). A total RNA library was generated using the TruSEQ stranded mRNA with RiboZero depletion (Illumina) for each sample. The samples were fitted with one of six different adapters with a different 6-base barcode for multiplexing. Completed libraries were run on an Illumina HiSEQ2500 with v4 chemistry on 2 replicate lanes for hamster and 1 lane for everything else of an 8 lane flow cell, generating 30-50 million high quality 50bp single-end reads per sample.

Multispecies RNA-Seq analyses: Kallisto version 0.42.4 was used to pseudo-align the raw RNA-Seq reads to genomes (see **Supplementary Table 1** for reference genome assemblies). We used default parameters bias correction, and 100 bootstrap replicates. Kallisto outputs consist of transcript abundance estimates in transcripts per million (TPM), which were used to determine gene expression.

Ancestral transcriptome reconstruction: We previously showed that genes with $TPM \geq 2.0$ are actively transcribed in endometrium while genes with $TPM < 2.0$ lack hallmarks of active transcription such as promoters marked with H3K4me3 (Wagner, Kin and Lynch, 2012, 2013). Based on these findings, we transformed values of transcript abundance estimates into discrete character states, such that genes with $TPM \geq 2.0$ were coded as expressed (state=1), genes with $TPM < 2.0$ were coded as not expressed (state=0), and genes without data in specific species coded as missing (state=?). The binary encoded endometrial gene expression dataset generally grouped species by phylogenetic relatedness, suggesting greater signal to noise ratio than raw transcript abundance estimates. Therefore, we used the binary encoded endometrial transcriptome dataset to reconstruct ancestral gene expression states and trace the evolution of gene expression gains ($0 \rightarrow 1$) and losses ($1 \rightarrow 0$) in the endometria across vertebrate phylogeny (**Figure 1A**). We used Mesquite (v2.75) with parsimony optimization to reconstruct ancestral gene expression states, and identify genes that gained or lost endometrial expression. Expression was classified as an unambiguous gain if a gene was not inferred as expressed at a particular ancestral node (state 0) but inferred as expressed (state 1) in a descendent of that node, and vice versa for the classification of a loss of endometrial expression. We thus identified 149 genes that unambiguously evolved endometrial expression in the stem-lineage of Placental mammals (**Supplementary Table 2**).

Pathway enrichments: We used WebGestalt v. 2019 (Liao *et al.*, 2019) to determine if the 149 identified genes were enriched in ontology terms using over-representation analysis (ORA). A key advantage of WebGestalt is that it allowed for the inclusion of a custom background gene list, which was the set of 21,750 genes for which we could reconstruct ancestral states, rather than all annotated protein-coding genes in the human genome. We used ORA to identify enriched terms for three pathway databases (KEGG, Reactome, and Wikipathway), the Human Phenotype Ontology database, and a custom database of genes implicated in preterm birth by GWAS. The preterm birth gene set was assembled from the NHGRI-EBI Catalog of published genome-wide association studies (GWAS Catalog), including genes implicated in GWAS with either the ontology terms “Preterm Birth” (EFO_0003917) or “Spontaneous Preterm Birth” (EFO_0006917), as well as two recent preterm birth GWAS (Warrington *et al.*, 2019; Sakabe *et al.*, 2020).

Immunohistochemistry (IHC)

Endometrial tissue from pregnant opossum (12.5 d.p.c.) was fixed in 10% neutral-buffered formalin, paraffin-embedded, sectioned at 4 μ m, and mounted on slides. Paraffin sections were dried at room temperature overnight and then baked for 12h at 50°C. Prior to immunostaining, de-paraffinization and hydration were done in xylene and graded ethanol to distilled water. During hydration, a 5mins blocking for endogenous peroxidase was done in 0.3% H₂O₂ in 95% ethanol. Antigen retrieval was performed in retrieval buffer pH6, using a pressure boiler microwave as a heat source with power set to full, allowing retrieval buffer to boil for 20mins, and then cooled in a cold water bath for 10mins. To stain sections, we used the Pierce Peroxidase IHC Detection Kit (Cat. 36000) following the manufacturers protocol. Briefly, uterine sections were incubated at 4°C overnight with polyclonal antibodies against HAND2 (Santa Cruz SC-9409), MUC1 (Novus Biologicals NB120-15481), p-ER α (Santa Cruz SC-12915), p-Erk1/2 (also known as MAPK1/2; Santa Cruz SC-23759-R) at 1:1000 dilution in blocking buffer. The next day sections were washed 3x in wash buffer, and incubated with HRP-conjugated rabbit anti-mouse IgG (H+L) secondary antibody (Invitrogen cat # 31450) at 1:10000 dilution in blocking buffer. After 30mins at 4°C slides were washed 3x in wash buffer. Slides were developed with 1x DAB/metal concentrate and stable peroxide buffer for 5mins, then rinsed 3x for 3mins in wash buffer, and mounted with Permount (SP15-100; Thermo Fisher Scientific).

Expression of *HAND2* and *IL15* at the maternal-fetal interface

We used previously published single-cell RNA-Seq (scRNA-Seq) data from the human first trimester maternal-fetal interface (Vento-Tormo *et al.*, 2018) to determine which cell types express *HAND2* and *IL15*. The dataset consists of transcriptomes for ~70,000 individual cells of many different cell types, including: three populations of tissue resident decidual natural killer cells (dNK1, dNK2, and dNK3), a population of proliferating natural killer cells (dNKp), type 2 and/or type 3 innate lymphoid cells (ILC2/ILC3), three populations of decidual macrophages (dM1, dM2, and dM3), two populations of dendritic cells (DC1 and DC2), granulocytes (Gran), T cells (TCells), maternal and lymphatic endothelial cells (Endo), two populations of epithelial glandular cells (Epi1 and Epi2), two populations of perivascular cells (PV1 and PV2), two endometrial stromal fibroblast populations (ESF1 and ESF2), and decidual stromal cells (DSCs), placental fibroblasts (fFB1), extravillous- (EVT), syncytiotrophoblasts (SCT), and villus- (VCT) cytotrophoblasts (**Figure 2A**). Data were not reanalyzed, rather previously analyzed data were accessed using the cellxgene website available at <https://maternal-fetal-cellxgene.s3.amazonaws.com>.

interface.cellgeni.sanger.ac.uk.

Expression of HAND2 and IL15 in human decidua cells

We used previously published IHC data for HAND2 and IL15 generated from pregnant human decidua as part of the Human Protein Atlas project (<http://www.proteinatlas.org/>; (Uhlén *et al.*, 2015)). Image/gene/data available from IL15 (<https://www.proteinatlas.org/ENSG00000164136-IL15/tissue>) and HAND2 (<https://www.proteinatlas.org/ENSG00000164107-HAND2/tissue>).

Functional genomic analyses of the *HAND2* and *IL15* loci

Gene expression data: We used previously published RNA-Seq and microarray gene expression data generated from human ESFs and DSCs that were downloaded from National Center for Biotechnology Information (NCBI) Short Read Archive (SRA) and processed remotely using Galaxy platform (<https://usegalaxy.org/>; Version 20.01) (Afgan *et al.*, 2018) for RNA-Seq data and GEO2R. RNA-Seq datasets were transferred from SRA to Galaxy using the Download and Extract Reads in FASTA/Q format from NCBI SRA tool (version 2.10.4+galaxy1). We used HISAT2 (version 2.1.0+galaxy5) (Kim, Langmead and Salzberg, 2015) to align reads to the Human hg38 reference genome using single- or paired-end options depending on the dataset and unstranded reads, and report alignments tailored for transcript assemblers including StringTie. Transcripts were assembled and quantified using StringTie (v1.3.6) (Pertea *et al.*, 2015, 2016), with reference file to guide assembly and the “reference transcripts only” option, and output count files for differential expression with DESeq2/edgeR/limma-voom. Differentially expressed genes were identified using DESeq2 (version 2.11.40.6+galaxy1) (Anders and Huber, 2010; Love, Huber and Anders, 2014). The reference file for StringTie guided assembly was wgEncodeGencodeBasicV33. GEO2R performs comparisons on original submitter-supplied processed data tables using the GEOquery (Davis and Meltzer, 2007) and limma (Smyth *et al.*, 2002) R packages from the Bioconductor project (<https://bioconductor.org/>; (Gentleman *et al.*, 2004)).

Datasets included gene expression profiles of primary human ESFs treated for 48 hours with control non-targeting, PGR-targeting (GSE94036), FOXO1-targeting (GSE94036) or NR2F2 (COUP-TFII)-targeting (GSE52008) siRNA prior to decidualization stimulus for 72 hours; transfection with GATA2-targeting siRNA was followed immediately by decidualization stimulus

(GSE108407). We also explored the expression of *HAND2* and *IL5* in the endometria of women with recurrent spontaneous abortion and infertility using a previously published dataset (GSE26787), as well as in the endometrium throughout the menstrual cycle (GSE4888) and basal plate throughout gestation (GSE5999); *HAND2* probe 220480_at, *IL15* probe 205992_s_at, *HAND2* expression from proliferative and mid-secretory endometria (GSE132713).

ChIP-Seq and open chromatin data: We used previously published ChIP-Seq data generated from human DSCs that were downloaded from National Center for Biotechnology Information (NCBI) Short Read Archive (SRA) and processed remotely using Galaxy (Afgan *et al.*, 2018). ChIP-Seq reads were mapped to the human genome (GRCh37/hg19) using HISAT2 (Kim, Langmead and Salzberg, 2015) with default parameters and peaks called with MACS2 (Zhang *et al.*, 2008; Feng *et al.*, 2012) with default parameters. Samples included PLZF (GSE112362), H3K4me3 (GSE61793), H3K27ac (GSE61793), H3K4me1 (GSE57007), PGR (GSE94038), the PGR A and B isoforms (GSE62475), NR2F2 (GSE52008), FOSL2 (GSE94038), FOXO1 (GSE94037), PolII (GSE94037), GATA2 (GSE108408), SRC-2/NCOA2 (GSE123246), AHR (GSE114552), ATAC-Seq (GSE104720) and DNase1-Seq (GSE61793). FAIRE-Seq peaks were downloaded from the UCSC genome browser and not re-called.

Chromosome interaction data: To assess chromatin looping, we utilized a previously published H3K27Ac HiChIP dataset from a normal hTERT-immortalized endometrial cell line (E6E7hTERT) and three endometrial cancer cell lines (ARK1, Ishikawa and JHUEM-14) (O'Mara, Spurdle and Glubb, 2019). This study identified 66,092 to 449,157 *cis* HiChIP loops (5 kb–2 Mb in length) per cell line, with a majority involving interactions of over 20 kb in distance; 35%–40% had contact with a promoter and these promoter-associated loops had a median span >200 kb. Contact data were from the original publication and not re-called for this study. Regions that interacted with the *IL15* promoter were searched for *HAND2* binding motifs using the JASPAR 2020 database of curated, non-redundant transcription factor binding profiles stored as position frequency matrices (PFMs) (Fornes *et al.*, 2020); *IL15* promoter interacting regions were searched for the *HAND2* matrix profile (matrix ID MA1638.1) (Dai and Cserjesi, 2002) using the 'Scan" option on the JASPAR website.

Cell culture and *HAND2* knockdown

Human hTERT-immortalized endometrial stromal fibroblasts (T-HESC; CRL-4003, ATCC) were grown in maintenance medium, consisting of Phenol Red-free DMEM (31053-028, Thermo Fisher Scientific), supplemented with 10% charcoal-stripped fetal bovine serum (CS-FBS; 12676029, Thermo Fisher Scientific), 1% L-glutamine (25030-081, Thermo Fisher Scientific), 1% sodium pyruvate (11360070, Thermo Fisher Scientific), and 1x insulin-transferrin-selenium (ITS; 41400045, Thermo Fisher Scientific). 2x10⁵ cells were plated per well of a 6-well plate and 18 hours later cells in 1750µl of Opti-MEM (31985070, Thermo Fisher Scientific) were transfected with 50nM of siRNA targeting *HAND2* (s18133; Silencer Select Pre-Designed siRNA; cat#4392420, Thermo Fisher Scientific) and 9µl of Lipofectamine RNAiMAX (133778-150, Invitrogen) in 250µl Opti-MEM. BlockIT Fluorescent Oligo (44-2926, Thermo Fisher Scientific) was used as a scrambled non-targeting RNA control. Cells were incubated in the transfection mixture for 6 hours. Then, cells were washed with PBS and incubated in the maintenance medium overnight. Cells in the control wells were checked under the microscope for fluorescence the next day. 48 hours post treatment cells were washed with PBS, trypsinized (0.05% Trypsin-EDTA; 15400-054, Thermo Fisher Scientific) and total RNA was extracted using RNeasy Plus Mini Kit (74134, QIAGEN) following the manufacturer's protocol. The knockdown experiment was done in 3 biological replicates. To test for the efficiency of the knockdown, cDNA was synthesized from 100-200ng RNA using Maxima H Minus First Strand cDNA Synthesis Kit (K1652, Thermo Fisher Scientific) following the manufacturer's protocol. qRT-PCR was performed using QuantiTect SYBR Green PCR (204143, QIAGEN). *HAND2* primers: fwd CACCAGCTACATCGCCTACC, rev ATTCGTTCAGCTCCTTCTTCC. *GAPDH* housekeeping gene was used for normalization; primers fwd AATCCCATCACCATCTTCCA, rev TGGACTCCACGACGTACTCA. Samples that showed >70% KD efficiency were used for RNA-Seq.

HAND2 knockdown RNA-Seq analysis

RNA from knockdown and control samples were DNase treated with TURBO DNA-free Kit (AM1907, Thermo Fisher Scientific) and RNA quality and quantity were assessed on 2100 Bioanalyzer (Agilent Technologies, Inc.). RNA-Seq libraries were prepared using TruSeq Stranded Total RNA Library Prep Kit with Ribo-Zero Human (RS-122-2201, Illumina Inc.) following manufacturer's protocol. Library quality and quantity were checked on 2100 Bioanalyzer and the pool of libraries was sequenced on Illumina HiSEQ4000 (single-end 50bp)

using manufacturer's reagents and protocols. Quality control, Ribo-Zero library preparation and Illumina sequencing were performed at the Genomics Facility at The University of Chicago.

All sequencing data were uploaded and analyzed on the Galaxy platform (<https://usegalaxy.org/>; Version 20.01). Individual reads for particular samples were concatenated using the "Concatenate datasets" tool (version 1.0.0). We used HISAT2 (version 2.1.0+galaxy5) (Kim, Langmead and Salzberg, 2015) to align reads to the Human hg38 reference genome using "Single-end" option, and reporting alignments tailored for transcript assemblers including StringTie. Transcripts were assembled and quantified using StringTie (v1.3.6) (Pertea *et al.*, 2015, 2016), with reference file to guide assembly and the "reference transcripts only" option, and output count files for differential expression with DESeq2/edgeR/limma-voom. Differentially expressed genes were identified using DESeq2 (version 2.11.40.6+galaxy1) (Anders and Huber, 2010; Love, Huber and Anders, 2014). The reference file for StringTie guided assembly was wgEncodeGencodeBasicV33. The volcano plot was generated using Blighe K., Rana S., and Lewis M. (2018): EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling (available at <https://github.com/kevinblighe/EnhancedVolcano>).

Trans-well migration assay

Choice of cells and culture: Human hTERT-immortalized ESFs (T-HESC) were selected as a model ESF cell line because they are proliferative, maintain hormone responsiveness and gene expression patterns characteristic of primary ESFs, and have been relatively well characterized (Krikun *et al.*, 2004). ESFs were cultured in the maintenance medium as described above in T75 flasks until ~80% confluent. Cryopreserved primary adult human CD56+ NK cells purified by immunomagnetic bead separation were obtained from ATCC (PCS-800-019) and cultured in RPMI-1640 containing 10% FBS and IL2 in T75 flasks for two days prior to trans-well migration assays. We used the immortalized first trimester extravillous trophoblast cell line HTR-8/SVneo (Graham *et al.*, 1993), because it maintains characteristics of extravillous trophoblasts and has previously been shown to be a good model of trophoblast migration and invasion (Iacob *et al.*, 2008; Paiva *et al.*, 2009; Hannan *et al.*, 2010). HTR-8/SVneo cells were obtained from ATCC (CRL-3271) and cultured in RPMI-1640 containing 5% FBS in T75 flasks for two days prior to trans-well migration assays.

3x10⁴ ESF cells were plated per well of a 24-well plate and 18 hours later cells were transfected in Opti-MEM with 10nM (per well) of siRNA targeting *HAND2* (s18133; Silencer Select Pre-Designed siRNA, cat#4392420; Thermo Fisher Scientific) or *IL15* (s7377; Silencer Select Pre-Designed siRNA, cat#4392420; Thermo Fisher Scientific) and 1.5µl (per well) of Lipofectamine RNAiMAX (133778-150; Invitrogen). As a negative control we used Silencer Select negative control No. 1 (4390843; Thermo Fisher Scientific). ESF cells were incubated in the transfection mixture for 6 hours. Then, ESF cells were washed with warm PBS and incubated in the maintenance medium overnight. Efficiency of the knockdown was confirmed 48 hours post-treatment by qPCR, media from each well was transferred to new 24-well plates and stored at 4°C. Total RNA from cells was extracted using RNeasy Plus Mini Kit (QIAGEN) following the manufacturer's protocol. cDNA was synthesized from 10ng RNA using Maxima H Minus First Strand cDNA Synthesis Kit following the manufacturer's protocol. qRT-PCR was performed using TaqMan Fast Universal PCR Master Mix 2X (4352042, Thermo Fisher Scientific), with primers for *HAND2* (Hs00232769_m1), *IL15* (Hs01003716_m1) and *Malat* (Hs00273907_s1) as a control housekeeping gene. Conditioned media from samples with >70% knockdown efficiency was used for trans-well migration assays.

Corning HTS Trans-well permeable supports were used for the trans-well migration assay (Corning, cat # CLS3398). Prior to the assays, conditioned media was warmed to room temperature and centrifuged for 3min at 1000 RPM to pellet any cells. For experiments using recombinant human IL15 (rhIL15), we added 10ng/ml (AbCam, ab259403) of rhIL15 to fresh, non-conditioned ESF media; 10ng/ml has previously been shown to induce migration of JEG-3 choriocarcinoma cells (Zygmunt *et al.*, 1998). For neutralizing antibody experiments, either 1µg/ml of anti-IL15 IgG (AbCam cat # MA5-23729) or control IgG (AbCam cat # 31903) were added to non-conditioned ESF media; 1µg/ml has previously been shown to neutralize ESF-derived proteins and inhibit AC-1M88 trophoblast cell migration in trans-well assays (Gellersen *et al.*, 2010, 2013). Plates were incubated with shaking at 37°C for 30min prior to initiation of migration assays. During antibody incubation, NK and HTR-8/SVneo cells were collected and resuspended in fresh ESF growth media. For the trans-well migration assay, 5x10⁶ of either NK or HTR-8/SVneo cells were added to each well of the upper chamber and either treatment or control media were added to the lower chambers. Plates were incubated at 37°C with 5% CO₂.

After 8 hours incubation, we removed the upper plate (containing remaining NK and




HTR-8/SVneo cells) and discarded non-migrated cells. 50 μ l from each well in the lower chamber was transferred into a single well of a 96-well opaque plate. We used the CellTiter-Glo luminescent cell viability assay (G7570 Promega) to measure luminescence, which is proportional to the number of live cells per well. Data are reported as effect sizes (mean differences) between treatment and control. Confidence intervals are bias-corrected and accelerated. The P-values reported are the likelihoods of the observed effect sizes, if the null hypothesis of zero difference is true and calculated from a two-sided permutation t-test (5000 reshuffles of the control and test labels). Cumming estimation plots and estimation statistics were calculated using DABEST R package (Ho *et al.*, 2019).

Acknowledgements

The authors would like to thank the following researchers for providing pregnant endometrial samples: G.P. Wagner (Yale University) – *Monodelphis domestica*; R. R. Behringer (The University of Texas MD Anderson Cancer Center) – *Carollia perspicillata*; B.C. Paria (Vanderbilt University School of Medicine) – *Mesocricetus auratus*; A.T. Fazleabas (Michigan State University) – *Papio anubis*; D. K. Merriman (University of Wisconsin Oshkosh) – *Ictidomys tridecemlineatus*. We are also grateful to A.M. Bamberger (University Hospital Eppendorf) for providing the trans-well migration assay protocol, D. Glubb (QIMR Berghofer Medical Research Institute) for assistance in interpreting the HiChIP assay data, R. Beaumont (University of Exeter Medical School) and R.M. Freathy (University of Exeter) for assistance with interpreting maternal-fetal birth weight GWAS data. This study was supported by a grant from the March of Dimes (March of Dimes Prematurity Research Center to principal investigator VJL) and a Burroughs Wellcome Fund Preterm Birth Initiative grant (1013760, to principal investigator VJL). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. VJL thanks the Department of Human Genetics at The University of Chicago for support during the planning and preliminary data generation phase of this work. MM thanks Michael Sulak for the help with editing the manuscript.

Table 1. Exemplar genes differentially expressed in ESFs upon siRNA-mediated *HAND2* knockdown. Mean, base mean expression level. FC, \log_2 fold change. SE, standard error in \log_2 fold change. WS, Wald statistic. P-value, Wald test P-value. Adj-P, Benjamini-Hochberg (BH) adjusted P-value. Function, function of gene inferred from WikiPathway 2019 human annotation. Association with preterm birth (PTB, HP:0001622) and premature rupture of membranes (PROM, HP:0001788) inferred from human phenotype ontology annotation.

Gene	Mean	FC	SE	WS	P-value	Adj-P	Function
<i>ARNT2</i>	44.47	1.20	0.28	4.30	1.73E-05	1.89E-03	AHR signaling / Circadian rhythm
<i>ARNTL</i>	42.52	-2.95	0.80	-3.67	2.46E-04	1.79E-02	AHR signaling / Circadian rhythm
<i>IL15</i>	63.31	7.98	1.49	5.37	7.91E-08	1.49E-05	Cell migration
<i>BMP4</i>	58.10	-8.34	1.25	-6.66	2.75E-11	8.28E-09	Deciduallization
<i>GSK3B</i>	69.23	-5.53	1.48	-3.73	1.94E-04	1.49E-02	Deciduallization
<i>HAND2</i>	136.06	-1.63	0.18	-8.85	9.10E-19	7.64E-16	Knocked down gene
<i>CHUK</i>	765.77	0.44	0.14	3.14	1.67E-03	7.94E-02	NF κ B pathway
<i>IkBKE</i>	9.91	-5.41	1.65	-3.27	1.06E-03	5.72E-02	NF κ B pathway
<i>MYD88</i>	95.02	-0.93	0.26	-3.55	3.88E-04	2.59E-02	NF κ B pathway
<i>NFκBIE</i>	106.63	-0.78	0.25	-3.08	2.10E-03	9.53E-02	NF κ B pathway
<i>RTKN2</i>	50.64	8.02	1.33	6.02	1.75E-09	4.05E-07	NF κ B pathway
<i>CRKL</i>	1278.85	-0.31	0.09	-3.31	9.21E-04	5.10E-02	PTB
<i>EOGT</i>	233.65	-2.77	0.80	-3.45	5.63E-04	3.38E-02	PTB
<i>LMNA</i>	773.24	6.97	1.57	4.44	8.92E-06	1.06E-03	PTB; PROM
<i>PEX11B</i>	181.31	10.05	1.12	9.00	2.28E-19	2.06E-16	PTB
<i>SERPINH1</i>	1654.90	-6.01	1.31	-4.59	4.40E-06	5.61E-04	PROM
<i>SLC17A5</i>	826.28	0.76	0.13	5.81	6.34E-09	1.40E-06	PTB
<i>ZMPSTE24</i>	2140.67	-0.19	0.05	-4.10	4.10E-05	3.92E-03	PTB; PROM



12.5 d.p.c. pregnant opossum endometrium. Data shown as square root (Sqrt) transformed Transcripts Per Million (TPM). The TPM=2 expression cutoff is shown as a red line; genes with $TPM \geq 2$ are shown as black bars.

Figure 1. Recruitment of HAND2-mediated anti-estrogenic signaling in the Eutherian endometrium.

A. Evolution of *HAND2* expression at the maternal-fetal interface. Amniotes phylogeny with horizontal branch lengths drawn proportional to the number of gene expression changes inferred by parsimony (most parsimonious reconstruction). Circles indicate *HAND2* expression in extant species and ancestral reconstructions. Black, expressed (state=1). White, not expressed (state=0). Inset legend shows the number of most gene expression changes from the root node to human (+ = gene expression gained; - = gene expression lost).

B. Cartoon model of estrogen signaling and *HAND2*-mediated anti-estrogenic signaling in the endometrium. The estrogen-mediated signaling network is suppressed by progesterone through the activation of *HAND2* and antagonists of canonical WNT/β-catenin mediated signaling pathways such as *DKK1*. In the proliferative phase of the reproductive cycle, estrogen acts through *ESR1* in stromal cells to increase the production of fibroblast growth factors (FGFs), which serve as paracrine signals leading to sustained proliferation of epithelial cells. Active estrogen signaling maintains epithelial expression of Mucin 1 (MUC1), a cell surface glycoprotein that acts as a barrier to implantation. During the receptive phase of the cycle, however, progesterone induces *HAND2* and *DKK1* expression in the endometrial stroma, inhibiting production of fibroblast growth factors (FGFs), suppressing epithelial proliferation and antagonizing estrogen-mediated expression of MUC1, thereby promoting uterine receptivity to implantation.

C. Upper, Timeline of pregnancy in *Monodelphis domestica* highlighting the histotrophic and placental phases. Bottom left, Immunohistochemistry showing phosphorylated ESR1 (pESR1), phosphorylated MAPK1/2 (pERK1/2), and MUC1 expression in paraffin embedded 12.5 d.p.c. pregnant opossum endometrium compared to control (IgG). PI, placenta. LE, luminal epithelium. ES, endometrial stroma. BV, blood vessels. Bottom right, expression of genes related to *HAND2*-mediated anti-estrogenic signaling in RNA-Seq from




Figure 2. Expression of *HAND2* at the maternal-fetal interface

- UMAP clustering of scRNA-Seq data from human first trimester maternal-fetal interface. Left, clusters colored according to inferred cell type. The ESF1, ESF2, and DSC clusters are highlighted. Right, cells within clusters are colored according to *HAND2* expression level.
- HAND2* protein expression in human pregnant decidua, with strong staining and localization in the nuclei of endometrial stromal cells (image color auto adjusted). Image credit: Human Protein Atlas.
- Regulatory landscape of the *HAND2* locus. Chromatin loops inferred from H3K27ac HiChIP, regions of open chromatin inferred from FAIRE-, DNasel, and ATAC-Seq, and the locations of histone modifications and transcription factor ChIP-Seq peaks are shown. The location of SNPs associated with gestation length is also shown (highlighted in gray). Note that the *HAND2* promoter forms a long-range loop to a region marked by H3K27ac and bound by PR, NR2F2 (COUP-TFII), GATA2, FOSL2, and FOXO1.
- HAND2* expression is up-regulated by in vitro decidualization of ESFs into DSC by cAMP/progesterone treatment, and down-regulated by siRNA-mediated knockdown of PR and GATA2, but not NR2F2 or FOXO1. n=3 per transcription factor knockdown.
- Relative expression of *HAND2* in the proliferative (n=6), early (n=4), middle (n=9), and late (n=9) secretory phases of the menstrual cycle. Note that outliers are excluded from the figure but not the regression; 95% CI is shown in gray.
- Relative expression of *HAND2* in the basal plate from mid-gestation to term (14-40 weeks); 95% CI is shown in gray. Inset, percent of up- (Up) and down-regulated (Down) genes between weeks 14-19 and 37-40 of pregnancy (FDR≤0.10).
- HAND2* expression is significantly down-regulated in the endometria of women with implantation failure (IF, n=5) and recurrent spontaneous abortion (RSA, n=5) compared to fertile controls (n=5), but is not differentially expressed in ESFs or DSCs from women with preeclampsia (PE, n=5) compared to healthy controls (n=5).




Figure 3. HAND2 regulates a distinct set of target genes, including IL15

A. Volcano plot of gene expression upon *HAND2* knockdown. Only genes that are significantly differentially expressed (DE) with $FDR \leq 0.10$ are shown. Genes with ≥ 1.5 -fold changes in expression are shown in red, others in blue.

B. UpSet plot showing intersection of genes DE by *HAND2*, *NR2F2*, *FOXO1*, *GATA2*, and *PR* knockdown (KD), and cAMP/MPA treatment. Inset plot shows number of DE genes for each treatment. Note large intersections for genes DE by cAMP/MPA/PR KD, GATA2 KD/PR KD, and cAMP/GATA2 KD/PR KD, but few intersections with *HAND2* KD and other TF KDs.

C. Pathways (purple) and human phenotype ontologies (pink) in which genes dysregulated upon *HAND2* KD are enriched.




Figure 4. Expression of *IL15* at the maternal-fetal interface

- UMAP clustering of scRNA-Seq data from human the first trimester maternal-fetal interface. Left, clusters colored according to inferred cell type. The ESF1, ESF2, and DSC clusters are highlighted. Inset, per cell expression of *HAND2* and *IL15* in ESF1s, ESF2s, and DSCs. Right, cells within clusters are colored according to *IL15* expression level.
- IL15 protein expression in human pregnant decidua, with strong cytoplasmic staining in endometrial stromal cells (image color auto adjusted). Image credit: Human Protein Atlas.
- Regulatory landscape of the *IL15* locus. Chromatin loops inferred from H3K27ac HiChIP, regions of open chromatin inferred from FAIRE-, DNasel, and ATAC-Seq, and the locations of histone modifications and transcription factor ChIP-Seq peaks are shown. The location of a SNP associated with gestation length / birth weight is also shown (highlighted in gray). Note that the *IL15* promoter forms many long-range loops to regions marked by H3K27ac and bound by PR, NR2F2, GATA2, FOSL2, FOXO1, and SRC2.
- IL15* expression is up-regulated by in vitro decidualization of ESFs into DSC by cAMP/progesterone treatment, and down-regulated by siRNA-mediated knockdown of PR and GATA2 but not NR2F2 or FOXO1. $n=3$ per transcription factor knockdown.
- Relative expression of *IL15* in the proliferative ($n=6$), early ($n=4$), middle ($n=9$), and late ($n=9$) secretory phases of the menstrual cycle. Note that outliers are excluded from the figure. Note that outliers are excluded from the figure but not the regression; 95% CI is shown in gray.
- Relative expression of *IL15* in the basal plate from mid-gestation to term (14-40 weeks); 95% CI is shown in gray. Inset, percent of up- (Up) and down-regulated (Down) genes between weeks 14-19 and 37-40 of pregnancy (FDR ≤ 0.10).
- IL15* expression is significantly up-regulated in DSCs from women with preeclampsia (PE, $n=5$) compared to healthy controls ($n=5$), while it is only marginally up-regulated in ESFs from the same patient group. It is also marginally down-regulated in the endometria of women with implantation failure (IF, $n=5$) and it is not differentially expressed in the endometria of women with recurrent spontaneous abortion (RSA, $n=5$) compared to fertile controls ($n=5$).




Figure 5. ESF-derived IL15 promotes NK and trophoblast migration in trans-well assays.

A. Cartoon of trans-well migration assay comparisons.

Cells that migrated to the lower chamber were quantified using the CellTiter-Glo luminescent cell viability assay after 8 hours.

B. Primary natural killer (NK) cells. Raw luminescence data (RLU) is shown in the upper panel, mean difference (effect size) in experiment minus control luminescence values are shown as a dots with the 95% confidence interval indicated by vertical bars in the lower panel; distribution estimated from 5000 bootstrap replicates. The mean difference between Media and media supplemented with recombinant human IL15 (Media+rhIL15) is 0.432 [95.0% CI: 0.376 – 0.492]; P=0.00. The mean difference between ESFs transiently transfected with non-targeting siRNA (NT siRNA) and *HAND2*-specific siRNAs (siHAND2) is 0.847 [95.0% CI: 0.774 – 0.936]; P=0.00. The mean difference between ESFs transiently transfected with NT siRNA and IL15-specific siRNAs (siL15) is -0.223 [95.0% CI: -0.256 – -0.192]; P=0.00. The mean difference between ESF media neutralized with either a non-specific antibody (IgG) or IL15-specific antibody (abIL15) is -0.106 [95.0% CI -0.147 – -0.067]; P=0.00. n=12.

C. Extravillous trophoblast cell line HTR-8/SVneo. Raw luminescence data (RLU) from cells in the lower chamber is shown in the upper panel, mean difference (effect size) in experiment minus control luminescence values are shown as a dots with the 95% confidence interval indicated by vertical bars in the lower panel; distribution estimated from 5000 bootstrap replicates. The mean difference between Media and Media+rhIL15 is 0.482 [95.0% CI: 0.193 – 0.701]; P=0.002.

The mean difference between ESFs transiently transfected with NT siRNA and siHAND2 is 0.463 [95.0% CI: 0.291 – 0.559]; P=0.00. The mean difference between ESFs transiently transfected with NT siRNA and siL15 is -0.598 [95.0% CI: -0.698 – 0.490]; P=0.00. The mean difference between ESF media neutralized with IgG or abIL15 is -0.267 [95.0% CI: -0.442 – -0.151]; P=0.0004. n=12.

D. Model of HAND2 functions in the endometrium. During the proliferative phase HAND2 inhibits *IL15*, and thus the migration of uNK and trophoblast cells. In the receptive phase, HAND2 activates *IL15*, which promotes migration of uNK and trophoblast cells. In the receptive phase and early pregnancy (EP), HAND2 suppresses estrogen signaling by down-regulating *FGFs* and directly binding and inhibiting the ligand-dependent transcriptional activation function of ESR1. During late pregnancy/term (LP/Term), reduced HAND2 expression mitigates its anti-estrogenic functions. Parturition signal unknown (??).

References

Abbott, B. D. *et al.* (1999) 'Adverse reproductive outcomes in the transgenic Ah receptor-deficient mouse', *Toxicology and Applied Pharmacology*. Academic Press Inc., 155(1), pp. 62–70. doi: 10.1006/taap.1998.8601.

Abegglen, L. M. *et al.* (2015) 'Potential mechanisms for cancer resistance in elephants and comparative cellular response to DNA Damage in Humans', *JAMA - Journal of the American Medical Association*. American Medical Association, 314(17), pp. 1850–1860. doi: 10.1001/jama.2015.13134.

Afgan, E. *et al.* (2018) 'The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update', *Nucleic Acids Research*. Narnia, 46(W1), pp. W537–W544. doi: 10.1093/nar/gky379.

Allavena, P. *et al.* (1997) 'IL-15 is chemotactic for natural killer cells and stimulates their adhesion to vascular endothelium', *Journal of Leukocyte Biology*. Federation of American Societies for Experimental Biology, 61(6), pp. 729–735. doi: 10.1002/jlb.61.6.729.

Allport, V. C. (2001) 'Human labour is associated with nuclear factor-kappaB activity which mediates cyclo-oxygenase-2 expression and is involved with the "functional progesterone withdrawal"', *Molecular Human Reproduction*. Oxford Academic, 7(6), pp. 581–586. doi: 10.1093/molehr/7.6.581.

Anders, S. and Huber, W. (2010) 'Differential expression analysis for sequence count data', *Genome Biology*. Genome Biol, 11(10). doi: 10.1186/gb-2010-11-10-r106.

Armstrong, D. L. *et al.* (2017) 'The core transcriptome of mammalian placentas and the divergence of expression with placental shape', *Placenta*. W.B. Saunders Ltd, 57, pp. 71–78. doi: 10.1016/j.placenta.2017.04.015.

Arthur, A. *et al.* (2018) 'Molecular Evolution of Genes Associated with Preeclampsia: Genetic Conflict, Antagonistic Coevolution and Signals of Selection', *Ashdin Publishing Journal of Evolutionary Medicine*, 6. doi: 10.4303/jem/236038.

Ashkar, A. A. *et al.* (2003) 'Assessment of requirements for IL-15 and IFN regulatory factors in uterine NK cell differentiation and function during pregnancy', *The Journal of Immunology*. The American Association of Immunologists, 171(6), pp. 2937–2944. doi: 10.4049/jimmunol.171.6.2937.

Bany, B. M., Scott, C. A. and Eckstrum, K. S. (2012) 'Analysis of uterine gene expression in interleukin-15 knockout mice reveals uterine natural killer cells do not play a major role in decidualization and associated angiogenesis', *Reproduction*. Reproduction, 143(3), pp. 359–375. doi: 10.1530/REP-11-0325.

Barber, E. M. and Pollard, J. W. (2003) 'The uterine NK cell population requires IL-15 but these cells are not required for pregnancy nor the resolution of a Listeria monocytogenes infection', *The Journal of Immunology*. The American Association of Immunologists, 171(1), pp. 37–46. doi: 10.4049/jimmunol.171.1.37.

Bartmann, C. *et al.* (2014) 'Quantification of the predominant immune cell populations in decidua throughout human pregnancy', *American Journal of Reproductive Immunology*. Am J Reprod Immunol, 71(2), pp. 109–119. doi: 10.1111/aji.12185.

Boretto, M. *et al.* (2017) 'Development of organoids from mouse and human endometrium

showing endometrial epithelium physiology and long-term expandability', *Development (Cambridge)*. Company of Biologists Ltd, 144(10), pp. 1775–1786. doi: 10.1242/dev.148478.

Brayer, K., Lynch, V. and Wagner, G. (2009) 'Evolution of physical interactions among the transcription factors HoxA-11 and FOXO1a during the evolution of pregnancy in mammals', *Integrative and Comparative Biology*. Oxford University Press (OUP), 49(Supplement 1), p. e21. doi: 10.1093/icb/icp002.

Brighton, P. J. *et al.* (2017) 'Clearance of senescent decidual cells by uterine natural killer cells in cycling human endometrium', *eLife*. eLife Sciences Publications Ltd, 6. doi: 10.7554/elife.31274.

Brockway, H. M. *et al.* (2019) 'Unique transcriptomic landscapes identified in idiopathic spontaneous and infection related preterm births compared to normal term births', *PLoS ONE*. Public Library of Science, 14(11). doi: 10.1371/journal.pone.0225062.

Bulmer, J. N. *et al.* (1991) 'Granulated lymphocytes in human endometrium: Histochemical and immunohistochemical studies', *Human Reproduction*. Oxford University Press, 6(6), pp. 791–798. doi: 10.1093/oxfordjournals.humrep.a137430.

Burke, S. D. *et al.* (2010) 'Uterine NK cells, spiral artery modification and the regulation of blood pressure during mouse pregnancy', *American Journal of Reproductive Immunology*. Am J Reprod Immunol, 63(6), pp. 472–481. doi: 10.1111/j.1600-0897.2010.00818.x.

Buzzio, O. L. *et al.* (2006) 'FOXO1A differentially regulates genes of decidualization', *Endocrinology*. Endocrinology, 147(8), pp. 3870–3876. doi: 10.1210/en.2006-0167.

Chekir, C. *et al.* (2006) 'Accumulation of advanced glycation end products in women with preeclampsia: Possible involvement of placental oxidative and nitrative stress', *Placenta*. Placenta, 27(2–3), pp. 225–233. doi: 10.1016/j.placenta.2005.02.016.

Chen, Y. *et al.* (2011) 'Effect of human endometrial stromal cell-derived conditioned medium on uterine natural killer (uNK) cells' proliferation and cytotoxicity', *American Journal of Reproductive Immunology*. Am J Reprod Immunol, 65(6), pp. 589–596. doi: 10.1111/j.1600-0897.2010.00955.x.

Dai, Y. S. and Cserjesi, P. (2002) 'The basic helix-loop-helix factor, HAND2, functions as a transcriptional activator by binding to E-boxes as a heterodimer', *Journal of Biological Chemistry*. J Biol Chem, 277(15), pp. 12604–12612. doi: 10.1074/jbc.M200283200.

Davis, S. and Meltzer, P. S. (2007) 'GEOquery: A bridge between the Gene Expression Omnibus (GEO) and BioConductor', *Bioinformatics*, 23(14), pp. 1846–1847. doi: 10.1093/bioinformatics/btm254.

Dunn, C. L., Critchley, H. O. D. and Kelly, R. W. (2002) 'IL-15 regulation in human endometrial stromal cells', *The Journal of Clinical Endocrinology & Metabolism*. The Endocrine Society, 87(4), pp. 1898–1901. doi: 10.1210/jcem.87.4.8539.

Eidem, H. R. *et al.* (2016) 'Comparing human and macaque placental transcriptomes to disentangle preterm birth pathology from gestational age effects', *Placenta*. W.B. Saunders Ltd, 41, pp. 74–82. doi: 10.1016/j.placenta.2016.03.006.

Elliot, M. G. (2017) 'Evolutionary origins of preeclampsia', *Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health*. Elsevier BV, 7, p. 56. doi: 10.1016/j.preghy.2016.10.006.

Erlebacher, A. (2013) 'Mechanisms of T cell tolerance towards the allogeneic fetus', *Nature Reviews Immunology*. Nature Publishing Group, pp. 23–33. doi: 10.1038/nri3361.

Felker, A. M. and Croy, B. A. (2016) 'Uterine natural killer cell partnerships in early mouse decidua basalis', *Journal of Leukocyte Biology*. Wiley-Blackwell, 100(4), pp. 645–655. doi: 10.1189/jlb.1HI0515-226R.

Feng, J. *et al.* (2012) 'Identifying ChIP-seq enrichment using MACS', *Nature Protocols*. Nat Protoc, 7(9), pp. 1728–1740. doi: 10.1038/nprot.2012.101.

Fornes, O. *et al.* (2020) 'JASPAR 2020: Update of the open-access database of transcription factor binding profiles', *Nucleic Acids Research*, 48(D1), pp. D87–D92. doi: 10.1093/nar/gkz1001.

Franco, H. L. *et al.* (2011) 'WNT4 is a key regulator of normal postnatal uterine development and progesterone signaling during embryo implantation and decidualization in the mouse', *The FASEB Journal*. Wiley, 25(4), pp. 1176–1187. doi: 10.1096/fj.10-175349.

Fraser, R. *et al.* (2015) 'Decidual natural killer cells regulate vessel stability: Implications for impaired spiral artery remodelling', *Journal of Reproductive Immunology*. Elsevier Ireland Ltd, 110, pp. 54–60. doi: 10.1016/j.jri.2015.04.003.

Fukuda, T. *et al.* (2015) 'HAND2-mediated proteolysis negatively regulates the function of estrogen receptor α', *Molecular Medicine Reports*. Spandidos Publications, 12(4), pp. 5538–5544. doi: 10.3892/mmr.2015.4070.

Funeshima, N. *et al.* (2005) 'Inhibition of allogeneic T-cell responses by dendritic cells expressing transduced indoleamine 2,3-dioxygenase', *Journal of Gene Medicine*. J Gene Med, 7(5), pp. 565–575. doi: 10.1002/jgm.698.

Gellersen, B. *et al.* (2010) 'Invasiveness of human endometrial stromal cells is promoted by decidualization and by trophoblast-derived signals', *Human Reproduction*, 25(4), pp. 862–873. doi: 10.1093/humrep/dep468.

Gellersen, B. *et al.* (2013) 'Human endometrial stromal cell-trophoblast interactions: Mutual stimulation of chemotactic migration and promigratory roles of cell Surface molecules CD82 and CEACAM11', *Biology of Reproduction*. Oxford University Press (OUP), 88(3). doi: 10.1095/biolreprod.112.106724.

Gellersen, B. and Brosens, J. (2003) 'Cyclic AMP and progesterone receptor cross-talk in human endometrium: A decidualizing affair', *Journal of Endocrinology*, 178(3), pp. 357–372. doi: 10.1677/joe.0.1780357.

Gentleman, R. C. *et al.* (2004) 'Bioconductor: open software development for computational biology and bioinformatics.', *Genome biology*. Genome Biol, 5(10). doi: 10.1186/gb-2004-5-10-r80.

Girling, J. E. (2002) 'The reptilian oviduct: A review of structure and function and directions for future research', *Journal of Experimental Zoology*. J Exp Zool, pp. 141–170. doi: 10.1002/jez.10105.

Godbole, G. *et al.* (2011) 'Decidualized endometrial stromal cell derived factors promote trophoblast invasion', *Fertility and Sterility*. Fertil Steril, 95(4), pp. 1278–1283. doi: 10.1016/j.fertnstert.2010.09.045.

Godbole, G. and Modi, D. (2010) 'Regulation of decidualization, interleukin-11 and interleukin-

15 by homeobox A10 in endometrial stromal cells', *Journal of Reproductive Immunology*. J Reprod Immunol, 85(2), pp. 130–139. doi: 10.1016/j.jri.2010.03.003.

Gomaa, M. F. *et al.* (2017) 'Uterine natural killer cells dysregulation in idiopathic human preterm birth: a pilot study', *Journal of Maternal-Fetal and Neonatal Medicine*. Taylor and Francis Ltd, 30(15), pp. 1782–1786. doi: 10.1080/14767058.2016.1224840.

Gomez-Lopez, N., Guilbert, L. J. and Olson, D. M. (2010) 'Invasion of the leukocytes into the fetal-maternal interface during pregnancy', *Journal of Leukocyte Biology*. Wiley, 88(4), pp. 625–633. doi: 10.1189/jlb.1209796.

González-Morán, M. G. (2015) 'Immunohistochemical localization of progesterone receptor isoforms and estrogen receptor alpha in the chicken oviduct magnum during development', *Acta Histochemica*. Elsevier GmbH, 117(8), pp. 681–687. doi: 10.1016/j.acthis.2015.10.003.

Graham, C. H. *et al.* (1993) 'Establishment and characterization of first trimester human trophoblast cells with extended lifespan', *Experimental Cell Research. Exp Cell Res*, 206(2), pp. 204–211. doi: 10.1006/excr.1993.1139.

Graham, C. H. and Lala, P. K. (1991) 'Mechanism of control of trophoblast invasion in situ', *Journal of Cellular Physiology*. J Cell Physiol, 148(2), pp. 228–234. doi: 10.1002/jcp.1041480207.

Greenhill, C. (2014) 'Reproductive endocrinology: Circadian clock involved in embryo implantation', *Nature Reviews Endocrinology*, p. 701. doi: 10.1038/nrendo.2014.174.

Guedes-Martins, L. *et al.* (2013) 'AGEs, contributors to placental bed vascular changes leading to preeclampsia', *Free Radical Research. Free Radic Res*, pp. 70–80. doi: 10.3109/10715762.2013.815347.

Guleria, I. and Pollard, J. W. (2000) 'The trophoblast is a component of the innate immune system during pregnancy', *Nature Medicine*. Nat Med, 6(5), pp. 589–593. doi: 10.1038/75074.

Hamilton, S. *et al.* (2012) 'Macrophages infiltrate the human and rat decidua during term and preterm labor: Evidence that decidual inflammation precedes labor', *Biology of Reproduction*. Oxford University Press (OUP), 86(2). doi: 10.1095/biolreprod.111.095505.

Hanna, J. *et al.* (2006) 'Decidual NK cells regulate key developmental processes at the human fetal-maternal interface', *Nature Medicine*. Nat Med, 12(9), pp. 1065–1074. doi: 10.1038/nm1452.

Hannan, N. J. *et al.* (2010) 'Models for study of human embryo implantation: Choice of cell lines?', *Biology of Reproduction*. Oxford University Press (OUP), 82(2), pp. 235–245. doi: 10.1095/biolreprod.109.077800.

Hao, K. *et al.* (2013) 'Possible role of the "IDO-ahr axis" in maternal-foetal tolerance', *Cell Biology International. Cell Biol Int*, 37(2), pp. 105–108. doi: 10.1002/cbin.10023.

Hayashi, K. *et al.* (2009) 'Wnt genes in the mouse uterus: Potential regulation of implantation', *Biology of Reproduction*. Oxford University Press (OUP), 80(5), pp. 989–1000. doi: 10.1095/biolreprod.108.075416.

Hazan, A. D. *et al.* (2010) 'Vascular-leukocyte interactions: Mechanisms of human decidual spiral artery remodeling in vitro', *American Journal of Pathology*. Elsevier Inc., 177(2), pp. 1017–1030. doi: 10.2353/ajpath.2010.091105.

Hiby, S. E. *et al.* (2010) 'Maternal activating KIRs protect against human reproductive failure

mediated by fetal HLA-C2', *Journal of Clinical Investigation*. American Society for Clinical Investigation, 120(11), pp. 4102–4110. doi: 10.1172/JCI43998.

Hill, J. P. (1936) 'V. The Development of the Monotremata.—Part I. The Histology of the Oviduct during Gestation. By Catherine J. Hill. B.Sc., Ph.D. Part II. The Structure of the Egg-shell.', *The Transactions of the Zoological Society of London*, 21(6), pp. 413–476. doi: 10.1111/j.1096-3642.1936.tb00458.x.

Ho, J. *et al.* (2019) 'Moving beyond P values: data analysis with estimation graphics', *Nature Methods*. Nature Publishing Group, 16(7), pp. 565–566. doi: 10.1038/s41592-019-0470-3.

Hou, Z. C. *et al.* (2012) 'Elephant transcriptome provides insights into the evolution of eutherian placentation', *Genome Biology and Evolution*, 4(5), pp. 713–725. doi: 10.1093/gbe/evs045.

Huyen, D. V. and Bany, B. M. (2011) 'Evidence for a conserved function of heart and neural crest derivatives expressed transcript 2 in mouse and human decidualization', *Reproduction. Reproduction*, 142(2), pp. 353–368. doi: 10.1530/REP-11-0060.

Iacob, D. *et al.* (2008) 'Decorin-mediated inhibition of proliferation and migration of the human trophoblast via different tyrosine kinase receptors', *Endocrinology*. Oxford Academic, 149(12), pp. 6187–6197. doi: 10.1210/en.2008-0780.

Jones, R. L. *et al.* (2006) 'TGF- β superfamily expression and actions in the endometrium and placenta', *Reproduction. Reproduction*, pp. 217–232. doi: 10.1530/rep.1.01076.

Kajihara, T., Brosens, J. J. and Ishihara, O. (2013) 'The role of FOXO1 in the decidual transformation of the endometrium and early pregnancy', *Medical Molecular Morphology*. Springer Tokyo, pp. 61–68. doi: 10.1007/s00795-013-0018-z.

Kato, S. *et al.* (1992) 'A far upstream estrogen response element of the ovalbumin gene contains several half-palindromic 5'-TGACC-3' motifs acting synergistically', *Cell. Cell*, 68(4), pp. 731–742. doi: 10.1016/0092-8674(92)90148-6.

Kato, S. *et al.* (1995) 'Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase', *Science. Science*, 270(5241), pp. 1491–1494. doi: 10.1126/science.270.5241.1491.

Kieckbusch, J. *et al.* (2014) 'MHC-dependent inhibition of uterine NK cells impedes fetal growth and decidual vascular remodelling', *Nature Communications*. Nature Publishing Group, 5. doi: 10.1038/ncomms4359.

Kim, D., Langmead, B. and Salzberg, S. L. (2015) 'HISAT: a fast spliced aligner with low memory requirements', *Nature Methods*. Nature Publishing Group, 12(4), pp. 357–360. doi: 10.1038/nmeth.3317.

King, A. *et al.* (1991) 'CD3- leukocytes present in the human uterus during early placentation: Phenotypic and morphologic characterization of the CD56 ++ population', *Developmental Immunology. Dev Immunol*, 1(3), pp. 169–190. doi: 10.1155/1991/83493.

Kitaya, K. *et al.* (2000) 'IL-15 expression at human endometrium and decidua', *Biology of Reproduction*. Oxford University Press (OUP), 63(3), pp. 683–687. doi: 10.1095/biolreprod63.3.683.

Kitaya, K., Yamaguchi, T. and Honjo, H. (2005) 'Central role of interleukin-15 in postovulatory recruitment of peripheral blood CD16(-) natural killer cells into human endometrium', *Journal of Clinical Endocrinology and Metabolism. J Clin Endocrinol Metab*, 90(5), pp. 2932–2940. doi:

10.1210/jc.2004-2447.

Klonisch, T. *et al.* (2001) 'Epidermal growth factor-like ligands and erbB genes in the peri-Implantation rabbit uterus and blastocyst', *Biology of Reproduction*. Oxford University Press (OUP), 64(6), pp. 1835–1844. doi: 10.1095/biolreprod64.6.1835.

Kosova, G. *et al.* (2015) 'Evolutionary forward genomics reveals novel insights into the genes and pathways dysregulated in recurrent early pregnancy loss', *Human Reproduction*, 30(3), pp. 519–529. doi: 10.1093/humrep/deu355.

Krikun, G. *et al.* (2004) 'A novel immortalized human endometrial stromal cell line with normal progestational response', *Endocrinology*, 145(5), pp. 2291–2296. doi: 10.1210/en.2003-1606.

LaBella, A. L. *et al.* (2019) *Accounting for diverse evolutionary forces reveals the mosaic nature of selection on genomic regions associated with human preterm birth*, *bioRxiv*. Cold Spring Harbor Laboratory. doi: 10.1101/816827.

Lappas, M., Permezel, M. and Rice, G. E. (2007) 'Advanced glycation endproducts mediate pro-inflammatory actions in human gestational tissues via nuclear factor- κ B and extracellular signal-regulated kinase 1/2', *Journal of Endocrinology. J Endocrinol*, 193(2), pp. 269–277. doi: 10.1677/JOE-06-0081.

Large, M. J. *et al.* (2014) 'The epidermal growth factor receptor critically regulates endometrial function during early pregnancy', *PLoS Genetics*. Edited by C. Stewart. Public Library of Science, 10(6), p. e1004451. doi: 10.1371/journal.pgen.1004451.

Lash, G. E., Robson, S. C. and Bulmer, J. N. (2010) 'Review: Functional role of uterine natural killer (uNK) cells in human early pregnancy decidua', *Placenta. Placenta*, 31(SUPPL.). doi: 10.1016/j.placenta.2009.12.022.

Laskarin, G. *et al.* (2006) 'Physiological role of IL-15 and IL-18 at the maternal-fetal interface', *Chemical Immunology and Allergy*. Basel: KARGER, pp. 10–25. doi: 10.1159/000087906.

Lee, K. Y. *et al.* (2007) 'Bmp2 is critical for the murine uterine decidual response', *Molecular and Cellular Biology*. American Society for Microbiology, 27(15), pp. 5468–5478. doi: 10.1128/MCB.00342-07.

Li, Q. *et al.* (2007) 'Bone morphogenetic protein 2 functions via a conserved signaling pathway involving Wnt4 to regulate uterine decidualization in the mouse and the human', *Journal of Biological Chemistry. J Biol Chem*, 282(43), pp. 31725–31732. doi: 10.1074/jbc.M704723200.

Li, Q. *et al.* (2011) 'The antiproliferative action of progesterone in uterine epithelium is mediated by Hand2', *Science. Science*, 331(6019), pp. 912–916. doi: 10.1126/science.1197454.

Li, Q. (2014) 'Transforming growth factor β signaling in uterine development and function', *Journal of Animal Science and Biotechnology*. BioMed Central Ltd. doi: 10.1186/2049-1891-5-52.

Liao, Y. *et al.* (2019) 'WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs', *Nucleic Acids Research*, 47(W1), pp. W199–W205. doi: 10.1093/nar/gkz401.

Lim, H., Dey, S. K. and Das, S. K. (1997) 'Differential expression of the erbB2 gene in the periimplantation mouse uterus: Potential mediator of signaling by epidermal growth factor-like growth factors', *Endocrinology. Endocrine Society*, 138(3), pp. 1328–1337. doi: 10.1210/endo.138.3.4991.

Lima, P. D. A. *et al.* (2014) 'Leukocyte driven-decidua angiogenesis in early pregnancy',

Cellular and Molecular Immunology. Chinese Soc Immunology, pp. 522–537. doi: 10.1038/cmi.2014.63.

Lindstrom, T. M. and Bennett, P. R. (2005) ‘The role of nuclear factor kappa B in human labour’, *Reproduction*. Reproduction, 130(5), pp. 569–581. doi: 10.1530/rep.1.00197.

Love, M. I., Huber, W. and Anders, S. (2014) ‘Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2’, *Genome Biology*. BioMed Central, 15(12), p. 550. doi: 10.1186/s13059-014-0550-8.

Lu, C. X. *et al.* (2016) ‘A novel HAND2 loss-of-function mutation responsible for tetralogy of Fallot’, *International Journal of Molecular Medicine*. Spandidos Publications, 37(2), pp. 445–451. doi: 10.3892/ijmm.2015.2436.

Lucas, E. S. *et al.* (2020) ‘Recurrent pregnancy loss is associated with a pro-senescent decidual response during the peri-implantation window’, *Communications Biology*. Nature Research, 3(1), p. 37. doi: 10.1038/s42003-020-0763-1.

Lynch, V. J. *et al.* (2009) ‘HoxA-11 and FOXO1A cooperate to regulate decidual prolactin expression: Towards inferring the core transcriptional regulators of decidual genes’, *PLoS ONE*. Edited by S. K. Dey, 4(9), p. e6845. doi: 10.1371/journal.pone.0006845.

Lynch, V. J. *et al.* (2015) ‘Ancient transposable elements transformed the uterine regulatory landscape and transcriptome during the evolution of mammalian pregnancy’, *Cell Reports*, 10(4), pp. 551–561. doi: 10.1016/j.celrep.2014.12.052.

Marinić, M. and Lynch, V. J. (2019) ‘Derivation of endometrial gland organoids from term post-partum placenta’, *bioRxiv*. biorxiv.org, (2017), p. 753780. doi: 10.1101/753780.

McConaha, M. E. *et al.* (2011) ‘Microarray assessment of the influence of the conceptus on gene expression in the mouse uterus during decidualization’, *Reproduction*. Reproduction, 141(4), pp. 511–527. doi: 10.1530/REP-10-0358.

Means, A. R. *et al.* (1975) ‘Estrogen induction of ovalbumin mRNA: Evidence for transcription control’, *Molecular and Cellular Biochemistry*. Kluwer Academic Publishers, pp. 33–42. doi: 10.1007/BF01732161.

Menon, R. *et al.* (2016) ‘Novel concepts on pregnancy clocks and alarms: Redundancy and synergy in human parturition’, *Human Reproduction Update*, pp. 535–560. doi: 10.1093/humupd/dmw022.

Mesiano, S. *et al.* (2002) ‘Progesterone withdrawal and estrogen activation in human parturition are coordinated by progesterone receptor A expression in the myometrium’, *Journal of Clinical Endocrinology and Metabolism*, 87(6), pp. 2924–2930. doi: 10.1210/jcem.87.6.8609.

Mestre-Citrinovitz, A. C. *et al.* (2015) ‘A suppressive antagonism evidences progesterone and estrogen receptor pathway interaction with concomitant regulation of Hand2, Bmp2 and ERK during early decidualization’, *PLoS ONE*. Public Library of Science, 10(4). doi: 10.1371/journal.pone.0124756.

Moffett-King, A. (2002) ‘Natural killer cells and pregnancy’, *Nature Reviews Immunology*. Nat Rev Immunol, pp. 656–663. doi: 10.1038/nri886.

Moffett, A. and Loke, Y. W. (2004) ‘The immunological paradox of pregnancy: A reappraisal’, *Placenta*. W.B. Saunders Ltd, 25(1), pp. 1–8. doi: 10.1016/S0143-4004(03)00167-X.

Moffett, A., Regan, L. and Braude, P. (2004) ‘Natural killer cells, miscarriage, and infertility’,

British Medical Journal. BMJ, pp. 1283–1285. doi: 10.1136/bmj.329.7477.1283.

Munn, D. H. *et al.* (1998) ‘Prevention of allogeneic fetal rejection by tryptophan catabolism’, *Science*. American Association for the Advancement of Science, 281(5380), pp. 1191–1193. doi: 10.1126/science.281.5380.1191.

Murata, H. *et al.* (2019) ‘Progesterin-induced heart and neural crest derivatives-expressed transcript 2 inhibits angiopoietin 2 via fibroblast growth factor 9 in human endometrial stromal cells’, *Reproductive Biology*. Elsevier Sp. z o.o., 19(1), pp. 14–21. doi: 10.1016/j.repbio.2019.02.005.

Murata, H. *et al.* (2020) ‘The transcription factor HAND2 up-regulates transcription of the IL15 gene in human endometrial stromal cells’, *Journal of Biological Chemistry*. J Biol Chem, p. jbc.RA120.012753. doi: 10.1074/jbc.RA120.012753.

Murphy, S. P. *et al.* (2005) ‘Uterine NK cells mediate inflammation-induced fetal demise in IL-10-null mice’, *The Journal of Immunology*. The American Association of Immunologists, 175(6), pp. 4084–4090. doi: 10.4049/jimmunol.175.6.4084.

Murphy, S. P. *et al.* (2009) ‘Evidence for participation of uterine natural killer cells in the mechanisms responsible for spontaneous preterm labor and delivery’, *American Journal of Obstetrics and Gynecology*. Mosby Inc., 200(3), pp. 308.e1-308.e9. doi: 10.1016/j.ajog.2008.10.043.

Ni, N. and Li, Q. (2017) ‘TGF β superfamily signaling and uterine decidualization’, *Reproductive Biology and Endocrinology*. BioMed Central Ltd. doi: 10.1186/s12958-017-0303-0.

O’Mara, T. A., Spurdle, A. B. and Glubb, D. M. (2019) ‘Analysis of promoter-associated chromatin interactions reveals biologically relevant candidate target genes at endometrial cancer risk loci’, *Cancers*. MDPI AG, 11(10), p. 1440. doi: 10.3390/cancers11101440.

Okada, H. *et al.* (2004) ‘Interleukin-1 inhibits interleukin-15 production by progesterone during in vitro decidualization in human’, *Journal of Reproductive Immunology*. Elsevier Ireland Ltd, 61(1), pp. 3–12. doi: 10.1016/j.jri.2003.10.002.

Okada, S. (2000) ‘Expression of interleukin-15 in human endometrium and decidua’, *Molecular Human Reproduction*. Mol Hum Reprod, 6(1), pp. 75–80. doi: 10.1093/molehr/6.1.75.

Olcese, J. (2012) ‘Circadian aspects of mammalian parturition: A review’, *Molecular and Cellular Endocrinology*. Mol Cell Endocrinol, 349(1), pp. 62–67. doi: 10.1016/j.mce.2011.06.041.

Olcese, J., Lozier, S. and Paradise, C. (2013) ‘Melatonin and the circadian timing of human parturition’, *Reproductive Sciences*. Reprod Sci, 20(2), pp. 168–174. doi: 10.1177/1933719112442244.

Oliver, E. A. *et al.* (2011) ‘Activation of the receptor for advanced glycation end products system in women with severe preeclampsia’, *Journal of Clinical Endocrinology and Metabolism*. J Clin Endocrinol Metab, 96(3), pp. 689–698. doi: 10.1210/jc.2010-1418.

Osman, I. *et al.* (2003) ‘Leukocyte density and pro-inflammatory cytokine expression in human fetal membranes, decidua, cervix and myometrium before and during labour at term’, *Molecular Human Reproduction*. Mol Hum Reprod, 41–45. doi: 10.1093/molehr/gag001.

Paiva, P. *et al.* (2009) ‘Interleukin 11 inhibits human trophoblast invasion indicating a likely role in the decidual restraint of trophoblast invasion during placentation’, *Biology of Reproduction*. Oxford University Press (OUP), 80(2), pp. 302–310. doi: 10.1095/biolreprod.108.071415.

Peng, S. *et al.* (2008) 'Dickkopf-1 secreted by decidual cells promotes trophoblast cell invasion during murine placentation', *Reproduction. Reproduction*, 135(3), pp. 367–375. doi: 10.1530/REP-07-0191.

Pepe, G. J. and Albrecht, E. D. (1995) 'Actions of placental and fetal adrenal steroid hormones in primate pregnancy', *Endocrine Reviews. Endocr Rev*, 16(5), pp. 608–648. doi: 10.1210/edrv-16-5-608.

Pertea, M. *et al.* (2015) 'StringTie enables improved reconstruction of a transcriptome from RNA-seq reads', *Nature Biotechnology. Nature Publishing Group*, 33(3), pp. 290–295. doi: 10.1038/nbt.3122.

Pertea, M. *et al.* (2016) 'Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie and Ballgown', *Nature Protocols. Nature Publishing Group*, 11(9), pp. 1650–1667. doi: 10.1038/nprot.2016.095.

Peters, G. A. *et al.* (2016) 'Inflammatory stimuli increase progesterone receptor-A stability and transpressive activity in myometrial cells', *Endocrinology. Endocrine Society*, 158(1), p. en.2016-1537. doi: 10.1210/en.2016-1537.

Pinto, R. M. *et al.* (1966) 'Influence of estradiol-17 β upon the oxytocic action of oxytocin in the pregnant human uterus', *American Journal of Obstetrics and Gynecology. Am J Obstet Gynecol*, 96(6), pp. 857–862. doi: 10.1016/0002-9378(66)90682-X.

Plunkett, J. *et al.* (2011) 'An evolutionary genomic approach to identify genes involved in human birth timing', *PLoS Genetics*. Edited by G. S. Barsh. Public Library of Science, 7(4), p. e1001365. doi: 10.1371/journal.pgen.1001365.

Ratajczak, C. K., Fay, J. C. and Muglia, L. J. (2010) 'Preventing preterm birth: The past limitations and new potential of animal models', *DMM Disease Models and Mechanisms. Company of Biologists Ltd*, 3(7–8), pp. 407–414. doi: 10.1242/dmm.001701.

Renaud, S. J. *et al.* (2017) 'Natural killer cell deficiency alters placental development in rats', *Biology of Reproduction. Oxford University Press (OUP)*, 96(1), pp. 145–158. doi: 10.1095/biolreprod.116.142752.

Renfree, M. B. (2010) 'Review: Marsupials: Placental mammals with a difference', *Placenta. Placenta*, 31(SUPPL.), pp. S21–S26. doi: 10.1016/j.placenta.2009.12.023.

Renfree, M. and Shaw, G. (2001) 'Reproduction in Monotremes and Marsupials', in *Encyclopedia of Life Sciences*. Chichester, UK: John Wiley & Sons, Ltd. doi: 10.1038/npg.els.0001856.

Rinaldi, S. F. *et al.* (2011) 'Anti-inflammatory mediators as physiological and pharmacological regulators of parturition', *Expert Review of Clinical Immunology. Expert Rev Clin Immunol*, pp. 675–696. doi: 10.1586/eci.11.58.

Rinaldi, S. F. *et al.* (2014) 'Immune cells and preterm labour: Do invariant NKT cells hold the key?', *Molecular Human Reproduction. Mol Hum Reprod*, 21(4), pp. 309–312. doi: 10.1093/molehr/gav002.

Rinehart, C. A., Lyn-Cook, J. R. B. D. and Kaufman, D. G. (1988) 'Gland Formation From Human Endometrial Epithelial Cells in vitro', *In Vitro Cellular and Developmental Biology*, 24(10).

Robson, A. *et al.* (2012) 'Uterine natural killer cells initiate spiral artery remodeling in human

pregnancy', *FASEB Journal*. FASEB J, 26(12), pp. 4876–4885. doi: 10.1096/fj.12-210310.

Roizen, J. *et al.* (2007) 'Oxytocin in the circadian timing of birth', *PLoS ONE*. PLoS One, 2(9). doi: 10.1371/journal.pone.0000922.

Sakabe, N. *et al.* (2020) 'Transcriptome and regulatory maps of decidua-derived stromal cells inform gene discovery in preterm birth', *bioRxiv*. Cold Spring Harbor Laboratory, p. 2020.04.06.017079. doi: 10.1101/2020.04.06.017079.

Scholzen, T. and Gerdes, J. (2000) 'The Ki-67 protein: From the known and the unknown', *Journal of Cellular Physiology*. J Cell Physiol, pp. 311–322. doi: 10.1002/(SICI)1097-4652(200003)182:3<311::AID-JCP1>3.0.CO;2-9.

Shen, L. *et al.* (2010) 'Transcription factor HAND2 mutations in sporadic Chinese patients with congenital heart disease', *Chinese Medical Journal*, 123(13), pp. 1623–1627. doi: 10.3760/cma.j.issn.0366-6999.2010.13.002.

Shindoh, H. *et al.* (2014) 'Requirement of heart and neural crest derivatives—expressed transcript 2 during decidualization of human endometrial stromal cells in vitro', *Fertility and Sterility*. Elsevier Inc., 101(6), pp. 1781-1790.e5. doi: 10.1016/j.fertnstert.2014.03.013.

Shynlova, O. *et al.* (2013) 'Myometrial immune cells contribute to term parturition, preterm labour and post-partum involution in mice', *Journal of Cellular and Molecular Medicine*, 17(1), pp. 90–102. doi: 10.1111/j.1582-4934.2012.01650.x.

Smith, R. *et al.* (2009) 'Patterns of plasma corticotropin-releasing hormone, progesterone, estradiol, and estriol change and the onset of human labor', *Journal of Clinical Endocrinology and Metabolism*. Endocrine Society, 94(6), pp. 2066–2074. doi: 10.1210/jc.2008-2257.

Smith, S. D. *et al.* (2009) 'Evidence for immune cell involvement in decidual spiral arteriole remodeling in early human pregnancy', *American Journal of Pathology*. American Society for Investigative Pathology Inc., 174(5), pp. 1959–1971. doi: 10.2353/ajpath.2009.080995.

Smyth, G. K. *et al.* (2002) 'limma: Linear models for microarray and RNA-Seq data user's guide'. Available at: <https://bioconductor.org/packages/release/bioc/vignettes/limma/inst/doc/usersguide.pdf>.

Sonderegger, S., Pollheimer, J. and Knöfler, M. (2010) 'Wnt signalling in implantation, decidualisation and placental differentiation - Review', *Placenta*. W.B. Saunders Ltd, 31(10), pp. 839–847. doi: 10.1016/j.placenta.2010.07.011.

Srivastava, D. *et al.* (1997) 'Regulation of cardiac mesodermal and neural crest development by the bHLH transcription factor, dHAND', *Nature Genetics*. Nat Genet, 16(2), pp. 154–160. doi: 10.1038/ng0697-154.

Šućurović, S. *et al.* (2020) 'Analysis of heart and neural crest derivatives-expressed protein 2 (HAND2)-progesterone interactions in peri-implantation endometrium', *Biology of Reproduction*. Oxford Academic, 102(5), pp. 1111–1121. doi: 10.1093/biolre/ioaa013.

Sun, Y. M. *et al.* (2016) 'A HAND2 loss-of-function mutation causes familial ventricular septal defect and pulmonary stenosis', *G3: Genes, Genomes, Genetics*. Genetics Society of America, 6(4), pp. 987–992. doi: 10.1534/g3.115.026518.

Swaggart, K. A., Pavlicev, M. and Muglia, L. J. (2015) 'Genomics of preterm birth', *Cold Spring Harbor Perspectives in Medicine*. Cold Spring Harbor Laboratory Press, 5(2), pp. a023127–a023127. doi: 10.1101/csdperspect.a023127.

Talbi, S. *et al.* (2006) 'Molecular phenotyping of human endometrium distinguishes menstrual cycle phases and underlying biological processes in normo-ovulatory women', *Endocrinology*. Endocrinology, 147(3), pp. 1097–1121. doi: 10.1210/en.2005-1076.

Tamura, M. *et al.* (2013) 'Overdosage of Hand2 causes limb and heart defects in the human chromosomal disorder partial trisomy distal 4q', *Human Molecular Genetics*, 22(12), pp. 2471–2481. doi: 10.1093/hmg/ddt099.

Thomson, A. J. *et al.* (1999) 'Leukocytes infiltrate the myometrium during human parturition: Further evidence that labour is an inflammatory process', *Human Reproduction*, 14(1), pp. 229–236. doi: 10.1093/humrep/14.1.229.

Turco, M. Y. *et al.* (2017) 'Long-term, hormone-responsive organoid cultures of human endometrium in a chemically defined medium', *Nature Cell Biology*, 19(5), pp. 568–577. doi: 10.1038/ncb3516.

Turco, M. Y. *et al.* (2018) 'Trophoblast organoids as a model for maternal–fetal interactions during human placentation', *Nature*. Nature Publishing Group, 564(7735), pp. 263–281. doi: 10.1038/s41586-018-0753-3.

Uhlén, M. *et al.* (2015) 'Tissue-based map of the human proteome', *Science*. American Association for the Advancement of Science, 347(6220), pp. 1260419–1260419. doi: 10.1126/science.1260419.

Varas Enriquez, P. J., McKerracher, L. J. and Elliot, M. G. (2018) 'Pre-eclampsia and maternal–fetal conflict', *Evolution, Medicine, and Public Health*, 2018(1), pp. 217–218. doi: 10.1093/emph/eoy029.

Varki, A. (2012) 'Nothing in medicine makes sense, except in the light of evolution', *Journal of Molecular Medicine*. J Mol Med (Berl), pp. 481–494. doi: 10.1007/s00109-012-0900-5.

Varki, N. *et al.* (2009) 'Heart disease is common in humans and chimpanzees, but is caused by different pathological processes', *Evolutionary Applications*. Wiley-Blackwell, 2(1), pp. 101–112. doi: 10.1111/j.1752-4571.2008.00064.x.

Varki, N. M. and Varki, A. (2015) 'On the apparent rarity of epithelial cancers in captive chimpanzees', *Philosophical Transactions of the Royal Society B: Biological Sciences*. Royal Society of London. doi: 10.1098/rstb.2014.0225.

Vento-Tormo, R. *et al.* (2018) 'Single-cell reconstruction of the early maternal–fetal interface in humans', *Nature*. Nature Publishing Group, 563(7731), pp. 347–353. doi: 10.1038/s41586-018-0698-6.

Verma, S. *et al.* (2000) 'Human decidual natural killer cells express the receptor for and respond to the cytokine Interleukin 15', *Biology of Reproduction*. Oxford University Press (OUP), 62(4), pp. 959–968. doi: 10.1095/biolreprod62.4.959.

Wagner, G. P., Kin, K. and Lynch, V. J. (2012) 'Measurement of mRNA abundance using RNA-seq data: RPKM measure is inconsistent among samples', *Theory in Biosciences*, 131(4), pp. 281–285. doi: 10.1007/s12064-012-0162-3.

Wagner, G. P., Kin, K. and Lynch, V. J. (2013) 'A model based criterion for gene expression calls using RNA-seq data', *Theory in Biosciences*. Springer, 132(3), pp. 159–164. doi: 10.1007/s12064-013-0178-3.

Wallace, A. E. *et al.* (2013) 'Decidual natural killer cell interactions with trophoblasts are

impaired in pregnancies at increased risk of preeclampsia', *American Journal of Pathology*. American Society for Investigative Pathology, 183(6), pp. 1853–1861. doi: 10.1016/j.ajpath.2013.08.023.

Wallace, A. E. *et al.* (2014) 'Increased angiogenic factor secretion by decidual natural killer cells from pregnancies with high uterine artery resistance alters trophoblast function', *Human Reproduction*, 29(4), pp. 652–660. doi: 10.1093/humrep/deu017.

Wang, Q. *et al.* (2013) 'Wnt6 Is essential for stromal cell proliferation during decidualization in mice', *Biology of Reproduction*. Oxford University Press (OUP), 88(1). doi: 10.1095/biolreprod.112.104687.

Warrington, N. M. *et al.* (2019) 'Maternal and fetal genetic effects on birth weight and their relevance to cardio-metabolic risk factors', *Nature Genetics*. Nature Publishing Group, 51(5), pp. 804–814. doi: 10.1038/s41588-019-0403-1.

Welsh, T. *et al.* (2012) 'Estrogen receptor (ER) expression and function in the pregnant human myometrium: Estradiol via ER α activates ERK1/2 signaling in term myometrium', *Journal of Endocrinology*. J Endocrinol, 212(2), pp. 227–238. doi: 10.1530/JOE-11-0358.

Wetendorf, M. and DeMayo, F. J. (2012) 'The progesterone receptor regulates implantation, decidualization, and glandular development via a complex paracrine signaling network', *Molecular and Cellular Endocrinology*, pp. 108–118. doi: 10.1016/j.mce.2011.10.028.

Williams, P. J. *et al.* (2009) 'Decidual leucocyte populations in early to late gestation normal human pregnancy', *Journal of Reproductive Immunology*. J Reprod Immunol, 82(1), pp. 24–31. doi: 10.1016/j.jri.2009.08.001.

Wilson, R. A. and Mesiano, S. A. (2020) 'Progesterone signaling in myometrial cells: role in human pregnancy and parturition', *Current Opinion in Physiology*. Elsevier Ltd, pp. 117–122. doi: 10.1016/j.cophys.2019.09.007.

Winn, V. D. *et al.* (2007) 'Gene expression profiling of the human maternal-fetal interface reveals dramatic changes between midgestation and term', *Endocrinology*. Endocrinology, 148(3), pp. 1059–1079. doi: 10.1210/en.2006-0683.

Ying, Y. and Zhao, G.-Q. (2000) 'Detection of multiple Bone Morphogenetic Protein messenger ribonucleic acids and their signal transducer, Smad1, during mouse decidualization', *Biology of Reproduction*. Oxford University Press (OUP), 63(6), pp. 1781–1786. doi: 10.1095/biolreprod63.6.1781.

Young, A. *et al.* (2002) 'Immunolocalization of proinflammatory cytokines in myometrium, cervix, and fetal membranes during human parturition at term', *Biology of Reproduction*. Oxford University Press (OUP), 66(2), pp. 445–449. doi: 10.1095/biolreprod66.2.445.

Zhang, J., Dunk, C. E. and Lye, S. J. (2013) 'Sphingosine signalling regulates decidual NK cell angiogenic phenotype and trophoblast migration', *Human Reproduction*, 28(11), pp. 3026–3037. doi: 10.1093/humrep/det339.

Zhang, Y. *et al.* (2008) 'Model-based analysis of ChIP-Seq (MACS)', *Genome Biology*. Genome Biol, 9(9). doi: 10.1186/gb-2008-9-9-r137.

Zhu, X. M. *et al.* (2009) 'Conditioned medium from human decidual stromal cells has a concentration-dependent effect on trophoblast cell invasion', *Placenta*. Placenta, 30(1), pp. 74–78. doi: 10.1016/j.placenta.2008.09.013.

Zygmunt, M. *et al.* (1998) 'Invasion of cytotrophoblastic (JEG-3) cells is up-regulated by interleukin-15 in vitro', *American Journal of Reproductive Immunology*. Blackwell Munksgaard, 40(5), pp. 326–331. doi: 10.1111/j.1600-0897.1998.tb00061.x.