

The abundant-centre is not all that abundant: a comment to Osorio-Olvera et al. 2020

Tad Dallas^{a, †, *}, Samuel Pironon^{b, *} and Luca Santini^{c, *}

^a*Department of Biological Sciences, Louisiana State University, Baton Rouge, USA*

^b*Royal Botanic Gardens, Kew, Richmond, UK*

^c*National Research Council, Institute of Research on Terrestrial Ecosystems (CNR-IRET), Via Salaria km 29.300, 00015, Monterotondo (Rome), Italy*

* All authors contributed equally

† Corresponding author: tad.a.dallas@gmail.com

¹ **Author contributions:** All authors contributed equally.

²

³ **Data accessibility:** R code is available on figshare at

⁴ <https://figshare.com/s/8fafdf780810e73d44623>.

⁵

⁶ **Funding:** TAD was supported by Louisiana State University.

⁷

⁸ **Competing interests:** The authors declare no competing interests.

⁹

¹⁰ **Abstract**

¹¹ Species abundance is expected to decrease from the centre towards the edge of their
¹² ecological niches (abundant niche-centre hypothesis). Recently, Osorio-Olvera
¹³ *et al.* (2020) reported strong support for the abundant niche-centre relationship
¹⁴ in North American birds. We demonstrate here that methodological decisions
¹⁵ strongly affected perceived support. Avoiding these issues casts doubt on conclu-
¹⁶ sions by Osorio-Olvera *et al.* and the putative support for the abundant niche-
¹⁷ centre hypothesis in North American birds.

18 The spatial distribution of abundance has long fascinated ecologists who searched
19 for general rules governing where species occur and the density at which they are
20 found (McGill *et al.*, 2007; Sagarin & Gaines, 2002). Particularly controversial
21 rules are the abundant-centre and abundant niche-centre hypotheses, which pre-
22 dict abundance to decrease gradually from the centre to the margins of species
23 geographic ranges and ecological niches respectively (Brown, 1984; Pironon *et al.*,
24 2017). Both theories have received mixed empirical support (Martínez-Meyer
25 *et al.*, 2013; Sagarin & Gaines, 2002; Dallas *et al.*, 2017) and limited theoretic-
26 al development (Osorio-Olvera *et al.*, 2019; Holt, 2019; Dallas & Santini, 2020).
27 Moreover, recent analyses highlighted that tests of these hypotheses were sensi-
28 tive to the quality of the input data and the methodological approach considered
29 (Santini *et al.*, 2019).

30 Osorio-Olvera *et al.* (2020) analyze data from the North American Breeding Bird
31 Survey (BBS) to test for a negative correlation between species abundance and the
32 distance to their climatic niche centroid. Counter to recent findings questioning
33 its generalizability (Sagarin & Gaines, 2002; Dallas *et al.*, 2017; Santini *et al.*,
34 2019), the authors claimed general support for the hypothesis and proposed that
35 the distance to species climatic niche centroid (quantified using minimum volume
36 ellipsoids) could represent a reliable and simple new metric to predict the current
37 and future distribution of species abundance. However, we discuss how serious
38 problems related to data quality, modelling choice, and presentation of the results,
39 prevent from making any reliable conclusion, and can greatly affect the perceived
40 support for the hypothesis.

41 First of all, many of the species considered in the study also occur well be-
42 yond the study area (e.g. *Ardea alba*, *Corvus corax*), and some only share a very
43 small portion of the range in the study area (e.g. *Thalasseus maximus*, *Aramus*

⁴⁴ *guarauna*). We calculated geographic and climatic niche overlap of a convex hull
⁴⁵ encompassing the BBS data with the BirdLife International data (BirdLife Inter-
⁴⁶ national, 2017) only considering the resident and breeding range, demonstrating
⁴⁷ a clear influence on the estimation of the geographic range and climatic niche
⁴⁸ boundaries, as well as their centroid distance (Figure 1). This subsequently af-
⁴⁹ fects the abundant niche-centre relationship, as discussed in Soberón *et al.* (2018),
⁵⁰ questioning the validity of the relationships estimated. Oddly, many of the bird
⁵¹ species whose geographic ranges are underestimated and whose niches have been
⁵² underestimated also exhibit significant negative abundant-centre relationships (e.g.
⁵³ *Tyrannus couchii*, *Thalasseus maximus*, *Glaucidium gnoma*), putatively support-
⁵⁴ ing the hypothesis.

⁵⁵ The authors found that the percentage of species range overlap (Table 1 in
⁵⁶ Osorio-Olvera *et al.* (2020)) had a non-significant effect on the correlation coeffi-
⁵⁷ cients (multivariate analysis), thus arguing that range overlap does not affect their
⁵⁸ conclusions. However, this may not account for the effect of niche truncation, as
⁵⁹ 1) geographic overlap does not necessarily translate into niche overlap (Fig. 1c,d),
⁶⁰ which is also why the authors estimate niche centres instead of geographic centres,
⁶¹ and 2) the location of niche centres are still biased towards climatic conditions of
⁶² the study area, which affects the calculation of centroids and distances (Fig. 1b).
⁶³ On a more fundamental level, testing if abundant-centre relationships differ as a
⁶⁴ function of range overlap does not address the influence of range overlap directly,
⁶⁵ but makes the assumption that as long as correlation coefficients do not differ as
⁶⁶ a function of range overlap, then the range centroid distances were estimated ap-
⁶⁷ propriately. This is not a clear test of the influence of range overlap, and risks the
⁶⁸ fallacy of *asserting the null*. We note that a biased estimation of the niche centre
⁶⁹ is supposed to matter in such an analysis, a non-significant difference suggests that

70 using high-quality data does not increase the support rate for the hypothesis.

71 The strongest support for the abundant niche-centre relationships comes from
72 Osorio-Olvera *et al.* (2020) estimating the species niche as a minimum volume
73 ellipsoid (MVE) by considering more than 4000 combinations of climatic variables,
74 including all 19 commonly-used bioclimatic variables together with the first 15
75 PCA components of a PCA based on the same bioclimatic variables. The authors
76 use every possible combination of two and three niche axes to estimate the niche.
77 We identify two main issues associated with this procedure.

78 First, the authors report results only for models showing significant abundant
79 niche-centre relationships, omitting non-significant correlations (Figure 2a). This
80 issue is not only present in the fit MVE models, but also in the 2 and 3 feature
81 models using convex hulls or MVEs. The effects of this are clear (Figure 2). By
82 including non-significant correlations, the mean abundant niche-centre relationship
83 across all model sets becomes weak ($\bar{\rho} \pm sd = -0.08 \pm 0.01$), and more species exhibit
84 significantly positive abundant niche-centre relationships (Figure 2). Including
85 these non-significant results is important, in our view, and strongly influences
86 the resulting perceived support for the abundant niche-centre pattern (Figure 2).
87 Presenting also non-significant results demonstrates that only between 37% and
88 45% of species have negative abundant-centre relationships, regardless of approach
89 used (see <https://figshare.com/s/8fadf780810e73d44623>), while the majority of the
90 estimated relationships are either positive or non-significant. Interestingly, this low
91 empirical support is consistent with previous findings (Dallas *et al.*, 2017; Pironon
92 *et al.*, 2017; Sagarin & Gaines, 2002; Santini *et al.*, 2019).

93 Second, while the authors train an average of 1,852 models per species to calcu-
94 late MVEs, they perform no form of model selection (i.e., excluding models based

95 on omission rate is thresholding, not model selection). This functionally treats
96 the poorest fit MVE and the best fit MVE per species as equivalent, provided
97 the model produced a significant abundant-centre relationship. This condition
98 results in between 1 and 3460 abundant-centre estimates for any given species,
99 introducing substantial bias in estimation of the distribution of abundant-centre
100 relationships. When non-significant results are included, and only best models are
101 retained, the overall pattern changes substantially (Figure 2). When only the best
102 fit models are considered, 115 out of 379 species (30%) had significant abundant
103 niche-centre relationships, with a mean correlation coefficient of -0.07. Some of
104 these best models had higher omission rates than what Osorio-Olvera *et al.* (2020)
105 considered. Removing these models reduces the number of species down to 303
106 species, of which 94 had significantly negative abundant niche-centre relationships
107 (31%), while 180 and 29 had non-significant (59%) or significantly positive (10%)
108 relationships, respectively (Figure 2c). It is not our assertion that abundant-centre
109 relationships do not exist. The negative relationships found by Osorio-Olvera *et al.*
110 (2020) support the idea of an abundant-centre, but do so in a misleading manner.

111 The study from Osorio-Olvera *et al.* (2020) highlights the timely need for dis-
112 entangling the complex relationship between species ecological niche, geographic
113 distribution and demographic performance (Holt, 2019; Bohner & Diez, 2020). Ex-
114 plaining the convergence and divergence of results of studies exploring occurrence
115 and abundance patterns is key for improving our understanding of biodiversity
116 and ability to predict its response to ongoing changes in the global environment.

117 References

118 BirdLife International (2017). Bird species distribution maps of the world. version
119 6.0. Birdlife International, Cambridge, United Kingdom, and Lynx Edicions,
120 Barcelona, Spain. Available from <http://datazone.birdlife.org/species/requestdis>
121 (accessed June 2017).

122 Blonder, B., Blonder, M.B., Rcpp, D. & Rcpp, L. (2015). Package hypervolume.
123 *Global Ecology and Biogeography*, 23, 595–609.

124 Bohner, T. & Diez, J. (2020). Extensive mismatches between species distributions
125 and performance and their relationship to functional traits. *Ecology Letters*, 23,
126 33–44.

127 Brown, J.H. (1984). On the relationship between abundance and distribution of
128 species. *The American Naturalist*, 124, 255–279.

129 Dallas, T., Decker, R.R. & Hastings, A. (2017). Species are not most abundant
130 in the centre of their geographic range or climatic niche. *Ecology letters*, 20,
131 1526–1533.

132 Dallas, T.A. & Santini, L. (2020). The influence of stochasticity, landscape struc-
133 ture, and species traits on abundant-centre relationships. *Ecography*.

134 Holt, R.D. (2019). Reflections on niches and numbers. *Ecography*, pp. 1–4.

135 Martínez-Meyer, E., Díaz-Porras, D., Peterson, A.T. & Yáñez-Arenas, C. (2013).
136 Ecological niche structure and rangewide abundance patterns of species. *Biology*
137 *Letters*, 9, 20120637.

138 McGill, B.J., Etienne, R.S., Gray, J.S., Alonso, D., Anderson, M.J., Benecha, H.K.,
139 Dornelas, M., Enquist, B.J., Green, J.L., He, F. *et al.* (2007). Species abundance

140 distributions: moving beyond single prediction theories to integration within an
141 ecological framework. *Ecology Letters*, 10, 995–1015.

142 Osorio-Olvera, L., Soberón, J. & Falconi, M. (2019). On population abundance
143 and niche structure. *Ecography*, 42, 1415–1425.

144 Osorio-Olvera, L., Yañez-Arenas, C., Martínez-Meyer, E. & Peterson, A.T. (2020).
145 Relationships between population densities and niche-centroid distances in
146 North American birds. *Ecology Letters*.

147 Pironon, S., Papuga, G., Villegas, J., Angert, A.L., García, M.B. & Thompson,
148 J.D. (2017). Geographic variation in genetic and demographic performance:
149 new insights from an old biogeographical paradigm. *Biological Reviews*, 92,
150 1877–1909.

151 Sagarin, R.D. & Gaines, S.D. (2002). The “abundant centre” distribution: to what
152 extent is it a biogeographical rule? *Ecology Letters*, 5, 137–147.

153 Santini, L., Pironon, S., Maiorano, L. & Thuiller, W. (2019). Addressing common
154 pitfalls does not provide more support to geographical and ecological abundant-
155 centre hypotheses. *Ecography*, 42, 696–705.

156 Soberón, J., Peterson, A.T. & Osorio-Olvera, L. (2018). A comment on "Species
157 are not most abundant in the center of their geographic range or climatic niche".
158 *Rethinking Ecology*, pp. 13–18.

¹⁵⁹ **Figures**




Figure 1: Mismatch in geographic and niche estimates between abundance data used in Osorio-Olvera *et al.* (2020) and the combined resident and breeding range species distributions, estimated IUCN range polygons. **a)** Lesser nighthawk (*Chordeiles acutipennis*) IUCN geographic range (in blue) and sample data to estimate the niche (in red); **b)** First two PCA axes of all bioclimatic variables showing environmental values considered in the study (red triangles) and those estimated considering the cells in the IUCN range (blue dots). The darker and larger triangle and circle represent the estimated centroids of the two hypervolumes; **c)** Distribution of geographic range overlap between convex hulls drawn around abundance estimates and the IUCN ranges for all species considered in the study; **d)** Distribution of niche overlap between convex hulls drawn around abundance estimates and grid cells within the IUCN ranges for all species in the study; **e)** Percentage of centroid displacement over the truncated niche; **f)** Percentage of centroid displacement over the full niche. Niche overlap and niche centroids were estimated using the hypervolume package Blonder *et al.* (2015).




Figure 2: Differences in support for the abundant niche-centre hypothesis for a set of North American birds as a function of analytical decisions. We reproduce the results of Osorio-Olvera *et al.* (2020), demonstrating largely significant and negative abundant-centre relationships (panel a). However, by including all correlation coefficients, instead of only the significant ones, support for abundant niche-centre relationships become weak and largely non-significant (panel b). We also include the results when only considering the best fit MVE models per species ('best fit MVE') when non-significant relationships were excluded (panel a) and included (panel b). Significance values (**** $p \leq 0.0001$, *** $p \leq 0.001$, ** $p \leq 0.01$, * $p \leq 0.05$, ns not significant), compare all other methods to the MVE < 1% case.