

SOFTWARE

gapseq: Informed prediction of bacterial metabolic pathways and reconstruction of accurate metabolic models

Johannes Zimmermann¹, Christoph Kaleta¹ and Silvio Waschina^{1,2*}

*Correspondence:

s.waschina@nutrinf.uni-kiel.de

²Christian-Albrechts-University
Kiel, Institute of Human Nutrition
and Food Science,
Nutriinformatics,
Heinrich-Hecht-Platz 10, 24118
Kiel, Germany

Full list of author information is
available at the end of the article

Abstract

Microbial metabolic processes greatly impact ecosystem functioning and the physiology of multi-cellular host organisms. The inference of metabolic capabilities and phenotypes from genome sequences with the help of reference biomolecular knowledge stored in online databases remains a major challenge in systems biology. Here, we present gapseq: a novel tool for automated pathway prediction and metabolic network reconstruction from microbial genome sequences. gapseq combines databases of reference protein sequences (UniProt, TCDB), in tandem with pathway and reaction databases (MetaCyc, KEGG, ModelSEED). This enables the prediction of an organism's metabolic capabilities from sequence homology and pathway topology criteria. By incorporating a novel LP-based gap-filling algorithm, gapseq facilitates the construction of genome-scale metabolic models that are suitable for metabolic phenotype predictions by using constraint-based flux analysis. We validated gapseq by comparing predictions to experimental data for more than 3,000 bacterial organisms comprising 14,895 phenotypic traits that include enzyme activity, energy sources, fermentation products, and gene essentiality. This large-scale phenotypic trait prediction test showed, that gapseq yields an overall accuracy of 81% and thereby outperforms other commonly used reconstruction tools. Furthermore, we illustrate the application of gapseq-reconstructed models to simulate biochemical interactions between microorganisms in multi-species communities. Altogether, gapseq is a new method that improves the predictive potential of automated metabolic network reconstructions and further increases their applicability in biotechnological, ecological, and medical research. gapseq is available at <https://github.com/jotech/gapseq>.

Keywords: Metabolic pathway analysis; Metabolic networks; Genome-scale metabolic models; Benchmark; Community simulation; Microbiome; Metagenome

1 Background

Anything you have to do repeatedly may be ripe for automation.

— Doug McIlroy

- Metabolism is central for organismal life. It provides metabolites and energy for all cellular processes. A majority of metabolic reactions are catalysed by enzymes, which are encoded in the genome of the respective organism. Those catalysed reactions form a complex metabolic network of numerous biochemical transformations, which the organism is presumably able to perform [1].
- In systems biology, the reconstruction of metabolic networks plays an essential role,

as the network represents an organism's capabilities to interact with its biotic and abiotic environment and to transform nutrients into biomass. Mathematical analysis has shown great potential for dissecting the functioning of metabolic networks on the level of topological, stoichiometric, and kinetic models [2], which together provide a wide array of methods [3]. Although different microbial metabolic modelling approaches exist, they can be summarised by a theoretical framework that provides a unifying view on microbial growth [4]. Metabolic models not only have demonstrated their ability to predict phenotypes on the level of cellular growth and gene knockouts, but also provide potential molecular mechanisms in form of gene and reaction activities, which can be validated experimentally [5]. Due to this predictive potential, genome-scale metabolic models have been applied to identify metabolic interactions between different organisms [6, 7, 8, 9, 10], to study host-microbiome interactions [11, 12, 13], to predict novel drug targets to fight microbial pathogens [14, 15], and for the rational design of microbial genotypes and growth-media conditions for the industrial production or degradation of biochemicals [16, 17]. Furthermore, recent advances in DNA-sequencing technologies have led to a vast increase in available genomic- and metagenomic sequences in databases [18], which further expands the applicability of genome-scale metabolic network reconstructions.

The reconstruction of metabolic networks links genomic content with biochemical reactions and therefore depends on sequence annotations and reaction databases, which are both crucial for overall network quality [19, 20]. A general problem in reconstructing metabolic networks occurs by an incorrect representation of the organism's physiology. First, inconsistencies in databases can lead to an incorporation of imbalanced reactions into the metabolic network, which may become responsible for incorrect energy production by futile cycles [20]. Second, many genes are lacking a functional annotation due to a lack of knowledge [21] and, thus, also the gene products cannot be integrated into the metabolic networks, which potentially lead to gaps in pathways. Third, the gap-filling of metabolic networks is frequently done by adding a minimum number of reactions from a reference database that facilitate growth under a chemically defined growth medium [22, 23, 24]. Such approaches miss further evidences potentially hidden in sequences and are biased towards the growth medium used for gap-filling. And fourth, the validation of predictions made by metabolic networks is so far only performed with smaller experimental data sets from model laboratory strains such as *Escherichia coli* K12 or *Bacillus subtilis* 168 and therefore the overall performance of many metabolic models is insufficiently assured.

Genome-scale metabolic network reconstructions are increasingly applied to simulate complex metabolic processes in microbial communities [25]. Such simulations are highly sensitive to the quality of the individual metabolic networks of the community members. This is because the accurate prediction of fermentation products and carbon source utilisation is crucial for the correct prediction of metabolic interactions since the substances produced by one organism may serve as resource for others [26]. Thus, in multi-species communities, the metabolic fluxes of organisms are intrinsically connected, which can lead to error propagation when one defective model affects otherwise correctly working models. As a consequence, the feasibility of community modeling intrinsically depends on the accuracy of the individual organismal models.

In this work, we present **gapseq** a novel software for pathway analysis and metabolic network reconstruction. The pathway prediction is based on multiple biochemistry databases that comprise information on pathway structures, the pathways' key enzymes, and reaction stoichiometries. Moreover, **gapseq** constructs genome-scale metabolic models that enable metabolic phenotype predictions as well as the application in simulations of community metabolism. Models are constructed using a manually curated reaction database that is free of energy-generating thermodynamically infeasible reaction cycles. As input, **gapseq** takes the organism's genome sequence in FASTA format, without the need for an additional annotation file. Topology as well as sequence homology to reference proteins inform the filling of network gaps, and the screening for potential carbon sources and metabolic products is done in a way that reduces the impact of growth medium definitions. Finally, we used large-scale experimental data sets to validate enzyme activity, carbon source utilisation, fermentation products, gene essentiality, and metabolite-cross feeding interactions in microbial communities.

2 Results

2.1 Biochemistry database and universal model

The pathway-, transporter, and complex prediction is based on a protein sequence database that is derived from UniProt as well as TCDB and consists in total of 130,671 unique sequences (111,542 reviewed unipac 0.9 clusters and 19,129 TCDB transporter) and also 1,131,132 unreviewed unipac 0.5 cluster that can be included optionally. In addition, the protein sequence database in **gapseq** can be updated to include new sequences from Uniprot and TCDB. For the construction of genome-scale metabolic network models we have built a biochemistry database, that is derived from the ModelSEED biochemistry database. In total, the resulting curated **gapseq** metabolism database comprises 14,287 reactions (including transporters) and 7,570 metabolites. All metabolites and reactions from the biochemistry database are incorporated in the universal model that **gapseq** utilises for the gap-filling algorithm. When removing all dead-end metabolites and corresponding reactions, the universal model comprises 10,194 reactions and 3,337 metabolites. It needs to be noted, that the current biochemistry database and the derived universal model represents bacterial metabolic functions and that, at the current version of **gapseq**, the database does not include archaea-specific reactions. However, those reactions and, thus, also the possibility to use **gapseq** for the reconstruction of archaeal models will be included in an later version of the software.

2.2 Agreement with enzymatic data (BacDive)

We used experimental data of active metabolic enzymes to compare the accuracy of model generation pipelines. In total, we compared 10,538 enzyme activities, comprising 30 unique enzymes, in 3,017 organisms. For all organisms, genome-scale metabolic models were constructed using three different pipelines (CarveMe[39], **gapseq**, ModelSEED[24]). **gapseq** models had with 6% the lowest false-negative rate compared to CarveMe (32%) and ModelSEED (28%). Correspondingly, **gapseq** showed with 53% also highest true positive rate compared to CarveMe (27%) and ModelSEED (30%), while the rates of false positive and true negative predictions

were comparable (Figure 1A). For this test, the most prominent EC numbers were the catalase, 1.11.1.6, accounting for 26% of the comparisons and the cytochrome oxidase, 1.9.3.1, accounting for 22%.

2.3 Validation of carbon source usage (ProTraits)

Growth predictions are essential for metabolic models. We checked the quality of model generation pipelines to predict the growth on different carbon sources. In summary, we compared 1,795 different growth prediction for 526 organism and 48 carbon sources (Figure 1B). *gapseq* outperformed the other methods in terms of false negatives (14% compared with 29% ModelSEED and 37% CarveMe) and true positives (45% compared with 31% ModelSEED and 23% CarveMe). ModelSEED showed fewer false positives (5% compared with 10% *gapseq* and 11% CarveMe) and more true negatives (35% compared with 30% *gapseq* and 30% CarveMe). *gapseq*, predicted most false positives for formate (29 times). This overestimate of formate as potential carbon source is likely due to the fact that we tested carbon source utilisation on the basis of electron transfer from the source to electron carriers (i.e. ubiquinol, menaquinol, or NADH), which is analogous to the experimental carbon source test of BIOLOG plates [46]. However, while it is known that formate can serve in fact as electron donor in a number of different bacteria [84], the role as source of carbon atoms for the synthesis of biomass components is limited to a few known methyloprophs [85].

Across all methods, the most accurately predicted carbon sources, with more than 100 tested organisms, were fructose (91% correct predictions), mannose (89%), or arginine (84%), whereby less good predictions were obtained for arabinose (29% correct predictions), dextrin (40%), or acetate (42%).





Figure 1: Results from enzyme activity and carbon source validations.

A) In total 10,538 enzyme activities (30 enzymes and 3,017 organisms) from experimental standardised experiments from the DSMZ BacDive database were compared for three different methods. B) The predictions of 1,795 carbon sources (48 unique carbon sources and 526 organisms) were validated with data from the ProTraits database.

124

2.4 Gene essentiality

125




We compared the ability of *gapseq* models to predict the essentiality of genes with predictions from ModelSEED and CarveMe reconstructions as well as with curated models for the same organisms (Figure 2). As expected, the curated models outperform all three automated reconstruction tools for most species and prediction metrics (namely precision, sensitivity, specificity, accuracy, and F1-score). Interestingly, for *Pseudomonas aeruginosa* the *gapseq* model shows better gene essentiality predictions in terms of sensitivity, accuracy, and F1-score than the curated model (Figure 2D). Compared to CarveMe, *gapseq* shows generally a higher sensitivity in essentiality predictions but, at the same time, a lower precision rate. This pattern is attributed to the fact, that *gapseq* models tend to predict more genes as essential than CarveMe, leading to a higher number of true positive (TP) predictions but also more false positives (FP). For most organisms and on the basis of most prediction metrics, *gapseq* outperforms network models that were reconstructed using ModelSEED.

2.5 Fermentation products

Anaerobic or facultative anaerobic bacteria utilise different fermentation pathways in order to extract energy from environmental compounds by chemical transformations in the absence of oxygen. We tested if the identity of fermentation products can be predicted by metabolic network model constructions obtained from *gapseq*, CarveMe, and ModelSEED for 18 different bacterial organisms (Figure 2).

The organisms were selected based on following criteria: (1) the organisms have a published RefSeq genome sequence [52], (2) are known anaerobic or facultative anaerobic organisms, and (3) the identity of fermentation products has been experimentally described and reported in primary literature (Suppl. table S2). Overall,




gapseq showed the highest number of true positive predictions (TP) with 36 TP predicted with the Minimize-Total-Flux (MTF) and 37 TP predicted with Flux-Variability-Analysis (FVA) which is substantially higher compared to CarveMe (8 TP with MTF, 10 TP with FVA) and ModelSEED (1 TP, 3 TP). The production of the short-chain-fatty-acids acetate, butyrate, and propionate was correctly predicted by gapseq in 78% of cases and thereby outcompetes CarveMe (9%) and ModelSEED (0%), which did not predict butyrate or propionate production for any organism tested. Moreover, gapseq correctly predicted homolactic fermentation by *Lactobacillus delbrueckii* and *Lactobacillus acidophilus*, which is dominated by lactate as fermentation end-product and also predicted heterolactic fermentation by *Bifidobacterium longum*. However, gapseq failed to predict lactate production of organisms that utilise different fermentation strategies, which also yield lactate (e.g. mixed-acid fermentation by *Escherichia coli*). Interestingly, the predicted quantities of fermentation product release is higher for true positive than for false negative predictions (Figure 3). This further suggests, that gapseq is able to predict the main fermentation products of bacterial organisms during anaerobic growth based on the organism's genome sequence.

167 2.6 Anaerobic food web of the gut microbiome

168 The prediction of metabolic interactions between microbial organisms is of special
169 interest in ecology, medicine, and biotechnology. So far, we showed the capacity
170 of **gapseq** on the level of individual models. In a next step, we simulated several
171 individual models together as a multi-species community to validate the potential
172 of **gapseq** in microbial community modelling. As sample application we selected
173 representative members of the human gut microbiome that are known to form an
174 anaerobic food web [64, 65]. Altogether, we employed 20 organisms and simulated
175 the combined growth in a shared environment for several time steps using the com-
176 munity modeling framework BacArena [68]. On the community level, simulations
177 using **gapseq** models captured all important substances, which are known to be
178 produced in the context of the food web (Figure 4). This included the production
179 of short chain fatty acids (acetate, propionate, butyrate), lactate, hydrogen, hy-
180 drogen sulfide (H₂S), methane, formate, and succinate. The formation of acetate,
181 formate, and hydrogen was most prevalent, which are also common end-products
182 of fermentation. Lactate, succinate, acetate, hydrogen, formate, and H₂S were fur-
183 ther metabolised by some community members (Figure 4). The predicted identity
184 of fermentation end-products and other by-products of metabolism was found to
185 be in line with literature information [64, 65, 86]. For example, the formation of
186 lactate was observed for *Lactobacillus acidophilus* and *Bifidobacterium longum*, and
187 butyrate was released by known butyrate producers, i.e. *Faecalibacterium praus-*
188 *nitzii*, *Anaerobutyricum hallii*, *Clostridium perfringens*, and *Coprococcus* spp.. Es-
189 pecially the main products of mixed acid fermentation (acetate, formate, hydrogen,
190 ethanol) were predicted for most members of the community which is in agreement
191 with what is known about common metabolic end products of many gut-dwelling
192 microorganisms [86]. Interestingly, for *Faecalibacterium prausnitzii* no acetate pro-
193 duction is reported [86], which was also observed in our simulations. Moreover, H₂S
194 was correctly predicted to be produced by *Desulfovibrio desulfuricans*. In general,
195 the anaerobic oxidation of fatty acids is not favored by the gut environment because
196 the host competes for the uptake of butyrate, propionate, and acetate, which serve
197 as energy source for colonic epithelial cells and are involved in many host functions
198 [87]. Therefore, the gut community lacks syntrophic organisms which are able to
199 anaerobically degrade butyrate [88]. In agreement with this, we found no microbial
200 uptake of butyrate in the community simulation. In contrast, lactate was predicted
201 to be produced and consumed by distinct community members. We found utilis-
202 ation of lactate by *Coprococcus comes*, *Megasphaera elsdenii*, and *Veillonella dispar*,
203 which is a known feature of these organisms [64]. In addition, succinate was cor-
204 rectly predicted to be used by *Bacteroides* species [86]. The formation of methane
205 is known to be limited to methanogenic archaea, and thus *Methanosarcina barkeri*
206 produced methane from acetate and hydrogen during our simulations.


207 For comparison, the community simulation were also performed using models re-
208 constructed with CarveMe and ModelSEED (Figure 4). In both cases, most of the
209 above-mentioned known metabolic cross-feeding interactions and end-products were
210 not predicted, for instance the production of the short chain fatty acids butyrate
211 and propionate was missing. In summary, **gapseq** models were able to recapitulate
212 the major interactions, which are described for microbial communities in the hu-
213 man gut. The overall consumption pattern and individual microbial contributions

214 were found to be in agreement with literature data. Taken together, the community
215 simulation results illustrate the capacity of **gapseq** to construct predictive models
216 for complex metabolic interaction networks comprising several different species.



217 2.7 Pathway prediction of soil and gut microorganisms

218 To demonstrate the pathway prediction capabilities of **gapseq**, we analysed two
219 communities of soil and gut microorganisms comprising 922 and 822 organisms,
220 respectively. The two communities could be separated from each other by differ-
221 ences in energy metabolism (Principal component analysis, Figure 5A). Here, most
222 variance was explained by subsystems of pathways that are involved in chemoau-
223 totrophic, respiratory, and fermentative processes including hydrogen production.
224 Out of 128 energy pathways, the presence of 40 pathways differed significantly
225 (Kolmogorov-Smirnov test, $P < 0.05$) between soil and gut microorganisms and
226 could be categorised into 12 subsystems (Figure 5B). In total, gut microorganisms
227 showed less variety in energy pathways than soil microorganisms. Only pathways
228 relevant for the formation of acetate, hydrogen, and lactate were predicted to be
229 enriched. In the case of all other energy subsystems, more pathways were predicted
230 for soil organisms, most prominently pathways relevant for aerobic and anaerobic
231 respiration as well as the tricarboxylic acid cycle (TCA). In summary, members of
232 the soil community showed a more versatile energy metabolisms, which potentially
233 indicates a higher energetic specialisation of gut microbes. This sample application
234 demonstrates how **gapseq** can facilitate the characterisation and comparison of mi-
235 crobial communities based on the analysis of the presence and absence of specific
236 metabolic pathways.



2.8 Model reconstructions for metagenomic assemblies

Genome-scale metabolic models can also be reconstructed on the basis of species-level genome bins (SGBs, [69]) assembled from shotgun metagenomic sequencing reads. Yet, genome assemblies from metagenomic material are more prone to errors, fragmentation, and sequence gaps than assemblies of isolated genomes [89], which can potentially cause gaps in the metabolic network reconstructions. We tested whether **gapseq** is able to identify and fill such gaps by comparing the models reconstructed for 127 SGBs from the human microbiome [69] to corresponding models of closely-related reference genomes that were assembled from DNA-sequencing of pure cultures (Figure S2).

As expected, we found a strong positive correlation between the SGBs' genome completion and their model similarity to their respective reference models (Spearman's rank correlation, $n = 127$, $P < 10^{-9}$). To estimate the quantitative effect of genome completion on the model similarity, a logarithmic function ($y(x) = c + b * \log(x)$) was fitted to the data ($R^2 = 0.71$, Figure S2). The fitted model indicated, that **gapseq** is able to reconstruct the underlying metabolic network of an organism even on the basis of incomplete and fragmented genomes. For instance, **gapseq** was on average able to recover 90% of the enzymatic reactions that are found in the reference models for SGBs with a predicted genome completion of only 80% (Figure S2).

2.9 Summary of validation tests

For each validation approach, predictions were compared to experimental data obtained from databases and literature to calculate prediction performance scores. The overall accuracy (proportion all correct prediction in relation to all predictions made) of model predictions with experimental data was 66% (CarveMe), 70%




Figure 6: **Summary of different validation tests.** The specificity and sensitivity for all compared methods are shown. This includes results from benchmarks concerning enzyme activities, energy sources, fermentation products, gene essentiality, and metabolite production/consumption in an anaerobic food web.

(ModelSEED), and 81% (gapseq)(Table 1). Sensitivity measures the proportion of correctly predicted positives, whereas specificity accounts for the accurate prediction of negatives. All approaches showed a high specificity > 0.7 with highest values for fermentation product and gene essentiality tests. Notably, gapseq showed the highest sensitivity over all tests (Figure 6). In summary, gapseq outperformed other methods in terms of accuracy and sensitivity while showing similar specificity.

3 Discussion

Here, we introduced gapseq - a new tool for metabolic pathway analysis and genome-scale metabolic network reconstruction. The novelty of gapseq lies in the combination of (i) a novel reaction prediction that is based both on genomic sequence homology as well as pathway topology, (ii) a profound curation of the reaction and transporter database to prevent thermodynamically infeasible reaction cycles, and (iii) a reaction evidence score-oriented gap-filling algorithm. In order to scrutinise gapseq metabolic models, we compared the models' network structures and predictions with large-scale experimental data sets, which were retrieved from publicly available databases. Furthermore, the ability of gapseq to predict bacterial phenotypes was compared to two other commonly used automatic reconstruction methods, namely, CarveMe [39] and ModelSEED [24] (Table 1). ModelSEED is also implemented in the KBASE online software platform [90].

Table 1: Summary of different methods that were compared in this work. Accuracy, sensitivity, and specificity scores are based on 14,895 tested phenotypes including energy sources, enzyme activity, fermentation products, gene essentiality, and anaerobic food web structure predictions.

Metric	CarveMe	gapseq	ModelSEED
<i>Implementation</i>			
Infrastructure	local	local	web service
Input (FASTA file)	protein	nucleotide	nucleotide
Programming languages	python	shell script, R	perl/javascript
Gap-fill solver	CPLEX	GLPK/CPLEX	not needed*
Gap-fill problem formulation	MILP	LP	MILP
<i>Performance</i>			
Accuracy	0.66	0.81	0.70
Sensitivity	0.34	0.73	0.32
Specificity	0.84	0.83	0.88
Model file quality**	0.32 ± 0.006	0.78 ± 0.004	0.39 ± 0.016

* Solver runs on ModelSEED server. No local solver is required.

** MEMOTE total score (± SD).

Crucial large-scale benchmarking of metabolic models

The quality of genome-scale metabolic networks can be assessed by comparing model predictions with experimental physiological data. The protocol by Thiele and Pals-son (2010) for the reconstruction of genome-scale metabolic networks recommends the quality assessment and manual network curation using data for (i) known secretion products (e.g. fermentation end-products), (ii) single-gene deletion mutant growth phenotypes (i.e. gene essentiality), and (iii) the utilisation of carbon/energy sources [20]. Tools for the automatic reconstruction of metabolic networks should also make use of such physiological data whenever available for benchmarking. Here, we tested our **gapseq** approach on the basis of all three recommended phenotypic data and compared the performance with CarveMe and ModelSEED. Additionally, we included two novel benchmark tests: The comparison of model predictions with (iv) the activity of specific enzymes known from experimental studies [49] and (v) metabolic interactions (food web) among microorganisms in a multi-species community within an anaerobic environment. Across all five benchmark tests, we could show that **gapseq** outperformed CarveMe and ModelSEED in terms of sensitivity while achieving specificity scores that are comparable to the other two tools (Figure 6).

Publicly available genome sequences of microorganisms, which can be subject for automated metabolic network reconstruction are massively increasing in number due to continuing advances in high-quality and high-throughput sequencing technologies [18]. This development is further fueled by the the increasing number of genome assemblies from metagenomic material [91]. In contrast, standardised phenotypic data for microorganisms remains a bottleneck for the validation of automated metabolic network reconstruction pipelines such as **gapseq**. As consequence, it is crucial for the future development of automated network reconstruction software to include possibly all available phenotypic data for benchmarking, especially data from non-model organisms. To benchmark **gapseq** in relation to CarveMe and ModelSEED using phenotypic data from mainly non-model organisms, we retrieved phenotypic data of enzyme activity for more than 3,000 organisms and carbon source utilisation for more than 500 organisms from online databases, which is, to our knowledge, the

yet largest phenotypic data set used for validation of automatically reconstructed metabolic networks. In this validation approach **gapseq** achieved the highest prediction accuracy among all three tools tested (Figure 1).

Hence, those results suggest that **gapseq** is a powerful new tool for the automated reconstruction of genome-scale metabolic network models. Moreover, the underlying reference protein sequences as well as the pathway database can readily be updated using online resources, which makes **gapseq** flexible to include future developments and findings in microbial metabolic physiology.

Automated network reconstructions for community modelling

While single organisms can be considered as the building blocks of microbial communities, individual metabolic models of organisms are the building blocks of *in silico* microbial community simulations. Therefore, genome-scale metabolic models are increasingly applied to predict the function of multi-species microbial communities [61, 92, 93]. To correctly infer metabolic interaction networks between different organisms, it is important that individual models accurately predict nutrient utilisation (e.g. carbon source) and metabolic end-products (e.g. fermentation products). In this study, the benchmarks for carbon source utilisation and fermentation end-product identity indicated that **gapseq** has the highest prediction performance compared to other reconstruction tools (Figure 1 and Figure 3).

To illustrate the applicability of **gapseq**-reconstructed metabolic models for the simulation of multi-species community metabolism, we generated models for microbial strains from the human gut microbiota and simulated their growth in a shared environment. Without further curation, the community simulation reproduced all important hallmarks of intestinal anaerobic food webs [64, 86]. Above all, short chain fatty acids (SCFA) were predicted to be the primary end products of fermentation. This prediction is important to represent intestinal metabolism, because SCFA are crucially involved in host physiology by affecting regulatory response in intestinal and immune cells [94, 95]. Furthermore, the simulation accurately predicted the exchange of metabolites between different members of the microbial community (Figure 4). Cross-feeding of metabolites and the formation of anaerobic food chains have been associated with a healthy microbiome [9, 96]. For instance, the cross-feeding of lactate has been reported to be vital for the early establishment of a healthy gut microbiota in infants [96]. Accordingly we observed the exchange of lactate between different bacterial species in the community simulations (Figure 4) and involved known lactate producers (e.g. *Enterococcus faecalis*) and consumers (e.g. *Megasphaera elsdenii*). This example illustrates that we are able to predict key features of the anaerobic food-web within the gastrointestinal microbiota using **gapseq** models. In addition to the ability to accurately model metabolic processes within existing microbial communities, **gapseq** will further promote the potential of metabolic modelling to predict how complex microbial communities can be modulated by targeted interventions. Specific interventions, which could for instance be predicted, are the introduction of new species to the community (i.e. probiotics) or microbiome-modulating compounds (prebiotics) to the environment. Predictions of potential intervention strategies that target the microbiome are of vast relevance

for biomedical research. Furthermore, metabolic interactions between microbiome members are difficult to detect *in vivo* due to the simultaneous production and uptake of metabolites. Thus, *in silico* predictions of metabolite cross-feeding interactions are highly valuable for hypothesis generation about the function and dynamics of microbial communities.

Taken together, the results obtained with **gapseq** suggest, that metabolic models which are reconstructed using **gapseq** are promising starting points to construct ecosystem-scale models of inter-species biochemical processes and to predict targeted strategies to modulate microbiome structure and function.

Pathway analysis of microbial communities

The construction of genome-scale metabolic models is based on metabolic networks that are inferred from genomic sequences in the context of biochemical databases [20]. Although, the reconstruction of metabolic networks is closely related to the prediction of metabolic pathways, metabolic modelling and pathway analysis are often treated separately [97]. In **gapseq**, the prediction of metabolic pathways is intrinsically tied to the reconstruction of metabolic networks and gap-filling. In addition, reaction, transporter, and pathway predictions can also be used to evaluate the functional capacities of microorganisms without the need of metabolic modelling. As an example for metabolic pathway analysis, we compared the predicted energy metabolism of two large microbial communities that occur in soil and the human gut. We could show that the predicted distribution of pathways differ between both communities based on the habitat, which usually accommodates the members of the respective community. Gut microorganisms showed a less versatile energy metabolism and a specialisation towards fermentation pathways, which lead to the formation of acetate, hydrogen, and lactate. Variations in pathways distributions between both communities may be explained by distinct evolutionary histories. The habitat of the diverse group of soil microorganisms more likely represents an open ecosystem, whereas the gut microbiome is directly constraint by a multi-cellular host that potentially affect microbial phenotypic traits [98]. In general, metabolic modelling should be accompanied by the analysis of pathways based on statistical methods [97] to compensate for additional assumptions, which are introduced in constraint-based metabolic flux modelling [4].

Limitations and outlook

gapseq requires 1-2h for the reconstruction of a single model, whereas ModelSEED and CarveMe operate faster (10min) on a standard desktop computer. Nonetheless, CarveMe needs as input gene sequences (protein or nucleotide), which has to be predicted first, and ModelSEED works as a web service, which can complicate the handling of large-scale reconstruction projects. In **gapseq**, pathways were predicted based on topology and sequence homology searches. However, the assignment of enzymatic function from sequence comparisons has been shown to potentially miss protein domain structures and thus can cause false annotations [99, 100]. In addition, **gapseq** uses many resources to find potential sequences for reactions in pathway databases. Together this might explain why although **gapseq** performed better than other methods on predicting positive phenotypes (function present),

it went head to head with regard to negative phenotype predictions (function not present). CarveMe takes a different approach when inferring function by taking care of functional regions (protein domains) [101], resulting in orthologous groups [102], which results in a slightly better specificity (true negative phenotype predictions) in benchmarks (Figure 6). Future developments of **gapseq** will address orthologous groups by using multiple inference methods. Furthermore, the integration of functional predictions coming from phylogenetic inference without the need of genomic sequences [103] might also be promising for further developments of **gapseq**.

Conclusion

We provide a new software tool called **gapseq** that is suitable for metabolic network analysis and metabolic model reconstruction. To enhance phenotype predictions, **gapseq** employs various data sources and a novel gap-filling procedure that reduces the impact of arbitrary growth medium requirements. We further brought together the so far largest benchmarking of genome-scale metabolic models, in which **gapseq** outperformed comparable alternative tools. With the increased model quality of automated network reconstructions, **gapseq** will provide new insights into the metabolic phenotypes of non-model and yet-uncultured bacteria whose genomes are assembled from metagenomic material. In this way, the models and their simulations allow predictions on the organisms' ecological role in their natural environments. Taken together, we consider **gapseq** as important contribution to the modelling of microbial communities in the age of the microbiome.


4 Methods

4.1 Program overview & source code availability

The source code is accessible and maintained at <https://github.com/jotech/gapseq>. The program is called by `./gapseq`, which is a wrapper script for the main modules. Important program calls are `./gapseq find` (pathway and reaction finder), `./gapseq find-transport` (transporter detection), `./gapseq draft` (draft model creation), `./gapseq fill` (gap-filling), or `./gapseq doall` to perform all in line. When ever necessary, method sections directly refer to config, data and source code files from the **gapseq** package, which contains the main sub-directory `src/` with source code files and `dat/`, which contains databases and also the sequence files in `dat/seq/`. Figure 7 shows an overview of the different **gapseq** modules.

4.2 Pathway and sequence databases

Pathways are considered as a list of reactions with enzyme names and EC numbers. Pathway definition were obtained from MetaCyc [27], KEGG [28], and ModelSEED [24]. For MetaCyc, PathwayTools [29] was used in combination with Python-Cyc to obtain pathway definitions [30] (`src/meta2pwy.py`). Information on Kegg pathways were retrieved directly from the KEGG homepage: reactions (<http://rest.kegg.jp/list/reaction>), and EC numbers (<http://rest.kegg.jp/link/pathway/ec>) and further processed (`src/kegg_pwy.R`). In case of ModelSEED, subsystem definition were obtained from the homepage: <http://modelseed.org/genomes/Annotations> (`src/seed_pwy.R`). In addition, manual defined and revised



pathways are stored in the file `dat/custom_pwy.tbl`.
 Sequence data needed for pathway prediction were downloaded from UniProt [31] for each reaction identified by EC number, enzyme name, or gene name. Both reviewed and unreviewed sequences are considered and stored as clustered UniPac sequences (`src/uniprot.sh`). To increase the sequence pool for a given reaction, alternative EC numbers from BRENDA [32] and from the Enzyme Nomenclature Committee <https://www.qmul.ac.uk/sbcs/iubmb/enzyme/> are integrated (`src/altec.R`, `dat/brenda_ec.csv`).

4.3 Pathway prediction

For each pathway selected from a pathway database (MetaCyc, KEGG, ModelSEED, custom), `gapseq` searches for sequence evidence and a pathway is defined as present if enough of its reactions were found to have sequence evidence. In more detail, sequence data (section 4.2) is used for homology search by *tblastn* [33] with the protein sequence as query and the genome as database. By default, a bitscore ≥ 200 and a coverage of at least 75% is needed for a match. For certain reactions, the user can define additional criteria, for example an identity of $\geq 75\%$ (`dat/exception.tbl`). In case of protein complexes with subunits, a more complex procedure is followed (section 4.4). Spontaneous reactions, which do not need an enzyme, were set to be present in any case. In general, a pathway is considered to be present if at least 80% of the reactions are found (`completenessCutoffNoHints` threshold). This pathway completeness threshold is lowered for pathways in following cases:

- 1 If the pathway contains key reactions, as it is defined for some pathways in MetaCyc, and all key reactions are found, then `completenessCutoff` of the total reactions needed to be found. We used a value of 2/3 for this threshold.
- 2 In cases in which no sequence data is available for specific reactions, the status of the reactions is set to "vague" and these reactions do not count as missing

if they account for less than `vagueCutoff` of the total reactions of a pathway.
We used a value of 1/3 for this threshold.
The pathway prediction algorithm is implemented in the bash shell script
`src/gapseq.find.sh`, which uses GNU parallel [34] and fastaindex/fastafetch from
exonerate [35].

4.4 Protein complex prediction

A problem with automatic sequence download for reactions (as FASTA files) comes
with protein complexes, for which a single blast hit may be not sufficient to predict
enzyme presence. In `gapseq`, subunits are detected by text matching in the FASTA
headers. Search terms are: "subunit", "chain", "polypeptide", "component", and
different numbering systems (roman, arabic, greek) are homogenised. To avoid ar-
tifacts in text matching, subunits that occur less than five times in the sequence
file are not considered, and in cases in which a subunit occurs almost exclusively
($\geq 66\%$) the other entries are not taken into account. All FASTA entries, which
could not be matched by text mining, or which are excluded because of the coverage,
are labeled 'undefined subunit' and do not add to the total amount of subunits.
For each recognised subunit, a blast search is done. A protein complex counts as
present if more than 50% of the subunits could be found, whereby the presence of
'undefined subunits' tip the balance if exactly 50% of the subunits were found. The
text matching with regular expressions is done with R's `stringr` [36] and `biostrings`
[37] as defined in `src/complex_detection.R`. The script is called from within the
shell script `src/gapseq.find.sh`.

4.5 Transporter prediction

For transporter search, sequence data from the Transporter Classification Database
is employed [38]. In addition, manual defined sequences can be defined in
`dat/seq/transporter.fasta`. The sequence set is reduced to a subset of trans-
porters that involve metabolites known to be produced or consumed by microor-
ganisms (`dat/sub2pwy.csv`). Subsequently, the genome is queried by the reduced
sequences using `tblastn` [33]. For each hit (default cutoffs: bitscore ≥ 200 and cov-
erage $\geq 75\%$), the transporter type (1. Channels and pores, 2. Electrochemical
potential-driven transporter, 3. Primary active transporters, 4. Group transloca-
tors) is determined using the TC number mentioned in the FASTA header. A
suitable candidate reaction is searched in the reaction database. If there is a hit for
a transporter of a substance but no candidate reaction for the respective transporter
type can be found, then other transporter types are considered. The transporter
search is done by the shell script `src/transporter.sh` that uses GNU parallel [34]
and fastaindex/fastafetch from exonerate [35].
Candidate transporters are selected from the reaction database by transporter type
and substance name. This is done by text search and is currently implemented
only for the ModelSEED namespace. From the ModelSEED reaction database
all reaction with the flag `is_transport = 1` are taken and the transporter type
is predicted by keywords: "channel", "pore" (1. Channels and pores); "uniport",
"symport", "antiport", "permease", "gradient" (2. Electrochemical potential-driven
transporters); "ABC", "ATPase", "ATP" (3. Primary active transporters); "PTS"

(4. Group translocators). If no transporter type could be identified by keywords, additional string matching is done for ATPases, proton/sodium antiporter, and PTS by considering the stoichiometry of the involved metabolites. The transported substance is identified as the substance that occurs on both sides of the reaction. In addition, reactions from the reaction database can be linked manually to substances and transporter types (`dat/seed_transporter_custom.tbl`). The text matching with regular expressions is done with `stringr` [36] (`src/seed_transporter.R`).

4.6 Biochemistry database curation and construction of universal metabolic model

For the construction of genome-scale metabolic network models, `gapseq` uses a reactions and metabolite database that is derived from the ModelSEED database [24] as from January 2018. In addition, 30 new reactions and 2 new metabolites were introduced to the `gapseq` biochemistry database (see suppl. table S1). All reactions and metabolites from the database were included for the construction of a full universal metabolic network model; an approach that is also used in CarveMe [39]. We curated the underlying biochemistry database in order to correct inconsistencies in reaction stoichiometries and reversibilities. Inconsistencies were identified by optimising the universal network model for ATP-production without any nutritional input to the model using flux balance analysis. In case of ATP-production, the flux distributions of such thermodynamically infeasible reaction cycles were investigated by cross-checking the involved reactions with literature information, the BRENDA database for enzymes [32], and the MetaCyc database [27]. Stoichiometries and reversibilities of erroneous reactions were corrected accordingly. This curation procedure was repeated until no thermodynamically infeasible and ATP-generating reaction cycles were observed.

Hits from the pathway prediction (4.3) and transporter prediction (4.5) are mapped to the `gapseq` reaction database using different common identifiers. A majority of reactions are directly matched via their corresponding Enzyme Commission (EC) system identifier [40] and Transporter Classification (TC) system identifier [38], respectively. For this mapping, also alternative EC-numbers for enzymatic reactions as defined in the BRENDA database [32] are considered. Moreover, the databases used for pathway and transporter predictions often provide cross-links to the reaction's KEGG ID, which is also assigned to most reactions in the `gapseq` database and used to match reactions. Additionally, the MNXref database [41] provides cross links between several biochemistry databases, which `gapseq` also utilises to translate hits from the pathway predictions to model reactions. Finally, a manual translation of enzyme names to model reactions is done for some reactions, which we identified as important reactions but which failed to match between the pathway databases (4.3) and the `gapseq` model reactions using other reaction identifiers (`dat/seed_Enzyme_Name_Reactions_Aliases.tsv`). The overall mapping is done by the function `getDBhit()` as defined in `./src/gapseq_find.sh`.

4.7 Model draft generation

A draft genome-scale metabolic model is constructed based on the results from the pathway and transporter predictions (see above). A reaction is added to the draft model if the corresponding enzyme/transporter was directly found or if the pathway

was predicted to be present (i.e. due to pathway completeness and key enzymes) in which the reaction participates. Additionally, spontaneous reactions as defined in the MetaCyc database as well as transport reaction of compounds, which are known to be able to cross cell membranes by means of diffusion (e.g. H_2), are directly added to every draft model. As part of the draft model construction **gapseq** adds a biomass reaction to the network that aims to describe the composition of molecular constituents that the organism needs to produce in order to form 1 g dry weight (1 gDW) of bacterial biomass. **gapseq** uses the biomass composition definition from the ModelSEED database for Gram-positive (`dat/seed_biomass.DT_gramPos.tsv`) and Gram-negative bacteria (`dat/seed_biomass.DT_gramNeg.tsv`). If no Gram-staining property is specified by the user, **gapseq** predicts the Gram-staining-dependent biomass reactions by finding the closest 16S-rRNA-gene neighbor using a **blastn** search against reference 16S-rRNA gene sequences from 4647 bacterial species with known Gram-staining properties that are obtained from the PROTRAITS database [42]. The model draft generation is done by the R script `src/generate_GSdraft.R`.

4.8 Gap-filling algorithm

gapseq provides a gap-filling algorithm that adds reactions to the model in order to enable biomass production (i.e. growth) and likely anabolic and catabolic capabilities. The algorithm uses the alignment statistics (i.e. the bitscore) from the pathway- and transporter prediction steps of **gapseq** (see above) to preferentially add reactions to the network, which have the highest genetic evidence. This approach is especially relevant in cases where the sequence similarity to known enzyme-coding reference genes was close to but did not reach the cutoff value b , which is required for a reaction to be included directly into the draft network. In contrast to the gap-filling algorithms described in previous works [43] and [39], which also use genetic evidence-weighted gap-filling, the gap-filling problem in **gapseq** is not formulated as Mixed Integer Linear Program (MILP) but as Linear Program (LP), and is derived from the parsimonious enzyme usage Flux Balance Analysis (pFBA) algorithm developed by Lewis *et al.*, 2010 [3]. Therefore, the alignment statistics (i.e. bitscore) are translated into weights for the corresponding model reactions and incorporated into the problem formulation:

$$\max: v_j - c \sum_{i \in R_{all}} w_i |v_i|, \quad (1)$$

$$w_i = \begin{cases} w_{min} & b_i \geq u \quad | \quad i \in R_{draft} \\ (b_i - u) \left(\frac{w_{min} - w_{max}}{u - l} \right) + w_{min} & l \leq b_i < u \\ w_{max} & b_i < l \end{cases}$$

s.t.

$$S \cdot v = 0$$

$$lb \leq v \leq ub$$

Where R_{all} is the set of all reaction in the universal model, R_{draft} are the reactions, which are already part of the draft network before gap-filling, v_j is the flux through the objective reactions (e.g. biomass production), v_i the flux through reaction i , w_i the weight for reaction i , v the flux vector for all reactions, and c a scalar factor that determines the contribution of the absolute reduction of weighted fluxes to the overall FBA solution (default: $c = 0.001$). Moreover, a maximum weight value w_{max} (default: 100) is assigned if the reaction's highest bitscore is smaller than a threshold l (default: 50). A minimum reaction weight w_{min} (default: 0.005) is assigned to reactions with a bitscore higher than u (default: 200) or if the reactions are already part of the draft model. S is the stoichiometric matrix and lb and ub the lower and upper flux bound vectors.

Two other LP-based gap-filling algorithms that incorporate reaction evidence scores have been formulated by Dreyfuss *et al.* (2013) [44] and Medlock *et al.* (2020) [45], respectively. These approaches require a definition of a minimum flux through the biomass reaction to ensure growth. The pFBA-derived LP formulation of **gapseq** (equation 1) includes the flux through the biomass/objective reaction v_j together with the reaction evidence scores in a single objective function.

In **gapseq** and following the solution of the LP (1), reactions carrying a flux and which are not part of the draft model are added to the network model. The algorithm is implemented in **src/gapfill4.R**.

4.9 Gap-filling of biomass, carbon sources, and fermentation products

Gap-filling of a draft model in **gapseq** requires only for the first step a user-defined growth medium that is ideally known to support growth of the organism of interest *in vivo*. If no growth medium is specified by the user, a complete medium (ALLmed) is chosen by **gapseq** (as done for the large-scale benchmarks of enzyme activity and carbon sources, cf. 4.11, 4.12). A set of common microbial growth media (e.g. LB, TSB, M9) is provided in the **gapseq** software directory **dat/medium/**. In addition, the user can provide a custom growth medium definition. The above described gap-filling algorithm is used to improve the generated draft model in four steps.

- 1 **Biomass production:** To ensure that the model is able to produce biomass under the given nutritional input (medium) the gap-filling algorithm is applied while the objective is defined as the flux through the biomass reaction. This step will add all missing reactions that are essential for *in silico* growth.
- 2 **Individual biomass components:** It is checked whether the model supports the biosynthesis of biomass components. Therefore, model is re-constrained to a M9-like minimal medium with a carbon source for which an exchange reactions is found (default: glucose if available). The objective function is set to the production of one biomass component at a time and the gap-fill algorithm is performed. This gap-filling step is repeated for each biomass component metabolite twice, with and without oxygen to potentially allow aerobic and anaerobic growth for facultative anaerobe species.
- 3 **Alternative energy sources:** **gapseq** attempts to gap-fill likely metabolic pathways, which enable the utilisation of alternative energy sources, which might not be part of the defined growth medium from step (1). To this end, the model is re-constrained to a M9-like minimal medium containing a single

carbon source of interest at the time. As objective function, the summed flux of artificial reactions that accept electrons from the electron carriers ubiquinol, menaquinol, or NADH is defined. This test can be considered as an *in silico* simulation of the commonly used BIOLOG carbon source utilisation test arrays [46] in which the colometric effect is coupled to a dehydrogenase [47]. This gap-filling step is performed for all metabolites defined in `dat/sub2pwy.csv`.

4 **Metabolic products:** Finally, the same list of compounds as for step (3), is used to check whether the network can be gap-filled to allow the formation of these metabolites given the original medium. For each compound the gap-filling algorithm is applied with the production of the focal compound as objective function.

While step (1) considers all reaction from the universal model as potential candidate reactions for gap-filling, steps (2-4) allow only the addition of candidate reactions to the model with a corresponding bitscore from the pathway prediction (4.3) higher than a threshold value b (default: 50). Thus, these so-termed '*core reactions*' represent only reactions, for which `gapseq` has found genomic sequence or pathway evidence. This approach for steps (2-4) is chosen to avoid the addition of biosynthetic capabilities to the model, which the organism presumably does not possess.

4.10 Formal and functional model file testing

The validity of genome-scale metabolic model files was checked with MEMOTE (0.10.2) [48]. For all models used in the anaerobic food web (4.16), the total MEMOTE score was computed for the respective SBML-Model files. MEMOTE was executed using the parameter `--skip test_find_metabolites_not_produced_with_open_bounds` and `--skip test_find_metabolites_not_consumed_with_open_bounds` since these tests do not contribute to the total MEMOTE score but require long computation time.

4.11 Validation with enzymatic data (BacDive)

The Bacterial Diversity Metadatabase (BacDive) [49] was used to obtain enzymatic activity data. For this purpose, a list of type strains IDs were downloaded using the advanced search. Afterwards the IDs were used to query the database via the R package BacDiveR (0.9.1) to obtain the data [50]. If the stored data contained non-zero entries for enzymatic activity and if a genome assembly was available on NCBI, the type strain was considered for the validation analysis. The respective genome assemblies were downloaded with `ncbi-genome-download` (<https://github.com/kbclin/ncbi-genome-download>). If multiple genomes were available for one type strain, '*representative*' and '*complete*' (NCBI tags) genomes were preferred and, in case there were still multiple candidate genomes available, the most complete genome was selected. Genome completeness was estimated by employing the software BUSCO (3.0.2) [51]. In total, 3017 type strain genomes were taken as input for ModelSEED (2.5.1), CarveMe (1.2.2), and `gapseq` to create metabolic models. The gap-filling parameters were set to default values for each program, i.e. a complete medium was assumed. The final test whether a reaction activity is covered by a model was done by checking if the corresponding reaction

is present in the model. This was done by matching enzymes and reactions via EC numbers. For CarveMe the vmh (<https://www.vmh.life>) and for ModelSEED and gapseq the ModelSEED (<http://modelseed.org>) reaction database was used to match reactions and EC numbers. For the EC numbers 3.1.3.1, 3.1.3.2, the corresponding reactions were the same, and thus unspecific, so that both EC numbers were not considered for the validation analysis. In general, the enzyme activities in the BacDive database have the form active ("+") or not active ("-") but some entries were ambiguous (e.g.: "+/-"). The ambiguous entries were omitted from the analysis.

4.12 Validation with carbon sources data (ProTraits)

Data for the validation of carbon source utilisation was obtained from the "atlas of prokaryotic traits" database (ProTraits) [42]. A tab-separated table with binarised predictions with a stringent threshold of precision of ≥ 0.95 were downloaded from <http://protraits.irb.hr/data.html>. For organisms which had at least one carbon source prediction, the corresponding genome was obtained from NCBI RefSeq [52] if available. In cases where a genome assembly was found, it was taken as input for ModelSEED, CarveMe, and gapseq to create metabolic models. The number of potential carbon sources was reduced to a subset for which a mapping from substance name to ModelSEED and CarveMe model namespace existed (dat/sub2pwy.csv). The tests for D-lyxose were removed because it was listed as all negative in ProTraits and also all compared pipelines predicted no utilisation. The main test whether a carbon source can be used by a model was done in a BIOLOG-like manner as described above (see 4.9). To this end, temporary reactions to recycle reduced electron carriers as carbon source utilisation indicators were added to the respective model. The objective for optimisation was set to maximise the flux through these recycling reactions. The exchange reactions were limited to a minimal medium with minerals and the focal potential carbon source. This theoretical approach tested, whether the model is able to pass electrons from the potential carbon source to electron carrier metabolites. A carbon source was predicted to be able to serve as energy source if the recycle reactions carried a positive flux.

4.13 Prediction of gene essentiality

To predict the essentiality of genes we performed *in silico* single gene deletion phenotype analysis for the network reconstructions of *Escherichia coli* str. K-12 substr. MG1655 (RefSeq assembly accession: GCF_000005845.2), *Bacillus subtilis* substr. *subtilis* str. 168 (GCF_000789275.1), *Shewanella oneidensis* MR-1 (GCF_000146165.2), *Pseudomonas aeruginosa* PAO1 (GCF_000006765.1), and *Mycoplasma genitalium* G37 (GCF_000027325.1). The analysis was performed on the basis of the models' Gene-Protein-Reaction (GPR) mappings and according to the protocol by Thiele and Palsson, 2010 [20]. To this end, the contingency tables of predicted growth/no growth phenotypes from the network models and experimentally determined growth phenotypes of gene deletion mutants were constructed. Genes were predicted to be conditionally essential under the given growth environment if the predicted growth rates of the models were below 0.01 hr^{-1} . The growth media compositions for growth predictions were defined as M9 with glucose as carbon-

and energy source for *E. coli*, lysogeny broth (LB) for *B. subtilis* and *S. oneidensis*, M9 with succinate as carbon and energy-source for *P. aeruginosa*, and a complete medium (all external metabolites available for uptake) for *M. genitalium*. Experimental data for gene essentiality was obtained from [53, 54, 55, 56, 57].

4.14 Fermentation product tests

The release of by-products from anaerobic metabolism was predicted using Flux Balance Analysis (FBA) coupled with a minimisation of total flux [58] to avoid fluxes that do not contribute to the objective function of the biomass production. In addition, Flux-Variability-Analysis (FVA) [59] was applied to predict the maximum fermentation product release of individual metabolites across all possible FBA solutions. Metabolites with a positive exchange flux (i.e. outflow) were considered as fermentation products. The analysis was performed for 18 different bacterial organisms, which (1) have a genome assembly available in the RefSeq database [52], (2) are known to grow in anaerobic environments, and (3) for which the fermentation products have been described in the literature based on anaerobic cultivation experiments (suppl. table S2). The gap-filling of the network models using gapseq, CarveMe, and ModelSEED as well as the simulations of anaerobic growth were all performed assuming the same growth medium that comprised several organic compounds (i.e. carbohydrates, polyols, nucleotides, amino acids, organic acids) as potential energy sources and nutrients for growth (see media file `dat/media/FT.csv` at the gapseq github repository).

Since the amount of fermentation product release depends on the organism's growth rate, we normalised the outflow of the individual fermentation products, which has the unit $mmol * gDW^{-1} * hr^{-1}$, by the predicted growth rate of the respective organism which has the unit hr^{-1} . Thus, we report the amount of fermentation product production in the quantity of the metabolite that is produced per unit of biomass: $mmol * gDW^{-1}$.

4.15 Pathway prediction of soil and gut microorganisms

The pathway analysis was done by comparing predicted pathways of soil and gut microorganisms. For this means, genomes were downloaded from a resource of reference soil organisms [60] and gut microbes [61]. The default parameter of gapseq were used for pathway prediction. The principal component analysis was done in R using the factoextra package [62]. For predicted pathways for soil and gut microorganisms, it was checked if samples belong to different distributions using a bootstrap version of the Kolmogorov-Smirnov test [63].

4.16 Anaerobic food web of the human gut microbiome

Representative bacterial organisms known to be relevant in the human intestinal cross-feeding of metabolites were selected based on the proposed food webs by Louis *et al.*, 2014 [64] and Rivera-Chavez *et al.*, 2015 [65]. The genomes of organisms were obtained from NCBI RefSeq [52] and metabolic models reconstructed using gapseq, carveme, and modelseed. A medium containing minerals, vitamins, amino acids, fermentation- and metabolic by-products (namely acetate, formate, lactate, butyrate, propionate, H_2 , CH_4 , ethanol, H_2S , succinate), and carbohydrates (glucose, fructose, arabinose, ribose, fucose, rhamnose, lactose) was used for gap-filling.

Furthermore, a published model of *Methanosarcina barkeri* was added to the community [66] to represent archaea that are also known to be part of anaerobic food webs [67]. All organisms of the modeled community and their respective genome assembly accession numbers are listed in supplementary table S3. All metabolic models were then simulated with BacArena [68] by using the described medium but without the fermentation and by-products, plus sulfite and 4-aminobenzoate which were needed for growth by the *M. barkeri* model. The community was simulated for five time steps (corresponding to 5 hours simulated time). The analysis of metabolite uptake and production were done after the third time step, for which all organisms were still growing exponentially.

4.17 Model reconstructions from metagenomic assemblies

4,930 species-level genome bins (SGBs) assembled from shotgun metagenome sequencing reads were obtained from the study of Pasolli *et al.*, 2019 [69]. Only those SGBs were considered for further analysis, which were already classified as bacteria on a species-level in the original publication by Pasolli *et al.* For each SGB, closely related reference assemblies from the RefSeq database [52] were identified by constructing a multi-locus phylogenetic tree using autoMLST (version as of April 7th 2020, [70]). RefSeq assemblies were considered as genomes from the same species-level taxonomic group as the focal SGB if their predicted MASH distance (D) [71] were below or equal to 0.05. This threshold was shown before to cluster bacterial genomes at the taxonomic level of species [71]. Only SGBs with 10 or more assigned reference assemblies were considered for further analysis, which yielded in total 127 SGBs. Metabolic models were reconstructed using **gapseq** for each SGB and their 10 closest reference assemblies (Suppl. Table S5).

Next, similarity of SGB models with their respective reference models was calculated using the following metabolic network similarity score T_{SGB} :

$$T_{SGB} = \frac{\sum_i a_i b_i}{\sum_i b_i}, \quad i \in R_{SGB_Ref}, \quad 0 \leq b_i \leq 1$$

(2)

with

$$a_i = \begin{cases} 0 & \text{if } i \notin R_{SGB} \\ 1 & \text{if } i \in R_{SGB} \end{cases}$$

R_{SGB_Ref} is the union set of reactions with associated genes that are part of the network models reconstructed for the ten reference genome assemblies of the focal SGB. R_{SGB} is the set of reactions part of the SGB's model reconstruction. b_i is the frequency of reaction i among the ten SGB's reference models.

Completion of the genome sequence of SGBs was estimated by using BUSCO (version 4.0.6, [51]) using the specific completion score.

4.18 Technical details

The pathway prediction part of **gapseq** is implemented as Bash shell script and the metabolic model generation part is written in R. Linear optimisation can be

performed with a different solvers (GLPK or CPLEX). Other requirements are
exonerate, bedtools, and barrnap. In addition, the following R packages are needed:
data.table [72], stringr [73], sybil [74], getopt [75], reshape2 [76], doParallel [77],
foreach [78], R.utils [79], stringi [80], glpkAPI [81], and BioStrings [82]. Models can
be exported as SBML [83] file using sybilSBML [74] or R data format (RDS) for
further analysis in R, for example with sybil [74] or BacArena [68].

Competing interests

The authors declare that they have no competing interests.

Author's contributions

JZ, CK, and SW conceptualized gapseq. JZ and SW developed the software and did the analysis. JZ, CK, and SW wrote the manuscript.

Acknowledgements

We thank Martin Sperfeld for fruitful comments and discussions during the developmental phase. The software was thankfully tested by Georgios Marinos, Shan Zhang, and Lena Best.

Availability of data and materials

gapseq is implemented in R and python and is freely available under the GNU General Public License (v3.0) on GitHub (<https://github.com/jotech/gapseq/>). All results presented in this manuscript were produced using the specific gapseq version 1.0 as archived on GitHub. The datasets used for model construction and validation purposes were obtained from publicly available databases and publications as cited at the respective parts of the manuscript.

Funding

CK and SW acknowledges support by the Collaborative Research Centre 1182 - "Origin and Function of Metaorganisms" - Deutsche Forschungsgemeinschaft and by the Cluster of Excellence 2167 - "Precision medicine in chronic inflammation" - Deutsche Forschungsgemeinschaft. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author details

¹Christian-Albrechts-University Kiel, Institute of Experimental Medicine, Research Group Medical Systems Biology, Michaelis-Str. 5, 24105 Kiel, Germany. ²Christian-Albrechts-University Kiel, Institute of Human Nutrition and Food Science, Nutriinformatics, Heinrich-Hecht-Platz 10, 24118 Kiel, Germany.

References

- Fell, D.A.: Systems properties of metabolic networks. In: Bar-Yam, Y. (ed.) Unifying Themes In Complex Systems, Volume 1, pp. 163–178. CRC Press, ??? (2003)
- Steuer, R.: Computational approaches to the topology, stability and dynamics of metabolic networks. *Phytochemistry* **68**(16), 2139–2151 (2007). doi:[10.1016/j.phytochem.2007.04.041](https://doi.org/10.1016/j.phytochem.2007.04.041). Dynamic Metabolic Networks
- Lewis, N.E., Hixson, K.K., Conrad, T.M., Lerman, J.A., Charusanti, P., Polpitiya, A.D., Adkins, J.N., Schramm, G., Purvine, S.O., Lopez-Ferrer, D., Weitz, K.K., Eils, R., König, R., Smith, R.D., Palsson, B.O.: Omic data from evolved e. coli are consistent with computed optimal growth from genome-scale models. *Molecular Systems Biology* **6**(1), 390 (2010). doi:[10.1038/msb.2010.47](https://doi.org/10.1038/msb.2010.47). <https://www.embopress.org/doi/pdf/10.1038/msb.2010.47>
- de Jong, H., Casagrande, S., Giordano, N., Cinquemani, E., Ropers, D., Geiselmann, J., Gouzé, J.-L.: Mathematical modeling of microbes: Metabolism, gene expression, and growth. *Journal of the Royal Society Interface* **14**(20170502) (2017). doi:[10.1098/rsif.2017.0502](https://doi.org/10.1098/rsif.2017.0502)
- Varma, A., Palsson, B.O.: Metabolic capabilities of escherichia coli ii. optimal growth patterns. *Journal of Theoretical Biology* **165**(4), 503–522 (1993). doi:[10.1006/jtbi.1993.1203](https://doi.org/10.1006/jtbi.1993.1203)
- Stolyar, S., Van Dien, S., Hillesland, K.L., Pinel, N., Lie, T.J., Leigh, J.A., Stahl, D.A.: Metabolic modeling of a mutualistic microbial community. *Molecular Systems Biology* **3**(1), 92 (2007). doi:[10.1038/msb4100131](https://doi.org/10.1038/msb4100131). <https://www.embopress.org/doi/pdf/10.1038/msb4100131>
- Zomorodi, A.R., Islam, M.M., Maranas, C.D.: d-optcom: Dynamic multi-level and multi-objective metabolic modeling of microbial communities. *ACS Synthetic Biology* **3**(4), 247–257 (2014). doi:[10.1021/sb4001307](https://doi.org/10.1021/sb4001307). PMID: 24742179. <https://doi.org/10.1021/sb4001307>
- Harcombe, W., Riehl, W., Dukovski, I., Granger, B., Betts, A., Lang, A., Bonilla, G., Kar, A., Leiby, N., Mehta, P., Marx, C., Segrè, D.: Metabolic resource allocation in individual microbes determines ecosystem interactions and spatial dynamics. *Cell Reports* **7**(4), 1104–1115 (2014). doi:[10.1016/j.celrep.2014.03.070](https://doi.org/10.1016/j.celrep.2014.03.070)
- Aden, K., Rehman, A., Waschina, S., Pan, W.-H., Walker, A., Lucio, M., Nunez, A.M., Bharti, R., Zimmermann, J., Bethge, J., Schulte, B., Schulte, D., Franke, A., Nikolaus, S., Schroeder, J.O., Vandeputte, D., Raes, J., Szymczak, S., Waetzig, G.H., Zeuner, R., Schmitt-Kopplin, P., Kaleta, C., Schreiber, S., Rosenstiel, P.: Metabolic functions of gut microbes associate with efficacy of tumor necrosis factor antagonists in patients with inflammatory bowel diseases. *Gastroenterology* (2019). doi:[10.1053/j.gastro.2019.07.025](https://doi.org/10.1053/j.gastro.2019.07.025)
- Koch, S., Kohrs, F., Lahmann, P., Bissinger, T., Wendschuh, S., Benndorf, D., Reichl, U., Klamt, S.: Redcom: A strategy for reduced metabolic modeling of complex microbial communities and its application for analyzing experimental datasets from anaerobic digestion. *PLOS Computational Biology* **15**(2), 1–32 (2019). doi:[10.1371/journal.pcbi.1006759](https://doi.org/10.1371/journal.pcbi.1006759)

11. Heinken, A., Thiele, I.: Systematic prediction of health-relevant human-microbial co-metabolism through a computational framework. *Gut Microbes* **6**(2), 120–130 (2015). doi:[10.1080/19490976.2015.1023494](https://doi.org/10.1080/19490976.2015.1023494). PMID: 25901891. <https://doi.org/10.1080/19490976.2015.1023494>
12. Pryor, R., Norvaisas, P., Marinos, G., Best, L., Thingholm, L.B., Quintaneiro, L.M., Haes, W.D., Esser, D., Waschina, S., Lujan, C., Smith, R.L., Scott, T.A., Martinez-Martinez, D., Woodward, O., Bryson, K., Laudes, M., Lieb, W., Houtkooper, R.H., Franke, A., Temmerman, L., Bjedov, I., Cochemé, H.M., Kaleta, C., Cabreiro, F.: Host-microbe-drug-nutrient screen identifies bacterial effectors of metformin therapy. *Cell* **178**(6), 1299–131229 (2019). doi:[10.1016/j.cell.2019.08.003](https://doi.org/10.1016/j.cell.2019.08.003)
13. Zimmermann, J., Obeng, N., Yang, W., Pees, B., Petersen, C., Waschina, S., Kisson, K.A., Aidley, J., Hoeppner, M.P., Bunk, B., Spröer, C., Leippe, M., Dierking, K., Kaleta, C., Schulenburg, H.: The functional repertoire contained within the native microbiota of the model nematode *Caenorhabditis elegans*. *The ISME Journal* **14**(1), 26–38 (2019). doi:[10.1038/s41396-019-0504-y](https://doi.org/10.1038/s41396-019-0504-y)
14. Oberhardt, M.A., Yizhak, K., Rupp, E.: Metabolically re-modeling the drug pipeline. *Current Opinion in Pharmacology* **13**(5), 778–785 (2013). doi:[10.1016/j.coph.2013.05.006](https://doi.org/10.1016/j.coph.2013.05.006). Anti-infectives • New technologies
15. Trawick, J.D., Schilling, C.H.: Use of constraint-based modeling for the prediction and validation of antimicrobial targets. *Biochemical Pharmacology* **71**(7), 1026–1035 (2006). doi:[10.1016/j.bcp.2005.10.049](https://doi.org/10.1016/j.bcp.2005.10.049). Special Issue on Antibacterials
16. Rau, M.H., Zeidan, A.A.: Constraint-based modeling in microbial food biotechnology. *Biochem. Soc. Trans.* **46**, 249–260 (2018). doi:[10.1042/BST20170268](https://doi.org/10.1042/BST20170268)
17. Park, J.H., Lee, S.Y.: Towards systems metabolic engineering of microorganisms for amino acid production. *Current Opinion in Biotechnology* **19**(5), 454–460 (2008). doi:[10.1016/j.copbio.2008.08.007](https://doi.org/10.1016/j.copbio.2008.08.007). Tissue, cell and pathway engineering
18. Loman, N.J., Pallen, M.J.: Twenty years of bacterial genome sequencing. *Nature Reviews Microbiology* **13**(12), 787–794 (2015). doi:[10.1038/nrmicro3565](https://doi.org/10.1038/nrmicro3565)
19. Wittig, U., De Beuckelaer, A.: Analysis and comparison of metabolic pathway databases. *Briefings in bioinformatics* **2**(2), 126–142 (2001)
20. Thiele, I., Palsson, B.O.: A protocol for generating a high-quality genome-scale metabolic reconstruction. *Nat Protoc* **5**(1), 93–121 (2010). doi:[10.1038/nprot.2009.203](https://doi.org/10.1038/nprot.2009.203)
21. Blaby-Haas, C.E., de Crécy-Lagard, V.: Mining high-throughput experimental data to link gene and function. *Trends in Biotechnology* **29**(4), 174–182 (2011). doi:[10.1016/j.tibtech.2011.01.001](https://doi.org/10.1016/j.tibtech.2011.01.001)
22. Thiele, I., Vlassis, N., Fleming, R.M.T.: fastgapfill: efficient gap filling in metabolic networks. *Bioinformatics* **30**(17), 2529–2531 (2014). doi:[10.1093/bioinformatics/btu321](https://doi.org/10.1093/bioinformatics/btu321)
23. Prigent, S., Frioux, C., Dittami, S.M., Thiele, S., Larhlami, A., Collet, G., Gutknecht, F., Got, J., Eveillard, D., Bourdon, J., Plewniak, F., Tonon, T., Siegel, A.: Meneco, a topology-based gap-filling tool applicable to degraded genome-wide metabolic networks. *PLOS Computational Biology* **13**(1), 1005276 (2017). doi:[10.1371/journal.pcbi.1005276](https://doi.org/10.1371/journal.pcbi.1005276)
24. Henry, C.S., DeJongh, M., Best, A.A., Frybarger, P.M., Linsay, B., Stevens, R.L.: High-throughput generation, optimization and analysis of genome-scale metabolic models. *Nature Biotechnology* **28**(9), 977–982 (2010). doi:[10.1038/nbt.1672](https://doi.org/10.1038/nbt.1672)
25. Kumar, M., Ji, B., Zengler, K., Nielsen, J.: Modelling approaches for studying the microbiome. *Nature Microbiology* **4**(8), 1253–1267 (2019). doi:[10.1038/s41564-019-0491-9](https://doi.org/10.1038/s41564-019-0491-9)
26. Phelan, V.V., Liu, W.-T., Pogliano, K., Dorrestein, P.C.: Microbial metabolic exchange—the chemotype-to-phenotype link. *Nat. Chem. Biol.* **8**, 26–35 (2012). doi:[10.1038/nchembio.739](https://doi.org/10.1038/nchembio.739)
27. Caspi, R., Billington, R., Fulcher, C.A., Keseler, I.M., Kothari, A., Krummenacker, M., Latendresse, M., Midford, P.E., Ong, Q., Ong, W.K., Paley, S., Subhraveti, P., Karp, P.D.: The metacyc database of metabolic pathways and enzymes. *Nucleic Acids Research* **46**(D1), 633–639 (2018). doi:[10.1093/nar/gkx935](https://doi.org/10.1093/nar/gkx935)
28. Kanehisa, M., Sato, Y., Furumichi, M., Morishima, K., Tanabe, M.: New approach for understanding genome variations in KEGG. *Nucleic Acids Research* **47**(D1), 590–595 (2018). doi:[10.1093/nar/gky962](https://doi.org/10.1093/nar/gky962). <http://oup.prod.sis.lan/nar/article-pdf/47/D1/D590/27436321/gky962.pdf>
29. Karp, P.D., Latendresse, M., Paley, S.M., Ong, M.K.Q., Billington, R., Kothari, A., Weaver, D., Lee, T., Subhraveti, P., Spaulding, A., Fulcher, C., Keseler, I.M., Caspi, R.: Pathway tools version 19.0: Integrated software for pathway/genome informatics and systems biology (2015). [1510.03964](https://doi.org/10.1510.03964)
30. SRI International: PythonCyc (2014). <https://github.com/latendresse/PythonCyc>
31. The UniProt Consortium: UniProt: the universal protein knowledgebase. *Nucleic Acids Research* **45**(D1), 158–169 (2016). doi:[10.1093/nar/gkw1099](https://doi.org/10.1093/nar/gkw1099). <http://oup.prod.sis.lan/nar/article-pdf/45/D1/D158/23819877/gkw1099.pdf>
32. Jeske, L., Placzek, S., Schomburg, I., Chang, A., Schomburg, D.: BRENDA in 2019: a European ELIXIR core data resource. *Nucleic Acids Research* **47**(D1), 542–549 (2018). doi:[10.1093/nar/gky1048](https://doi.org/10.1093/nar/gky1048). <http://oup.prod.sis.lan/nar/article-pdf/47/D1/D542/27437170/gky1048.pdf>
33. Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., Madden, T.L.: Blast+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009). doi:[10.1186/1471-2105-10-421](https://doi.org/10.1186/1471-2105-10-421)
34. Tange, O.: Gnu parallel - the command-line power tool. *login: The USENIX Magazine*, 42–47 (2011)
35. Slater, G.S.C., Birney, E.: Automated generation of heuristics for biological sequence comparison. *BMC Bioinformatics* **6**, 31 (2005). doi:[10.1186/1471-2105-6-31](https://doi.org/10.1186/1471-2105-6-31)
36. Wickham, H.: Stringr: Simple, Consistent Wrappers for Common String Operations. (2019). R package version 1.4.0. <https://CRAN.R-project.org/package=stringr>
37. Pagès, H., Aboyoun, P., Gentleman, R., DebRoy, S.: Biostrings: Efficient Manipulation of Biological Strings. (2019). R package version 2.50.2
38. Saier, M.H., Reddy, V.S., Tamang, D.G., Vastermark, A.: The transporter classification database. *Nucleic Acids Research* **42**(D1), 251–258 (2013). doi:[10.1093/nar/gkt1097](https://doi.org/10.1093/nar/gkt1097)
39. Machado, D., Andrejev, S., Tramontano, M., Patil, K.R.: Fast automated reconstruction of genome-scale metabolic models for microbial species and communities. *Nucleic Acids Research* **46**(15), 7542–7553 (2018).

- doi:[10.1093/nar/gky537](https://doi.org/10.1093/nar/gky537). <http://oup.prod.sis.lan/nar/article-pdf/46/15/7542/25689981/gky537.pdf>
40. Webb, E.C., *et al.*: Enzyme Nomenclature 1992. Recommendations of the Nomenclature Committee of the International Union of Biochemistry and Molecular Biology on the Nomenclature and Classification of Enzymes. vol. Ed. 6. Academic Press, ??? (1992)
41. Bernard, T., Bridge, A., Morgat, A., Moretti, S., Xenarios, I., Pagni, M.: Reconciliation of metabolites and biochemical reactions for metabolic networks. *Brief Bioinform* **15**(1), 123–135 (2014). doi:[10.1093/bib/bbs058](https://doi.org/10.1093/bib/bbs058)
42. Brbić, M., Piškorec, M., Vidulin, V., Kriško, A., Šmuc, T., Supek, F.: The landscape of microbial phenotypic traits and associated genes. *Nucleic Acids Research*, 964 (2016). doi:[10.1093/nar/gkw964](https://doi.org/10.1093/nar/gkw964)
43. Benedict, M.N., Mundy, M.B., Henry, C.S., Chia, N., Price, N.D.: Likelihood-based gene annotations for gap filling and quality assessment in genome-scale metabolic models. *PLOS Computational Biology* **10**(10), 1–14 (2014). doi:[10.1371/journal.pcbi.1003882](https://doi.org/10.1371/journal.pcbi.1003882)
44. Dreyfuss, J.M., Zucker, J.D., Hood, H.M., Ocasio, L.R., Sachs, M.S., Galagan, J.E.: Reconstruction and validation of a genome-scale metabolic model for the filamentous fungus *Neurospora crassa* using FARM. *PLoS Computational Biology* **9**(7), 1003126 (2013). doi:[10.1371/journal.pcbi.1003126](https://doi.org/10.1371/journal.pcbi.1003126)
45. Medlock, G.L., Papin, J.A.: Guiding the refinement of biochemical knowledgebases with ensembles of metabolic networks and machine learning. *Cell Systems* **10**(1), 109–1193 (2020). doi:[10.1016/j.cels.2019.11.006](https://doi.org/10.1016/j.cels.2019.11.006)
46. Smalla, K., Wachtendorf, U., Heuer, H., Liu, W.-t., Forney, L.: Analysis of biologic substrate utilization patterns by microbial communities. *Applied and Environmental Microbiology* **64**(4), 1220–1225 (1998). <https://aem.asm.org/content/64/4/1220.full.pdf>
47. Bochner, B.R.: Global phenotypic characterization of bacteria. *FEMS Microbiology Reviews* **33**(1), 191–205 (2009). doi:[10.1111/j.1574-6976.2008.00149.x](https://doi.org/10.1111/j.1574-6976.2008.00149.x)
48. Lieven, C., Beber, M.E., Olivier, B.G., Bergmann, F.T., Ataman, M., Babaei, P., Bartell, J.A., Blank, L.M., Chauhan, S., Correia, K., Diener, C., Dräger, A., Ebert, B.E., Edirisinghe, J.N., Faria, J.P., Feist, A.M., Fengos, G., Fleming, R.M.T., García-Jiménez, B., Hatzimanikatis, V., van Helvoirt, W., Henry, C.S., Hermjakob, H., Herrgård, M.J., Kaafarani, A., Kim, H.U., King, Z., Klamt, S., Klipp, E., Koehorst, J.J., König, M., Lakshmanan, M., Lee, D.-Y., Lee, S.Y., Lee, S., Lewis, N.E., Liu, F., Ma, H., Machado, D., Mahadevan, R., Maia, P., Mardinoglu, A., Medlock, G.L., Monk, J.M., Nielsen, J., Nielsen, L.K., Nogales, J., Nookaew, I., Palsson, B.O., Papin, J.A., Patil, K.R., Poolman, M., Price, N.D., Resendis-Antonio, O., Richelle, A., Rocha, I., Sánchez, B.J., Schaap, P.J., Sheriff, R.S.M., Shoaie, S., Sonnenschein, N., Teusink, B., Vilaca, P., Vik, J.O., Wodke, J.A.H., Xavier, J.C., Yuan, Q., Zakhartsev, M., Zhang, C.: Memote for standardized genome-scale metabolic model testing. *Nat. Biotechnol.* **38**, 272–276 (2020). doi:[10.1038/s41587-020-0446-y](https://doi.org/10.1038/s41587-020-0446-y)
49. Reimer, L.C., Vetscinova, A., Carbasse, J.S., Söhngen, C., Gleim, D., Ebeling, C., Overmann, J.: BacDive in 2019: bacterial phenotypic data for High-throughput biodiversity analysis. *Nucleic Acids Research* **47**(D1), 631–636 (2018). doi:[10.1093/nar/gky879](https://doi.org/10.1093/nar/gky879). <http://oup.prod.sis.lan/nar/article-pdf/47/D1/D631/27436018/gky879.pdf>
50. Leinweber, K.: BacDiveR: A Programmatic Interface For BacDive, The DSMZ's Bacterial Diversity Metadatabase. (2019). R package version 0.6.0. <https://github.com/TIBHannover/BacDiveR>
51. Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V., Zdobnov, E.M.: BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**, 3210–2 (2015). doi:[10.1093/bioinformatics/btv351](https://doi.org/10.1093/bioinformatics/btv351)
52. Sayers, E.W., Beck, J., Brister, J.R., Bolton, E.E., Canese, K., Comeau, D.C., Funk, K., Ketter, A., Kim, S., Kimchi, A., Kitts, P.A., Kuznetsov, A., Lathrop, S., Lu, Z., McGarvey, K., Madden, T.L., Murphy, T.D., O'Leary, N., Phan, L., Schneider, V.A., Thibaud-Nissen, F., Trawick, B.W., Pruitt, K.D., Ostell, J.: Database resources of the national center for biotechnology information. *Nucleic Acids Research* **48**(D1), 9–16 (2019). doi:[10.1093/nar/gkz899](https://doi.org/10.1093/nar/gkz899)
53. Zhu, B., Stülke, J.: SubtiWiki in 2018: from genes and proteins to functional network annotation of the model organism *Bacillus subtilis*. *Nucleic Acids Research* **46**(D1), 743–748 (2017). doi:[10.1093/nar/gkx908](https://doi.org/10.1093/nar/gkx908). <https://academic.oup.com/nar/article-pdf/46/D1/D743/23162685/gkx908.pdf>
54. Monk, J.M., Lloyd, C.J., Brunk, E., Mih, N., Sastry, A., King, Z., Takeuchi, R., Nomura, W., Zhang, Z., Mori, H., *et al.*: iM1515, a knowledgebase that computes *Escherichia coli* traits. *Nature biotechnology* **35**(10), 904 (2017)
55. Turner, K.H., Wessel, A.K., Palmer, G.C., Murray, J.L., Whiteley, M.: Essential genome of *Pseudomonas aeruginosa* in cystic fibrosis sputum. *Proceedings of the National Academy of Sciences* **112**(13), 4110–4115 (2015)
56. Price, M.N., Wetmore, K.M., Waters, R.J., Callaghan, M., Ray, J., Liu, H., Kuehl, J.V., Melnyk, R.A., Lamson, J.S., Suh, Y., *et al.*: Mutant phenotypes for thousands of bacterial genes of unknown function. *Nature* **557**(7706), 503 (2018)
57. Glass, J.I., Assad-Garcia, N., Alperovich, N., Yooseph, S., Lewis, M.R., Maruf, M., Hutchison, C.A., Smith, H.O., Venter, J.C.: Essential genes of a minimal bacterium. *Proceedings of the National Academy of Sciences* **103**(2), 425–430 (2006)
58. Holzhütter, H.-G.: The principle of flux minimization and its application to estimate stationary fluxes in metabolic networks. *European journal of biochemistry* **271**(14), 2905–2922 (2004)
59. Mahadevan, R., Schilling, C.H.: The effects of alternate optimal solutions in constraint-based genome-scale metabolic models. *Metab Eng* **5**(4), 264–276 (2003). doi:[10.1016/j.ymben.2003.09.002](https://doi.org/10.1016/j.ymben.2003.09.002)
60. Choi, J., Yang, F., Stepanauskas, R., Cardenas, E., Garoutte, A., Williams, R., Flater, J., Tiedje, J.M., Hofmockel, K.S., Gelder, B., Howe, A.: Strategies to improve reference databases for soil microbiomes. *The ISME Journal* **11**(4), 829–834 (2016). doi:[10.1038/ismej.2016.168](https://doi.org/10.1038/ismej.2016.168)
61. Magnusdottir, S., Heinken, A., Kutt, L., Ravcheev, D.A., Bauer, E., Noronha, A., Greenhalgh, K., Jäger, C., Baginska, J., Wilmes, P., Fleming, R.M.T., Thiele, I.: Generation of genome-scale metabolic reconstructions for 773 members of the human gut microbiota. *Nature Biotechnology* (2016). doi:[10.1038/nbt.3703](https://doi.org/10.1038/nbt.3703)

62. Kassambara, A., Mundt, F.: Factoextra: Extract and Visualize the Results of Multivariate Data Analyses. (2019). R package version 1.0.6. <https://CRAN.R-project.org/package=factoextra>
63. Sekhon, J.S.: Multivariate and propensity score matching software with automated balance optimization: The Matching package for R. *Journal of Statistical Software* **42**(7), 1–52 (2011)
64. Louis, P., Hold, G.L., Flint, H.J.: The gut microbiota, bacterial metabolites and colorectal cancer. *Nat. Rev. Microbiol.* **12**, 661–72 (2014). doi:[10.1038/nrmicro3344](https://doi.org/10.1038/nrmicro3344)
65. Rivera-Chávez, F., Bäuml, A.J.: The pyromaniac inside you: Salmonella metabolism in the host gut. *Annual Review of Microbiology* **69**(1), 31–48 (2015). doi:[10.1146/annurev-micro-091014-104108](https://doi.org/10.1146/annurev-micro-091014-104108). PMID: 26002180
66. Feist, A.M., Scholten, J.C.M., Palsson, B.O., Brockman, F.J., Ideker, T.: Modeling methanogenesis with a genome-scale metabolic reconstruction of methanosarcina barkeri. *Mol Syst Biol* **2**, 2006–0004 (2006). doi:[10.1038/msb4100046](https://doi.org/10.1038/msb4100046)
67. Sieber, J.R., McInerney, M.J., Gunsalus, R.P.: Genomic insights into syntrophy: The paradigm for anaerobic metabolic cooperation. *Annual Review of Microbiology* **66**(1), 429–452 (2012). doi:[10.1146/annurev-micro-090110-102844](https://doi.org/10.1146/annurev-micro-090110-102844). PMID: 22803797. <https://doi.org/10.1146/annurev-micro-090110-102844>
68. Bauer, E., Zimmermann, J., Baldini, F., Thiele, I., Kaleta, C.: Bacarena: Individual-based metabolic modeling of heterogeneous microbes in complex communities. *PLOS Computational Biology* **13**(5), 1–22 (2017). doi:[10.1371/journal.pcbi.1005544](https://doi.org/10.1371/journal.pcbi.1005544)
69. Pasolli, E., Asnicar, F., Manara, S., Zolfo, M., Karcher, N., Armanini, F., Beghini, F., Manghi, P., Tett, A., Ghensi, P., Collado, M.C., Rice, B.L., DuLong, C., Morgan, X.C., Golden, C.D., Quince, C., Huttenhower, C., Segata, N.: Extensive unexplored human microbiome diversity revealed by over 150,000 genomes from metagenomes spanning age, geography, and lifestyle. *Cell* **176**(3), 649–66220 (2019). doi:[10.1016/j.cell.2019.01.001](https://doi.org/10.1016/j.cell.2019.01.001)
70. Alanjary, M., Steinke, K., Ziemert, N.: AutoMLST: an automated web server for generating multi-locus species trees highlighting natural product potential. *Nucleic Acids Research* **47**(W1), 276–282 (2019). doi:[10.1093/nar/gkz282](https://doi.org/10.1093/nar/gkz282)
71. Ondov, B.D., Treangen, T.J., Melsted, P., Mallonee, A.B., Bergman, N.H., Koren, S., Phillippy, A.M.: Mash: fast genome and metagenome distance estimation using MinHash. *Genome Biology* **17**(1) (2016). doi:[10.1186/s13059-016-0997-x](https://doi.org/10.1186/s13059-016-0997-x)
72. Dowle, M., Srinivasan, A.: Data.table: Extension of 'data.frame'. (2019). R package version 1.12.6. <https://CRAN.R-project.org/package=data.table>
73. Wickham, H.: Stringr: Simple, Consistent Wrappers for Common String Operations. (2019). R package version 1.4.0. <https://CRAN.R-project.org/package=stringr>
74. Gelius-Dietrich, G., Desouki, A.A., Fritzemeier, C.J., Lercher, M.J.: Sybil—efficient constraint-based modelling in R. *BMC Syst Biol* **7**, 125 (2013). doi:[10.1186/1752-0509-7-125](https://doi.org/10.1186/1752-0509-7-125)
75. Davis, T.L., Day, A.: Getopt: C-Like 'getopt' Behavior. (2019). R package version 1.20.3. <https://CRAN.R-project.org/package=getopt>
76. Wickham, H.: Reshaping data with the reshape package. *Journal of Statistical Software* **21**(12), 1–20 (2007)
77. Corporation, M., Weston, S.: doParallel: Foreach Parallel Adaptor for the 'parallel' Package. (2019). R package version 1.0.15. <https://CRAN.R-project.org/package=doParallel>
78. Microsoft, Weston, S.: Foreach: Provides Foreach Looping Construct. (2019). R package version 1.4.7. <https://CRAN.R-project.org/package=foreach>
79. Bengtsson, H.: R.utils: Various Programming Utilities. (2019). R package version 2.9.2. <https://CRAN.R-project.org/package=R.utils>
80. Gagolewski, M.: R Package Stringi: Character String Processing Facilities. (2019). <http://www.gagolewski.com/software/stringi/>
81. Gelius-Dietrich, G.: glpkAPI: R Interface to C API of GLPK. (2018). R package version 1.3.1. <https://CRAN.R-project.org/package=glpkAPI>
82. Pagès, H., Aboyoun, P., Gentleman, R., DebRoy, S.: Biostrings: Efficient Manipulation of Biological Strings. (2019). R package version 2.54.0
83. Bornstein, B.J., Keating, S.M., Jouraku, A., Hucka, M.: Libsbml: an api library for sbml. *Bioinformatics* **24**(6), 880–881 (2008). doi:[10.1093/bioinformatics/btn051](https://doi.org/10.1093/bioinformatics/btn051)
84. Cook, G.M., Greening, C., Hards, K., Berney, M.: Chapter one - energetics of pathogenic bacteria and opportunities for drug development. In: Poole, R.K. (ed.) *Advances in Bacterial Pathogen Biology. Advances in Microbial Physiology*, vol. 65, pp. 1–62. Academic Press, ??? (2014). doi:[10.1016/bs.ampbs.2014.08.001](https://doi.org/10.1016/bs.ampbs.2014.08.001). <http://www.sciencedirect.com/science/article/pii/S0065291114000022>
85. Goldberg, I., Rock, J., Ben-Bassat, A., Mateles, R.: Bacterial yields on methanol, methylamine, formaldehyde, and formate. *Biotechnology and bioengineering* **18**(12), 1657–1668 (1976)
86. Oliphant, K., Allen-Vercoe, E.: Macronutrient metabolism by the human gut microbiome: major fermentation by-products and their impact on host health. *Microbiome* **7**, 91 (2019). doi:[10.1186/s40168-019-0704-8](https://doi.org/10.1186/s40168-019-0704-8)
87. Ríos-Covián, D., Ruas-Madiedo, P., Margolles, A., Gueimonde, M., de Los Reyes-Gavilán, C.G., Salazar, N.: Intestinal short chain fatty acids and their link with diet and human health. *Front Microbiol* **7**, 185 (2016)
88. Ziels, R.M., Nobu, M.K., Sousa, D.Z.: Elucidating syntrophic butyrate-degrading populations in anaerobic digesters using stable-isotope-informed genome-resolved metagenomics. *mSystems* **4**(4) (2019). doi:[10.1128/mSystems.00159-19](https://doi.org/10.1128/mSystems.00159-19)
89. Nurk, S., Meleshko, D., Korobeynikov, A., Pevzner, P.A.: metaSPAdes: a new versatile metagenomic assembler. *Genome Research* **27**(5), 824–834 (2017). doi:[10.1101/gr.213959.116](https://doi.org/10.1101/gr.213959.116)
90. Arkin, A.P., Cottingham, R.W., Henry, C.S., Harris, N.L., Stevens, R.L., Maslov, S., Dehal, P., Ware, D., Perez, F., Canon, S., Sneddon, M.W., Henderson, M.L., Riehl, W.J., Murphy-Olson, D., Chan, S.Y., Kamimura, R.T., Kumari, S., Drake, M.M., Brettin, T.S., Glass, E.M., Chivian, D., Gunter, D., Weston, D.J., Allen, B.H., Baumohl, J., Best, A.A., Bowen, B., Brenner, S.E., Bun, C.C., Chandonia, J.-M., Chia, J.-M., Colasanti, R., Conrad, N., Davis, J.J., Davison, B.H., DeJongh, M., Devoid, S., Dietrich, E., Dubchak, I.,

- Edirisinghe, J.N., Fang, G., Faria, J.P., Frybarger, P.M., Gerlach, W., Gerstein, M., Greiner, A., Gurtowski, J., Haun, H.L., He, F., Jain, R., Joachimiak, M.P., Keegan, K.P., Kondo, S., Kumar, V., Land, M.L., Meyer, F., Mills, M., Novichkov, P.S., Oh, T., Olsen, G.J., Olson, R., Parrello, B., Pasternak, S., Pearson, E., Poon, S.S., Price, G.A., Ramakrishnan, S., Ranjan, P., Ronald, P.C., Schatz, M.C., Seaver, S.M.D., Shukla, M., Sutormin, R.A., Syed, M.H., Thomason, J., Tintle, N.L., Wang, D., Xia, F., Yoo, H., Yoo, S., Yu, D.: KBase: The united states department of energy systems biology knowledgebase. *Nature Biotechnology* **36**(7), 566–569 (2018). doi:[10.1038/nbt.4163](https://doi.org/10.1038/nbt.4163)
91. Quince, C., Walker, A.W., Simpson, J.T., Loman, N.J., Segata, N.: Shotgun metagenomics, from sampling to analysis. *Nature Biotechnology* **35**(9), 833–844 (2017). doi:[10.1038/nbt.3935](https://doi.org/10.1038/nbt.3935)
92. Kim, W.J., Kim, H.U., Lee, S.Y.: Current state and applications of microbial genome-scale metabolic models. *Current Opinion in Systems Biology* **2**, 10–18 (2017). doi:[10.1016/j.coisb.2017.03.001](https://doi.org/10.1016/j.coisb.2017.03.001). Regulatory and metabolic networks • Cancer and systemic diseases
93. Graspentner, S., Waschina, S., Künzel, S., Twisselmann, N., Rausch, T.K., Cloppenborg-Schmidt, K., Zimmermann, J., Viemann, D., Herting, E., Göpel, W., Baines, J.F., Kaleta, C., Rupp, J., Härtel, C., Pagel, J.: Gut Dysbiosis With Bacilli Dominance and Accumulation of Fermentation Products Precedes Late-onset Sepsis in Preterm Infants. *Clinical Infectious Diseases* **69**(2), 268–277 (2018). doi:[10.1093/cid/ciy882](https://doi.org/10.1093/cid/ciy882). <https://academic.oup.com/cid/article-pdf/69/2/268/28893438/ciy882.pdf>
94. Byndloss, M.X., Olsan, E.E., Rivera-Chávez, F., Tiffany, C.R., Cevallos, S.A., Lokken, K.L., Torres, T.P., Byndloss, A.J., Faber, F., Gao, Y., *et al.*: Microbiota-activated ppar- γ signaling inhibits dysbiotic enterobacteriaceae expansion. *Science* **357**(6351), 570–575 (2017). doi:[10.1126/science.aam9949](https://doi.org/10.1126/science.aam9949)
95. Smith, P.M., Howitt, M.R., Panikov, N., Michaud, M., Gallini, C.A., Bohlooly-y, M., Glickman, J.N., Garrett, W.S.: The microbial metabolites, short-chain fatty acids, regulate colonic treg cell homeostasis. *Science* **341**(6145), 569–573 (2013). doi:[10.1126/science.1241165](https://doi.org/10.1126/science.1241165)
96. Pham, V.T., Lacroix, C., Braegger, C.P., Chassard, C.: Early colonization of functional groups of microbes in the infant gut. *Environmental Microbiology* **18**(7), 2246–2258 (2016). doi:[10.1111/1462-2920.13316](https://doi.org/10.1111/1462-2920.13316)
97. García-Campos, M.A., Espinal-Enríquez, J., Hernández-Lemus, E.: Pathway analysis: state of the art. *Frontiers in physiology* **6**, 383 (2015). doi:[10.3389/fphys.2015.00383](https://doi.org/10.3389/fphys.2015.00383)
98. Foster, K.R., Schluter, J., Coyte, K.Z., Rakoff-Nahoum, S.: The evolution of the host microbiome as an ecosystem on a leash. *Nature* **548**(7665), 43–51 (2017). doi:[10.1038/nature23292](https://doi.org/10.1038/nature23292)
99. Bateman, A., Birney, E., Durbin, R., Eddy, S.R., Howe, K.L., Sonnhammer, E.L.L.: The Pfam Protein Families Database. *Nucleic Acids Research* **28**(1), 263–266 (2000). doi:[10.1093/nar/28.1.263](https://doi.org/10.1093/nar/28.1.263). <http://oup.prod.sis.lan/nar/article-pdf/28/1/263/9895152/280263.pdf>
100. Galperin, M.Y., Koonin, E.V.: Sources of systematic error in functional annotation of genomes: domain rearrangement, non-orthologous gene displacement and operon disruption. *In silico biology* **1**(1), 55–67 (1998)
101. El-Gebali, S., Mistry, J., Bateman, A., Eddy, S.R., Luciani, A., Potter, S.C., Qureshi, M., Richardson, L.J., Salazar, G.A., Smart, A., Sonnhammer, E.L., Hirsh, L., Paladin, L., Piovesan, D., Tosatto, S.C., Finn, R.D.: The Pfam protein families database in 2019. *Nucleic Acids Research* **47**(D1), 427–432 (2018). doi:[10.1093/nar/gky995](https://doi.org/10.1093/nar/gky995). <http://oup.prod.sis.lan/nar/article-pdf/47/D1/D427/27436497/gky995.pdf>
102. Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S.K., Cook, H., Mende, D.R., Letunic, I., Rattei, T., Jensen, L.J., von Mering, C., Bork, P.: eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids Res.* **47**, 309–314 (2019). doi:[10.1093/nar/gky1085](https://doi.org/10.1093/nar/gky1085)
103. Douglas, G.M., Maffei, V.J., Zaneveld, J., Yurgel, S.N., Brown, J.R., Taylor, C.M., Huttenhower, C., Langille, M.G.I.: PICRUSt2: An improved and extensible approach for metagenome inference. *bioRxiv* (2019). doi:[10.1101/672295](https://doi.org/10.1101/672295)

Additional Files

Table S1 — New reactions and metabolites added to biochemistry database.
see file: Table_S1.xlsx


Table S2 — Organisms included in fermentation product validation test.
see file: Table_S2.xlsx

Table S4 — References for substance production and consumption in anaerobic gut communities (see Supplementary Figure S1).
see file: Table_S4.ods

Table S5 — 127 Species-level genome bins (SGBs) from Pasolli *et al.*, 2019 [69] and 1270 mapped reference genome assemblies from RefSeq.
see file: Table_S5.ods

Table S3 — Organisms used in modelling of the anaerobic food web of the human gut microbiome.

RefSeq Assembly	Organism name	Reconstruction method
GCF_000173975.1	<i>Anaerobutyricum hallii</i> DSM 3353	gapseq / modelseed / carveme
GCF_000025985.1	<i>Bacteroides fragilis</i> NCTC 9343	gapseq / modelseed / carveme
GCF_001314975.1	<i>Bacteroides thetaiotaomicron</i>	gapseq / modelseed / carveme
GCF_000196555.1	<i>Bifidobacterium longum</i> subsp. <i>longum</i> JCM 1217	gapseq / modelseed / carveme
GCF_000157975.1	<i>Blautia hydrogenotrophica</i> DSM 10507	gapseq / modelseed / carveme
GCF_000013285.1	<i>Clostridium perfringens</i> ATCC 13124	gapseq / modelseed / carveme
GCF_003434235.1	<i>Coprococcus catus</i>	gapseq / modelseed / carveme
GCF_000155875.1	<i>Coprococcus comes</i> ATCC 27758	gapseq / modelseed / carveme
GCF_000154425.1	<i>Coprococcus eutactus</i> ATCC 27759	gapseq / modelseed / carveme
GCF_000189295.2	<i>Desulfovibrio desulfuricans</i> ND132	gapseq / modelseed / carveme
GCF_000391485.2	<i>Enterococcus faecalis</i> EnGen0107	gapseq / modelseed / carveme
GCF_000005845.2	<i>Escherichia coli</i> str. K-12 substr. MG1655	gapseq / modelseed / carveme
GCF_000162015.1	<i>Faecalibacterium prausnitzii</i> A2-165	gapseq / modelseed / carveme
GCF_003047065.1	<i>Lactobacillus acidophilus</i>	gapseq / modelseed / carveme
GCF_001304715.1	<i>Megasphaera elsdenii</i> 14-14	gapseq / modelseed / carveme
GCF_000195895.1	<i>Methanosarcina barkeri</i> str. <i>Fusaro</i>	manually curated (BiGG-ID: iAF692)[66]
GCF_000144405.1	<i>Prevotella melaninogenica</i> ATCC 25845	gapseq / modelseed / carveme
GCF_900101355.1	<i>Ruminococcus bromii</i>	gapseq / modelseed / carveme
GCF_000006945.2	<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar <i>Typhimurium</i> str. LT2	gapseq / modelseed / carveme
GCF_900637515.1	<i>Veillonella dispar</i>	gapseq / modelseed / carveme






Figure S2 — Similarity of **gapseq** models reconstructed for 127 species-level genome bins (SGBs) from metagenomes compared to models reconstructed for reference genomes (RefSeq Prokaryotic Genomes). The x-axis represents the genome assembly completion of SGBs estimated using the BUSCO software version 4.0.6 [51]. The line shows the result of non-linear regression using a logarithmic function of form $y(x) = c + b * \log(x)$. Sequences of SGBs were obtained from Pasolli *et al.*, 2019 [69].