

# Deficiency in cytosine DNA methylation leads to high chaperonin expression and tolerance to aminoglycosides in *Vibrio cholerae*

André Carvalho<sup>1,2</sup>, Didier Mazel<sup>1\*</sup>, Zeynep Baharoglu<sup>1\*</sup>

<sup>1</sup> Département Génomes et Génétique, Institut Pasteur, UMR3525, CNRS, Unité Plasticité du Génome Bactérien, Paris, France

<sup>2</sup> Sorbonne Université, Collège doctoral, F-75005 Paris, France

\* corresponding authors, [mazel@pasteur.fr](mailto:mazel@pasteur.fr), [zeynep.baharoglu@pasteur.fr](mailto:zeynep.baharoglu@pasteur.fr)

# ABSTRACT

Antibiotic resistance has become a major global issue. Understanding the molecular mechanisms underlying microbial adaptation to antibiotics is of keen importance to fight Antimicrobial Resistance (AMR). Aminoglycosides are a class of antibiotics that target the small subunit of the bacterial ribosome, disrupting translational fidelity and increasing the levels of misfolded proteins in the cell. In this work, we investigated the role of VchM, a DNA methyltransferase, in the response of the human pathogen *Vibrio cholerae* to aminoglycosides. VchM is a *V. cholerae* specific orphan m5C DNA methyltransferase that generates cytosine methylation at 5'-RCCGGY-3' motifs. We show that deletion of *vchM*, although causing a growth defect in absence of stress, allows *V. cholerae* cells to cope with aminoglycoside stress at both sub-lethal and lethal concentrations of these antibiotics. Through transcriptomic and genetic approaches, we show that *groESL-2* (a specific set of chaperonin-encoding genes located on the second chromosome of *V. cholerae*), are upregulated in cells lacking *vchM* and are needed for the tolerance of *vchM* mutant to lethal aminoglycoside treatment, likely by fighting aminoglycoside-induced misfolded proteins. Interestingly, preventing VchM methylation of the four RCCGGY sites located in *groESL-2* region, leads to a higher expression of these genes in WT cells, showing that VchM modulates the expression of these chaperonins in *V. cholerae* directly through DNA methylation.

## AUTHOR SUMMARY

Bacteria are organisms with a remarkable ability to adapt to several stress conditions, including to the presence of antibiotics. The molecular mechanisms underlying such adaptation lead, very often, to phenomena like antimicrobial tolerance and resistance, responsible for the frequent failure of antibiotic treatment. The study of these molecular mechanisms is thus an important tool to understand development of antimicrobial resistance in bacteria. In this work, we show that abrogating cytosine DNA methylation in *Vibrio cholerae* increases its tolerance to aminoglycosides, a class of antibiotics that cause protein misfolding. DNA methylation is known to affect gene expression and regulate several cellular processes in bacteria. Here we provide evidence that DNA methylation also has a more direct role in controlling antibiotic susceptibility in bacteria. Consequently, the study of bacterial DNA methyltransferases and DNA methylation should not be overlooked when addressing the problem of antimicrobial tolerance/resistance.

## INTRODUCTION

In the past decades, the over/misuse and large-scale production of antibiotics has created a serious ecological problem with important consequences for the emergence of antimicrobial resistance (AMR). In fact, a large proportion of the antibiotics ingested are released intact in the environment (1, 2) and found at trace levels or as gradients in various environments (3, 4). Hence, in these environments, one can find the presence of very low

doses of drugs commonly referred as subMIC, i.e. under the MIC (Minimal Inhibitory Concentration). Although not enough to kill or prevent the growth of bacterial populations, subMIC doses of antibiotics are proposed to work as signaling molecules (5) and trigger important stress mechanisms that often result in development of antibiotic resistance (4, 6–8). We have previously shown that subMIC of antibiotics, such as aminoglycosides, trigger common and specific stress responses in Gram-negative bacteria (9, 10).

Aminoglycosides (AGs) are positively charged molecules that bind 16S rRNA at the 30S ribosomal subunit and negatively affect translation. Specifically, AGs (e.g. tobramycin, streptomycin, kanamycin, gentamicin and neomycin) are known to disrupt translational fidelity and increase the levels of mistranslation, i.e. the misincorporation of certain amino acids in proteins (11, 12). In turn, high levels of mistranslation result in the production and accumulation of aberrant proteins in the cell, which contribute to the collapse of important cell processes and ultimately lead to cell death (13, 14).

*V. cholerae* is a water-borne gram-negative bacterium, human pathogen and the causative agent of cholera disease. As part of its life cycle, *V. cholerae* often transits between the human gut and the external environment where it can find low doses of antibiotics. During our studies to better understand adaption of *V. cholerae* to aminoglycosides (15), we observed that a mutant of a *V. cholerae*'s specific DNA methyltransferase (*vca0198* - VchM) was less susceptible to aminoglycosides than its isogenic WT strain, suggesting that DNA methylation could play a role in *V. cholerae* adaptation to AGs. *vchM* codes for an Orphan m5C DNA methyltransferase that causes DNA methylation at 5'-RCCGGY-3' motifs (16). DNA methylation is catalyzed by enzymes called

DNA methyltransferases (DNA MTases) that transfer a methyl group from S-adenosyl-L-methionine (SAM) to adenine and cytosine in specific DNA motifs (17, 18). As a result, one can find the existence of small amounts of N6-methyl-adenine (6mA), C5-methyl-cytosine (5mC) and N4-methyl-cytosine (4mC) in the DNA of both eukaryotes and prokaryotes. In bacteria, the existence of such modified DNA bases have been shown to play a critical role in processes such as protection against invasive DNA, DNA replication and repair, cell cycle regulation and control of gene expression (19–23).

While it was previously proposed that VchM plays a role in the cell envelope stress response of *V. cholerae* (23), no link between this DNA MTase and antibiotic stress has yet been established. Here, we show that deletion of *vchM* (although causing a growth defect in absence of stress) allows *V. cholerae* cells to better deal with the effect of aminoglycosides. In fact, not only the *vchM* mutant is a better competitor during growth in presence of subMIC doses of aminoglycosides, it is also more tolerant to killing by lethal doses of these antibiotics. Transcriptome analysis of a  $\Delta vchM$  strain revealed the upregulation of *groESL-2* genes, a specific set of chaperonin-encoding genes located on the second chromosome of *V. cholerae*. High expression of *groESL-2* genes (but not of chromosome one *groESL-1* homologues) determines the higher tolerance of  $\Delta vchM$  to lethal AG treatment, suggesting a new and specific role of *groESL-2* in managing AG-mediated proteotoxic stress. Interestingly, we observed the presence of four VchM motifs in *groESL-2* region. Preventing methylation of all these sites in the WT strain by disrupting such motifs results in increased expression of these genes. Intriguingly, the high expression of *groESL-2* does not seem to contribute to the competitive advantage of the  $\Delta vchM$  strain

grown under subMIC AG which suggests the involvement of additional players in the global response of  $\Delta vchM$  to aminoglycosides.

## RESULTS

### ***V. cholerae* cells lacking *vchM* cope better with subMIC doses of AGs**

In order to explore a possible role of *vchM* in the response of *V. cholerae* O1 El Tor N16961 to aminoglycosides, we constructed an in-frame deletion mutant of *vchM* by allelic replacement with an antibiotic resistance cassette, and compared its growth to the isogenic wild-type (WT) strain, in rich media, with or without increasing concentrations of subMIC tobramycin (Fig 1). As previously described (23), this mutant exhibits a reduced doubling rate when grown in monoculture in antibiotic free rich media. However, the difference in growth between WT and  $\Delta vchM$  strains observed in absence of antibiotics becomes gradually more negligible with increasing concentrations of subMIC TOB. At higher concentrations (90% of the MIC),  $\Delta vchM$  even displays a clear advantage over the WT (Fig 1A). Importantly, a  $\Delta vchM$  strain harboring a low-copy number plasmid with *vchM* gene under the control of its own promoter behaves as the WT strain in absence of tobramycin and even slightly worse in presence of higher doses of this drug (Fig 1A), showing that the observed growth phenotypes are due to the absence of *vchM*.

Next, we asked whether the growth phenotype observed in monocultures was translatable to a higher relative fitness in co-cultures in the presence of subMIC doses of tobramycin and other AGs. For that, we competed both WT and  $\Delta vchM$  strains (both  $lacZ^+$ ) with an isogenic  $\Delta lacZ$  mutant (initial ratio of 1:1), in MH or MH supplemented with subMIC concentrations (50% MIC) of the aminoglycosides tobramycin (TOB), gentamicin (GEN) and neomycin (NEO). We assessed relative fitness by plating cultures after 20 hours of growth and counting the final proportion of  $lacZ^+/lacZ^-$  colonies. Competition of WT against the  $lacZ^-$  mutant served as a control to account for any effect of  $lacZ$  deletion on growth. Supporting the previous results in monocultures,  $\Delta vchM$  is outcompeted by the  $lacZ^-$  mutant in MH ( $\approx 10$ -fold difference) (Fig 1B). More importantly, in presence of low concentrations of aminoglycosides,  $\Delta vchM$  is either equally competitive or even displays a clear growth advantage over the reference strain (Fig 1B). Additionally, in order to test whether these results hold for drugs other than aminoglycosides we performed competitions in the presence of chloramphenicol (CAM) and the beta-lactam carbenicillin (CARB). Unlike AGs, the presence of low concentrations of these drugs did not increased the relative fitness of the  $\Delta vchM$  mutant.

Altogether, these results confirm that lack of  $vchM$  in *V. cholerae* negatively impacts growth in antibiotic-free media (23) but confers a selective advantage to *V. cholerae* in presence of subMIC doses of AGs (Fig 1). In order to test if deleting  $vchM$  affects the MIC of these drugs, we measured the MIC of both WT and  $\Delta vchM$  mutant and found no difference (Table 1).

**Table 1.** MICs ( $\mu\text{g/ml}$ ) of the different antibiotics tested for *V. cholerae* N16961 WT and  $\Delta vchM$  strains

| Strain        | TOB | GEN | NEO | CAM | CARB |
|---------------|-----|-----|-----|-----|------|
| WT            | 1-2 | 1   | 4   | 1   | 15   |
| $\Delta vchM$ | 1-2 | 1   | 4   | 1   | 7.5  |

TOB, tobramycin; GEN, gentamicin; NEO, neomycin; CAM, chloramphenicol; CARB, carbenicillin

## **VchM deficiency promotes higher tolerance to lethal doses of aminoglycosides**

It has been previously shown that certain mutations affect functions conferring bacterial populations a tolerant phenotype towards a specific drug (24). Such bacterial populations can transiently withstand lethal doses of that drug without necessarily any impact on the MIC of the population (24).

To continue exploring the different susceptibility of the  $\Delta vchM$  strain to aminoglycosides we assessed the survival rate of *V. cholerae* WT and  $\Delta vchM$  strains during treatment with lethal doses of tobramycin and gentamicin at 20x and 10x the MIC, respectively (Fig 2). Given the inherent growth defect of VchM deficiency, we performed time-dependent killing curves on stationary phase cells, where both WT and  $\Delta vchM$  strains are no longer actively growing, excluding a possible link between growth rate and aminoglycoside lethality as previously shown (25). Strikingly, survival to both antibiotics was increased 10-1000 fold in the  $\Delta vchM$  mutant, suggesting that the absence of VchM allows *V. cholerae* to transiently withstand lethal doses of these aminoglycosides (Fig 2).

One crucial aspect that determines the efficacy of aminoglycoside treatment is the uptake of these drugs by the bacterial cell. This process is energy dependent and requires a threshold membrane potential (26). We used a previously reported assay that measures cell fluorescence after incubation with fluorescent-marked neomycin (neo-cy5) (27) as a proxy for aminoglycoside uptake. We did not observe any difference in fluorescence between WT and  $\Delta vchM$  mutant strains (S1 Fig). Thus, differential uptake of aminoglycosides is unlikely the reason for the increased tolerance to these drugs in  $\Delta vchM$ .

# **A specific set of chaperonins is upregulated in $\Delta vchM$ cells**

To understand the high tolerance to aminoglycosides observed in  $\Delta vchM$ , we performed RNA-seq on stationary phase cells of WT and  $\Delta vchM$  strains grown in rich, stress-free media. The analysis of the transcriptome of  $\Delta vchM$  *V. cholerae* O1 El Tor N16961 strain reveals the significant upregulation (fold change  $\geq 2$ ,  $p < 0.01$ ) and downregulation (fold change  $\leq -2$ ,  $p < 0.01$ ) of 68 and 53 genes, respectively (S1 Table). Among the differentially expressed, we found four genes directly involved in protein folding to be upregulated in  $\Delta vchM$  strain. Those are the molecular chaperones GroEL and co-chaperonins GroES (Table 2). In many bacterial species, GroEL and its co-chaperonin GroES form a molecular machine essential for folding of large newly synthesized proteins also helping re-folding of proteins damaged by proteotoxic stress (28). Interestingly, overexpression of GroES and GroEL

195 proteins was found to promote short-term tolerance to aminoglycoside-induced protein  
196 misfolding in *E. coli* (29).

**Table 2.** Protein folding and stabilization genes upregulated in  $\Delta vchM$  (fold change > 2, p-value < 0.01)

| Locus   | Name    | Fold change<br>( $\Delta vchM$ /WT) | Annotation                 |
|---------|---------|-------------------------------------|----------------------------|
| vc2665  | groEL-1 | <b>2.24</b>                         | Chaperonin, 60-kDa subunit |
| vca0820 | groEL-2 | <b>3.54</b>                         | Chaperonin, 60-kDa subunit |
| vc2664  | groES-1 | <b>2.60</b>                         | Chaperonin, 10-kDa subunit |
| vca0819 | groES-2 | <b>6.78</b>                         | Chaperonin, 10-kDa subunit |

197

198 *V. cholerae* is one of, at least, seven *Vibrio* species harboring two copies of *groES* –  
199 *groEL* (*groESL*) bicistronic operons (30). Whereas *groESL-1* is encoded in chromosome 1  
200 (*vc2664-vc2665*), *groESL-2* is located in chromosome 2 (*vca0819-0820*) (Fig 3A) (30). Based  
201 on our RNA-seq data, the latter manifested a larger fold change (Table 2). In order to  
202 confirm differential expression of these genes in  $\Delta vchM$ , we measured *groES-1* and *groES*-  
203 2 relative gene expression in exponential and stationary phase cells of WT and mutant  
204 strains, using digital qRT-PCR with the housekeeping *gyrA* gene as reference (31). The  
205 results confirm a higher induction of *groES-2* genes in both exponential and stationary  
206 phase  $\Delta vchM$  cells with a fold change (over the WT) of ca. 10X and 5X, respectively (Fig 3B).  
207 However, *groES-1* fold change was only slightly increased in exponential and unnoticeable  
208 in stationary phase.

209 Induction of *groESL* genes is usually associated to perturbations in proteostasis  
210 which leads to activation of the heat-shock response (32). Indeed, expression of both  
211 *groESL-1* and *groESL-2* is controlled by the heat-shock alternative sigma factor RpoH in *V.*

*cholerae* (33). However, the upregulation of *groESL-2* genes in  $\Delta vchM$  cells is likely independent of heat-shock activation as i) we do not observe any other genes of the heat-shock regulon being upregulated in the mutant (for example, *dnaKJ/grpE*, *clpB*, *ibpAB*) and ii) expression of *groESL-1* is not as increased as expression of *groESL-2* (Fig 3B).

Altogether, these results confirm that in absence of *VchM*, expression of *groESL-2* genes is markedly increased in *V. cholerae* and suggest that regulation of *groESL-2* operon in the  $\Delta vchM$  mutant can be independent of heat-shock response.

## **Deletion of *groESL-2* operon abolishes $\Delta vchM$ high tolerance to lethal doses of tobramycin**

In bacteria that harbor a single copy of this operon, GroESL are essential proteins for cell viability (34). However, possible redundancy between *groESL-1* and *groESL-2* could allow for the deletion of one or the other operon in *V. cholerae*. Thus, we attempted to delete *groESL-1* and *groESL-2* from *V. cholerae* WT and  $\Delta vchM$  strains. While  $\Delta groESL-2$  and  $\Delta vchM groESL-2$  strains were easily obtained, we could not manage to delete *groESL-1* in any background despite several attempts. Moreover, deletion of *groESL-2* did not affect the growth of *V. cholerae* in rich medium (Fig 4A). Respectively, GroES-1 and GroEL-1 share 80% and 87% amino acid identity with the only and essential GroES and GroEL proteins of *E. coli*, but lower (66% and 76%) amino acid identity with GroES-2 and GroEL-2 (S2 Fig). These observations suggest that i) GroESL-1 (but not GroESL-2) is essential for *V. cholerae* viability and ii) GroESL-1 is probably the main housekeeping chaperonin system while the divergent GroESL-2 could act synergistically in response to high levels of misfolding or having specific

substrates upon protein damage caused by specific stresses. Surprisingly, competition of  $\Delta groESL-2$  with a lacZ<sup>-</sup> strain shows that loss of these proteins is not detrimental for growth of *V. cholerae* in presence of subMIC TOB (S3A Fig). Similarly, survival of the  $\Delta groESL-2$  strain to lethal doses of TOB does not differ from that of the WT (S3B Fig). However, these genes are intrinsically highly expressed in  $\Delta vchM$  strain, where they may confer a selective advantage in presence of AG stress. In this case, deletion of *groESL-2* in  $\Delta vchM$  background would affect the mutant's tolerance. Indeed, when we compared the survival to lethal AG treatment of  $\Delta vchM$  to that of a  $\Delta vchM groESL-2$  double mutant, we found that the absence of *groESL-2* abolishes high tolerance to tobramycin and gentamicin in  $\Delta vchM$  (Fig 4B), without affecting growth in absence of stress (Fig 4A). These results show that the higher expression of *groESL-2* is required for the high tolerance of the  $\Delta vchM$  mutant to lethal AG treatment.

We then tested whether the high tolerance of this mutant relies on general higher levels of chaperonins or if it specifically linked to GroESL-2 chaperonins. We thus tried to complement the  $\Delta vchM groESL-2$  mutant by ectopically expressing *groESL-1* or *groESL-2* and assessed survival to lethal doses of AGs. Strikingly, only overexpression of *groESL-2* is able to promote survival levels similar to those observed in  $\Delta vchM$  (Fig 4C), suggesting a specific role for GroESL-2 in managing AG-mediated proteotoxic stress in *V. cholerae* cells lacking VchM. Interestingly, we observed no difference in the relative fitness of  $\Delta vchM$  and  $\Delta vchM groESL-2$  mutants in competitions in presence of subMIC doses of tobramycin, which shows that *groESL-2* it is not implicated in  $\Delta vchM$  higher relative fitness during growth in subMIC AGs (Fig 4D).

## 255 VchM controls *groESL-2* expression through direct DNA methylation

256           Knowing the role of VchM in regulating gene expression in *V. cholerae* (23), we asked  
 257 whether VchM controls *groESL* expression directly through DNA methylation. VchM  
 258 methylates the first cytosine in 5'-RCCGGY-3' motifs (16). This prompted us to search for  
 259 such motifs in both *groESL* operons. While we couldn't detect any of these sites along the  
 260 *groESL-1* locus, we found a total of four VchM motifs in *groESL-2* region: motif #1 within the  
 261 5' UTR of the operon, 47 bp away from the initiation codon; motif #2 is within the coding  
 262 region of *groES-2* while motifs #3 and #4 are located within the coding region of *groEL-2*  
 263 (Fig 5A). We hypothesized that the methylation state of these motifs could modulate the  
 264 transcription of *groESL-2* genes. To test this, we generated a mutant by replacing all RCCGGY  
 265 motifs in *groESL-2* region by non-consensus motifs but maintaining the amino acid  
 266 sequence of GroESL-2 proteins intact (Fig 5A, mut#1-4). Additionally, we created a mutant  
 267 where only the RCCGGY #1 was altered in order to investigate if this site, for being in the  
 268 regulatory region of this operon, had a stronger contribution in modulating gene expression  
 269 (Fig 5A, mut#1). We then measured *groES-2* expression in both mutants and observed that  
 270 disruption of RCCGGY #1 lead to a very weak increase in *groES-2* expression relative to the  
 271 WT, while disruption of all four sites led to a significantly higher expression of this gene (Fig  
 272 5B). We additionally tested *groEL-2* expression and observed similar results (S4 Fig).  
 273 Supporting our hypothesis that this regulation is methylation-dependent, we did not  
 274 observe any difference in *groES-2* or *groEL-2* expression when we mutated sites #1-4 in the  
 275  $\Delta vchM$  background (S4 Fig). It is worth mentioning that, in these experiments, the  
 276 expression of *groES-2* in the  $\Delta vchM$  strain was consistently higher than in the WT mut#1-4

(Fig 5B) suggesting that an additional factor, in synergy with the methylation of RCCGGY sites, may control expression of *groES-2*. Nonetheless, overall these results show that a specific set of chaperonin encoding genes is under the control of DNA cytosine methylation in *V. cholerae*, linking DNA methylation to modulation of chaperonin expression and tolerance to antibiotics.

## DISCUSSION

Antimicrobial resistance (AMR) is currently one of the biggest threats to global health (35). It is thus urgent not only to find new and alternative ways to fight bacterial infections but also to understand how bacteria adapt to the presence of antibiotics and study the molecular mechanisms they use to circumvent antibiotic action.

In this study, we establish a previously unknown link between VchM-mediated DNA methylation and aminoglycoside susceptibility in the human pathogen *V. cholerae*. VchM is a relatively understudied orphan DNA methyltransferase only found in *V. cholerae* species, known to methylate the first cytosine at 5'-RCCGGY-3' DNA motifs (16, 23). VchM is necessary for the optimal growth of *V. cholerae*, both *in vitro* and *in vivo*, and it was shown to repress the expression of a gene important for cell envelope stability through direct DNA methylation (23).

Here we show that despite the growth defect in stress-free medium, cells lacking VchM are also less susceptible to aminoglycoside toxicity. Specifically, we show that these

cells have a higher relative fitness in presence of low AG concentrations. The reason for this can be inferred from the growth curves in presence of subMIC TOB (Fig 1A) where it is clear that small increments in TOB concentration lead to a higher toxicity in the WT strain when compared to the *ΔvchM*. Moreover, even though the MIC values for the tested AGs are the same in both strains, *ΔvchM* displays a higher tolerance to lethal aminoglycoside treatment (Fig 2).

Aminoglycosides are a well-known class of antimicrobial drugs that cause disruption of the translation process and consequently protein misfolding (14, 36). The exact mechanism underlying the bactericidal activity of aminoglycosides has been subject of debate in the literature (37) but it is generally accepted that killing by AGs involves i) the uptake of the AG into the cytoplasm (11, 38, 39) and ii) membrane disruption mediated by insertion of misfolded proteins in the membrane as consequence of AG binding to the ribosomes and disruption of translational fidelity (11–13, 40). Indeed, mechanisms modulating aminoglycoside tolerance/resistance in different bacterial species (in exponential or stationary phase) have been shown to be associated either to AG uptake (41–44) or to translational fidelity and proteostasis (14, 29, 40, 45). Our results revealed a higher relative abundance of *groESL-2* transcripts in bacterial cells lacking VchM, which led us to hypothesize that such increased expression of these chaperonins could underlie the high tolerance to AGs observed in this mutant, as it had been previously observed in *E. coli* (29). In fact, we show that stationary phase cells lacking both *vchM* and *groESL-2* genes have similar or even lower tolerance to lethal AG treatment compared to the WT strain. However, we could not observe a significant increase in tolerance upon overexpression of *groESL-2* in

the WT strain (S5 Fig), suggesting that high *groESL-2* levels alone do not explain the high tolerance to lethal AG treatment. Instead, it is possible that high levels of GroESL-2 chaperone system counteract AG-mediated misfolding of specific substrates present only in cells devoid of VchM.

*V. cholerae* harbors two copies of *groESL* operon in its genome, thus belonging to the group of 30% of bacterial species that contains multiple copies of these chaperonins (46, 47). An interesting question to ask is whether these extra copies of chaperonins are functionally redundant or have a more specialized role in the cell, as it had been observed for *Myxococcus xanthus* (46–48). Supporting the latter hypothesis, we show here that the high tolerance observed in  $\Delta vchM$  is dependent on the high expression of *groESL-2* but not on the high expression of *groESL-1* (Fig 4B). Amino acid identity comparison between these proteins suggests that Vc GroESL-1 is likely the orthologue of the housekeeping GroESL of *E. coli* whereas Vc GroESL-2, thought to have appeared by duplication in *V. cholerae* (30), differ equally from both (S2 Fig). Thus, we speculate that *V. cholerae* GroESL-2 constitutes an alternative chaperone system capable of helping the folding of specific substrates important for survival to specific stresses. Interestingly, even though essential for the higher tolerance to lethal aminoglycoside treatment, *groESL-2* is not involved in the increased relative fitness of the  $\Delta vchM$  mutant in presence of subMIC doses of aminoglycosides (Fig 4C). This suggests that the mechanisms operating in  $\Delta vchM$  cells that increase their relative fitness during growth in subMIC AGs are not the same that increase their tolerance to lethal doses of these drugs. In fact, it has been recently shown that the type of translation errors occurring at lower streptomycin (another AG) concentrations differ from those found in

high concentrations of this aminoglycoside, with the latter being associated to a higher misfolding propensity (12). Thus, it seems plausible that, in  $\Delta vchM$  cells, the higher expression of *groESL-2* is likely to be more important at high concentrations of AGs, when the abundance of misfolded proteins tend to increase. The mechanisms driving  $\Delta vchM$  higher relative fitness at lower doses of aminoglycosides remain to be elucidated in future work.

DNA methylation controls gene expression through modulation of protein-DNA interactions (49). In most of the cases, the methylated base interferes with the binding of transcription factors and/or the RNA polymerase at the regulatory region of a gene, affecting transcription (50–52). However, there is also evidence that the presence of methylated DNA bases that occur along the coding region of genes could also directly affect their expression in bacteria, even though the precise mechanism is still unknown (20, 22, 23). In eukaryotes, cytosine methylation tends to repress gene expression. A recent study shedding light on how cytosine methylation affect DNA mechanical properties shows that cytosine methylation stabilizes the DNA helix and slows transcription in eukaryotic cells (53). Thus, a similar m5C-mediated transcriptional hindrance is likely to happen also in prokaryotes. Here we support this view by showing that abrogation of VchM-dependent methylation of cytosines at the four RCCGGY motifs in *groESL-2* region increased its expression in WT cells (Fig 5B). However, this is unlikely the sole mechanism responsible for the high expression of *groESL-2* genes in  $\Delta vchM$  cells, as this mutant has even higher expression levels of *groESL-2*. It is possible that the pleiotropic effects resulting from VchM

deficiency also affect, indirectly, the expression of these genes through regulation of a specific transcription factor.

Our work shows that a *V. cholerae* deletion mutant of the orphan DNA methyltransferase VchM have a general higher tolerance towards aminoglycosides. It remains to be explored whether *V. cholerae* WT cells can modulate VchM expression and, consequently, alter the levels of cytosine methylation. Bisulfite sequencing analysis of *V. cholerae* genome shows that all cytosines within RCCGGY motifs were methylated in *V. cholerae*, during exponential and stationary phases, with the exception of three of these sites which had been previously shown to be constantly undermethylated in this species (54). However, these studies were conducted in cells cultured in LB stress-free media or collected from frozen rabbit cecal fluid, and thus may not reflect the m5C profile of *V. cholerae* during other stress conditions. Moreover, bisulfite sequencing allows for cytosine methylation analysis of the total population at a specific time and thus it is not suitable to detect potential transient changes in small subpopulations of cells. Such changes could be mediated, for example, by altering the levels of VchM through gene expression. Little is known about *vchM* regulation but it was recently shown that the *V. cholerae* quorum sensing low density transcriptional regulator AphA is able to bind the *vchM* region (55) leaving the possibility that *vchM* may be regulated by quorum sensing. Moreover, *vchM* was previously found to be differentially expressed between different stages of human infection (56), suggesting the possibility that modulation of cytosine methylation levels can be adaptative during *V. cholerae*'s life cycle. In line with our work, lowering VchM levels could lead to a trade-off, where low m5C levels

would be detrimental for fitness in stress-free contexts, but highly advantageous in presence of specific stress conditions, such as antibiotic exposure.

## MATERIALS AND METHODS

### Strains, media and culture conditions

*V. cholerae* was routinely cultured at 37°C in Mueller-Hinton (MH) medium. Plasmids were introduced in *V. cholerae* by electrotransformation. Strains containing the pSC101 plasmid were grown in presence of 100 µg/mL carbenicillin for plasmid maintenance. All *V. cholerae* mutant strains are derived from *Vibrio cholerae* serotype O1 biotype El Tor strain N16961 hapR+. Mutants were constructed by homologous recombination after natural transformation or with a conjugative suicide plasmid as previously described (15, 57–59). Primers, strains and plasmids used in this study, and their constructions, are listed in Table S2. For routine cloning we used chemically competent *E. coli* One Shot® TOP10 (Invitrogen). All strains and plasmids were confirmed by sanger sequencing.

### Mutation of RCCGGY sites #1-4 in *groESL-2* region

In order to mutate all four RCCGGY sites present in *groESL-2* (*vca0819-0820*) region we generated a DNA fragment (S2 Table) with these sites containing the following nucleotide changes: #1- ACCGGC changed to ATCGGC; #2- ACCGGC changed to ACGGGC; #3- GCCGGC changed to GCGGGC and #4- ACCGGC changed to ACGGGC. This fragment was then

introduced in *V. cholerae* at the endogenous locus by allelic replacement as described in Table S2.

### **Growth curves**

Overnight cultures from single colonies were diluted 1:100 in Mueller-Hinton (MH) rich media or MH + subMIC antibiotics at different concentrations, in 96-well microplates. OD<sub>600</sub> was measured in a Tecan Infinite plate reader at 37°C, with agitation for 20 hours. Measurements were taken every 10 minutes.

### **MIC determination**

MICs were determined by microtiter broth dilution method (60) with an initial inoculum size of 10<sup>5</sup> CFUs/mL. The MIC was interpreted as the lowest antibiotic concentration preventing visible growth.

### **Neo-Cy5 uptake**

Quantification of fluorescent neomycin (Neo-cy5) uptake was performed as described (61). Neo-cy5 is an aminoglycoside coupled to the fluorophore Cy5, and has been shown to be active against Gram- bacteria (27). Briefly, overnight cultures were diluted 100-fold in rich MOPS (Teknova EZ rich defined medium). When the bacterial cultures reached an OD<sub>600</sub> of

0.25, they were incubated with 0.4  $\mu$ M of Cy5 labeled Neomycin for 15 minutes at 37°C. 10  $\mu$ L of the incubated culture were then used for flow cytometry, diluting them in 250  $\mu$ L of PBS before reading fluorescence. Flow cytometry experiments were performed as described (62). For each experiment, 100000 events were counted on the Miltenyi MACSquant device.

### Competitions experiments

Overnight cultures from single colonies of lacZ<sup>-</sup> and lacZ<sup>+</sup> strains were washed in PBS (Phosphate Buffer Saline) and mixed 1:1 (500 $\mu$ L + 500 $\mu$ L). At this point 100 $\mu$ L of the mix were serial diluted and plated in MH agar supplemented with X-gal (5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside) at 40  $\mu$ g/mL to assess T0 initial 1:1 ratio. At the same time, 10  $\mu$ L from the mix were added to 2 mL of MH or MH supplemented with subMIC tobramycin at 0.6 $\mu$ g/mL and incubated with agitation at 37°C for 20 hours. Cultures were then diluted and plated in MH agar plates supplemented with X-gal. Plates were incubated overnight at 37°C and the number of blue and white CFUs was assessed. Competitive index was calculated by dividing the number of blue CFUs (lacZ<sup>+</sup> strain) by the number of white CFUs (lacZ<sup>-</sup> strain).

### Survival assays

Bacterial cultures from single colonies were cultured at 37°C for 16 h with agitation in 10 mL of MH medium. Aliquots from these cultures were removed, serial diluted and plated in MH agar plates to assess CFUs formation prior antibiotic treatment (T0). In addition, 5 mL

of these aliquots were subjected to antibiotic treatment and incubated with agitation at 37°C. At the indicated time points, 500uL of these cultures were collected, washed in PBS, serial diluted and plated in MH agar plates. The plates were then incubated overnight at 37°C. Survival at each time point was determined by dividing the number of CFUs/mL at that time point by the number of CFUs/mL prior treatment. Antibiotics were used at the following final concentrations: 20 µg/mL Tobramycin (TOB) and 10 µg/mL Gentamicin (GEN). Experiments were repeated at least two to three times.

## Digital qRT-PCR

For RNA extraction, overnight cultures of three biological replicates of strains of interest were diluted 1:1000 in MH media and grown with agitation at 37°C until an OD<sub>600</sub> of 0.3 (exponential phase) or an OD<sub>600</sub> of 1.0 or 2.0 (stationary phase). 0.5 mL of these cultures were centrifuged and supernatant removed. Pellets were homogenized by resuspension with 1.5 mL of cold TRIzol™ Reagent. Next, 300 µL chloroform were added to the samples following mix by vortexing. Samples were then centrifuged at 4°C for 10 minutes. Upper (aqueous) phase was transferred to a new 2mL tube and mixed with 1 volume of 70% ethanol. From this point, the homogenate was loaded into a RNeasy Mini kit (Qiagen) column and RNA purification proceeded according to the manufacturer's instructions. Samples were then subjected to DNase treatment using TURBO DNA-free Kit (Ambion) according to the manufacturer's instructions. RNA concentration of the samples was

measured with NanoDrop™ spectrophotometer and diluted to a final concentration of 1-10 ng/μL.

qRT-PCR reactions were prepared with 1 μL of diluted RNA samples using the qScript™ XLT 1-Step RT-qPCR ToughMix (Quanta Biosciences, Gaithersburg, MD, USA) within Sapphire chips. Digital PCR was conducted on a Naica Geode (programmed to perform the sample partitioning step into droplets, followed by the thermal cycling program suggested in the user's manual. Primer and probe sequences used in digital qRT-PCR reaction are listed in Table S3. Image acquisition was performed using the Naica Prism3 reader. Images were then analyzed using Crystal Reader software (total droplet enumeration and droplet quality control) and the Crystal Miner software (extracted fluorescence values for each droplet). Values were normalized against expression of the housekeeping gene *gyrA* as previously described (31).

## RNA-seq

For RNA extraction, overnight cultures of three biological replicates of WT and *ΔvchM* strains were diluted 1:100 in MH medium and grown with agitation at 37°C until cultures reach an OD<sub>600</sub> of 2.0. Total RNA extraction, library preparation, sequencing and analysis were performed as previously described (63). The data for this RNA-seq study has been submitted in the GenBank Sequence Read Archive (SRA) under project number PRJNA509113.

## ACKNOWLEDGMENTS

We are thankful to Manon Lang for her valuable help with the neo-cy5 uptake experiments and Sebastian Aguilar Pierlé for all the help with RNAseq analysis. We also thank Evelyne Krin for help with molecular cloning procedures and João Gama for helpful comments on the manuscript.

## FIGURE CAPTIONS

**Fig 1. *V. cholerae* N16961  $\Delta vchM$  is less susceptible to subMIC aminoglycosides** **A.** Growth curves in absence (MH) or presence of subMIC doses of tobramycin. Bars represent SD (n=3) **B.** *In vitro* competitions of WT and mutant strains against isogenic  $\Delta lacZ$  reference strain in absence or presence of different antibiotics at subMIC concentrations (TOB, 0.6  $\mu\text{g/ml}$ ; GEN, 0.5  $\mu\text{g/ml}$ ; NEO, 2.0  $\mu\text{g/ml}$ ; CAM, 0.4  $\mu\text{g/ml}$ ; CARB, 2.5  $\mu\text{g/ml}$ ). Box plots indicate the median and the 25<sup>th</sup> and 75<sup>th</sup> percentiles; whiskers indicate the min and max values (n=6).

**Fig 2.  $\Delta vchM$  strain is more tolerant to lethal aminoglycoside treatment.** Survival of stationary-phase WT and  $\Delta vchM$  cells exposed to lethal doses of tobramycin (TOB) **(A)**, and gentamicin (GEN) **(B)**. Survival represents the number of bacteria (CFU/mL) after treatment divided by the initial number of bacteria prior treatment. Means and SD are represented, n=3.

**Fig 3. *groESL-2* operon is upregulated in  $\Delta vchM$  strain.** **A.** Schematic representation of both *groESL* operons in *V. cholerae*. The four RCCGGY sites present along the *groESL-2* region are represented by the inverted orange triangles. **B.** Fold change ( $\Delta vchM$ /WT) of the relative

expression levels of *groES-1* and *groES-2* in cultures at exponential phase (Exp, OD<sub>600</sub> ≈ 0.3) or stationary phase (Stat, OD<sub>600</sub> ≈ 1.8-2.0). Means and SD are represented, n=3.

**Fig 4. *groESL-2* is needed for the increased tolerance of  $\Delta vchM$  to lethal AG treatment. A.**

Growth curves in MH medium. Means and SD are represented, n=3. **B.** Survival of stationary-phase WT and  $\Delta groESL-2$  cells exposed to lethal aminoglycoside treatment for 7 hours. Box plots indicate the median and the 25<sup>th</sup> and 75<sup>th</sup> percentiles; whiskers indicate the min and max values (n=6 from two independent experiments). **C.** Survival (after 7 hours AG treatment) of  $\Delta vchM groESL-2$  double mutant harboring an empty plasmid or a plasmid expressing either *groESL-1* or *groESL-2*, relative to survival of the  $\Delta vchM$  with the control plasmid. Box plots indicate the median and the 25<sup>th</sup> and 75<sup>th</sup> percentiles; whiskers indicate the min and max values (n=6 from two independent experiments). In **B** and **C** statistically significant differences were determined using Friedman's test with Dunn's post-hoc test for multiple comparisons. \* P<0.05, \*\* P<0.01, ns = not significant. **D.** *In vitro* competitions of  $\Delta vchM$  and  $\Delta vchM groESL-2$  double mutant strains against isogenic  $\Delta lacZ$  reference strain in absence or presence of subMIC TOB, 0.6 µg/ml; Error bars indicate SD (n=6).

**Fig 5. Disrupted VchM sites in *groESL-2* region leads to increased gene expression in the**

**WT. A.** Schematic representation of mutants with abrogated VchM sites. **B.** Relative expression of *groES-2* in the different strains at OD<sub>600</sub> of 1.0. Box plots indicate the median and the 25<sup>th</sup> and 75<sup>th</sup> percentiles; whiskers indicate the min and max values (n= 5). Statistical significance was determined by Kruskal-Wallis test with Dunn's post-hoc test for multiple comparisons. \* P<0.05, \*\* P<0.01, ns = not significant

## REFERENCES

1. Liu YC, Huang WK, Huang TS, Kunin CM. 1999. Detection of antimicrobial activity in urine for epidemiologic studies of antibiotic use. *J Clin Epidemiol*.
2. Haggard BE, Bartsch LD. 2009. Net Changes in Antibiotic Concentrations Downstream from an Effluent Discharge. *J Environ Qual*.
3. Fick J, Söderström H, Lindberg RH, Phan C, Tysklind M, Larsson DGJ. 2009. Contamination of surface, ground, and drinking water from pharmaceutical production. *Environ Toxicol Chem*.
4. Andersson DI, Hughes D. 2014. Microbiological effects of sublethal levels of antibiotics. *Nat Rev Microbiol*.
5. Davies J, Spiegelman GB, Yim G. 2006. The world of subinhibitory antibiotic concentrations. *Curr Opin Microbiol* 9:445–453.
6. Wistrand-Yuen E, Knopp M, Hjort K, Koskiniemi S, Berg OG, Andersson DI. 2018. Evolution of high-level resistance during low-level antibiotic exposure. *Nat Commun*.
7. Jørgensen KM, Wassermann T, Jensen PØ, Hengzuang W, Molin S, Høiby N, Ciofu O. 2013. Sublethal ciprofloxacin treatment leads to rapid development of high-level ciprofloxacin resistance during long-term experimental evolution of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*.
8. Gullberg E, Cao S, Berg OG, Ilbäck C, Sandegren L, Hughes D, Andersson DI. 2011. Selection of resistant bacteria at very low antibiotic concentrations. *PLoS Pathog*.

- 547 9. Baharoglu Z, Mazel D. 2011. *Vibrio cholerae* triggers SOS and mutagenesis in  
548 response to a wide range of antibiotics: A route towards multiresistance. *Antimicrob*  
549 *Agents Chemother.*
- 550 10. Baharoglu Z, Babosan A, Mazel D. 2014. Identification of genes involved in low  
551 aminoglycoside-induced SOS response in *Vibrio cholerae*: A role for transcription  
552 stalling and Mfd helicase. *Nucleic Acids Res.*
- 553 11. Davis BD. 1987. Mechanism of bactericidal action of aminoglycosides. *Microbiol Rev.*
- 554 12. Wohlgemuth I, Garofalo R, Samatova E, Günenç AN, Lenz C, Urlaub H, Rodnina M V.  
555 2021. Translation error clusters induced by aminoglycoside antibiotics. *Nat Commun.*
- 556 13. Kohanski MA, Dwyer DJ, Wierzbowski J, Cottarel G, Collins JJ. 2008. Mistranslation of  
557 Membrane Proteins and Two-Component System Activation Trigger Antibiotic-  
558 Mediated Cell Death. *Cell* 135:679–690.
- 559 14. Ling J, Cho C, Guo LT, Aerni HR, Rinehart J, Söll D. 2012. Protein Aggregation Caused  
560 by Aminoglycoside Action Is Prevented by a Hydrogen Peroxide Scavenger. *Mol Cell*  
561 48:713–722.
- 562 15. Negro V, Krin E, Pierlé SA, Chaze T, Gianetto QG, Kennedy SP, Matondo M, Mazel D,  
563 Baharoglu Z. 2019. RadD contributes to R-Loop avoidance in Sub-MIC tobramycin.  
564 *MBio.*
- 565 16. Banerjee S, Chowdhury R. 2006. An orphan {DNA} (cytosine-5-)-methyltransferase in  
566 *Vibrio cholerae* 152:1055–1062.

- 567 17. Fujimoto D, Srinivasan PR, Borek E. 1965. On the Nature of the Deoxyribonucleic Acid  
568 Methylases. Biological Evidence for the Multiple Nature of the Enzymes.  
569 Biochemistry.
- 570 18. Casadesús J, Low D. 2006. Epigenetic Gene Regulation in the Bacterial World.  
571 Microbiol Mol Biol Rev.
- 572 19. Sánchez-Romero MA, Casadesús J. 2020. The bacterial epigenome. Nat Rev  
573 Microbiol.
- 574 20. Kumar S, Karmakar BC, Nagarajan D, Mukhopadhyay AK, Morgan RD, Rao DN. 2018.  
575 N4-cytosine DNA methylation regulates transcription and pathogenesis in  
576 Helicobacter pylori. Nucleic Acids Res.
- 577 21. Estibariz I, Overmann A, Ailloud F, Krebs J, Josenhans C, Suerbaum S. 2019. The core  
578 genome m5C methyltransferase JHP1050 (M.Hpy99III) plays an important role in  
579 orchestrating gene expression in Helicobacter pylori. Nucleic Acids Res 47:2336–  
580 2348.
- 581 22. Militello KT, Simon RD, Qureshi M, Maines R, van Horne ML, Hennick SM, Jayakar SK,  
582 Pounder S. 2012. Conservation of Dcm-mediated cytosine DNA methylation in  
583 Escherichia coli. FEMS Microbiol Lett.
- 584 23. Chao MC, Zhu S, Kimura S, Davis BM, Schadt EE, Fang G, Waldor MK. 2015. A Cytosine  
585 Methyltransferase Modulates the Cell Envelope Stress Response in the Cholera  
586 Pathogen 11:e1005666.

- 587 24. Brauner A, Fridman O, Gefen O, Balaban NQ. Distinguishing between resistance,  
588 tolerance and persistence to antibiotic treatment. Nat Publ Gr.
- 589 25. Haugan MS, Løbner-Olesen A, Frimodt-Møller N. 2019. Comparative activity of  
590 ceftriaxone, ciprofloxacin, and gentamicin as a function of bacterial growth rate  
591 probed by Escherichia coli chromosome replication in the mouse peritonitis model.  
592 Antimicrob Agents Chemother.
- 593 26. Nakae R, Nakae T. 1982. Diffusion of aminoglycoside antibiotics across the outer  
594 membrane of Escherichia coli. Antimicrob Agents Chemother.
- 595 27. Sabeti Azad M, Okuda M, Cyrenne M, Bourge M, Heck MP, Yoshizawa S, Fourmy D.  
596 2020. Fluorescent Aminoglycoside Antibiotics and Methods for Accurately  
597 Monitoring Uptake by Bacteria. ACS Infect Dis.
- 598 28. Hartl FU, Hayer-Hartl M. 2002. Protein folding. Molecular chaperones in the cytosol:  
599 From nascent chain to folded protein. Science (80- ) 295:1852–1858.
- 600 29. Goltermann L, Good L, Bentin T. 2013. Chaperonins fight aminoglycoside-induced  
601 protein misfolding and promote short-term tolerance in Escherichia coli. J Biol Chem.
- 602 30. Chowdhury N, Kingston JJ, Brian Whitaker W, Carpenter MR, Cohen A, Fidelma Boyd  
603 E. 2014. Sequence and expression divergence of an ancient duplication of the  
604 chaperonin groESEL operon in Vibrio species. Microbiol (United Kingdom) 160:1953–  
605 1963.
- 606 31. Lo Scrudato M, Blokesch M. 2012. The regulatory network of natural competence

607 and transformation of *Vibrio cholerae*. PLoS Genet.

608 32. Richter K, Haslbeck M, Buchner J. 2010. The Heat Shock Response: Life on the Verge  
609 of Death. Mol Cell.

610 33. Slamti L, Livny J, of bacteriology WMK. 2007. Global gene expression and phenotypic  
611 analysis of a *Vibrio cholerae* {rpoH} deletion mutant.

612 34. Fayet O, Ziegelhoffer T, Georgopoulos C. 1989. The groES and groEL heat shock gene  
613 products of *Escherichia coli* are essential for bacterial growth at all temperatures. J  
614 Bacteriol 171:1379–1385.

615 35. O’Neill J. 2015. Tackling a Crisis for the Health and Wealth of Nations. Rev Antimicrob  
616 Resist.

617 36. Magnet S, Blanchard JS. 2005. Molecular insights into aminoglycoside action and  
618 resistance. Chem Rev 105:477–497.

619 37. Baquero F, Levin BR. 2021. Proximate and ultimate causes of the bactericidal action  
620 of antibiotics. Nat Rev Microbiol.

621 38. Ezraty B, Vergnes A, Banzhaf M, Duverger Y, Huguenot A, Brochado AR, Su SY,  
622 Espinosa L, Loiseau L, Py B, Typas A, Barras F. 2013. Fe-S cluster biosynthesis controls  
623 uptake of aminoglycosides in a ROS-less death pathway. Science (80- ).

624 39. Bruni GN, Kralj JM. 2020. Membrane voltage dysregulation driven by metabolic  
625 dysfunction underlies bactericidal activity of aminoglycosides. Elife.

626 40. Zou J, Zhang W, Zhang H, Zhang XD, Peng B, Zheng J. 2018. Studies on aminoglycoside

627           susceptibility identify a novel function of KsgA to secure translational fidelity during  
628           antibiotic stress. *Antimicrob Agents Chemother* 62:1–13.

629   41.   Shan Y, Lazinski D, Rowe S, Camilli A, Lewis K. 2015. Genetic basis of persister  
630           tolerance to aminoglycosides in *Escherichia coli*. *MBio* 6:1–10.

631   42.   Allison KR, Brynildsen MP, Collins JJ. 2011. Metabolite-enabled eradication of  
632           bacterial persisters by aminoglycosides. *Nature*.

633   43.   McKay SL, Portnoy DA. 2015. Ribosome hibernation facilitates tolerance of  
634           stationary-phase bacteria to aminoglycosides. *Antimicrob Agents Chemother*  
635           59:6992–6999.

636   44.   Hall CW, Farkas E, Zhang L, Mah TF. 2019. Potentiation of Aminoglycoside Lethality  
637           by C4-Dicarboxylates Requires RpoN in Antibiotic-Tolerant *Pseudomonas aeruginosa*.  
638           *Antimicrob Agents Chemother*.

639   45.   Ji X, Zou J, Peng H, Stolle AS, Xie R, Zhang H, Peng B, Mekalanos JJ, Zheng J. 2019.  
640           Alarmone Ap4A is elevated by aminoglycoside antibiotics and enhances their  
641           bactericidal activity. *Proc Natl Acad Sci U S A*.

642   46.   Kumar CMS, Mande SC, Mahajan G. 2015. Multiple chaperonins in bacteria—novel  
643           functions and non-canonical behaviors. *Cell Stress Chaperones* 20:555–574.

644   47.   Goyal K, Qamra R, Mande SC. 2006. Multiple gene duplication and rapid evolution in  
645           the groEL Gene: Functional implications. *J Mol Evol*.

646   48.   Wang Y, Zhang W yan, Zhang Z, Li J, Li Z feng, Tan Z gao, Zhang T tian, Wu Z hong, Liu

647 H, Li Y zhong. 2013. Mechanisms Involved in the Functional Divergence of Duplicated  
648 GroEL Chaperonins in *Myxococcus xanthus* DK1622. *PLoS Genet*.

649 49. Casadesus J, Low D. 2006. Epigenetic Gene Regulation in the Bacterial World.  
650 *Microbiol Mol Biol Rev*.

651 50. Hernday AD, Braaten BA, Low DA. 2003. The mechanism by which DNA adenine  
652 methylase and PapI activate the Pap epigenetic switch. *Mol Cell*.

653 51. Cota I, Sánchez-Romero MA, Hernández SB, Pucciarelli MG, García-Del Portillo F,  
654 Casadesús J. 2015. Epigenetic Control of *Salmonella enterica* O-Antigen Chain Length:  
655 A Tradeoff between Virulence and Bacteriophage Resistance. *PLoS Genet*.

656 52. Wion D, Casadesús J. 2006. N6-methyl-adenine: an epigenetic signal for DNA-protein  
657 interactions. *Nat Rev Microbiol*.

658 53. Rausch C, Zhang P, Casas-Delucchi CS, Daiß JL, Engel C, Coster G, Hastert FD, Weber  
659 P, Cardoso MC. 2021. Cytosine base modifications regulate DNA duplex stability and  
660 metabolism. *Nucleic Acids Res*.

661 54. Dalia AB, Lazinski DW, Camilli A. 2013. Characterization of undermethylated sites in  
662 *vibrio cholerae*. *J Bacteriol*.

663 55. Haycocks JRJ, Warren GZL, Walker LM, Chlebek JL, Dalia TN, Dalia AB, Grainger DC.  
664 2019. The quorum sensing transcription factor AphA directly regulates natural  
665 competence in *Vibrio cholerae*. *PLoS Genet*.

666 56. LaRocque RC, Harris JB, Dziejman M, Li X, Khan AI, Faruque ASG, Faruque SM, Nair

667 GB, Ryan ET, Qadri F, Mekalanos JJ, Calderwood SB. 2005. Transcriptional profiling of  
668 *Vibrio cholerae* recovered directly from patient specimens during early and late  
669 stages of human infection. Infect Immun.

670 57. Baharoglu Z, Krin E, Mazel D. 2012. Connecting environment and genome plasticity  
671 in the characterization of transformation-induced {SOS} regulation and carbon  
672 catabolite control of the *Vibrio cholerae* integron integrase. J Bacteriol 194:1659–  
673 1667.

674 58. Baharoglu Z, Krin E, Mazel D. 2013. RpoS Plays a Central Role in the SOS Induction by  
675 Sub-Lethal Aminoglycoside Concentrations in *Vibrio cholerae*. PLoS Genet.

676 59. Val ME, Skovgaard O, Ducos-Galand M, Bland MJ, Mazel D. 2012. Genome  
677 engineering in *Vibrio cholerae*: A feasible approach to address biological issues. PLoS  
678 Genet.

679 60. Name P, Date R. 2010. Methods for Dilution Antimicrobial Suceptibility tests for  
680 Bacteria that Grow Aerobically; Approved Standard- Seventh Edition. Clin Lab Stand  
681 Inst.

682 61. Lang M, Krin E, Korlowski C, Sismeiro O, Varet H, Coppée J-Y, Mazel D, Baharoglu Z.  
683 2020. Sleeping ribosomes: bacterial signaling triggers RaiA mediated persistence to  
684 aminoglycosides. bioRxiv 2020.11.27.401281.

685 62. Baharoglu Z, Bikard D, Mazel D. 2010. Conjugative {DNA} transfer induces the  
686 bacterial {SOS} response and promotes antibiotic resistance development through  
687 integron activation. {PLoS} Genet 6:e1001165.

63. Krin E, Pierlé SA, Sismeiro O, Jagla B, Dillies MA, Varet H, Irazoki O, Campoy S, Rouy Z, Cruveiller S, Médigue C, Coppée JY, Mazel D. 2018. Expansion of the SOS regulon of *Vibrio cholerae* through extensive transcriptome analysis and experimental validation. BMC Genomics.

## SUPPORTING INFORMATION

**S1 Table. Differentially regulated genes in  $\Delta vchM$  strain**

**S2 Table. Strains, plasmids and primers used in this study**

**S3 Table. Primer and probe sequences used in digital qRT-PCR**

**S1 Fig. Neo-cy5 uptake is not increased in  $\Delta vchM$ .** Percentage of neo-cy5 positive cells analyzed by flow cytometry after incubation with fluorescent marked neomycin. Means and SD are represented, n=3.

**S2 Fig. Comparison of GroESL proteins from *E. coli* and *V. cholerae*.** Amino acid identity between GroES and GroEL proteins of *E. coli* MG1655 (*Eco*) and *V. cholerae* O1 El Tor N16961 (*Vch*) computed by BLASTP. Values represent percentage identity between proteins.


**S3 Fig. Deletion of *groESL-2* does not increase susceptibility to tobramycin. A.** *In vitro* competitions of WT and  $\Delta groESL-2$  strains against isogenic  $\Delta lacZ$  reference strain in absence or presence of tobramycin (TOB) at 0.6  $\mu\text{g/ml}$ ; n=3, error bars indicate SD. **B.** Survival of

stationary-phase WT and  $\Delta groESL-2$  cells exposed to 20X MIC of tobramycin. n=3, error bars indicate SD.

**S4 Fig. Mutation of VchM sites in *groESL-2* region fails to affect gene expression of the operon in absence of VchM.** Relative expression of *groES-2* and *groEL-2* genes in the indicated strains grown at OD<sub>600</sub> 1.0. n=3, error bars indicate SD.

**S5 Fig. Overexpression of *groESL-2* in the WT does not increase tolerance to tobramycin.** Survival (after 7 hours TOB treatment) of WT strain carrying a control plasmid or a plasmid overexpressing *groESL-2* genes. n=6, error bars indicate SD.


718 **Figure 1.**



719

720 **Figure 2.**


721




722

723


**Figure 3**



**Figure 4**



747 **Figure 5**



748