

1 **A B73 x Palomero Toluqueño mapping population reveals local adaptation in** 2 **Mexican highland maize**

3 Sergio Perez-Limón^{1,2}, Meng Li², G. Carolina Cintora-Martinez^{1,3}, M Rocio Aguilar-Rangel^{1,4},
4 M. Nancy Salazar-Vidal^{1,5}, Eric González-Segovia^{1,6}, Karla Blöcher-Juárez^{1,7}, Alejandro
5 Guerrero-Zavala¹, Benjamin Barrales-Gamez¹, Jessica Carcaño-Macias¹, Denise E. Costich⁸,
6 Jorge Nieto-Sotelo⁹, Octavio Martinez de la Vega¹, June Simpson¹, Matthew B. Hufford¹⁰,
7 Jeffrey Ross-Ibarra¹¹, Sherry Flint-Garcia⁵, Luis Diaz-Garcia¹², Rubén Rellán-Álvarez^{1, 13},
8 Ruairidh J. H. Sawers^{1,2*}

9 ¹ Laboratorio Nacional de Genómica para la Biodiversidad/Unidad de Genómica Avanzada,
10 Centro de Investigación y de Estudios Avanzados, Instituto Politécnico Nacional (CINVESTAV-
11 IPN), Irapuato, C.P. 36821, Guanajuato, México.

12 ² Department of Plant Science, The Pennsylvania State University, State College, PA, USA.

13 ³present address: Max Planck Institute for Plant Breeding Research, Cologne 50829 Germany.

14 ⁴present address: Corteva Agriscience, Tlajomulco de Zúñiga, Jalisco, México, 45560

15 ⁵present address: U.S. Department of Agriculture - Agricultural Research Service Plant Genetics
16 Research Unit, University of Missouri, Columbia, MO, United States

17 ⁶ Department of Botany, University of British Columbia, Vancouver, British Columbia, Canada.

18 ⁷present address: Institute of Molecular Biology (IMB), Mainz 55128, Germany.

19 ⁸ Institute for Genomic Diversity, Cornell University, Ithaca, NY

20 ⁹ Jardín Botánico, Instituto de Biología, Universidad Nacional Autónoma de México, Coyoacán,
21 C.P. 04510, Ciudad de México, México.

22 ¹⁰ Department of Ecology, Evolution, and Organismal Biology, Iowa State University

23 ¹¹ Department of Evolution and Ecology, Center for Population Biology, and Genome Center,

24 UC Davis

25 ¹² Campo Experimental Pabellón-INIFAP. Carretera Aguascalientes-Zacatecas, km. 32.5. 20660.

26 Pabellón de Arteaga, Aguascalientes, México

27 ¹³ Department of Molecular and Structural Biochemistry, North Carolina State University

28

29 * Author for correspondence: rjs6686@psu.edu

30

31

32

ABSTRACT

Generations of farmer selection have produced a unique collection of traditional maize varieties adapted to the environmental challenges of the central Mexican highlands. In addition to agronomic and cultural value, Mexican highland maize represents a good system for the study of local adaptation and acquisition of adaptive phenotypes under cultivation. In this study, we characterized a recombinant inbred line population derived from the cross of the B73 reference line and the Mexican highland maize variety Palomero Toluqueño. Evaluation over multiple years in lowland and highland field sites in Mexico identified genomic regions linked to yield components and putatively adaptive morphological traits. A region on chromosome 7 associated with ear weight showed antagonistic allelic effects in lowland and highland fields, suggesting a trade-off consistent with local adaptation. We identified several alleles of highland origin associated with characteristic highland traits, including reduced tassel branching, increased stem pigmentation and the presence of stem macrohairs. The oligogenic architecture of characteristic morphological traits supports their role in adaptation, suggesting they have arisen from consistent directional selection acting at distinct points across the genome. We discuss these results in the context of the origin of phenotypic novelty during selection, commenting on the role of *de novo* mutation and the acquisition of adaptive variation by gene flow from endemic wild relatives.

INTRODUCTION

Climatic trends and a need to reduce the level of agronomic inputs have fostered interest in the development of crop varieties that are not only high yielding but *resilient* - *i.e.* their performance is stable in the face of diverse, and potentially unpredictable, environmental challenges. One approach to enhance the cultivated genepool for greater resilience is to explore diversity at the extremes of a crop's distribution (Emon *et al.*, 2015; Dwivedi *et al.*, 2016; Corrado & Rao, 2017; Sousaraei *et al.*, 2021). Thousands of years of effort and care by the world's traditional farming communities have generated a rich diversity of landrace varieties, collectively adapted to a far broader ecological range than modern breeding material (Bellon *et al.*, 2018). Crop landraces serve to illustrate the mechanisms whereby plants can adapt to environmental stress as well as representing a valuable source of adaptive variation in their own right.

Strong directional selection imposed by prevailing conditions tends to produce highly specialized forms that perform well in their home environment, but relatively poorer in other locations, a process referred to as *local adaptation*. Local adaptation is defined formally as superior performance of local genotypes in their native environment versus non-local genotypes (Anderson *et al.*, 2013; Mitchell-Olds *et al.*, 2007; Hall *et al.*, 2010; Anderson *et al.*, 2011). Concomitantly, the average performance of a locally adapted variety over a range of environments may be poorer than that of a generalist that maintains a reasonable level of performance in all environments - *stability* in the context of plant breeding. Experimentally, the best demonstration of local adaptation is the *reciprocal transplant* experiment, in which varieties of interest are evaluated in a series of common gardens covering the range of their home environments. Both local adaptation and stability can be described with reference to *genotype x environment interaction* (GEI), *i.e.* the degree to which the relative performance of a given

variety compared with others depends on environmental conditions (Juenger, 2013; Assmann, 2013; Scheiner, 1993; El-Soda *et al.*, 2014).

By definition, all varieties will likely suffer reduced performance when challenged by environmental stress. GEI describes variety-specific deviations from the environmental main effect: some varieties suffer more than average, while others are better able to mitigate the impact of the stress. In extreme cases, the relative performance of varieties changes between environments, a scenario referred to as *rank changing* GEI. While stress is often considered with respect to a single sub-optimal factor, the same framework applies equally to the complex pattern of challenges presented by different localities. It can be seen that rank changing GEI underpins local adaptation, as defined above. With the advent of comparative genomics and greater understanding of the physiology and cell biology of environmental responses, it has become feasible to begin to characterize the genetic basis of local adaptation (Lovell *et al.*, 2021). Two principal modes of gene action have been proposed to drive rank changing GEI, namely *conditional neutrality* and *antagonistic pleiotropy*. Under conditional neutrality, a given genetic variant is linked to phenotypic change in some environments but not others. A complementary suite of conditionally neutral loci would, theoretically, be sufficient to generate rank changing GEI. Under antagonistic pleiotropy, the sign of the effect of a given variant changes between environments, *e.g.* a beneficial allele in one environment becomes deleterious in another, with a behaviour at a single variant that directly mirrors the whole genotype pattern of GEI. In practice, both behaviors will typically contribute to GEI and, indeed, classification of any given variant will be specific to the environments under consideration (Fournier-Level *et al.*, 2011). In addition, distinguishing conditional neutrality from antagonistic pleiotropy may be limited by statistical power in any given design. To date, studies of local adaptation in wild barley *Hordeum*

spontaneum (Verhoeven et al., 2004; Verhoeven et al., 2008), the annual grass *Avena barbata* (Gardner & Latta, 2006; Latta et al., 2007; Latta et al., 2010), the model plant *Arabidopsis thaliana* (Weinig et al., 2003; Fournier-Level et al., 2011) and monkey flower *Mimulus guttatus* (Hall et al., 2010; Lowry et al., 2009) have predominantly found cases of conditional neutrality. That said, examples of antagonistic pleiotropy do exist, although mostly limited to plant model organisms such as *Arabidopsis thaliana* (Scarcelli et al., 2007; Todesco et al., 2010), monkey flower *Mimulus guttatus* (Hall et al., 2010) and *Boechera stricta* (Anderson et al., 2013). An important consequence of the genetic architecture of local adaptation is the degree to which the specialist is constrained by trade-offs that impose an unavoidable cost of poor performance outside of the home environment. In terms of plant breeding, there are analogous implications with regard to how extensively a given variety can be used and how robust yields will be in the face of unpredictable or changing environmental conditions.

In addition to their intrinsic value, crop landraces provide an excellent system to study local adaptation, especially with regard to the rapid change required over the relatively short time frame of domestication. Landraces are dynamic populations, each with a unique identity shaped by biotic and abiotic stresses, crop management, seed handling and consumer preferences. As such, landraces are the product of both direct and indirect farmer selection, natural selection in the face of the local environment and exchange through traditional seed flow networks (Louette et al., 1997; Cleveland & Soleri, 2007; Mercer et al., 2008; Mercer & Perales, 2019). Typically, they are cultivated under low-input conditions and produce a modest but stable yield (Zeven, 1998; Breseghello & Coelho, 2013; Dwivedi et al., 2016). The sustained association of a given landrace population with a given locality results in local adaptation, in the same way it is seen in wild populations, demonstrable by reciprocal transplantation (Janzen et al., 2021).

Maize (*Zea mays* ssp. *mays*) was domesticated from balsas teosinte (*Zea mays* subsp. *parviglumis*) (Matsuoka *et al.*, 2002), about 9000 years ago, in the basin of the Balsas River in Mexico (Piperno *et al.*, 2007). After domestication, maize dispersed and was successfully established in different environments throughout the Americas and, eventually, across the world. In Mexico alone, 59 different native landraces of maize have been described, grown from sea level to 3400 m.a.s.l., in a range of environments, from semi-desert to regions with high humidity and temperature (CONABIO, 2018). One of the environments colonized by early maize was the central highlands of Mexico. The central Mexican highlands are characterized by low atmospheric pressure and temperature, high UV-B radiation, seasonal precipitation, presence of early frosts and low phosphorus availability due to the volcanic origin of the soil (Bellon *et al.*, 2005; Körner, 2007; Mercer *et al.*, 2008; Espinosa-Calderón *et al.*, 2011; Galván-Tejada *et al.*, 2014). Previous work has highlighted the impact of low temperature on unadapted maize varieties, which when grown at low temperatures and exposed to high light intensity, suffer metabolic lesions in chlorophyll synthesis, leading to increased photodamage and chlorophyll turnover (McWilliam & Naylor, 1967). Interestingly, these cold stress-induced symptoms were not observed in highland maize. Highland maize varieties have to mature and complete the grain filling before the first frosts (Alvarado-Beltrán *et al.*, 2019). In warmer lowland conditions, highland material is precocious, flowering in as little as 40 to 50 days.

In the Mexican highlands, farmers have adapted their management practices to improve their chances of obtaining a successful harvest (CIMMYT & Bjarnason, 1994; Eagles & Lothrop, 1994). To maximize the length of the growing season, farmers sow early, before the onset of the annual rains. Traditionally, seeds are deep planted (10 - 25 cm) to benefit from residual soil humidity and to protect from damage from late frosts. This practice allows varieties

that require 160-180 days to reach maturity to be grown in areas with a frost-free season of 90-120 days. The volcanic soils of the Mexican highlands have low pH, restricting the availability of phosphate to the plant (Bayuelo-Jiménez *et al.*, 2011). Although displaying enhanced phosphorus use efficiency (Bayuelo-Jiménez & Ochoa-Cadavid, 2014), Mexican highland landraces tend to show restricted root growth (CIMMYT & Bjarnason, 1994; Eagles & Lothrop, 1994). To compensate for weak root development and prevent lodging, plants may be hilled (piling of soil around the base of the plant) up to three times during vegetative growth.

Palomero Toluqueño (PT) is a popcorn distributed in the highlands of the Mexican Central Plateau, notably in the valley of Toluca, at elevations from ~2100 to ~2900 m.a.s.l. (Fig. 1A. (Wellhausen *et al.*, 1951; Ruiz Corral *et al.*, 2008; Perales & Golicher, 2014)). Although present day cultivation is limited (<https://www.biodiversidad.gob.mx/diversidad/proyectoMaices>), PT is considered ancestral to the broader Mexican highland maize group and a progenitor of more productive modern highland landraces (Reif *et al.*, 2006; Arteaga *et al.*, 2016). PT has a relatively small genome and was selected as the target of the first landrace maize genome sequencing study (Vega-Arreguín *et al.*, 2009; Vielle-Calzada *et al.*, 2009). Subsequent work has continued to explore gene expression variation in PT (Aguilar-Rangel *et al.*, 2017; Crow *et al.*, 2020) and characterize wild-relative introgression through generation of additional genomic sequence (Gonzalez-Segovia *et al.*, 2019). In contrast, B73 is an elite inbred line developed in 1972 by Iowa State University as part of the breeding program for the US maize corn belt. It was highly prized for its ability to form high yielding hybrids and soon became a key ancestral female line in today's global germplasm pool (Troyer, 1999; Iowa State University, 2009). B73 was selected for the

first genome assembly in maize (Schnable *et al.*, 2009) and remains as the primary reference genome (Jiao *et al.*, 2017).

In this work, we characterize a mapping population generated from the cross of B73 and PT. We demonstrate local adaptation in PT and characterize associated genetic architecture by reciprocal transplant and two-site evaluation of our mapping population. We identified Quantitative Trait Loci (QTL) linked to phenology, morphology and yield components, including evidence to QTL x environment interaction (QEI). Overall, morphological QTL were stable across environments with either little QEI or mild scaling effects. We observed stronger QEI associated with yield components, including an example of antagonistic pleiotropy on chromosome (chr) 7, indicating a single locus fitness trade-off. We found evidence for relatively complex genetic architectures associated with putatively adaptive morphological traits. We discuss the implications of these results with respect to the origin of adaptive variation during rapid local adaptation in cultivated species.

MATERIALS AND METHODS

Plant material

To generate the biparental mapping population, an F₁ was generated from the cross between the reference inbred line B73 and pollen pooled from several individuals of Palomero Toluqueño (PT), an open pollinated landrace endemic of the Mexican highlands. The accession used was MEXI5 (CIMMYTMA-002233) obtained from the International Center for Maize and Wheat Improvement (CIMMYT) seed bank, originally collected near the city of Toluca, Mexico State (19.286184N, -99.570871W) at 2597 m.a.s.l. A single B73xPT F₁ individual was crossed as male to multiple B73 ears to generate a large BC₁ population, capturing a single haplotype of PT.

The BC₁ was then self-pollinated 5 generations to form a BC₁S₅ RIL population, with an average of 25% of their genome from PT, and 75% of B73. 120 different families were advanced as independent pedigrees from BC₁S₁ to BC₁S₅. The same initial crossing strategy was used to generate material from the cross between B73 and the open-pollinated Conico/Celaya accession Michoacán 21 (Mi21; CIMMYTMA-001872). B73xMi21 stocks were further backcrossed to B73 with phenotypic selection for sheath pubescence and a segregating BC₅S₁ stock produced. The progenitor B73xPT and B73xMi21 F₁ individuals described here are the same as those used in a previous report to derive introgression stocks segregating the *Inv4m* inversion polymorphism (Crow *et al.*, 2020).

DNA preparation and genotyping

DNA was extracted from 100 B73xPT Recombinant Inbred Lines and four different B73 and PT individuals using isopropanol extraction. 50 mg of leaf tissue were harvested for each plant in a 2.0 mL tube and then frozen to -80 C. The frozen tissue was ground in a Qiagen TissueLyser II (Cat. ID: 85300) with a 30 Hz frequency for 30 s. After grinding, 300 µL of UEB1 (250 mM NaCl, 200 mM Tris pH 7.5, 25 mM EDTA, 0.5%SDS) buffer were added and the solution was mixed in a Thermomixer at 38 C for 10 minutes. 2µL of PureLink RNase were added and the mix was left incubating for 30 min. After incubation, samples were separated by centrifugation at 14,000 rpm for 10 min at room temperature. 250 µL of supernatant was recovered and collected in a 1.5 mL tube. 40 µL of 3 M sodium acetate, pH 5.2 and 450 µL of isopropanol were added per tube, and samples incubated for 20 min at 4 C. A further centrifugation step was performed (14000 rpm, 10 min, room temperature) and the supernatant was discarded. Pellets were washed twice with 250 µL of 70% ethanol. The supernatant was discarded, and the pellet was left to dry

for 30 min. When the pellet was dry, it was resuspended in 100 μ L of milliQ water. DNA was quantified by spectroscopy and adjusted to a concentration of 20 ng/ μ L. DNA was genotyped at the SAGA (Servicio de Análisis Genético para la Agricultura, <https://seedsofdiscovery.org/about/genotyping-platform/>) laboratory in CIMMyT by DArTSeq (Edet *et al.*, 2018) , generating $\sim$ 30,000 short length reads per sample.

Processing of short read genotyping data and construction of the genetic map

Short read DNA sequences generated DArT-Seq were aligned to the v4 B73 reference genome (Jiao *et al.*, 2017) using seqmap (Jiang & Wong, 2008). Sequences that aligned to more than one physical position in the reference genome or that did not align were discarded. Genotype and SNP calling were performed with TASSEL 5 (Bradbury *et al.*, 2007). SNP calls were transformed to an ABH format, A assigned to B73 and B to PT. Sites for which the parental genotype was missing, ambiguous or heterozygous were removed. SNP calls were processed using Genotype-Corrector (Miao *et al.*, 2018), which considerably increased the contiguity of haplotypes among chromosomes. A set of 2, 067 polymorphic markers were selected for further analysis. The ABH genotype file was visualized using R/ABHgenotypeR (Reuscher & Furuta, 2016). Linked markers with shared patterns of segregation were identified with findDupMarkers function of R/qtl package (Broman *et al.*, 2003). Removing redundant makers reduced the final set to 918 polymorphic markers. The linkage map was built using the R/ASmap::mstmap (Taylor & Butler, 2017) under the Kosambi map function. Five individuals from a B73xMi21 BC₅S₁ family segregating sheath pubescence were genotyped using the same DArT-Seq platform as part of a project described in (Gonzalez-Segovia *et al.*, 2019).

Field Evaluation

The BC₁S₅ population was evaluated in the highlands during 2015, 2016, 2018 and 2019 at 2610 m.a.s.l. in Metepec (MT; Mean average temperature: 12.4 °C; mean annual precipitation: 809 mm; Andosolic soil), Mexico State, and in the lowlands during 2015 and 2016 at 54 m.a.s.l. in Valle de Banderas (VB. Mean average temperature: 25.8 °C; mean annual precipitation: 1173 mm; Regosolic soil), Nayarit (Table S1; Fig. S6). BC₁S₅ families were evaluated in single-row 15 plant plots with 15 plants in 3 randomized complete blocks in MT and two blocks in VB. B73 and PT parents were inserted randomly in each block during 2015 and 2016. Weeds and insects were controlled by chemical methods as needed. The VB field site was provided with a ferti-irrigation. System. The MT field site was rain-fed with supplemental sprinkler irrigation after planting and when needed.

Data preparation and trait estimation

Preparation of trait data and QTL mapping was performed in R Statistics (Robinson *et al.*, 2010; McCarthy *et al.*, 2012; R Core Team, 2019). Data collected from single row plots were collapsed to a single value per plot: plot medians were taken for traits scored on multiple individuals; plot level traits such as stand count or flowering time were unchanged. Data was trimmed to remove outliers per trait/location (VB or MT) using R/graphics::boxplot default criteria. Continuous traits (ASI, DTA, DTS, ED, EH, EL, EW, PH, TKN, TKW, and TL) were further adjusted on a per block basis to a spline fitted using R/stats::smooth.spline against row number to reduce spatial variation at the sub-block scale. Spline fitting was not applied to any block containing less than 50 plots. The final dataset contained 4 years of data for location MT (123 genotypes from one block in 2015, 105 genotypes from three blocks in 2016, 140 genotypes from two blocks in

2018, and 110 genotypes from one block in 2019), and two years of data for location VB (123 genotypes from one block in 2015, and 117 genotypes from two blocks in 2016). For each continuous phenotypic trait, a mixed linear model was fitted using restricted maximum-likelihood with R/lme4::lmer. To fit the model, a location-year variable was generated to represent the location by year combinations, and a location-year-block variable was generated to represent all location, year, and block combinations, such that:

$$y_{ijkm} = \mu + E_i + G_j + GE_{ij} + Y_k + B_m + \varepsilon_{ijkm}$$

where the response variable y_{ijkm} is a function of the overall mean (μ), fixed effect of location (E_i), random effect of genotype (G_j), genotype by location interaction (GE_{ij}), location-year term (Y_k), location-year-block term (B_m), and the residual. BLUP values for the genotypic effect (G) and genotype by location interactions (GEI) were extracted using R/lme4::ranef. We calculated BLUP values for each genotype and location combination (G+GEI) by adding genotypic BLUPs and GEI BLUPs (Olivoto *et al.*, 2019). We also calculated fitted values by adding BLUPs to the appropriate means for data visualization and downstream analyses using natural units: for genotypic effect, fitted values were calculated by adding genotypic BLUPs to the grand mean; fitted values for each genotype and location combination were calculated by adding G+GEI BLUPs to the location mean. The significance of the environment effect was evaluated by comparing the full model with location effect and the reduced model without location effect using the likelihood ratio test for continuous phenotypic traits (Table 1). For phenotypic traits with count and scale data, two-group Wilcoxon tests were conducted to evaluate the difference between the two locations (Table 1).

QTL mapping

The BLUPs for continuous traits, the medians for count traits and the mode for semi-quantitative scale traits were used as phenotypic inputs for QTL mapping. Phenotypic scores were selected/combined to perform four distinct analyses: 1) GEN: the genotype main effect (G) of the mixed linear model for continuous traits and the median/mode across all plots for other traits; 2) VB: G+GxE term for VB for continuous traits and the VB median/mode for other traits; 3) MT: G+GxE term for MT for continuous traits and the MT median/mode for other traits; 4) GEI: the difference between MT and VB GxE BLUPs for continuous traits and the difference between MT and VB median/mode for other traits.

Individual QTLs were detected using single QTL scan and Multiple QTL Mapping (MQM) with R/qtl::scanone (default options; Haley-Knott regression. Broman *et al.*, 2003) and R/qtl::MQM (default options. 100 autocofactors, step.size = 1, window.size = 25. Arends *et al.*, 2010), respectively. Genome-wide LOD significance thresholds were established at $\alpha = 0.05$ by 1000 permutations of scanone and MQM and scanone models. Individually QTL were combined in an additive multi-QTL model with R/qtl::makeqtl and their positions refined with R/qtl::refineqtl. The function R/qtl::addqtl was used to detect additional QTLs in a multi-QTL context with a LOD threshold of 3 LOD considered significant. The final multi-QTL model was applied using R/qtl::fitqtl (Haley-Knott regression) to obtain the refined position and variance explained. Significance levels of the full model and the component QTL terms were obtained from the drop-one ANOVA table. Bayes confidence intervals were obtained from the fitqtl model. The effect size and effect plots of each individual term of the full model were obtained with R/qtl::effectplot.

Elevation eGWAS

We performed an environmental genome-wide association analysis (eGWAS) to measure the association between genetic loci and the elevation of native environment for landrace accessions across Mexico, as previously described (Gates *et al.*, 2019). The data set consisted of 1, 830 Mexican maize landrace accessions from the CIMMyT Maize Germplasm Bank with elevation data, genotyped for 440, 000 single nucleotide polymorphisms (SNPs; Romero Navarro *et al.*, 2017; Gates *et al.*, 2019). We used a linear model to fit the genotypic data and elevation to the landrace data. The first five eigenvectors of the genetic relationship matrix were included in the linear model to control for the population structure. The top 1000 SNPs with the strongest association with elevation were selected and used in the downstream analysis.

Data availability statement

PT (CIMMYTMA-002233) and Mi21 (CIMMYTMA-001872) accessions are available directly from CIMMyT (www.cimmyt.org). All other materials are available on request subject to costs of propagation and export if outside of Mexico. Derived material is covered by the same CIMMyT MTA as the progenitor parents. Supplemental files available at <https://doi.org/10.6084/m9.figshare.16608517.v1>. *Phenotypic data*: BLUPs and fitted values estimated for diverse traits for 97 B73xPT BC₁S₅ families. *Genetic map*: genetic map of the B73xPT BC₁S₅ mapping population. *Altitude eGWAS*: results contains the results of the eGWAS analysis. *QTL LOD profile*: LOD profile of the multi-QTL models for different traits for each set of phenotypic data. *Effect Plots*: estimated effect of the QTLs detected for a set of phenotypic data using fitted values. *Reaction norms*: contains the reaction norms estimated for all the QTLs detected in VB and MT phenotypic sets using fitted values.

RESULTS

The stress of the highland environment limits maize growth and productivity

To characterize the genetic architecture of highland adaptation in Mexican native maize (Fig. 1A), we crossed the highland landrace Palomero Toluqueño (PT) to the US reference inbred line B73 and derived 120 BC₁S₅ families. When generating the BC₁, we used a single F₁ individual as a male to pollinate several B73 females, ensuring that a single PT haplotype was captured from the open-pollinated donor accession. As a consequence our mapping population was bi-allelic, *i.e.* segregating for B73 and a single PT allele at any given locus. The final BC₁S₅ families carried ~25% PT genome in a B73 background, with homozygosity > 98%. BC₁S₅ families were genotyped using the DArT Seq platform (<http://www.diversityarrays.com/>) and a final set of 918 markers were selected and used to generate a linkage map for quantitative trait locus (QTL) mapping.

We evaluated the 120 B73 x PT BC₁S₅ families and parents in Mexican lowland (Valle de Banderas, Nayarit at 54 m.a.s.l.) and highland (Metepc, Mexico State, at 2610 m.a.s.l.) field sites. Lowland trials were conducted during the dry season from November to March with supplemental irrigation. Highland trials were conducted in a rain-fed field in the standard highland cycle from April to November. We collected data on a range of phenological, morphological and agronomic traits (Table 1). B73 and PT parents showed a classic pattern of rank-changing GEI for yield components across the two locations, demonstrating adaptation of PT to the highland environment (Fig. 1B, 1-S5). The negative impact of the highland environment on B73 was dramatic, while PT was more stable across the two sites. Across all of the BC₁S₅ genotypes, there was a significant environmental effect on 9 of 19 traits (Fig. 1C; 1-S1; Table 1). Average flowering (days to anthesis, DTA and days to silking, DTS), measured in

chronological days, was greatly delayed in the highland site by 71 days. Overall, plants in the highlands were shorter in stature (plant height, PH) and produced smaller ears (ear diameter, ED; ear height, EH; ear weight, EW) bearing fewer grains (total kernel number, TKN) (Fig. 1C). Average total kernel weight per plant (TKW) dropped from 36.5 g to 15.3 g, a reduction of 58 %, from the lowland to highland field (Fig.1 C).

Segregation in the BC₁S₅ reflects GEI seen in the B73 and PT parents

Having characterized the main effect of the highland environment, we explored GEI among the BC₁S₅ families. For certain traits, such as EH, there was a clear environmental effect (Fig. 2A), but little evidence of GEI between the parents nor among the lines (Fig. 2B). In contrast, yield components such as EW and TKW showed a strong environmental effect (Fig. 1C), GEI between parents, and extensive rank-changing GEI among BC₁S₅ families (Fig. 2C, D; Fig. 2-S1). Taking EW as a proxy for yield, many BC₁S₅ families were more stable with respect to location than the two parents, although the majority were inferior to the better parent in either site. That said, we did observe a small number of families that performed as well as, or better, than the parents in both locations.

QEI associated with yield components indicates local adaptation at the locus level

Extensive GEI for yield components could be associated with either conditional effects or antagonistic pleiotropic at individual QTL (Fig. 3A). To explore the genetic architecture underlying GEI for yield components in our BC₁S₅ families, we performed a QTL analysis, using a series of trait combinations to allow us to determine QTL effects in the lowland and highland sites, as well as to identify QEI. Across the four phenotypic sets, we identified 44 different QTLs

where eighteen were significant under the GxE analysis, mirroring the GEI observed at the level of individual lines (Table 2).

We identified a total of 14 QTL for the yield components: ear length (qEL3, qEL4, qEL7, and qEL8), ear diameter (qED4, qED5, and qED8), kernel row number (qKRN1 and qKRN8), kernels per row (qKPR8), ear weight (qEW7 and qEW8), and total kernel number (qTKN7, qTKN8, and qTKN9) (Fig. 3C, Table 2). QTL for the primary yield component EW on chromosomes (chr) 7 and 8 showed the B73 alleles to be associated with higher EW in the lowlands, and the PT alleles associated with higher EW in the highlands (Fig. 3C). Statistical support for GEI was detected at EW7 but not EW8. There was some support for antagonistic pleiotropy at EW7, although evidence was strongest for a highland conditional effect (Fig. 3D, E, F).

PT alleles at major flowering time QTL on chromosome 8 and 6 accelerate flowering

We detected 13 unique QTLs for flowering time traits (DTA, DTS, ASI) some of which were detected across all analyses (*e.g.* qDTA8b, detected in GEN, VB, MT and GEI sets). Clusters of flowering QTL were found on both chromosomes 8 and 6 (Table 2). qDTA8b, qDTS8 and qASI8 were consistently detected across the GEN, MT, VB and GEI sets, and qDTA6, qDTS6 were detected in the GEN, MT and GEI sets. Mirroring the parental difference, the PT allele at qDTA8b, qDTS8, qDTA6 and qDTS6 accelerates flowering. Flowering QTL have been consistently detected on Chr 8 in maize-teosinte mapping populations and maize diversity panels in the context of differences between temperate and tropical material (Jiang *et al.*, 1999; Chardon *et al.*, 2004; Buckler *et al.*, 2009; Coles *et al.*, 2010; Bouchet *et al.*, 2013; Xu *et al.*, 2017; Guo *et al.*, 2018). The QTLs on chromosome 8 are in the vicinity of the well-characterized flowering

loci *vgt1* and *ZEA CENTRORADIALIS 8 (Zcn8)*. The locus *vgt1* corresponds to a non-coding region of ~ 2 kb that regulates *ZmRap2.7*, an *APETALA-2* like gene located ~ 70 kb downstream (Salvi *et al.*, 2007); *Zcn8* is the florigen gene of maize and has a central role in mediating flowering (Meng *et al.*, 2011; Guo *et al.*, 2018). Polymorphisms in *vgt1* and *Zcn8* have previously been associated with flowering time variation associated with both adaptation to latitude and altitude (Salvi *et al.*, 2007; Ducrocq *et al.*, 2008; Buckler *et al.*, 2009; Romero Navarro *et al.*, 2017; Guo *et al.*, 2018). Given their close proximity, it was not possible to confidently separate the potential effects of *vgt8* and *Zcn8* in our population, and we consider it possible that the combined effect of variation in these two loci underlies our Chr 8 QTL.

The QTL qDTA6 and qDTS6 are in close proximity to the gene *Peamt2* (Zm00001eb294690, chromosome 6 ~ 166.5 MB), an ortholog of the *Arabidopsis XIPTL1* gene which encodes a phosphoethanolamine N-methyltransferase (PEAMT). PEAMT catalyzes the transformation of phosphocholine to phosphatidylcholine (PC), (Cruz-Ramírez *et al.*, 2004; Sánchez Martínez, 2018). The balance between PC and its precursors has been associated with the timing of flowering in *Arabidopsis* (Nakamura *et al.*, 2014) and previously implicated in early flowering in Mexican highland maize (Rodríguez-Zapata *et al.*, 2021). Nevertheless, fine mapping and metabolic studies would be needed to confirm the possible role of variation of *Peamt2* in flowering time variation

Identification of QTL linked to characteristic tassel and stem traits

PT displays a number of putatively adaptive morphological traits that are characteristic of the Mexican highland group as a whole (Fig. 1A, Eagles & Lothrop, 1994; Gonzalez-Segovia *et al.*, 2019). To gain insight into the targets and mechanism of selection during local adaptation, we

collected data on tassel (male inflorescence) morphology, stem pigmentation and stem pubescence from our BC₁S₅ families (Table 1).

The PT tassel is large but unbranched with respect to B73 or typical Mexican lowland landraces (Fig. 4A, B). Across the BC₁S₅ population, tassel length (TL) and tassel branch number (TBN) showed a mild reduction in the highlands compared with the lowland environment, with little GEI between parents or among families (Fig. 4C-F). We identified two QTL linked to TL (qTL1 and qTL2) and two linked to TBN (qTBN2 and qTBN7; Table 2). In common with observations at the whole genotype level, QTL effects for tassel traits were constant in the two environments, and there was no indication of QEI (Perez-Limon et al. 2021, QTL reaction norms). For qTL1, the effect was not aligned with the parental difference, with the PT allele being linked to shorter TL. For TBN, the PT allele at qTBN7 was linked to less branching, while the PT allele at qTBN2 was linked to greater branching (Figure 4H). The largest TBN effect was associated with qTBN7 that co-localized with a previously reported TBN QTL (Xu *et al.*, 2017; Gonzalez-Segovia *et al.*, 2019) and the *Ramosa1* (*Ra1*, Zm00001d020430, chromosome 7 at 113.57 MB) candidate gene (Sigmon & Vollbrecht, 2010; Fig. 3G, H).

PT displays strong stem pigmentation in comparison to the non-pigmented stem of B73 (Fig. 1A, 5A). We detected two QTL for pigment intensity (qINT2 and qINT10) with no evidence of QEI (Perez-Limon et al. 2021, QTL reaction norms; Table 2). The qINT2 interval colocalizes with a QTL previously reported in a Palomero Toluqueño x Reventador F₂ mapping population (Gonzalez-Segovia *et al.*, 2019). Pigment QTL were linked to the well-characterized basic helix-loop-helix (bHLH) regulators of anthocyanin biosynthesis *b1* (Zm00001d000236 chr 2, 198.2 MB) and *r1* (Zm00001d026147, chr10 139.78 MB. Dooner & Kermicle, 1976; Radicella *et al.*, 1992; Selinger *et al.*, 1998; Selinger & Chandler, 1999, 2001; Chatham & Juvik,

2021). It has been demonstrated that functional variation at *b1* is driven by varying patterns of upstream transposon insertion, leading to differences in pigmentation patterns. For example, *B-Bolivia* induces the biosynthesis of anthocyanin in both vegetative tissue and the aleurone of the grain, while the *B-Mex7* allele, which was identified from the Mexican highland landrace Cacahuazintle, induces pigment in the sheath margins of the sheath (Chandler *et al.*, 1989; Radicella *et al.*, 1992; Selinger & Chandler, 1999).

PT, in common with other Mexican highland landraces, exhibits pronounced stem pubescence (Fig. 5A). Although stem macrohairs were present in the BC₁S₅ families, no single family reached the level of pubescence seen in the PT parent, suggesting a complex genetic architecture. Furthermore, the reduced vigor of the BC₁S₅ families in the highland location was associated with poor expression of the pubescence trait and difficulty in scoring. Using a semi-quantitative scale for evaluation, we identified four QTL linked to stem macrohairs, located on chromosomes 3, 7, 8, and 9 (Fig. 5). The QTL interval qMH9 included the *macrohairless1* (*mhl1*, bin 9.04) locus that has previously been linked to the production of leaf blade macrohairs in temperate inbred maize (Moose *et al.*, 2004). The qMH9 region also coincided with a previously reported region of introgression from the highland teosinte *Zea mays* ssp. *mexicana* (itself typically pubescent) to Mexican highland maize (Hufford *et al.*, 2013; Gonzalez-Segovia *et al.*, 2019; Calfee *et al.*, 2021). This region has been characterized as a chromosomal inversion of ~3 MB that displays patterns of selection in highland maize populations (Calfee *et al.*, 2021). The qMH9 interval was relatively large (~12 cM, estimated to cover ~100 Mb) and inspection of the LOD profile suggested the possible presence of two peaks (Fig. 5B). Although presented here as a single QTL, there may in fact be two linked factors.

For all macrohair QTL, the PT allele was associated with greater stem pubescence. We previously reported difficulty in mapping sheath macrohairs in a PT x lowland landrace F₂ population because nearly all plants were scored as pubescent in a simple qualitative evaluation (Gonzalez-Segovia *et al.*, 2019). We interpreted this previous observation to indicate the action of several partially dominant factors, each individually sufficient to trigger the production of stem macrohairs. To further test this hypothesis, we extracted the effect of the PT allele at each macrohair QTL in turn, fixing the other loci as B73 (Fig. 5C). Consistent with genetic redundancy, the PT allele at any macrohair QTL was sufficient to promote a degree of stem pubescence ($p < 0.01$ for all three QTL compared to families carrying B73 alleles at all qMH loci). Although limited by the size of our population and the qualitative nature of our phenotyping, we could detect a significant difference between families carrying PT alleles at several macrohair loci and those carrying the PT allele at only one of the loci ($p = 0.26$; Fig. 5D). Unfortunately, no family carried PT alleles at all four of the loci (this is not unexpected in a BC₁ population of 120 families). Although the four macrohair loci were individually sufficient to induce stem macrohair production, we hypothesise that their combined effect (and potentially that of additional loci) is necessary to approach the levels of pubescence of the PT parent.

In parallel with generation of the BC₁S₅ population, we also produced pubescent near isogenic lines (NILs) by phenotypic selection and recurrent backcrossing to B73. Here, we initially used several different Mexican highland landrace donors. Material generated from the PT relative Conico (accession Michoacan 21) consistently showed the greatest pubescence and was prioritized for backcrossing and genotypic analysis. A BC₅S₁ family showed 3:1 segregation of pubescent to glabrous plants, indicating the action of a single, dominant locus (we did not attempt to distinguish degrees of pubescence in this evaluation, and we do not exclude partial

dominance or an additive effect). We selected two strongly pubescent and three strongly glabrous individuals for genotyping using the DArT-Seq platform. The pubescent individuals carried a large block of Mi21 introgression across chr 3 that was absent from glabrous plants (Fig. 3-S1). Introgression carried in the pubescent NIL spanned the qMHP3 interval identified in the B73xPT population, providing an independent line of evidence for a QTL in this location. There was no evidence of significant Mi21 introgression on chr 7, 8 or 9 in these BC₅S₁ individuals, supporting our previous conclusion that macrohair QTL are individually sufficient to promote a degree of stem macrohair production. The BC₅S₁ family provides a good starting point towards fine mapping and cloning of qMHP3.

Comparison of B73xPT QTL and broader landrace diversity

To compare QTL detected in our B73xPT BC₁S₅ population to the broader diversity present in Mexican highland maize, we performed an environmental-genome wide association study (eGWAS) using a previously genotyped panel of 1830 geo-referenced Mexican landrace accessions (Romero Navarro *et al.*, 2017; Gates *et al.*, 2019; Fig. 6A; 4-S1). We selected the top 1000 SNPs most significantly associated with elevation and compared their physical location with the location of our QTL. The strongest environmental association was detected on chromosome 4 at the previously reported inversion polymorphism *Inv4m* (Romero Navarro *et al.*, 2017; Crow *et al.*, 2020). Although qED4, detected in our QTL mapping, co-localized with this region on chromosome 4 (Fig. 6B), we found no signal linking *Inv4m* to additional yield components or evidence of a previously reported flowering time effect in our QTL analysis. Further high confidence SNPs were found on chromosomes 2, 3, 5, 7, 8 and 10 (Fig. S4). A high confidence SNP on chr 7 (7_17794242) was identified adjacent to the GEI peak associated with

qEW7 (Fig. 6C). This region was also identified in a previous experiment to map yield and harvest index QTL in a comparison of lowland and highland Mexican maize (Jiang *et al.*, 1999). In other cases, for example qTKW8 (Fig. 6D), there was no strong correspondence between the eGWAS hits and the location of the QTL. Although we would not draw strong conclusions from differences between eGWAS and QTL results, those cases where they do overlap provide compelling candidates for functional study. For example, the SNP 7_17794242 lies within a gene (Zm00001d019117) encoding a putative transmembrane protein that has been shown in temperate maize to be differentially expressed in response to salt, cold and UV (Makarevitch *et al.*, 2015).

DISCUSSION

Evaluation of a B73xPT mapping population in lowland and high elevation field sites identified QTL associated with both morphological and yield components traits. Although showing plasticity, the genetic architecture of morphological traits was conserved across environments and we saw little evidence of GEI. Indeed, characteristic highland traits such as pigmentation and pubescence were actually easier to evaluate in the lowland field as a result of the overall greater vigor of the plants. In contrast, we saw greater evidence of GEI with respect to yield components, with individual BC₁S₅ genotypes showing the signature of local adaptation and others stability across our two test environments. This broad trend of greater stability of morphological traits rather than yield components is consistent with a previous study mapping maize adaptation across four elevations (Jiang *et al.*, 1999).

In total, across the different phenotypic sets, we detected 44 unique QTLs, eighteen of which present a significant QEI where the majority (17) are examples of conditional neutrality,

while only one (qEW7) was associated with statistical support for antagonistic pleiotropy. In a review of genetic architecture in 37 studies, the authors estimated that ~60% of the QTLs detected displayed QEI, but that there was only evidence for antagonistic pleiotropy in ~2% of cases (Des Marais *et al.*, 2013). This broad trend was reflected in a recent multisite mapping experiment in switchgrass (*Panicum virgatum* L.) in which the majority of QTL associated with adaptive traits showed conditional positive effects in their home environment with little or no detectable effect or cost in other environments (Lowry *et al.*, 2019). Detecting antagonistic pleiotropy requires higher statistical power than identification of conditional effects (*i.e.* the latter are typically supported by a failure to detect an effect in certain environments), resulting in a potential bias in the classification of QEI (Anderson *et al.*, 2011). In our study, the statistical power necessary to dissect QEI is limited by the size of the mapping population and the number of trials and locations evaluated. QTL effects for biomass, yield components changed in both magnitude and direction over location, suggesting antagonistic pleiotropy, and additional evaluation of our material may provide more evidence of this. We would also point out the extensive nature of the management employed at the lowland (Valle de Banderas) site; this site might be considered as an “ideal” environment in comparison to the highland site, exposing the plants to few of the stresses traditionally faced in tropical lowland fields. Any buffering of the potential costs of highland variants in the lowland site by management would push the genetic architecture from antagonistic pleiotropy towards conditionality.

We identified several QTLs that could be confidently associated with strong candidate genes. For stem pigmentation, qPINT2 and qPINT10 correspond well to the loci *b1* and *r1*, respectively. The qPINT2 locus had the greatest effect of the two (explaining ~40% variance) with the PT allele promoting pigmentation. The *b1* gene encodes a bHLH transcription factor

that regulates the temporal and tissue-specific expression of genes that produce anthocyanins in maize (Ludwig *et al.*, 1989; Petroni *et al.*, 2000; Chatham & Juvik, 2021). Interestingly, *b1* was also identified in a mapping cross between lowland and highland teosinte, the latter showing the stem pigmentation also seen in highland maize (Lauter *et al.*, 2004). Several independently derived *B1* alleles have been linked to stem pigmentation (Dooner & Kermicle, 1976; Radicella *et al.*, 1992; Selinger *et al.*, 1998; Selinger & Chandler, 1999, 2001; Chatham & Juvik, 2021), indicating the ready production of functional diversity at this locus and implicating convergent selection (Stern, 2013) for pigmentation among highland *Zea*, supporting an adaptive role (Doebley, 1984; Lauter *et al.*, 2004). Further sequencing of *b1* alleles from highland maize will shed greater light on patterns of diversity and the origin of different alleles. Stem pigmentation, unlike stem pubescence, is also shared with South American highland maize (Janzen *et al.*, 2021). Dark red pigmentation in the stem can help the plant to absorb more solar radiation and keep the plant warmer in a cold environment and might also protect DNA from damage due to higher UV-B radiation in the highlands (Barthakur, 1974; Eagles & Lothrop, 1994; Casati & Walbot, 2005) - although it is unclear why such protection might be required more so in the stem than in the leaf blades.

The flowering QTL qDTA8, qDTS8 and qASI8 overlap a ~10Mb region that contains the two well-characterized flowering genes *Vgt1* and *Zcn8*. This region and/or these genes have been reproducibly detected in linkage- and association- mapping studies of maize flowering time (Chardon *et al.*, 2004; Buckler *et al.*, 2009; Steinhoff *et al.*, 2012; Li *et al.*, 2016; Romero Navarro *et al.*, 2017), temperate adaptation (Ducrocq *et al.*, 2008; Bouchet *et al.*, 2013; Guo *et al.*, 2018; Castelletti *et al.*, 2020) and adaptation to the Mexican Highlands (Gates *et al.*, 2019; Janzen *et al.*, 2021; Wang *et al.*, 2021). An early flowering *vgt1* allele from northern germplasm

has previously been associated with a miniature transposon (MITE) insertion, although the absence of the MITE alone did not explain late flowering *vgt1* alleles (Buckler *et al.*, 2009). In a *Zcn8* association study using maize and teosinte, the haplotype associated with earliest flowering (A-Del) was hypothesised to have originated in highland teosinte (*Zea mays* ssp. *mexicana*) and to have moved to cultivated maize by introgression (Guo *et al.*, 2018). Interestingly, in this same study the authors report Palomero Toluqueño to carry both the MITE-associated allele of *vgt1* and the A-Del haplotype of *Zcn8*. Although we have not sequenced the *Vgt1* and *Zcn8* alleles present in our mapping population and available genome sequence data do not provide good coverage in this region, our linkage mapping results are consistent with this previous association analysis.

We identified QTLs associated with sheath pubescence on chr 3, 7, 8 and 9. Our QTL on chr 9 is consistent with previous observations co-localizing i) a leaf blade macrohair mutation in temperate maize, ii) a stem pubescence QTL in *mexicana* teosinte, iii) introgression from *mexicana* to highland maize and iv) a ~ 3 Mb inversion that displays patterns of selection in mexican highland maize (Moose *et al.*, 2004; Lauter *et al.*, 2004; Hufford *et al.*, 2013; Gonzalez-Segovia *et al.*, 2019; Calfee *et al.*, 2021). Identification of a PT allele linked to stem pubescence in this same region of chr 9 adds further support to the hypothesis of adaptive introgression at this locus (Wilkes, 1972; Gonzalez-Segovia *et al.*, 2019). In this context, it is interesting to note that all macrohair QTL identified appeared to be sufficient on their own to induce stem pubescence, although their combined action would be needed to approach the level of pubescence of the PT parent. Limitations of population size and the semi-quantitative nature of our evaluation prevent strong conclusions concerning additivity or interactions among macrohair QTL. Nonetheless, our data suggest that qMH9 is just one of a number of contributing loci, the

origin of the trait being a complex mix of wild-relative introgression and *de novo* mutation. Fine mapping and molecular cloning of the genes underlying macrohair QTL would allow a far more detailed view of the history of the stem pubescent trait and associated genetic variants in Mexican highland maize. Pubescence extends the boundary layer around the stem and could act as protection from cold by preventing heat loss or conserve water by minimizing transpiration (Chalker-Scott, 1999; Schuepp, 1993). We did not observe any strong correlation between pubescence and yield components. However, further experiments making use of the range of pubescence in our inbred families or derived NILs, might have the power to detect more subtle effects in either controlled conditions or large-scale highland yield trials.

The rich diversity of Mexican landrace maize is closely tied to local adaptation. Yet, this same specialization places these varieties at risk from future climate change (Mercer & Perales, 2010)(Bellon *et al.*, 2011) (Romero *et al.*, 2020). In a study to project landrace distribution under different climate change scenarios, PT was the landrace identified as the most vulnerable (Ureta *et al.*, 2012), although, as the authors note, models based on current distribution and climate do not take into account the full range of environmental, biotic and cultural factors that impact diversity and distribution. In the specific case of PT, limited yield potential in comparison to more modern landraces is likely to see it abandoned by farmers (<https://www.biodiversidad.gob.mx/diversidad/proyectoMaices>). That said, PT has contributed to the broader Mexican highland group (Reif *et al.*, 2006; Warburton *et al.*, 2008)(Arteaga *et al.*, 2016) and locally-adapted alleles will likely be conserved. The fate of the Mexican highland maize group will be influenced by their degree of resilience and ability to adapt to climate change. Although we found some evidence of antagonistic pleiotropy, PT and PT allele effects were largely stable and GEI was driven by plasticity associated with B73. As such, our data

would support cautious optimism that highland varieties might maintain current levels of productivity in the face of future climate change. However, if climate change results in the expansion of lower elevation varieties to the highlands, the home site advantage of traditional highland landraces might be eroded. Ultimately, the conservation of maize diversity, along with the responsible and equitable use of this unique resource, will be informed by a greater understanding of the physiological and mechanistic basis of local adaptation. With an increase in mapping resolution (for example by using larger or more diverse populations) and the availability of high quality landrace genome assemblies, it will be possible to take important further steps towards defining not just the genetic architecture but also the genes and genetic variants that underlie local adaptation in maize landraces.

REFERENCES

- Aguilar-Rangel MR, Chávez Montes RA, González-Segovia E, Ross-Ibarra J, Simpson JK, Sawers RJH. 2017.** Allele specific expression analysis identifies regulatory variation associated with stress-related genes in the Mexican highland maize landrace Palomero Toluqueño. *PeerJ* **5**: e3737.
- Alvarado-Beltrán G, López-Sánchez H, Santacruz-Varela A, Muñoz-Orozco A, Valadez-Moctezuma E, Gutiérrez-Espinosa MA, López PA, Gil-Muñoz A, de Dios Guerrero-Rodríguez J, Taboada-Gaytán OR. 2019.** Morphological variability of native maize (*Zea mays* L.) of the west highland of Puebla and east highland of Tlaxcala, Mexico. *Revista de la Facultad de Ciencias Agrarias UNCuyo* **51**: 217–234.
- Anderson JT, Lee C-R, Rushworth CA, Colautti RI, Mitchell-Olds T. 2013.** Genetic trade-offs and conditional neutrality contribute to local adaptation. *Molecular ecology* **22**: 699–708.
- Anderson JT, Willis JH, Mitchell-Olds T. 2011.** Evolutionary genetics of plant adaptation. *Trends in genetics: TIG* **27**: 258–266.
- Arends D, Prins P, Jansen RC, Broman KW. 2010.** R/qtl: high-throughput multiple QTL mapping. *Bioinformatics* **26**: 2990–2992.
- Arteaga MC, Moreno-Letelier A, Mastretta-Yanes A, Vázquez-Lobo A, Breña-Ochoa A, Moreno-Estrada A, Eguiarte LE, Piñero D. 2016.** Genomic variation in recently collected

maize landraces from Mexico. *Genomics data* **7**: 38–45.

Assmann SM. 2013. Natural Variation in Abiotic Stress and Climate Change Responses in Arabidopsis : Implications for Twenty-First-Century Agriculture. *International journal of plant sciences* **174**: 3–26.

Barthakur N. 1974. Temperature differences between two pigmented types of corn plants. *International journal of biometeorology* **18**: 70–75.

Bayuelo-Jiménez JS, Gallardo-Valdéz M, Pérez-Decelis VA, Magdaleno-Armas L, Ochoa I, Lynch JP. 2011. Genotypic variation for root traits of maize (*Zea mays* L.) from the Purhepecha Plateau under contrasting phosphorus availability. *Field crops research* **121**: 350–362.

Bayuelo-Jiménez JS, Ochoa-Cadavid I. 2014. Phosphorus acquisition and internal utilization efficiency among maize landraces from the central Mexican highlands. *Field crops research* **156**: 123–134.

Bellon MR, Hodson D, Bergvinson D, Beck D, Martinez-Romero E, Montoya Y. 2005. Targeting agricultural research to benefit poor farmers: Relating poverty mapping to maize environments in Mexico. *Food policy* **30**: 476–492.

Bellon MR, Hodson D, Hellin J. 2011. Assessing the vulnerability of traditional maize seed systems in Mexico to climate change. *Proceedings of the National Academy of Sciences of the United States of America* **108**: 13432–13437.

Bellon MR, Mastretta-Yanes A, Ponce-Mendoza A, Ortiz-Santamaría D, Oliveros-Galindo O, Perales H, Acevedo F, Sarukhán J. 2018. Evolutionary and food supply implications of ongoing maize domestication by Mexican. *Proceedings. Biological sciences / The Royal Society* **285**.

Bouchet S, Servin B, Bertin P, Madur D, Combes V, Dumas F, Brunel D, Laborde J, Charcosset A, Nicolas S. 2013. Adaptation of maize to temperate climates: mid-density genome-wide association genetics and diversity patterns reveal key genomic regions, with a major contribution of the Vgt2 (ZCN8) locus. *PloS one* **8**: e71377.

Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. 2007. TASSEL: software for association mapping of complex traits in diverse samples. *Bioinformatics* **23**: 2633–2635.

Breseghele F, Coelho ASG. 2013. Traditional and modern plant breeding methods with examples in rice (*Oryza sativa* L.). *Journal of agricultural and food chemistry* **61**: 8277–8286.

Broman KW, Wu H, Sen S, Churchill GA. 2003. R/qt1: QTL mapping in experimental crosses. *Bioinformatics* **19**: 889–890.

Buckler ES, Holland JB, Bradbury PJ, Acharya CB, Brown PJ, Browne C, Ersoz E, Flint-Garcia S, Garcia A, Glaubitz JC, et al. 2009. The genetic architecture of maize flowering time. *Science* **325**: 714–718.

689 **Calfee E, Gates D, Lorant A, Taylor Perkins M, Coop G, Ross-Ibarra J. 2021.** Selective
690 sorting of ancestral introgression in maize and teosinte along an elevational cline. *bioRxiv*:
691 2021.03.05.434040.

692 **Casati P, Walbot V. 2005.** Differential accumulation of maysin and rhamnosylisoorientin in
693 leaves of high-altitude landraces of maize after UV-B exposure. *Plant, cell & environment* **28**:
694 788–799.

695 **Castelletti S, Coupel-Ledru A, Granato I, Palaffre C, Cabrera-Bosquet L, Tonelli C,**
696 **Nicolas SD, Tardieu F, Welcker C, Conti L. 2020.** Maize adaptation across temperate climates
697 was obtained via expression of two florigen genes. *PLoS genetics* **16**: e1008882.

698 **Chandler VL, Radicella JP, Robbins TP, Chen J, Turks' D. 1989.** Two Regulatory Genes of
699 the Maize Anthocyanin Pathway Are Homologous: Isolation of B Utilizing R Genomic
700 Sequences. *The Plant Cell*.

701 **Chardon F, Virlon B, Moreau L, Falque M, Joets J, Decousset L, Murigneux A, Charcosset**
702 **A. 2004.** Genetic architecture of flowering time in maize as inferred from quantitative trait loci
703 meta-analysis and synteny conservation with the rice genome. *Genetics* **168**: 2169–2185.

704 **Chatham LA, Juvik JA. 2021.** Linking anthocyanin diversity, hue, and genetics in purple corn.
705 *G3* **11**.

706 **CIMMYT, Bjarnason MS. 1994.** The Subtropical, Midaltitude, and Highland Maize
707 Subprogram. **viii**.

708 **Cleveland DA, Soleri D. 2007.** Extending Darwin's Analogy: Bridging Differences in Concepts
709 of Selection between Farmers, Biologists, and Plant Breeders. *Economic botany* **61**: 121–136.

710 **Coles ND, McMullen MD, Balint-Kurti PJ, Pratt RC, Holland JB. 2010.** Genetic control of
711 photoperiod sensitivity in maize revealed by joint multiple population analysis. *Genetics* **184**:
712 799–812.

713 **CONABIO. 2018.** Maíz. *Biodiversidad Mexicana*.

714 **Corrado G, Rao R. 2017.** Towards the Genomic Basis of Local Adaptation in Landraces.
715 *Diversity* **9**: 51.

716 **Crow T, Ta J, Nojoomi S, Aguilar-Rangel MR, Torres Rodríguez JV, Gates D, Rellán-**
717 **Álvarez R, Sawers R, Runcie D. 2020.** Gene regulatory effects of a large chromosomal
718 inversion in highland maize. *PLoS genetics* **16**: e1009213.

719 **Cruz-Ramírez A, López-Bucio J, Ramírez-Pimentel G, Zurita-Silva A, Sánchez-Calderon**
720 **L, Ramírez-Chávez E, González-Ortega E, Herrera-Estrella L. 2004.** The xip1 mutant of
721 Arabidopsis reveals a critical role for phospholipid metabolism in root system development and
722 epidermal cell integrity. *The Plant cell* **16**: 2020–2034.

723 **Des Marais DL, Hernandez KM, Juenger TE. 2013.** Genotype-by-Environment Interaction

- 724 and Plasticity: Exploring Genomic Responses of Plants to the Abiotic Environment. *Annual*
725 *review of ecology, evolution, and systematics* **44**: 5–29.
- 726 **Doebley JF. 1984.** Maize Introgression Into Teosinte-A Reappraisal. *Annals of the Missouri*
727 *Botanical Garden. Missouri Botanical Garden* **71**: 1100–1113.
- 728 **Dooner HK, Kermicle JL. 1976.** Displaced and tandem duplications in the long arm of
729 chromosome 10 in maize. *Genetics* **82**: 309–322.
- 730 **Ducrocq S, Madur D, Veyrieras J-B, Camus-Kulandaivelu L, Kloiber-Maitz M, Prestler T,**
731 **Ouzunova M, Manicacci D, Charcosset A. 2008.** Key impact of Vgt1 on flowering time
732 adaptation in maize: evidence from association mapping and ecogeographical information.
733 *Genetics* **178**: 2433–2437.
- 734 **Dwivedi SL, Ceccarelli S, Blair MW, Upadhyaya HD, Are AK, Ortiz R. 2016.** Landrace
735 Germplasm for Improving Yield and Abiotic Stress Adaptation. *Trends in plant science* **21**: 31–
736 42.
- 737 **Eagles HA, Lothrop JE. 1994.** Highland Maize from Central Mexico—Its Origin,
738 Characteristics, and Use in Breeding Programs. *Crop Science* **34**: 11–19.
- 739 **Edet OU, Gorafi YSA, Nasuda S, Tsujimoto H. 2018.** DArTseq-based analysis of genomic
740 relationships among species of tribe Triticeae. *Scientific reports* **8**: 16397.
- 741 **El-Soda M, Malosetti M, Zwaan BJ, Koornneef M, Aarts MGM. 2014.** Genotype ×
742 environment interaction QTL mapping in plants: lessons from Arabidopsis. *Trends in plant*
743 *science* **19**: 390–398.
- 744 **Emon RM, Islam MM, Halder J, Fan Y. 2015.** Genetic diversity and association mapping for
745 salinity tolerance in Bangladeshi rice landraces. *The Crop Journal* **3**: 440–444.
- 746 **Espinosa-Calderón A, Robledo-Tadeo M, Montiel NG, Macías MS, Vargas JV, Caballero**
747 **AP, Hernández FC, Carrillo GV, Rodríguez Montalvo FA. 2011.** ‘V-55 A’, A MAIZE
748 VARIETY OF YELLOW GRAIN FOR MEXICAN HIGHLANDS. *Nueva Variedad Rev. Fitotec.*
749 *Mex. Vol* **34**: 149–150.
- 750 **Fournier-Level A, Korte A, Cooper MD, Nordborg M, Schmitt J, Wilczek AM. 2011.** A
751 map of local adaptation in Arabidopsis thaliana. *Science* **334**: 86–89.
- 752 **Galván-Tejada NC, Peña-Ramírez V, Mora-Palomino L, Siebe C. 2014.** Soil P fractions in a
753 volcanic soil chronosequence of Central Mexico and their relationship to foliar P in pine trees.
754 *Journal of Plant Nutrition and Soil Science* **177**: 792–802.
- 755 **Gardner KM, Latta RG. 2006.** Identifying loci under selection across contrasting environments
756 in Avena barbata using quantitative trait locus mapping. *Molecular ecology* **15**: 1321–1333.
- 757 **Gates DJ, Runcie D, Janzen GM, Navarro AR, Willcox M, Sonder K, Snodgrass SJ,**
758 **Rodríguez-Zapata F, Sawers RJH, Rellán-Álvarez R, et al. 2019.** Single-gene resolution of

759 locally adaptive genetic variation in Mexican maize. *bioRxiv*: 706739.

760 **Gonzalez-Segovia E, Pérez-Limon S, Cíntora-Martínez GC, Guerrero-Zavala A, Janzen**
761 **GM, Hufford MB, Ross-Ibarra J, Sawers RJH. 2019.** Characterization of introgression from
762 the teosinte *Zea mays* ssp. *mexicana* to Mexican highland maize. *PeerJ* 7: e6815.

763 **Guo L, Wang X, Zhao M, Chen Q, Doebley JF, Tian F, Huang C, Li C, Li D, Yang CJ, et**
764 **al. 2018.** Stepwise cis-Regulatory Changes in ZCN8 Contribute to Maize Flowering-Time
765 Adaptation | Elsevier Enhanced Reader. *Current biology: CB*: 3005–3015.

766 **Hall MC, Lowry DB, Willis JH. 2010.** Is local adaptation in *Mimulus guttatus* caused by trade-
767 offs at individual loci? *Molecular ecology* 19: 2739–2753.

768 **Hufford MB, Lubinsky P, Pyhäjärvi T, Devengenzo MT, Ellstrand NC, Ross-Ibarra J.**
769 **2013.** The genomic signature of crop-wild introgression in maize. *PLoS genetics* 9: e1003477.

770 **Iowa State University. 2009.** Researchers provide understanding to maize genome sequence.
771 *Science Daily*.

772 **Janzen GM, Aguilar-Rangel MR, Cíntora-Martínez C, Blöcher-Juárez KA, González-**
773 **Segovia E, Studer AJ, Runcie DE, Flint-Garcia SA, Rellán-Álvarez R, Sawers RJH, et al.**
774 **2021.** Demonstration of local adaptation of maize landraces by reciprocal transplantation.
775 *bioRxiv*: 2021.03.25.437076.

776 **Jiang C, Edmeades GO, Armstead I, Lafitte HR, Hayward MD, Hoisington D. 1999.**
777 Genetic analysis of adaptation differences between highland and lowland tropical maize using
778 molecular markers. *TAG. Theoretical and applied genetics. Theoretische und angewandte*
779 *Genetik* 99: 1106–1119.

780 **Jiang H, Wong WH. 2008.** SeqMap: mapping massive amount of oligonucleotides to the
781 genome. *Bioinformatics* 24: 2395–2396.

782 **Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, Campbell MS, Stein JC, Wei X,**
783 **Chin C-S, et al. 2017.** Improved maize reference genome with single-molecule technologies.
784 *Nature* 546: 524–527.

785 **Juenger TE. 2013.** Natural variation and genetic constraints on drought tolerance. *Current*
786 *opinion in plant biology* 16: 274–281.

787 **Körner C. 2007.** The use of ‘altitude’ in ecological research. *Trends in ecology & evolution* 22:
788 569–574.

789 **Latta RG, Gardner KM, Johansen-Morris AD. 2007.** Hybridization, recombination, and the
790 genetic basis of fitness variation across environments in *Avena barbata*. *Genetica* 129: 167–177.

791 **Latta RG, Gardner KM, Staples DA. 2010.** Quantitative trait locus mapping of genes under
792 selection across multiple years and sites in *Avena barbata*: epistasis, pleiotropy, and genotype-
793 by-environment interactions. *Genetics* 185: 375–385.

794 **Lauter N, Gustus C, Westerbergh A, Doebley J. 2004.** The inheritance and evolution of leaf
795 pigmentation and pubescence in teosinte. *Genetics* **167**: 1949–1959.

796 **Li Y-X, Li C, Bradbury PJ, Liu X, Lu F, Romay CM, Glaubitz JC, Wu X, Peng B, Shi Y, et**
797 **al. 2016.** Identification of genetic variants associated with maize flowering time using an
798 extremely large multi-genetic background population. *The Plant journal: for cell and molecular*
799 *biology* **86**: 391–402.

800 **Louette D, Charrier A, Berthaud J. 1997.** In Situ conservation of maize in Mexico: Genetic
801 diversity and Maize seed management in a traditional community. *Economic botany* **51**: 20–38.

802 **Lovell JT, MacQueen AH, Mamidi S, Bonnette J, Jenkins J, Napier JD, Sreedasyam A,**
803 **Healey A, Session A, Shu S, et al. 2021.** Genomic mechanisms of climate adaptation in
804 polyploid bioenergy switchgrass. *Nature* **590**: 438–444.

805 **Lowry DB, Hall MC, Salt DE, Willis JH. 2009.** Genetic and physiological basis of adaptive
806 salt tolerance divergence between coastal and inland *Mimulus guttatus*. *The New phytologist*
807 **183**: 776–788.

808 **Lowry DB, Lovell JT, Zhang L, Bonnette J, Fay PA, Mitchell RB, Lloyd-Reilley J, Boe AR,**
809 **Wu Y, Rouquette FM Jr, et al. 2019.** QTL × environment interactions underlie adaptive
810 divergence in switchgrass across a large latitudinal gradient. *Proceedings of the National*
811 *Academy of Sciences of the United States of America* **116**: 12933–12941.

812 **Ludwig SR, Habera LF, Dellaporta SL, Wessler SR. 1989.** Lc, a member of the maize R gene
813 family responsible for tissue-specific anthocyanin production, encodes a protein similar to
814 transcriptional activators and contains the myc-homology region. *Proceedings of the National*
815 *Academy of Sciences of the United States of America* **86**: 7092–7096.

816 **Makarevitch I, Waters AJ, West PT, Stitzer M, Hirsch CN, Ross-Ibarra J, Springer NM.**
817 **2015.** Transposable elements contribute to activation of maize genes in response to abiotic stress.
818 *PLoS genetics* **11**: e1004915.

819 **Matsuoka Y, Vigouroux Y, Goodman MM, Sanchez G. J, Buckler E, Doebley J. 2002.** A
820 single domestication for maize shown by multilocus microsatellite genotyping. *Proceedings of*
821 *the National Academy of Sciences* **99**: 6080–6084.

822 **McCarthy DJ, Chen Y, Smyth GK. 2012.** Differential expression analysis of multifactor RNA-
823 Seq experiments with respect to biological variation. *Nucleic acids research* **40**: 4288–4297.

824 **McWilliam JR, Naylor AW. 1967.** Temperature and plant adaptation. I. Interaction of
825 temperature and light in the synthesis of chlorophyll in corn. *Plant physiology* **42**: 1711–1715.

826 **Meng X, Muszynski MG, Danilevskaya ON. 2011.** The FT-like ZCN8 Gene Functions as a
827 Floral Activator and Is Involved in Photoperiod Sensitivity in Maize. *The Plant cell* **23**: 942–960.

828 **Mercer K, Martínez-Vásquez Á, Perales HR. 2008.** Asymmetrical local adaptation of maize
829 landraces along an altitudinal gradient. *Evolutionary applications* **1**: 489–500.

830 **Mercer KL, Perales HR. 2010.** Evolutionary response of landraces to climate change in centers
831 of crop diversity. *Evolutionary Applications* **3**: 480–493.

832 **Mercer KL, Perales H. 2019.** Structure of local adaptation across the landscape: flowering time
833 and fitness in Mexican maize (*Zea mays* L. subsp. *mays*) landraces. *Genetic resources and crop*
834 *evolution* **66**: 27–45.

835 **Miao C, Fang J, Li D, Liang P, Zhang X, Yang J, Schnable JC, Tang H. 2018.** Genotype-
836 Corrector: improved genotype calls for genetic mapping in F and RIL populations. *Scientific*
837 *reports* **8**: 10088.

838 **Mitchell-Olds T, Willis JH, Goldstein DB. 2007.** Which evolutionary processes influence
839 natural genetic variation for phenotypic traits? *Nature reviews. Genetics* **8**: 845–856.

840 **Moose SP, Lauter N, Carlson SR. 2004.** The maize macrohairless1 locus specifically promotes
841 leaf blade macrohair initiation and responds to factors regulating leaf identity. *Genetics* **166**:
842 1451–1461.

843 **Nakamura Y, Andrés F, Kanehara K, Liu Y-C, Dörmann P, Coupland G. 2014.** Arabidopsis
844 florigen FT binds to diurnally oscillating phospholipids that accelerate flowering. *Nature*
845 *communications* **5**: 3553.

846 **Olivoto T, Lúcio ADC, Silva JAG, Marchioro VS, Souza VQ, Jost E. 2019.** Mean
847 performance and stability in multi-environment trials I: Combining features of AMMI and BLUP
848 techniques. *Agronomy journal* **111**: 2949–2960.

849 **Perales H, Golicher D. 2014.** Mapping the diversity of maize races in Mexico. *PloS one* **9**:
850 e114657.

851 **Petroni K, Cominelli E, Consonni G, Gusmaroli G, Gavazzi G, Tonelli C. 2000.** The
852 developmental expression of the maize regulatory gene *Hopi* determines germination-dependent
853 anthocyanin accumulation. *Genetics* **155**: 323–336.

854 **Piperno DR, Moreno JE, Iriarte J, Holst I, Lachniet M, Jones JG, Ranere AJ, Castanzo R.**
855 **2007.** Late Pleistocene and Holocene environmental history of the Iguala Valley, Central Balsas
856 Watershed of Mexico. *Proceedings of the National Academy of Sciences of the United States of*
857 *America* **104**: 11874–11881.

858 **Radicella JP, Brown D, Tolar LA, Chandler VL. 1992.** Allelic diversity of the maize B
859 regulatory gene: different leader and promoter sequences of two B alleles determine distinct
860 tissue specificities of anthocyanin production. *Genes & development* **6**: 2152–2164.

861 **R Core Team. 2019.** R: A Language and Environment for Statistical Computing.

862 **Reif JC, Warburton ML, Xia XC, Hoisington DA, Crossa J, Taba S, Muminović J, Bohn**
863 **M, Frisch M, Melchinger AE. 2006.** Grouping of accessions of Mexican races of maize
864 revisited with SSR markers. *TAG. Theoretical and applied genetics. Theoretische und*
865 *angewandte Genetik* **113**: 177–185.

866 **Reuscher S, Furuta T. 2016.** ABHgenotypeR: Easy Visualization of ABH Genotypes.

867 **Robinson MD, McCarthy DJ, Smyth GK. 2010.** edgeR: a Bioconductor package for
868 differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–140.

869 **Rodríguez-Zapata F, Barnes AC, Blöcher-Juárez KA, Gates D, Kur A, Wang L, Janzen**
870 **GM, Jensen S, Estévez-Palmas JM, Crow T, et al. 2021.** Teosinte introgression modulates
871 phosphatidylcholine levels and induces early maize flowering time. *bioRxiv*: 2021.01.25.426574.

872 **Romero Navarro JA, Willcox M, Burgueño J, Romay C, Swarts K, Trachsel S, Preciado E,**
873 **Terron A, Delgado HV, Vidal V, et al. 2017.** A study of allelic diversity underlying flowering-
874 time adaptation in maize landraces. *Nature genetics* **49**: 476–480.

875 **Romero AA, Rivas AIM, Díaz JDG, Mendoza MÁP, Salas ENN, Blanco JL, Álvarez ACC.**
876 **2020.** Crop yield simulations in Mexican agriculture for climate change adaptation. *Atmósfera*.

877 **Ruiz Corral JA, Durán Puga N, Sánchez González J de J, Ron Parra J, González Eguiarte**
878 **DR, Holland JB, Medina García G. 2008.** Climatic Adaptation and Ecological Descriptors of
879 42 Mexican Maize Races. *Crop science* **48**: 1502–1512.

880 **Salvi S, Sponza G, Morgante M, Tomes D, Niu X, Fengler KA, Meeley R, Ananiev EV,**
881 **Svitashev S, Bruggemann E, et al. 2007.** Conserved noncoding genomic sequences associated
882 with a flowering-time quantitative trait locus in maize. *Proceedings of the National Academy of*
883 *Sciences of the United States of America* **104**: 11376–11381.

884 **Sánchez Martínez ES. 2018.** [Functional characterization of Zea mays Xipotl
885 (phosphoethanolamine N-methyltransferase, PEAMT) family genes] =Caracterización funcional
886 de la familia de genes Xipotl (fosfoetanolamina N-metiltransferasa, PEAMT) de Zea mays.

887 **Scarcelli N, Cheverud JM, Schaal BA, Kover PX. 2007.** Antagonistic pleiotropic effects
888 reduce the potential adaptive value of the FRIGIDA locus. *Proceedings of the National Academy*
889 *of Sciences of the United States of America* **104**: 16986–16991.

890 **Scheiner SM. 1993.** Genetics and Evolution of Phenotypic Plasticity. *Annual review of ecology*
891 *and systematics* **24**: 35–68.

892 **Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, Liang C, Zhang J, Fulton**
893 **L, Graves TA, et al. 2009.** The B73 maize genome: complexity, diversity, and dynamics.
894 *Science* **326**: 1112–1115.

895 **Selinger DA, Chandler VL. 1999.** Major recent and independent changes in levels and patterns
896 of expression have occurred at the b gene, a regulatory locus in maize. *Proceedings of the*
897 *National Academy of Sciences of the United States of America* **96**: 15007–15012.

898 **Selinger DA, Chandler VL. 2001.** B-Bolivia, an Allele of the Maize b1 Gene with Variable
899 Expression, Contains a High Copy Retrotransposon- Related Sequence Immediately Upstream1.
900 *Plant Physiology*: 1363–1379.

901 **Selinger DA, Lisch D, Chandler VL. 1998.** The maize regulatory gene B-Peru contains a DNA
902 rearrangement that specifies tissue-specific expression through both positive and negative
903 promoter elements. *Genetics* **149**: 1125–1138.

904 **Sigmon B, Vollbrecht E. 2010.** Evidence of selection at the ramosa1 locus during maize
905 domestication. *Molecular ecology* **19**: 1296–1311.

906 **Sousaraei N, Torabi B, Mashaieghi K, Soltani E, Mousavizadeh SJ. 2021.** Variation of seed
907 germination response to temperature in tomato landraces: An adaptation strategy to
908 environmental conditions. *Scientia horticultrae* **281**: 109987.

909 **Steinhoff J, Liu W, Reif JC, Della Porta G, Ranc N, Würschum T. 2012.** Detection of QTL
910 for flowering time in multiple families of elite maize. *TAG. Theoretical and applied genetics. Theoretische und angewandte Genetik* **125**: 1539–1551.

912 **Stern DL. 2013.** The genetic causes of convergent evolution. *Nature reviews. Genetics* **14**: 751–
913 764.

914 **Taylor J, Butler D. 2017.** R Package ASMap: Efficient Genetic Linkage Map Construction and
915 Diagnosis. *Journal of Statistical Software, Articles* **79**: 1–29.

916 **Todesco M, Balasubramanian S, Hu TT, Traw MB, Horton M, Epple P, Kuhns C,**
917 **Sureshkumar S, Schwartz C, Lanz C, et al. 2010.** Natural allelic variation underlying a major
918 fitness trade-off in *Arabidopsis thaliana*. *Nature* **465**: 632–636.

919 **Troyer AF. 1999.** Background of U.S. Hybrid Corn. *Crop science* **39**:
920 crops1999.0011183X003900020001x.

921 **Ureta C, Martínez-Meyer E, Perales HR, Álvarez-Buylla ER. 2012.** Projecting the effects of
922 climate change on the distribution of maize races and their wild relatives in Mexico. *Global*
923 *change biology* **18**: 1073–1082.

924 **Vega-Arreguín JC, Ibarra-Laclette E, Jiménez-Moraila B, Martínez O, Vielle-Calzada JP,**
925 **Herrera-Estrella L, Herrera-Estrella A. 2009.** Deep sampling of the Palomero maize
926 transcriptome by a high throughput strategy of pyrosequencing. *BMC genomics* **10**: 299.

927 **Verhoeven KJF, Poorter H, Nevo E, Biere A. 2008.** Habitat-specific natural selection at a
928 flowering-time QTL is a main driver of local adaptation in two wild barley populations.
929 *Molecular ecology* **17**: 3416–3424.

930 **Verhoeven KJF, Vanhala TK, Biere A, Nevo E, van Damme JMM. 2004.** The genetic basis
931 of adaptive population differentiation: A quantitative trait locus analysis of fitness traits in two
932 wild barley populations from contrasting habitats. *Evolution; international journal of organic*
933 *evolution* **58**: 270.

934 **Vielle-Calzada J-P, Martínez de la Vega O, Hernández-Guzmán G, Ibarra-Laclette E,**
935 **Alvarez-Mejía C, Vega-Arreguín JC, Jiménez-Moraila B, Fernández-Cortés A, Corona-**
936 **Armenta G, Herrera-Estrella L, et al. 2009.** The Palomero genome suggests metal effects on

- 937 domestication. *Science* **326**: 1078.
- 938 **Wang L, Josephs EB, Lee KM, Roberts LM, Rellán-Álvarez R, Ross-Ibarra J, Hufford**
 939 **MB. 2021.** Molecular Parallelism Underlies Convergent Highland Adaptation of Maize
 940 Landraces. *Molecular biology and evolution*.
- 941 **Warburton ML, Reif JC, Frisch M, Bohn M, Bedoya C, Xia XC, Crossa J, Franco J,**
 942 **Hoisington D, Pixley K, et al. 2008.** Genetic Diversity in CIMMYT Nontemperate Maize
 943 Germplasm: Landraces, Open Pollinated Varieties, and Inbred Lines. *Crop Science* **48**: 617–624.
- 944 **Weinig C, Dorn LA, Kane NC, German ZM, Halldorsdottir SS, Ungerer MC, Toyonaga Y,**
 945 **Mackay TFC, Purugganan MD, Schmitt J. 2003.** Heterogeneous selection at specific loci in
 946 natural environments in *Arabidopsis thaliana*. *Genetics* **165**: 321–329.
- 947 **Wellhausen EJ, Roberts LM, Hernandez-X. H. 1951.** *Razas de Maíz en México, su origen,*
 948 *características y distribución* (PC Mangelsdorf, Ed.). Secretaría de Agricultura y Ganadería de
 949 México D. F.; Fundación Rockefeller.
- 950 **Wilkes HG. 1972.** Maize and its wild relatives. *Science* **177**: 1071–1077.
- 951 **Xu G, Wang X, Huang C, Xu D, Li D, Tian J, Chen Q, Wang C, Liang Y, Wu Y, et al. 2017.**
 952 Complex genetic architecture underlies maize tassel domestication. *The New phytologist* **214**:
 953 852–864.
- 954 **Zeven AC. 1998.** Landraces: A review of definitions and classifications. *Euphytica/ Netherlands*
 955 *journal of plant breeding* **104**: 127–139.
- 956 **FUNDING SOURCES**
- 957 Consejo Nacional de Ciencia y Tecnología [Mexico] (FOINS-2016-01)
 958 Consejo Nacional de Ciencia y Tecnología [Mexico] (CB-2015-01 254012)
 959 Secretaria de Agricultura y Desarrollo Rural (Ministry of Agriculture and Rural Development:
 960 SADER) of the Government of Mexico under the MasAgro (Sustainable Modernization of
 961 Traditional Agriculture) initiative
 962 National Science Foundation [USA] (No. 1546719)
 963 UC-MEXUS (CN-15-1476)
 964 RJHS is supported by the USDA National Institute of Food and Agriculture and Hatch
 965 Appropriations under Project #PEN04734 and Accession #1021929.

Table 1: Description of phenotypic traits measured at two locations. The significance of location effect was calculated using a Wilcoxon test for count and scale data (i.e., TBN, KPR, KRN, P_INT and Hair_score), and likelihood ratio test was conducted for the other continuous traits. Note: *: $p < 0.05$; **: $p < 0.001$; *: $p < 0.0001$.**

Phenotypic traits	Description	Unit	Mean (MT)	Mean (VB)	Significance of E
DTA	Days to anthesis	Day	131.07	59.69	< 0.0001***
DTS	Days to silking	Day	131.30	60.15	< 0.0001***
ASI	Anthesis-silking interval	Day	0.46	0.30	0.5650
PH	Plant height	cm	122.72	148.65	0.0546
TBN	Tassel branch number	Count	4	5	0.1796
EW	Ear weight	g	19.60	42.69	0.0019**
EL	Ear length	cm	7.30	8.75	0.0860
ED	Ear diameter	cm	3.18	3.83	0.0083**
EH	Ear height	cm	49.68	65.30	0.0260*
KRN	Number of kernel rows	Count	14	17	0.0157*

KPR	Number of kernels per row	Count	13	20	0.0275*
TKW	Total kernel weight	g	15.31	36.50	0.0002**
TKN	Total kernel number	Count	94.44	210.14	0.0001**
TL	Tassel length	mm	21.65	28.35	0.0573
P_INT	Anthocyanin pigment intensity	visual 0-4 code scale	2	0	0.5303
MH	Macrohair score	Hair pattern ≤ 1, hair score =0; Hair pattern >1, hair score = hair density	0	0	---

971
972
973

Table 2: QTLs detected in the B73 x PT BC₁S₅ RIL population for the phenotypes (Trait, abbreviated as in Table 1) and datasets (Set Detected) analyzed. Marker column describes the marker linked to the QTL, chr: chromosome; Pos: genetic position of the QTL LOD peak (cM); P_pos: physical position of the QTL LOD peak (MB, Reference genome v4), SI: Support Interval of the QTL (MB); GxE: indicates if the QTL is detected in the GxE data set; %VE: additive variance explained by the QTL in the multi-QTL model.

QTL	Marker	Chr	pos	p_pos	Set Detected	SI	GxE	% VE
qASI1	1_53029437	1	30.46	53.03	VB	35.2 - 292.18		10.13
qASI2	2_241675850	2	77.91	241.68	GEN, VB	240.52 - 244.41		10.88 - 14.54
qASI3	3_229390098	3	65.63	229.39	GEN	11.85 - 234.87		7.29
qASI8	8_135484500	8	24	135.48	GEN, VB, MT	134.32 - 164.79		15.96 - 18
qDTA1	1_286172395	1	94.35	286.17	VB, GxE	2.81 - 306.46	*	6.07 - 10.19
qDTA6	6_166664744	6	22	166.66	GEN, MT, GxE	165.39 - 168.82	*	7.5 - 13.53
qDTA7	7_165928844	7	48.37	165.93	MT	0.84 - 173.75		5.51
qDTA8a	8_21391040	8	0	21.39	MT	25.12 - 97.57		9.45
qDTA8b	8_153580487	8	27.5	153.58	GEN, VB, MT, GxE	148.95 - 161.29	*	9.67 - 21.23
qDTS1	1_12171293	1	12.47	12.17	GxE	2.81 - 306.46	*	5.18
qDTS6	6_166506716	6	23	166.51	GEN, MT, GxE	165.39 - 168.12	*	7.59 - 15.18
qDTS7	7_163657186	7	48	163.66	MT	15.91 - 169.56		8.16
qDTS8	8_161289005	8	31	161.29	GEN, MT, GxE	133.04 - 169.05	*	11.17 - 16.07
qED4	4_179539232	4	41.65	179.54	GEN, VB	163.36 - 196.26		14.69 - 14.96
qED5	5_190812247	5	31.54	190.81	VB	1.48 - 198.95		9.2
qED8	8_97570666	8	7.81	97.57	GxE	21.39 - 112.91	*	14.86
qEH1	1_177987239	1	53.68	177.99	GEN, MT	162.88 - 198.3		11.87 - 14.99
qEH7	7_135399817	7	33.26	135.4	GEN, VB, MT	131.23 - 144.37		14.94 - 16.76
qEL3	3_190627575	3	43.33	190.63	MT	11.5 - 234.87		13.39
qEL4	4_74666833	4	23.02	74.67	GxE	18.12 - 161.42	*	12.34

qEL7	7_172341845	7	55	172.34	VB	166.94 - 181.12		12.16
qEL8	8_175513459	8	43.5	175.51	GEN, VB, GxE	172.04 - 179.51	*	9.25 - 14.81
qEW7	7_20110508	7	16.9	20.11	GxE	9.14 - 28.5	*	14.33
qEW8	8_110436593	8	13.45	110.44	GxE	88.36 - 170.14	*	11.6
qMH3	3_156124621	3	28.83	156.12	GEN, VB	11.85 - 161.43		9.43 - 11.07
qMH7	7_155328976	7	43.1	155.33	GEN, MT	152.38 - 164.52		14.21 - 17.8
qMH8	8_122414801	8	18.71	122.41	VB, GxE	115.01 - 123.81	*	15.42 - 17.91
qMH9	9_65716548	9	12.5	65.72	GEN, VB	20.72 - 124.18		16.61 - 19
qKPR8	8_135484500	8	24	135.48	GxE	115.01 - 164.79	*	18.86
qKRN1	1_161556632	1	49.5	161.56	VB	8.68 - 296.87		12.63
qKRN8	8_133038186	8	23.42	133.04	GxE	82.76 - 164.79	*	16.39
qP_INT2	2_19456739	2	25.61	19.46	GEN, VB, MT	18.31 - 25.87		22.12 - 40.03
qP_INT10	10_11796399 5	10	19.49	117.96	GEN, VB, MT	116.16 - 138.71		9.47 - 13.37
qPH1a	1_197051908	1	56.5	197.05	GEN, VB, MT	176 - 199.71		18.88 - 20.43
qPH1b	1_293830327	1	100.1 3	293.83	VB	283.08 - 299.44		12.26
qPH8	8_149541560	8	27.58	149.54	VB	142.92 - 164.79		13.69
qPH10	10_13688363 9	10	26	136.88	GxE	10.27 - 143.76	*	11.17
qTBN2	2_141142660	2	40.88	141.14	GEN	70.9 - 184.93		13.24
qTBN7	7_121548061	7	24.52	121.55	GEN, VB, GxE	112.71 - 121.69	*	15.86 - 33.17
qTKN7	7_15912522	7	15.35	15.91	MT	9.14 - 121.69		17.09
qTKN8	8_126886782	8	20.81	126.89	GxE	82.76 - 170.04	*	12.86
qTKN9	9_124180939	9	19.6	124.18	VB	111.77 - 135.89		16.39
qTL1	1_227715124	1	72.51	227.72	GxE	224.77 - 244.86	*	19.73
qTL2	2_230998297	2	70.19	231	GEN, MT	0.94 - 241.68		11.56 - 11.62

Table S1: Environmental characteristics of the experimental sites in Metepec (MT) and Valle de Banderas (VB) located in Mexico.

Field site	Valle de Banderas (VB) Lowlands	Metepec (MT) Highlands
Elevation (m.a.s.l.)	54	2610
State	Nayarit	Mexico State
Latitude	20.784	19.223
Longitude	-105.244	-99.547
Temperature Min/Mean/Max (°C)	22.5/25.8/28.4	12.3/12.4/17.1
Precipitation (mm)	1173	809
Type of soil	Regosol	Andosol




Figure 1. The highland environment impacts maize growth and productivity. A) A representative highland cultivated maize field at 3000 m.a.s.l. near the Nevado de Toluca volcano, State of Mexico, Mexico (19.121702, -99.660812). B) Representative ears of the US adapted inbred line B73 and the Mexican highland landrace Palomero Toluqueño (PT) grown at sea level (red bar; 54 m.a.s.l.; Valle de Banderas [VB], Nayarit, Mexico) or in the highlands (blue bar; 2610 m.a.s.l.; Metepec [MT], State of Mexico, Mexico). C) Effect of the highland environment on plant performance. Distribution of trait values for 122 B73xPT BC₁S₅ lines grown in the lowland (VB; red) and highland (MT; blue) field sites. Trait codes: DTA - days to anthesis (days); PH - plant height (cm); ED - ear diameter (cm); EW - ear dry weight (g); TKN - total kernel number; TKW - total kernel weight (g). Fitted values for each genotype and location combination were used to generate violin plots and were estimated by adding BLUPs of the G+GEI to the estimated location mean. Boxes represent the interquartile range with the horizontal line representing the median and whiskers representing 1.5 times the interquartile ranges. The shape of the violin plot represents probability density of data at different values along the y-axis.




Figure 2. Extensive GEI was observed for yield components. A) Distribution of *ear height* (EH, cm) in low (VB) and high elevation (MT) field sites. Boxes represent the interquartile range with the horizontal line representing the median and whiskers representing 1.5 times the interquartile range. The shape of the violin plot represents probability density of data at different values along the y-axis. B) Reaction norm plot for EH, showing little GEI. Values shown are G + GEI deviations from the field site average. Line segments connect values for each RIL genotype in the two field sites. B73 (yellow) and PT (green) parental values are shown. C), D) as B, showing extensive rank-changing GEI associated with *ear weight* (EW, g) and *total kernel weight* (TKW, g), respectively.




Figure 3. QEI interactions contribute to local adaptation. A,B) Schematic of QEI under models of A) *conditional* effect in one environment but not another or B) *antagonistic pleiotropy* showing a change in the sign of the QTL effect between environments. C) Heatmap representation of the standardized median G + GEI value for all families with a given genotype (B73 or PT) in lowland (VB) and highland (MT) sites, for the named QTL (see Table 2). Asterisks indicate QTL identified in the GEI model. Lowercase letters indicate Tukey means groups. D) LOD support for a conditional *ear weight* (EW) QTL (qEW7) on the short arm of chromosome (chr) 7. The QTL is well supported by data from the highland site (MT, blue trace) but not the lowland site (VB, red trace), and is captured by a multiQTL model for GxE (black trace). E) Scatter plot of EW in highland (MT) against lowland (VB) fields. Each RIL is represented by a single point, colored by genotype at qEW7 (yellow, B73; green, PT). RILs shown in F below are labelled. The solid line shows a linear fit through all points. Box plots parallel to the vertical and horizontal axes show the distribution by genotype in MT and VB, respectively. Boxes represent the interquartile range with the horizontal line representing the median, and whiskers extending 1.5 times the interquartile range. F) Ears of RILs LANMLR17B044 (#44) and LANMLR17B011 (#11) produced in lowland (red bar) and highland (blue bar) fields, showing marked differences in stability with respect to field. Points below the panel indicate QTL genotype.




Figure 4. A major QTL for tassel branch number co-localizes with the *Ramosa1* gene. In comparison with typical maize varieties (A), tassel branching is strongly reduced in Mexican highland maize (B). C) Distribution of tassel length (TL, cm) in low (VB) and high (MT) field sites. Boxes represent the interquartile range with the horizontal line representing the median and whiskers representing 1.5 times the interquartile range. The shape of the violin plot represents probability density of data at different values along the y-axis. D) Reaction norm plot for TL. Values shown are G + GxE deviations from the field site average. Line segments connect values for each RIL genotype in the two field sites. B73 (yellow) and PT (green) parental values are shown. E, F) as C and D for tassel branch number (TBN). For F, the plot shows the median for each genotype in each field. G) LOD support (multQTL model, G main effect) for a qTBN7 that co-localizes with the *Ramosa1* (*Ra1*) candidate gene. Red shading indicates a drop 2 LOD interval around the peak marker. H) Effect of the chr 7 TBN QTL showing trait values for families carrying B73 (yellow) or PT (green) alleles in lowland (VB) or highland (MT) field sites.




Figure 5. Stem macrohair production is promoted by multiple QTL. A) Mexican highland maize is characterized by extensive sheath pubescence. B) QTLs linked to macrohair score (MH) on chromosomes (chr) 3, 7, 8 and 9. Trace shows LOD support for the macrohair trait in the lowland field (VB), in the genotype main effect (G) or for GEI (GxE). Teosinte introgression on chr 9 reported by Hufford *et al.*, 2013 that includes the *mhl1* locus is marked by a black bar. C) QTL effect shown as the proportion (shown by circle diameter in the main plot) of RILs scored for different hair score values in a given genotypic class (B73 allele, yellow; PT allele, green). Panels show the effect of allele substitution at the stated QTL in the subset of RILs for which the other QTLs are fixed as B73 and the cumulative effect of increasing the number of PT alleles at qMH 3, 7, 8 or 9. Points below the panel indicate QTL genotype. D) Glabrous (*glab*) and pubescent (*hair*) near-isogenic siblings generated by selection for pubescent plants through five generations of backcrossing of a Mexican Conico highland landrace to B73.




Figure 6. Colocalization of QTL with SNPs showing elevational variation in Mexican landrace maize. A) Geographic distribution of Mexican maize landraces. The color gradient represents elevation of the associated sampling location of maize landrace accessions. B) Trace showing support (LOD, L) for an *ear diameter* (ED) QTL across chromosome 4 (chr 4; physical distance) and SNPs (red points) significantly ($-\log_{10}p$, p) associated with elevation in Mexican maize landraces. LOD profile drawn using physically anchored genetic markers and trait values from the lowland site (VB). The gray rectangle indicates the position of the previously characterized *inv4m* inversion polymorphism. C, D) as B, showing support for *ear weight* (EW) and *total kernel number* (TKN) QTL on chromosomes 7 and 8, respectively. LOD profiles associated with trait GEI (GxE) values.




Figure S1 - Supplemental 1 Distribution of plant phenotypic traits for B73xPT recombinant inbred lines grown in VB or MT field sites (trait descriptions were shown in Table 1). Fitted values for each genotype and location combination were used to generate violin plots and were estimated by adding BLUPs of the G+GEI to the estimated location mean. Median values were used to generate the violin plots for TBN, KPR, KRN. Boxes represent the interquartile range with the horizontal line representing the median and whiskers representing 1.5 times the interquartile ranges. The shape of the violin plot represents probability density of data at different values along the y-axis.




Figure S2 - Supplemental 1 Reaction norm plots of plant phenotypic traits for B73xPT recombinant inbred lines grown in VB or MT field sites (trait descriptions were shown in Table 1). Values shown are the sum of Best Linear Unbiased Prediction (BLUP) for genotype and genotype by location interaction effects for each genotype in two field sites. Gray line segments connect values for each RIL genotype in the two field sites. B73 and PT parental values are shown in thick yellow and green lines respectively.




Figure S3. Pubescent segregants from a B73xMi21 BC₅S₁ family contain Mi21 introgression on chromosome 3. Three glabrous (glab) and two pubescent (hair) individuals genotyped with DaRT-SEQ.




Figure S4. Elevation eGWAS Manhattan plot




Figure S5 - Supplemental 1: Reaction norms of B73 (green line) and Palomero Toluqueño landrace (yellow line) grown in VB and MT field sites (trait descriptions were shown in Table 1). The fitted values were estimated by adding BLUPs of the G+GEI to the estimated location mean. Palomero Toluqueño values were obtained by evaluating multiple heterozygous individuals from MEXI5 accession from CIMMYT.




Figure 8 - Supplemental 1: Monthly and accumulated precipitation, monthly average temperature and monthly and accumulated estimated Grow Degree Day patterns for VB and MT field sites for the years where evaluation was performed (2014-2015, 2015-2016 for VB field

site; 2015, 2016, 2018 and 2019 for MT field site). The precipitation (in mm) and monthly mean temperature (°C) data was obtained from CONAGUA (National Commission on Water) database (<https://smn.conagua.gob.mx/es/climatologia/informacion-climatologica/informacion-estadistica-climatologica>; estation #15266 for MT site; estation #18030 for VB site) and temperature data was complemented with <https://wu-next-ibm.wunderground.com/> where needed. Grow Degree Days estimation was performed with the formula $GDD = (T_{mean} - 10) * n$ where T_{mean} is the monthly average temperature and n is the number of days accumulated in that month. The blue areas describe the typical growing season in winter in VB site (november-april) and summer in MT site (march-october).