

1 **Androglobin, a chimeric mammalian globin, is required for male fertility**

2

3 Anna Keppner¹, Miguel Correia¹, Sara Santambrogio², Teng Wei Koay¹, Darko Maric¹,
4 Carina Osterhof³, Denise V Winter⁴, Angèle Clerc¹, Michael Stumpe⁵, Frédéric
5 Chalmel⁶, Sylvia Dewilde⁷, Alex Odermatt⁴, Dieter Kressler⁵, Thomas Hankeln³, Roland
6 H. Wenger², David Hoogewijs^{1*}

7

8 ¹ Section of Medicine, Department of Endocrinology, Metabolism and Cardiovascular
9 system (EMC), University of Fribourg, Switzerland; ² Institute of Physiology, University
10 of Zurich, Switzerland; ³ Institute for Organismic and Molecular Evolutionary Biology,
11 University of Mainz, Germany; ⁴ Department of Pharmaceutical Sciences, University of
12 Basel, Switzerland, ⁵ Department of Biology, University of Fribourg, Switzerland, ⁶
13 University of Rennes, Inserm, EHESP, Irset (Institut de recherche en santé,
14 environnement et travail) - UMR_S 1085, France, ⁷ Department of Biomedical Sciences,
15 University of Antwerp, Belgium

16

17 *Corresponding author: David Hoogewijs

18 Email: david.hoogewijs@unifr.ch

19 **Abstract**

20 Spermatogenesis is a highly specialised process, involving multiple dedicated pathways
 21 and regulatory check-points. Defects ultimately lead to male sub-fertility or sterility, and
 22 numerous aspects of mammalian sperm formation remain unknown. The predominant
 23 expression of the latest globin family member, androglobin (Adgb) in mammalian testis
 24 tissue prompted us to assess its physiological function in spermatogenesis. Adgb
 25 knockout mice display male infertility, reduced testis weight, impaired maturation of
 26 elongating spermatids, abnormal sperm shape and ultrastructural defects in microtubule
 27 and mitochondrial organisation. Epididymal sperm from Adgb knockout animals display
 28 multiple flagellar malformations including coiled, bifide or shortened flagella, and
 29 erratic acrosomal development. Following immunoprecipitation and mass spectrometry,
 30 we could identify septin 10 (Sept10) as interactor of Adgb. The Sept10-Adgb interaction
 31 was confirmed both *in vivo* using testis lysates, and *in vitro* by reciprocal co-
 32 immunoprecipitation experiments. Furthermore, absence of Adgb leads to
 33 mislocalisation of Sept10 in sperm, indicating defective manchette and sperm annulus
 34 formation. Finally, *in vitro* data suggest that Adgb contributes to Sept10 proteolysis in a
 35 calmodulin (CaM)-dependent manner. Collectively, our results provide evidence that
 36 Adgb is essential for murine spermatogenesis and further suggest that interdependence
 37 between Adgb and Sept10 is required for sperm head shaping via the manchette and
 38 proper flagellum formation.

39 **Introduction**

40 Spermatogenesis is a complex and tightly regulated process, involving germ cell
 41 differentiation, haploid cell formation and sperm maturation and elongation (Hermo et
 42 al. 2010a, b, c, d, e, Neto et al. 2016). In a first proliferative step, germ cells will either
 43 self-renew to maintain the pool of active progenitor cells, or commit to become
 44 spermatogonia. Spermatogonia then undergo mitosis, thereby generating primary
 45 spermatocytes, that will go through a first round of meiosis, forming secondary
 46 spermatocytes, and a second round of meiosis to generate spermatids (Mecklenburg and
 47 Hermann 2016). At this stage, DNA replication is arrested and all the genetic material
 48 (RNA) required for spermatid maturation and further development is stored in a
 49 specialized compartment known as the chromatoid body (Peruquetti 2015). During this
 50 maturation process, known as spermiogenesis, the spermatids undergo deep
 51 morphological changes, including the formation of the acrosome (the sperm head) from
 52 the Golgi apparatus before migration of the latter to the cytoplasmic droplet (Khawar,
 53 Gao, and Li 2019), condensation of the nucleus, and the sperm flagellum formation from
 54 the centriole, around which mitochondria will migrate to form the midpiece, the sperm
 55 annulus and the mobile tail (Hermo et al. 2010a, b, c, d, e). These different steps
 56 ultimately lead to male fertility and are governed by numerous factors, including
 57 hormonal, translational, post-translational and epigenetic events. Dysfunctions in any of
 58 these regulatory processes will have consequences on sperm quality, sperm motility, or
 59 fecundation capacity (Neto et al. 2016). However, numerous physiological and
 60 pathological aspects of spermatogenesis are still not fully understood, and about 30% of
 61 sperm abnormalities are still idiopathic (Fainberg and Kashanian 2019). Moreover,
 62 several studies suggest that semen quality and male fertility have declined in Western
 63 countries over the past 50 years, further urging the need of a better global understanding

64 of male gamete formation (Levine et al. 2017). This led in recent years to the
65 identification of a growing number of genes implicated in spermatogenesis and
66 appearing causative for partial or complete male infertility upon mutation (Bracke et al.
67 2018a).

68 Globins are small globular metallo-proteins, which have the capacity to reversibly bind
69 gaseous ligands via a typical 8 alpha-helical structure in which a heme prosthetic group
70 can be embedded. In mammals, five globin types exist: the well-established hemoglobin
71 (Hb) and myoglobin (Mb), neuroglobin (Ngb) in neuronal cells, cytoglobin (Cygb)
72 ubiquitously expressed in fibroblasts, and the more recently identified androglobin
73 (Adgb), predominantly expressed in mammalian testis tissue (Keppner et al. 2020).

74 Adgb is a chimeric protein containing an N-terminal calpain-like cysteine protease
75 domain, a central permuted functional globin domain (Bracke, Hoogewijs, and Dewilde
76 2018b), interrupted by a potential calmodulin (CaM)-binding IQ motif and a large 700
77 amino acid long C-terminal tail of unknown identity (Hoogewijs et al. 2012). Decreased
78 mRNA expression levels in infertile versus fertile men (Platts et al. 2007) suggest a
79 potential role of Adgb in spermatogenesis. Gene regulation and expression studies
80 further suggest an association of Adgb with ciliogenesis including flagellum formation
81 (Koay et al. 2021). However, the *in vivo* function of Adgb remains unexplored. In this
82 study we investigated the physiological function of Adgb during murine spermatogenesis
83 by generating and analyzing Adgb knockout mice. We show that Adgb is mainly
84 expressed in late steps of spermiogenesis, that it locates to the acrosome, the sperm
85 flagellum, the annulus and the midpiece, and that it is crucial for male fertility and sperm
86 formation. Furthermore, we demonstrate that Adgb interacts with septin10 (Sept10), and
87 that co-localization occurs from the first steps of acrosome formation onwards within the

- 88 sperm neck and annulus. Finally, *in vitro* data suggest that Adgb contributes to Sept10
- 89 proteolysis in a CaM-dependent manner.

90 Results

91 *Adgb* knockout mice display male infertility

92 A gene-trap strategy, provided by the Knockout Mouse Project (KOMP)(Skarnes et al.
93 2011), was applied to target exons 13 and 14 of the *Adgb* gene (**Fig S1**). The correct
94 targeting of ES cells was verified first by long-range PCR (**Fig. S1**), and second by
95 Southern blotting (**Fig. S1**). The targeted *Adgb*^{tm1a(KOMP)Wtsi} allele (*Adgb* tm1a mice),
96 generated on a C57BL/6N background, displays a gene-trap DNA cassette, which was
97 inserted into the twelfth intron of the *Adgb* gene. The gene trap consists of a splice
98 acceptor site, an internal ribosome entry site, a β -galactosidase reporter sequence, and a
99 neomycin resistance sequence. Breeding of *Adgb* tm1a mice with ubiquitously expressed
100 CMV Cre-deleter mice allowed generation of mice deficient for exons 13 and 14 but still
101 expressing the β -galactosidase reporter (*Adgb* tm1b mice) (**Fig. S1**). Furthermore,
102 mating of *Adgb* tm1a mice with Flp-deleter mice enabled the generation of conditional
103 floxed mice (*Adgb* tm1c) (**Fig. S1**). These mice were further crossed with CMV Cre-
104 deleter mice to generate the full knockout animals (*Adgb* tm1d) (**Fig. S1**), that were used
105 for all downstream applications if not otherwise stated. Genotyping was performed by
106 regular PCR (**Fig. S1**), and revealed no significant differences in the Mendelian
107 distribution of offspring for *Adgb* tm1d animals following interbreeding of heterozygous
108 parents (380 pups: +/+, n= 101; +/-, n= 168; -/-, n=111; $X^2 = 5.62$, $p > 0.05$, ns). The
109 genetic ablation of *Adgb* expression was further verified by RT-qPCR (**Fig. 1A**) and
110 immunoblotting (**Fig. 1B,C**). While female knockout mice displayed no fertility issues,
111 male knockout mice never generated offspring, indicative for infertility. Full penetrance
112 male infertility was also observed in homozygous tm1a and tm1b male mice, whereas
113 homozygous tm1c animals showed normal fertility (data not shown). Accordingly, the
114 testis weight was significantly reduced in knockouts (**Fig. 1D**) associated with decreased

115 serum testosterone levels (**Fig. 1E**). Stage-specific histological examination of
 116 seminiferous tubules of control animals revealed normal architecture, normal
 117 spermatogenic maturation steps, and the presence of mature sperm with flagella
 118 extending into the lumen (**Fig. 1F**). In contrast, in knockout animals, despite the
 119 presence of meiotic events (stages X-XII, **Fig. 1F**), no flagella could be observed during
 120 the spermiation stage (stage VII-VIII, **Fig. 1F**). The absence of mature sperm was
 121 accompanied by abnormally shaped heads, trapped stage 16 spermatids within the
 122 epithelium and the presence of cytoplasmic material filling the lumen of the tubules (**Fig.**
 123 **1F**). Within the cauda epididymis, knockout animals displayed accumulations of residual
 124 bodies, cytoplasmic material, shed germ cells and occasional abnormally shaped sperm
 125 heads, but an overall absence of mature sperm as compared to wildtype animals (**Fig.**
 126 **1G**). No differences were detected in knockout testes at mRNA levels of nitric oxide
 127 synthases 1-3 (NOS1-3) or superoxide dismutases 1-3 (SOD1-3), and the ratio of
 128 Bax/Bcl2 was unchanged compared to wildtype testes, suggesting no increase in
 129 apoptotic events or oxidative stress (**Fig. S2**).

130

131 *Absence of Adgb interferes with the maturation of elongating spermatids*

132 To determine the temporal expression pattern of Adgb during spermatogenesis, wildtype
 133 embryos and pups at different post-natal ages, corresponding to the stages of the first
 134 wave of spermatogenesis during puberty, were dissected and analysed by RT-qPCR.
 135 Whereas the expression of Adgb nearly remained undetectable until post-natal day 21
 136 (corresponding to the stage of round spermatids), Adgb mRNA levels drastically
 137 increased to reach a peak at post-natal day 25, coinciding with the first elongating
 138 spermatids (**Fig. 2A**). Bulk and single-cell RNA sequencing analysis in mouse and
 139 human datasets available at the ReproGenomics Viewer resource (Darde et al. 2015,

140 Darde et al. 2019) confirmed the conserved expression pattern of ADGB in spermatids
 141 (Green et al. 2018, Jégou et al. 2017, Lukassen et al. 2018, Wang et al. 2018) (**Fig. S3**).
 142 Accordingly, Adgb protein expression equally reached its peak at post-natal days 26 to
 143 28 (**Fig. 2B,C**). This finding was further confirmed by propidium iodide staining and
 144 FACS sorting on testis lysates of the different genotypes. While no variations could be
 145 detected for phases 2C (spermatogonia, secondary spermatocytes, testicular somatic
 146 cells), S-phase (pre-meiotic spermatogonia), 4C (primary spermatocytes) and 1C (round
 147 spermatids), an abnormal accumulation of elongating and elongated spermatids (phase
 148 H) could be detected in knockout animals, suggesting a blockade in the elongation
 149 process (**Fig. 2D**). Additionally, immunofluorescence (**Fig. 2E**), mRNA *in situ*
 150 hybridization (**Fig. 2F**) and X-gal (**Fig. 2G**) stainings of testis sections from wildtype
 151 and knockout mice confirmed the presence of Adgb within layers containing post-
 152 meiotic cells, and further intensifying towards the lumen and mature sperm (**Fig. 2E-G**).
 153 Moreover, in mature sperm, Adgb expression could be visualized within the midpiece
 154 and along the whole flagellum by both X-gal (**Fig. 2F**) and immunofluorescence (**Fig.**
 155 **2G**) stainings.

156
 157 *Adgb is required for proper assembly of microtubules and positioning of mitochondria*

158 To gain additional insights into the origin of male infertility, cauda epididymal sperm
 159 was collected from both wildtype and knockout mice, and visualized under a
 160 microscope. While wildtype sperm appeared normal, very few knockout spermatozoa
 161 were found and displayed various defects of the head and/or flagellum structure,
 162 including shortened or bifide tails, loopings of the flagellum, and immature acrosomal
 163 structures (**Fig. 3A**). Knockout sperm acrosome structure appeared partially conserved,
 164 with sperm displaying either normal acrosomes, or various defects in shape or staining

intensity, or total absence (**Fig. 3B**). TEM ultrastructural analysis of testis sections further revealed misshaped heads with nuclear inclusions, disorganised axonemes failing to arrange around the flagella, misaligned microtubules, defective manchette elongation, and chaotic mitochondria within the forming midpieces of elongating spermatids in knockout samples (**Fig. 3C**).

The Adgb-dependent transcriptome reveals dysregulation of multiple spermiogenesis genes

To understand the molecular consequences of loss of Adgb in the testis, we performed RNA-sequencing experiments on total testis RNAs from wildtype and knockout mice at post-natal day 25. An elaborate set of significant differentially expressed genes (74 genes upregulated and 204 downregulated) was identified, underscoring the crucial function of Adgb in spermatogenesis (**Fig S4, Dataset S1**). Functional analysis based on Gene Ontology term enrichments confirmed that many of these genes are related to sperm head, acrosome reaction, acrosomal membrane, sperm motility, spermatid development and spermatid differentiation in line with the pronounced structural changes in spermatids during spermiogenesis. Intriguingly, a more refined Ingenuity pathway-based analysis (IPA) of the differentially expressed gene set provided a link to testosterone synthesis as deregulated pathway, as evidenced by reduced 17 β hsd3 and Lhcgr mRNA levels in knockout mice and consistent with the observed decrease in serum testosterone levels (**Fig. 1E, Fig. S4**).

Adgb interacts and co-localizes with Sept10

To obtain more insights into the physiological function of Adgb, we explored the Adgb-dependent interactome. Total protein extracts from wildtype testes were

immunoprecipitated (IP) with anti-Adgb antibody or IgG and subsequently submitted to mass spectrometry (MS) analysis to reveal potential interacting proteins of Adgb. Among the specifically enriched proteins, there were various members of the septin family of proteins, such as Sept10, Sept11, Sept2, and Sept7 (**Fig. 4A and Dataset S2**). Particular focus was put on Sept10 for further downstream experiments given its strong enrichment combined with high abundance in the immunoprecipitation. To confirm the interaction between Adgb and Sept10, reciprocal co-immunoprecipitation (co-IP) experiments were performed (**Fig. 4B,C**) on tissue extracts from wildtype and knockout testes (**Fig. 4B**) and in HEK293 cells overexpressing full-length ADGB (**Fig. 4C**) and Sept10. The results demonstrate that Adgb and Sept10 interact both *in vivo* (**Fig. 4B**) and *in vitro* (**Fig. 4C**), whereas in testis lysates of Adgb-deficient mice, no Sept10 precipitation was observed (**Fig. 4B**). Endogenous Sept10 protein levels were equal in testis lysates of Adgb-deficient and wildtype mice. Similar observations were made for Sept11, Sept7, and Sept2, as well as for other septins that are crucial for spermatogenesis, including Sept8, Sept9 and Sept14 (**Fig. S5**). We next investigated whether the interaction with Sept10 occurs at the N-terminal or the C-terminal portion of Adgb (**Fig. 4D,E**). Following co-overexpression of Adgb deletion constructs with SEPT10 and subsequent co-immunoprecipitation, immunoblotting revealed that both parts of ADGB interact with SEPT10 (**Fig. 4D,E**), and that this interaction remained intact also upon deletion of the coiled-coil domain of ADGB (**Fig. S6**). Consistent with a functional interaction, the temporal expression profiles of SEPT10 and ADGB substantially overlapped as illustrated by analysis of bulk and single-cell RNA sequencing datasets of mouse and human RNA (**Fig. S3**) as well as by RT-qPCR and immunoblotting of mouse tissue samples (**Fig. S7**). The localization of Adgb and Sept10 was assessed in microdissected tubules and in epididymal sperm by

immunofluorescence. Co-localization of Adgb and Sept10 was visible during the Golgi phase in the acrosomal granule, during manchette and sperm tail formation (**Fig. 5A**), and at the level of the sperm annulus in mature wildtype sperm (**Fig. 5B**). A moderate Sept10 staining was also observed in the neck region of mature sperm (**Fig. 5B**). In knockout epididymal sperm, only a single signal, likely corresponding to the annulus, was observed and displayed abnormal migration, indicating defective manchette or microtubule formation (**Fig. 5B**). The migration of the annulus drives mitochondrial placement along the forming mid-piece (Toure et al. 2011). Since knockout animals displayed abnormal ultrastructural mitochondria organisation (**Fig. 3C**), CoxIV staining was performed on wildtype and knockout epididymal sperm. As expected, a robust staining was observed along the whole midpiece in wildtype sperm, whereas mitochondria were barely visible and formed a cloudy structure around the neck region of knockout sperm (**Fig. 5B**).

228

229 *ADGB contributes to Sept10 proteolytic cleavage in vitro*

To analyse the functional consequences of the SEPT10-ADGB interaction we transiently co-overexpressed both proteins in HEK293 cells. Intriguingly, apart from the intact form of overexpressed SEPT10 at 60 kDa, increased levels of a lower band of 37 kDa were consistently detected in the presence of exogenous ADGB (**Fig. 6A**), in a dose-dependent manner (**Fig. 6A**). Immunoblotting with a V5-antibody upon co-overexpression of a C-terminal V5-tagged SEPT10 with ADGB (**Fig. 6B**) as well as the presence of this band upon SEPT10/ADGB co-IP (**Fig. 6B, Fig. S8**) further support its origin as proteolytic SEPT10 product. To investigate a potential influence of the globin domain this experiment was repeated under normoxic and hypoxic conditions (0.2% O₂) but no differences were observed upon exposure to hypoxic conditions (**Fig. 6C**). To

240 determine a potential role of calmodulin we constructed a deletion mutant lacking the IQ
241 domain. Notably, transient overexpression of IQ-mutant ADGB resulted in considerably
242 reduced appearance of the 37 kDa band relative to WT ADGB (**Fig. 6D**). This finding
243 prompted us to experimentally validate the suspected CaM-ADGB interaction. Whereas
244 co-IP experiments following overexpression of full-length ADGB did not interact with
245 CaM under basal experimental conditions in HEK293 cells (data not shown), a truncated
246 construct covering the globin and IQ domains displayed robust ADGB-CaM interaction
247 (**Fig. 6E**). Consistently, MS analysis of proteins that were present in the IP of the
248 overexpressed, isolated globin domain revealed a prominent enrichment of endogenous
249 CaM (**Fig. S9 and Dataset S3**). Importantly, individual or double mutation of the
250 proximal histidine (HisF8) or distal glutamine (GluE7), critical residues in the globin
251 domain for heme coordination, did not alter the interaction, suggesting that the ADGB-
252 CaM interaction occurs independently of heme incorporation (**Fig. 6F, Fig. S10**). As an
253 additional layer of support for the ADGB-CaM interaction, chimeric Gal4 DNA-binding
254 domain and VP16 transactivation domain-based fusion constructs were generated for
255 mammalian 2-hybrid (M2H) luciferase reporter gene assays (**Fig. 6G,H**). These M2H
256 assays performed in 2 different cell lines, HEK293 and A375, revealed up to 7.5-fold
257 increase in luciferase activity upon co-transfection of both chimeric proteins compared to
258 single construct transfections, providing independent evidence that ADGB interacts with
259 CaM. To further assess the potential O₂-dependency of the CaM-ADGB interaction, we
260 repeated the M2H assays in A375 cells under hypoxic conditions. Exposure to hypoxic
261 conditions (0.2% O₂) did not alter luciferase activity, while luciferase activity of an *EPO*
262 promoter-driven reporter gene increased, suggesting that the ADGB-CaM interaction is
263 O₂-independent (**Fig. 6H**). This result is consistent with the maintained ADGB-CaM
264 interaction following mutation of critical residues in the globin domain in co-IP

265 experiments and unchanged ADGB-dependent cleavage of SEPT10 under hypoxic
266 conditions. Collectively, these *in vitro* data suggest a scenario in which exogenous
267 ADGB proteolytically contributes to cleavage of exogenous SEPT10 in an O₂-
268 independent but CaM-dependent manner.

269 Discussion

270 In the present study, we aimed to explore the testicular function of Adgb by generating
 271 and analysing Adgb^{-/-} constitutive knockout mice. Our results demonstrate that Adgb is
 272 required for late stages of spermatogenesis and male fertility, as evidenced by a total lack
 273 of functional sperm in knockout animals. Also, a reduction in testis weight, reduced
 274 testosterone production and dysregulated testosterone-associated genes such as Lhcgr
 275 and 17βhsd3 were observed. This finding is remarkable as Adgb displays little to no
 276 expression levels in the testosterone-producing Leydig cells. Our data further show that
 277 Adgb is indispensable for proper microtubule formation, mitochondria localisation and
 278 annulus positioning. Spermatid elongation is marked by profound morphological
 279 changes, which start with the formation of the acrosome, concomitant with the first
 280 expression of Adgb. Adgb mRNA and protein are detectable from post-natal day 21
 281 onwards, the stage at which the acrosomal granule is formed (Clermont and Leblond
 282 1955, Khawar, Gao, and Li 2019). Accordingly, a strong signal was visible within the
 283 acrosomal granule, and Adgb could be detected in all steps of spermatogenesis within
 284 the acrosome, including in mature sperm. Moreover, the defects in head shape and
 285 acrosome structure visible in Adgb mutant mice strongly suggest a function of Adgb in
 286 the maturation of the acrosome.

287 To understand the molecular mechanisms by which Adgb may exert its function during
 288 spermatogenesis we analysed the interactome of Adgb by LC-MS/MS after
 289 immunoprecipitation of the latter in testicular lysates. Strikingly, several members of the
 290 septin family ranked in the top hits. Septins are conserved GTPases that have the ability
 291 to form large oligomers and filamentous polymers and which associate with cell
 292 membranes and with the cytoskeleton. They serve as scaffolds for the proper localization
 293 of intracellular proteins via their diffusion barrier-forming characteristic. All septins

display a conserved structure formed by an N-terminal proline-rich polybasic region which interacts with membrane phospholipids, a central GTP-binding domain, a C-terminal region of unknown function (the septin unique element) further flanked by a C-terminal tail containing α -helical coiled coils of varying sizes enabling the polymer-forming protein-protein interactions typical of septins (Dolat, Hu, and Spiliotis 2014). Thirteen septins have been described so far in mammals (Sept1-12 and Sept14), further sub-classified into 4 distinct families depending on their biochemical and biophysical properties (Kinoshita 2003). Sept10 belongs to the Sept6 group (containing Sept6, Sept8, Sept10, Sept14, as well as Sept11, the latter also present in the Adgb interactome), which - unlike the other septin members - are catalytically inactive and constitutively bound to GTP (Sirajuddin et al. 2009). In sperm, Sept1, 4, 6, 7, and 12 have been localized to the sperm annulus, where they polymerize to a filamentous structure, forming a barrier between the midpiece and the principal piece of the spermatozoon (Toure et al. 2011). In this study, we focused on Sept10, and detected a strong signal in acrosomal granules in S7 spermatids, in the migrating annulus of S12 spermatids as well as in the annulus and neck region of mature sperm together with Adgb, suggesting a very early interaction between the two proteins, resulting in a missing signal (likely the annulus) upon absence of Adgb in knockouts. It is noteworthy that two independently generated mouse models for Sept4^{-/-} as well as Sept12^{+/-} mice are male infertile and harbour disorganized sperm mitochondria (Ihara et al. 2005, Kissel et al. 2005, Lin et al. 2009). Moreover, Sept4^{-/-} mice display a bent sperm tail and absence of annulus (Ihara et al. 2005, Kissel et al. 2005), whereas Sept12^{+/-} mice exhibit broken acrosomes, misshaped nuclei and increased apoptosis of germ cells (Lin et al. 2009). Correspondingly, SEPT12 mutations have been described in infertile men displaying abnormal sperm including defective annulus with a bent tail (Kuo et al. 2012). Additionally, defective sperm head morphology and DNA

integrity have recently been reported for two different SEPT14 missense mutations (Wang et al. 2019, Lin et al. 2020). Previous findings proposed a septin ring assembly as octameric filaments at the annulus, classically composed of septins 12-7-6-2-2-6-7-12 or 12-7-6-4-4-6-7-12, in which septins 12 and 7 and septins 6 and 2/4 are connected through their GTP-binding domain (G-interface), whereas septins 7 and 6 and the two central septins 2/4 interact via their NC-termini (NC-interface) (Sirajuddin et al. 2007, Kuo et al. 2015). It was further proposed that septins from the same sub-group can substitute each other (Sellin et al. 2011). It is therefore tempting to speculate that Sept10 might replace Sept6 in the ring, thus Adgb's localisation at the annulus. The identification of Sept7 and Sept2 as additional Adgb-interacting proteins is in favour of this assumption. Interestingly, another ring-like septin structure was recently described at the sperm neck, composed of Sept12 which complexes together with Sept1, Sept2, Sept10 and Sept11. Two mutations of Sept12 identified in patients disrupted the complex, leading to unstable head-tail junctions and defective connecting piece formation. Strikingly, the mutation of Sept12 and the subsequent disruption of the complex led to loss of Sept10 signal in the annulus (Shen et al. 2020) as also observed in Adgb-deficient mice. Accordingly, our data suggest an interdependence between Adgb and Sept10, which is required for the maintenance of the annulus, head shaping and proper mitochondrial localisation.

Septins have also been identified as components of cilia, which are hair-like microtubular protrusions at the surface of various cell types. Sept2 forms a diffusion barrier at the base of the cilium, impeding ciliary formation through loss of Sonic Hedgehog signalling when depleted (Hu et al. 2010). Sept2/7/9 form a complex that associates with the ciliary axoneme, thereby regulating ciliary length (Ghossoub et al. 2013). Accordingly, septin association with the cytoskeleton and particularly with

344 microtubular structures has been extensively studied (Spiliotis and Nakos 2021), and
345 numerous other cilia-related proteins participate in sperm flagellum formation.
346 Furthermore, most ciliopathies include male infertility and immotile sperm due to
347 defective axonemal organisation (Brown and Witman 2014). Likewise, Adgb knockout
348 mice display aberrant microtubule arrangements, thus it cannot be excluded that a
349 scaffolding and simultaneous regulatory action between Adgb and Sept10 is necessary to
350 support microtubular structure. Moreover, a recent study reported the consistent presence
351 of Adgb in the ciliomes of three distinct evolutionary ancestral taxa, further suggesting a
352 conserved function related to microtubular organisation and likely flagella formation
353 (Sigg et al. 2017). In line with this study, our recent *in vitro* investigations have
354 demonstrated that Adgb is transcriptionally regulated by FoxJ1, a master regulator of
355 ciliogenesis (Koay et al. 2021).

356 A robust Adgb-Sept10 interaction was also detected by co-immunoprecipitation
357 experiments, both in testis lysates as well as in HEK293 cells overexpressing ADGB and
358 SEPT10. Unexpectedly, interaction was still observed upon truncating the N-terminal
359 and C-terminal portions of ADGB. Furthermore, specific mutation of the coiled-coil
360 domain of either ADGB or SEPT10 did not disrupt the interaction, suggesting that
361 interaction of ADGB and SEPT10 occurs at various locations along the two proteins and
362 that the proteins may entangle around each other. This close interaction may serve as
363 targeting mechanism for the proteolytic processing of SEPT10 that we could observe
364 upon ectopic expression of both proteins. The unique chimeric domain structure of Adgb
365 with an N-terminal calpain-like protease domain and an IQ motif within a rearranged
366 globin domain suggests a CaM-mediated regulation. Indeed, we could demonstrate that
367 ADGB interacts with CaM via its IQ binding motif, and that the IQ binding motif seems
368 pivotal in the observed proteolytic cleavage of SEPT10. CaM is a versatile protein,

which can interact with a broad range of target proteins and act in a wide variety of cellular pathways and processes (Villalobo et al. 2018). Calmodulin-bound calcium represents a crucial activator of proteolytic activity of Ca^{2+} -dependent calpain proteases (Villalobo, González-Muñoz, and Berchtold 2019, Bähler and Rhoads 2002). Most interestingly, calpain-dependent proteolytic cleavage of septins is not unprecedented: it was shown that Sept5 is a substrate of both calpain-1 and calpain-2 in platelets, where the cleavage triggers secretion of chemokine-containing granules (Randriamboavonjy et al. 2012). Additionally, it was reported that small modifications in septin structures may have profound consequences on the protein's function. For example, a single point mutation within the N-terminus of Sept2 leads to homomeric filament formation without including any other septin type (Kim et al. 2012). Furthermore, the PKA-dependent phosphorylation of Sept12 at a single site leads to the disruption of the Sept12 filament, which dissociates from Sept7-6-2 and Sept7-6-4 complexes (Shen et al. 2017). Finally, SUMOylation failures are associated with aberrant filament formation for Sept6, 7 and 11, which are still able to interact with other endogenous septin members, remarkably in a stronger way than when properly SUMOylated (Ribet et al. 2017). It is therefore conceivable that CaM-dependent Adgb-mediated proteolytic cleavage of Sept10 may be a prerequisite for proper Sept10 function or localization within the sperm neck or annulus.

The presence of an oxygen-binding globin might be beneficial in a strongly hypoxic environment such as the testis. Indeed, testicular interstitial oxygen levels are very low, and tissue oxygen pO_2 is estimated to represent only 20% of the testicular artery pO_2 (Free, Schluntz, and Jaffe 1976), whereby the microvasculature controls the oxygen supply to the testicular tissue and thereby indirectly also controlling the oxygen reaching the large seminiferous tubules by diffusion, only (Reyes et al. 2012). We previously

394 demonstrated heme hexa-coordination of Adgb (Bracke, Hoogewijs, and Dewilde
395 2018b), a unique characteristic also reported for mammalian Cygb and Ngb, rather
396 associated with functionality beyond canonical oxygen transport and storage. Similar to
397 the postulated cytoprotective functions against ROS of Cygb and Ngb (Burmester and
398 Hankeln 2009, Burmester and Hankeln 2014, Randi et al. 2020, Keppner et al. 2020), it
399 is tempting to speculate that Adgb might be involved in detoxification of harmful ROS
400 as male infertility is affected by ROS, and spermatozoa are particularly sensitive to
401 ROS-induced damage. The lack of Adgb-dependent differential regulation of redox
402 sensitive genes on transcriptome or single gene level between wildtype and Adgb-
403 deficient bulk testis tissue lysates argues against, but does not exclude, an anti-oxidative
404 function of Adgb. In line with these observations, the absence of any O₂-dependent
405 effect of our *in vitro* findings on ADGB-dependent SEPT10 cleavage suggests that the
406 proteolytic activity of Adgb is independent of a functional globin domain. Although we
407 reported heme incorporation on recombinantly expressed human protein (Bracke,
408 Hoogewijs, and Dewilde 2018b), Adgb orthologs in more basal organisms lack the
409 crucial proximal His, suggesting that heme-binding might not be the most prominent
410 characteristic feature of Adgb. Establishing a mechanistic explanation for the chimeric
411 domain structure of the Adgb protein will remain a major challenge for the future.

412

413 In conclusion our study is the first to demonstrate a functional role for androglobin, the
414 fifth mammalian globin. We present convincing *in vivo* evidence that Adgb is required
415 for murine spermatogenesis. Interdependence between Adgb and Sept10 is necessary for
416 sperm head shaping via the manchette and proper flagellum formation. *In vitro* data
417 demonstrated CaM binding to ADGB and suggested that ADGB contributes to
418 proteolytical cleavage of SEPT10 in a CaM-dependent manner. Our work provides a

419 crucial contribution to the characterization of the physiological role of this novel

420 enigmatic chimeric globin type.

421

422 **Materials and methods**

423 *Animals, ethics statement and genotyping*

424 All experimental procedures and animal maintenance followed Swiss federal guidelines
 425 and the study was revised and approved by the “*Service de la sécurité alimentaire et des*
 426 *affaires vétérinaires*” (SAAV) of the canton of Fribourg, Switzerland (license number
 427 2017_16_FR). Animals were housed in rooms with a 12 hour/12 hour light/dark cycle,
 428 controlled temperature and humidity levels, and had free access to food and water.
 429 Interbreeding of heterozygous animals was performed to obtain wildtype (+/+),
 430 heterozygous (Tg/+ for tm1a and tm1b, or +/- for tm1d), and homozygous/knockout
 431 (Tg/Tg for tm1a and tm1b, or -/- for tm1d) littermates, that were experimentally used, if
 432 not otherwise stated, between 3 to 9 months of age. Genotyping of tm1a and tm1b
 433 animals was performed using the following primers (**Fig. S1**): F1 5’-
 434 CCGTGCCCAGCTATATGAGT-3’; R1 5’-CACAAACGGGTTCTTCTGTAGTCC-3’;
 435 R2 5’-CCAGCGGTGTTCTTTCTTA-3’. Primers for tm1d genotyping were the
 436 following (**Fig. S1**): F1, R2, and R4 5’-ACTGATGGCGAGCTCAGACC-3’. PCR
 437 amplification was performed for 36 cycles of 1 min at 95°C, 1 min at 56°C and 1 min at
 438 72°C. The PCR products were separated by electrophoresis on 2% agarose gels and
 439 visualized by ethidium bromide staining.

440

441 *Gene targeting and knockout mouse generation*

442 The Adgb^{tm1a(KOMP)Wtsi} (tm1a) strain was generated by blastocyst microinjection of
 443 embryonic stem (ES) cell clone EPD0707_3_H06, provided by the Knockout Mouse
 444 Project (KOMP) (Skarnes et al. 2011). Correct targeting of the Adgb locus was verified
 445 prior to microinjection by long-range PCR using primers 5’S 5’-
 446 CTGTACACTGGTTGTACACTGGTACAACACTG-3’; 5’AS 5’-

447 GGACTAACAGAAGAACCCGTTGTG-3'; 3'S 5'-
 448 CACACCTCCCCCTGAACCTGAAAC-3'; 3'AS 5'-
 449 GTACTTGATTGGACGATGATCCAAG-3' (**Fig. S1**), generating a band of 6.7 kb for
 450 5' primers, and 5.1 kb for 3' primers (**Fig. S1**). Targeted clones were confirmed by
 451 Southern blot analysis using a hybridization probe that targets exon 13 (**Fig. S1**)
 452 revealing a band of 4.1 kb (wildtype) or 3.2 kb (tm1a allele) following digestion of
 453 genomic DNA with *PvuII*, and a band of 2.8 kb (wildtype) or 2.4 kb (tm1a allele)
 454 following digestion of genomic DNA with *PstI* (**Fig. S1**). Chimeric mice were bred with
 455 C57BL/6-Tyr^{c-Brd} mice, and germline transmission in the F1 offspring was verified by
 456 PCR using primers F1, R1 and R2 (**Fig. S1**). The mice were further bred to C57BL/6N-
 457 Hprt^{Tg(CMV-cre)Brd/Wtsi} transgenic mice expressing the Cre allele to delete exons 13 and 14
 458 and the neo cassette to generate the Adgb^{tm1b(KOMP)Wtsi} strain (tm1b), or to C57BL/6N-
 459 Gt(ROSA)26Sor^{tm1(FLP1)Dym/Wtsi} transgenic mice expressing the Flp recombinase to delete
 460 the whole transgene cassette, thereby generating the Adgb^{tm1c(KOMP)Wtsi} strain. The latter
 461 were further crossed with C57BL/6N-Hprt^{Tg(CMV-cre)Brd/Wtsi} transgenic mice to delete
 462 exons 13 and 14, thereby generating the Adgb^{tm1d(KOMP)Wtsi} (tm1d, knockout) strain. The
 463 Cre recombinase allele was bred out before any experiments were performed.

464

465 *RNA extraction and RT-qPCR*

466 Testes were frozen in liquid nitrogen and stored at -80°C. Tissues were homogenized
 467 using a TissueLyser (Qiagen, Valencia, CA, USA). Subsequent RNA isolation and
 468 cDNA synthesis were performed as described previously (Keppner et al. 2019). In brief,
 469 RNA was extracted using an RNeasy Mini Kit (Qiagen,) and reverse-transcription (RT)
 470 was performed with 1.5 µg of total RNA and PrimeScript reverse transcriptase (Takara
 471 Bio Inc, Kusatsu, Japan). RT-quantitative (q)-PCR was performed on a CFX96 C1000

472 real-time PCR cycler (Bio-Rad Laboratories, Hercules, CA) using SYBRgreen PCR
473 master mix (Kapa Biosystems, London, UK). 21.5 ng of cDNA were loaded, and each
474 sample was run as duplicate. mRNA levels were normalized to β -actin as previously
475 described (De Backer et al. 2021). Primer sequences are displayed in **Table S1**.

476

477 *Cell culture and transfection*

478 HEK293 and A375 (ATCC CRL-1619) cells were maintained in Dulbecco's Minimum
479 Essential Media (DMEM) (Gibco, Life Technologies, Carlsbad, CA, USA), containing
480 L-glutamine, supplemented with 10% heat-inactivated fetal bovine serum FBS (PAN
481 Biotech, Aidenbach, Germany) and 100 Units/mL penicillin/100 μ g/mL streptomycin
482 (Gibco, Life Technologies, Carlsbad, CA, USA). Both cell lines were incubated in a
483 humidified 5% CO₂ atmosphere at 37°C and were routinely subcultured after
484 trypsinization. For hypoxic experiments, cells were seeded out in 6-well plates or 100-
485 mm culture dishes. The subsequent day, hypoxia experiments were carried out at 0.2%
486 O₂ and 5% CO₂ in a gas-controlled glove box (InvivoO2 400, Baker Ruskinn, Bridgend,
487 UK) for 24 hours. Transfection of HEK293 cells was performed using calcium-
488 phosphate (Jordan, Schallhorn, and Wurm 1996), with 2 μ g of plasmid DNA for regular
489 immunoblotting experiments, and 5 μ g of plasmid DNA for immunoprecipitation
490 experiments. Briefly, the DNA was diluted in sterile water and mixed with 250 mM
491 CaCl₂. 25 μ M chloroquine was added to the cells and allowed to incubate for a minimum
492 of 20 min. Prewarmed 37°C 2x HBS buffer pH 7.05 (NaCl 280 mM, KCl 10 mM,
493 Na₂HPO₄ 1.5 mM, D-glucose 12mM, HEPES 50mM) was added to the DNA solution
494 (50% v/v), and the transfection mixture was added dropwise to the cells. The medium
495 was replaced after 6 hours. Transfection of A375 cells was performed using JetOptimus

(Polyplus-transfection SA, Illkirch-Graffenstaden, France) according to the manufacturer's instructions.

498

499 *SDS-PAGE and immunoblotting*

500 Tissues were homogenized as described (Keppner et al. 2019). Cells were lysed in triton
501 buffer (Tris-HCl [20 mM, pH 7.4], NaCl [150 mM], triton X-100 [1%]), left on ice for
502 15 min, centrifuged, and the proteins were quantified by Bradford assay.
503 25 µg of proteins were separated by SDS-PAGE on 10% gels, and proteins were
504 electrotransferred to nitrocellulose membranes (Amersham Hybond-ECL, GE
505 Healthcare, Chicago, IL, USA). The membranes were incubated overnight at 4°C with
506 primary antibody (**Table S2**) and for 1 hour with donkey anti-rabbit or anti-mouse IgG
507 HRP-conjugated secondary antibody (1:5000, Amersham, Buckinghamshire, UK). All
508 antibodies were diluted in TBS-tween (1%) and dried milk (1%). The signal was
509 revealed using ECL Prime (Amersham, Buckinghamshire, UK) on a C-DiGit Western
510 blot scanner (LI-COR Biosciences), and quantified using ImageStudio program (LI-COR
511 Biosciences, Lincoln, NE, USA). The polyclonal anti-Adgb antibody was custom-made
512 (Proteintech Group Inc, Rosemont, IL, USA). A fusion protein immunogen raised
513 against the 409-745 amino acid region of mouse Adgb was used for the immunisation of
514 two rabbits over a period of 102 days. The antibodies in immune sera were affinity-
515 purified. Pre-bleeds, test bleeds and purified antibodies were tested and validated by
516 immunoblotting on wildtype and knockout testis extracts.

517

518 *Immunoprecipitation*

519 For immunoprecipitation (IP) of subsequent LC-MS/MS and immunoblotting analyses, 4
520 and 2 mg of proteins were used, respectively. The protein lysates were first pre-cleared

for 24 hours at 4°C with G-sepharose beads (GE Healthcare, Chicago, IL, USA) coupled to rabbit IgG (Bethyl Laboratories Inc, Montgomery, TX, USA). Samples were incubated overnight at 4°C with 2 µg primary antibody (**Table S2**) or 2 µg rabbit IgG, followed by 4 hours with G-sepharose beads, then washed 2 times with wash buffer (Tris-HCl [20 mM, pH 7.4], NaCl [300 mM], LAP [1 mM]), and 3 times with equilibration buffer (Tris-HCl [20 mM, pH 7.4], NaCl [150 mM], LAP [1 mM]). Samples were eluted by boiling for 5 min at 95°C in 2x sample buffer, and separated from the beads by centrifugation.

529

530 *LC-MS/MS analysis*

Washed IP beads were incubated with *Laemmli* sample buffer and proteins were reduced with 1 mM DTT for 10 min at 75°C and alkylated using 5.5 mM iodoacetamide for 10 min at room temperature. Protein samples were separated by SDS-PAGE on 4-12% gradient gels (ExpressPlus, Genscript, New Jersey, NJ, USA). Each gel lane was cut into 6 equal slices, the proteins were in-gel digested with trypsin (Promega, Madison, WI, USA), and the resulting peptide mixtures were processed on StageTips (Rappsilber, Mann, and Ishihama 2007, Shevchenko et al. 2006).

LC-MS/MS measurements were performed on a Q Exactive Plus mass spectrometer (Thermo Fisher Scientific, Waltham, MA, USA) coupled to an EASY-nLC 1000 nanoflow HPLC (Thermo Fisher Scientific). HPLC-column tips (fused silica) with 75 µm inner diameter were packed with Reprosil-Pur 120 C18-AQ, 1.9 µm (Dr. Maisch GmbH, Ammerbuch, Germany) to a length of 20 cm. A gradient of solvents A (0.1% formic acid in water) and B (0.1% formic acid in 80% acetonitrile in water) with increasing organic proportion was used for peptide separation (loading of sample with 0% B; separation ramp: from 5-30% B within 85 min). The flow rate was 250 nL/min

546 and for sample application 650 nL/min. The mass spectrometer was operated in the data-
547 dependent mode and switched automatically between MS (max. of 1×10^6 ions) and
548 MS/MS. Each MS scan was followed by a maximum of ten MS/MS scans using
549 normalized collision energy of 25% and a target value of 1000. Parent ions with a charge
550 state form $z = 1$ and unassigned charge states were excluded from fragmentation. The
551 mass range for MS was $m/z = 370-1750$. The resolution for MS was set to 70,000 and for
552 MS/MS to 17,500. MS parameters were as follows: spray voltage 2.3 kV; no sheath and
553 auxiliary gas flow; ion-transfer tube temperature 250°C. The MS raw data files were
554 uploaded into the MaxQuant software version 1.6.2.10 for peak detection, generation of
555 peak lists of mass error corrected peptides, and for database searches (Tyanova, Temu,
556 and Cox 2016). A full-length UniProt mouse (based on UniProt FASTA version April
557 2016) or human database (UniProt FASTA version March 2016) additionally containing
558 common contaminants, such as keratins and enzymes used for in-gel digestion, was used
559 as reference. Carbamidomethylcysteine was set as fixed modification and protein amino-
560 terminal acetylation and oxidation of methionine were set as variable modifications.
561 Three missed cleavages were allowed, enzyme specificity was trypsin/P, and the MS/MS
562 tolerance was set to 20 ppm. The average mass precision of identified peptides was in
563 general less than 1 ppm after recalibration. Peptide lists were further used by MaxQuant
564 to identify and relatively quantify proteins using the following parameters: peptide and
565 protein false discovery rates, based on a forward-reverse database, were set to 0.01,
566 minimum peptide length was set to 7, minimum number of peptides for identification
567 and quantitation of proteins was set to one which must be unique. The ‘match-between-
568 run’ option (0.7 min) was used.
569
570

571 *Propidium iodide staining and flow cytometry*

572 Preparation of germ cell suspensions was achieved as described (Jeyaraj, Grossman, and
573 Petrusz 2003). Briefly, decapsulated testes were incubated in 0.5 mg/mL collagenase
574 type IV in PBS, washed with PBS, and incubated in 1 µg/mL DNase and 1 µg/mL
575 trypsin. Soybean trypsin inhibitor was added, the suspension was filtered, washed in
576 PBS, fixed with 70% ethanol and stored at 4°C. DNA staining using propidium iodide
577 was performed as described (Krishnamurthy et al. 2000). Propidium iodide-stained cells
578 were analysed in a FACScan flow cytometer (Becton-Dickinson Immunocytometry, San
579 Jose, CA, USA). Cell populations were selected based on their DNA content, and their
580 relative numbers were calculated using Summit (Cytomation, CO, USA).

581

582 *Histological analyses and immunofluorescence*

583 Testes were fixed in 4% paraformaldehyde and embedded in paraffin. Preparation of
584 sections and H&E staining was performed as described (Keppner et al. 2015). Pictures
585 were taken using a Nikon Eclipse microscope (Nikon Corporation, Tokyo, Japan).
586 For immunofluorescence, testes were fixed in 4% paraformaldehyde (PFA) for at least 1
587 week, and incubated in 30% sucrose for another week. The testes were embedded in
588 Optimal Cutting Temperature compound (O.C.T. Tissue-Tek, Sakura Finetek, Tokyo,
589 Japan), and 5 µm thick sections were cut on a cryotome. For seminiferous tubule
590 dissections and stainings, slides were prepared as previously described (Kotaja et al.
591 2004). For sperm stainings, cauda epididymal sperm was retrieved and diluted in PBS. A
592 drop of the suspension was smeared on glass sections, and fixed by drying for 15 min
593 and by 4% PFA for 20 min. The slides were blocked in 10% normal goat serum and
594 0.5% triton X-100 for 1 hour. Testis sections and sperm slides were incubated overnight
595 at 4°C with primary antibodies (**Table S5**) in 5% normal goat serum and 0.25% triton X-

100. The slides were washed with PBS (3x10 min), and incubated with secondary Alexa Fluor 488 or 594 coupled goat anti-mouse or anti-rabbit IgG (1:300, Invitrogen, Waltham, MA, USA) for 1 hour, washed again with PBS (3x10 min), and counterstained with Sudan Black (0.1%) for autofluorescence quenching. Slides were mounted with fluoromount mounting medium containing DAPI (SouthernBiotech, Birmingham, AL, USA), and visualized using a Nikon Eclipse fluorescent microscope (Nikon Corporation).

In situ hybridization via RNAscope

RNAscope *in situ* hybridization was performed using BaseScope Detection Reagent Kit v2 RED (Advanced Cell Diagnostics Inc, Newark, CA, USA, Cat. No. 323900) according to the manufacturer's instructions. H₂O₂ treatment, antigen retrieval and protease treatment were performed on 5 µm-thick sections prior to hybridization with probes for Adgb (BA-Mm-Adgb-3zz-st, Advanced Cell Diagnostics, Cat. No. 862141), DapB as negative control (BA-DapB-3zz, Advances Cell Diagnostics, Cat. No. 701011) and Ppib as positive control (Ba-Mm-Ppib-3zz, Advanced Cell Diagnostics, Cat. No. 701071) at 40°C for 2 hours followed by eight amplification steps. The signal was revealed with Fast Red, the sections were counterstained with Gill's hematoxylin no. 1 and mounted with VectaMount (Vector Laboratories, Burlingame, CA, USA). The sections were visualized on a light microscope.

Electron microscopy

All electron microscopy experiments were performed at the Electron Microscopy Platform of the University of Lausanne, Switzerland.

620 Mouse testes were fixed in 2.5% glutaraldehyde solution (EMS, Hatfield, PA, USA) in
621 phosphate buffer (PB 0.1 M [pH 7.4]) for 1 hour at room temperature and postfixed in a
622 fresh mixture of osmium tetroxide 1% (EMS) with 1.5% of potassium ferrocyanide
623 (Sigma, St. Louis, MO, USA) in PB buffer for 1 hour at room temperature. The samples
624 were then washed twice in distilled water and dehydrated in acetone solution (Sigma) at
625 graded concentrations (30% - 40 min; 50% - 40 min; 70% - 40 min; 100% - 2x1 hour).
626 This was followed by infiltration in in Epon resin (EMS, Hatfield, PA, USA) at graded
627 concentrations (Epon 33% in acetone-4 hours; Epon 66% in acetone-4 hours; Epon
628 100%-2x8 hours) and finally polymerized for 48 hours at 60°C in an oven. Ultrathin
629 sections of 50 nm thick were cut using a Leica Ultracut (Leica Mikrosysteme GmbH,
630 Vienna, Austria), picked up on a copper slot grid 2x1 mm (EMS, Hatfield, PA, USA)
631 coated with a polystyrene film (Sigma, St Louis, MO, USA). Sections were post-stained
632 with uranyl acetate (Sigma, St Louis, MO, USA) 4% in H₂O for 10 min, rinsed several
633 times with H₂O followed by Reynolds lead citrate in H₂O (Sigma, St Louis, MO, USA)
634 for 10 min and rinsed several times with H₂O. Micrographs were taken with a
635 transmission electron microscope FEI CM100 (FEI, Eindhoven, The Netherlands) at an
636 acceleration voltage of 80kV with a TVIPS TemCamF416 digital camera (TVIPS
637 GmbH, Gauting, Germany).

638

639 *RNA-seq library preparation and transcriptome sequencing*

640 Total RNA from 2 independent samples of wildtype and Adgb^{-/-} testis was extracted
641 using the mirVana miRNA-Kit according to manufacturer's instructions (Life
642 Technologies, Carlsbad, US). Prior to library construction, RNA quality was assessed
643 using an Agilent 2100 Bioanalyzer and the Agilent RNA 6000 Nano Kit (Agilent
644 Technologies, Santa Clara, CA, USA). RNA was quantified using Qubit RNA BR Assay

645 Kit (Invitrogen, Waltham, MA, USA). Libraries were prepared starting from 1000 ng of
646 total RNA using the RNA Sample Prep Kit v2 (Illumina Inc, San Diego, CA, USA)
647 including a poly-A selection step following the manufacturer's instructions and
648 sequenced as 2 x 100 nt paired-end reads using an Illumina HiSeq 2500. Library
649 preparation and sequencing were performed by the NGS Core Facility of the Department
650 of Biology, Johannes-Gutenberg University (Mainz, Germany). RNA-Seq data are
651 available at the European Nucleotide Archive under accession number PRJEB46499.

652 *Differential gene expression, GO term annotation and pathway enrichment analyses*

653 Raw sequences were pre-processed to remove low quality reads and residual Illumina
654 adapter sequences using BBduk from the BBtools suite
655 (<https://sourceforge.net/projects/bbmap/>). The overall sequencing quality and the
656 absence of adapter contamination were evaluated with FastQC. Mapping was performed
657 with HISAT2 and quantification of gene expression was done using StringTie.
658 Differentially expressed genes were determined using DESeq2. Genes were considered
659 differentially expressed when presenting $|\text{fold change}| > 2$ and false discovery rate
660 (FDR)-corrected p-value ≤ 0.1 . GO term enrichment analyses were performed using
661 WebGestalt 2019 using the Overrepresentation Enrichment Analysis method, requiring a
662 BH-corrected p-value ≤ 0.05 and a minimum enrichment of 4 genes for term/pathway.
663 Enrichment in Canonical Pathways were performed with Qiagen's Ingenuity Pathway
664 Analysis (IPA, Qiagen, Hilden, Germany), Core analysis tool using bias-corrected z-
665 score (when applicable) and BH-corrected p-values ≤ 0.05 .

666

667

668

669

670 *Cloning and construction of expression plasmids*

671 Generation of pLenti6-ADGB was described before (Bracke, Hoogewijs, and Dewilde
672 2018b), pLenti6-SEPT10-V5 was purchased from DNASU (clone ID HsCD00943271,
673 DNASU Plasmid Repository, Arizona State University, AZ, USA). All additional
674 recombinant genes were cloned into pFLAG-CMVTM-6a expression vector (Sigma)
675 unless otherwise specified. All genes were amplified by PCR using Phusion High-
676 Fidelity DNA polymerase (Thermo Fisher Scientific). Recombinant *SEPT10* with N-
677 terminal FLAG tag and C-terminal myc tag was constructed by amplifying and ligating
678 SEPT10 coding sequence with in-primer designed myc tag (EQKLISEEDL) into the
679 expression vector, in-frame with the N-terminal FLAG tag. A glycine-serine (GSG)
680 linker was added between the last codon of *SEPT10* and the first codon of the myc tag.
681 Truncated ADGB proteins consisting of the calpain-like domain, 350-residue
682 uncharacterized domain and globin domain (N-terminal mutant), or the 700-residue
683 uncharacterized region (C-terminal mutant) domain were designed with GFP tags at both
684 N- and C-termini. The N-terminal mutant ADGB was amplified between codons of
685 residues 58 – 968 in ADGB while the C-terminal mutant ADGB was amplified between
686 codons of residues 969 – 1667. Amplicons were designed with 5'- and 3'- overhangs
687 compatible with two customized GFP amplicons designed to anneal at the 5'- and 3'-
688 ends of the genes. Glycine-serine linkers (GGSGGGGSGG) were added to bridge the
689 GFP tags and the truncated ADGB proteins. Similarly, N-terminally GFP-tagged isolated
690 ADGB globin was cloned by amplifying and ligating amplicons of the *ADGB* globin
691 gene downstream to a *GFP* gene with complementary overhangs, with the glycine-serine
692 linker added between the two proteins. With the same construction, GFP-tagged ADGB
693 globin domains with a single mutation on the proximal histidine in helix F codon 8
694 (H824G) or the distal glutamine in helix E codon 7 (Q792G), or both (H824G/Q792G)

were cloned by amplifying and ligating the globin gene using primers designed to carry the mutated codon sequence. ADGB Δ IQ and ADGB Δ CCD were constructed by amplifying designed *ADGB* amplicons with compatible overhangs and were ligated in-frame to generate a *ADGB* gene with deletions in the desired domains. For mammalian 2-hybrid assays, Gal4-CaM and VP16-globin domain were cloned into a pcDNA3.0 expression vector. Gal4 DNA-binding domain or VP16 transactivation domain sequences were amplified with the in-primer designed glycine-serine linker at the 3'-end of the amplicons. The coding sequence of *CALM3* and *ADGB* globin domain were amplified with complementary 5'-end and ligated to the Gal4 and VP16 sequences, respectively to generate the fusion genes.

Reporter gene assays

For mammalian 2-hybrid assays 2.15×10^5 HEK293 or 4×10^5 A375 cells were transiently transfected with 1 μ g firefly luciferase reporter plasmid (5xGAL4-TATA-luciferase, Addgene, 46756) (Sun et al. 1994) and 500 ng or 200 and 300 ng chimeric Gal4 and VP16 fusion protein vectors, respectively, in 12- or 6-well format using CaCl₂ or JetOptimus. To control for differences in transfection efficiency and extract preparation, 25 ng or 50 ng pRL-SV40 *Renilla* luciferase reporter vector (Promega, Madison, WI, USA) was co-transfected, respectively for HEK293 and A375 cells. Cultures were evenly split onto 12-well plates 24 hours after transfection for A375 cells. For hypoxia control experiments, 4×10^5 A375 cells were transiently co-transfected with 500 ng firefly luciferase *EPO* reporter plasmid (Storti et al. 2014) and 50 ng pRL-SV40 *Renilla* luciferase reporter vector. Luciferase activities of duplicate wells were determined using the Dual Luciferase Reporter Assay System (Promega) as described before (Schörg et al. 2015). Reporter activities were expressed as relative firefly/*Renilla*

720 luciferase activities. All reporter gene assays were performed at least 3 times
721 independently.

722

723 *Testosterone quantification*

724 Serum testosterone levels were determined as described previously with minor adaptations
725 (Strajhar et al. 2016). Briefly, for solid-phase extraction (SPE), each serum sample (100
726 μ L) was mixed with protein precipitation solution (100 μ L, 0.8 M zinc sulphate in
727 water/methanol; 50/50 v/v) containing 33 nM deuterium-labeled testosterone (D2) as
728 internal standard. Prior SPE, all samples were diluted to a final volume of 1 mL with water
729 and incubated in a thermoshaker (10 min at 4 °C, 1300 rpm). Following incubation,
730 samples were centrifuged for 10 min at 16000 g at 4°C, and supernatants (950 μ L) were
731 transferred to Oasis HBL SPE (1 cc) cartridges (Waters, Milford, MA, USA),
732 preconditioned with methanol and water (3 x 1 mL each). Samples were washed with water
733 (2 x 1 mL) and methanol/water (2 x 1 mL, 10/90 v/v). Testosterone was eluted with
734 methanol (2 x 0.75 mL), evaporated to dryness (3 hours, 35°C) and reconstituted in
735 methanol (25 μ L, 10 min, 4°C, 1300 rpm). Testosterone content was analyzed by ultra-
736 performance liquid chromatography-MS/MS (UPLC-MS/MS) using an Agilent 1290
737 Infinity II UPLC coupled to an Agilent 6495 triple quadrupole mass spectrometer
738 equipped with a jet-stream electrospray ionization interface (Agilent Technologies).
739 Analyte separation was achieved using a reverse-phase column (1.7 μ m, 2.1 mm x 150
740 mm; Acquity UPLC BEH C18; Waters). Data acquisition and quantitative analysis was
741 performed by MassHunter (Version B.10.0. Build 10.0.27, Agilent Technologies).

742

743

744

745 *Statistical analysis*

746 All values were presented as mean $\pm$ standard error of the mean (SEM). Differences in
 747 means between two groups were analyzed with unpaired 2-tailed Student's t-test (Fig.
 748 1A,C,D,E; Fig. 6H right graph; Fig. S4B,C) and those among multiple groups with one-
 749 way ANOVA followed by Tukey posthoc test (Fig. 2A; Fig. 6D,G,H left graph). All
 750 statistics were performed with GraphPad Prism software 7.05. Values of $p \leq 0.05$ were
 751 considered statistically significant.

752 **Acknowledgements**

753 We thank Christine Roulin for technical assistance. We thank Damien De Bellis from the
754 Electron Microscopy Platform of the University of Lausanne for EM section preparation
755 and image acquisition. This work was supported by the Swiss National Science
756 Foundation to DH (grant 31003A_173000) and the German Research Foundation to DH
757 (HO 5837/1-1) and TH (HA 2103/9-1).

758

759 **Competing interests**

760 The authors declare no competing interests.

761

762 **References**

- 763 Bähler, Martin, and Allen Rhoads. 2002. "Calmodulin signaling via the IQ motif." *FEBS*
764 *Letters* 513 (1):107-113. doi: 10.1016/s0014-5793(01)03239-2.
- 765 Bracke, A., D. Hoogewijs, and S. Dewilde. 2018b. "Exploring three different expression
766 systems for recombinant expression of globins: *Escherichia coli*, *Pichia pastoris*
767 and *Spodoptera frugiperda*." *Anal Biochem* 543:62-70. doi:
768 10.1016/j.ab.2017.11.027.
- 769 Bracke, An, Kris Peeters, Usha Punjabi, David Hoogewijs, and Sylvia Dewilde. 2018a.
770 "A search for molecular mechanisms underlying male idiopathic infertility."
771 *Reproductive BioMedicine Online* 36 (3):327-339. doi:
772 10.1016/j.rbmo.2017.12.005.
- 773 Brown, Jason M., and George B. Witman. 2014. "Cilia and Diseases." *BioScience* 64
774 (12):1126-1137. doi: 10.1093/biosci/biu174.
- 775 Burmester, T., and T. Hankeln. 2014. "Function and evolution of vertebrate globins."
776 *Acta Physiol (Oxf)* 211 (3):501-514. doi: 10.1111/apha.12312.

777 Burmester, Thorsten, and Thomas Hankeln. 2009. "What is the function of
778 neuroglobin?" *Journal of Experimental Biology* 212 (10):1423-1428. doi:
779 10.1242/jeb.000729.

780 Clermont, Y., and C. P. Leblond. 1955. "Spermiogenesis of man, monkey, ram and other
781 mammals as shown by the 'periodic acid-schiff' technique." *American Journal*
782 *of Anatomy* 96 (2):229-253. doi: 10.1002/aja.1000960203.

783 Darde, Thomas A., Estelle Lecluze, Aurélie Lardenois, Isabelle Stévant, Nathan Alary,
784 Frank Tüttelmann, Olivier Collin, Serge Nef, Bernard Jégou, Antoine D. Rolland,
785 Frédéric Chalmel, and Janet Kelso. 2019. "The ReproGenomics Viewer: a multi-
786 omics and cross-species resource compatible with single-cell studies for the
787 reproductive science community." *Bioinformatics* 35 (17):3133-3139. doi:
788 10.1093/bioinformatics/btz047.

789 Darde, Thomas A., Olivier Sallou, Emmanuelle Becker, Bertrand Evrard, Cyril
790 Monjeaud, Yvan Le Bras, Bernard Jégou, Olivier Collin, Antoine D. Rolland,
791 and Frédéric Chalmel. 2015. "The ReproGenomics Viewer: an integrative cross-
792 species toolbox for the reproductive science community." *Nucleic Acids*
793 *Research* 43 (W1):W109-W116. doi: 10.1093/nar/gkv345.

794 De Backer, Joey, Darko Maric, Matthias Bosman, Sylvia Dewilde, and David
795 Hoogewijs. 2021. "A reliable set of reference genes to normalize oxygen-
796 dependent cytoglobin gene expression levels in melanoma." *Scientific Reports*
797 11 (1). doi: 10.1038/s41598-021-90284-6.

798 Dolat, Lee, Qicong Hu, and Elias T. Spiliotis. 2014. "Septin functions in organ system
799 physiology and pathology." *Biological Chemistry* 395 (2):123-141. doi:
800 10.1515/hsz-2013-0233.

801 Fainberg, Jonathan, and James A. Kashanian. 2019. "Recent advances in understanding
802 and managing male infertility." *F1000Research* 8:670. doi:
803 10.12688/f1000research.17076.1.

804 Free, M. J., G. A. Schluntz, and R. A. Jaffe. 1976. "Respiratory Gas Tensions in Tissues
805 and Fluids of the Male Rat Reproductive Tract1." *Biology of Reproduction* 14
806 (4):481-488. doi: 10.1095/biolreprod14.4.481.

807 Ghossoub, Rania, Qicong Hu, Marion Failler, Marie-Christine Rouyez, Benjamin
808 Spitzbarth, Serge Mostowy, Uwe Wolfrum, Sophie Saunier, Pascale Cossart, W.
809 James Nelson, and Alexandre Benmerah. 2013. "Septins 2, 7, and 9 and MAP4
810 co-localize along the axoneme in the primary cilium and control ciliary length."
811 *Journal of Cell Science* 126:2583-2594. doi: 10.1242/jcs.111377.

812 Green, Christopher Daniel, Qianyi Ma, Gabriel L. Manske, Adrienne Niederriter Shami,
813 Xianing Zheng, Simone Marini, Lindsay Moritz, Caleb Sultan, Stephen J.
814 Gurczynski, Bethany B. Moore, Michelle D. Tallquist, Jun Z. Li, and Saher Sue
815 Hammoud. 2018. "A Comprehensive Roadmap of Murine Spermatogenesis
816 Defined by Single-Cell RNA-Seq." *Developmental Cell* 46 (5):651-667.e10. doi:
817 10.1016/j.devcel.2018.07.025.

818 Hermo, Louis, R. Marc Pelletier, Daniel G. Cyr, Charles E. Smith, and R. Marc Pelletier.
819 2010a. "Surfing the wave, cycle, life history, and genes/proteins expressed by
820 testicular germ cells. Part 1: Background to spermatogenesis, spermatogonia, and
821 spermatocytes." *Microscopy Research and Technique* 73 (4):241-278. doi:
822 10.1002/jemt.20783.

823 Hermo, Louis, R. Marc Pelletier, Daniel G. Cyr, Charles E. Smith, and R. Marc Pelletier.
824 2010b. "Surfing the wave, cycle, life history, and genes/proteins expressed by
825 testicular germ cells. Part 2: Changes in spermatid organelles associated with

- 826 development of spermatozoa." *Microscopy Research and Technique* 73 (4):279-
827 319. doi: 10.1002/jemt.20787.
- 828 Hermo, Louis, R. Marc Pelletier, Daniel G. Cyr, Charles E. Smith, and R. Marc Pelletier.
829 2010c. "Surfing the wave, cycle, life history, and genes/proteins expressed by
830 testicular germ cells. Part 3: Developmental changes in spermatid flagellum and
831 cytoplasmic droplet and interaction of sperm with the zona pellucida and egg
832 plasma membrane." *Microscopy Research and Technique* 73 (4):320-363. doi:
833 10.1002/jemt.20784.
- 834 Hermo, Louis, R. Marc Pelletier, Daniel G. Cyr, Charles E. Smith, and R. Marc Pelletier.
835 2010d. "Surfing the wave, cycle, life history, and genes/proteins expressed by
836 testicular germ cells. Part 4: Intercellular bridges, mitochondria, nuclear
837 envelope, apoptosis, ubiquitination, membrane/voltage-gated channels,
838 methylation/acetylation, and transcription factors." *Microscopy Research and*
839 *Technique* 73 (4):364-408. doi: 10.1002/jemt.20785.
- 840 Hermo, Louis, R. Marc Pelletier, Daniel G. Cyr, Charles E. Smith, and R. Marc Pelletier.
841 2010e. "Surfing the wave, cycle, life history, and genes/proteins expressed by
842 testicular germ cells. Part 5: Intercellular junctions and contacts between germs
843 cells and Sertoli cells and their regulatory interactions, testicular cholesterol, and
844 genes/proteins associated with more than one germ cell generation." *Microscopy*
845 *Research and Technique* 73 (4):409-494. doi: 10.1002/jemt.20786.
- 846 Hoogewijs, D., B. Ebner, F. Germani, F. G. Hoffmann, A. Fabrizius, L. Moens, T.
847 Burmester, S. Dewilde, J. F. Storz, S. N. Vinogradov, and T. Hankeln. 2012.
848 "Androglobin: a chimeric globin in metazoans that is preferentially expressed in
849 Mammalian testes." *Mol Biol Evol* 29 (4):1105-1114. doi:
850 10.1093/molbev/msr246.

851 Hu, Q., L. Milenkovic, H. Jin, M. P. Scott, M. V. Nachury, E. T. Spiliotis, and W. J.
852 Nelson. 2010. "A Septin Diffusion Barrier at the Base of the Primary Cilium
853 Maintains Ciliary Membrane Protein Distribution." *Science* 329 (5990):436-439.
854 doi: 10.1126/science.1191054.

855 Ihara, Masafumi, Ayae Kinoshita, Shuichi Yamada, Hiromitsu Tanaka, Ai Tanigaki,
856 Ayumi Kitano, Motohito Goto, Kazutoshi Okubo, Hiroyuki Nishiyama, Osamu
857 Ogawa, Chiaki Takahashi, Shigeyoshi Itohara, Yoshitake Nishimune, Makoto
858 Noda, and Makoto Kinoshita. 2005. "Cortical Organization by the Septin
859 Cytoskeleton Is Essential for Structural and Mechanical Integrity of Mammalian
860 Spermatozoa." *Developmental Cell* 8 (3):343-352. doi:
861 10.1016/j.devcel.2004.12.005.

862 Jégou, B., S. Sankararaman, A. D. Rolland, D. Reich, and F. Chalmel. 2017. "Meiotic
863 Genes Are Enriched in Regions of Reduced Archaic Ancestry." *Molecular
864 Biology and Evolution* 34 (8):1974-1980. doi: 10.1093/molbev/msx141.

865 Jeyaraj, D. Antony, Gail Grossman, and Peter Petrusz. 2003. *Reproductive Biology and
866 Endocrinology* 1 (1):48. doi: 10.1186/1477-7827-1-48.

867 Jordan, M., A. Schallhorn, and F. M. Wurm. 1996. "Transfecting Mammalian Cells:
868 Optimization of Critical Parameters Affecting Calcium-Phosphate Precipitate
869 Formation." *Nucleic Acids Research* 24 (4):596-601. doi: 10.1093/nar/24.4.596.

870 Keppner, Anna, Ditte Andreassen, Anne-Marie Mérillat, Julie Bapst, Camille Ansermet,
871 Qing Wang, Marc Maillard, Sumedha Malsure, Antoine Nobile, and Edith
872 Hummler. 2015. "Epithelial Sodium Channel-Mediated Sodium Transport Is Not
873 Dependent on the Membrane-Bound Serine Protease CAP2/Tmprss4." *Plos One*
874 10 (8):e0135224. doi: 10.1371/journal.pone.0135224.

- 875 Keppner, Anna, Darko Maric, Miguel Correia, Teng Wei Koay, Ilaria M. C. Orlando,
876 Serge N. Vinogradov, and David Hoogewijs. 2020. "Lessons from the post-
877 genomic era: Globin diversity beyond oxygen binding and transport." *Redox*
878 *Biology* 37:101687. doi: 10.1016/j.redox.2020.101687.
- 879 Keppner, Anna, Darko Maric, Chloé Sergi, Camille Ansermet, Damien De Bellis, Denise
880 V. Kratschmar, Jérémie Canonica, Petra Klusonova, Robert A. Fenton, Alex
881 Odermatt, Gilles Crambert, David Hoogewijs, and Edith Hummler. 2019.
882 "Deletion of the serine protease CAP2/Tmprss4 leads to dysregulated renal water
883 handling upon dietary potassium depletion." *Scientific Reports* 9 (1). doi:
884 10.1038/s41598-019-55995-x.
- 885 Khawar, Muhammad Babar, Hui Gao, and Wei Li. 2019. "Mechanism of Acrosome
886 Biogenesis in Mammals." *Frontiers in Cell and Developmental Biology* 7 (195).
887 doi: 10.3389/fcell.2019.00195.
- 888 Kim, Moshe S., Carol D. Froese, Hong Xie, and William S. Trimble. 2012. "Uncovering
889 Principles That Control Septin-Septin Interactions." *Journal of Biological*
890 *Chemistry* 287 (36):30406-30413. doi: 10.1074/jbc.M112.387464.
- 891 Kinoshita, M. 2003. "Assembly of Mammalian Septins." *Journal of Biochemistry* 134
892 (4):491-496. doi: 10.1093/jb/mvg182.
- 893 Kissel, Holger, Maria-Magdalena Georgescu, Sarit Larisch, Katia Manova, Gary R.
894 Hunnicutt, and Hermann Steller. 2005. "The Sept4 Septin Locus Is Required for
895 Sperm Terminal Differentiation in Mice." *Developmental Cell* 8 (3):353-364.
896 doi: 10.1016/j.devcel.2005.01.021.
- 897 Koay, Teng Wei, Carina Osterhof, Ilaria M. C. Orlando, Anna Keppner, Daniel Andre,
898 Schayan Yousefian, María Suárez Alonso, Miguel Correia, Robert Markworth,
899 Johannes Schödel, Thomas Hankeln, and David Hoogewijs. 2021. "Androglobin

900 gene expression patterns and FOXJ1-dependent regulation indicate its functional
 901 association with ciliogenesis." *Journal of Biological Chemistry* 296:100291. doi:
 902 10.1016/j.jbc.2021.100291.

903 Kotaja, Noora, Sarah Kimmins, Stefano Brancorsini, Didier Hentsch, Jean-Luc Vonesch,
 904 Irwin Davidson, Martti Parvinen, and Paolo Sassone-Corsi. 2004. "Preparation,
 905 isolation and characterization of stage-specific spermatogenic cells for cellular
 906 and molecular analysis." *Nature Methods* 1 (3):249-254. doi:
 907 10.1038/nmeth1204-249.

908 Krishnamurthy, Hanumanthappa, Natalia Danilovich, Carlos R. Morales, and M. Ram
 909 Sairam. 2000. "Qualitative and Quantitative Decline in Spermatogenesis of the
 910 Follicle-Stimulating Hormone Receptor Knockout (FORKO) Mouse1." *Biology*
 911 *of Reproduction* 62 (5):1146-1159. doi: 10.1095/biolreprod62.5.1146.

912 Kuo, Yung-Che, Ying-Hung Lin, Hau-Inh Chen, Ya-Yun Wang, Yu-Wei Chiou, Hsi-
 913 Hui Lin, Hsien-An Pan, Ching-Ming Wu, Shih-Ming Su, Chao-Chin Hsu, and
 914 Pao-Lin Kuo. 2012. "SEPT12 mutations cause male infertility with defective
 915 sperm annulus." *Human Mutation* 33 (4):710-719. doi: 10.1002/humu.22028.

916 Kuo, Yung-Che, Yi-Ru Shen, Hau-Inh Chen, Ying-Hung Lin, Ya-Yun Wang, Yet-Ran
 917 Chen, Chia-Yih Wang, and Pao-Lin Kuo. 2015. "SEPT12 orchestrates the
 918 formation of mammalian sperm annulus by organizing SEPT12-7-6-2/-4 core
 919 complexes." *Journal of Cell Science* 128:923-934. doi: 10.1242/jcs.158998.

920 Levine, Hagai, Niels Jørgensen, Anderson Martino-Andrade, Jaime Mendiola, Dan
 921 Weksler-Derri, Irina Mindlis, Rachel Pinotti, and Shanna H. Swan. 2017.
 922 "Temporal trends in sperm count: a systematic review and meta-regression
 923 analysis." *Human Reproduction Update* 23 (6):646-659. doi:
 924 10.1093/humupd/dmx022.

925 Lin, Ying-Hung, Yung-Ming Lin, Ya-Yun Wang, I. Shing Yu, Yi-Wen Lin, Yun-Han
926 Wang, Ching-Ming Wu, Hsien-An Pan, Shin-Chih Chao, Pauline H. Yen, Shu-
927 Wha Lin, and Pao-Lin Kuo. 2009. "The Expression Level of Septin12 Is Critical
928 for Spermiogenesis." *The American Journal of Pathology* 174 (5):1857-1868.
929 doi: 10.2353/ajpath.2009.080955.

930 Lin, Yu-Hua, Chia-Yen Huang, Chih-Chun Ke, Ya-Yun Wang, Tsung-Hsuan Lai,
931 Hsuan-Che Liu, Wei-Chi Ku, Chying-Chyuan Chan, and Ying-Hung Lin. 2020.
932 "ACTN4 Mediates SEPT14 Mutation-Induced Sperm Head Defects."
933 *Biomedicines* 8 (11):518. doi: 10.3390/biomedicines8110518.

934 Lukassen, S., E. Bosch, A. B. Ekici, and A. Winterpacht. 2018. "Characterization of
935 germ cell differentiation in the male mouse through single-cell RNA
936 sequencing." *Scientific Reports* 8 (1). doi: 10.1038/s41598-018-24725-0.

937 Mecklenburg, Jennifer M., and Brian P. Hermann. 2016. "Mechanisms Regulating
938 Spermatogonial Differentiation." *Results Probl Cell Differ* 58:253-287. doi:
939 10.1007/978-3-319-31973-5_10.

940 Neto, Filipe Tenorio Lira, Phil Vu Bach, Bobby B. Najari, Philip S. Li, and Marc
941 Goldstein. 2016. "Spermatogenesis in humans and its affecting factors."
942 *Seminars in Cell & Developmental Biology* 59:10-26. doi:
943 10.1016/j.semcdb.2016.04.009.

944 Peruquetti, Rita Luiza. 2015. "Perspectives on mammalian chromatoid body research."
945 *Animal Reproduction Science* 159:8-16. doi: 10.1016/j.anireprosci.2015.05.018.

946 Platts, Adrian E., David J. Dix, Hector E. Chemes, Kary E. Thompson, Robert Goodrich,
947 John C. Rockett, Vanesa Y. Rawe, Silvina Quintana, Michael P. Diamond,
948 Lillian F. Strader, and Stephen A. Krawetz. 2007. "Success and failure in human

949 spermatogenesis as revealed by teratozoospermic RNAs." *Human Molecular*
950 *Genetics* 16 (7):763-773. doi: 10.1093/hmg/ddm012.

951 Randi, E. B., B. Vervaet, M. Tsachaki, E. Porto, S. Vermeulen, M. T. Lindenmeyer, L.
952 T. T. Thuy, C. D. Cohen, O. Devuyst, A. D. Kistler, C. Szabo, N. Kawada, T.
953 Hankeln, A. Odermatt, S. Dewilde, R. H. Wenger, and D. Hoogewijs. 2020. "The
954 Antioxidative Role of Cytochrome in Podocytes: Implications for a Role in
955 Chronic Kidney Disease." *Antioxid Redox Signal* 32 (16):1155-1171. doi:
956 10.1089/ars.2019.7868.

957 Randriamboavonjy, Voahanginirina, Johann Isaak, Amro Elgheznavy, Frank Pistrosch,
958 Timo Frömel, Xiaoke Yin, Klaus Badenhoop, Heinrich Heide, Manuel Mayr, and
959 Ingrid Fleming. 2012. "Calpain inhibition stabilizes the platelet proteome and
960 reactivity in diabetes." *Blood* 120 (2):415-423. doi: 10.1182/blood-2011-12-
961 399980.

962 Rappsilber, J., M. Mann, and Y. Ishihama. 2007. "Protocol for micro-purification,
963 enrichment, pre-fractionation and storage of peptides for proteomics using
964 StageTips." *Nat Protoc* 2 (8):1896-906. doi: 10.1038/nprot.2007.261.

965 Reyes, Juan G., Jorge G. Farias, Sebastián Henríquez-Olavarrieta, Eva Madrid, Mario
966 Parraga, Andrea B. Zepeda, and Ricardo D. Moreno. 2012. "The Hypoxic
967 Testicle: Physiology and Pathophysiology." *Oxidative Medicine and Cellular*
968 *Longevity* 2012:1-15. doi: 10.1155/2012/929285.

969 Ribet, David, Serena Boscaini, Clothilde Cauvin, Martin Siguier, Serge Mostowy,
970 Arnaud Echard, and Pascale Cossart. 2017. "SUMOylation of human septins is
971 critical for septin filament bundling and cytokinesis." *Journal of Cell Biology*
972 216 (12):4041-4052. doi: 10.1083/jcb.201703096.

973 Schörg, Alexandra, Sara Santambrogio, James L. Platt, Johannes Schödel, Maja T.
974 Lindenmeyer, Clemens D. Cohen, Katrin Schrödter, David R. Mole, Roland H.
975 Wenger, and David Hoogewijs. 2015. "Destruction of a distal hypoxia response
976 element abolishes trans-activation of the PAG1 gene mediated by HIF-independent
977 chromatin looping." *Nucleic Acids Research* 43 (12):5810-5823. doi:
978 10.1093/nar/gkv506.

979 Sellin, Mikael E., Linda Sandblad, Sonja Stenmark, Martin Gullberg, and Douglas R.
980 Kellogg. 2011. "Deciphering the rules governing assembly order of mammalian
981 septin complexes." *Molecular Biology of the Cell* 22 (17):3152-3164. doi:
982 10.1091/mbc.e11-03-0253.

983 Shen, Y. R., H. Y. Wang, Y. C. Tsai, Y. C. Kuo, S. R. Wu, C. Y. Wang, and P. L. Kuo.
984 2020. "The SEPT12 complex is required for the establishment of a functional
985 sperm head-tail junction." *Molecular Human Reproduction* 26 (6):402-412. doi:
986 10.1093/molehr/gaaa031.

987 Shen, Yi-Ru, Han-Yu Wang, Yung-Che Kuo, Shih-Chuan Shih, Chun-Hua Hsu, Yet-
988 Ran Chen, Shang-Rung Wu, Chia-Yih Wang, and Pao-Lin Kuo. 2017. "SEPT12
989 phosphorylation results in loss of the septin ring/sperm annulus, defective sperm
990 motility and poor male fertility." *PLOS Genetics* 13 (3):e1006631. doi:
991 10.1371/journal.pgen.1006631.

992 Shevchenko, A., H. Tomas, J. Havlis, J. V. Olsen, and M. Mann. 2006. "In-gel digestion
993 for mass spectrometric characterization of proteins and proteomes." *Nat Protoc* 1
994 (6):2856-60. doi: 10.1038/nprot.2006.468.

995 Sigg, Monika Abedin, Tabea Menchen, Chanjae Lee, Jeffery Johnson, Melissa K.
996 Jungnickel, Semil P. Choksi, Galo Garcia, Henriette Busengdal, Gerard W.
997 Dougherty, Petra Pennekamp, Claudius Werner, Fabian Rentzsch, Harvey M.

998 Florman, Nevan Krogan, John B. Wallingford, Heymut Omran, and Jeremy F.
999 Reiter. 2017. "Evolutionary Proteomics Uncovers Ancient Associations of Cilia
1000 with Signaling Pathways." *Developmental Cell* 43 (6):744-762.e11. doi:
1001 10.1016/j.devcel.2017.11.014.

1002 Sirajuddin, M., M. Farkasovsky, E. Zent, and A. Wittinghofer. 2009. "GTP-induced
1003 conformational changes in septins and implications for function." *Proceedings of*
1004 *the National Academy of Sciences* 106 (39):16592-16597. doi:
1005 10.1073/pnas.0902858106.

1006 Sirajuddin, Minhajuddin, Marian Farkasovsky, Florian Hauer, Dorothee Kühlmann, Ian
1007 G. Macara, Michael Weyand, Holger Stark, and Alfred Wittinghofer. 2007.
1008 "Structural insight into filament formation by mammalian septins." *Nature* 449
1009 (7160):311-315. doi: 10.1038/nature06052.

1010 Skarnes, William C., Barry Rosen, Anthony P. West, Manousos Koutsourakis, Wendy
1011 Bushell, Vivek Iyer, Alejandro O. Mujica, Mark Thomas, Jennifer Harrow, Tony
1012 Cox, David Jackson, Jessica Severin, Patrick Biggs, Jun Fu, Michael Nefedov,
1013 Pieter J. de Jong, A. Francis Stewart, and Allan Bradley. 2011. "A conditional
1014 knockout resource for the genome-wide study of mouse gene function." *Nature*
1015 474 (7351):337-342. doi: 10.1038/nature10163.

1016 Spiliotis, Elias T., and Konstantinos Nakos. 2021. "Cellular functions of actin- and
1017 microtubule-associated septins." *Current Biology* 31 (10):R651-R666. doi:
1018 10.1016/j.cub.2021.03.064.

1019 Storti, F., S. Santambrogio, L. M. Crowther, T. Otto, I. Abreu-Rodriguez, M. Kaufmann,
1020 C. J. Hu, C. Dame, J. Fandrey, R. H. Wenger, and D. Hoogewijs. 2014. "A novel
1021 distal upstream hypoxia response element regulating oxygen-dependent

1022 erythropoietin gene expression." *Haematologica* 99 (4):e45-e48. doi:

1023 10.3324/haematol.2013.102707.

1024 Strajhar, P., Y. Schmid, E. Liakoni, P. C. Dolder, K. M. Rentsch, D. V. Kratschmar, A.

1025 Odermatt, and M. E. Liechti. 2016. "Acute Effects of Lysergic Acid

1026 Diethylamide on Circulating Steroid Levels in Healthy Subjects." *J*

1027 *Neuroendocrinol* 28 (3):12374. doi: 10.1111/jne.12374.

1028 Sun, P., H. Enslen, P. S. Myung, and R. A. Maurer. 1994. "Differential activation of

1029 CREB by Ca²⁺/calmodulin-dependent protein kinases type II and type IV

1030 involves phosphorylation of a site that negatively regulates activity." *Genes Dev*

1031 8 (21):2527-39. doi: 10.1101/gad.8.21.2527.

1032 Toure, Aminata, Baptiste Rode, Gary R. Hunnicutt, Denise Escalier, and Gérard Gacon.

1033 2011. "Septins at the annulus of mammalian sperm." *Biological Chemistry* 392

1034 (8-9):799-803. doi: 10.1515/bc.2011.074.

1035 Tyanova, S., T. Temu, and J. Cox. 2016. "The MaxQuant computational platform for

1036 mass spectrometry-based shotgun proteomics." *Nat Protoc* 11 (12):2301-2319.

1037 doi: 10.1038/nprot.2016.136.

1038 Villalobo, Antonio, María González-Muñoz, and Martin W. Berchtold. 2019. "Proteins

1039 with calmodulin-like domains: structures and functional roles." *Cellular and*

1040 *Molecular Life Sciences* 76 (12):2299-2328. doi: 10.1007/s00018-019-03062-z.

1041 Villalobo, Antonio, Hiroaki Ishida, Hans J. Vogel, and Martin W. Berchtold. 2018.

1042 "Calmodulin as a protein linker and a regulator of adaptor/scaffold proteins."

1043 *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1865 (3):507-

1044 521. doi: 10.1016/j.bbamcr.2017.12.004.

1045 Wang, Mei, Xixi Liu, Gang Chang, Yidong Chen, Geng An, Liying Yan, Shuai Gao,

1046 Yanwen Xu, Yueli Cui, Ji Dong, Yuhan Chen, Xiaoying Fan, Yuqiong Hu, Ke

1047 Song, Xiaohui Zhu, Yun Gao, Zhaokai Yao, Shuhui Bian, Yu Hou, Jiahao Lu,
1048 Rui Wang, Yong Fan, Ying Lian, Wenhao Tang, Yapeng Wang, Jianqiao Liu,
1049 Lianming Zhao, Luyu Wang, Zhaoting Liu, Renpei Yuan, Yujia Shi, Boqiang
1050 Hu, Xiulian Ren, Fuchou Tang, Xiao-Yang Zhao, and Jie Qiao. 2018. "Single-
1051 Cell RNA Sequencing Analysis Reveals Sequential Cell Fate Transition during
1052 Human Spermatogenesis." *Cell Stem Cell* 23 (4):599-614. doi:
1053 10.1016/j.stem.2018.08.007.

1054 Wang, Ya-Yun, Tsung-Hsuan Lai, Mei-Feng Chen, Hui-Ling Lee, Pao-Lin Kuo, and
1055 Ying-Hung Lin. 2019. "SEPT14 Mutations and Teratozoospermia: Genetic
1056 Effects on Sperm Head Morphology and DNA Integrity." *Journal of Clinical*
1057 *Medicine* 8 (9):1297. doi: 10.3390/jcm8091297.

1058

Fig.1




Figure 1. Validation of the knockout model and testicular phenotype.

(A) Relative mRNA expression levels of Adgb in testes of wildtype (+/+) and knockout mice (-/-) (n=6 per genotype; p=0.000008). (B) Representative immunoblot for Adgb in testis lysates from wildtype (+/+) and knockout mice (-/-) (n=6 per genotype) and (C) corresponding protein quantification. Tubulin was used as loading control. p=0.0007. (D) Testis weight (g) in Adgb wildtype (+/+), heterozygous (+/-) and knockout (-/-) mice (n=8-13 per genotype). p=0.000003. (E) Serum testosterone levels (nM) in Adgb wildtype (+/+) and knockout (-/-) mice (n=8 per genotype). p=0.041. (F) Representative hematoxylin and eosin (H&E) stained sections of testes from Adgb wildtype (+/+) and knockout mice (-/-) at the different stages of spermatogenesis. Heads (H), tails (T), residual bodies (R) cytoplasmic bulges (C), meiosis (M), elongating spermatids (ES), round spermatids (RS), stage 16 spermatids (S16) and abnormal heads (A) are indicated. Scale bar represents 50 μ m. (G) Representative H&E stained sections of epididymides from Adgb wildtype (+/+) and knockout mice (-/-). Heads (H), tails (T), cytoplasmic bodies (C),

residual bodies (R), germ cells (G), abnormal heads (A) are shown. Note the empty lumen (EL) in knockout mice. Scale bar represents 50 μm . ** $p < 0.01$, *** $p < 0.001$.

Fig.2




Figure 2. Testicular Adgb expression pattern and localization.

(A) Relative mRNA expression levels of Adgb in testes of wildtype mice during embryonic development (E) and early post-natal (P) life (n=3-4 per condition). (B) Representative immunoblot for Adgb in testis lysates from wildtype mice at different post-natal (P) ages (n=3 per condition) and (C) corresponding protein quantification. Tubulin was used as loading control. (D) Flow cytometric analysis of spermatogenic cell populations following propidium iodide staining in Adgb wildtype (+/+, white circles, n=4), heterozygous (+/-, grey circles, n=5) and knockout (-/-, black

circles, n=3) testes. H: elongating and elongated spermatids; 1C: round spermatids; 2C: spermatogonia, secondary spermatocytes, testicular somatic cells; S: spermatogonia synthesizing DNA; 4C: primary spermatocytes. $p=0.00024$. **(E)** Representative pictures of Adgb protein (red fluorescence) detection in testes of wildtype (+/+) and knockout (-/-) animals. Left panels 20x magnification, right panels 40x magnification, scale bars represent 100 μm and 50 μm , respectively. Nuclei were stained with DAPI. **(F)** Representative pictures of Adgb mRNA *in situ* hybridization in testes from wildtype (+/+) and knockout (-/-) animals. Left panels 20x magnification; right panels, 40x magnification; scale bars represent 100 μm and 50 μm , respectively. Positive (ctrl +, PPIB) and negative (ctrl -, DapB) control sections are shown. **(G)** Representative pictures of b-galactosidase activity (X-gal staining) in testes from Tm1b wildtype (+/+) and Tm1b heterozygous (tg/+) mice, and isolated spermatozoa from Tm1b heterozygous (tg/+) mice. Left panels, 20x magnification; middle panels, 40x magnification; right panel, 60x magnification; scale bars represent 100 μm , 50 μm and 20 μm , respectively. Spermatozoa were counterstained with nuclear fast red. **(H)** Representative picture of Adgb protein (green fluorescence) in a single spermatozoon from wildtype (+/+) mice (left panel), and negative control (secondary antibody only, right panel). Scale bar represents 20 μm , nuclei were stained with DAPI. ** $p < 0.001$.

Fig.3




Figure 3. Defective spermatogenesis is associated with flagellar malformation in Adgb knockout mice.

(A) Representative pictures of cauda epididymis sperm from wildtype (+/+) and Adgb knockout animals (-/-). Scale bar represents 20 μ m. (B) Representative pictures of PNA-stained cauda epididymis sperm from wildtype (+/+) and Adgb knockout animals (-/-). Nuclei were stained with DAPI. Scale bar represents 20 μ m. (C) Representative TEM pictures from wildtype (+/+, left panels) and knockout (-/-, right panels) testes. Misshaped sperm heads (hash), axonemes (arrows),

mitochondria (arrowheads) and microtubules (asterisks), nuclear inclusions (ampersand), and abnormal manchette elongation (dollar) are shown. Scale bar lengths are indicated on each picture.

Fig.4




Figure 4. Adgb and Sept10 interact *in vivo* and *in vitro*.

(A) Proteins of the septin family are specifically enriched in the Adgb immuno-precipitation (IP). The iBAQ (intensity-based absolute quantification) values of each Adgb IP (triplicate) and IgG control IP (duplicate) were log2 transformed and normalized against the median value. Missing values were imputed before the mean values of the Adgb and IgG control IPs were calculated. The normalized abundance of each protein detected in the Adgb IP ($\log_2$ Adgb (mean)) is plotted against its specific enrichment compared to the IgG control IP ($\log_2$ (Adgb

(mean) / IgG (mean)). Adgb and septins are highlighted as blue and red dots, respectively, in the christmas tree plot representation. **(B)** Representative immunoblot of Adgb and Sept10 in testis lysates from wildtype (+/+) and knockout (-/-) mice following co-immunoprecipitation of Adgb and Sept10. **(C-E)** Representative immunoblots of ADGB and Sept10 in protein lysates of HEK293 cells (co-)transfected with full-length ADGB **(C)**, N-ter ADGB **(D)** and C-ter ADGB **(E)** and Sept10 following co-immunoprecipitation of ADGB and Sept10.

Fig.5




Figure 5. Adgb and Sept10 co-localize in developing acrosome, sperm neck and annulus.

(A) Representative pictures of Adgb protein (green fluorescence) and Sept10 (red fluorescence) in elongating spermatids (stage 7 [S7] upper panels, stage 12 [S12] middle panels, and stage 15 [S15] lower panels) after stage-specific tubule dissection of wildtype testes. Nuclei were stained with DAPI. Negative control (secondary antibodies only) is shown on the right. Scale bar represents 10 μ m. Sperm neck (arrowhead) and annulus (arrow) are highlighted. (B) Representative pictures of Sept10 (left panels, green fluorescence) and CoxIV (middle panels, red fluorescence) proteins in wildtype (+/+) and knockout (-/-) epididymal sperm. Sections were counterstained with DAPI. Negative control (secondary antibodies only) are shown (right panels). Scale bar represents 10 μ m.

Fig.6




Figure 6. Adgb contributes to *in vitro* CaM-dependent Sept10 cleavage.

(A) Representative immunoblots of ADGB and SEPT10 in protein lysates of HEK293 cells co-transfected with plasmids encoding SEPT10 and two dose-dependent amounts of full-length ADGB. **(B)** Representative immunoblot of ADGB and V5 in protein lysates of HEK293 cells

(co-)transfected with full-length ADGB and a C-terminally V5-tagged SEPT10 construct following co-immunoprecipitation of ADGB and V5-SEPT10. **(C)** Representative immunoblots of ADGB, V5, HIF-1 α and PHD2 in protein lysates of HEK293 cells (co-)transfected with full-length ADGB and V5-SEPT10 following exposure to normoxic (20% O₂) and hypoxic conditions (0.2% O₂) for 24 hours. HIF-1 α and PHD2 were used as positive controls for hypoxia. **(D)** Representative immunoblot of ADGB and V5 in protein lysates of HEK293 cells (co-)transfected with full-length ADGB, V5-SEPT10 and an ADGB-IQ deletion mutant, and corresponding protein quantification of cleaved/full-size SEPT10 ratio (n=3-6 independent experiments). Ponceau S protein staining was used as loading control. **(E)** Representative immunoblot of GFP and V5 in protein lysates of HEK293 cells (co-)transfected with a truncated construct of the globin domain of ADGB (spanning the IQ domain) (GFP-Globin) and a V5-tagged CaM (V5-Calmodulin) following immunoprecipitation of GFP. **(F)** Representative immunoblot of GFP and V5 in protein lysates of HEK293 cells (co-)transfected with GFP-Globin, a GFP-Globin construct with mutation of the proximal heme-binding histidine (GFP-Globin-F8(Gly)), a GFP-Globin construct with mutation of the distal glutamine (GFP-Globin-E7(Gly)) and V5-CaM following immunoprecipitation of GFP. **(G, H)** Mammalian-2 hybrid assays in HEK293 cells under normoxic conditions **(G)** and A375 cells under normoxic and hypoxic (0.2% O₂) conditions **(H)** (n=3-5 independent experiments). HEK293 and A375 cells were transiently transfected with fusion protein vectors based on a Gal4 DNA binding domain fused to calmodulin (Gal4-CaM) and a VP16 activation domain fused to the ADGB globin domain comprising the IQ domain (VP16-Globin), a Gal4 response element-driven firefly luciferase reporter, and a *Renilla* luciferase control vector. Increasing transfection amounts for the Gal4-CaM fusion protein were employed. Following transfection, A375 cells were incubated under normoxic (20% O₂) or hypoxic (0.2% O₂) conditions, and luciferase reporter gene activities were determined 24 hours later. Single construct transfections served as negative controls. The single Gal4-CaM control condition is not displayed in A375 cells exposed to hypoxia, due to its hypoxic regulation. An Epo hypoxia response element-driven firefly luciferase construct served as hypoxic control. * p< 0.05, ** p< 0.01, *** p< 0.001.

Supplementary Information for

Androglobin, a chimeric mammalian globin, is required for male fertility.

Anna Keppner¹, Miguel Correia¹, Sara Santambrogio², Teng Wei Koay¹, Darko Maric¹,
Carina Osterhof³, Denise V Winter⁴, Angèle Clerc¹, Michael Stumpe⁵, Frédéric Chalmel⁶,
Sylvia Dewilde⁷, Alex Odermatt⁴, Dieter Kressler⁵, Thomas Hankeln³, Roland H.
Wenger², David Hoogewijs^{1*}

¹ Section of Medicine, Department of Endocrinology, Metabolism and Cardiovascular system (EMC), University of Fribourg, Switzerland; ² Institute of Physiology, University of Zurich, Switzerland; ³ Institute for Organismic and Molecular Evolutionary Biology, University of Mainz, Germany; ⁴ Department of Pharmaceutical Sciences, University of Basel, Switzerland, ⁵ Department of Biology, University of Fribourg, Switzerland, ⁶ University of Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, France, ⁷ Department of Biomedical Sciences, University of Antwerp, Belgium

*Corresponding author: David Hoogewijs
Email: david.hoogewijs@unifr.ch

This section includes:

Figures S1 to S10
Tables S1 to S2
Legends for datasets S1 to S3
SI References
Source data files are provided for Figures 1, 2, 4B, 4C, 4D, 4E, 6A, 6B, 6C, 6D, 6E, 6F, S1, S5A, 5B, S5C, S5D, S5E, S5F, S5G, S6, S7, S8, S10

Other supplementary materials for this manuscript include the following:

Datasets S1 to S3

Fig.S1




Figure S1. Generation of *Adgb* knockout mice. (A) Schematic representation of the wild-type, Tm1a (gene-trap), Tm1b (following breeding with Cre-deleter mice), Tm1c (following breeding with FLP-deleter mice) and Tm1d (knockout) (following breeding of Tm1c mice with Cre-deleter mice) alleles. The position of the probe (asterisk), the primers (5'S, 5'AS, 3'S and 3'AS) and restriction sites for the ES cell screening by PCR and Southern blot are shown. FRT sites flank the gene-trap construct, containing the lacZ gene and the neomycin resistance cassette (neo), whereas loxP sites flank the neo-cassette and exons 13 and 14 of the *Adgb* gene. Position of the PCR primers for the genotyping of the different mouse lines are shown (F1 and R1-4). (B) Representative PCR-based analysis of targeted ES cells using primers 5'S and 5'AS (left panel) (M: marker, 1 and 2: positive and negative clones respectively), and primers 3'S and 3'AS (right panel) (M: marker, 1 and 2: negative and positive clones respectively). (C) Southern blot analysis of targeted ES cell clones using the exon 13 probe (asterisk) and following digestion with *PvuII* and *PstI* (M: marker, 1-5: different clones tested positive for both 5' and 3' PCR reactions). (D) Genotyping of Tm1a wild-type (+/+), positive for both 5' and 3' PCR reactions.

heterozygous (tg/+) and homozygous (tg/tg) mice (upper panel), M: marker, Tm1b wild-type (+/+), heterozygous (tg/+) and homozygous (tg/tg) mice (middle panel), and Tm1d wild-type (+/+), heterozygous (+/-) and homozygous (-/-, knockout) mice (lower panel).

Fig.S2




Figure S2. Adgb KO mice do not display changes in NOS, SOD and apoptotic gene expression. Relative mRNA expression levels of (A) NOS1, (B) NOS2, (C) NOS3, (D) SOD1, (E) SOD2, (F) SOD3, and (G) ratio of Bax and Bcl2 in testis lysates from Adgb wildtype (+/+, n=6) and knockout (-/-, n=5) mice.

Fig.S3




Figure S3. Temporal Adgb and Sept10 expression profiles based on single-cell and bulk RNA-seq datasets. (A, C, E, G): Adgb; (B, D, F, H): Sept10. Data were obtained from Lukassen et al. 2018 (mouse, panels A and B), Green et al. 2018 (mouse, panels C and D), Jégou et al. 2017 (human, panels E and F), and Wang et al.

2018 (human, panels G and H), respectively (Green et al. 2018, Jégou et al. 2017, Lukassen et al. 2018, Wang et al. 2018).

Fig.S4




Figure S4. Volcano plot of differentially expressed genes in *Adgb* knockout mice testis samples and validation by RT-qPCR. (A) X-axis represents the log2 fold change and y-axis represents log10 of the adjusted P-value. Genes are assigned with specific colors after DESeq2 analysis: gray (not significant [NS]), green $|\text{Log2FC}|>1$, blue (adjusted $P<0.05$), or red ($|\text{Log2FC}|>1$ and adjusted $P<0.05$). *Adgb* is indicated. (B) Relative mRNA expression levels of *17βhds3* in testes of wildtype (+/+) and knockout mice (-/-) (n=3-4 per genotype) on post-natal day 24. (C) Relative mRNA expression levels of *Lhcgr* in testes of wildtype (+/+) and knockout mice (-/-) (n=3-4 per genotype) on post-natal day 24. * $p<0.05$.

Fig.S5




Figure S5. The protein expression levels of Sept2, 7, 8, 9, 10, 11 and 14 are unaffected in Adgb KO testis. Representative immunoblots of testis lysates from

Adgb wildtype (+/+, n=6-12) and knockout (-/-, n=6-11) mice for (A) Sept2, (B) Sept7, (C) Sept8, (D) Sept9, (E) Sept10, (F) Sept11, and (G) Sept14, and the corresponding protein quantifications for (H) Sept2, (I) Sept7, (J) Sept8, (K) Sept9, (L) Sept10, (M) Sept11, and (N) Sept14. Ponceau S protein staining was used as loading control.

Fig.S6




Figure S6. The interaction between ADGB and SEPT10 is maintained despite mutation of the coiled-coil domains. Representative immunoblots of ADGB and Sept10 in protein lysates of HEK293 cells (co-)transfected with CCD mutant ADGB and CCD mutant SEPT10 following co-immunoprecipitation of ADGB and SEPT10.

Fig.S7




Figure S7. Sept10 temporal expression profile on mRNA and protein levels. (A) Relative mRNA expression levels of Sept10 in testes of wildtype mice during the first wave of spermatogenesis at indicated post-natal (P) days. **(B)** Representative immunoblot for Sept10 in testis lysates from wildtype mice at indicated post-natal (P) ages (n=2 per condition) and **(C)** corresponding protein quantification. Ponceau S protein staining was used as loading control.

Fig.S8




Figure S8. Reciprocal co-immunoprecipitation (coIP) of ADGB and V5-SEPT10 from Fig 7B. Representative immunoblot of ADGB and V5 in protein lysates of HEK293 cells (co-)transfected with full-length ADGB and a C-terminally V5-tagged SEPT10 construct following co-immunoprecipitation of ADGB and V5-SEPT10.

Fig.S9




Figure S9. ADGB globin immunoprecipitation (IP) vs GFP control IP.

Calmodulin (CALM1) is among the proteins that are prominently enriched in the anti-GFP IP of cells expressing the GFP-tagged globin domain of ADGB. The iBAQ values of the two IPs were log2 transformed and normalized against the median value. The normalized abundance of each protein detected in the globin IP (log2 globin) of cells expressing the GFP-globin construct is plotted against its specific enrichment compared to the control IP (log2 (globin / control)) from cells expressing the GFP-linker-GFP construct. ADGB and CALM1 are highlighted as blue and red dots, respectively, in the christmas tree plot representation.

Fig.S10




Figure S10. Double mutation of the key heme-binding residues does not abrogate interaction of ADGB and CaM. Representative immunoblot of GFP and V5 in protein lysates of HEK293 cells (co-)transfected with GFP-Globin, a GFP-Globin construct with a double mutation of the proximal heme-binding histidine and the distal heme-binding glutamine (GFP-Globin-F8/E7(Gly)) and V5-CaM following immunoprecipitation of GFP.

Table S1. List of mouse primers used for RT-qPCR

Gene	Forward primer 5'-3'	Reverse primer 5'-3'
Adgb	TTCCAACAGAAACACATTTTGTCCA	TCCATTACTATGCTCATTTCCCCT
NOS1	CAGGCAAATCCCAAGCCTATG	CAAAGCACAGCCGAATTTCTC
NOS2	ATTCACAGCTCATCCGGTACG	GGATCTTGACCATCAGCTTGC
NOS3	CATTTTCGGACTCACATTGCG	TTGGTCAACCGAACGAAGTG
SOD1	GTGATTGGGATTGCGCAGTA	TGGTTGAGGGTAGCAGATGAGT
SOD2	TTAACGCGCAGATCATGCA	GGTGGCGTTGAGATTGTTCA
SOD3	CATGCAATCTGCAGGGTACAA	AGAACCAAGCCGGTGATCTG
Bax	GCGTGGTTGCCCTCTTCTACTTTG	AGTCCAGTGTCCAGCCCATGATG
Bcl2	AAGGGCTTCACACCCAAATCT	CTTCTACGTCTGCTTGGCTTTGA
Sept10	GGCCTCATGCGACGAGATAA	CCAATTCCAGTCTCCCCAC
17βhsd3	ATT TTA CCA GAG AAG ACA TCT	GGG GTC AGC ACC TGA ATA ATG
Lhcgr	CGA CGC TAA TCT CGC TGG AG	CGT AAT CCC AGC CAC TGA GTT
Actin	GAGCGTGGCTACAGCTTCAC	GGCATAGAGGTCTTTACGGAT G

Table S2. List of antibodies used throughout the study.

Antibody	Reference	Species	Application/dilution
anti-mAdgb	Proteintech, custom made against region 409-745 of mAdgb	Rabbit	Immunoblot 1/200 IF 1/300
anti-hAdgb N-ter	Atlas Antibodies, HPA036340	Rabbit	Immunoblot 1/500
anti-hAdgb C-ter	OriGene, TA330746	Rabbit	Immunoblot 1/500
anti-Sept10	Proteintech, 12420-1-AP	Rabbit	Immunoblot 1/500 IF 1/300
anti-Sept2	Proteintech, 60075-1-Ig	Mouse	Immunoblot 1/500
anti-Sept7	Proteintech, 13818-1-AP	Rabbit	Immunoblot 1/500
anti-Sept8	Proteintech, 11769-1-AP	Rabbit	Immunoblot 1/500
anti-Sept9	Proteintech, 10769-1-AP	Rabbit	Immunoblot 1/500
anti-Sept11	Proteintech, 14672-1-AP	Rabbit	Immunoblot 1/500
anti-Sept14	Proteintech, 24590-1-AP	Rabbit	Immunoblot 1/500
anti-GFP	Proteintech, 50430-2-AP	Rabbit	Immunoblot 1/1000
anti-V5	Invitrogen, 46-0705	Mouse	Immunoblot 1/1000
anti-CoxIV	Abcam, ab14744	Rabbit	IF 1/300
anti-HIF-1 α	BD Transduction laboratories, 610958	Mouse	Immunoblot 1/500
anti-PHD2	Novus Biologicals, NB100-137	Rabbit	Immunoblot 1/1000
anti- α -tubulin	Santa Cruz, TU-02	Mouse	Immunoblot 1/1000

Datasets

Dataset S1. Differentially regulated genes in wild-type versus Adgb knockout testes. Genes with a significant >2-fold induction or reduction are displayed.

Dataset S2. Raw MS data of the Adgb IP vs IgG control IP.

Dataset S3. Raw MS data of the ADGB globin IP vs GFP control IP.

SI references

- Green, Christopher Daniel, Qianyi Ma, Gabriel L. Manske, Adrienne Niederriter Shami, Xianing Zheng, Simone Marini, Lindsay Moritz, Caleb Sultan, Stephen J. Gurczynski, Bethany B. Moore, Michelle D. Tallquist, Jun Z. Li, and Saher Sue Hammoud. 2018. "A Comprehensive Roadmap of Murine Spermatogenesis Defined by Single-Cell RNA-Seq." *Developmental Cell* 46 (5):651-667.e10. doi: 10.1016/j.devcel.2018.07.025.
- Jégou, B., S. Sankararaman, A. D. Rolland, D. Reich, and F. Chalmel. 2017. "Meiotic Genes Are Enriched in Regions of Reduced Archaic Ancestry." *Molecular Biology and Evolution* 34 (8):1974-1980. doi: 10.1093/molbev/msx141.
- Lukassen, S., E. Bosch, A. B. Ekici, and A. Winterpacht. 2018. "Characterization of germ cell differentiation in the male mouse through single-cell RNA sequencing." *Scientific Reports* 8 (1). doi: 10.1038/s41598-018-24725-0.
- Wang, Mei, Xixi Liu, Gang Chang, Yidong Chen, Geng An, Liying Yan, Shuai Gao, Yanwen Xu, Yueli Cui, Ji Dong, Yuhao Chen, Xiaoying Fan, Yuqiong Hu, Ke Song, Xiaohui Zhu, Yun Gao, Zhaokai Yao, Shuhui Bian, Yu Hou, Jiahao Lu, Rui Wang, Yong Fan, Ying Lian, Wenhao Tang, Yapeng Wang, Jianqiao Liu, Lianming Zhao, Luyu Wang, Zhaoting Liu, Renpei Yuan, Yujia Shi, Boqiang Hu, Xiulian Ren, Fuchou Tang, Xiao-Yang Zhao, and Jie Qiao. 2018. "Single-Cell RNA Sequencing Analysis Reveals Sequential Cell Fate Transition during Human Spermatogenesis." *Cell Stem Cell* 23 (4):599-614. doi: 10.1016/j.stem.2018.08.007.

Source data files

Source data files Figure 1. Full uncropped immunoblots for Adgb and tubulin.

Source data files Figure 2. Full uncropped immunoblots for Adgb and tubulin.

Source data files Figure 4B and 4C. Full uncropped immunoblots for Adgb and Sept10.

Source data files Figure 4D. Full uncropped immunoblots for ADGB and SEPT10.

Source data files Figure 4E. Full uncropped immunoblots for ADGB and SEPT10.

Source data files Figure 6A and 6B. Full uncropped immunoblots for ADGB, V5 and SEPT10.

Source data files Figure 6C and 6D. Full uncropped immunoblots for ADGB, PHD2, HIF-1 α and V5.

Source data files Figure 6E. Full uncropped immunoblots for GFP and V5.

Source data files Figure 6F. Full uncropped immunoblots for GFP and V5.

Source data files Figure S1. Full uncropped DNA gels and southern blot.

Source data files Figure S5A, B, C. Full uncropped immunoblots for Sept2, Sept7, Sept8 and Ponceau S staining.

Source data files Figure S5D, E, F. Full uncropped immunoblots for Sept9, Sept10, Sept11 and Ponceau S staining.

Source data files Figure S5G. Full uncropped immunoblots for Sept14 and Ponceau S staining.

Source data files Figure S6. Full uncropped immunoblots for ADGB and SEPT10.


Source data files Figure S7B. Full uncropped immunoblots for Sept10 and Ponceau S staining.

Source data files Figure S8. Full uncropped immunoblots for ADGB and V5.


Source data files Figure S10. Full uncropped immunoblots for GFP and V5.

Source files Fig. 1B

Adgb




Tubulin




Source files Fig. 2B

Adgb




Tubulin



Source file for Fig. 4B

Adgb



Sept10





Fig. 4C

IP Adgb IB Adgb



IP Adgb IB Sept10






Fig. 4C

IP Sept10 IB Adgb




IP Sept10 IB Sept10



Source files for Fig. 4D

IP ADGB IB ADGB



IP ADGB IB SEPT10






Fig. 4D

IP SEPT10 IB ADGB




IP SEPT10 IB SEPT10



Source files for Fig. 4E

IP ADGB IB ADGB



IP ADGB IB SEPT10






Fig. 4E

IP SEPT10 IB ADGB




IP SEPT10 IB SEPT10



Source files for Fig. 6A

ADGB



Sept10






Fig. 6B

IP ADGB IB ADGB



IP ADGB IB V5



Source files for Fig. 6C






Fig. 6D




Source files for Fig. 6E


GFP




V5




GFP



V5



Source files for Fig. 6F



Source files for Fig. S1

Fig. S1B




Fig. S1C

Southern blot





Fig. S1D



Source files for Fig. S5 A

Sept2



Ponceau





Fig. S5 B

Sept7



Ponceau






Fig. S5 C

Sept8




Ponceau



Source files for Fig. S5 D

Sept9



Ponceau





Fig. S5 E

Sept10



Ponceau






Fig. S5 F

Sept11




Ponceau




Source file for Fig. S5 G

Sept14




Ponceau



Source files for Fig. S6

IP ADGB IB ADGB



IP ADGB IB SEPT10






Fig. S6

IP SEPT10 IB ADGB




IP SEPT10 IB SEPT10




Source file for Fig. S7B

Sept10




Ponceau




Source file for Fig. S8

IP ADGB IB ADGB




IP ADGB IB V5




Source files for Fig. S10


Input IB GFP




Input IB V5




IP GFP IB GFP



IP GFP IB V5



IgG IB GFP



IgG IB V5

