

1 **Nicotinic modulation insecticides act on diverse** 2 **receptor subtypes with distinct subunit compositions**

3 Wanjun Lu^{1#}, Zhihan Liu^{1#}, Xinyu Fan^{1#}, Xinzhong Zhang^{2#}, Xiaomu Qiao¹ and
4 Jia Huang^{1*}

5

6 ¹Ministry of Agriculture Key Laboratory of Molecular Biology of Crop Pathogens
7 and Insects, Institute of Insect Sciences, Zhejiang University, Hangzhou 310058,
8 China;

9 ²Tea Research Institute, Chinese Academy of Agricultural Sciences, Hangzhou
10 310008, China

11

12 #These authors contributed equally.

13 *Jia Huang. **Email:** huangj@zju.edu.cn

14

15 **Author Contributions:** J.H. designed the study. W.L., Z.L., X.F. and X.Z.
16 performed experiments. X.M. and J.H. analyzed data and wrote the manuscript.

17

Keywords: neonicotinoid, nicotinic acetylcholine receptor, resistance, insecticide, *Drosophila*

Abstract

Insect nicotinic acetylcholine receptors (nAChRs) are ligand gated ion channels mainly expressed in the central nervous system of insects. They are the directed targets of nicotinic modulation insecticides including neonicotinoids, the most widely used insecticides in the world. However, the resistance development from pests and the negative impacts on the pollinators affect their applications and create demand for the alternatives. Thus, it is very important to understand the mode of action of these insecticides at the molecular level, which is actually unclear for more than 30 years. In this study, we systematically examined the susceptibility of ten *Drosophila melanogaster* nAChR subunits mutants against eleven nicotinic modulation insecticides. Our results showed that there are several subtypes of nAChRs with distinct subunits compositions that are responsible for the toxicity of different insecticides, respectively. At least three of them are the major molecular targets of seven structurally similar neonicotinoids *in vivo*. On the other hand, the spinosyns may exclusively act on the $\alpha 6$ homomeric nAChR but not any other heteromeric pentamers. Behavioral assays

using thermogenetic tools further confirmed the bioassay results and support the idea that receptor activation rather than inhibition leads to the insecticidal effects of neonicotinoids. The present findings reveal native nAChR subunit interactions with various insecticides and have important implications for resistance management and the development of novel insecticides targeting this important ion channels.

Author Summary

The neonicotinoids and spinosyns make up about 27% of the insecticides by world market value. Novel insecticides like sulfoxaflor, flupyradifurone and triflumezopyrim are developed as alternatives due to the negative effects of neonicotinoids on pollinators. Although all act via insect nicotinic acetylcholine receptors, the mode of action is unclear. Our work shows that these insecticides act on diverse receptor subtypes with distinct subunit compositions. This finding

could lead to the development of more selective insecticides to control pests with minimal effects on beneficial insects.

Main Text

Introduction

Chemical insecticides have been widely used to control pests in agriculture, horticulture, forestry, homes and cities. They have also played a vital role in preventing the spread of human and animal vector-borne diseases. However, Insecticide resistance is a serious worldwide problem for invertebrate pest control, with more than 600 different insect and mite species having become resistant to at least one insecticide. In addition, there is at least one documented case of resistance for more than 335 insecticides/acaricides[1]. Therefore, there is great demand for effective insecticide resistance management (IRM) and development of new pest control compounds. To address both issues, we need to know the mode of action of insecticides: the process of how an insecticide works at a molecular level [2].

A complete understanding of the mode of action of an insecticide requires knowledge of how it affects a specific target site within an organism. Although most insecticides have multiple biological effects, toxicity is usually attributed to a

single major effect. For many insecticides, however, the exact molecular targets remain elusive. In order to ascribe whether a candidate protein is indeed the target for an insecticidal effect *in vivo*, it is not sufficient to demonstrate an *in vitro* biochemical interaction between an insecticide and a protein. Genetic evidence demonstrating an effect due to mutation of the candidate target is critical before it is possible to conclude that a given protein is the target of an insecticide.

The neonicotinoids (acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram, thiacloprid, and thiamethoxam) are remarkably effective to control agricultural pests, ectoparasites and arthropod vectors [3]. They are taken up by the roots or leaves and translocated to all parts of the plant due to high systemic activity, making them effectively toxic to wide range of sap-feeding and foliar feeding insects. Thus, neonicotinoids account for 24% of the global insecticide market, the largest market share of all chemical classes[1]. They act selectively on insect nicotinic acetylcholine receptors (nAChR) as agonists compared with the mammalian-selective nicotine. The spinosyns are a naturally derived, unique family of macrocyclic lactones which act on insect nAChR in an allosteric fashion. Besides, the sulfoximine sulfoxaflor, butenolide flupyradifurone and mesoionic triflumezopyrim are three newly developed insecticides which are also nAChR competitive modulators [4]. It is expected that the market of all above nAChR

targeting insecticides which show excellent insect to mammalian selectivity, will continue to grow. However, the molecular targets of neonicotinoids and other nAChR modulators remain unclear, mainly because we do not know the structure and assembly of native nAChRs in insects [5].

The cation-selective nAChRs are members of the Cys-loop ligand gated ion channel superfamily responsible for rapid excitatory neurotransmission. The functional nAChRs are homo- or heteromeric pentamers of structurally related subunits arranged around a central ion-conducting pore[6]. Each subunits has a extracellular N-terminal domain which contains six distinct regions (loops A–F) involved in ligand binding, four C-terminal transmembrane segments (TM1–TM4) and an intracellular loop between TM3 and TM4. nAChRs are divided into α -subunits possessing two adjacent cystine residues in loop C, while those subunits without this motif are termed non- α subunits. In vertebrates, 17 nAChR subunits have been identified, which can co-assemble to generate a diverse family of nAChR subtypes with different pharmacological properties and physiological functions. Insects have fewer nAChR subunits (10–12 subunits) according to the available genome data. Although co-immunoprecipitation studies have indicated potential associations of several subunits, the exact subunits composition of native insect nAChRs remains unknown[5]. Unlike the

vertebrate counterparts, heterologous expression of genuine arthropod α and β subunits has not been successful until recently two groups found that three ancillary proteins are essential for robust expression of arthropod nAChR heteromers [7, 8]. Thus for a long time, researchers used hybrid receptors with insect α subunits and mammalian/avian β subunits to study the interaction of insecticides and receptors. Such alternatives may not faithfully reflect all features of the native nAChRs [9].

In this study, we systematically examined the effects of total ten (seven α and three β) *Drosophila melanogaster* subunit mutants against eleven different nAChR targeting insecticides. We found that there are multiply subtypes of receptors with distinct subunits compositions which are responsible for the toxicity of different insecticides, respectively. Artificial activation/inhibition of subunit-expressing neurons also mimicked insecticides poisoning symptoms in pests. The elucidation of molecular targets of these economically important agrochemicals and the assembly of native nAChRs will be very helpful for resistance management and ecotoxicological evaluation on beneficial insects like predators and pollinators.

Results

Generation of nAChR β 1^{R81T} mutant

We got all 10 nAChRs knock-out mutants from Yi Rao's lab and found that KO of α 4 and β 1 was homozygous lethal. Thus we used a point mutation (T227M) allele of α 4 (*redeye*, *rye*) in bioassays, which is a dominant-negative mutation to cause reduced sleep phenotype in flies [10]. An R81T mutation of the nAChR β 1 was found in neonicotinoids-resistant peach aphids and later in cotton aphids [11, 12], so we introduced the homologous mutation into the β 1 of *Drosophila melanogaster* with CRISPR-Cas9-mediated homology-directed repair (HDR). The design of gRNA target site and HDR template was shown and the screen of successful R81T knock-in was accreted under imidacloprid selection pressure and confirmed by direct DNA sequencing (Figure 1 and S1).

nAChR mutants showed distinct resistance to multiple insecticides

We tested the effects of 10 nAChR mutants and some heterozygous mutants against 11 insecticides (Figure 2 and Table S1-11). The α 1 mutant showed moderate levels of resistance to imidacloprid, thiacloprid, acetamiprid and triflumezopyrim, the LC₅₀ resistance ratio (RR) is about 13.5 - 88.0. Its

heterozygous mutant also showed low levels of resistance to these insecticides. Besides, it showed a low but statistically significant increases of RR (2.7 - 3.7) to thiamethoxam, clothianidine, dinotefuran and nitenpyram. The $\alpha 2$ mutant also showed similar levels of resistance (17.2 - 48.5 in the terms of RR) to imidacloprid, thiacloprid and triflumezopyrim. For the $\alpha 3$ mutant, it showed small RR increases (2.7 - 5.5) to thiamethoxam, clothianidine, dinotefuran, nitenpyram, sulfoxaflor and flupyradifurone. The $\alpha 4$, $\alpha 5$, $\alpha 6$, $\alpha 7$ and $\beta 3$ mutants are sensitive to almost all the tested insecticides, the obvious exception is that the $\alpha 6$ homozygous mutant is resistant to spinetoram with a RR of 42.8 but the heterozygous mutant is close to the wild type (RR 1.2). The $\beta 1$ mutant exhibited medium to high resistance to all insecticides (23.9 - 398.3 in the terms of RR) except spinetoram, and its heterozygous mutant showed small increases of RR for most insecticides. The resistance profile of $\beta 2$ mutant is similar to that of $\alpha 1$ mutant, with 13.0 - 84.3 folds RR increases to imidacloprid, thiacloprid, acetamiprid and triflumezopyrim.

Both $\alpha 1$ and $\beta 1$ mutants showed variable resistance to multiply insecticides, thus we generated a $\alpha 1/\beta 1$ double mutant with recombination. However, the eggs laid by this combined mutant can not hatch for further experiments. A

recent paper also generated a $\beta 1$ R81T *Drosophila* and found that it has serious defects in reproduction and locomotion [13], however, the $\beta 1$ mutant we made here did not show any significant fitness cost (Figure S2). The sequences of $\alpha 5$, $\alpha 6$ and $\alpha 7$ are very close and show high similarity to the vertebrate nAChR $\alpha 7$ subunit. They can also form heteromeric ion channels in vitro with different combinations like $\alpha 5/\alpha 6$, $\alpha 5/\alpha 7$ and $\alpha 5/\alpha 6/\alpha 7$ [14]. Since only the $\alpha 6$ mutant showed resistance to spinetoram, we then wonder whether there is a genetic redundancy among these evolutionarily conserved gene. However, the $\alpha 5/\alpha 7$ double mutant was still sensitive to spinetoram (Table S11), indicating that the $\alpha 6$ homomeric channel could be the solo target for spinosyns.

Hyperactivating/silencing nAChR-expressing neurons mimics insecticides poisoning symptoms

The way insects react when they are exposed to neonicotinoids, sulfoxaflor, flupyradifurone and spinosyns are similar. The early-onset behaviors including hyperactivity, convulsion, uncoordinated movements, leg extension and tremors. At higher doses, these excitatory symptoms can induce severe tremors and complete paralysis that lead to death [15-17]. We then wondered whether

artificial activation of *nAChR*-expressing neurons would induce insecticides-like poisoning symptoms. Thus, we used the thermosensitive cation channel *Drosophila* TRPA1 to acutely hyper-stimulate these neurons with all available *nAChR* KI-Gal4 strains [18]. We found that expressing *trpA1* in *nAChR* $\alpha 1^{2A-GAL4}$, *nAChR* $\alpha 2^{2A-GAL4}$, *nAChR* $\alpha 3^{2A-GAL4}$, *nAChR* $\alpha 6^{2A-GAL4}$ and *nAChR* $\beta 2^{2A-GAL4}$ neurons strongly induced hyperactivity behavior at 32 °C, and eventually led to paralysis (Figure 3A, Video 1), which is similar to the above-mentioned symptoms. However, activation of *nAChR* $\beta 3^{2A-GAL4}$ neurons did not show any behavioral defects. These results parallel the above bioassay data that the deletion of $\alpha 1$, $\alpha 2$, $\alpha 3$, $\alpha 6$ and $\beta 2$ caused medium to high resistances to these insecticides respectively. Therefore, thermogenetic activation of some *nAChR*-expressing neurons in a short time window phenocopies the action of insecticides in target pests, which demonstrates that *in vivo* pharmacological activation of these subunits-containing nAChRs leads to toxicity and finally death.

The poisoning symptoms associated with triflumezopyrim is distinct from other nicotinic modulation insecticides since it inhibits rather than activates insect nAChRs. There is no any neuro-excitatory symptoms after treatment of triflumezopyrim, on the contrary, it induces lethargic poisoning characterized by

slow but coordinated leg movements and insects became less responsive to stimuli over time [19]. Thus, we chose to use *UAS-Shibire^{ts}* to inhibit *nAChR*-expressing neurons [20]. As expected, *nAChR* α 1^{2A-GAL4}, *nAChR* α 2^{2A-GAL4} and *nAChR* β 2^{2A-GAL4} neurons produced a “sluggish” behavior rather than hyperactivity (Figure 3B). The flies exhibited almost no translational or rotational body movement (Video 1). Silencing of *nAChR* α 3^{2A-GAL4} and *nAChR* α 6^{2A-GAL4} neurons also produced similar behaviors, further confirming that the α 3- and α 6-containing nAChRs can not be blocked by triflumezopyrim, otherwise both mutants would show resistance in bioassays.

Expression patterns of nAChRs in KO mutants

We confirmed that the KO coding regions were not detected or barely detectable with real-time PCR quantification (Figure S2). There was no big difference of expression levels of each subunits in these mutant flies, except that the β 3 was relatively less transcribed than other genes. For the α 1 heterozygous mutant, the mRNA levels of all subunits were almost same as the wild type control.

Discussion

216 The Insecticide Resistance Action Committee (IRAC) classifies neonicotinoids,
 217 sulfoximines, butenolides and mesoionics into sub-groups 4A, 4C, 4D and 4E
 218 respectively, according to their chemical similarity relations. However, our results
 219 clearly showed that sulfoxaflor and flupyradifurone may specifically act on the
 220 same nAChR subtype which consists of $\alpha 3$ and $\beta 1$ subunits (Figure 4A), albeit
 221 their big differences in chemical structures. More importantly, we found that the
 222 neonicotinoids act on distinct nAChR subtypes and such selectivity is not
 223 dependent on the aromatic heterocyclic (A), or the electron-withdrawing nitro or
 224 cyano moiety (X-Y) which is considered the key toxophore. Interestingly, the ring
 225 systems and the R_2 substituents in the open-chain structures are the determining
 226 factors (Figure 4). For example, the $\alpha 1$, $\alpha 2$, $\beta 1$ and $\beta 2$ mutants showed similar
 227 levels of resistance to imidacloprid and thiacloprid (both have a five-membered
 228 ring), indicating that they mainly act on the same $\alpha 1/\alpha 2/\beta 1/\beta 2$ pentamer (Figure
 229 4B). This is consistent with previous ex vivo recording results [21] and the two
 230 recent reconstituted studies in which both drugs act as partial agonists on the
 231 $\alpha 1/\alpha 2/\beta 1/\beta 2$ nAChR [7, 8]. Acetamiprid is structurally similar to thiacloprid with
 232 the cyanoimine pharmacophore, but the acyclic configuration changes its
 233 molecular target in vivo. It may act on the $\alpha 1/\beta 1/\beta 2$ nAChR and again the
 234 electrophysiological studies had already indicated that acetamiprid is nearly a full

agonist [21] and its potency on the recombinant louse $\alpha 1/\alpha 2/\beta 1/\beta 2$ nAChR is about 860 fold lower than that of thiacloprid [8]. Although thiamethoxam has a six-membered-ring, it is a pro-drug without intrinsic nAChR activity until metabolized to the active form clothianidine in plants and insects [22]. Therefore, thiamethoxam, clothianidine, dinotefuran and nitenpyram can be considered as the same type which have the N-methy substitution in the R₂ position and mainly act on the $\alpha 1/\alpha 3/\beta 1$ nAChR (Figure 4B). The neonicotinoids are traditionally divided into nitroimines (NNO₂), nitromethylenes (CHNO₂) or cyanoimines (NCN), but our findings proposed a new classification according to their major nAChR subtypes targets.

Despite the widespread use of neonicotinoids for almost four decades, the first and only field-evolved target-site resistance mutation (R81T in nAChR $\beta 1$) was reported in 2011 and it has only been found in two species to date [11, 12]. Such unusual phenomenon can be partially explained by our findings that the seven neonicotinoids have at least three distinct molecular targets in vivo. To some extent, the continuous use of different neonicotinoids is a kind of spontaneous insecticides rotations, which has been proven to be effective in mitigating or delaying resistance. New nicotine mimic insecticides like sulfoxaflor and flupyradifurone mainly act on another nAChR subtype which is distinct from

neonicotinoids (Figure 4), indicating their potential use in insecticides resistance management.

Electrophysiological studies with native tissues or recombinant receptors showed that low concentrations of neonicotinoids can block nAChR while higher concentrations cause receptor activation[7, 23]. Therefore, it is still unclear whether the insecticidal activity is the consequence of nAChR inhibition or activation in vivo. We found that transient artificial activation rather than inhibition of *nAChR*-expressing neuron is sufficient to induce neonicotinoids-like poisoning symptoms in flies (Figure 3). Thus, the overall effect of neonicotinoids is neuronal depolarizing by activation of nAChR which is more physiologically relevant.

Triflumezopyrim is the first member of a new class of mesoionic insecticides, which act via inhibition of the orthosteric binding site of the nAChR [19]. We found that the $\alpha 1/\alpha 2/\beta 1/\beta 2$ nAChR could be its major target like imidacloprid and thiacloprid, all these mutants showed high resistance to triflumezopyrim (Figure 4A). This is consistent with radioligand binding results in which triflumezopyrim potently displacing [³H]imidacloprid with a K_i value of 43 nM using the membrane preparations from the aphid [19]. Thermogenetic inhibition neurons expressing $\alpha 1$, $\alpha 2$ and $\beta 2$ also mimic the lethargic intoxication symptoms (Figure 3B). Thus, in order to maintain the durability and effectiveness of this new powerful tool for

control of hopper species in rice, it is very critical to avoid repeated use of triflumezopyrim with imidacloprid and thiacloprid.

The spinosyns including spinosad and spinoteram have been shown to act a population of nAChR that are not targeted by neonicotinoids, the binding site is also distinct to the orthosteric site [4]. The $\alpha 6$ subunit has been proposed as the main target of spinosyns since the resistance to spinosad in many insects is associated with loss-of-function mutations in the $\alpha 6$ gene [24], however, whether other subunits are involved is still unknown. We used spinoteram in bioassays and the results strongly indicated that spinosyns may specifically act on the $\alpha 6$ homomeric nAChR but not any other subtypes (Figure 4), which is consistent with a recent report using spinosad [25]. Thermogenetic activation of $\alpha 6$ -expressing neurons also induced spinosyns-like poisoning symptoms in flies.

Our current knowledge about the subunit composition of insect nAChRs is very limited. Immunoprecipitation data with subunit-specific antibodies showed that the *Drosophila* $\alpha 3$ and $\beta 1$ co-assemble within the same receptor complex [26]. Further studies from the same group indicated that $\alpha 1/\alpha 2/\beta 2$ and $\beta 1/\beta 2$ may co-assemble into the same receptor complex respectively [27]. Similar studies using the brown planthopper suggested that there are two populations of nAChRs which contain the *Drosophila* equivalent subunits combinations

$\alpha 1/\alpha 2/\beta 1$ and $\alpha 3/\beta 1/\beta 2$, respectively [28]. These previous findings are partially confirmed by the present results, as the $\alpha 3/\beta 1$, $\alpha 1/\alpha 3/\beta 1$, $\alpha 1/\beta 1/\beta 2$ and $\alpha 1/\alpha 2/\beta 1/\beta 2$ could be the major receptor subtypes for the tested insecticides, indicating that the $\beta 1$ subunit could be an indispensable component for all heteromeric pentamers (Figure 4). Besides, we noticed that for some insecticides, different subunits mutations contribute in an asymmetrical manner to resistance (Figure 4A). Therefore, there could be functional redundancy between some α -type subunits and we can not exclude the existence of other potential receptor subtypes such as $\alpha 1/\beta 1$ and $\alpha 3/\beta 1/\beta 2$. The diversity of insect nAChRs and their druggability make them remain an extremely important target for insecticides development.

Growing evidence indicates that sublethal doses of neonicotinoids like imidacloprid, thiamethoxam and clothianidin negatively affect wild and managed bees which are important pollinators in ecosystems and agriculture [29-31]. They reduce reproduction and colony development, perhaps by impairing foraging, homing and nursing behaviors of bees [32]. These severe sublethal effects have led to heavy restrictions on the use of above three neonicotinoids in Europe to protect pollinators [33]. The sulfoxaflor and flupyradifurone are potential alternatives for neonicotinoids, however, their risk to bees is controversial [34-

36]. Therefore, it is critical to understand the mode of action of these insecticides inside bees. Since most *Drosophila* nAChR subunit genes (except $\alpha 5$ and $\beta 3$) have one-to-one orthologs in the honeybee and bumblebee genomes [7], the expression and assembly of receptors could be conserved between flies and bees, suggesting that our results will enable further studies about the ecotoxicology and risk assessment for these nAChR modulators.

Materials and Methods

Insecticides

Imidacloprid (600g/LSC, Bayer CropScience, Germany), thiamethoxam (70%GZ, Syngenta, China), clothianidin (48%SC, HeNan Hansi crop protection, China), dinotefuran (20%SG, Mitsui Chemicals, Japan), nitenpyram (30%WG, ZinGrow, China), acetamiprid (20%SP, Noposion, China), thiacloprid (40%SC, Limin Chemical, China), sulfoxaflor (22%SC, Dow AgroSciences, USA), flupyradifurone (17%SC, Bayer CropScience, Germany), triflumezopyrim (10%SC, DuPont, USA), spinetoram (60g/LSC, Dow AgroSciences, USA) and triton X-100 (Sangon Biotech, China) were purchased commercially.

Fly strains

Flies were maintained and reared on conventional cornmeal-agar-molasses medium at 25 ± 1 °C, $60\% \pm 10\%$ humidity with a photoperiod of 12 hours light: 12 hours night. For experiments using *UAS-trpA1* and *UAS-Shibire^{ts}* transgenes, flies were reared at 21 °C. The following strains were sourced from the Bloomington Stock Center (Indiana University): *vas-cas* (#51323), *UAS-trpA1* (#26263), *UAS-Shibire^{ts}* (44222). All nAChR KO mutants and KI-Gal4 strains were gifts from Dr. Yi Rao (Deng et al., 2019) (Peking University). The *w¹¹¹⁸* used for outcrossing was used as wide-type for insecticide bioassays.

We generated the nAChR β 1^{R81T} mutant by CRISPR/Cas9 genome editing. The gRNA sequence (3L:4433329~4433352, ATCAAACGTTTGGTTAACTTTAG) was designed with flyCRISPR Target Finder (<https://flycrispr.org/target-finder/>) and cloned into the pDCC6 plasmid (addgene #59985). A 110 bp ssODN (single-strand oligodeoxynucleotide) was customer synthesized as the donor template to replace the targeted genomic region. This ssODN contained three nucleotides changes with two (CG to AC) conferring the R81T mutation and one synonymous

mutation (G to A) to prevent the re-cleavage from Cas9 after incorporation. Both gRNA plasmid and ssODN were microinjected into the embryos of *vas-cas* flies (BL #51323). The crossing and selection scheme was shown in the Figure S1.

Insecticide bioassays

3–5 day old and uniform size adult females were used in insecticide bioassays to assess the susceptibility of different fly strains. The testing method was modified from the IRAC susceptibility test method 026 (<https://irac-online.org/methods/>). Briefly, the required serial dilutions of insecticide solution are prepared in 200g/L sucrose using formulated insecticides. Approximately 5ml of insecticide solution is required for each concentration. A piece of dental wick (1cm) is placed in a standard *Drosophila* vials and treated with 800 µL 20% aqueous sucrose with or without insecticide. The vials were kept upside down until all flies became active to avoid flies getting trapped in the dental wick. The bioassay was assessed after 48 h, dead flies as well as seriously affected flies displaying no coordinated movement, that were unable to walk up the vial, or unable to get to their feet were cumulatively scored as 'affected'. The lethal concentrations LC₅₀ were calculated by probit analysis using the Polo Plus software (LeOra Software,

Berkeley, CA, USA). Non-linear log dose-response curves were generated in Graphpad Prism 8.21 (Graphpad Software Inc., La Jolla, CA, USA).

Thermogenetic activation and silencing assays

Flies for TRPA1-mediated thermogenetic activation and Shibire-mediated silencing experiments were collected upon eclosion and reared in vials containing standard food medium at 21 °C for 5-8 days. For thermogenetic activation with the *UAS-trpA1* transgene, 10 flies were transferred to new empty vials by gently inspiration, and then the assays were performed at 23 °C and 32 °C for 10 minutes. The percentage of paralysis behavior, in which the animal lies on its back with little effective movement of the legs and wings, was measured. For silencing assays, *UAS-Shibire^{ts}* transgene was used and flies were also transferred to fly vials at 23 °C and 32 °C for 10 minutes.

Real-time quantitative PCR

The relative transcription levels of *nAChRs* in different KO mutants were examined using real-time quantitative PCR performed with an CFX96TM Real-Time PCR System (Bio-rad, Hercules, USA). Total RNA was isolated with Trizol reagent according to the manufacturer's instructions. Residual genomic DNA was removed by RQ1 RNase-Free DNase (Promega). Total RNA was reverse transcribed to cDNA with the EasyScript First-Strand cDNA Synthesis SuperMix (Transgene, Beijing, China). qPCR with gene-specific primers was performed with the ChamQ Universal SYBR qPCR Master Mix (Vazyme, Nanjing, China) to investigate relative expression levels of different *nAChRs*. The *RpL32* (ribosomal protein L32) was used as an internal control. Relative expression of *nAChRs* were normalized to the reference (*RpL32*) using the $2^{-\Delta\Delta CT}$ method.

Fecundity and development assays

10 pairs of freshly emerged couples of wild type control and $\beta 1^{R81T}$ mutant were transferred into vials containing normal food for 72 hr. These flies were then transferred into a new dish which is used for egg-laying assay. The numbers of egg laid in each dishes were recorded after 24 hr. To calculate the larvae to pupae rate, 60 second-instar larvae were collected and transferred into a new

vial as one group. The numbers of pupae in each vial were recorded after 7 days in an incubator. Each genotypes were repeated for at least three times with duplicates.

Climbing assay

About three-day-old male flies were collected with CO₂ anesthesia into groups of 10, and then allowed to recover for 2 days. A climbing tube consisted of two vials with 90 mm height and 20 mm diameter. The flies were filmed for 30 s with a SONY HDR-CX900E camera. The climbing index (percentage of flies in the upper half of the vial) were determined at 5 s intervals, after the flies had been tapped down to the bottom of the vials.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (31802019, 32072496) and Zhejiang Provincial Fund for Distinguished Young Scholars (LR19C140002). We thank Yi Rao (Peking University) for fly stocks.

References

1. Sparks TC, Crossthwaite AJ, Nauen R, Banba S, Cordova D, Earley F, et al. Insecticides, biologics and nematicides: Updates to IRAC's mode of action classification - a tool for resistance management. *Pestic Biochem Physiol.* 2020;167:104587. Epub 2020/06/13. doi: 10.1016/j.pestbp.2020.104587. PubMed PMID: 32527435.
2. Casida JE, Bryant RJ. The ABCs of pesticide toxicology: amounts, biology, and chemistry. *Toxicol Res (Camb).* 2017;6(6):755-63. doi: 10.1039/c7tx00198c. PubMed PMID: 30090540; PubMed Central PMCID: PMC6062263.
3. Jeschke P, Nauen R, Beck ME. Nicotinic Acetylcholine Receptor Agonists: A Milestone for Modern Crop Protection. *Angew Chem Int Edit.* 2013;52(36):9464-85. doi: 10.1002/anie.201302550. PubMed PMID: WOS:000323474600015.
4. Crossthwaite AJ, Bigot A, Camblin P, Goodchild J, Lind RJ, Slater R, et al. The invertebrate pharmacology of insecticides acting at nicotinic acetylcholine receptors. *J Pestic Sci.* 2017;42(3):67-83. Epub 2017/08/20. doi: 10.1584/jpestics.D17-019. PubMed PMID: 30363948; PubMed Central PMCID: PMC6183333.
5. Matsuda K, Ihara M, Sattelle DB. Neonicotinoid Insecticides: Molecular Targets, Resistance, and Toxicity. *Annu Rev Pharmacol Toxicol.* 2020;60:241-55. Epub 2020/01/10. doi: 10.1146/annurev-pharmtox-010818-021747. PubMed PMID: 31914891.
6. Casida JE. Neonicotinoids and Other Insect Nicotinic Receptor Competitive Modulators: Progress and Prospects. *Annu Rev Entomol.* 2018;63:125-44. Epub 2018/01/13. doi: 10.1146/annurev-ento-020117-043042. PubMed PMID: 29324040.

- 438 7. Ihara M, Furutani S, Shigetou S, Shimada S, Niki K, Komori Y, et al. Cofactor-
439 enabled functional expression of fruit fly, honeybee, and bumblebee nicotinic receptors
440 reveals picomolar neonicotinoid actions. *Proc Natl Acad Sci U S A*. 2020;117(28):16283-
441 91. Epub 2020/07/03. doi: 10.1073/pnas.2003667117. PubMed PMID: 32611810;
442 PubMed Central PMCID: PMCPMC7368294.
- 443 8. Rufener L, Kaur K, Sarr A, Aaen SM, Horsberg TE. Nicotinic acetylcholine
444 receptors: Ex-vivo expression of functional, non-hybrid, heteropentameric receptors
445 from a marine arthropod, *Lepeophtheirus salmonis*. *PLoS Pathog*. 2020;16(7):e1008715.
446 Epub 2020/07/28. doi: 10.1371/journal.ppat.1008715. PubMed PMID: 32716968;
447 PubMed Central PMCID: PMCPMC7419010.
- 448 9. Ihara M, Buckingham SD, Matsuda K, Sattelle DB. Modes of Action, Resistance
449 and Toxicity of Insecticides Targeting Nicotinic Acetylcholine Receptors. *Curr Med Chem*.
450 2017;24(27):2925-34. Epub 2017/02/09. doi: 10.2174/0929867324666170206142019.
451 PubMed PMID: 28176635.
- 452 10. Shi M, Yue Z, Kuryatov A, Lindstrom JM, Sehgal A. Identification of Redeye, a new
453 sleep-regulating protein whose expression is modulated by sleep amount. *Elife*.
454 2014;3:e01473. Epub 2014/02/06. doi: 10.7554/eLife.01473. PubMed PMID: 24497543;
455 PubMed Central PMCID: PMCPMC3912633.
- 456 11. Bass C, Puinean AM, Andrews M, Cutler P, Daniels M, Elias J, et al. Mutation of a
457 nicotinic acetylcholine receptor beta subunit is associated with resistance to
458 neonicotinoid insecticides in the aphid *Myzus persicae*. *Bmc Neuroscience*. 2011;12. doi:
459 Artn 51
460 10.1186/1471-2202-12-51. PubMed PMID: WOS:000291969900001.

- 461 12. Shi XG, Zhu YK, Xia XM, Qiao K, Wang HY, Wang KY. The mutation in nicotinic
462 acetylcholine receptor beta 1 subunit may confer resistance to imidacloprid in *Aphis*
463 *gossypii* (Glover). *J Food Agric Environ*. 2012;10(2):1227-30. PubMed PMID:
464 WOS:000305310800064.
- 465 13. Homem RA, Buttery B, Richardson E, Tan Y, Field LM, Williamson MS, et al.
466 Evolutionary trade - offs of insecticide resistance — The fitness costs associated with
467 target - site mutations in the nAChR of *Drosophila melanogaster*. *Molecular Ecology*.
468 2020;29(14):2661-75. doi: 10.1111/mec.15503.
- 469 14. Lansdell SJ, Collins T, Goodchild J, Millar NS. The *Drosophila* nicotinic
470 acetylcholine receptor subunits $\alpha 5$ and $\alpha 7$ form functional homomeric and
471 heteromeric ion channels. *BMC Neurosci*. 2012;13:73. Epub 2012/06/26. doi:
472 10.1186/1471-2202-13-73. PubMed PMID: 22727315; PubMed Central PMCID:
473 PMCPMC3444433.
- 474 15. Salgado VL. Studies on the mode of action of spinosad: Insect symptoms and
475 physiological correlates. *Pesticide Biochemistry and Physiology*. 1998;60(2):91-102. doi:
476 DOI 10.1006/pest.1998.2332. PubMed PMID: WOS:000074891100003.
- 477 16. Watson GB, Loso MR, Babcock JM, Hasler JM, Letherer TJ, Young CD, et al. Novel
478 nicotinic action of the sulfoximine insecticide sulfoxaflor. *Insect Biochem Molec*.
479 2011;41(7):432-9. doi: 10.1016/j.ibmb.2011.01.009. PubMed PMID:
480 WOS:000293041200003.
- 481 17. Nauen R, Jeschke P, Velten R, Beck ME, Ebbinghaus-Kintscher U, Thielert W, et al.
482 Flupyradifurone: a brief profile of a new butenolide insecticide. *Pest Management*
483 *Science*. 2015;71(6):850-62. doi: 10.1002/ps.3932. PubMed PMID:
484 WOS:000354035100009.

- 485 18. Hamada FN, Rosenzweig M, Kang K, Pulver SR, Ghezzi A, Jegla TJ, et al. An
486 internal thermal sensor controlling temperature preference in *Drosophila*. *Nature*.
487 2008;454(7201):217-20. doi: 10.1038/nature07001. PubMed PMID: 18548007; PubMed
488 Central PMCID: PMC2730888.
- 489 19. Cordova D, Benner EA, Schroeder ME, Holyoke CW, Jr., Zhang W, Pahutski TF, et
490 al. Mode of action of triflumezopyrim: A novel mesoionic insecticide which inhibits the
491 nicotinic acetylcholine receptor. *Insect Biochem Mol Biol*. 2016;74:32-41. Epub
492 2016/05/01. doi: 10.1016/j.ibmb.2016.04.008. PubMed PMID: 27130855.
- 493 20. Kitamoto T. Conditional disruption of synaptic transmission induces male-male
494 courtship behavior in *Drosophila*. *Proc Natl Acad Sci U S A*. 2002;99(20):13232-7. doi:
495 10.1073/pnas.202489099. PubMed PMID: 12239352; PubMed Central PMCID:
496 PMC130616.
- 497 21. Tan J, Galligan JJ, Hollingworth RM. Agonist actions of neonicotinoids on nicotinic
498 acetylcholine receptors expressed by cockroach neurons. *Neurotoxicology*.
499 2007;28(4):829-42. doi: 10.1016/j.neuro.2007.04.002. PubMed PMID:
500 WOS:000248875500014.
- 501 22. Nauen R, Ebbinghaus-Kintscher U, Salgado VL, Kausmann M. Thiamethoxam is a
502 neonicotinoid precursor converted to clothianidin in insects and plants. *Pesticide*
503 *Biochemistry and Physiology*. 2003;76(2):55-69. doi: 10.1016/s0048-3575(03)00065-8.
- 504 23. Salgado VL. Selective actions of insecticides on desensitizing and non-
505 desensitizing nicotinic acetylcholine receptors in cockroach (*Periplaneta americana*)
506 neurons. *Pest Management Science*. 2021;77(8):3663-72. doi: 10.1002/ps.6396.
507 PubMed PMID: WOS:000643154400001.

- 508 24. Zuo YY, Xue YX, Wang ZY, Ren X, Aioub AAA, Wu YD, et al. Knockin of the G275E
509 mutation of the nicotinic acetylcholine receptor (nAChR) alpha 6 confers high levels of
510 resistance to spinosyns in *Spodoptera exigua*. *Insect Science*. 2021. doi: 10.1111/1744-
511 7917.12922. PubMed PMID: WOS:000651046300001.
- 512 25. Perry T, Chen W, Ghazali R, Yang YT, Christesen D, Martelli F, et al. Role of
513 nicotinic acetylcholine receptor subunits in the mode of action of neonicotinoid,
514 sulfoximine and spinosyn insecticides in *Drosophila melanogaster*. *Insect Biochem Mol*
515 *Biol*. 2021;131:103547. Epub 2021/02/07. doi: 10.1016/j.ibmb.2021.103547. PubMed
516 PMID: 33548485.
- 517 26. Chamaon K, Schulz R, Smalla KH, Seidel B, Gundelfinger ED. Neuronal nicotinic
518 acetylcholine receptors of *Drosophila melanogaster*: the alpha-subunit dalpha3 and the
519 beta-type subunit ARD co-assemble within the same receptor complex. *FEBS Lett*.
520 2000;482(3):189-92. Epub 2000/10/12. doi: 10.1016/s0014-5793(00)02057-3. PubMed
521 PMID: 11024458.
- 522 27. Chamaon K, Smalla KH, Thomas U, Gundelfinger ED. Nicotinic acetylcholine
523 receptors of *Drosophila*: three subunits encoded by genomically linked genes can co-
524 assemble into the same receptor complex. *Journal of Neurochemistry*. 2002;80(1):149-
525 57. doi: DOI 10.1046/j.0022-3042.2001.00685.x. PubMed PMID:
526 WOS:000173059900016.
- 527 28. Li J, Shao Y, Ding Z, Bao H, Liu Z, Han Z, et al. Native subunit composition of two
528 insect nicotinic receptor subtypes with differing affinities for the insecticide
529 imidacloprid. *Insect Biochem Mol Biol*. 2010;40(1):17-22. Epub 2009/12/17. doi:
530 10.1016/j.ibmb.2009.12.003. PubMed PMID: 20005950.

- 531 29. Tsvetkov N, Samson-Robert O, Sood K, Patel HS, Malena DA, Gajiwala PH, et al.
532 Chronic exposure to neonicotinoids reduces honey bee health near corn crops. *Science*.
533 2017;356(6345):1395-7. Epub 2017/07/01. doi: 10.1126/science.aam7470. PubMed
534 PMID: 28663503.
- 535 30. Woodcock BA, Bullock JM, Shore RF, Heard MS, Pereira MG, Redhead J, et al.
536 Country-specific effects of neonicotinoid pesticides on honey bees and wild bees.
537 *Science*. 2017;356(6345):1393-5. Epub 2017/07/01. doi: 10.1126/science.aaa1190.
538 PubMed PMID: 28663502.
- 539 31. Rundlof M, Andersson GK, Bommarco R, Fries I, Hederstrom V, Herbertsson L, et
540 al. Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature*.
541 2015;521(7550):77-80. doi: 10.1038/nature14420. PubMed PMID: 25901681.
- 542 32. Crall JD, Switzer CM, Oppenheimer RL, Ford Versypt AN, Dey B, Brown A, et al.
543 Neonicotinoid exposure disrupts bumblebee nest behavior, social networks, and
544 thermoregulation. *Science*. 2018;362(6415):683-6. Epub 2018/11/10. doi:
545 10.1126/science.aat1598. PubMed PMID: 30409882.
- 546 33. Cressey D. Neonics Vs Bees. *Nature*. 2017;551(7679):156-8. PubMed PMID:
547 WOS:000414734200022.
- 548 34. Siviter H, Brown MJF, Leadbeater E. Sulfoxaflor exposure reduces bumblebee
549 reproductive success. *Nature*. 2018;561(7721):109-12. doi: 10.1038/s41586-018-0430-6.
550 PubMed PMID: 30111837.
- 551 35. Haas J, Zaworra M, Glaubitz J, Hertlein G, Kohler M, Lagojda A, et al. A
552 toxicogenomics approach reveals characteristics supporting the honey bee (*Apis*
553 *mellifera* L.) safety profile of the butenolide insecticide flupyradifurone. *Ecotoxicol*

554 Environ Saf. 2021;217:112247. Epub 2021/04/27. doi: 10.1016/j.ecoenv.2021.112247.
555 PubMed PMID: 33901780.

556 36. Tosi S, Nieh JC, Brandt A, Colli M, Fourrier J, Giffard H, et al. Long-term field-
557 realistic exposure to a next-generation pesticide, flupyradifurone, impairs honey bee
558 behaviour and survival. Commun Biol. 2021;4(1):805. Epub 2021/06/30. doi:
559 10.1038/s42003-021-02336-2. PubMed PMID: 34183763; PubMed Central PMCID:
560 PMCPMC8238954.

561
562

Figures




Figure 1. Generation of the nAChR β 1^{R81T} mutant by CRISPR/Cas9 genome

editing. (A) Schematic of the nAChR β 1 locus and the sequence of donor construct. The boxes represent exons and the coding regions are shown in blue. The gRNA sequence is indicated in red and the codon for amino acid substitution (CGT to ACT) is highlighted in green. One synonymous mutation (G to A) is also introduced in the PAM region (in yellow) to prevent the re-cleavage from Cas9 after successful integration. (B) The sequence comparison between wild type and point mutation flies. The nucleotides replaced are highlighted in green and yellow boxes.




Figure 2. Non-linear Log-dose mortality data for tested insecticides against ten *Drosophila* nAChR homozygous mutants including eight null alleles and two point mutation alleles ($\alpha 4^{T227M}$ and $\beta 1^{R81T}$). Mortality (0-1 means 0-100% in terms of percentage) of control and mutant female adults after 48 hour exposure to

increasing concentrations of insecticides. Error bars represent standard deviations.




Figure 3. The effects of artificial neuronal activation and inhibition in various *nAChR*-expressing neurons. (A) Thermogenetic activation of five *nAChR*-expressing neurons using *UAS-trpA1* induced paralysis behavior. (B) Thermogenetic silencing of five *nAChR*-expressing neurons using *UAS-Shibire^{ts}* decreased activity. n = 30-50.




Figure 4. The resistance patterns of tested insecticides on different nAChR mutants (A) and the proposed target receptor subtypes for neonicotinoids and others (B). Various resistance ratios are grouped and represented as different

600 colors in the heatmap. Thiamethoxam is considered as a prodrug of clothianidine
601 and not listed in the structural formula.
602

603 **Video 1.** The effects of thermogenetic activation and inhibition in *nAChR α 1*-
604 expressing neurons. The following transgenes were used: *nAChR α 1^{2A-GAL4}* >
605 *UAS-trpA1*; *nAChR α 1^{2A-GAL4}* > *UAS-Shibire^{ts}*. Other *nAChR* KI-Gal4 strains like
606 *nAChR α 2^{2A-GAL4}*, *nAChR α 3^{2A-GAL4}*, *nAChR α 6^{2A-GAL4}* and *nAChR β 2^{2A-GAL4}* also
607 produced similar behaviors when stimulated under 32 °C, these videos are not
608 shown.

Supplementary Information for

Nicotinic modulation insecticides act on diverse receptor subtypes with distinct subunit compositions

Wanjun Lu, Zhihan Liu, Xinyu Fan, Xinzhong Zhang, Xiaomu Qiao and Jia Huang*

* Jia Huang

huangj@zju.edu.cn

This PDF file includes:

Figures S1 to S3

Tables S1 to S12




Figure S1 The crossing schemes to establish the *nAChRβ1^{R81T}* knock-in line. The HDR event was isolated by imidacloprid selection and confirmed by PCR. The *vas-Cas9* (3XP3 RFP) was removed by the absence of red fluorescence in eyes.




Figure S2 Effects of *nAChRβ1^{R81T}* point mutation on number of eggs laid (A), pupation rate of larvae (B) and negative geotaxis behavior (C).




Figure S3 Expression patterns of the nAChR genes in different KO mutants.

Table S1 Log dose probit mortality data and resistance ratios for imidacloprid

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	8.6	7.5-9.9	26.6	21.5-35.9	1.0	1.0
$\alpha 1^{-/-}$	296.7	161.4-519.4	14555	5396.3-85114.0	34.5	547.2
$\alpha 1^{+/-}$	150.4	74.7-272.7	935.5	456.0-5243.1	17.5	35.2
$\alpha 2^{-/-}$	417.0	254.8-686.0	3113.5	1549.4-13500.0	48.5	117.0
$\alpha 2^{+/-}$	52.5	38.7-69.6	114.0	82.2-259.1	6.1	4.3
$\alpha 3^{-/-}$	10.7	9.7-11.7	16.7	14.7-21.1	1.2	0.6
$\alpha 4^{T227M}$	22.9	19.2-27.2	58.0	45.2-86.9	2.7	2.2
$\alpha 5^{-/-}$	9.6	7.5-12.1	27.5	19.9-50.2	1.1	1.0
$\alpha 6^{-/-}$	7.8	6.2-10.0	28.2	19.9-49.7	0.9	1.1
$\alpha 7^{-/-}$	22.8	20.4-25.5	43.6	37.1-55.8	2.7	1.6
$\beta 1^{R81T}$	565.7	407.3-781.4	2885.7	1835.2-6036.1	65.8	108.5
$\beta 1^{R81T/+}$	87.3	73.8-103.4	210.3	165.0-312.2	10.2	7.9
$\beta 2^{-/-}$	725.4	513.6-1027.5	4702.2	2840.2-10564.0	84.3	176.8
$\beta 2^{+/-}$	24.7	21.6-28.4	42.3	35.1-60.5	2.9	1.6
$\beta 3^{-/-}$	13.1	11.6-14.7	27.5	23.1-35.6	1.5	1.0

Table S2 Log dose probit mortality data and resistance ratios for thiacloprid

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	30.4	28.3-32.6	49.3	44.6-56.7	1.0	1.0
$\alpha I^{-/-}$	2674.9	2007.8-3553.3	8940.1	6082.5-17994.0	88.0	181.3
$\alpha I^{+/-}$	89.2	67.1-116.2	182.2	134.3-394.2	2.9	3.7
$\alpha 2^{-/-}$	1031.0	783.7-1384.3	14812.0	8693.3-31445.0	33.9	300.4
$\alpha 2^{+/-}$	37.3	31.6-43.9	86.2	68.2-126.3	1.2	1.7
$\alpha 3^{-/-}$	36.9	31.0-43.8	93.4	73.1-138.0	1.2	1.9
$\alpha 4^{T227M}$	27.0	16.7-40.5	106.2	63.4-357.9	0.9	2.2
$\alpha 5^{-/-}$	20.0	16.9-23.6	46.7	36.9-68.4	0.7	0.9
$\alpha 6^{-/-}$	26.1	22.6-29.9	78.5	63.8-105.1	0.9	1.6
$\alpha 7^{-/-}$	40.2	33.7-47.5	95.3	75.1-142.7	1.3	1.9
βI^{R81T}	1342.7	802.5-2229.7	8533.6	4418.4-32285.0	44.2	173.1
$\beta I^{R81T/+}$	40.3	26.3-62.2	106.9	67.4-458.1	1.3	2.2
$\beta 2^{-/-}$	1677.8	922.8-3425.5	21015.0	8759.4-98473.0	54.3	401.8
$\beta 2^{+/-}$	11.8	8.7-14.4	32.5	24.6-58.3	0.4	0.7
$\beta 3^{-/-}$	39.4	33.5-46.4	88.0	70.0-128.4	1.3	1.8

Table S3 Log dose probit mortality data and resistance ratios for acetamiprid

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	9.7	7.4-13.1	22.0	15.5-50.7	1.0	1.0
$\alpha I^{-/-}$	131.3	118.0-146.2	239.4	204.7-306.1	13.5	10.9
$\alpha I^{+/-}$	43.6	31.0-59.2	90.2	64.7-240.8	4.5	4.1
$\alpha 2^{-/-}$	15.2	11.4-20.5	37.9	26.2-89.1	1.6	1.7
$\alpha 3^{-/-}$	16.9	14.7-19.5	43.1	34.5-61.8	1.7	2.0
$\alpha 4^{T227M}$	9.4	8.2-10.9	17.3	14.2-24.8	1.0	0.8
$\alpha 5^{-/-}$	10.1	8.8-11.5	23.9	19.7-32.0	1.0	1.1
$\alpha 6^{-/-}$	22.6	18.5-28.0	63.7	46.6-109.0	2.3	2.9
$\alpha 7^{-/-}$	27.4	18.8-41.2	55.3	38.0-204.6	2.8	2.5
βI^{R81T}	231.5	214.7-249.5	397.7	355.5-465.5	23.9	18.1
$\beta I^{R81T/+}$	31.2	27.4-35.4	50.5	43.0-67.6	3.2	2.3
$\beta 2^{-/-}$	126.5	109.6-146.4	238.3	195.2-340.1	13.0	10.8
$\beta 2^{+/-}$	24.9	22.1-28.3	38.0	32.4-51.8	2.6	1.7
$\beta 3^{-/-}$	17.7	13.5-23.3	55.6	38.2-111.7	1.8	2.5

Table S4 Log dose probit mortality data and resistance ratios for thiamethoxam

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	8.2	7.6-8.9	15.3	13.4-18.4	1.0	1.0
$\alpha 1^{-/-}$	30.5	22.7-41.7	65.8	46.6-158.5	3.7	4.3
$\alpha 2^{-/-}$	7.9	6.8-9.2	14.9	12.2-21.2	1.0	1.0
$\alpha 3^{-/-}$	32.9	29.5-36.4	71.2	61.6-87.1	4.0	4.7
$\alpha 4^{T227M}$	4.1	2.6-5.9	9.9	6.6-34.8	0.5	0.6
$\alpha 5^{-/-}$	7.3	6.6-8.0	11.3	10.0-13.6	0.9	0.7
$\alpha 6^{-/-}$	10.8	10.0-11.6	16.7	14.8-20.4	1.3	1.1
$\alpha 7^{-/-}$	8.2	7.2-9.3	16.5	13.7-22.1	1.0	1.1
$\beta 1^{R81T}$	1935.5	1756.0-2142.6	4783.9	4047.9-5980.8	236.0	312.7
$\beta 1^{R81T/+}$	18.3	12.9-25.0	38.4	27.4-97.2	2.2	2.5
$\beta 2^{-/-}$	12.8	11.0-14.9	25.8	20.9-37.0	1.6	1.7
$\beta 3^{-/-}$	11.8	8.9-20.0	22.6	15.4-121.0	1.4	1.5

Table S5 Log dose probit mortality data and resistance ratios for clothianidin

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	3.5	3.1-3.9	8.4	7.0-10.9	1.0	1.0
$\alpha 1^{-/-}$	10.3	9.2-11.4	20.5	17.5-26.1	2.9	2.4
$\alpha 2^{-/-}$	4.7	4.3-5.2	8.5	7.4-10.5	1.3	1.0
$\alpha 3^{-/-}$	11.5	8.1-16.7	25.9	17.5-78.1	3.3	3.1
$\alpha 4^{T227M}$	2.5	2.1-2.9	6.3	4.9-9.5	0.7	0.8
$\alpha 5^{-/-}$	2.8	2.5-3.2	4.4	3.8-5.7	0.8	0.5
$\alpha 6^{-/-}$	5.1	4.6-5.7	9.4	8.1-11.9	1.5	1.1
$\alpha 7^{-/-}$	6.1	5.3-6.9	17.3	14.1-23.2	1.7	2.1
$\beta 1^{R81T}$	969.4	730.6-1278.8	4940.0	3300.4-9182.5	277.0	588.1
$\beta 1^{R81T}/+$	13.4	11.8-15.2	21.2	18.1-28.7	3.8	2.5
$\beta 2^{-/-}$	3.4	3.0-4.0	6.2	5.2-8.6	1.0	0.7
$\beta 3^{-/-}$	7.6	7.0-8.4	13.5	11.7-16.9	2.2	1.6

Table S6 Log dose probit mortality data and resistance ratios for dinotefuran

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	5.5	5.2-5.9	11.4	10.2-13.0	1.0	1.0
$\alpha 1^{-/-}$	15.0	13.6-17.0	25.9	21.7-34.8	2.7	2.3
$\alpha 2^{-/-}$	7.2	6.5-7.9	12.3	10.7-15.4	1.3	1.1
$\alpha 3^{-/-}$	30.4	27.6-33.2	55.0	48.4-66.0	5.5	4.8
$\alpha 3^{+/-}$	16.2	14.1-18.7	29.2	24.2-41.3	2.9	2.6
$\alpha 4^{T227M}$	3.7	3.2-4.3	7.0	5.7-10.2	0.7	0.6
$\alpha 5^{-/-}$	6.5	6.0-7.2	9.7	8.6-11.8	1.2	0.9
$\alpha 6^{-/-}$	8.7	7.8-9.8	16.7	14.1-21.7	1.6	1.5
$\alpha 7^{-/-}$	7.3	6.7-8.0	10.5	9.5-12.2	1.3	0.9
$\beta 1^{R81T}$	2190.9	1790.6-2702.1	4803.3	3659.6-7985.8	398.3	421.3
$\beta 1^{R81T/+}$	9.3	8.1-10.6	15.6	13.0-22.2	1.7	1.4
$\beta 2^{-/-}$	10.3	9.0-11.8	17.6	14.7-24.7	1.9	1.5
$\beta 3^{-/-}$	11.4	8.0-20.6	23.7	15.4-201.0	2.1	2.1

Table S7 Log dose probit mortality data and resistance ratios for nitenpyram

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	19.8	17.9-22.2	36.6	30.8-48.9	1.0	1.0
$\alpha I^{-/-}$	54.4	43.5-69.5	115.2	85.4-220.8	2.7	3.1
$\alpha 2^{-/-}$	15.1	13.4-17.0	33.1	27.7-43.0	0.8	0.9
$\alpha 3^{-/-}$	53.7	44.8-64.3	147.3	113.2-223.5	2.7	4.0
$\alpha 4^{T227M}$	13.4	11.2-15.7	28.9	23.0-44.3	0.7	0.8
$\alpha 5^{-/-}$	16.0	14.5-17.5	24.7	22.0-29.9	0.8	0.7
$\alpha 6^{-/-}$	19.4	16.1-24.8	35.4	27.7-61.1	1.0	1.0
$\alpha 7^{-/-}$	13.6	12.2-15.2	25.9	22.0-33.2	0.7	0.7
βI^{R8IT}	3629.0	2906.8-4770.6	11394.0	7852.2-21070.0	183.3	311.3
$\beta I^{R8IT/+}$	30.3	26.6-34.7	51.4	43.2-70.4	1.5	1.4
$\beta 2^{-/-}$	29.6	25.8-34.0	52.2	43.4-72.6	1.5	1.4
$\beta 3^{-/-}$	17.1	15.5-19.0	29.6	25.5-37.5	0.9	0.8

Table S8 Log dose probit mortality data and resistance ratios for flupyradifurone

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	38.8	36.1-41.8	74.6	66.3-87.5	1.0	1.0
$\alpha 1^{-/-}$	59.2	52.7-66.6	121.6	102.2-157.6	1.5	1.6
$\alpha 2^{-/-}$	53.6	49.0-59.7	100.1	85.4-127.9	1.4	1.3
$\alpha 3^{-/-}$	108.1	98.4-119.0	171.0	150.4-208.7	2.8	2.3
$\alpha 4^{T227M}$	33.0	28.5-38.3	64.3	52.3-92.5	0.9	0.9
$\alpha 5^{-/-}$	34.9	26.8-46.2	88.8	62.3-187.3	0.9	1.2
$\alpha 6^{-/-}$	55.3	33.2-99.2	129.7	79.8-1051.9	1.4	1.7
$\alpha 7^{-/-}$	62.5	57.1-68.5	127.0	110.4-153.7	1.6	1.7
$\beta 1^{R81T}$	842.0	767.5-924.5	2151.2	1849.8-2616.5	21.7	28.8
$\beta 1^{R81T/+}$	81.6	70.7-94.4	151.8	124.8-214.9	2.1	2.0
$\beta 2^{-/-}$	60.7	53.1-69.3	102.9	86.5-140.9	1.6	1.4
$\beta 3^{-/-}$	54.5	46.2-64.5	91.2	74.5-137.6	1.4	1.2

Table S9 Log dose probit mortality data and resistance ratios for sulfoxaflor

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	59.4	56.1-62.8	90.6	83.9-99.9	1.0	1.0
$\alpha 1^{-/-}$	54.8	50.0-60.3	85.3	75.3-103.2	0.9	0.9
$\alpha 2^{-/-}$	82.0	73.9-91.0	142.7	122.8-181.6	1.4	1.6
$\alpha 3^{-/-}$	207.8	185.6-236.3	306.0	263.2-402.5	3.5	3.4
$\alpha 4^{T227M}$	73.0	50.8-108.0	166.2	110.7-594.5	1.2	1.8
$\alpha 5^{-/-}$	55.7	50.7-61.3	88.5	77.9-107.6	0.9	1.0
$\alpha 6^{-/-}$	63.0	57.3-69.0	96.6	86.0-115.7	1.1	1.1
$\alpha 7^{-/-}$	56.6	51.5-62.1	88.2	78.0-106.2	1.0	1.0
$\beta 1^{R81T}$	948.5	726.2-1245.3	2280.2	1628.6-4643.7	16.0	25.2
$\beta 1^{R81T/+}$	116.3	102.9-131.0	175.1	152.0-222.1	2.0	1.9
$\beta 2^{-/-}$	50.3	46.4-54.8	76.2	67.6-91.7	0.8	0.8
$\beta 3^{-/-}$	64.2	59.7-68.1	89.7	83.9-98.4	1.1	1.0

Table S10 Log dose probit mortality data and resistance ratios for triflumezopyrim

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	28.2	24.7-31.9	237.0	184.0-328.0	1.0	1.0
$\alpha I^{-/-}$	922.1	756.1-1066.0	12856.0	9343.9-19626.0	32.7	54.2
$\alpha 2^{-/-}$	484.4	266.6-985.3	13397.0	3076.9-71026.0	17.2	56.5
$\alpha 3^{-/-}$	36.0	24.7-47.1	448.5	278.0-1005.9	1.3	1.9
$\alpha 4^{T227M}$	18.2	13.8-22.8	73.4	51.4-141.1	0.6	0.3
$\alpha 5^{-/-}$	18.0	1.0-15.2	43.3	34.2-63.5	0.6	0.2
$\alpha 6^{-/-}$	28.7	24.3-34.0	142.5	105.2-219.3	1.0	0.6
$\alpha 7^{-/-}$	30.0	24.5-37.0	238.1	157.0-440.0	1.1	1.0
βI^{R81T}	4349.2	3096.0-6661.3	33601	17279.0-117310.0	154.0	141.8
$\beta 2^{-/-}$	668.9	431.0-913.5	5797.4	3728.1-12256.0	23.7	24.5
$\beta 3^{-/-}$	28.2	24.7-31.9	237.0	184.0-327.9	1.5	0.7

Table S11 Log dose probit mortality data and resistance ratios for spinetora

Strain	LC ₅₀ (mg/L)	95% CL	LC ₉₅ (mg/L)	95% CL	Resistance ratio	
					LC ₅₀	LC ₉₅
Control	1.3	1.2-1.4	2.5	2.2-3.1	1.0	1.0
$\alpha 1^{-/-}$	1.8	1.2-2.6	4.0	2.7-13.5	1.4	1.6
$\alpha 2^{-/-}$	0.8	0.7-0.9	1.8	1.5-2.3	0.6	0.7
$\alpha 3^{-/-}$	1.2	1.0-1.3	2.6	2.2-3.5	0.9	1.0
$\alpha 4^{T227M}$	1.8	1.3-2.5	4.4	3.0-11.1	1.4	1.8
$\alpha 5^{-/-}$	1.4	1.2-1.6	4.0	3.2-5.6	1.1	1.6
$\alpha 6^{-/-}$	55.6	45.8-68.1	134.3	101.8-218.1	42.8	53.7
$\alpha 6^{+/-}$	1.6	1.4-1.8	3.4	2.9-4.5	1.2	1.4
$\alpha 7^{-/-}$	1.8	1.6-2.2	4.8	3.8-6.7	1.4	1.9
$\beta 1^{R81T}$	0.8	0.7-0.9	2.1	1.7-2.8	0.6	0.8
$\beta 2^{-/-}$	0.8	0.7-0.9	1.3	1.1-1.7	0.6	0.5
$\beta 3^{-/-}$	1.4	1.2-1.6	2.9	2.4-3.8	1.1	1.2
$\alpha 5^{-/-}; \alpha 7^{-/-}$	2.2	1.8-2.5	4.5	3.6-6.6	1.7	1.8

Table S12 Primers used in qPCR analysis

Primer	Forward Sequences (5'-3')	Reverse Sequences (5'-3')
Ribosomal protein L32	GACGCTTCAAGGGACAGTATCTG	AAACGCGGTTCTGCATGAG
nAChR α1	TACGTTCGAGAAGCCCTACG	GGGAGCCTGCAGGATAATCA
nAChR α2	GGCCGCACGCAAAAAGTATC	CAAATGTGCCCACCAAGGATG
nAChR α3	CTGTCCGGAAGTCCACAAGG	GTCGGAGCCTGCAGGATAAT
nAChR α4	CAGACGAAATAGCCGCCGTC	ACCACAACTGCCAACGTGA
nAChR α5	AGACAATGCCGGCTACTTCC	CAGCCAGCACAAAAACACGA
nAChR α6	GTGGAACGACTACAATCTGCG	AAGATACCAGGGGGCACGTA
nAChR α7	CCAATGTGCTCGATATAGACGATG	CTGTTATCCAACGCAGCTCCT
nAChR β1	GCAAATCCTGGCTGTTGTGC	ACGGTTAGCAGAGTTAACAGAGTT
nAChR β2	GCGTGACAGCATCAGCG	AGAGCCAGAGAAAGAAGCGG
nAChR β3	GGCCTGTTCACGAACTACGA	CAGACCTCGCTGGACTTCAA