

Colonial choanoflagellate isolated from Mono Lake harbors a microbiome

Hake, K. H.^{1, 2}, West, P.T.³, McDonald, K.⁴, Laundon, D.⁵, Feng, C.¹, Burkhardt, P.^{5, 6},
Richter, D.J.⁷, Banfield, J.F.³, and King, N.^{1,*}

5

¹Howard Hughes Medical Institute and Department of Molecular and Cell Biology,
University of California, Berkeley, CA, USA

10

² Present address: Calico Life Sciences, South San Francisco, CA, USA
³ Department of Environmental Science, Policy, & Management, University of California,
Berkeley, CA, USA

15

⁴ Electron Microscopy Laboratory, University of California, Berkeley, CA, USA

15

⁵ Marine Biological Association of the United Kingdom, Plymouth, United Kingdom

20

⁶ Sars International Centre for Molecular Marine Biology, University of Bergen, Bergen,
Norway

20

⁷ Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Barcelona, Spain

25

*Send correspondence to nking@berkeley.edu

ABSTRACT

30 Choanoflagellates offer key insights into bacterial influences on the origin and early evolution of animals. Here we report the isolation and characterization of a new colonial choanoflagellate species, *Barrosiga monosierra*, that, unlike previously characterized species, harbors a stable microbiome. *B. monosierra* was isolated from Mono Lake, California and forms large spherical colonies that are more than an order of
35 magnitude larger than those formed by the closely related *Salpingoeca rosetta*. By designing fluorescence *in situ* hybridization probes from metagenomic sequences, we found that *B. monosierra* colonies are colonized by members of the halotolerant and closely related *Saccharospirillaceae* and *Oceanospirillaceae*, as well as purple sulfur bacteria (*Ectothiorhodospiraceae*) and non-sulfur *Rhodobacteraceae*. This relatively
40 simple microbiome in a close relative of animals presents a new experimental model for investigating the evolution of stable interactions among eukaryotes and bacteria.

IMPORTANCE

45 The animals and bacteria of Mono Lake (California) have evolved diverse strategies for surviving the hypersaline, alkaline, arsenic-rich environment. We sought to investigate whether the closest living relatives of animals, the choanoflagellates, exist among the relatively limited diversity of organisms in Mono Lake. We repeatedly isolated members of a single species of choanoflagellate, which we have named *Barrosiga monosierra*, suggesting that it is a stable and abundant part of the ecosystem.
50 Characterization of *B. monosierra* revealed that it forms large spherical colonies that each contain a microbiome, providing an opportunity to investigate the evolution of stable physical associations between eukaryotes and bacteria.

OBSERVATION

A newly identified choanoflagellate species forms large colonies that contain a microbiome

60 Choanoflagellates are the closest living relatives of animals and, as such, provide insights into the origin of key features of animals, including animal multicellularity and cell biology (1, 2). Over a series of four sampling trips to Mono Lake, California (Fig. 1A; Table S1) we collected single-celled choanoflagellates and large spherical
65 choanoflagellate colonies, many of which were hollow (Fig. 1B) and resembled the blastula stage of animal development. In colonies and single cells, each cell had the typical collar complex observed in other choanoflagellates: an apical flagellum surrounded by a collar of microvilli (1, 2). In these “rosette” colonies, the cells were
70 oriented with the basal pole of each cell pointing inwards and the apical flagellum facing out (Fig. 1). To study the Mono Lake choanoflagellates in greater detail, we established clonal strains from ten independent isolates, two of which were each started from a single-celled choanoflagellate and the remaining eight of which were each started from a single colony (Table S1). The two strains started from single-celled choanoflagellates, isolates ML1.1 and ML1.2, took on the colonial morphology observed in the other
75 isolates after culturing in the laboratory, suggesting that the colonies and single cells isolated from Mono Lake could belong to the same species. We are aware of no prior reports of choanoflagellates having been cultured from any alkaline soda lake, including Mono Lake.

80 The 18S rDNA genes for six of the Mono Lake isolates were sequenced and found to be >99% identical (Table S1). Phylogenetic analysis confirmed that all of the isolates are members of a single choanoflagellate species (Fig. S1). In further phylogenetic analyses based on 18S rDNA and two protein-coding genes from isolate ML2.1 (Fig. 1C) (3), we found that its closest relatives are the emerging model
85 choanoflagellate *S. rosetta* (4–8) and *Microstomoeca roanoka* (3, 10). The phylogenetic distance separating the Mono Lake species from its closest relatives is similar to the distance separating other choanoflagellate genera. Therefore, we propose the name *Barrosiga monosierra*, with the genus name inspired by esteemed choanoflagellate researcher Prof. Barry Leadbeater and the species name inspired by the location of
90 Mono Lake in the Sierra Nevada mountain range. (See Supplemental Methods for further details and a formal species description.)

95 Although *B. monosierra* and *S. rosetta* form rosette-shaped spherical colonies, they differ greatly in size. *S. rosetta* colonies range from 10-30 µm in diameter while *B. monosierra* forms among the largest choanoflagellate colonies observed (1, 11), with a single culture exhibiting colony sizes spanning from 10-120 µm in diameter (Fig. 1D-F). Unlike the rosettes of *S. rosetta*, in which the basal poles of cells are closely apposed in the rosette center (5, 7), cells in large *B. monosierra* rosettes form a shell on the surface of a seemingly hollow sphere. Inside the ostensibly hollow sphere, a branched network
100 of extracellular matrix connects the basal poles of all cells (Fig. S2.)

Upon staining *B. monosierra* with the DNA dye Hoechst 33342, we observed the expected toroidal nuclei in each choanoflagellate cell (12), but were surprised to detect Hoechst-positive material in the interior of *B. monosierra* colonies (Fig. 2A, A').

105 Transmission electron microscopy revealed the presence of 1 μ m and smaller cells with diverse morphologies bounded by cell walls in the centers of rosettes (Fig. 2B, B'; Fig. S3). Together, these observations led us to hypothesize that the centers of *B. monosierra* colonies contain bacteria.

110 By performing hybridization chain reaction fluorescence *in situ* hybridization (HCR-FISH (13–15)) with a broad-spectrum probe of bacterial 16S rRNA, EUB338 (16), we confirmed that the cells in the center of colonies are bacteria (Fig. 2C). A second probe that specifically targeted 16S rRNA sequences from Gammaproteobacteria, GAM42a, revealed that the majority of the bacteria inside the colonies are 115 Gammaproteobacteria (Fig. 2C')(17). Finally, by incubating *B. monosierra* cultures with fluorescently labeled D-amino acids, which are specifically incorporated into the cell walls of growing bacteria, we found that the bacteria in *B. monosierra* colonies are alive and growing (Fig. S4)(18). Therefore, *B. monosierra* contains a microbiome (19).

120 To visualize the spatial distribution of choanoflagellate and bacterial cells in a representative colony, we generated a 3D reconstruction from serial sections imaged by TEM. The colony contained 70 choanoflagellate cells that were tightly packed, forming a largely continuous monolayer of cells (Fig. 2D). As observed by immunofluorescence microscopy (Figs. 2A and 2A'), all cells were highly polarized and oriented with their 125 apical flagella and collars extending away from the centroid of the rosette. Many cells were connected by fine intercellular bridges (Fig. S5) that have been previously observed in other colonial choanoflagellates, including *S. rosetta* (5, 20).

130 The 3D reconstruction also revealed at least 200 bacterial cells in the center of the rosette (Fig. 2D', 2D''), some of which were physically associated with and wrapped around the choanoflagellate ECM (Fig. S6). A small number of bacterial cells were observed between the lateral surfaces of choanoflagellate cells, although it was not possible to determine whether they were entering or exiting the colony (Fig. 2D''; Fig. 135 S7). Colonies failed to incorporate bacteria-sized bovine serum albumin (BSA)-coated latex microspheres (0.2 μ m and 1 μ m) into their centers, suggesting that environmental bacteria may not be capable of passively accessing the centers of *B. monosierra* colonies (Fig. S8).

Gammaproteobacteria and Alphaproteobacteria in the *B. monosierra* microbiome

140 We next sought to identify which bacteria comprise the microbiomes of *B. monosierra*. To identify candidate bacteria for which to design FISH probes, we first sequenced and assembled metagenomes and 16S rDNA sequences from choanoflagellate-enriched samples and from environmental bacteria-enriched samples. 145 These samples were derived from two co-cultures of *B. monosierra* with Mono Lake bacteria, ML2.1E and ML2.1G (Fig. S9), with the enrichment for choanoflagellates or bacteria performed by centrifugation. A total of 24 different bacterial species were

150 identified via two complementary bioinformatic approaches (EukRep Metagenomic Analysis and EMIRGE 16S rRNA Analysis; Table S2), of which 22 species were present in fractions enriched with *B. monosierra* colonies (Table S3). The phylogenetic relationships among these and other bacterial species were determined based on analysis of highly conserved ribosomal proteins and 16S rDNA sequences (Fig. 2E and S10).

155 The 22 bacterial species detected in cultures with *B. monosierra* may have co-sedimented with the *B. monosierra* colonies due their community-structure densities (e.g. biofilms), a transient association with the choanoflagellate colonies (e.g. as prey), or through a stable association with the choanoflagellate colonies. Upon investigation by FISH microscopy, we detected ten or eleven of these species in the centers of *B. monosierra* colonies (Table S4, Fig. S11). (The uncertainty regarding the precise number of choanoflagellate-associated bacterial species stems from the inability to disambiguate 16S rDNA sequences corresponding to one or two of the species.) Of these microbiome bacteria, nine were Gammaproteobacteria from the families *Oceanospirillaceae* (Fig. S11A; OceaML1, OceaML2, OceaML3, OceaML4, OceaML4), 160 *Ectothiorhodospiraceae* (Fig. S11B; EctoML1, EctoML2, EctoML3, EctoML4), and *Saccharospirillaceae* (Fig. S11C; SaccML), matching our original observation that the majority of the bacteria were Gammaproteobacteria (Fig. 2C, C'). The remaining species was a *Roseinatronobacter* sp. (RoseML; Alphaproteobacteria) (Fig. S11D). The microbiome bacteria exhibited an array of morphologies, from long and filamentous to 170 rod shaped (Fig. S3 and Fig. S12). Intriguingly, with the exception of OceaML3, which was exclusively detected inside *B. monosierra* colonies of ML2.1E (Fig. S13), all other microbiome species identified in this study were detected both inside and outside the colonies.

175 In animals, the microbiome often contains a core set of host-adapted bacteria that are present in many or all individuals in a host species, as well as a more flexible set of bacteria that may be found in only a subset of individuals (21–23). To identify core members of the *B. monosierra* microbiome, we measured the frequency with which colonies of ML2.1EC contained or lacked a number of representative microbiome 180 bacteria (Fig. S14A) and estimated the abundance of each species relative to the total microbiome (Fig. S14B). Only one bacterium tested, OceaML1, was found in the microbiome of all *B. monosierra* colonies (Fig. S14A). The other most frequently observed members of the microbiome were SaccML (93.3% of colonies), EctoML3 (91.8% of colonies) and EctoML1 (82.4% of colonies; Fig. S14A). Two other 185 Gammaproteobacterial species, OceaML2 and EctoML2, were found in ~50 - 60% of colonies, while the Alphaproteobacterium RoseML was found in only 13.9% of rosettes.

190 The most common resident of the *B. monosierra* microbiome, OceaML1, was also the most abundant, representing on average 66.4% of the total bacterial load per colony (Fig. S14B). Other abundant bacteria, some found in >80% of colonies, represented approximately smaller percentages of the average bacterial biomass in the microbiomes in which they were found. For example, SaccML was found in 93.3% of microbiomes but represented only 30.3% of total bacteria in the *B. monosierra* rosettes

195 in which it was found. EctoML1, which was found in 82.4% of *B. monosierra* rosettes represented less than 10% of the bacteria in the bacteria in which was detected. Thus, only OceaML1 appears to be a core member of the *B. monosierra* microbiome. Other symbionts detected in *B. monosierra* (e.g. SaccML and OceaML2-4) are close relatives of OceaML1 and may engage in similar metabolic or developmental functions in their interactions with *B. monosierra* (24).

200

Discussion

205 Interactions with bacteria are essential to choanoflagellate nutrition and life history. Bacteria are the primary food source for choanoflagellates, and the choanoflagellate *S. rosetta* responds to different secreted bacterial cues to undergo either multicellular developmental or mating (25–27). We report the isolation and characterization of a new choanoflagellate species, *B. monosierra*, that forms large colonies and contains a microbiome consisting of at least ten different bacterial 210 symbionts. To our knowledge, this is the first example of a stable interaction between choanoflagellates and ectosymbiotic bacteria. Future studies will be important to determine how the *B. monosierra* colonies and their bacterial symbionts interact.

215 We detected 10 – 11 bacterial species from four different families (*Saccharospirillaceae*, *Oceanospirillaceae*, *Ectothiorhodospiraceae*, and *Rhodobacteraceae*) in the *B. monosierra* microbiome. Comparisons with other 220 symbioses suggest that their interactions with *B. monosierra* may relate to metabolism and detoxification of environmental sulfur. Such functions have been reported for *Oceanospirillaceae* species that are symbionts of diverse marine animals, from corals to snails (28–35) and for *Ectothiorhodospiraceae* that are symbionts of diverse ciliates and animals (36–39).

225 Although the composition of the animal gut microbiome often varies between individuals in a species, many host species harbor a core set of microbes, with which they participate in stable metabolic interactions and may coevolve (22, 23). Indeed, choanoflagellates express homologs of Toll-like receptors that, in animals, mediate interactions with gut bacteria to maintain homeostasis (10). In *B. monosierra*, OceaML1 230 was found in 100% of rosettes assessed and was the most abundant bacterium in the microbiome, suggesting that its interactions with *B. monosierra* may be essential for *B. monosierra* biology. Interestingly, the related bacterium *Endozooicomonas* sp. (Order Oceanospirillales), is a core member of the gut microbiome of the ascidian *Ciona intestinalis*, where it represents up to 54% of the bacterial biomass (31).

235 *B. monosierra* and its associated microbiome provide a unique opportunity to characterize a new symbiotic interaction between a single choanoflagellate species and a microbial community. Due to the phylogenetic relevance of choanoflagellates, this symbiotic relationship has the potential to illuminate the ancestry of and mechanisms underlying stable associations between animals and bacteria, colonization of organisms

240 by diverse microbial communities, and insights into one of the most complex animal-bacterial interactions, the animal gut microbiome.

245 **Data Availability:** Genbank accession numbers for bacterial 16S rDNA sequences are listed in Supplementary Data File 1. Sequences for *B. monosierra* 18S rDNA, EFL, and Hsp90 (Fig. 1C) can be found in Supplementary Data File 2, and have been assigned GenBank accession numbers MW838180, MW979373 and MW979374, respectively. 18S sequences for different *B. monosierra* strains (Fig. S1, Table S1) have been assigned GenBank accession numbers MZ015010-MZ015015. The assembled *B. monosierra* genome sequence is currently being uploaded to GenBank and an accession number will be available prior to publication. All relevant input and output data from the phylogenetic trees presented in Fig. 1C and Fig. S1 are available via FigShare (<https://doi.org/10.6084/m9.figshare.14474214>).

250
255 **Author contributions:** *P.W. performed metagenomic analysis, K.M. did the transmission electron microscopy, D.L. and P.B. did the TEM-reconstruction, and C.F. helped culture *B. monosierra*. K.H. performed all other experiments and analysis. D.J.R. originally isolated *B. monosierra* and contributed to manuscript editing. J.B. and N.K. contributed to project leadership, experimental design, figure design, writing, and editing.*

260

Acknowledgements

265 We thank Tarja Hoffmeyer for help with establishing some of the original *B. monosierra* cultures. We thank Emil Ruff and Julia Schwartzman for assistance with early CARD-FISH experiments through the Physiology Course and Microbial Diversity Course at the Marine Biological Laboratory in Woods Hole. We thank Michael VanNieuwenhze for the fluorescent D-amino acids. Frank Nitsche provided advice on the formulation of an artificial Mono Lake culture medium. We thank the following for helpful discussions and 270 research support: Reef Aldayafleh, Cédric Berney, David Booth, Ben Larson, Monika Sigg, Laura Wetzel, and Arielle Woznica. We thank Alain Garcia De Las Bayanos and Karen Carniol for valuable feedback on the manuscript. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. DGE 1106400 and DGE 1752814. D.J.R. received the support of a 275 fellowship from "la Caixa" Foundation (ID 100010434) with the fellowship code LCF/BQ/PI19/11690008.

Supplement Contents

280

Text S1: Supplemental Materials and Methods

Figure S1: Independent choanoflagellate isolates from Mono Lake are the same species

285

Figure S2: Six-gene phylogenetic tree of *B. monosierra* and other choanoflagellates.

Figure S3. *B. monosierra* colonies contain an extracellular matrix (ECM).

Figure S4: Bacterial symbionts in *B. monosierra* colonies exhibit a range of morphologies.

Figure S5: Bacteria inside *B. monosierra* colonies are alive and growing.

290

Figure S6: Intercellular bridges connect cells in *B. monosierra* colonies.

Figure S7: Bacterial symbionts physically associate with and wrap around the choanoflagellate ECM.

Figure S8: Bacteria are found wedged between the lateral surfaces of *B. monosierra* cells.

295

Figure S9: *B. monosierra* colonies cannot be passively penetrated by sub-micron particles.

Figure S10: Schematic describing the establishment of *B. monosierra* strains.

Figure S11: Phylogenetic analysis of 16S rRNA tree of EMIRGE sequences.

Figure S12: Diverse bacteria are found in the microbiome of *B. monosierra*.

300

Figure S13: *B. monosierra* symbionts exhibit filamentous and rod morphologies.

Figure S14: The symbiont OceaML3 is present exclusively inside rosettes.

Figure S15: OceaML1 is a core member of the *B. monosierra* microbiome.

305

Table S1: Date, location, phenotype, and isolate designations for *Barrosiga monosierra* isolates.

Table S2: Nine genera of bacteria identified in *B. monosierra* cultures through two independent analyses: comparison of ribosomal proteins detected through metagenomic assembly and by 16s rRNA assembly and analysis.

Table S3: Predicted targets of HCR-FISH probes based on 16S rRNA sequences.

310

Bacteria in bold were identified inside *B. monosierra* colonies.

Table S4: Full length probes, with spacer and initiator sequences, used for HCR-FISH.

Table S5: Growth Media Recipes.




Figure 1

Figure 1. A new colony-forming choanoflagellate isolated from Mono Lake.

(A) Choanoflagellates were collected from two sampling sites (asterisks) near the shore of Mono Lake, California. (Modified from map at monolake.org.) (B) *B. monosierra* forms large colonies (DIC image). Scale bar = 50 μ m. (C) *B. monosierra* (shown in bold) is a craspedid choanoflagellate closely related to *S. rosetta* and *Microstomoeca roanoka*. Phylogeny based on sequences of 3 genes: 18S rDNA, EFL and HSP90. Metazoa (7 species) were collapsed to save space. Bayesian posterior probabilities are indicated above each internal branch, and maximum likelihood bootstrap values below. (A '-' value indicates a bifurcation lacking support or not present in one of the two reconstructions.) (D-E) Two representative colonies reveal the extremes of the *B. monosierra* colony size range (D, 58 μ m diameter; E, 19 μ m diameter; scale bar = 20 μ m). In *B. monosierra* colonies, each cell is oriented with its apical flagellum (white; labeled with anti-tubulin antibody) and the apical collar of microvilli (red; stained with phalloidin) pointing out. Nuclei (cyan) were visualized with the DNA-stain Hoechst 33342. (F) Colonies of *B. monosierra* span from 10 μ m in diameter, a size comparable to that of small *S. rosetta* colonies, to 120 μ m, over an order of magnitude larger. Diameters of *B. monosierra* and *S. rosetta* colonies were plotted as a violin plot; median indicated as thick black line. Diameters of representative colonies indicated as colored bars behind violin plot (D, red bar; E, blue bar).




Figure 2

345 **Figure 2. *B. monosierra* colonies are filled with bacteria.**

(A, A') The center of a representative *B. monosierra* colony, shown as a maximum intensity projection (A) and optical z-section (A'), contains DNA (revealed by Hoechst 33342 staining; cyan). Apical flagella were labeled with anti-tubulin antibody (white); microvilli were stained with phalloidin (red). Hoechst 33342 staining (cyan) revealed the 350 toroidal choanoflagellate nuclei along the colony perimeter and an amorphous cloud of DNA sitting within the central cavity formed by the monolayer of choanoflagellate cells. (B-B') Thin section through a representative *B. monosierra* colony, imaged by transmission electron microscopy (TEM), revealed the presence of small cells in the 355 central cavity. (B') Inset (box; panel B') reveals that the interior cells are each surrounded by a cell wall. (C-C'') The small cells inside *B. monosierra* colonies are bacteria, as revealed by hybridization with a broad spectrum 16S rRNA probe (C, green) and a probe targeting Gammaproteobacteria (C', red). Choanoflagellate nuclei and bacterial nucleoids were revealed by staining with Hoechst (C'', cyan). (C'') Merge of panels C – C''. Scale bar for all = 5 μ m. (D-D'') 3D reconstruction of a 70-cell *B. 360 monosierra* choanoflagellate colony from transmission electron micrographs of serial ultrathin sections revealed that the bacteria are closely associated with and wrapped around the ECM inside the colony. (D) Whole colony view. (D') Cut-away view of colony center. Color code: cell bodies (cyan); microvilli (orange); flagella (green); bacteria (red); 365 ECM (white); intercellular bridges (yellow, see also Fig. S5); filopodia (purple). (D'') Reducing the opacity of the choanoflagellate cell renderings revealed the presence of bacteria positioned between the lateral surfaces of choanoflagellate cells (brackets, see also Fig. S7). (E) Unrooted phylogenetic tree based on 16 concatenated ribosomal 370 protein sequences representing bacterial diversity modified from (40), illustrated to indicate the phylogenetic placement of bacteria co-cultured from Mono Lake with *B. monosierra*. The bacteria belonged to four major classes: Spirochaetia, Alphaproteobacteria, Gammaproteobacteria, and Bacteroidetes, however the bacteria found associated with *B. monosierra* colonies came only from Alphaproteobacteria and Gammaproteobacteria. Circles represent the phylogenetic placement of non-symbionts (white) and symbionts (*Oceanospirillaceae* sp., magenta; *Saccharospirillaceae* sp., green; *Ectothiorhodospiraceae* sp., blue; *Roseinatronobacter* sp., orange). See also 375 Figs. S10 and S11.

References

1. Leadbeater BSC. 2015. *The Choanoflagellates: Evolution, Ecology, and Biology*. Cambridge University Press, Cambridge, UK.
2. Brunet T, King N. 2017. The origin of animal multicellularity and cell differentiation. *Dev Cell* 43:124–140.
3. Carr M, Richter DJ, Fozouni P, Smith TJ, Jeuck A, Leadbeater BSC, Nitsche F. 2017. A six-gene phylogeny provides new insights into choanoflagellate evolution. *Mol Phylogenet Evol* 107:166–178.
4. Booth DS, Szmidt-Middleton H, King N. 2018. Transfection of choanoflagellates illuminates their cell biology and the ancestry of animal septins. *Mol Biol Cell* <https://doi.org/10.1091/mbc.e18-08-0514>.
5. Dayel MJ, Alegado RA, Fairclough SR, Levin TC, Nichols SA, McDonald K, King N. 2011. Cell differentiation and morphogenesis in the colony-forming choanoflagellate *Salpingoeca rosetta*. *Dev Biol* 357:73–82.
6. Fairclough SR, Chen Z, Kramer E, Zeng Q, Young S, Robertson HM, Begovic E, Richter DJ, Russ C, Westbrook MJ, others. 2013. Premetazoan genome evolution and the regulation of cell differentiation in the choanoflagellate *Salpingoeca rosetta*. *Genome Biol* 14:R15.
7. Levin TC, Greaney AJ, Wetzel L, King N, Sánchez Alvarado A. 2014. The rosetteless gene controls development in the choanoflagellate *S. rosetta*. *eLife* 3.
8. Wetzel LA, Levin TC, Hulett RE, Chan D, King GA, Aldayafleh R, Booth DS, Sigg MA, King N. 2018. Predicted glycosyltransferases promote development and prevent spurious cell clumping in the choanoflagellate *S. rosetta*. *eLife* <https://doi.org/10.7554/eLife.41482>.
9. Schiwitza S, Arndt H, Nitsche F. 2018. Four new choanoflagellate species from extreme saline environments: Indication for isolation-driven speciation exemplified by highly adapted Craspedida from salt flats in the Atacama Desert (Northern Chile). *Eur J Protistol* 66:86–96.
10. Richter DJ, Fozouni P, Eisen M, King N. 2018. Gene family innovation, conservation and loss on the animal stem lineage. *eLife* 7:e34226.
11. Larson BT, Ruiz-Herrero T, Lee S, Kumar S, Mahadevan L, King N. 2019. Biophysical principles of choanoflagellate self-organization. *bioRxiv* 659698.
12. Burkhardt P, Grønborg M, McDonald K, Sulur T, Wang Q, King N. 2014. Evolutionary insights into premetazoan functions of the neuronal protein Homer. *Mol Biol Evol* msu178.

13. Choi HMT, Calvert CR, Husain N, Huss D, Barsi JC, Deverman BE, Hunter RC, Kato M, Lee SM, Abelin ACT, Rosenthal AZ, Akbari OS, Li Y, Hay BA, Sternberg PW, Patterson PH, Davidson EH, Mazmanian SK, Prober DA, van de Rijn M, Leadbetter JR, Newman DK, Readhead C, Bronner ME, Wold B, Lansford R, Sauka-Spengler T, Fraser SE, Pierce NA. 2016. Mapping a multiplexed zoo of mRNA expression. *Development* 143:3632–3637.
14. DePas WH, Starwalt-Lee R, Van Sambeek L, Kumar SR, Gradinaru V, Newman DK. 2016. Exposing the three-dimensional biogeography and metabolic states of pathogens in cystic fibrosis sputum via hydrogel embedding, clearing, and rRNA labeling. *MBio* 7:e00796---16.
15. Pernthaler A, Pernthaler J, Amann R. 2002. Fluorescence in situ hybridization and catalyzed reporter deposition for the identification of marine bacteria. *Appl Environ Microbiol* 68:3094–3101.
16. Amann RI, Binder BJ, Olson RJ, Chisholm SW, Devereux R, Stahl DA. 1990. Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations. *Appl Environ Microbiol* 56:1919–1925.
17. Manz W, Amann R, Ludwig W, Wagner M, Schleifer KH. 1992. Phylogenetic Oligodeoxynucleotide Probes for the Major Subclasses of Proteobacteria: Problems and Solutions. *Syst Appl Microbiol* 15:593–600.
18. Kuru E, Hughes HV, Brown PJ, Hall E, Tekkam S, Cava F, De Pedro MA, Brun V, Vannieuwenhze MS. 2012. In situ probing of newly synthesized peptidoglycan in live bacteria with fluorescent D-amino acids. *Angew Chemie - Int Ed* 51:12519–12523.
19. WHIPPS JM. 1991. Effects of Mycoparasites on Sclerotia-Forming Fungi, p. 129–140. *In* BEEMSTER, ABR, BOLLEN, GJ, GERLAGH, M, RUISEN, MA, SCHIPPERS, B, TEMPEL, A (eds.), *Biotic Interactions and Soil-Borne Diseases*. Elsevier.
20. Laundon D, Larson BT, McDonald K, King N, Burkhardt P. 2019. The architecture of cell differentiation in choanoflagellates and sponge choanocytes. *PLOS Biol* 17:e3000226.
21. Shapira M. 2016. Gut Microbiotas and Host Evolution: Scaling Up Symbiosis. *Trends Ecol Evol* 31:539–549.
22. Roeselers G, Mittge EK, Stephens WZ, Parichy DM, Cavanaugh CM, Guillemain K, Rawls JF. 2011. Evidence for a core gut microbiota in the zebrafish. *ISME J* 5:1595–1608.
23. Huse SM, Ye Y, Zhou Y, Fodor AA. 2012. A core human microbiome as viewed through 16S rRNA sequence clusters. *PLoS One* 7.

24. Doolittle WF, Booth A. 2017. It's the song, not the singer: an exploration of holobiosis and evolutionary theory. *Biol Philos* 32:5–24.
25. Alegado RA, Brown LW, Cao S, Dermenjian RK, Zuzow R, Fairclough SR, Clardy J, King N. 2012. A bacterial sulfonolipid triggers multicellular development in the closest living relatives of animals. *Elife* 1:e00013.
26. Woznica A, Cantley AM, Beemelmanns C, Freinkman E, Clardy J, King N. 2016. Bacterial lipids activate, synergize, and inhibit a developmental switch in choanoflagellates. *Proc Natl Acad Sci* 113:7894–7899.
27. Woznica A, Gerdt JP, Hulett RE, Clardy J, King N. 2017. Mating in the closest living relatives of animals is induced by a bacterial chondroitinase. *Cell* 170:1175–1183.
28. Bayer T, Neave MJ, Alsheikh-Hussain A, Aranda M, Yum LK, Mincer T, Hughen K, Apprill A, Voolstra CR. 2013. The microbiome of the red sea coral *Stylophora pistillata* is dominated by tissue-associated *Endozoicomonas* bacteria. *Appl Environ Microbiol* 79:4759–4762.
29. Beinart RA, Nyholm S V, Dubilier N, Girguis PR. 2014. Intracellular Oceanospirillales inhabit the gills of the hydrothermal vent snail *Alviniconcha* with chemosynthetic, γ -Proteobacterial symbionts. *Environ Microbiol Rep* 6:656–664.
30. Carlos C, Torres TT, Ottoboni LMM. 2013. Bacterial communities and species-specific associations with the mucus of Brazilian coral species. *Sci Rep* 3.
31. Dishaw LJ, Flores-Torres J, Lax S, Gemayel K, Leigh B, Melillo D, Mueller MG, Natale L, Zucchetti I, De Santis R, Pinto MR, Litman GW, Gilbert JA. 2014. The gut of geographically disparate *Ciona intestinalis* harbors a core microbiota. *PLoS One* 9:e93386.
32. Goffredi SK, Orphan VJ, Rouse GW, Jahnke L, Embaye T, Turk K, Lee R, Vrijenhoek RC. 2005. Evolutionary innovation: A bone-eating marine symbiosis. *Environ Microbiol* 7:1369–1378.
33. Kurahashi M, Yokota A. 2007. *Endozoicomonas elysicola* gen. nov., sp. nov., a γ -proteobacterium isolated from the sea slug *Elysia ornata*. *Syst Appl Microbiol* 30:202–206.
34. Morrow KM, Moss AG, Chadwick NE, Liles MR. 2012. Bacterial Associates of Two Caribbean Coral Species Reveal Species-Specific Distribution and Geographic Variability. *Appl Environ Microbiol* 78:6438–6449.
35. Vezzulli L, Pezzati E, Huete-Stauffer C, Pruzzo C, Cerrano C. 2013. 16SrDNA Pyrosequencing of the Mediterranean Gorgonian *Paramuricea clavata* Reveals a Link among Alterations in Bacterial Holobiont Members, Anthropogenic Influence

and Disease Outbreaks. PLoS One 8:e67745.

36. Bayer C, Heindl NR, Rinke C, Lücker S, Ott JA, Bulgheresi S. 2009. Molecular characterization of the symbionts associated with marine nematodes of the genus *Robbea*. Environ Microbiol Rep 1:136–144.
37. Payne SH, Loomis WF. 2006. Retention and loss of amino acid biosynthetic pathways based on analysis of whole-genome sequences. Eukaryot Cell 5:272–276.
38. Rinke C, Schmitz-Esser S, Stoecker K, Nussbaumer AD, Molnar DA, Vanura K, Wagner M, Horn M, Ott JA, Bright M. 2006. “*Candidatus Thiobios zoothamnicoli*”, an ectosymbiotic bacterium covering the giant marine ciliate *Zoothamnium niveum*. Appl Environ Microbiol 72:2014–2021.
39. Tian RM, Wang Y, Bougouffa S, Gao ZM, Cai L, Bajic V, Qian PY. 2014. Genomic analysis reveals versatile heterotrophic capacity of a potentially symbiotic sulfur-oxidizing bacterium in sponge. Environ Microbiol 16:3548–3561.
40. Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN, Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman DA, Finstad KM, Amundson R, Thomas BC, Banfield JF. 2016. A new view of the tree of life. Nat Microbiol 1:nmicrobiol201648.