

Synergistic block of SARS-CoV-2 infection by combined drug inhibition of the host entry factors PIKfyve kinase and TMPRSS2 protease

Alex J.B. Kreutzberger^{a,b}, Anwesha Sanyal^{a,b}, Ravi Ojha^c, Jesse D. Pyle^d, Olli Vapalahti^{c,e,f},
Giuseppe Balistreri^{c,e} and Tom Kirchhausen^{a,b,g,#}

^a Department of Cell Biology, Harvard Medical School, 200 Longwood Av, Boston, MA 02115, USA

^b Program in Cellular and Molecular Medicine, Boston Children's Hospital, 200 Longwood Av, Boston, MA 02115, USA

^c Department of Virology, Faculty of Medicine, University of Helsinki, Helsinki, Finland

^d Program in Virology, Harvard Medical School, 200 Longwood Ave, Boston, MA 02115, USA

^e Department of Veterinary Biosciences, University of Helsinki, Helsinki, Finland

^f Virology and Immunology, Helsinki University Hospital Diagnostic Center (HUSLAB), Helsinki, Finland

^g Department of Pediatrics, Harvard Medical School, 200 Longwood Ave, Boston, MA 02115, USA

Running Head: Synergistic block of SARS-CoV-2 infection

#Address correspondence to Tom Kirchhausen kirchhausen@crystal.harvard.edu

ABSTRACT

Repurposing FDA-approved inhibitors able to prevent infection by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) could provide a rapid path to establish new therapeutic options to mitigate the effects of coronavirus disease 2019 (COVID-19). Proteolytic cleavages of the spike S protein of SARS-CoV-2, mediated by the host cell proteases cathepsin and TMPRSS2, alone or in combination, are key early activation steps required for efficient infection. The PIKfyve kinase inhibitor apilimod interferes with late endosomal viral traffic, and through an ill-defined mechanism prevents *in vitro* infection through late endosomes mediated by cathepsin. Similarly, inhibition of TMPRSS2 protease activity by camostat mesylate or nafamostat mesylate prevents infection mediated by the TMPRSS2-dependent and cathepsin-independent pathway. Here, we combined the use of apilimod with camostat mesylate or nafamostat mesylate and found an unexpected ~5-10-fold increase in their effectiveness to prevent SARS-CoV-2 infection in different cell types. Comparable synergism was observed using both, a chimeric vesicular stomatitis virus (VSV) containing S of SARS-CoV-2 (VSV-SARS-CoV-2) and SARS-CoV-2 virus. The substantial ~5-fold or more decrease of half maximal effective concentrations (EC₅₀ values) suggests a plausible treatment strategy based on the combined use of these inhibitors.

IMPORTANCE

Infection by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is causing the coronavirus disease 2019 (COVID-2019) global pandemic. There are ongoing efforts to uncover effective antiviral agents that could mitigate the severity of the

47 disease by controlling the ensuing viral replication. Promising candidates include small
48 molecules that inhibit the enzymatic activities of host proteins, thus preventing SARS-
49 CoV-2 entry and infection. They include Apilimod, an inhibitor of PIKfyve kinase and
50 camostat mesylate and nafamostat mesylate, inhibitors of TMPRSS2 protease. Our
51 research is significant for having uncovered an unexpected synergism in the effective
52 inhibitory activity of apilimod used together with camostat mesylate or with nafamostat
53 mesylate.

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection has caused the global pandemic known as coronavirus disease 2019 (COVID-2019). Currently there is no widespread use of an antiviral agent against the disease, but several candidates have been identified (1-6). Apilimod is currently in a clinical trial for the prevention of SARS-CoV-2 infections in the United States of America (ClinicalTrials.gov Identifier: NCT04446377). The TMPRSS2 protease inhibitor camostat mesylate has been tested with hospitalized COVID-19 patients in Denmark (7) and is in a clinical trial on adult COVID-19 patients in France (ClinicalTrials.gov Identifier: NCT04608266), while the higher affinity TMPRSS2 protease inhibitor, nafamostat mesylate, is being used in COVID-19 related trials in Russia (ClinicalTrials.gov Identifier: NCT04623021), South Korea (ClinicalTrials.gov Identifier: NCT04418128) and Japan (8).

Targeting the entry route of SARS-CoV-2 has been particularly challenging because there appear to be at least two different pathways for virus entry into cells (1). SARS-CoV-2 entry is mediated by the virus spike protein (9-11) and requires the receptor ACE2 in the host cell (1, 12, 13). Upon engagement of the ACE2 receptor, the spike protein catalyzes fusion of the viral membrane envelope with a host cell membrane to release the contents of the virus into the cytoplasm of host cells. For the spike protein to facilitate this reaction, it must first be cleaved by a host cell protease (1, 10). This can be accomplished by different host proteases including TMPRSS2 and TMPRSS4 (14), Factor Xa (15, 16), and by cathepsins during endocytosis (1). The protease inhibitors E-64 and camostat mesylate target cathepsin and TMPRSS2 and inhibit

SARS-CoV-2 infection (1), but their low (micromolar) affinities have made them unlikely candidates for clinical use. Nafamostat mesylate inhibits TMPRSS2 with a high (nanomolar) affinity (2, 6). Apilimod has been shown to function in the endosomal pathway by inhibiting PIKfyve kinase causing a defect in viral trafficking prior to entry (3) but how this relates to the cathepsin or TMPRSS2 protease dependent pathways has not been investigated.

Using a chimeric vesicular stomatitis virus (VSV) in which the attachment and fusion glycoprotein G is replaced by the spike S protein of SARS-CoV-2 (VSV-SARS-CoV-2) (17) we investigate inhibition of infection in different cell types known to contain different levels of cathepsin and TMPRSS2 proteases. We tested combinations of the protease inhibitors E-64 (cathepsin), camostat mesylate and nafamostat mesylate (TMPRSS2) and the lipid kinase inhibitor apilimod (PIKfyve kinase) on VSV-SARS-CoV-2 infection of multiple cell types. We observed a 5-fold synergistic effect in the infection of VSV-SARS-CoV-2 by simultaneous inhibition of TMPRSS2 and PIKfyve kinase. Furthermore, the synergistic inhibitory effects on infection by VSV-SARS-CoV-2 or a clinical isolate of SARS-CoV-2 were observed with nanomolar concentrations of camostat mesylate and apilimod. This finding suggests that a combination of inhibitors that target two different host factors in the entry pathway of SARS-CoV-2 will likely be more effective than targeting either alone.

RESULTS AND DISCUSSION

VSV-SARS2-CoV-2 infection is partially prevented by PIKfyve kinase or TMPRSS2

protease inhibitors

It has been proposed that two distinct host proteases, TMPRSS2 and cathepsin facilitate different SARS2-CoV-2 viral entry routes - cell surface or endosomal - and their abundance in different cell types may influence the entry pathway (1). To test this, we employed a panel of inhibitors on SARS-CoV-2 infection in African green monkey kidney epithelial derived VeroE6 cells poorly expressing TMPRSS2, VeroE6 cells stably expressing ectopic TMPRSS2, as well as human colon carcinoma derived Caco-2 cells and human lung derived Calu-3 cells naturally expressing TMPRSS2.

We first established the relative importance of the cathepsin-dependent route for infection of VSV-SARS-CoV-2 by determining the effect of the cathepsin inhibitor E-64 on expression of eGFP mediated by VSV-SARS-CoV-2. As summarized in the plots in Fig. 1A, we found a cell type dependence in the extent of infection block with a half maximal effective concentration (EC_{50}) in the ~5-10 μ M range. Notably, only Vero E6 cells treated with E-64 displayed a full infection block, the result expected for cells expressing cathepsin but no TMPRSS2; in contrast, cells expressing cathepsin and TMPRSS2 (VeroE6 + TMPRSS2, Caco-2) displayed a partial block, while Calu-3 cells with minimal expression of cathepsin L (18) did not respond, as expected, to E-64 treatment (1).

We extended the analysis and studied the inhibitory effect by the PIKfyve kinase inhibitor apilimod on VSV-SARS-CoV-2 infection (Fig. 1 B). We found that all cell types shown to be sensitive to E-64 also responded to treatment with apilimod with $EC_{50} \sim 10$ nM whereas Calu-3, which is insensitive to E-64 (19), did not respond. Although apilimod doesn't inhibit cathepsin B or L (20), these observations are consistent with potential modulation of the endosomal availability of cathepsin by the activity of PIKfyve kinase (21, 22).

A similar analysis, to document the reduction of viral infection using camostat mesylate to block the protease activity of TMPRSS2 (Fig. 1C) or nafamostat mesylate (Fig. 1D) showed, as anticipated, no response in Vero E6 cells lacking TMPRSS2, partial inhibition in Vero E6 or Caco-2 cells expressing TMPRSS2 or full inhibition in Calu-3 cells naturally expressing TMPRSS2 but insensitive to cathepsin inhibition.

These observations are consistent with previous results (1, 2, 6, 19) that infection by VSV-SARS-CoV-2 occurred through two complementary entry pathways with different importance depending on the cell type, one depending on the proteolytic activity of cathepsin and the second relying on the proteolytic activity of TMPRSS2. As a negative control for these experiments, we included infection by the parental VSV (Fig. 1E), whose ability to infect host cells is known to be independent of the enzymatic activities of cathepsin, TMPRSS2, or PIKfyve kinase (1-4, 6). As expected, none of the inhibitors for these enzymes influenced the extent of VSV infectivity in any of the cells used. From

these results we could also exclude potential cytotoxic effects by the compounds in the concentration range used.

Synergistic prevention of VSV-SARS2-CoV-2 infection by combined use of PIKfyve kinase and TMPRSS2 protease inhibitors

Since the cathepsin and TMPRSS2-dependent activation of SARS-CoV-2 correspond to complementary entry pathways thought to act independent of each other, we expected an additive inhibitory effect upon their simultaneous inhibition in cells that express both proteases. Indeed, VSV-CoV-2 infection of Vero E6 cells expressing cathepsin and TMPRSS2 (VeroE6 +TMPRSS2) were equally inhibited ($EC_{50} \sim 2 \mu M$) by camostat mesylate in the absence or increasing concentrations of E-64 up to $20 \mu M$, the concentration at which E-64 maximally blocked VSV-CoV-2 infection (Fig. 2A). We used SynergyFinder 2.0 (23) to compare the combined response obtained experimentally with the expected outcome calculated by the Bliss synergy-scoring model. Using this reference model, that considers multiplicative effect of single drugs as if they acted independently (23, 24), we obtained an overall δ -score of 6 indicative of an additive interaction between camostat and E-64. At combined high levels of camostat and E64, these protease inhibitors also showed weak concentration dependence, consistent with the prediction of a published model (25). In contrast, combined use of camostat mesylate and apilimod led to enhanced inhibition of VSV-CoV-2 infection with a ~ 2 -fold decrease in the EC_{50} of camostat mesylate (from 1 to $0.4 \mu M$) (Fig. 2B), suggestive of a synergy effect.

We carried a similar set of experiments using Calu-3 cells known to be deficient in Cathepsin-L (18, 26) and which are poorly susceptible to infection of filoviruses mediated by the cathepsin-dependent infection route (18). These cells are insensitive to inhibition by of VSV-SARS-CoV-2 and SARS-CoV-2 infection by E-64 (1, 19, 27, 28) (Fig.1B) and apilimod ((19, 29), Fig. 1C). While presence of E-64 did not affect the inhibition profile of camostat mesylate (Fig. 2C), we detected a ~ 5-fold decrease from 1 μ M to 0.2 μ M in the EC₅₀ of apilimod in cells simultaneously treated with 10 μ M E-64 (Fig. 2D). This result was unexpected, given that infection of the Calu-3 cells by SARS-CoV-2 didn't appear to be affected by apilimod ((29) and Fig. 1C).

We extended the synergy analysis to verify the effects upon simultaneous use of variable amounts of apilimod and nafamostat mesylate, another inhibitor of TMPRSS2. Our data also indicates enhanced infection inhibition for nafamostat mesylate with a ~ 20-fold decrease of EC₅₀ from 0.02 to 0.001 μ M (Fig. 3A, central panel). Simultaneous treatment with apilimod and increasing amounts of nafamostat mesylate led to an analogous enhanced potency of apilimod inhibition with a ~10-fold decrease of EC₅₀ from 0.01 to 0.001 μ M (Fig. 3A, right panel). Formal evaluation for synergism using the Bliss reference model was consistent with a synergy δ -score of 41 (a score greater than 10 indicates synergy (23)).

Comparable strong synergistic effects (δ -score of 34) by combined use of apilimod and nafamostat mesylate were observed in Vero E6 cells expressing TMPRSS2 (Fig. 3B, central and right panels) infected with VSV-SARS-CoV-2 D614G whose spike S protein

includes the point mutation known to increase infectivity of the native SARS-CoV-2 (30). While this mutation has no effect on the virus entry mechanism or sensitivity to proteases, it stabilizes the stalk region of the spike, which otherwise tend to fall apart after furin cleavage between S1 and S2 (30-33).

Taken together, these synergy results highlight the unexpected non-additive involvement of PIKfyve kinase activity for the functional effectiveness of TMPRSS2 to mediate viral entry along the TMPRSS2 route.

Synergistic prevention of SARS2-CoV-2 infection by combined use of PIKfyve kinase and TMPRSS2 protease inhibitors

To determine whether SARS-CoV-2 virus also display an enhanced block of infection upon combined use of apilimod and camostat mesylate we infected Caco-2 cells ectopically expressing hACE2, the main receptor for SARS-CoV-2. This cell line was created as a way to enhance the susceptibility of Caco-2 to infection by SARS-CoV-2 since the parental cells express low endogenous levels of ACE2 (34); successful infection was scored 18 hrs post infection by the appearance of viral N protein using immunofluorescence microscopy (Fig. 4, left panel). The inhibitory EC₅₀ of camostat mesylate decreased ~ 6-fold from 0.6 to 0.1 μ M when used together with apilimod, (Fig. 4, right panel). This observation extends our results from VSV-SARS-CoV-2 chimeras to SARS-CoV-2 virus, and illustrates that combined chemical inhibition of PIKfyve kinase and TMPRSS2 protease activities is likely to also prevent SARS-CoV-2 infection with strong synergy.

211

212 **Final remarks**

213 According to our present understanding of early steps necessary for successful
 214 infection, SARS-CoV-2 would enter cells using two main redundant routes, each
 215 requiring cleavage of the viral S spike protein, one dependent on the proteolytic
 216 activities of members of the cathepsin family, the other requiring proteolysis by
 217 TMPRSS2 or similar transmembrane-serine proteases. Expression level of these
 218 proteases depends on cell type and hence their relative importance to support
 219 successful infection would hinge in part on their relative expression. Cathepsin is
 220 primarily found in late endosomes or lysosomes and requires low pH for optimal
 221 enzymatic activity; for these reasons it has been proposed that SARS-CoV-2 entry
 222 occurs from the endolysosomal compartment (35, 36). In contrast, TMPRSS2 is thought
 223 to be at the cell surface (37) and its optimal proteolytic activity is pH independent (38);
 224 hence it has been inferred that TMPRSS2 cleavage of spike S protein occurs on virions
 225 at the plasma membrane from which viral entry is then assumed to occur (1).

226

227 Not surprising for protease inhibitors with different targets, combined inhibition of the
 228 enzymatic activities of cathepsin and TMPRSS2 by the respective protease inhibitors E-
 229 64 and camostat mesylate led to additive prevention of infection by
 230 SARS-CoV-2 (1) and by the VSV-SARS-CoV-2 chimera (this study).

231

232 Apilimod specifically inhibits PIKfyve kinase thereby blocking accumulation of PI(3,5)P₂,
 233 a key phosphoinositide required to modulate the function of a number of proteins

involved in late endosomal traffic (3, 21, 39, 40). While it is now well established that interference with PIKfyve kinase prevents infection of a selected group of viruses including Ebola (20), VSV-Ebola (3), Marburg (20), VSV-SARS-CoV-2 (3) and native SARS-CoV-2 (3-5), the molecular mechanism events responsible for the interference remain to be determined. It has been shown that Apilimod has anti-Ebola infection synergy with clomiphene cytrate (δ -score of 23.8; re-calculated from data in Fig. 1B of (41) using the Bliss reference model (23)) and also with other drugs that interfere, by unknown mechanisms, with Ebola entry in late endosomes (41). We have now shown synergistic inhibition of SARS-CoV-2 infection by combined use of apilimod and camostat or apilimod and nafamostat mesylate, even in cells as diverse as Vero (which have an active cathepsin entry pathway) and Calu-3 (which lack it).

Prevailing models for the cellular location of TMPRSS2-activated SARS-CoV-2 entry favor fusion at the cell surface rather than from endosomal compartments, because TMPRSS2 activity is pH independent. Ebola or SARS-CoV-2 viruses accumulate in EEA1 containing early endosomes (3, 20), a behavior consistent with the disruption of endolysosomal traffic that occurs when PIKfyve kinase activity is inhibited by genetic or pharmacological means (21, 40, 42). It is therefore possible that changes in the endosomal milieu, combined with redirection of SARS-CoV-2 and/or TMPRSS2 traffic in the presence of apilimod, might make infection more dependent on TMPRSS2 pharmacological inactivation than in its absence. Combined presence of apilimod and camostat or nafamostat mesylate would manifest as synergy of these compounds to prevent infection. These observations would also be consistent with favored entry of

TMPRSS2-activated SARS-CoV-2 from internal membrane compartments rather than from the cell surface.

The strong synergy we described in the laboratory setting for single cell types upon combined use of the PIKfyve kinase and TMPRSS2 inhibitors highlights the potential clinical advantage of using a similar strategy to ameliorate the viral load and potentially reduce the risk to COVID-19 in patients infected with SARS-CoV-2 based on simultaneous inhibition of these enzymes. Variants of apilimod and camostat, rendered more soluble by mesylate modification, are currently in clinical trials as orally administered inhibitors dispensed alone for various indications, including COVID-19 (7, 43, 44). More soluble nafamostat mesylate has been administered intravenously to treat COVID-19 patients (8). We do not yet know how efficiently these inhibitors reach nasopharyngeal and lung tissues; nevertheless, we surmise that combined use of these PIKfyve and TMPRSS2 inhibitors might also recapitulate the ~5-10-fold increase in inhibitor efficiency that we observe in the laboratory setting, offering considerable advantage for clinical application.

FIGURE LEGENDS

Fig. 1. Protease inhibitors E-64, apilimod, camostat mesylate or nafamostat mesylate prevent infection by VSV-SARS-CoV-2 but not by VSV.

(A) Schematic of infectivity assay for cells pretreated for 1 h or not with the inhibitors, subsequently infected with VSV-SARS-CoV-2 for 1 h in the presence or absence of inhibitors. The cells were incubated for another 7 h in the presence or absence of inhibitors and then fixed; the percentage of cells expressing eGFP measured by spinning disc confocal microscopy.

(B-F) Quantification of the number of infected cells from three independent experiments, each determined from 5 fields of view containing 80-200 cells per experiment (error bars show SEM) for the indicated cell types. Infected Vero **(B)** or Vero + TMPRSS2 cells **(C)** were analyzed 8 hpi using 0.5 µg/mL VSV-SARS-CoV-2 RNA. Infected Caco-2 **(D)** or Calu-3 cells **(E)** were analyzed 8 hpi using 5 µg/mL VSV-SARS-CoV-2 RNA. Cells infected with 0.075 µg/mL VSV RNA **(F)** were analyzed 6 hpi. In each case, these virus concentrations and conditions of infection corresponded to an MOI of ~ 0.5.

Fig. 2. Synergistic inhibition of VSV-SARS2-CoV-2 infection by combined use of apilimod and camostat mesylate.

Data are from infection results using VSV-SARS2-CoV-2 obtained with different cell types in the absence or combined presence at increasing concentrations of E-64 and camostat mesylate **(A, C)** or apilimod and camostat mesylate **(B, D)**. Representative maximum-Z projections views (left panels) are from whole-cell volume images obtained with optical sections separated by 0.5 µm using spinning disk confocal microscopy; cells

were infected with 0.5 µg/mL viral RNA VSV-SARS-CoV-2 and imaged 8 hpi. Scale bar: 50 µm. Corresponding quantifications of infection (right panels) are shown in the plots. Each point corresponds to one independent experiment; the data represent results from 5 fields of view containing 80-200 cells per experiment. Estimated EC₅₀'s are indicated. Significance in the difference of EC₅₀ was determined from fitting replicated experiments and using an unpaired t-test; P-values were 0.02 for **(B)** and 0.002 for **(D)**; there was no statistically significant difference in the EC₅₀ values for the experiments reported in **(A)** or **(C)**.

Fig. 3. Synergistic inhibition of VSV-SARS2-CoV-2 infection by combined use of apilimod and nafamostat mesylate.

Data are from infection results obtained with VeroE6 + TPRSS2 cells in the absence or combined presence at increasing concentrations of apilimod and nafamostat mesylate. Representative maximum-Z projections views (left panels) are from whole-cell volume images obtained with optical sections separated by 0.5 µm using spinning disk confocal microscopy; cells were infected with 0.5 µg/mL viral RNA VSV-SARS-CoV-2 **(A)** or 0.2 µg/mL VSV-SARS-CoV-2 D614G viral RNA **(B)** and imaged 8 hpi. Scale bar: 50 µm. Corresponding quantifications of infection (right panels) are shown in the plots. Each point corresponds to one independent experiment; the data represent results from 5 fields of view containing 80-200 cells per experiment. Estimated EC₅₀'s are indicated. Both plots in the central and right panels used the same data.

Fig. 4. Synergistic inhibition of SARS2-CoV-2 infection by combined use of apilimod and camostat mesylate.

Natural SARS-CoV-2 infection inhibition in Caco-2 + hACE2 cells (MOI = 0.5) by camostat mesylate in the absence or combined presence of apilimod. Representative examples of images of fixed samples stained with an antibody specific for N (red) and with Hoechst DNA stain (cyan) to identify the nuclei obtained using wide field epifluorescence microscopy (left panels) and corresponding quantifications of infection (right panels) are shown. Each data points were from 4 independent experiments, representing results from 6,000-10,000 cells per experiment. Estimated EC₅₀'s are shown. Significance in the difference (P-value of 0.01) of EC₅₀ was determined using an unpaired t-test after a least square non-linear fitting of the data curves from replicated experiments.

MATERIAL AND METHODS

Materials

These reagents were purchased as indicated: Dulbecco modified Eagles Medium (DMEM) supplemented with 4.5 g/L glucose, L-glutamine, and sodium pyruvate (Corning, Inc., cat. 10-013-CV), fetal bovine serum (FBS, Atlanta Biologicals, cat. S11150H), penicillin-streptomycin 100x solution (VWR, cat. 45000-652), camostat mesylate (Sigma-Aldrich, cat. SML0057), E-64 (Santa Cruz Biotechnology, cat. sc-201276), nafamostat mesylate (Cayman Chemical Company, cat. #14837), apilimod (MedChem Express, cat. HY-14644), dimethyl sulfoxide (DMSO, Sigma-Aldrich, cat. 26855), Round cover glass #1.5, 25 mm (Cellvis cat. C8-1.5H-N), Polydimethylsiloxane (PDMS, SYLGARD from Krayden, cat. DC4019862), isopropyl alcohol (VWR, cat. 9080-03/MK303108), potassium hydroxide (Sigma-Aldrich, cat. 484016), wheat germ agglutinin (WGA)-Alexa647 (Invitrogen, cat. W32466), paraformaldehyde (Sigma-Aldrich, cat. P6148), sodium phosphate dibasic (Thermo Fisher Scientific, cat. BP329-1), potassium phosphate monobasic (Sigma-Aldrich, cat. P5379), sodium chloride (EMD Millipore, cat. SX0420-5), and potassium chloride (Thermo Fisher Scientific, cat. P217-500), TRIS (Goldbio, cat. T-400-500), EDTA (Sigma-Aldrich, cat. E5134), sucrose (Sigma-Aldrich, cat. S0389), fetal calf serum (FCS, Hyclone, cat. SH30073.03), bovine serum albumin (GE Healthcare), Triton-X (Thermo Fischer, cat. 28314), Hoechst DNA stain (Thermo Fischer, cat. 62249), L-glutamine (Sigma, cat. G7513), Hoechst DNA dye (cat. H6021, Sigma-Aldrich), and Alexa 647 fluorescently labelled goat anti-rabbit antibody (Thermo Fisher, cat. A32733).

Purification of VSV-SARS-CoV-2 chimeras

Generation of recombinant VSV (Indiana serotype) expressing eGFP (VSV-eGFP) and VSV-eGFP chimeras whose glycoprotein G was replaced with either the wild type spike S protein of SARS-CoV-2 Wuhan-Hu-1 strain (VSV-SARS-CoV-2) or with the point mutant D614G (VSV-SARS-CoV-2 D614G) was done as described (17). Briefly, VSV was grown by infection of BSRT7/5 and VSV-SARS-CoV-2 chimeras were grown by infection of MA104 cells. Cells were grown in 12-18 150 mm dishes and infected at an MOI of 0.01. Virus-containing supernatant was collected at 48 hours post infection. The supernatant was clarified by low-speed centrifugation at $1,000 \times g$ for 10 min at room temp. An initial pellet of virus and extracellular particles was generated by centrifugation in a Ti45 fixed-angle rotor at $30,000 \times g$ (25,000 rpm) for 2 hours at 4°C . The pellet was resuspended overnight in 1X NTE (100 mM NaCl, 10 mM Tris-HCl pH 7.4, 1 mM EDTA) at 4°C . The resuspended pellet was layered on top of a 15% sucrose-NTE cushion and subjected to ultracentrifugation in a SW55 swinging bucket rotor at $110,000 \times g$ (35,000 rpm) for 2 hours at 4°C . The supernatant was aspirated and virus pellet resuspended in 1X NTE overnight at 4°C . The resuspended virus pellet was separated on a 15-45% sucrose-NTE gradient by ultracentrifugation in a SW55 swinging bucket rotor at $150,000 \times g$ (40,000 rpm) for 1.5 hours at 4°C . The predominant light-scattering virus band was observed in the lower third of the gradient and was extracted by side puncture of the gradient tube. Extracted virus was diluted in 1X NTE and collected by ultracentrifugation in a Ti60 fixed-angle rotor at $115,000 \times g$ (40,000 rpm) for 2 hours at 4°C . The final pellet was re-suspended overnight in 1X NTE in a volume of 0.2 to 0.5 mL depending on the size of the pellet and stored at 4°C for use in subsequent imaging experiments.

376

377 **Isolation and propagation of SARS-CoV-2**

378 Human samples were obtained under the Helsinki University Hospital laboratory
379 research permit 30 HUS/32/2018 § 16. Isolation of SARS-CoV-2 from a COVID-19
380 Briefly, a nasopharyngeal swab in 500 µl of Copan UTM® Universal Transport Medium
381 was inoculated on Calu-3 cells (P1) and incubated for 1 h at 37°C, after which the
382 inoculum was removed and replaced with Minimum Essential Medium supplemented
383 with 2% FBS, L-glutamine, penicillin and streptomycin. Virus replication was determined
384 by RT-PCR for SARS-CoV-2 RdRP (45), and the infectious virus collected 48h after
385 inoculation. The P1 stock was propagated once (P2) in VeroE6+TMPRSS2, sequenced
386 and stored at -80 °C. Virus stocks were stored in DMEM, 2% FCS, 2 mM L-glutamine,
387 1% penicillin-streptomycin.

388

389 **Cell Culture**

390 VeroE6 (ATCC CRL-1586), Caco-2 (ATCC HTB-37), and Calu-3 cells (ATCC HTB-55)
391 were purchased from ATTC. VeroE6+TMPRSS2 cells were a gift from Siyan Ding (14).
392 VeroE6, VeroE6+TMPRSS2, Caco-2, and Calu-3 cells were maintained in DMEM
393 supplemented with 25 mM HEPES, pH 7.4, 10% fetal bovine serum and 1% penicillin-
394 streptomycin. VeroE6 and VeroE6+TMPRSS2 cells were grown at 37°C and 5% CO₂
395 and split at a ratio of 1:10 every 3-4 days when cells were at ~90% confluency. Caco-2
396 cells were grown at 37°C and 5% CO₂ and split at a ratio of 1:5 every 3-4 days when
397 cells were at ~95% confluency. Calu-3 cells were grown at 37°C and 7% CO₂ and split
398 at a ratio of 1:3 every 5-6 days when cells were at ~95% confluency. BSR-T7 cells

were derived from BHC cells (46) and grown in DMEM supplemented with 10% FBS and 1% penicillin-streptomycin. BSR-T7 were grown at 37°C and 5% CO₂ and split at a ratio of 1:20 every 2-3 days when cells were at ~90% confluency. MA104 cells (ATCC, CRL-2378.1) were grown in Media 199 supplemented with 10% FBS and 1% penicillin-streptomycin. MA104 cells were grown at 37°C and 5% CO₂ and split at a ratio of 1:3 every 2 days when cells were at ~90% confluency. The media was changed in all cell types every 2 days and regularly tested for presence of mycoplasma.

Caco-2 cells stably expressing human ACE2 were generated by transduction with third generation lentivirus pLenti7.3/V5 DEST ACE2-EmGFP (prepared by the cloning facility Dream-Lab, Institute of Biotechnology, University of Helsinki, Helsinki, Finland; the expression of Emerald GFP is driven by the SV40 promoter. Cells expressing eGFP were isolated by FACS. These cells were grown in DMEM media supplemented with 10% fetal calf serum, 1% penicillin-streptomycin, and 2 mM L-glutamine. Cells were kept at 37°C, 5% CO₂, and split every 2-4 days when ~90% confluent.

Infection protocol for VSV and VSV-SARS-CoV-2

Polydimethylsiloxane was cured by vigorously mixing with curing reagent at a ratio of 1 to 10. PDMS was poured into 10 cm petri dish (5 g of PDMS per plate) and incubated at 90 °C for ~4 hours. Plates were removed and stored at room temperature until use. PDMS was removed from petri dish using a razor blade. 3 mm holes were punched into PDMS which was cut with a blade to be ~6-10 mm. Glass slides were cleaned by sonication first in isopropanol for 20 min then in 0.5M potassium hydroxide for 20 min

followed by extensive washing in Milli-q water. Glass was dried in an oven 60 °C for 30 minutes then bonded to the PDMS by exposing PDMS and glass to air plasma at 750 mTorr, 30 W for 2 minutes using a PDC-001 plasma cleaner (Harrick Plasma) then firmly pressing glass and PDMS together. This was followed by placing bonded glass and PDMS in an oven at 90 °C for 20 minutes. The glass cover slips mounted with a PDMS well were then placed in 70% ethanol for 10 minutes to sterilize prior to use for cell culture.

The day prior to the experiment cells were plated in PDMS wells on the glass slide stored in a 6 well plate at a density to achieve ~70% confluence the day of the experiment. On the day of the experiment cells were incubated with the desired inhibitor concentration for 1 hour. Media was then removed and virus that had been diluted into media containing the indicated inhibitor concentration was added to the well in a volume of 10 µL. Media was left in the 6 well plate outside of the PDMS well at a level less than the height of the PDMS well to maintain humidity and prevent evaporation. After the virus was incubated with the cells for 1 hour the cells were washed with media containing the indicated inhibitor and then the well was filled with fresh media. In all experiment's cells were kept at 37°C and 5% (for Vero, Vero+TMRPSS2, and Caco-2 cells) or 7% CO₂ (for Calu-3 cells) and media was pre warmed to 37°C. At 6 hours after initiating the infection with VSV or 8 hours after initiating the infection with VSV-SARS-CoV-2 media was removed and the cells were stained by adding 5µg/mL WGA-Alexa647 in PBS (137 mM NaCl, 2.7 mM KCl, 8 mM Na₂HPO₄, 2 mM KH₂PO₄, pH 7.4) for 30 seconds at room temperature. Cells were then

washed with sterile PBS 2 times and then fixed with 4% paraformaldehyde in PBS.

Infected cells were imaged using a spinning disk confocal microscope with a 40X oil objective with a pixel size of 0.33 μm where 20 optical planes were taken at 0.5 μm apart for every field of view (47). Cells were considered infected when they displayed a cytosolic eGFP fluorescence signal with a relative intensity of 1.4 times that of the background of uninfected cells. Example images are max intensity projections of the cell volume where the outline (white) line was obtained by tracing the WGA-Alexa647 signal outlining the cell.

Infection protocol for SARS-CoV-2

All experiments with SARS-CoV-2 were performed in BSL3 facilities at the University of Helsinki with appropriate institutional permits. Virus samples were obtained under the Helsinki University Hospital laboratory research permit 30 HUS/32/2018 § 16. Virus titers were determined by plaque assay in VeroE6+TMPRSS2 cells. Cells in DMEM, supplemented with 10% FBS, 2 mM L-glutamine, 1% penicillin-streptomycin, 20 mM HEPES (pH 7.2) were seeded 48 h before treatment at a density of 15,000 cells per well in 96-well imaging plates (PerkinElmer cat. 6005182). Inhibitors, or DMSO control, were either added 60 min prior to infection, or added 90 min post infection at a multiplicity of infection (MOI) 0.5 plaque forming units per cell. Infections were carried for 20 h at 37 °C and 5% CO₂. Cells were then fixed with 4% paraformaldehyde in PBS for 30 min at room temperature before being processed for immuno-detection of viral N protein, automated fluorescence imaging and image analysis. Briefly, viral NP was detected with an in house developed rabbit polyclonal antibody (34) counterstained with Alexa-647-conjugated goat

anti rabbit secondary antibody, and nuclear staining done using Hoechst DNA dye. Automated fluorescence imaging was done using an epifluorescence high-content Molecular Device Image-Xpress Nano microscope equipped with a 20x objective and a 4.7Mpixel CMOS camera (pixel size 0.332 μm). Image analysis was performed with the software CellProfiler-3 (www.cellprofiler.org). Automated detection of nuclei was obtained using the Otsu algorithm inbuilt in the software. To automatically identify infected cells, an area surrounding each nucleus (5 pixels expansion of the nuclear area) was used to estimate the fluorescence intensity of the viral NP immuno labeled protein, using an intensity threshold such that less than 0.01% of positive cells were detected in non-infected wells.

Statistical analysis

The significance of response (synergy δ -score) upon combined use of two drugs (Fig. 3) was calculated using the Bliss reference model for combination of two drugs incorporated in the stand-alone web-application SynergyFind v2 (23). This model assumes a stochastic process in which the effects of two drugs act independently. δ -score values between -10 and 10 suggest an additive interaction; δ -scores larger than 10 suggest the interaction is synergistic.

The EC_{50} values in Figures 2 B, C, and Figure 4 were obtained from replicate determinations calculated with least-square nonlinear curve fitting using Igor Pro (WaveMetrics). An unpaired T-test was then used to determine the statistical significance in the difference in the values of EC_{50} of camostat mesylate as a function of apilimod.

AUTHOR CONTRIBUTIONS

We thank Sean Whelan for comments and suggestions. Alex J.B. Kreutzberger carried out all the experiments in Figs. 1-3. Ravi Ojha and Giuseppe Balistreri carried out the experiments in Fig. 4. Anwesha Sanyal maintained the cells lines and assisted with infectivity assays. Ravi Ojha generated Caco2-ACE2+EmGFP cells and propagated SARS-CoV-2 virus under the supervision of Olli Vapalahti. Jesse Pyle generated the stocks of VSV-SARS-CoV-2 chimeras. We thank Tegye John Vadakkan for maintaining the spinning disc confocal microscope. This research was supported by a NIH Maximizing Investigators' Research Award (MIRA) GM130386, by research grants from the Danish Technical University and SANA and unrestricted funds from IONIS to T.K; Harvard Virology Program, NIH training Grant T32 AI07245 postdoctoral fellowship to A.J.B.K.; by an Academy of Finland research grant 336490 by the Jane and Aatos Erkkö Foundation, by the EU Horizon 2020 program VEO 874735, and by Helsinki University Hospital Funds TYH2018322 to O.V; by a research grant from the Academy of Finland grant 318434 and private funds supporting COVID-19 research to G.B.; by the University of Helsinki graduate program in Microbiology and Biotechnology to R.O. Tom Kirchhausen and Alex J.B. Kreutzberger were responsible for the overall design of the study; Tom Kirchhausen and Alex Kreutzberger drafted the manuscript; all authors commented on the manuscript.

510 REFERENCES


- 511 1. Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, Schiergens
512 TS, Herrler G, Wu NH, Nitsche A, Muller MA, Drosten C, Pohlmann S. 2020. SARS-
513 CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically
514 Proven Protease Inhibitor. *Cell* 181:271-280 e8.
- 515 2. Hoffmann M, Schroeder S, Kleine-Weber H, Muller MA, Drosten C, Pohlmann S. 2020.
516 Nafamostat Mesylate Blocks Activation of SARS-CoV-2: New Treatment Option for
517 COVID-19. *Antimicrob Agents Chemother* 64.
- 518 3. Kang YL, Chou YY, Rothlauf PW, Liu Z, Soh TK, Cureton D, Case JB, Chen RE,
519 Diamond MS, Whelan SPJ, Kirchhausen T. 2020. Inhibition of PIKfyve kinase prevents
520 infection by Zaire ebolavirus and SARS-CoV-2. *Proc Natl Acad Sci U S A* 117:20803-
521 20813.
- 522 4. Ou X, Liu Y, Lei X, Li P, Mi D, Ren L, Guo L, Guo R, Chen T, Hu J, Xiang Z, Mu Z,
523 Chen X, Chen J, Hu K, Jin Q, Wang J, Qian Z. 2020. Characterization of spike
524 glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-
525 CoV. *Nat Commun* 11:1620.
- 526 5. Riva L, Yuan S, Yin X, Martin-Sancho L, Matsunaga N, Pache L, Burgstaller-
527 Muehlbacher S, De Jesus PD, Teriete P, Hull MV, Chang MW, Chan JF, Cao J, Poon
528 VK, Herbert KM, Cheng K, Nguyen TH, Rubanov A, Pu Y, Nguyen C, Choi A,
529 Rathnasinghe R, Schotsaert M, Miorin L, Dejoze M, Zwaka TP, Sit KY, Martinez-
530 Sobrido L, Liu WC, White KM, Chapman ME, Lendy EK, Glynne RJ, Albrecht R,
531 Ruppin E, Mesecar AD, Johnson JR, Benner C, Sun R, Schultz PG, Su AI, Garcia-Sastre
532 A, Chatterjee AK, Yuen KY, Chanda SK. 2020. Discovery of SARS-CoV-2 antiviral
533 drugs through large-scale compound repurposing. *Nature* 586:113-119.
- 534 6. Yamamoto M, Kiso M, Sakai-Tagawa Y, Iwatsuki-Horimoto K, Imai M, Takeda M,
535 Kinoshita N, Ohmagari N, Gohda J, Semba K, Matsuda Z, Kawaguchi Y, Kawaoka Y,
536 Inoue JI. 2020. The Anticoagulant Nafamostat Potently Inhibits SARS-CoV-2 S Protein-
537 Mediated Fusion in a Cell Fusion Assay System and Viral Infection In Vitro in a Cell-
538 Type-Dependent Manner. *Viruses* 12.
- 539 7. Gunst JD, Staerke NB, Pahus MH, Kristensen LH, Bodilsen J, Lohse N, Dalgaard LS,
540 Bronnum D, Frobert O, Honge B, Johansen IS, Monrad I, Erikstrup C, Rosendal R,
541 Vilstrup E, Mariager T, Bove DG, Offersen R, Shakar S, Cajander S, Jorgensen NP,
542 Sritharan SS, Breining P, Jespersen S, Mortensen KL, Jensen ML, Kolte L, Frattari GS,
543 Larsen CS, Storgaard M, Nielsen LP, Tolstrup M, Saedder EA, Ostergaard LJ, Ngo HTT,
544 Jensen MH, Hojen JF, Kjolby M, Sogaard OS. 2021. Efficacy of the TMPRSS2 inhibitor
545 camostat mesilate in patients hospitalized with Covid-19-a double-blind randomized
546 controlled trial. *EClinicalMedicine* doi:10.1016/j.eclinm.2021.100849:100849.
- 547 8. Doi K, Ikeda M, Hayase N, Moriya K, Morimura N, Group C-US. 2020. Nafamostat
548 mesylate treatment in combination with favipiravir for patients critically ill with Covid-
549 19: a case series. *Crit Care* 24:392.
- 550 9. Huang Y, Yang C, Xu XF, Xu W, Liu SW. 2020. Structural and functional properties of
551 SARS-CoV-2 spike protein: potential antiviral drug development for COVID-19. *Acta*
552 *Pharmacol Sin* 41:1141-1149.
- 553 10. Shang J, Wan Y, Luo C, Ye G, Geng Q, Auerbach A, Li F. 2020. Cell entry mechanisms
554 of SARS-CoV-2. *Proc Natl Acad Sci U S A* 117:11727-11734.

11. Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, Graham BS, McLellan JS. 2020. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 367:1260-1263.
12. Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. 2020. Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science* 367:1444-1448.
13. Yang J, Petitjean SJL, Koehler M, Zhang Q, Dumitru AC, Chen W, Derclaye S, Vincent SP, Soumillion P, Alsteens D. 2020. Molecular interaction and inhibition of SARS-CoV-2 binding to the ACE2 receptor. *Nat Commun* 11:4541.
14. Zang R, Gomez Castro MF, McCune BT, Zeng Q, Rothlauf PW, Sonnek NM, Liu Z, Brulois KF, Wang X, Greenberg HB, Diamond MS, Ciorba MA, Whelan SPJ, Ding S. 2020. TMPRSS2 and TMPRSS4 promote SARS-CoV-2 infection of human small intestinal enterocytes. *Sci Immunol* 5.
15. Du L, Kao RY, Zhou Y, He Y, Zhao G, Wong C, Jiang S, Yuen KY, Jin DY, Zheng BJ. 2007. Cleavage of spike protein of SARS coronavirus by protease factor Xa is associated with viral infectivity. *Biochem Biophys Res Commun* 359:174-9.
16. Frydman GH, Streiff MB, Connors JM, Piazza G. 2020. The Potential Role of Coagulation Factor Xa in the Pathophysiology of COVID-19: A Role for Anticoagulants as Multimodal Therapeutic Agents. *TH Open* 4:e288-e299.
17. Case JB, Rothlauf PW, Chen RE, Liu Z, Zhao H, Kim AS, Bloyet LM, Zeng Q, Tahan S, Droit L, Ilagan MXG, Tartell MA, Amarasinghe G, Henderson JP, Miersch S, Ustav M, Sidhu S, Virgin HW, Wang D, Ding S, Corti D, Theel ES, Fremont DH, Diamond MS, Whelan SPJ. 2020. Neutralizing Antibody and Soluble ACE2 Inhibition of a Replication-Competent VSV-SARS-CoV-2 and a Clinical Isolate of SARS-CoV-2. *Cell Host Microbe* 28:475-485 e5.
18. Gonzalez-Hernandez M, Muller A, Hoenen T, Hoffmann M, Pohlmann S. 2019. Calu-3 cells are largely resistant to entry driven by filovirus glycoproteins and the entry defect can be rescued by directed expression of DC-SIGN or cathepsin L. *Virology* 532:22-29.
19. Dittmar M, Lee JS, Whig K, Segrist E, Li M, Kamalia B, Castellana L, Ayyanathan K, Cardenas-Diaz FL, Morrissey EE, Truitt R, Yang W, Jurado K, Samby K, Ramage H, Schultz DC, Cherry S. 2021. Drug repurposing screens reveal cell-type-specific entry pathways and FDA-approved drugs active against SARS-Cov-2. *Cell Rep* 35:108959.
20. Nelson EA, Dyal J, Hoenen T, Barnes AB, Zhou H, Liang JY, Michelotti J, Dewey WH, DeWald LE, Bennett RS, Morris PJ, Guha R, Klumpp-Thomas C, McKnight C, Chen YC, Xu X, Wang A, Hughes E, Martin S, Thomas C, Jahrling PB, Hensley LE, Olinger GG, Jr., White JM. 2017. The phosphatidylinositol-3-phosphate 5-kinase inhibitor apilimod blocks filoviral entry and infection. *PLoS Negl Trop Dis* 11:e0005540.
21. Baranov MV, Bianchi F, Schirmacher A, van Aart MAC, Maassen S, Muntjewerff EM, Dingjan I, Ter Beest M, Verdoes M, Keyser SGL, Bertozzi CR, Diederichsen U, van den Bogaart G. 2019. The Phosphoinositide Kinase PIKfyve Promotes Cathepsin-S-Mediated Major Histocompatibility Complex Class II Antigen Presentation. *iScience* 11:160-177.
22. Gayle S, Landrette S, Beeharry N, Conrad C, Hernandez M, Beckett P, Ferguson SM, Mandelkern T, Zheng M, Xu T, Rothberg J, Lichenstein H. 2017. Identification of apilimod as a first-in-class PIKfyve kinase inhibitor for treatment of B-cell non-Hodgkin lymphoma. *Blood* 129:1768-1778.
23. Ianevski A, Giri AK, Aittokallio T. 2020. SynergyFinder 2.0: visual analytics of multi-drug combination synergies. *Nucleic Acids Res* 48:W488-W493.


24. Bliss C. 1939. The Toxicity Of Poisons Applied Jointly. *Ann Appl Biol* doi:10.1111/j.1744-7348.1939.tb06990.x:585–615.
25. Padmanabhan P, Desikan R, Dixit NM. 2020. Targeting TMPRSS2 and Cathepsin B/L together may be synergistic against SARS-CoV-2 infection. *PLoS Comput Biol* 16:e1008461.
26. Park JE, Li K, Barlan A, Fehr AR, Perlman S, McCray PB, Jr., Gallagher T. 2016. Proteolytic processing of Middle East respiratory syndrome coronavirus spikes expands virus tropism. *Proc Natl Acad Sci U S A* 113:12262-12267.
27. Murgolo N, Therien AG, Howell B, Klein D, Koeplinger K, Lieberman LA, Adam GC, Flynn J, McKenna P, Swaminathan G, Hazuda DJ, Olsen DB. 2021. SARS-CoV-2 tropism, entry, replication, and propagation: Considerations for drug discovery and development. *PLoS Pathog* 17:e1009225.
28. Mykytyn AZ, Breugem TI, Riesebosch S, Schipper D, van den Doel PB, Rottier RJ, Lamers MM, Haagmans BL. 2021. SARS-CoV-2 entry into human airway organoids is serine protease-mediated and facilitated by the multibasic cleavage site. *Elife* 10.
29. J K, Uckelely Z, P D, M S, S B, PY L. 2020. Host Cell Proteases Drive Early or Late SARS-CoV-2 Penetration. *bioRxiv* doi:<https://doi.org/10.1101/2020.12.22.423906>.
30. Plante JA, Liu Y, Liu J, Xia H, Johnson BA, Lokugamage KG, Zhang X, Muruato AE, Zou J, Fontes-Garfias CR, Mirchandani D, Scharton D, Bilello JP, Ku Z, An Z, Kalveram B, Freiberg AN, Menachery VD, Xie X, Plante KS, Weaver SC, Shi PY. 2020. Spike mutation D614G alters SARS-CoV-2 fitness. *Nature* doi:10.1038/s41586-020-2895-3.
31. Gobeil SM, Janowska K, McDowell S, Mansouri K, Parks R, Manne K, Stalls V, Kopp MF, Henderson R, Edwards RJ, Haynes BF, Acharya P. 2021. D614G Mutation Alters SARS-CoV-2 Spike Conformation and Enhances Protease Cleavage at the S1/S2 Junction. *Cell Rep* 34:108630.
32. Yurkovetskiy L, Wang X, Pascal KE, Tomkins-Tinch C, Nyalile TP, Wang Y, Baum A, Diehl WE, Dauphin A, Carbone C, Veinotte K, Egri SB, Schaffner SF, Lemieux JE, Munro JB, Rafique A, Barve A, Sabeti PC, Kyratsous CA, Dudkina NV, Shen K, Luban J. 2020. Structural and Functional Analysis of the D614G SARS-CoV-2 Spike Protein Variant. *Cell* 183:739-751 e8.
33. Zhang L, Jackson CB, Mou H, Ojha A, Peng H, Quinlan BD, Rangarajan ES, Pan A, Vanderheiden A, Suthar MS, Li W, Izard T, Rader C, Farzan M, Choe H. 2020. SARS-CoV-2 spike-protein D614G mutation increases virion spike density and infectivity. *Nat Commun* 11:6013.
34. Cantuti-Castelvetri L, Ojha R, Pedro LD, Djannatian M, Franz J, Kuivanen S, van der Meer F, Kallio K, Kaya T, Anastasina M, Smura T, Levanov L, Szirovicza L, Tobi A, Kallio-Kokko H, Osterlund P, Joensuu M, Meunier FA, Butcher SJ, Winkler MS, Mollenhauer B, Helenius A, Gokce O, Teesalu T, Hepojoki J, Vapalahti O, Stadelmann C, Balistreri G, Simons M. 2020. Neuropilin-1 facilitates SARS-CoV-2 cell entry and infectivity. *Science* 370:856-860.
35. Bayati A, Kumar R, Francis V, McPherson PS. 2021. SARS-CoV-2 infects cells following viral entry via clathrin-mediated endocytosis. *J Biol Chem* doi:10.1016/j.jbc.2021.100306:100306.
36. Khan N, Chen X, Geiger JD. 2020. Role of Endolysosomes in Severe Acute Respiratory Syndrome Coronavirus-2 Infection and Coronavirus Disease 2019 Pathogenesis: Implications for Potential Treatments. *Front Pharmacol* 11:595888.

37. Uhlen M, Fagerberg L, Hallstrom BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson A, Kampf C, Sjostedt E, Asplund A, Olsson I, Edlund K, Lundberg E, Navani S, Szigartyo CA, Odeberg J, Djureinovic D, Takanen JO, Hober S, Alm T, Edqvist PH, Berling H, Tegel H, Mulder J, Rockberg J, Nilsson P, Schwenk JM, Hamsten M, von Feilitzen K, Forsberg M, Persson L, Johansson F, Zwahlen M, von Heijne G, Nielsen J, Ponten F. 2015. Proteomics. Tissue-based map of the human proteome. *Science* 347:1260419.
38. Glowacka I, Bertram S, Muller MA, Allen P, Soilleux E, Pfefferle S, Steffen I, Tsegaye TS, He Y, Gnirss K, Niemeyer D, Schneider H, Drosten C, Pohlmann S. 2011. Evidence that TMPRSS2 activates the severe acute respiratory syndrome coronavirus spike protein for membrane fusion and reduces viral control by the humoral immune response. *J Virol* 85:4122-34.
39. de Lartigue J, Polson H, Feldman M, Shokat K, Tooze SA, Urbe S, Clague MJ. 2009. PIKfyve regulation of endosome-linked pathways. *Traffic* 10:883-93.
40. Kim GH, Dayam RM, Prashar A, Terebiznik M, Botelho RJ. 2014. PIKfyve inhibition interferes with phagosome and endosome maturation in macrophages. *Traffic* 15:1143-63.
41. Dyall J, Nelson EA, DeWald LE, Guha R, Hart BJ, Zhou H, Postnikova E, Logue J, Vargas WM, Gross R, Michelotti J, Deilulis N, Bennett RS, Crozier I, Holbrook MR, Morris PJ, Klumpp-Thomas C, McKnight C, Mierzwa T, Shinn P, Glass PJ, Johansen LM, Jahrling PB, Hensley LE, Olinger GG, Jr., Thomas C, White JM. 2018. Identification of Combinations of Approved Drugs With Synergistic Activity Against Ebola Virus in Cell Cultures. *J Infect Dis* 218:S672-S678.
42. Soares AC, Ferreira A, Marien J, Delay C, Lee E, Trojanowski JQ, Moechars D, Annaert W, De Mueynck L. 2021. PIKfyve activity is required for lysosomal trafficking of tau aggregates and tau seeding. *J Biol Chem* 296:100636.
43. Billich A. 2007. Drug evaluation: apilimod, an oral IL-12/IL-23 inhibitor for the treatment of autoimmune diseases and common variable immunodeficiency. *IDrugs* 10:53-9.
44. Uno Y. 2020. Camostat mesilate therapy for COVID-19. *Intern Emerg Med* 15:1577-1578.
45. Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, Bleicker T, Brunink S, Schneider J, Schmidt ML, Mulders DG, Haagmans BL, van der Veer B, van den Brink S, Wijsman L, Goderski G, Romette JL, Ellis J, Zambon M, Peiris M, Goossens H, Reusken C, Koopmans MP, Drosten C. 2020. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* 25.
46. Buchholz UJ, Finke S, Conzelmann KK. 1999. Generation of bovine respiratory syncytial virus (BRSV) from cDNA: BRSV NS2 is not essential for virus replication in tissue culture, and the human RSV leader region acts as a functional BRSV genome promoter. *J Virol* 73:251-9.
47. Cocucci E, Aguet F, Boulant S, Kirchhausen T. 2012. The first five seconds in the life of a clathrin-coated pit. *Cell* 150:495-507.


A




B




C




D







E




F






A VSV-SARS-CoV-2 : VeroE6 + TMPRSS2



B VSV-SARS-CoV-2 D614G : VeroE6 + TMPRSS2



SARS-CoV-2 : Caco-2 + hACE2

