

The molecular mechanism and evolutionary divergence of caspase 3/7-regulated gasdermin E activation

3

4

5

6 Hang Xu^{1,2,3}, Zihao Yuan^{1,2}, Kunpeng Qin^{1,2,3}, Shuai Jiang^{1,2,3*}, Li Sun^{1,2,3*}

7

8

9

10 ¹CAS and Shandong Province Key Laboratory of Experimental Marine Biology,
11 Institute of Oceanology; CAS Center for Ocean Mega-Science, Chinese Academy of
12 Sciences, Qingdao, China

13 ²Laboratory for Marine Biology and Biotechnology, Pilot National Laboratory for
14 Marine Science and Technology, Qingdao, China

15 ³University of Chinese Academy of Sciences, Beijing, China

16

17 *To whom correspondence should be addressed

18 Email: sjiang@qdio.ac.cn; lsun@qdio.ac.cn

19

Abstract

Caspase (CASP) is a family of proteases involved in cleavage and activation of gasdermin, the executor of pyroptosis. In human, CASP3 and CASP7 recognize the same consensus motif DxxD, which is present in gasdermin E (GSDME). However, human GSDME is cleaved by CASP3 but not by CASP7. The underlying mechanism of this observation is unclear. In this study, we identified a pyroptotic pufferfish GSDME that was cleaved by both pufferfish CASP3/7 and human CASP3/7. Domain swapping between pufferfish and human CASP and GSDME showed that the GSDME C-terminus and the CASP7 p10 subunit determined the cleavability of GSDME by CASP7. p10 contains a key residue that governs CASP7 substrate discrimination. This key residue is highly conserved in vertebrate CASP3 and in most vertebrate (except mammalian) CASP7. In mammals, the key residue is conserved in non-primate (e.g., mouse) but not in primate. However, mouse CASP7 cleaved human GSDME but not mouse GSDME. These findings revealed the molecular mechanism of CASP7 substrate discrimination and the divergence of CASP3/7-mediated GSDME activation in vertebrate. These results also suggested that mutation-mediated functional alteration of CASP probably enabled the divergence and specialization of different CASP members in the regulation of complex cellular activities in mammals.

Key words

Immunity; Pyroptosis; Gasdermin E; Caspase 3; Caspase 7; Evolution

Introduction

Pyroptosis represents a form of programmed cell death that provokes robust inflammatory immune response (Bergsbaken, Fink, & Cookson, 2009; Tsuchiya et al., 2019). Gasdermin (GSDM) serves as the direct executioner of pyroptosis. GSDM forms transmembrane pores to permeabilize the cytoplasmic membrane, which leads to the release of pro-inflammatory cytokines and, if these transmembrane pores persist, pyroptosis (Kovacs & Miao, 2017; J. Shi, Gao, & Shao, 2017; Xia et al., 2021). Human has six GSDM (HsGSDM) family members, HsGSDMA-E, and HsPJVK (Tamura et al., 2007). All HsGSDMs (except for HsPJVK) adopt a two-domain architecture, the N-terminal (NT) pore-forming domain and the C-terminal (CT) autoinhibitory domain (De Schutter et al., 2021; Shao, 2021). Proteolytic cleavage of HsGSDM to remove the autoinhibitory CT domain enables the binding of the lipophilic NT fragment to the cell membrane, where the NT fragments oligomerize and form transmembrane pores to induce osmotic cell lysis (Kuang et al., 2017; X. Liu et al., 2016).

Among the signaling pathways that activate GSDM-mediated pyroptosis, caspase (CASP) 1/4/5/11-mediated GSDMD activation is well documented. In human, HsGSDMD is specifically cleaved by the inflammatory HsCASP1 through multiple inflammasome signaling pathways. HsCASP4/5 also cleave HsGSDMD in response to lipopolysaccharide (LPS) stimulation. After cleavage by HsCASP1/4/5, the pore-forming HsGSDMD-NT fragment becomes unconstrained and triggers pyroptosis (He et al., 2015; Kayagaki et al., 2015; J. Shi et al., 2015). Structural and biochemical analysis reveal that HsCASP1/4 β III/ β III' sheet forms an exosite that interacts with the hydrophobic pocket of HsGSDMD-CT domain, rendering HsGSDMD cleavage independent of CASP recognition of the tetrapeptide motif FLTD (K. Wang et al., 2020). When the tetrapeptide FLTD is mutated to AAAD, HsCASP1/4 cleavage of HsGSDMD is not affected. The residues that form the hydrophobic pocket of HsGSDMD-CT domain are not conserved in HsGSDME, suggesting that HsGSDME cannot be cleaved by HsCASP1/4 via interaction with an exosite (Z. Liu et al., 2020).

Actually, human HsGSDME is specifically cleaved by HsCASP3 at the consensus motif DMPD, which is required for HsCASP3-mediated HsGSDME proteolysis. Mutation of either the Asp residue of the DMPD tetrapeptide motif leads to cleavage resistance of HsGSDME (Rogers et al., 2017; Wang et al., 2017), indicating that HsCASP3-mediated HsGSDME activation paradigm is distinct from that of HsGSDMD. It is intriguing that although HsCASP7 shares the same recognition motif (DxxD) with HsCASP3, HsCASP7 cannot cleave HsGSDME (Agniswamy, Fang, & Weber, 2009; Slee, Adrain, & Martin, 2001). The molecular mechanism underlying the HsCASP3/7 cleavage discrimination of HsGSDME is unknown.

Historically, HsCASP3 and HsCASP7, which are 54% identical, are known to possess similar structure and share overlapping protein substrate repertoire, and are considered to have functionally redundant roles in regulating cell death (Crawford & Wells, 2011; Kumar, 2007; Y. Shi, 2002). Both HsCASP3 and HsCASP7 consist of two polypeptide subunits named p20 and p10 (1:1 ratio), which assemble to form active heterotetramer (Timmer & Salvesen, 2007). HsCASP3/7 contain highly conserved QACRG and SHG motifs in p20 subunit, and SWR and GSWF motifs in p10 subunit, which participate in the catalytic reaction and substrate binding, respectively (Boyce, Degterev, & Yuan, 2004; Cohen, 1997). In the apoptosis signaling pathway, death receptor-mediated extrinsic pathway and mitochondria-mediated intrinsic pathway ultimately converge to the activation of HsCASP3/7 (Budd, Tanneti, Lishnak, & Lipton, 2000; Wajant, 2002). Active HsCASP3/7 cleave a series of protein substrates, including poly(ADP-ribose) polymerase and DNA fragmentation factor 45, which causes chromatin fragmentation and leads to apoptosis (Erener et al., 2012; Wolf, Schuler, Echeverri, & Green, 1999; Zheng et al., 1998). Although HsCASP3 and HsCASP7 exert almost indistinguishable proteolytic specificity to certain polypeptides, there exist functional differences between these two HsCASPs in cleaving some substrates (Brentnall, Rodriguez-Menocal, De Guevara, Cepero, & Boise, 2013; Demon et al., 2009). For instance, HsCASP3 cleaves Bid much more efficiently than HsCASP7, while HsCASP7 cleaves cochaperone p23 more efficiently than HsCASP3

(Slee et al., 2001; Walsh et al., 2008). These findings support the notion that HsCASP3 and HsCASP7 are enzymatically similar but functionally non-redundant.

Different from HsGSDME that is specifically cleaved by HsCASP3, teleost GSDME is cleaved via various modes (Angosto-Bazarra et al., 2022; Yuan, Jiang, Qin, & Sun, 2022). In this study, we identified a functional GSDME (named TrGSDME) from pufferfish *Takifugu rubripes*. We found that TrGSDME was specifically cleaved by both HsCASP3/7 and TrCASP3/7, whereas HsGSDME was cleaved by TrCASP3/7 and HsCASP3, but not by HsCASP7. We examined the underlying mechanism of this observation. We found that the GSDME-CT and the CASP7 p10 were critical for CASP7 cleavage of GSDME. By a series of site-directed mutagenesis, we discovered a previously unrecognized key residue (S234 in HsCASP7) in p10 that was responsible for the discriminative cleavage of HsGSDME by HsCASP7.

Results

Pufferfish GSDME is specifically cleaved by both human and fish CASP3/7

It has long been observed that although human caspase (HsCASP) 3 and 7 share the same consensus recognition motif DxxD, HsCASP3, but not HsCASP7, cleaves human GSDME (HsGSDME) (at the site of DMPD) (Poreba, Strozyk, Salvesen, & Drag, 2013; Wang et al., 2017). The underlying molecular mechanism is unknown. In this study, we found that a pufferfish *Takifugu rubripes* GSDME (designated TrGSDME), which belongs to the GSDMEa lineage of teleost GSDME (Figure-figure supplement 1), was specifically cleaved by both HsCASP3 and 7 (Figure 1A and B). This intriguing observation promoted us to explore the mechanism of HsCASP7 substrate discrimination. We first examined whether TrGSDME could be cleaved by pufferfish CASP (TrCASP) 3/7. The active forms of TrCASP3/7 were prepared (Figure 1C), both exhibited high proteolytic specificity and activity towards the tetrapeptide DEVD (Figure 1D and E), which is the conserved recognition motif of HsCASP3/7. When

incubated with TrGSDME, both TrCASP3 and TrCASP7 cleaved TrGSDME into the NT and CT fragments, similar to that cleaved by HsCASP3/7, in a dose dependent manner (Figure 1F and G). Accordingly, TrCASP3/7-mediated TrGSDME cleavage was inhibited by the CASP3 inhibitor (Z-DEVD-FMK) and the pan-CASP inhibitor (Z-VAD-FMK) (Figure 1H). Based on the molecular mass of the cleaved NT and CT products, we inferred that the tetrapeptide motif ²⁵⁵DAVD²⁵⁸ in the vicinity of the linker region of TrGSDME may be the recognition site of CASP3/7. Indeed, the D255R and D258A mutants of TrGSDME were resistant to TrCASP3/7 and HsCASP3/7 cleavage (Figure 1I and J, Figure 1-figure supplement 2). Taken together, these results demonstrated that TrGSDME was cleaved by fish and human CASP3/7 in a manner that depended on the specific sequence of DAVD (Figure 1K).

CASP3/7-cleaved TrGSDME is functionally activated and induces pyroptosis.

We next examined whether TrGSDME, like HsGSDME, is a functional pyroptosis inducer. For this purpose, HEK293T cells were transfected with mCherry-tagged full length (FL) or NT/CT domain of TrGSDME. The results revealed that TrGSDME-FL and -CT were abundantly expressed in the cells, whereas TrGSDME-NT expression was barely detectable (Figure 2A, Figure 2-figure supplement 1). No significant morphological change or LDH release was observed in cells expressing TrGSDME-FL or -CT (Figure 2B and C). By contrast, cells expressing TrGSDME-NT showed necrotic cell death with massive LDH release and positive Sytox Green staining (Figure 2B and C, Figure 2-figure supplement 2). TrGSDME-NT-induced cell death exhibited osmotic cell membrane swelling, a typical feature of pyroptosis (Figure 2D). Time-lapse imaging showed that TrGSDME-NT triggered rapid cell swelling and membrane rupture, which led to release of cytoplasmic contents and eventually cell death (Figure 2E, Figure 1-video 1). To examine the pyroptotic activity of CASP3/7-cleaved TrGSDME, TrCASP3/7 and TrGSDME were overexpressed in HEK293T cells (Figure 2F and G). The cells expressing TrCASP3, TrCASP7 or TrGSDME alone had no apparent morphological change or LDH release, whereas the cells co-expressing

TrGSDME and TrCASP3 or TrGSDME and TrCASP7 underwent pyroptosis, accompanying with massive LDH release, membrane rupture, and TrGSDME cleavage (Figure 2G-I, Figure 2-figure supplement 3A). Consistently, the presence of the CASP3 inhibitor and pan-CASP inhibitor hampered TrGSDME-mediated pyroptosis (Figure 2J and K). Mutation of the cleavage site (D255R and D258A) inhibited pyroptosis and significantly decreased LDH release and TrGSDME cleavage (Figure 2L and M, Figure 2-figure supplement 3B and C). These results indicated that TrCASP3/7 cleavage was required to activate TrGSDME-mediated pyroptosis.

GSDME-CT domain mediates the recognition by CASP7.

Since it is known that human GSDME is cleaved by CASP3 but not by CASP7, the observation that TrGSDME was cleaved by both human and fish CASP3/7 intrigued us to explore the underlying mechanism. We found that unlike TrCASP1, both HsCASP3 and 7 exhibited proteolytic activity towards the consensus CASP3/7 recognition motif, DMPD, in HsGSDME, but HsCASP7 failed to cleave HsGSDME (Figure 3A and B, Figure 3-figure supplement 1 and 2). Structure analysis revealed that compared to TrGSDME, HsGSDME possesses two additional regions with the possibility to form a loop (261-266 aa) and an α -helix (281-296 aa), respectively (Figure 3C). We tested whether deletion of these two regions could confer HsCASP7 cleavage on HsGSDME. The results showed that similar to the wild type HsGSDME, the loop deletion mutant (Δ 261-266) and the α -helix deletion mutant (Δ 281-296) were cleaved by HsCASP3 but not by HsCASP7 (Figure 3D). Since the NT and CT domains play different roles in GSDM structure maintenance, we constructed GSDME chimeras consisting of HsGSDME-NT plus TrGSDME-CT (named HsNT-TrCT), or TrGSDME-NT plus HsGSDME-CT (named TrNT-HsCT) (Figure 3E). Compared with wild type HsGSDME and TrGSDME, chimeric HsNT-TrCT was cleaved by HsCASP3/7, whereas TrNT-HsCT was cleaved only by HsCASP3 (Figure 3F-I), suggesting that the CT domain determined the cleavability of GSDME by HsCASP7.

The p10 subunit determines the substrate specificity of CASP7.

Since as shown above, unlike HsCASP7, TrCASP7 was able to cleave HsGSDME (Figure 3H), we compared the sequences of TrCASP7 and HsCASP7. The two CASPs share 66.24% sequence identity. In these CASPs, the catalytic motifs SHG and QACRG in the p20 subunit and the substrate binding motifs SWR and GSWF in the p10 subunit are conserved (Figure 4A). To explore their substrate discrimination mechanism, we constructed CASP7 chimeras consisting of HsCASP7 p20 plus TrCASP7 p10 (named Hsp20-Trp10) or TrCASP7 p20 plus HsCASP7 p10 (named Trp20-Hsp10) (Figure 4B, Figure 4-figure supplement 1A). Compared to the wild types (HsCASP7 and TrCASP7), the chimeras exhibited comparable enzymatic activities towards the tetrapeptide substrates DAVD and DMPD (Figure 4C). Like the wild types, both chimeras cleaved TrGSDME and HsNT-TrCT (Figure 4D and E). By contrast, the Hsp20-Trp10 chimera cleaved HsGSDME, whereas the Trp20-Hsp10 chimera did not cleave HsGSDME (Figure 4F). Similar cleavage pattern was observed towards TrNT-HsCT (Figure 4G). These observations indicated that the p10 subunit dictated the cleavage specificity of CASP7 toward GSDME.

The S234 of p10 is the key to HsCASP7 discrimination on HsGSDME.

Comparative analysis of the p10 sequences of TrCASP7 and HsCASP7 identified 13 non-conserved residues. To examine their potential involvement in GSDME cleavage, a series of site-directed mutagenesis was performed to swap the non-conserved residues between HsCASP7 p10 and TrCASP7 p10 (Figure 5A, Figure 4-figure supplement 1B and C). Of the 13 swaps created, S234N conferred on HsCASP7 the ability to cleave HsGSDME (Figure 5B), whereas its corresponding swap in TrCASP7 (N245S) markedly reduced the ability of TrCASP7 to cleave HsGSDME (Figure 5C). None of the 13 swaps affected the ability of HsCASP7 or TrCASP7 to cleave TrGSDME (Figure 5D, E). HsCASP7-S234N cleaved HsGSDME in a dose-dependent manner (Figure 5F), and the cleavage was not enhanced by the additional mutation of S247N&I248V (Figure 5G). Additionally, HsCASP7 and HsCASP7-S234N exhibited similar cleavage

capacities against other CASP7 substrates, such as PARP1 and gelsolin (Figure 5-figure supplement 1) (Erener et al., 2012; Walsh et al., 2008). Previous studies showed that human CASP1/4 cleaved GSDMD through exosite interaction (Z. Liu et al., 2020; K. Wang et al., 2020). However, this exosite is not conserved in HsCASP7, in which the corresponding region forms a coil, not a β sheet (Figure 5-figure supplement 1). Structural modeling showed that the residues Q276, D278, and H283 are close to the region corresponding to the HsCASP1/4 exosite (Figure 5-figure supplement 1). Since these three residues are also not conserved between HsCASP7 p10 subunit and TrCASP7 p10 subunit, they were selected for mutagenesis to examine whether there existed an exosite interaction between HsCASP7 and HsGSDME. The mutation results showed that, similar to the wild type, the mutant Q276W&D278E&H283S was unable to cleave HsGSDME (Figure 5G), suggesting that, unlike human GSDMD, HsGSDME cleavage by CASPs probably did not involve exosite interaction. In contrast to HsGSDME, TrGSDME cleavage by HsCASP7 was not affected by the mutation of S234N, S234N plus S247N&I248V, or Q276W&D278E&H283S (Figure 5H).

All the above observed GSDME cleavages by CASP7 were verified in a cellular system. When co-expressed in HEK293T cells, HsGSDME was cleaved by TrCASP7 but not by HsCASP7, while TrGSDME was cleaved by both TrCASP7 and HsCASP7 (Figure 5-figure supplement 3). Additionally, both HsGSDME and TrGSDME were cleaved by HsCASP7-S234N as well as TrCASP7-N245S (Figure 5I and J). An apparent dose effect was observed in the cleavage of HsGSDME by HsCASP7-S234N (Figure 5K).

Divergent and GSDME-independent evolution of CASP7 leads to the loss of GSDME-activation function in mammalian CASP7.

Given the importance of N234 in p10 for HsCASP7 cleavage of HsGSDME, we analyzed the sequence conservation of this locus in CASP3/7. For CASP3, a highly conserved Asn residue (corresponding to HsCASP7 S234) was found immediately after the SWR motif in vertebrates (Figure 6A and B). In HsCASP3, this conserved Asn is

at position 208. We examined whether this residue was functionally essential to HsCASP3. Compared to the wild type, the N208S mutant exhibited much weaker cleavage of HsGSDME (Figure 6C, Figure 4-figure supplement 1D). Similar weakened cleavage was also observed in HEK293T cells co-expressing HsCASP3-N208S and HsGSDME (Figure 6D). These results indicated that the Asn residue was also critical for HsCASP3 cleavage of HsGSDME.

For CASP7, an Asn at the position corresponding to HsCASP7 S234 is highly conserved in teleost, amphibians, reptiles and birds, but not in mammals (Figure 6A). In mammals, the corresponding Asn is present in most non-primate species, such as mouse *Mus musculus* and bovine *Bos taurus*, but is replaced by Ser in primate, such as human *H. sapiens*, gorillas *Gorilla gorilla*, and chimpanzee *Pan troglodytes* (Figure 6E, Figure 6-figure supplement 1). We examined whether mouse CASP7 (MmCASP7) could cleave GSDME. The results showed that MmCASP7 cleaved HsGSDME in a dose-dependent manner, and the cleaving ability was abrogated by N234S mutation (Figure 6G and H, Figure 4-figure supplement 1E). By contrast, neither MmCASP7 nor MmCASP7-N234S was able to cleave mouse GSDME (MmGSDME) (Figure 6I). Mouse CASP3 (MmCASP3), however, cleaved MmGSDME efficiently (Figure 6-figure supplement 2). These results indicated the existence of an intra-species barrier for GSDME cleavage by CASP7 in some mammals, suggesting independent evolutions of CASP7 and GSDME.

Discussion

GSDME is an ancient member of the GSDM family existing ubiquitously in vertebrate from teleost to mammals (Broz, Pelegrin, & Shao, 2020; De Schutter et al., 2021). In human, GSDME is specifically cleaved by CASP3 at the consensus motif DMPD, but it is not cleaved by CASP7, which recognizes the same DxxD motif. CASP3 cleavage releases the pyroptosis-inducing NT fragment from the association of the inhibitory CT fragment and switches cell death from apoptosis to pyroptosis (Rogers et al., 2017; Wang et al., 2017). In this study, we found that different from human GSDME, a

pufferfish GSDME belonging to the teleost GSDMEa lineage was specifically cleaved by both CASP3 and CASP7 at the site of DAVD to liberate the pyroptosis-inducing NT fragment. In teleost, there generally exist two GSDME orthologs, named GSDMEa and GSDMEb. Comparative study on the functions of GSDMEa and GSDMEb is scarce, and it remains to be explored whether clear physiological roles are played by these two orthologs in fish. As executors of pyroptosis, teleost GSDMEs are activated via different mechanisms. In tongue sole *Cynoglossus semilaevis*, GSDMEb is preferably cleaved by CASP1 to trigger pyroptosis (Jiang, Gu, Zhao, & Sun, 2019); in zebrafish *Danio rerio*, GSDMEb is activated by caspy2 (CASP5 homolog) through the NLR family pyrin domain containing 3 (NLRP3) inflammasome signaling pathway (Li et al., 2020; Z. Wang et al., 2020). These observations indicate that similar to mammalian GSDMD, teleost GSDMEb is activated by inflammatory CASPs. In contrast, teleost GSDMEa is cleaved mainly by apoptotic CASPs. In zebrafish, GSDMEa is cleaved by CASP3 to generate the pyroptotic NT fragment (Wang et al., 2017). In turbot *Scophthalmus maximus*, GSDMEa is bi-directionally regulated by CASP3/7, which activate GSDMEa, and CASP6, which inactivates GSDMEa (Xu, Jiang, Yu, Yuan, & Sun, 2022). In common carp *Cyprinus carpio haematopterus*, GSDMEa is cleaved by CASP3 and induces pyroptosis (Zhao et al., 2022). The complicated scenario of GSDME-mediated pyroptosis signaling in fish is likely due to the reason that, unlike mammals that have multiple GSDM members to induce pyroptosis under different conditions, fish have only GSDME to induce pyroptotic cell death. Regulation by different CASPs may represent a mechanism that enables fish GSDME to execute the orders of different pyroptotic signals.

It is intriguing that, although HsCASP3 and 7 were indistinguishable in proteolysis towards the consensus tetrapeptide, they differed remarkably in the cleavage of HsGSDME and TrGSDME. HsCASP3, but not HsCASP7, cleaved HsGSDME, whereas both HsCASP3 and 7 cleaved TrGSDME at the same site. NT/CT domain swapping between HsGSDME and TrGSDME showed that chimeric GSDMEs containing the TrGSDME-CT domain were cleaved by human as well as pufferfish

CASP7 regardless of the source of the NT, whereas chimeric GSDMEs containing the HsGSDME-CT domain were resistant to HsCASP7. These results indicate that the CT domain, which is well accepted to exert an inhibitory effect on the pore-forming NT domain (Z. Liu et al., 2019; J. Shi et al., 2015), is the target of HsCASP7 discrimination, and hence determines the cleavability of GSDME by HsCASP7. Recently, Wang and Liu studied the molecular mechanism of human GSDMD recognition by CASP1/4, and showed that GSDMD-CT interacted with CASP1/4 exosites through binding to a hydrophobic pocket, which enhanced the recognition by CASP1/4 and contributed to tetrapeptide sequence-independent cleavage of GSDMD (Z. Liu et al., 2020; K. Wang et al., 2020). Different from GSDMD, we found that for TrGSDME, mutation of either the P4 (D255) or P1 (D258) residue of the consensus motif ²⁵⁵DAVD²⁵⁸ made it resistant to CASP3/7, implying a lack of DAVD-independent cleavage mechanism. Similarly, a previous study observed that human and mouse GSDME harboring cleavage site (P4 or P1) mutation resisted CASP3 cleavage (Wang et al., 2017). These results suggest that, compared to GSDMD, GSDME has a distinct enzyme-substrate engagement mode that, as demonstrated in the present study, involves GSDME-CT.

Unlike HsCASP7, which was unable to cleave HsGSDME, TrCASP7 effectively cleaved HsGSDME. The swapping of p20 between HsCASP7 and TrCASP7 revealed that the two catalytic motifs in p20, i.e., SHG and QACRG, were not involved in HsCASP7 discrimination on GSDME. SWR and GSWF are known to be responsible for CASP substrate binding (Chai et al., 2001; Riedl et al., 2001). However, in our study, we found that these two motifs are conserved in human and pufferfish CASP7, and they had no apparent effect on the substrate discrimination of HsCASP7. By contrast, the chimeric HsCASP7 containing the p10 subunit of TrCASP7 acquired the ability to cleave HsGSDME, indicating an important role of p10. By a series of residue swapping and mutation analyses, we discovered that the residue in the position of 245 (in TrCASP7) or 234 (in HsCASP7) of the p10 subunit was the key that determined CASP7 cleavage of HsGSDME. Sequence analysis revealed that N245 is highly conserved in the CASP7 of teleost, amphibians, reptiles and birds. In contrast, this Asn

residue was changed to Ser in the CASP7 of mammals, especially primates, resulting in divergence of CASP7 from CASP3 in substrate recognition and cleavage. Since, unlike non-mammalian vertebrates, mammals possess multiple pyroptotic GSDM members (GSDMA, B, C, D, and E), the functional divergence of CASP7 from CASP3 may have occurred as a result of specific and non-redundant regulation of programmed cell death mediated by different executor molecules. One of the physiological consequences is that CASP7 frees itself from the GSDME-mediated pathway and thus is able to be engaged in other cellular processes. Another physiological consequence is that GSDME activation is limited to CASP3 cleavage, thus restricting GSDME activity to situations more specific, such as that inducing CASP3 activation. More physiological consequences of CASP3/7 divergence in GSDME activation need to be explored in future studies. With the exception of primates, the conserved Asn is present in a large number of mammals, including mice. Mouse CASP7 was able to cleave HsGSDME, further supporting the importance of the Asn in CASP7 recognition and cleavage. The inability of mouse CASP7 to cleave mouse GSDME is consistent with a previous report that CASP7 deletion had little effect on mouse GSDME-mediated pyroptosis (Sarhan et al., 2018). These results suggest mutually independent evolution of GSDME and CASP7 in mice, which likely has distanced GSDME and CASP7 from each other. Similarly, human CASP7 and GSDME may also have undergone independent evolution, which leads to the disengagement of GSDME from the substrate relationship with CASP7.

In conclusion, we identified a teleost GSDME that can be cleaved by fish and human CASP3/7 to trigger pyroptosis. The GSDME-CT domain and the CASP7 p10 subunit are critical in the determination of CASP7 cleavage of GSDME. Within the p10 subunit, a single residue plays a key role in CASP7 substrate recognition and cleavage. Our results reveal the molecular basis of the functional divergence of CASP7 and CASP3, and suggest separate evolutions of CASP7 and GSDME in mammals. These findings add new insights into CASP-regulated GSDME activation in lower and higher vertebrates.

Materials and Methods

Animal, ethics, and cell line

Clinically healthy pufferfish (*Takifugu rubripes*) were obtained from a local fish farm. In the laboratory, the fish were maintained at 19-20°C in aerated seawater as reported previously (Xu et al., 2022). For euthanization, the fish were immersed in excess tricaine methane sulfonate (Sigma, St. Louis, MO). The animal experiments were approved by the Ethics Committee of institute of Oceanology, Chinese Academy of Sciences. HEK293T cells (ATCC, Rockville, MD, USA) were cultured in DMED medium (Corning, NY, USA) supplemented with 10% fetal bovine serum (ExCell Bio, Shanghai, China) at 37°C in a 5% CO₂ incubator.

Sequence analysis

The sequences of 742 CASP3 and 758 CASP7 used in this study were downloaded from NCBI Orthologs. TrCASP7 and HsCASP7 sequences were aligned with Clustal W program (www.ebi.ac.uk/clustalw/) and visualized with ESPript 3.0 (<http://esprict.ibcp.fr/ESPript/cgi-bin/ESPript.cgi/>) (Robert & Gouet, 2014). Conservation of S/N234 in CASP3/7 was analyzed via Weblogo3 (<https://weblogo.threeplusone.com/>) (Crooks, Hon, Chandonia, & Brenner, 2004), and the sequences of CASP3/7 used are shown in Table 1 and Table 2. The phylogenetic analysis of GSDME was performed as reported previously (Xu et al., 2022). The phylogenetic relationship of major mammalian clades was estimated based on the Mammals birth-death node-dated completed trees in VertLife (Upham, Esselstyn, & Jetz, 2019). The phylogenetic tree was subsequently viewed and edited in iTOL (Letunic & Bork, 2021). The icons representing phylogeny clades were retrieved from the PhyloPic (<http://www.phylopic.org/>), with the detailed credentials provided in Table 3.

Antibodies and immunoblotting

Monoclonal anti-mouse (ab215191) and anti-human (ab221843) GSDME antibodies were purchased from Abcam (Cambridge, MA). Antibodies against β -actin (AC026), Flag (AE005), Myc (AE010), His (AE003), and mCherry (AE002) were purchased from Abclonal (Wuhang, China). Mouse polyclonal antibody against TrGSDME was prepared as reported previously (Xu et al., 2022). Anti-serum (1:1000 dilution) was used for immunoblotting. Immunoblotting was performed as reported previously (Zhao & Sun, 2022). Briefly, the samples were fractionated in 12% SDS-polyacrylamide gels (GenScript, Piscataway, NJ, USA). The proteins were transferred to NC membranes, immunoblotted with appropriate antibodies and visualized using an ECL kit (Sparkjade Biotechnology Co. Ltd., Shandong, China).

Gene cloning and mutagenesis

Total RNA was extracted from pufferfish kidney using the FastPure Cell/Tissue Total RNA Isolation Kit V2 (Vazyme Biotech Co. Ltd., Nanjing, China). cDNA synthesis was performed with Revert Aid First Strand cDNA Synthesis Kit (Thermo Fisher Scientific, Waltham, MA, United States). The coding sequences (CDSs) of pufferfish GSDME (Accession number XP_029701077.1) and CASP3/7 were amplified by PCR. The CDSs of mouse GSDME and CASP3/7 were synthesized by Sangon Biotech (Shanghai, China). Site-directed mutagenesis was performed using Hieff Mut Site-Directed Mutagenesis Kit (Yeasen, Shanghai, China). Truncates of TrGSDME were created as reported previously (Xu et al., 2022) and subcloned into pmCherry-N1 (Clontech, Mountain View, CA, USA). For the construction of GSDME and CASP7 chimera, the CDSs of GSDME NT/CT and CASP7 p10/p20 were obtained by PCR and ligated. The constructs of the truncate and chimeric proteins are shown in Table 4. All sequences/constructs were verified by sequencing analysis. The primers used are listed in Table 5.

Plasmids and transient expression

For expression in mammalian cell expression, the CDSs of GSDME (pufferfish, human and mouse) were subcloned into pCS2-3×Flag (Wang et al., 2017) or pmCherry-N1 and CDSs encoding CASP (pufferfish, human and mouse) were subcloned into p-CMV-Myc (Clontech, Mountain View, CA, USA) or pCS2-Myc (Wang et al., 2017). All plasmids were prepared with endotoxin-free plasmid kit (Sparkjade Biotechnology Co. Ltd., Shandong, China). For transient expression, HEK293T cells were cultured in 96- or 24-well plates (Corning, NY, USA) overnight, and then transfected with 100 ng/well (96-well plates) or 500 ng/well (24-well plates) indicated plasmids using Liopfectamine 3000 (Invitrogen, USA) for 24 h or as specified. For GSDME processing, the plasmids of GSDME and CASP were co-transfected in HEK293T cells as described above and cultured continuously for 24 h or as specified. Cells were harvested and lysed with RIPA Lysis Buffer (Beyotime, Shanghai, China) for immunoblotting. The primer used are shown in Table 5.

Protein purification

Recombinant GSDMEs and CASPs are all soluble and were purified as described previously (Jiang, Zhou, Sun, Zhang, & Sun, 2020; Xu et al., 2022). Briefly, the CDSs of TrGSDME and CASP variants were each cloned into pET30a (+), and the CDSs of HsGSDME, MmGSDME, and chimeric GSDME were each cloned into pET28a-SUMO. The recombinant plasmids were introduced into *Escherichia coli* Transetta (DE3) (TransGen, Beijing, China) by transformation. The transformants were cultured in Luria broth (LB) at 37°C until logarithmic growth phase. Isopropyl-β-D-thiogalactopyranoside (0.3 mM) was added to the medium, followed by incubation at 16°C for 20 h. Bacteria were harvested and lysed, and the supernatant was collected for protein purification with Ni-NTA columns (GE Healthcare, Uppsala, Sweden). The proteins were dialyzed with PBS at 4 °C and concentrated.

CASP activity

To measure their proteolytic activity, recombinant CASPs were incubated separately with various colorimetric substrates at 37°C for 2 h as described previously (Jiang et al., 2020; Xu et al., 2022), and then monitored for released pNA at OD405. To compare their substrate preference, TrCASP3/7 and HsCASP3/7 were incubated with Ac-DAVD-pNA or Ac-DMPD-pNA (Science Peptide Biological Technology Co., Ltd, shanghai, China) at 37°C for 30 min, and the released pNA was measured at OD405. The CASP activity in cells transfected TrCASP3/7 was determined as described previously (Xu et al., 2022).

GSDME cleavage by CASPs

GSDME cleavage assay was performed as described previously (Jiang et al., 2020; Xu et al., 2022). Briefly, recombinant GSDME was incubated with 1 U of recombinant HsCASP1, 2, 3, 6, 7, 8 and 9 (Enzo Life Sciences, Villeurbanne, France) at 37°C for 2 h in the reaction buffer (50 mM Hepes (pH 7.5), 3 mM EDTA, 150 mM NaCl, 0.005% (v/v) Tween 20, and 10 mM DTT). TrGSDME, HsGSDME or their chimeras were treated with 1 U of TrCASP3/7, HsCASP3/7, or their chimeras for 2 h as above in reaction buffer. After incubation, the samples were boiled in 5 x Loading Buffer (GenScript, Piscataway, NJ, USA) and subjected to SDS-PAGE or immunoblotting with indicated antibody.

Cell death examination

Cell death examination by microscopy was performed as reported previously (Jiang et al., 2020; Xu et al., 2022). Briefly, HEK293T cells were plated into 35 mm glass-bottom culture dishes (Nest Biotechnology, Wuxi, China) at about 60% confluency and subjected to the indicated treatment for 24 h. To examine cell death morphology, propidium iodide (PI) (Invitrogen, Carlsbad, CA, USA) and DiO (Solarbio, Beijing, China) were added to the culture medium, and the cells were observed with a Carl Zeiss LSM 710 confocal microscope (Carl Zeiss, Jena, Germany). To video the cell death process, the cells were transfected with 1 µg pmCherry-N1 vector expressing

TrGSDME-NT for 16 h, and cell death was recorded using the above microscope. Cell death measured by lactate dehydrogenase (LDH) assay was performed as reported previously (Xu et al., 2022).

Statistical analysis

The Student's t test and one-way analysis of variance (ANOVA) were used for comparisons between groups. Statistical analysis was performed with GraphPad Prism 7 software. Statistical significance was defined as $P < 0.05$.

Acknowledgments

We thank Prof. Feng Shao (National institute of Biological Sciences, Beijing, China) for providing human CASP3/7 and GSDME expression vectors. This work was supported by the grants from the Science & Technology Innovation Project of Laoshan Laboratory (LSKJ202203000), the Youth Innovation Promotion Association CAS (2021204), the National Natural Science Foundation of China (41876175), and the Taishan Scholar Program of Shandong Province (2018 and 2021).

Author Contributions

Hang Xu: Methodology; Investigation; Visualization; Writing—original draft. ;**Zihao Yuan:** Investigation. **Kunpeng Qin:** Investigation; **Shuai Jiang:** Conceptualization; Methodology; Supervision; Writing—review & editing. **Li Sun:** Conceptualization; Supervision; Writing—review & editing.

Competing Interest Statement

The authors declare that they have no competing interests.

References

- Agniswamy, J., Fang, B., & Weber, I. T. (2009). Conformational similarity in the activation of caspase-3 and -7 revealed by the unliganded and inhibited structures of caspase-7. *Apoptosis*, 14(10), 1135-1144. doi:10.1007/s10495-009-0388-9
- Angosto-Bazarra, D., Alarcon-Vila, C., Hurtado-Navarro, L., Banos, M. C., Rivers-Auty, J., & Pelegrin, P. (2022). Evolutionary analyses of the gasdermin family suggest conserved roles in infection response despite loss of pore-forming functionality. *BMC Biol*, 20(1), 9. doi:10.1186/s12915-021-01220-z
- Bergsbaken, T., Fink, S. L., & Cookson, B. T. (2009). Pyroptosis: host cell death and inflammation. *Nat Rev Microbiol*, 7(2), 99-109. doi:10.1038/nrmicro2070
- Boyce, M., Degterev, A., & Yuan, J. (2004). Caspases: an ancient cellular sword of Damocles. *Cell Death Differ*, 11(1), 29-37. doi:10.1038/sj.cdd.4401339
- Brentnall, M., Rodriguez-Menocal, L., De Guevara, R. L., Cepero, E., & Boise, L. H. (2013). Caspase-9, caspase-3 and caspase-7 have distinct roles during intrinsic apoptosis. *BMC Cell Biol*, 14, 32. doi:10.1186/1471-2121-14-32
- Broz, P., Pelegrin, P., & Shao, F. (2020). The gasdermins, a protein family executing cell death and inflammation. *Nat Rev Immunol*, 20(3), 143-157. doi:10.1038/s41577-019-0228-2
- Budd, S. L., Tenneti, L., Lishnak, T., & Lipton, S. A. (2000). Mitochondrial and extramitochondrial apoptotic signaling pathways in cerebrocortical neurons. *Proc Natl Acad Sci U S A*, 97(11), 6161-6166. doi:10.1073/pnas.100121097
- Chai, J., Shiozaki, E., Srinivasula, S. M., Wu, Q., Datta, P., Alnemri, E. S., & Shi, Y. (2001). Structural basis of caspase-7 inhibition by XIAP. *Cell*, 104(5), 769-780. doi:10.1016/s0092-8674(01)00272-0
- Cohen, G. M. (1997). Caspases: the executioners of apoptosis. *Biochem J*, 326 (Pt 1), 1-16. doi:10.1042/bj3260001

- Crawford, E. D., & Wells, J. A. (2011). Caspase substrates and cellular remodeling. *Annu Rev Biochem*, 80, 1055-1087. doi:10.1146/annurev-biochem-061809-121639
- Crooks, G. E., Hon, G., Chandonia, J. M., & Brenner, S. E. (2004). WebLogo: a sequence logo generator. *Genome Res*, 14(6), 1188-1190. doi:10.1101/gr.849004
- De Schutter, E., Roelandt, R., Riquet, F. B., Van Camp, G., Wullaert, A., & Vandenabeele, P. (2021). Punching Holes in Cellular Membranes: Biology and Evolution of Gasdermins. *Trends Cell Biol*, 31(6), 500-513. doi:10.1016/j.tcb.2021.03.004
- Demon, D., Van Damme, P., Vanden Berghe, T., Deceuninck, A., Van Durme, J., Verspurten, J., . . . Vandenabeele, P. (2009). Proteome-wide substrate analysis indicates substrate exclusion as a mechanism to generate caspase-7 versus caspase-3 specificity. *Mol Cell Proteomics*, 8(12), 2700-2714. doi:10.1074/mcp.M900310-MCP200
- Erener, S., Petrilli, V., Kassner, I., Minotti, R., Castillo, R., Santoro, R., . . . Hottiger, M. O. (2012). Inflammasome-activated caspase 7 cleaves PARP1 to enhance the expression of a subset of NF-kappaB target genes. *Mol Cell*, 46(2), 200-211. doi:10.1016/j.molcel.2012.02.016
- He, W. T., Wan, H., Hu, L., Chen, P., Wang, X., Huang, Z., . . . Han, J. (2015). Gasdermin D is an executor of pyroptosis and required for interleukin-1beta secretion. *Cell Res*, 25(12), 1285-1298. doi:10.1038/cr.2015.139
- Jiang, S., Gu, H., Zhao, Y., & Sun, L. (2019). Teleost Gasdermin E Is Cleaved by Caspase 1, 3, and 7 and Induces Pyroptosis. *J Immunol*, 203(5), 1369-1382. doi:10.4049/jimmunol.1900383
- Jiang, S., Zhou, Z., Sun, Y., Zhang, T., & Sun, L. (2020). Coral gasdermin triggers pyroptosis. *Sci Immunol*, 5(54). doi:10.1126/sciimmunol.abd2591
- Kayagaki, N., Stowe, I. B., Lee, B. L., O'Rourke, K., Anderson, K., Warming, S., . . . Dixit, V. M. (2015). Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. *Nature*, 526(7575), 666-671. doi:10.1038/nature15541

553 Kovacs, S. B., & Miao, E. A. (2017). Gasdermins: Effectors of Pyroptosis. *Trends Cell*
554 *Biol*, 27(9), 673-684. doi:10.1016/j.tcb.2017.05.005

555 Kuang, S., Zheng, J., Yang, H., Li, S., Duan, S., Shen, Y., . . . Li, J. (2017). Structure
556 insight of GSDMD reveals the basis of GSDMD autoinhibition in cell pyroptosis.
557 *Proc Natl Acad Sci U S A*, 114(40), 10642-10647. doi:10.1073/pnas.1708194114

558 Kumar, S. (2007). Caspase function in programmed cell death. *Cell Death Differ*, 14(1),
559 32-43. doi:10.1038/sj.cdd.4402060

560 Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: an online tool for
561 phylogenetic tree display and annotation. *Nucleic Acids Res*, 49(W1), W293-
562 W296. doi:10.1093/nar/gkab301

563 Li, J. Y., Wang, Y. Y., Shao, T., Fan, D. D., Lin, A. F., Xiang, L. X., & Shao, J. Z.
564 (2020). The zebrafish NLRP3 inflammasome has functional roles in ASC-
565 dependent interleukin-1 β maturation and gasdermin E-mediated pyroptosis. *J*
566 *Biol Chem*, 295(4), 1120-1141. doi:10.1074/jbc.RA119.011751

567 Liu, X., Zhang, Z., Ruan, J., Pan, Y., Magupalli, V. G., Wu, H., & Lieberman, J. (2016).
568 Inflammasome-activated gasdermin D causes pyroptosis by forming membrane
569 pores. *Nature*, 535(7610), 153-158. doi:10.1038/nature18629

570 Liu, Z., Wang, C., Yang, J., Chen, Y., Zhou, B., Abbott, D. W., & Xiao, T. S. (2020).
571 Caspase-1 Engages Full-Length Gasdermin D through Two Distinct Interfaces
572 That Mediate Caspase Recruitment and Substrate Cleavage. *Immunity*, 53(1), 106-
573 114 e105. doi:10.1016/j.immuni.2020.06.007

574 Liu, Z., Wang, C., Yang, J., Zhou, B., Yang, R., Ramachandran, R., . . . Xiao, T. S.
575 (2019). Crystal Structures of the Full-Length Murine and Human Gasdermin D
576 Reveal Mechanisms of Autoinhibition, Lipid Binding, and Oligomerization.
577 *Immunity*, 51(1), 43-49 e44. doi:10.1016/j.immuni.2019.04.017

578 Poreba, M., Strozyk, A., Salvesen, G. S., & Drag, M. (2013). Caspase Substrates and
579 Inhibitors. *Cold Spring Harbor Perspectives in Biology*, 5(8), a008680-a008680.
580 doi:10.1101/cshperspect.a008680

- Riedl, S. J., Renatus, M., Schwarzenbacher, R., Zhou, Q., Sun, C., Fesik, S. W., . . .
Salvesen, G. S. (2001). Structural basis for the inhibition of caspase-3 by XIAP.
Cell, 104(5), 791-800. doi:10.1016/s0092-8674(01)00274-4
- Robert, X., & Gouet, P. (2014). Deciphering key features in protein structures with the
new ENDscript server. *Nucleic Acids Res*, 42(Web Server issue), W320-324.
doi:10.1093/nar/gku316
- Rogers, C., Fernandes-Alnemri, T., Mayes, L., Alnemri, D., Cingolani, G., & Alnemri,
E. S. (2017). Cleavage of DFNA5 by caspase-3 during apoptosis mediates
progression to secondary necrotic/pyroptotic cell death. *Nat Commun*, 8, 14128.
doi:10.1038/ncomms14128
- Sarhan, J., Liu, B. C., Muendlein, H. I., Li, P., Nilson, R., Tang, A. Y., . . . Poltorak, A.
(2018). Caspase-8 induces cleavage of gasdermin D to elicit pyroptosis during
Yersinia infection. *Proc Natl Acad Sci U S A*, 115(46), E10888-E10897.
doi:10.1073/pnas.1809548115
- Shao, F. (2021). Gasdermins: making pores for pyroptosis. *Nat Rev Immunol*, 21(10),
620-621. doi:10.1038/s41577-021-00602-2
- Shi, J., Gao, W., & Shao, F. (2017). Pyroptosis: Gasdermin-Mediated Programmed
Necrotic Cell Death. *Trends Biochem Sci*, 42(4), 245-254.
doi:10.1016/j.tibs.2016.10.004
- Shi, J., Zhao, Y., Wang, K., Shi, X., Wang, Y., Huang, H., . . . Shao, F. (2015). Cleavage
of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature*,
526(7575), 660-665. doi:10.1038/nature15514
- Shi, Y. (2002). Mechanisms of caspase activation and inhibition during apoptosis. *Mol*
Cell, 9(3), 459-470. doi:10.1016/s1097-2765(02)00482-3
- Slee, E. A., Adrain, C., & Martin, S. J. (2001). Executioner caspase-3, -6, and -7
perform distinct, non-redundant roles during the demolition phase of apoptosis. *J*
Biol Chem, 276(10), 7320-7326. doi:10.1074/jbc.M008363200
- Tamura, M., Tanaka, S., Fujii, T., Aoki, A., Komiyama, H., Ezawa, K., . . . Shiroishi,
T. (2007). Members of a novel gene family, Gsdm, are expressed exclusively in

the epithelium of the skin and gastrointestinal tract in a highly tissue-specific manner. *Genomics*, 89(5), 618-629. doi:10.1016/j.ygeno.2007.01.003

Timmer, J. C., & Salvesen, G. S. (2007). Caspase substrates. *Cell Death Differ*, 14(1), 66-72. doi:10.1038/sj.cdd.4402059

Tsuchiya, K., Nakajima, S., Hosojima, S., Thi Nguyen, D., Hattori, T., Manh Le, T., . . . Suda, T. (2019). Caspase-1 initiates apoptosis in the absence of gasdermin D. *Nat Commun*, 10(1), 2091. doi:10.1038/s41467-019-09753-2

Upham, N. S., Esselstyn, J. A., & Jetz, W. (2019). Inferring the mammal tree: Species-level sets of phylogenies for questions in ecology, evolution, and conservation. *PLoS Biol*, 17(12), e3000494. doi:10.1371/journal.pbio.3000494

Wajant, H. (2002). The Fas signaling pathway: more than a paradigm. *Science*, 296(5573), 1635-1636. doi:10.1126/science.1071553

Walsh, J. G., Cullen, S. P., Sheridan, C., Luthi, A. U., Gerner, C., & Martin, S. J. (2008). Executioner caspase-3 and caspase-7 are functionally distinct proteases. *Proc Natl Acad Sci U S A*, 105(35), 12815-12819. doi:10.1073/pnas.0707715105

Wang, K., Sun, Q., Zhong, X., Zeng, M., Zeng, H., Shi, X., . . . Ding, J. (2020). Structural Mechanism for GSDMD Targeting by Autoprocessed Caspases in Pyroptosis. *Cell*, 180(5), 941-955 e920. doi:10.1016/j.cell.2020.02.002

Wang, Y., Gao, W., Shi, X., Ding, J., Liu, W., He, H., . . . Shao, F. (2017). Chemotherapy drugs induce pyroptosis through caspase-3 cleavage of a gasdermin. *Nature*, 547(7661), 99-103. doi:10.1038/nature22393

Wang, Z., Gu, Z., Hou, Q., Chen, W., Mu, D., Zhang, Y., . . . Yang, D. (2020). Zebrafish GSDMEb Cleavage-Gated Pyroptosis Drives Septic Acute Kidney Injury In Vivo. *J Immunol*, 204(7), 1929-1942. doi:10.4049/jimmunol.1901456

Wolf, B. B., Schuler, M., Echeverri, F., & Green, D. R. (1999). Caspase-3 is the primary activator of apoptotic DNA fragmentation via DNA fragmentation factor-45/inhibitor of caspase-activated DNase inactivation. *J Biol Chem*, 274(43), 30651-30656. doi:10.1074/jbc.274.43.30651

- Xia, S., Zhang, Z., Magupalli, V. G., Pablo, J. L., Dong, Y., Vora, S. M., . . . Wu, H. (2021). Gasdermin D pore structure reveals preferential release of mature interleukin-1. *Nature*. doi:10.1038/s41586-021-03478-3
- Xu, H., Jiang, S., Yu, C., Yuan, Z., & Sun, L. (2022). GSDMEa-mediated pyroptosis is bi-directionally regulated by caspase and required for effective bacterial clearance in teleost. *Cell Death Dis*, 13(5), 491. doi:10.1038/s41419-022-04896-5
- Yuan, Z., Jiang, S., Qin, K., & Sun, L. (2022). New insights into the evolutionary dynamic and lineage divergence of gasdermin E in metazoa. *Front Cell Dev Biol*, 10, 952015. doi:10.3389/fcell.2022.952015
- Zhao, Y., & Sun, L. (2022). Bacillus cereus cytotoxin K triggers gasdermin D-dependent pyroptosis. *Cell Death Discov*, 8(1), 305. doi:10.1038/s41420-022-01091-5
- Zhao, Y., Zhang, J., Qiao, D., Gao, F., Gu, Y., Jiang, X., . . . Kong, X. (2022). CcGSDMEa functions the pore-formation in cytomembrane and the regulation on the secretion of IL-1beta in common carp (Cyprinus carpio haematopterus). *Front Immunol*, 13, 1110322. doi:10.3389/fimmu.2022.1110322
- Zheng, T. S., Schlosser, S. F., Dao, T., Hingorani, R., Crispe, I. N., Boyer, J. L., & Flavell, R. A. (1998). Caspase-3 controls both cytoplasmic and nuclear events associated with Fas-mediated apoptosis in vivo. *Proc Natl Acad Sci U S A*, 95(23), 13618-13623. doi:10.1073/pnas.95.23.13618

Figure legends

Figure 1. Cleavage of TrGSDME by caspases. (A, B) TrGSDME was treated with 1 U of different HsCASP3/7 for 2 h and then subjected to SDS-PAGE (A) and immunoblotting (B). (C) SDS-PAGE analysis of purified TrCASP3/7. The p10 and p20 subunits are indicated. (D) TrCASP3/7 cleavage of different colorimetric CASP substrates was monitored by measuring released pNA. The values are the means $\pm$ SD of three replicates. (E) TrCASP3/7 (0.25-8U) were incubated with Ac-DEVD-pNA, and time-dependent release of pNA was measured. (F, G) SDS-PAGE (F) and immunoblotting (G) analysis of TrGSDME cleavage by TrCASP3/7 (0.25-4U). (H) TrGSDME cleavage by TrCASP3/7 was determined in the presence or absence of 20 μ M Z-DEVD-FMK or Z-VAD-FMK. (I, J) TrGSDME wild type (WT) and mutants (D255R or D258A) were incubated with TrCASP3/7 for 2 h, and the cleavage was determined by SDS-PAGE (I) and immunoblotting (J). (K) A schematic of TrGSDME cleavage by TrCASP3/7. The arrow indicates cleavage site. For all panels, FL, full length; NT, N-terminal fragment; CT, C-terminal fragment.

Figure 2. The pyroptotic activity of TrGSDME. (A-C) HEK293T cells were transfected with C-terminally mCherry-tagged TrGSDME full length (FL) or truncate (NT or CT) for 24 h, and then analyzed for TrGSDME expression (A), morphological change (B), and LDH release (C). Scale bars, 50 μ m (A) and 20 μ m (B). (D) HEK293T cells were transfected with the backbone vector or vector expressing Myc-tagged TrGSDME-NT for 24 h. The cell nuclei and membrane were stained with PI and DiO, respectively. Scale bar, 10 μ m. (E) HEK293T cells were transfected with mCherry-tagged TrGSDME-NT for 16 h, and the progression of cell death was shown by time-lapse microscopic imaging. Scale bar, 10 μ m. (F) HEK293T cells were transfected with TrCASP3/7 for 24 h, and the proteolytic activity of TrCASP3/7 was assessed by treatment with fluorogenic Ac-DEVD-AFC. (G) Phase-contrast images of HEK293T cells transfected with the indicated vectors for 24 h. Scale bar, 20 μ m. (H, I) LDH release (H) and TrGSDME cleavage and TrCASP3/7 expression (I) in the above

transfected cells were determined. For panel (I), FL, full-length; CT, C-terminal fragment. **(J, K)** HEK293T cells co-expressing TrGSDME and TrCASP3/7 for 24 h in the presence or absence (control) of 10 μ M Z-VAD-FMK or Z-DEVD-FMK were subjected to microscopy (J) and LDH measurement (K). Scale bar, 20 μ m. **(L, M)** HEK293T cells expressing TrCASP3/7 plus TrGSDME or TrCASP3/7 plus the D255R/D258A mutant were subjected to microscopy (L) and LDH measurement (M). Scale bar, 20 μ m. For panels (C, F, H, K and M), values are the means of three experimental replicates and shown as means $\pm$ SD. $**P < 0.01$, $***P < 0.001$.

Figure 3. GSDME-CT domain determines the recognition by CASP7. **(A)** HsGSDME was treated with 1 U of various HsCASPs for 2 h, and the products were analyzed by immunoblotting with anti-HsGSDME-CT antibody. **(B)** HsCASP3/7 and TrCASP3/7 were incubated with Ac-DAVD-pNA or Ac-DMPD-pNA at 37°C for 30 min, and the proteolytic activity was determined. Data are expressed as the means $\pm$ SD of three replicates. **(C)** Structure analysis of the linker region from HsGSDME and TrGSDME with mouse GSDMA3 (PDB:5b5r) as the template. Two regions that may form α -helix and loop are indicated. **(D)** HsGSDME wild type (WT) and mutants (Δ 261-266 and Δ 281-296) were treated with or without 1 U of HsCASP3/7 for 2 h, and the cleaved fragments were subjected to immunoblotting with anti-HsGSDME-CT antibody. **(E)** Schematics of the chimeric GSDME constructs. **(F-I)** TrGSDME (F), HsNT-TrCT (G), HsGSDME (H) and TrNT-HsCT (I) were incubated with 1 U of TrCASP3/7 or HsCASP3/7 for 2 h, and the cleavage was determined by immunoblotting with anti-TrGSDME or anti-HsGSDME-CT antibodies. For panels (A, D, and F-I): FL, full length; NT, N-terminal fragment; CT, C-terminal fragment.

Figure 4. The p10 subunit of CASP7 is essential for discrimination on GSDME. **(A)** Sequence alignment of HsCASP7 and TrCASP7 with HsCASP7 (PDB:1K86) as the template. α -helices and β -strands are indicated. The motifs involved in catalytic reaction (SHG and QACRG) and substrate binding (SWR and GSWF) are boxed in red. **(B)** Schematics of CASP7 chimeras. **(C)** The proteolytic activities of TrCASP7, HsCASP7 and their chimeras were determined by cleaving Ac-DAVD-pNA and Ac-

DMPD-pNA. Data are expressed as the means $\pm$ SD of three replicates. **(D-G)** TrGSDME (D), HsNT-TrCT (E), HsGSDME (F) or TrNT-HsCT (G) were incubated with TrCASP7, HsCASP7 and their chimeras for 2 h. The products were assessed by immunoblotting with anti-TrGSDME or anti-HsGSDME-CT antibodies. For panels (D-G): FL, full length; NT, N-terminal fragment; CT, C-terminal fragment.

Figure 5. Functional importance of the non-conserved residues in the p10 of HsCASP7 and TrCASP7. **(A)** Sequence alignment of the p10 region in HsCASP7 and TrCASP7. The arrows indicate non-conserved residues. **(B, C)** HsGSDME was treated with wild type (WT) or mutant HsCASP7 (B) or TrCASP7 (C) for 2 h. The cleavage was determined by immunoblotting with anti-HsGSDME-CT antibody. 205TME, HsCASP7 with TME insertion at position 205; 213 Δ TME, TrCASP7 with TME deleted at position 213. **(D, E)** TrGSDME was incubated with wild type (WT) or mutant HsCASP7 (D) or TrCASP7 (E) for 2 h. The cleavage was determined by immunoblotting with anti-TrGSDME antibody. **(F)** HsGSDME was incubated with HsCASP7 or the S234N mutant (0.25-4U) for 2 h. HsGSDME cleavage was analyzed as above. **(G, H)** HsGSDME (G) and TrGSDME (H) were treated with wild type or mutant HsCASP7 for 2 h. The cleavage was determined as above. **(I, J)** TrGSDME (I) or HsGSDME (J) was co-expressed with wild type or mutant TrCASP7/HsCASP7 for 24 h. GSDME cleavage, CASP7 and β -actin were determined as above. **(K)** HsGSDME was co-expressed with HsCASP7 or HsCASP7-S234N for 48 h. Cleavage of GSDME was detected as above. For panels (B-K): FL, full length; NT, N-terminal fragment; CT, C-terminal fragment.

Figure 6. Conservation and functional importance of S/N234 in CASP3/7. **(A)** Weblogo analysis of the conservation of S/N234 in CASP3/7 in Mammalia, Aves, Reptilia, Amphibia and Osteichthyes. **(B)** Sequence alignment of HsCASP3/7. The S/N residues are indicated by red arrow. **(C)** HsGSDME was treated with HsCASP3 or HsCASP3-N208S for 2 h, and then subjected to immunoblotting with anti-HsGSDME-CT antibody. **(D)** HsGSDME was co-expressed in HEK293T cells with different doses of HsCASP3 or HsCASP3-N208S for 24 h. The cell lysates were immunoblotted to

detect HsGSDME, CASP3, and β -actin. **(E)** Sequence alignment of the S/N234 region in the CASP7 of primate (shaded cyan) and non-primate mammals. The S/N residues are indicated by red arrow. Asterisks indicate identical residues. **(F)** Alignment of human and mouse CASP7 sequences. The S/N234 residues are indicated by arrow. **(G)** HsGSDME was treated with different units of HsCASP7 or MmCASP7 for 2 h, and the cleavage was assessed by immunoblotting with anti-HsGSDME-CT antibody. **(H)** HsGSDME was incubated with different units of MmCASP7 or its mutant (N234S) for 2 h, and HsGSDME cleavage was analyzed as above. **(I)** MmCASP7 was treated with HsCASP7, MmCASP7 or MmCASP7-N234S for 2 h. The cleavage was determined by immunoblotting with anti-MmGSDME-NT antibody. For panels (C, D, and G-I), FL, full length; NT, N-terminal fragment; CT, C-terminal fragment.

Figure 1-figure supplement 1. Phylogenetic analysis of the TrGSDME used in this study. The phylogenetic tree was constructed using 15 fish GSDMEa (red) and 51 fish GSDMEb (blue) collected from NCBI databank. The TrGSDME used in this study is indicated by a red star.

Figure 1-figure supplement 2. Cleavage of TrGSDME mutants by HsCASP3/7. TrGSDME wild type (WT) and mutants (D255R and D258A) were treated with or without (-) HsCASP3/7 for 2 h, and the cleavage was determined by immunoblotting with anti-His antibody. FL, full length; CT, cleaved C-terminal region.

Figure 2-figure supplement 1. Ectopic expression of TrGSDME-FL and truncates. HEK293T cells were transfected with the backbone vector or the vector expressing mCherry-tagged TrGSDME full length (FL) or truncates (NT or CT) for 24 h. The cell lysate was then immunoblotted with antibody against mCherry or β -actin.

Figure 2-figure supplement 2. The cell death-inducing ability of TrGSDME-FL and truncates. HEK293T cells were transfected with the backbone vector or the vector expressing mCherry-tagged TrGSDME full length (FL) or truncate (NT or CT) for 24 h. The cells were then stained with Sytox Green and examined with a microscope. Scale bar, 30 μ m.

Figure 2-figure supplement 3. Activation of TrGSDME-mediated pyroptosis by TrCASP3/7. (A) HEK293T cells were transfected with the backbone vector or the vector expressing the indicated protein for 24 h and then stained with Sytox Green. Scale bars, 30 μ m. **(B, C)** HEK293T cells expressing wild type (WT) or mutant TrGSDME together with or without (control) TrCASP3 or TrCASP7 were subjected to Sytox Green staining (B) and TrGSDME cleavage analysis by immunoblot with antibodies against mCherry (upper), Myc (middle), and actin (lower) (C). Scale bar, 30 μ m.

Figure 3-figure supplement 1. Cleavage of HsGSDME by caspases. Recombinant HsGSDME was incubated with or without (-) HsCASP1, 2, 3, 6, 7, 8, and 9 for 2 h, and the cleavage was determined by SDS-PAGE. M, molecular markers. FL, full length; NT, N-terminal fragment; CT, C-terminal fragment.

Figure 3-figure supplement 2. Proteolytic activity of TrCASP1. TrCASP1 was incubated with colorimetric Ac-YVAD-pNA, Ac-DAVD-pNA, and Ac-DMPD-pNA for 1 h, and the cleavage was determined by measuring the release of pNA. Values are the means of triplicate experiments and shown as means $\pm$ SD..

Figure 4-figure supplement 1. Preparation of recombinant proteins. (A) SDS-PAGE analysis of purified Hsp20-Trp10 and Trp20-Hsp10. **(B, C)** Purified TrCASP7 (B) and HsCASP7 (C) were subjected to SDS-PAGE (upper panel) and immunoblotting with anti-His tag antibody (lower panel). **(D)** HsCASP3 and its mutant (N208S) were analyzed by SDS-PAGE. **(E)** SDS-PAGE analysis of purified MmCASP7 and its mutant (N234S). In all panels, the p10 and p20 subunits are indicated.

Figure 5-figure supplement 1. Cleavage of poly (ADP-ribose) polymerase 1 (PARP1) and gelsolin by HsCASP7. HEK293T cells were transfected with C-terminal Flag-tagged PARP1 (left panel) or gelsolin (right panel) for 24 h. The cell lysates were incubated with or without (-) different units of HsCASP7 and HsCASP7-S234N for 2 h and then subjected to immunoblotting with anti-Flag antibody.

Figure 5-figure supplement 2. Structural analysis of HsCASP1/4/7. (A) The structures of HsCASP1 (PDB: 6kn0, gray), HsCASP4 (PDB: 6kmz, green), and HsCASP7 (PDB: 6kn0, purple) are superimposed. The β III/ β III' sheets of HsCASP1/4 are indicated by arrows; the residues Q276, D278, and H283 of HsCASP7 are shown in cyan. **(B)** The sequence alignment of HsCASP1/4 β III/ β III' sheet with the corresponding region of HsCASP7.

Figure 5-figure supplement 3. GSDME cleavage by CASP7 in cellular system. TrGSDME (A) or HsGSDME (B) was co-expressed with different doses of TrCASP7/HsCASP7 for 48 h in HEK293T cells. The expression/cleavage of

TrGSDME/HsGSDME and the expression of TrCASP7/HsCASP7 were determined by immunoblotting. β -actin was used as a loading control.

Figure 6-figure supplement 1. Conservation of S/N234 in mammalian CASP7. The four conserved amino acids are shown in LOGOs, with height representing the relative proportion of the corresponding amino acid at that position. The S/N234 is presented as the fourth LOGO and indicated with a red star.

Figure 6-figure supplement 2. The cleavage of MmGSDME by MmCASP3. (A) SDS-PAGE analysis of purified MmCASP3. **(B)** MmGSDME was treated with 1U of MmCASP3 for 2 h and then subjected to SDS-PAGE.

Legends for Figure 1-video 1:

Figure 1-video 1 (separate file). TrGSDME-NT induces pyroptosis of HEK293T cells. HEK293T cells were transfected with the vector expressing mCherry-tagged TrGSDME-NT, and time-lapse images of the cells were recorded with a confocal microscope.

Tables

Table 1. The CASP3 sequences used for WebLogo analysis in this study.

Table 2. The CASP7 sequences used for WebLogo analysis in this study.

Table 3. The credits for the pictures used in this study.

Table 4. The constructs of the truncate and chimeric proteins.

Table 5. Primers used in this study.

Table 1. The CASP3 sequences used for WebLogo analysis in this study.

Class	Species	Accession Number
Mammalia	<i>Homo sapiens</i>	P42574
	<i>Mus musculus</i>	P70677
	<i>Sus scrofa</i>	Q95ND5
	<i>Canis lupus familiaris</i>	Q8MKI5
	<i>Saimiri boliviensis</i>	NP_001266895.1
	<i>Equus caballus</i>	NP_001157433.1
	<i>Mesocricetus auratus</i>	Q60431
	<i>Pan troglodytes</i>	XP_009446811.1
	<i>Ovis aries</i>	XP_014960045.1
	<i>Capra hircus</i>	NP_001273018.1
	<i>Pan paniscus</i>	XP_003823509.1
	<i>Ailuropoda melanoleuca</i>	XP_002914062.1
	<i>Oryctolagus cuniculus</i>	Q8MJC3
	<i>Loxodonta africana</i>	XP_023410784.1
	<i>Bos taurus</i>	Q08DY9
	<i>Macaca fascicularis</i>	Q2PFV2
	<i>Delphinapterus leucas</i>	XP_022426661.1
	<i>Aotus nancymaae</i>	XP_012327995.1
	<i>Vulpes vulpes</i>	XP_025874057.1

	<i>Heterocephalus glaber</i>	EHB15384.1
	<i>Bos mutus grunniens</i>	XP_010834058.1
	<i>Balaenoptera acutorostrata</i>	XP_007177921.2
	<i>Odobenus rosmarus divergens</i>	XP_012422221.1
	<i>Tursiops truncatus</i>	XP_033704658.1
	<i>Saimiri boliviensis</i>	AAV74267.1
	<i>Theropithecus gelada</i>	XP_025242051.1
	<i>Rhinopithecus roxellana</i>	XP_010376231.2
	<i>Chlorocebus sabaeus</i>	XP_007998582.1
	<i>Arvicanthis niloticus</i>	XP_034376775.1
	<i>Castor canadensis</i>	JAV41953.1
	<i>Urocitellus parryii</i>	XP_026256383.1
	<i>Neogale vison</i>	XP_044081570.1
	<i>Lynx rufus</i>	XP_046947013.1
	<i>Canis lupus dingo</i>	XP_025327613.1
	<i>Balaenoptera acutorostrata scammoni</i>	XP_007177921.1
	<i>Balaenoptera musculus</i>	XP_036694636.1
	<i>Vicugna pacos</i>	XP_006198038.1
	<i>Camelus bactrianus</i>	XP_010951604.1
	<i>Eptesicus fuscus</i>	XP_008150114.1
	<i>Pteropus alecto</i>	XP_006916432.1
	<i>Ochotona princeps</i>	XP_004579072.1
	<i>Equus quagga</i>	XP_046504640.1
	<i>Talpa occidentalis</i>	XP_037382298.1
	<i>Manis pentadactyla</i>	XP_036750218.1
	<i>Gracilinanus agilis</i>	XP_044536929.1
Aves	<i>Corvus hawaiiensis</i>	XP_048160132.1
	<i>Taeniopygia guttata</i>	XP_030127125.3

<i>Pyrgilauda ruficollis</i>	XP_041335801.1
<i>Hirundo rustica</i>	XP_039921773.1
<i>Onychostruthus taczanowskii</i>	XP_041283907.1
<i>Passer montanus</i>	XP_039555833.1
<i>Molothrus ater</i>	XP_036238378.1
<i>Oxyura jamaicensis</i>	XP_035179722.1
<i>Melopsittacus undulatus</i>	XP_030904860.1
<i>Strigops habroptila</i>	XP_030349011.1
<i>Aythya fuligula</i>	XP_032043173.1
<i>Lonchura striata domestica</i>	OWK64686.1
<i>Geospiza fortis</i>	XP_005415590.1
<i>Camarhynchus parvulus</i>	XP_030803986.1
<i>Serinus canaria</i>	XP_030094646.1
<i>Manacus vitellinus</i>	XP_008923018.1
<i>Zonotrichia albicollis</i>	XP_005483896.1
<i>Apteryx rowi</i>	XP_025939844.1
<i>Numida meleagris</i>	XP_021249041.1
<i>Gallus gallus</i>	NP_990056.1
<i>Tyto alba</i>	XP_042663613.1
<i>Lagopus leucura</i>	XP_042726563.1
<i>Pygoscelis adeliae</i>	XP_009328950.1
<i>Calypte anna</i>	XP_030305365.1
<i>Aquila chrysaetos chrysaetos</i>	XP_029861401.1
<i>Haliaeetus leucocephalus</i>	XP_010568301.1
<i>Tyto alba</i>	XP_032854868.1
<i>Hirundo rustica</i>	XP_039921766.1
<i>Cygnus atratus</i>	XP_035413755.1
<i>Lepidothrix coronata</i>	XP_017664322.1

	<i>Amazona aestiva</i>	KQK77628.1
	<i>Chaetura pelagica</i>	XP_010000752.1
	<i>Antrostomus carolinensis</i>	XP_010171751.1
	<i>Nipponia nippon</i>	XP_009473093.1
	<i>Pelecanus crispus</i>	XP_009479867.1
	<i>Leptosomus discolor</i>	XP_009958677.1
	<i>Dryobates pubescens</i>	XP_009909179.1
	<i>Manacus vitellinus</i>	XP_008923018.1
	<i>Parus major</i>	XP_015479949.1
	<i>Falco rusticolus</i>	XP_037254769.1
	<i>Numida meleagris</i>	XP_021249038.1
Reptilia	<i>Pelodiscus sinensis</i>	XP_006128559.1
	<i>Gopherus evgoodei</i>	XP_030419000.1
	<i>Chelonia mydas</i>	XP_043402007.1
	<i>Dermochelys coriacea</i>	XP_043369528.1
	<i>Mauremys reevesii</i>	XP_039396469.1
	<i>Terrapene carolina triunguis</i>	XP_029766549.1
	<i>Trachemys scripta elegans</i>	XP_034627282.1
	<i>Python bivittatus</i>	XP_025018785.1
	<i>Pogona vitticeps</i>	XP_020658385.1
	<i>Zootoca vivipara</i>	XP_034967628.1
	<i>Podarcis muralis</i>	XP_028600040.1
	<i>Sceloporus undulatus</i>	XP_042323800.1
	<i>Lacerta agilis</i>	XP_033017209.1
	<i>Varanus komodoensis</i>	XP_044305785.1
	<i>Notechis scutatus</i>	XP_026522951.1
	<i>Pantherophis guttatus</i>	XP_034257179.1
	<i>Probothrops mucrosquamatus</i>	XP_015687286.1

	<i>Eremias argus</i>	QCQ80689.1
	<i>Sceloporus undulatus</i>	XP_042323800.1
	<i>Gekko japonicus</i>	XP_015272894.1
	<i>Lacerta agilis</i>	XP_033017209.1
	<i>Notechis scutatus</i>	XP_026522951.1
	<i>Pseudonaja textilis</i>	XP_026553662.1
	<i>Sphaerodactylus townsendi</i>	XP_048364960.1
	<i>Thamnophis sirtalis</i>	XP_013910221.1
	<i>Crotalus tigris</i>	XP_039186723.1
	<i>Thamnophis elegans</i>	XP_032080611.1
	<i>Crocodylus porosus</i>	XP_019390228.1
	<i>Chelonoidis abingdonii</i>	XP_032656285.1
	<i>Chrysemys picta bellii</i>	XP_005282030.1
	<i>Mauremys mutica</i>	XP_044873973.1
Amphibia	<i>Xenopus laevis</i>	NP_001081225.1
	<i>Bufo gargarizans</i>	XP_044145804.1
	<i>Rana temporaria</i>	XP_040189928.1
	<i>Bufo bufo</i>	XP_040274658.1
	<i>Geotrypetes seraphini</i>	XP_033799250.1
	<i>Microcaecilia unicolor</i>	XP_030047368.1
	<i>Nanorana parkeri</i>	XP_018421176.1
	<i>Cynops orientalis</i>	AFN55260.1
	<i>Xenopus tropicalis</i>	NP_001120900.1
	<i>Rhinatrema bivittatum</i>	XP_029441076.1
	<i>Geotrypetes seraphini</i>	XP_033799250.1
Osteichthyes	<i>Paralichthys olivaceus</i>	XP_019958716.1
	<i>Danio rerio</i>	AWP39888.1
	<i>Dicentrarchus labrax</i>	ABC70997.1

<i>Larimichthys crocea</i>	NP_001290322.1
<i>Scophthalmus maximus</i>	AVW89178.1
<i>Cynoglossus semilaevis</i>	XP_016894801.1
<i>Cyprinus carpio</i>	XP_018965718.1
<i>Cyprinodon variegatus</i>	XP_015245088.1
<i>Cyprinodon tularosa</i>	XP_038127805.1
<i>Takifugu rubripes</i>	NP_001027871.1
<i>Salmo salar</i>	NP_001133393.1
<i>Oncorhynchus mykiss</i>	BAU69680.1
<i>Monopterus albus</i>	XP_020451082.1
<i>Nothobranchius furzeri</i>	KAF7225424.1
<i>Miichthys miiuy</i>	AHG06618.1
<i>Anguilla japonica</i>	AYC61977.1
<i>Scleropages Formosus</i>	XP_018591577.1
<i>Poecilia reticulata</i>	XP_008417983.1
<i>Poecilia formosa</i>	XP_007549682.1
<i>Oplegnathus fasciatus</i>	AFM09714.1
<i>Collichthys lucidus</i>	TKS72762.1
<i>Notothenia coriiceps</i>	XP_010794822.1
<i>Poeciliopsis prolifica</i>	JAO80483.1
<i>Haplochromis burtoni</i>	XP_005944370.2
<i>Cyprinus carpio</i>	AGU12796.1
<i>Cyprinus carpio</i>	AGU12795.1
<i>Electrophorus electricus</i>	XP_026884624.2
<i>Amphiprion melanopus</i>	AEA08874.1
<i>Archocentrus centrarchus</i>	XP_030591190.1
<i>Bagarius yarrelli</i>	TSN95700.1
<i>Oryzias melastigma</i>	XP_024138660.1

<i>Oryzias latipes</i>	NP_001098140.1
<i>Siniperca chuatsi</i>	ADK47519.1
<i>Kryptolebias marmoratus</i>	XP_017278585.1
<i>Astyanax mexicanus</i>	XP_022539261.1
<i>Miichthys miiuy</i>	AHG06618.1
<i>Xiphophorus couchianus</i>	XP_027872889.1
<i>Oreochromis niloticus</i>	ADJ57601.1
<i>Lutjanus peru</i>	QBY35776.1
<i>Mugil incilis</i>	QDK54780.1

837

838

839 **Table 2. The CASP7 sequences used for the WebLogo analysis in this study.**

Class	Species	Accession Number
Mammalia	<i>Homo sapiens</i>	P55210
	<i>Mus musculus</i>	P97864
	<i>Mesocricetus auratus</i>	P55214
	<i>Bos taurus</i>	XP_002698555.1
	<i>Pan troglodytes</i>	XP_003825650.1
	<i>Macaca fascicularis</i>	XP_011822901.1
	<i>Oryctolagus cuniculus</i>	XP_002718761.1
	<i>Ovis aries</i>	XP_042094975.1
	<i>Capra hircus</i>	XP_005698554.1
	<i>Ailuropoda melanoleuca</i>	XP_034519188.1
	<i>Saimiri boliviensis boliviensis</i>	XP_034519188.1
	<i>Equus caballus</i>	XP_014588811.1
	<i>Delphinapterus leucas</i>	XP_022424428.1
	<i>Sus scrofa</i>	XP_020928977.1
	<i>Pan paniscus</i>	XP_003825650.1
	<i>Loxodonta africana</i>	XP_010587277.1
	<i>Canis lupus familiaris</i>	XP_005637795.1
	<i>Oryctolagus cuniculus</i>	XP_002718761.1
	<i>Vulpes vulpes</i>	XP_025849552.1
	<i>Heterocephalus glaber</i>	XP_004838949.1
	<i>Uroditellus parryi</i>	XP_026244955.1
	<i>Equus asinus</i>	XP_014723324.1
	<i>Tachyglossus aculeatus</i>	XP_038614071.1
	<i>Camelus bactrianus</i>	XP_010968130.1
	<i>Balaenoptera acutorostrata scammoni</i>	XP_007173176.1
	<i>Myodes glareolus</i>	XP_048279394.1
	<i>Dipodomys spectabilis</i>	XP_042545561.1

	<i>Delphinapterus leucas</i>	XP_022424428.1
	<i>Eptesicus fuscus</i>	XP_028005288.1
	<i>Bison bison bison</i>	XP_010857610.1
	<i>Pteropus alecto</i>	XP_006920912.1
	<i>Rhinopithecus bieti</i>	XP_017747992.1
	<i>Balaenoptera musculus</i>	XP_036685506.1
	<i>Saimiri boliviensis boliviensis</i>	XP_010331177.1
	<i>Myotis brandtii</i>	XP_005874576.1
	<i>Ursus americanus</i>	XP_045659135.1
	<i>Neogale vison</i>	XP_044096500.1
	<i>Vulpes lagopus</i>	XP_041601933.1
	<i>Equus quagga</i>	XP_046510739.1
	<i>Oryctolagus cuniculus</i>	XP_002718761.1
	<i>Galemys pyrenaicus</i>	KAG8517409.1
	<i>Manis pentadactyla</i>	XP_036754817.1
	<i>Gracilinanus agilis</i>	XP_044521530.1
Aves	<i>Corvus hawaiiensis</i>	XP_048166914.1
	<i>Taeniopygia guttata</i>	XP_030132303.3
	<i>Pyrgilauda ruficollis</i>	XP_041317496.1
	<i>Hirundo rustica</i>	XP_039927077.1
	<i>Onychostruthus taczanowskii</i>	XP_041271849.1
	<i>Passer montanus</i>	XP_039565350.1
	<i>Molothrus ater</i>	XP_036241577.1
	<i>Oxyura jamaicensis</i>	XP_035186324.1
	<i>Turdus rufiventris</i>	KAF4791228.1
	<i>Melopsittacus undulatus</i>	XP_005154576.1
	<i>Strigops habroptila</i>	XP_030344254.1
	<i>Aythya fuligula</i>	XP_032047208.1

<i>Lonchura striata domestica</i>	XP_021381698.1
<i>Geospiza fortis</i>	XP_005416425.1
<i>Camarhynchus parvulus</i>	XP_030807062.1
<i>Serinus canaria</i>	XP_009085424.1
<i>Manacus vitellinus</i>	XP_029814357.1
<i>Zonotrichia albicollis</i>	XP_005481147.1
<i>Apteryx rowi</i>	XP_025925619.1
<i>Numida meleagris</i>	XP_021255526.1
<i>Gallus gallus</i>	XP_040530979.1
<i>Tyto alba</i>	XP_032844696.2
<i>Lagopus leucura</i>	XP_042719653.1
<i>Pygoscelis adeliae</i>	XP_009324216.1
<i>Corvus brachyrhynchos</i>	XP_008632648.1
<i>Calypte anna</i>	XP_030309482.1
<i>Apteryx mantelli mantelli</i>	XP_013797196.1
<i>Apaloderma vittatum</i>	XP_009876249.1
<i>Haliaeetus leucocephalus</i>	XP_010563034.1
<i>Tyto alba</i>	XP_032844693.2
<i>Lonchura striata domestica</i>	OWK56987.1
<i>Cygnus atratus</i>	XP_035403725.1
<i>Cyanistes caeruleus</i>	XP_023785639.1
<i>Amazona aestiva</i>	KQK80332.1
<i>Mesitornis unicolor</i>	XP_010180183.1
<i>Melopsittacus undulatus</i>	XP_005154576.1
<i>Athene cunicularia</i>	XP_026707509.1
<i>Camarhynchus parvulus</i>	XP_030807062.1
<i>Gallus gallus</i>	XP_421764.3
<i>Chaetura pelagica</i>	XP_010006222.1

	<i>Ficedula albicollis</i>	XP_005048826.1
	<i>Serinus canaria</i>	XP_009085424.1
	<i>Cuculus canorus</i>	XP_009568871.1
	<i>Corvus cornix cornix</i>	XP_039410202.1
	<i>Nipponia nippon</i>	XP_009474328.1
	<i>Pelecanus crispus</i>	XP_009483464.1
	<i>Balearica regulorum gibbericeps</i>	XP_010295946.1
	<i>Manacus vitellinus</i>	XP_017924402.1
	<i>Phalacrocorax carbo</i>	XP_009508983.1
	<i>Parus major</i>	XP_015489222.1
	<i>Falco rusticolus</i>	XP_037256573.1
	<i>Corvus hawaiiensis</i>	XP_048166914.1
Reptilia	<i>Pelodiscus sinensis</i>	XP_006135034.1
	<i>Gopherus evgoodei</i>	XP_030424754.1
	<i>Chelonia mydas</i>	XP_037760318.1
	<i>Dermochelys coriacea</i>	XP_038265833.1
	<i>Mauremys reevesii</i>	XP_039402417.1
	<i>Terrapene carolina triunguis</i>	XP_026506928.1
	<i>Trachemys scripta elegans</i>	XP_034632909.1
	<i>Python bivittatus</i>	XP_007441954.1
	<i>Pogona vitticeps</i>	XP_020635566.1
	<i>Zootoca vivipara</i>	XP_034988803.1
	<i>Podarcis muralis</i>	XP_028586060.1
	<i>Sceloporus undulatus</i>	XP_042313006.1
	<i>Anolis carolinensis</i>	XP_008112942.1
	<i>Varanus komodoensis</i>	KAF7253514.1
	<i>Lacerta agilis</i>	XP_033005524.1
	<i>Notechis scutatus</i>	XP_026526250.1

	<i>Notechis scutatus</i>	XP_026526259.1
	<i>Pantherophis guttatus</i>	XP_034296056.1
	<i>Python bivittatus</i>	XP_007441954.1
	<i>Protobothrops mucrosquamatus</i>	XP_015676047.1
	<i>Zootoca vivipara</i>	XP_034988803.1
	<i>Podarcis muralis</i>	XP_028586060.1
	<i>Sceloporus undulatus</i>	XP_042313006.1
	<i>Anolis carolinensis</i>	XP_003223489.1
	<i>Varanus komodoensis</i>	KAF7253514.1
	<i>Varanus komodoensis</i>	XP_044295172.1
	<i>Lacerta agilis</i>	XP_033005524.1
	<i>Notechis scutatus</i>	XP_026526250.1
	<i>Pantherophis guttatus</i>	XP_034296056.1
	<i>Protobothrops mucrosquamatus</i>	XP_015676047.1
	<i>Pseudonaja textilis</i>	XP_026560589.1
	<i>Sphaerodactylus townsendi</i>	XP_048360798.1
	<i>Crotalus tigris</i>	XP_039208641.1
	<i>Thamnophis elegans</i>	XP_032081709.1
	<i>Chelonoidis abingdonii</i>	XP_032631043.1
	<i>Mauremys mutica</i>	XP_044881213.1
	<i>Chrysemys picta bellii</i>	XP_042711209.1
	<i>Trachemys scripta elegans</i>	XP_034632910.1
	<i>Crocodylus porosus</i>	XP_019411759.1
Amphibia	<i>Xenopus laevis</i>	BAA94748.1
	<i>Lithobates catesbeianus</i>	ACO51875.1
	<i>Bufo gargarizans</i>	XP_044152860.1
	<i>Rana temporaria</i>	XP_040217514.1
	<i>Bufo bufo</i>	XP_040291577.1

	<i>Geotrypetes seraphini</i>	XP_033799427.1
	<i>Microcaecilia unicolor</i>	XP_030058570.1
	<i>Nanorana parkeri</i>	XP_018419981.1
	<i>Cynops orientalis</i>	AFN55259.1
	<i>Xenopus tropicalis</i>	NP_001016299.1
	<i>Xenopus tropicalis</i>	CAJ82745.1
	<i>Rhinatrema bivittatum</i>	XP_029466303.1
Osteichthyes	<i>Paralichthys olivaceus</i>	XP_019965790.1
	<i>Danio rerio</i>	AWP39893.1
	<i>Dicentrarchus labrax</i>	CBN81450.1
	<i>Larimichthys crocea</i>	XP_010740374.1
	<i>Scophthalmus maximus</i>	XP_047183439.1
	<i>Cynoglossus semilaevis</i>	XP_024912944.1
	<i>Oncorhynchus kisutch</i>	XP_020329675.1
	<i>Cyprinus carpio</i>	XP_042591656.1
	<i>Cyprinodon variegatus</i>	XP_015245976.1
	<i>Fundulus heteroclitus</i>	XP_012710613.2
	<i>Cyprinodon tularosa</i>	XP_029690759.1
	<i>Takifugu rubripes</i>	XP_029690759.1
	<i>Salmo salar</i>	XP_014012537.2
	<i>Oncorhynchus mykiss</i>	QWC93456.1
	<i>Ictalurus punctatus</i>	XP_017338665.1
	<i>Monopterus albus</i>	XP_020460887.1
	<i>Nothobranchius furzeri</i>	KAF7204369.1
	<i>Alosa alosa</i>	XP_048088911.1
	<i>Alosa sapidissima</i>	XP_041939757.1
	<i>Anabarrilius grahami</i>	ROI81868.1
	<i>Salvelinus alpinus</i>	XP_023842909.1

<i>Salvelinus alpinus</i>	XP_023861943.1
<i>Gadus morhua</i>	XP_030195895.1
<i>Labrus bergylta</i>	XP_020506858.1
<i>Toxotes jaculatrix</i>	XP_040923277.1
<i>Collichthys lucidus</i>	TKS89629.1
<i>Notothenia coriiceps</i>	XP_010765636.1
<i>Melanotaenia boesemani</i>	XP_041830322.1
<i>Clarias magur</i>	KAF5896512.1
<i>Anabas testudineus</i>	XP_026207209.1
<i>Chelmon rostratus</i>	XP_041817955.1
<i>Anguilla anguilla</i>	XP_035260198.1
<i>Trematomus bernacchii</i>	XP_033975792.1
<i>Pimephales promelas</i>	KAG1954935.1
<i>Bagarius yarrelli</i>	TSK14754.1
<i>Seriola dumerili</i>	XP_022598445.1
<i>Xiphophorus hellerii</i>	XP_032429481.1
<i>Sebastes umbrosus</i>	XP_037611938.1
<i>Parambassis ranga</i>	XP_028285718.1
<i>Oryzias melastigma</i>	KAF6724972.1
<i>Megalops cyprinoides</i>	XP_036387401.1
<i>Salarias fasciatus</i>	XP_029954337.1
<i>Salvelinus namaycush</i>	XP_038866282.1

840

841

842 **Table 3. The credits for the pictures used in this study.**

Icon	Figure Origin
Monotremata	Becky-Barnes
Didelphimorphia	Daniel-Stadtmauer
Diprotodontia	Gavin-Prideaux
Afrosoricida	Mo-Hassan
Macroscelidea	uncredited
Pilosa	Xavier-A-Jenkins
Primates	T-Michael-Keesey
Rodentia	Jiro-Wada
Lagomorpha	Margot-Michaud
Eulipotyphla	Becky-Barnes
Perissodactyla	Mercedes-Yrayzoz-vectorized-by-T-Michael-Keesey
Artiodactyla	DFoidl-modified-by-T-Michael-Keesey
Carnivora	Chlo-Schmidt
Pholidota	Steven-Traver
Chiroptera	Margot-Michaud
<i>Homo sapiens</i>	NASA
Avian	Ferran Sayol
Reptilia	Gabriela Palomo-Munoz
Amphibia	Yusan Yang
Actinopterygii	Milton Tan

843

844 **Table 4. The constructs of the truncate and chimeric proteins.**

Truncate/chimeric protein	Amino acid sequence
TrGSDME-NT	1-258 aa of TrGSDME
TrGSDME-CT	259-474 aa of TrGSDME
HsNT-TrCT	1-270 aa of HsGSDME + 259-274 aa of TrGSMDE
TrNT-HsCT	1-258 aa of TrGSDME + 271-496 aa of HsGSDME
Hsp20-Trp10	1-198 aa of HsCASP7 + 207-313 aa of TrCASP7
Trp20-Hsp10	1-206 aa of TrCASP7 + 199-303 aa of HsCASP7

845

846

847

848 **Table 5. Primers used in this study.**

Primer	Sequence (5'-3')
Clone primers	
TrCASP3 forward	ATGTCGGCCAACGGACC
TrCASP3 reverse	GGAAAAATACATCTCTTTGGTCAGC
TrCASP7 forward	ATGCAGATGGCTGGAGAACC
TrCASP7 reverse	GTAAAGTACAGTTCTTTTGTGTCAGCATCG
TrGSDME forward	ATGTTTTCCAAGGCCACGG
TrGSDME reverse	ATCGATAAAATCCGTTTCAGACTTTG
Recombination primers	
TrCASP3 forward	TAAGAAGGAGATATACATATGATGT CGGCCAACGGACC
TrCASP3 reverse	GTGGTGGTGGTGGTGGTCTCGAGGGA AAAATACATCTCTTTGGTCAGC
TrCASP7 forward	TAAGAAGGAGATATACATATGATGCA GATGGCTGGAGAACC
TrCASP7 reverse	GTGGTGGTGGTGGTGGTCTCGAGGTAA AGTACAGTTCTTTTGTGTCAGCATCG

TrGSDME forward	TAAGAAGGAGATATACATATGATGTTT TCCAAGGCCACGG
TrGSDME reverse	GTGGTGGTGGTGGTGGTCTCGAGATCGA TAAAATCCGTTTCAGACTTTG
TrGSDME-D255R forward	TGGAATCCCCGAGAGCGGTGGATGG TCGTTGTCT
TrGSDME-D255R reverse	AGACAACGACCATCCACCGCTCTCGG GGATTCCCA
TrGSDME-D258A forward	AGACAACGACCATCCACCGCTCTCGG GGATTCCCA
TrGSDME-D258A reverse	AACGACCATCCACCGCTCTCGGGGATT CCCA
HsNT-TrCT forward	TAAGAAGGAGATATACATATGATGTT TGCCAAAGCAACCAGG
HsNT-TrCT reverse	GTGGTGGTGGTGGTGGTCTCGAGATCG ATAAAATCCGTTTCAGACTTTG
TrNT-HsCT forward	TAAGAAGGAGATATACATATGATGTT TTCCAAGGCCACGG
TrNT-HsCT reverse	GTGGTGGTGGTGGTGGTCTCGAGTGAA TGTTCTCTGCCTAAAGC
Hsp20-Trp10 forward	TAAGAAGGAGATATACATATGATGG CAGATGATCAGGGCTG
Hsp20-Trp10 reverse	GTGGTGGTGGTGGTGGTCTCGAGGTTA AAGTACAGTTCTTTTGTTCAGCATC
Trp20-Hsp10 forward	TAAGAAGGAGATATACATATGATGG CTGGAGAACCCACTGAG
Trp20-Hsp10 reverse	GTGGTGGTGGTGGTGGTCTCGAGTTGA CTGAAGTAGAGTTCCTTGGTGA

MmGSDME-N234S forward	GTTATTACTCATGGAGGAGCCCAGG GAAAG
MmGSDME-N234S reverse	CTTTCCTGGGCTCCTCCATGAGTA ATAAC
HsGSDME-Δ261-266 forward	CCAGGATGGACATCCATTTGCGGAG CTGC
HsGSDME-Δ261-266 reverse	CAAATGGATGTCCATCCTGGGAAGAT ATCCCAT
HsGSDME-Δ281-296 forward	CCTGGTCTTT GACATGCCAGATGCTG CGCA
HsGSDME-Δ281-296 reverse	CTGGCATGTCAAAGACCAGGGGGTCC AGGTAGA
Overexpression preimers	
TrGSDME-FL forward	TCAGATCTCGAGCTCAAGCTTATGTTT TCCAAGGCCACGG
TrGSDME-FL reverse	CATGGTGGCGACCGGTGGATCATCGA TAAAATCCGTTTCAGACTTTG
TrGSDME-NT forward	TCAGATCTCGAGCTCAAGCTTATGTTT TCCAAGGCCACGG
TrGSDME-NT reverse	CATGGTGGCGACCGGTGGATCGTCTAC AGCATCAGGAGACTCC
TrGSDME-CT forward	TCAGATCTCGAGCTCAAGCTTATGGGCC GGTGCCTGGA
TrGSDME-CT reverse	CATGGTGGCGACCGGTGGATCATCGA TAAAATCCGTTTCAGACTTTG
TrCASP7 forward	TTCTGAAGAGGACTTGAATTCAATGG CTGGAGAACCCACTGAG
TrCASP7 reverse	ACGACTCACTATAGTTCTAGATCAGT

	TAAAGTACAGTTCTTTTGTGTCAGC
TrCASP3 forward	TTCTGAAGAGGACTTGAATTCAAATG
	TCGGCCAACGGACC
TrCASP3 reverse	ACGACTCACTATAGTTCTAGAGGAA
	AAATACATCTCTTTGGTCAGC
HsGSDME forward	CAAGCTTGCGGCCGCGAATTCAATGT
	TTGCCAAAGCAACCAGG
HsGSDME reverse	ACGACTCACTATAGTTCTAGATCATG
	AATGTTCTCTGCCTAAAGC
HsCASP7 forward	TTCTGAAGAGGACTTGAATTCAATGG
	CAGATGATCAGGGCTG
HsCASP7 reverse	ACGACTCACTATAGTTCTAGACTATTG
	ACTGAAGTAGAGTTCCTTGGTG
HsCASP3 forward	TTCTGAAGAGGACTTGAATTCAATGG
	AGAACACTGAAAACCTCAGTGG
HsCASP3 reverse	ACGACTCACTATAGTTCTAGATTAGTG
	ATAAAAATAGAGTTCTTTTGTGAGC
MmCASP7 forward	TTCTGAAGAGGACTTGAATTCAATGA
	CCGATGATCAGGACTGTGC
MmCASP7 reverse	ACGACTCACTATAGTTCTAGATCAAC
	GGCTGAAGTACAGCTCTTTGG

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5






Figure 6


A




E




I




B



C




D



F



G



H




Figure 1-figure supplement 1




Figure 1-figure supplement 2




Figure 2-figure supplement 1




Figure 2-figure supplement 2




Figure 2-figure supplement 3




Figure 3-figure supplement 1




Figure 3-figure supplement 2




Figure 4-figure supplement 1




Figure 5-figure supplement 1




Figure 5-figure supplement 2




Figure 5-figure supplement 3




Figure 6-figure supplement 1




Figure 6-figure supplement 2

