

No evidence for highly pathogenic avian influenza virus H5N1 (clade 2.3.4.4b) in the Antarctic region during the austral summer 2022/23

Simeon Lisovski¹, Anne Günther², Meagan Dewar³, David Ainley⁴, Rodrigo Arce^{5,6}, Grant Ballard⁷, Josabel Belliure⁸, Thierry Boulinier⁹, Ashley Bennison¹⁰, Craig Cary¹¹, Paulo Catry¹², Augustin Clessin⁸, Maelle Connan¹³, Aidan Cox⁷, Juan Cristina⁵, Megan Elrod⁷, Irene Ferreira^{5,6}, Zoe Fowler¹⁴, Amandine Gamble^{15,16}, Joaquin Hurtado^{5,6}, Dennis Jongsomjit⁷, Mathilde Lejeune⁹, Amélie Lescroël⁷, Amy Li⁷, Virginia Morandini¹⁷, Gonzalo Moratorio^{4,6,18}, Teresa Militão¹², Pilar Moreno^{5,6}, Paula Perbolianachis^{5,6}, Jean Pennycook⁴, Annie E. Schmidt⁷, Lucy Smyth¹³, Alvaro Soutullo¹⁹, Jérémy Tornos⁹, Arvind Varsani²⁰, Ulrike Herzsich^{1,21,22}, Martin Beer², Michelle Wille²³

¹ Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Polar Terrestrial Environmental Systems, 14473 Potsdam, Germany.

² Friedrich-Loeffler-Institute, Greifswald–Insel Riems, Germany.

³ Future Regions Research Centre, Federation University Australia

⁴ H.T. Harvey & Associates Ecological Consultants, Los Gatos, CA 95032, USA

⁵ Laboratorio de Virología Molecular, Centro de Investigaciones Nucleares, Facultad de Ciencias, Universidad de la República, Igua 4225, Montevideo 11400, Uruguay

⁶ Laboratorio de Evolución Experimental de Virus, Institut Pasteur de Montevideo, Mataojo 2020, Montevideo 11400, Uruguay

⁷ Point Blue Conservation Science. Petaluma, CA 94924, USA

⁸ National Science Museum (MNCN-CSIC)

⁹ Centre d'Ecologie Fonctionnelle et Evolutive (CEFE), CNRS, Université Montpellier, EPHE, IRD, Montpellier, France

¹⁰ British Antarctic Survey, High Cross, Madingley, Cambridge, United Kingdom

¹¹ Thermophile Research Unit, Te Aka Mātuatua-School of Science, Te Whare Wānanga o Waikato, University of Waikato, Hamilton, Aotearoa-New Zealand

¹² MARE – Marine and Environmental Sciences Centre / ARNET - Aquatic Research Network, Ispa – Instituto Universitário, Portugal

¹³ Department of Zoology, Marine Apex Predator Research Unit (MAPRU), Institute for Coastal and Marine Research, Nelson Mandela University, Gqeberha 6001, South Africa

¹⁴ Department of Agriculture, Bypass Road, Stanley, Falkland Islands

- 37 ¹⁵ School of Biodiversity, One Health & Veterinary Medicine, University of Glasgow, United
38 Kingdom
- 39 ¹⁶ Department of Public & Ecosystem Health, Cornell University, New York, United States
- 40 ¹⁷ Departamento de Ecología Evolutiva, Museo Nacional de Ciencias Naturales, CSIC,
41 28006 Madrid, Spain
- 42 ¹⁸ Centro de Innovación en Vigilancia Epidemiológica, Institut Pasteur Montevideo, Mataojo
43 2020, Montevideo 11400, Uruguay
- 44 ¹⁹ Departamento de Ecología y Gestion Ambiental, Centro Universitario Regional del Este,
45 Universidad de la Republica, Montevideo, Uruguay
- 46 ²⁰ Biodesign Center for Fundamental and Applied Microbiomics, Center for Evolution and
47 Medicine, School of Life Sciences, Arizona State University, Tempe, AZ, USA
- 48 ²¹ Institute of Biochemistry and Biology, University of Potsdam, Potsdam, Germany
- 49 ²² Institute of Environmental Science and Geography, University of Potsdam, Potsdam, Germany
- 50 ²³ Centre for Pathogen Genomics, Department of Microbiology and Immunology, at the Peter
51 Doherty Institute for Infection and Immunity, The University of Melbourne, Melbourne, Victoria,
52 Australia

53

54 Abstract

55 The current highly pathogenic avian influenza H5N1 panzootic has profound impacts on
56 wild birds. Herein, we compiled H5N1 surveillance from Antarctica and Sub-Antarctic
57 Islands to ascertain whether HPAI was present in this region. Observations and HPAI
58 testing indicated no incursion of the virus during the austral summer 2022/23.

59 Main text

60 The increasing intensity of highly pathogenic avian influenza virus (HPAIV) H5N1
61 clade 2.3.4.4b outbreaks have had a profound impact on poultry and wildlife (1). Wild
62 bird movements have underpinned the rapid spread of this virus that swept across most
63 continents within two years (2). Compared to previous HPAIV subtypes and clades,
64 H5N1 2.3.4.4b has significantly improved replication in wild birds (3), and increased
65 fitness through continuous reassortments (4). Also, the virus is able to infect an
66 unusually broad variety of wild bird species (3), which has likely facilitated this spread. In
67 addition to their role as viral spreaders, wild birds are suffering huge losses following
68 mass mortality events, and the scale of mortality amongst wild birds is likely in the
69 millions rather than tens of thousands reported (5). Thus, the recent panzootic is a
70 serious conservation concern for a large range of wild bird species.

71 Due to the absence of waterfowl species that migrate to the Antarctic and sub-
72 Antarctic islands, the incursion risk of HPAI in notably these southernmost regions had
73 previously been considered low. Still, waterfowl are present in northern fringes of the
74 Southern Ocean.



75 *Figure 1: Seabird migration routes connect the world's oceans with Antarctic and sub-Antarctic*
76 *breeding sites. The south polar skua (Catharacta maccormicki) provides an example of trans-*
77 *hemispheric migrations (brown arrows, top left image). Brown skuas (Catharacta antarcticus),*
78 *Giant Petrels (Macronectes giganteus & halli, dark arrows, bottom image) and a whole range of*
79 *albatross species remain south but can be seen in waters of South America, southern Africa,*
80 *New Zealand, and Australia. Map made with Natural Earth.*

Moreover, the millions of seabirds that breed in the Antarctic region, and known migration and post-breeding dispersal routes establish links and thereby substantial global connectivity (Figure 1), including with regions of recent HPAIV H5N1 outbreaks. And despite the perceived remoteness, LPAIV viruses and antibodies against these viruses have previously been detected in various seabird species nesting at sites along the Antarctic Peninsula and South Shetland Islands, with viral genomes illustrating high phylogenetic relationships to viruses circulating on other continents (6, 7). As a result, experts have considered the risk of incursion of the recent panzootic HPAI H5 viruses into the Antarctic region in 2022/23 summer season to be high (8).

To identify a possible incursion of H5N1 into in Antarctic region during the summer season 2022/23, we sampled migratory seabirds at different locations across Antarctica and in sub-Antarctic areas (Figure 2), and collated observation data from researchers working in and around seabird colonies. In particular, we aimed to collect information pertaining to suspicious signs of unusual mortality and known clinical signs of HPAIV infection including loss of coordination and balance, trembling head and body, lethargy, respiratory distress, and conjunctivitis (8). Across all locations, sample collection was done in accordance with institutional animal ethics approval and sample testing was performed with national frameworks, with details available in the technical annex.

Overall, sampling and observational efforts span from early November to late March, comprising the entire austral breeding season in 2022/23, and include a large range of species (i.e., penguins, gulls, skuas, and petrels; see technical annex for more information) and locations. All swab samples collected were tested negative for avian influenza virus and thus for HPAI. Several suspicious observations of dead wild birds were recorded on the Falkland Islands (Gentoo penguin *Pygoscelis papua*, Cattle egret *Bubulcus ibis*), and South Georgia (Wandering albatrosses *Diomedea exulans*) and confirmed negative for HPAI based on qPCR testing. Taken together, and with the general lack of suspicious signs of HPAIV, we conclude that there is no evidence of HPAIV H5N1 clade 2.3.4.4b incursion into the Antarctic region during the austral summer 2022/23.

Obviously, incursion risk of HPAIV is contingent on a combination of factors. Most importantly, that (i) host species get in contact with HPAIV before travelling into the Antarctic regions, (ii) can migrate with an infection, and (iii) have contact with further susceptible species which could be the starting point of a new epizootic. While HPAIV (including H5N1) activity has occurred in South African seabirds since 2017, HPAIV did

not reach South America until ~Oct 2022, and Patagonia and Tierra del Fuego until early 2023 (2, 9, 10). Furthermore, while mainly ducks were previously central to HPAIV outbreaks, the global panzootic includes a large range of bird species, and increasingly novel species groups like seabirds (11). Notably in breeding colonies of seabirds, there have been extensive outbreaks in Europe in 2022 and 2023, and the southward expansion of HPAIV H5N1 in South America was largely driven through outbreaks in coastal seabirds, including pelicans and cormorants, as well as spill-overs to coastal marine mammals with the deaths of >20,000 South American sea lions (3, 9, 12, 13). Millions of seabirds and marine mammals will again return to the Antarctic region during the austral summer (Oct 2023 - Mar 2024), and with H5N1 outbreaks now occurring very close to the Falkland Island and the Antarctic peninsula. Thus, the risk of viral incursion will be even higher compared to 2022 (14).



Figure 2: (top) Sampling locations for qPCR analysis and the detection of H5N1 2.3.4.4b, as well as locations with intensive observational efforts to identify signs of HPAIV infections within breeding bird communities. (bottom) Northern giant petrels and Brown skuas scavenging on an Antarctic fur seal carcass, showing inter-species interactions with the potential for HPAI virus transmission (photo taken on South Georgia by Paulo Catry).

136

137 The consequences of viral incursion(s) into Southern Ocean wildlife are unclear
138 but based on observations from other regions, will likely have devastating effects.
139 Critically, densities of seabird colonies are very, facilitating the transmissions between
140 individuals (15). Further prospecting movements of potential recruits, predator-prey
141 interactions between bird species (e.g., skuas, penguins, and sheathbills), as well as
142 species scavenging on dead seabirds and mammals, may promote rapid spread of the
143 virus between colonies. The interaction between seabirds and marine mammals in the
144 Antarctic region may also result in further transmissions to mammals and facilitate the
145 adaptation of the virus in mammalian species. Finally, most animals of the Southern
146 Ocean are endemic to the region, such that mass mortality events in Antarctica due to
147 HPAI H5 will cause a very real conservation concern for many species.

148 Detecting H5N1 incursion(s) into the Antarctic region is highly relevant and
149 surveys for mortality and sampling should therefore be prioritized during the austral
150 summer 2023/24. These activities should be undertaken with consideration of the
151 potentially zoonotic risks of (emerging) HPAIV H5 risks (see details in 8). Importantly,
152 activities should not promote virus transmission within colonies nor the spatial spread
153 between colonies and species. Nevertheless, sampling, and detailed analysis of
154 lineages and virus phenotype will provide crucial information needed to assess risks and
155 respond to future wild bird outbreaks.

Acknowledgements

We acknowledge logistical and financial support from, the Uruguayan Antarctic Institute (SL, UH, AS, RA, JC, IF, JH, GM, PP, PM), Government South Georgia and South Sandwich Islands (AB), Antarctica New Zealand (CC), Spanish Research Agency (VM), French Polar Institute (ECOPATH-1151; TB, ML, JT), ANR (ECOPATHS ANR-21-CE35-0016 and REMOVE_DISEASE ANR-21-BIRE-006-01; TB, ML, JT), Biodiversa and Water JPI under the BiodivRestore ERA-NET Cofund (GA N°101003777; TB, MC, ML, PC), National Nature Reserve of Terres Australes et Antarctique Françaises, the South African Departments of Science and Innovation (National Research Foundation - South African National Antarctic Programme project no. 129226; MC) and of Forestry, Fisheries and the Environment, National Science Foundation award ANT 1935870 and ANT 2040199 (GB, DA, ME, AC AS, AL, AL, AV, JP), the Royal Society (Newton International Fellowship NIF\R1\211869; AGa), the UK Department for Environment, Food & Rural Affairs (Darwin Plus Grant DPLUS167; AGa), Falklands Conservation, the Wildlife Conservation Society, the South Atlantic Environmental Research Institute, Falkland Islands Government (PC), FCT – Portugal through the strategic project UIDB/04292/2020 (MARE) and LA/P/0069/2020 (ARNET) and DivRestore/0012/2020 (REMOVE_DISEASE project). We thank Teresa Militão, Corisande Abiven, Katharina Reusch, Louis Pole-Evans for fieldwork support. MB and AG received financial support by the European Union Horizon Europe research program (grant agreement “Kappa-Flu No. 101084171”).

References

1. Wille M, Barr IG. Resurgence of avian influenza virus. *Science*. 2022;376(6592):459-60.
2. Adlhoch C, Fusaro A, Gonzales JL, Kuiken T, Mirnaviciute G, Niqueux EE, et al. Avian influenza overview March - April 2023. *Efsa Journal*. 2023 Jun;21(6).
3. Abolnik C, Phiri T, Peyrot B, de Beer R, Snyman A, Roberts D, et al. The Molecular Epidemiology of Clade 2.3.4.4B H5N1 High Pathogenicity Avian Influenza in Southern Africa, 2021–2022. *Viruses*; 2023.
4. Kandeil A, Patton C, Jones JC, Jeevan T, Harrington WN, Trifkovic S, et al. Rapid evolution of A(H5N1) influenza viruses after intercontinental spread to North America. *Nat Commun*. 2023 May 29;14(1):3082.
5. WOA. 2023 [cited; Available from: <https://wahis.woah.org/>

- 189 6. Hurt AC, Vijaykrishna D, Butler J, Baas C, Maurer-Stroh S, Silva-de-la-Fuente MC, et
190 al. Detection of Evolutionarily Distinct Avian Influenza A Viruses in Antarctica. *Mbio*.
191 2014 May-Jun;5(3).
- 192 7. de Seixas MMM, de Araujo J, Krauss S, Fabrizio T, Walker D, Ometto T, et al. H6N8
193 avian influenza virus in Antarctic seabirds demonstrates connectivity between South
194 America and Antarctica. *Transboundary and Emerging Diseases*. 2022 Oct 21.
- 195 8. Dewar M, Wille M, Gamble A, Vanstreels R, Boulonier T, Smith A, et al. The Risk of A
196 vian Influenza in the Southern Ocean: A practical guide. 2022.
- 197 9. FAO. Global AIV with zoonotic potential situation update. 2023 July 2023 [cited;
198 Available from:
199 [https://www.fao.org/ag/againfo/programmes/en/empres/Global_AIV_Zoonotic_Update](https://www.fao.org/ag/againfo/programmes/en/empres/Global_AIV_Zoonotic_Update/situation_update.html)
200 [e/situation_update.html](https://www.fao.org/ag/againfo/programmes/en/empres/Global_AIV_Zoonotic_Update/situation_update.html)
- 201 10. Breed A, Dewar ML, Dodyk L, Kuiken T, Matus R, Serafini PP, et al. Southward
202 expansion of high pathogenicity avian influenza H5 in wildlife in South America:
203 estimated impact on wildlife populations, and risk of incursion into Antarctica; 2023.
- 204 11. Qiqi Y, Ben W, Phillipe L, Lu D, Tong M, Wiebe RA, et al. Synchrony of Bird
205 Migration with Avian Influenza Global Spread; Implications for Vulnerable Bird
206 Orders. *bioRxiv*. 2023:2023.05.22.541648.
- 207 12. Mariana L, Alejandra G-G, Breno M-S, Diana J, Patricia B, Carlos C-M, et al. Highly
208 pathogenic avian influenza A (H5N1) in marine mammals and seabirds in Peru.
209 *bioRxiv*. 2023:2023.03.03.531008.
- 210 13. Gunther A, Krone O, Svansson V, Pohlmann A, King J, Hallgrimsson GT, et al.
211 Iceland as Stepping Stone for Spread of Highly Pathogenic Avian Influenza Virus
212 between Europe and North America. *Emerging Infectious Diseases*. 2022
213 Dec;28(12):2383-8.
- 214 14. Dewar M, Vanstreels R, Boulonier T, Cary C, Gamble A, Clessin A, et al. Risk
215 Assessment of Highly Pathogenic Avian Influenza in the Southern Ocean Report
216 prepared for Scientific Committee on Antarctic Research; 2023.
- 217 15. Boulonier T. Avian influenza spread and seabird movements between colonies.
218 *Trends in Ecology & Evolution*. 2023;38(5):391-5.

219

220