

Gene expression imputation across multiple brain regions reveals schizophrenia risk throughout development.

Author List:

Laura M. Huckins^{1,2}, Amanda Dobbyn^{1,2}, Douglas M. Ruderfer³, Gabriel Hoffman^{1,4}, Weiqing Wang^{1,2}, Antonio Pardinas⁵, Veera M Rajagopal^{6,15,16}, Thomas D. Als^{6,15,16}, Hoang Nguyen^{1,2}, Kiran Girdhar^{1,2}, James Boocock⁷, Panos Roussos^{1,2}, Menachem Fromer^{1,2}, Robin Kramer⁸, Enrico Domencini⁹, Eric Gamazon^{3,14}, Shaun Purcell^{1,2,4}, CommonMind Consortium, the Schizophrenia Working Group of the Psychiatric Genomics Consortium, iPSYCH-GEMS Schizophrenia Working Group, Ditte Demontis^{6,15,16}, Anders D. Børglum^{6,15,16}, James Walters⁵, Michael O'Donovan⁵, Patrick Sullivan^{10,11}, Michael Owen⁵, Bernie Devlin¹², Solveig K. Sieberts¹³, Nancy Cox^{3,14}, Hae Kyung Im¹⁴, Pamela Sklar^{1,2,4}, Eli A. Stahl^{1,2}

Author Affiliations:

1. Division of Psychiatric Genomic, Icahn School of Medicine at Mount Sinai, NYC, NY;
2. Department of Genetics and Genomics, Icahn School of Medicine at Mount Sinai, NYC, NY;
3. Vanderbilt University Medical Center, Nashville, TN;
4. Icahn Institute for Genomics and Multiscale Biology, Icahn School of Medicine at Mount Sinai, NYC, NY, USA;
5. MRC Centre for Neuropsychiatric Genetics and Genomics, Cardiff University, UK;
6. Department of Biomedicine, Aarhus University, Aarhus, Denmark
7. Department of Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, USA;
8. Human Brain Collection Core, National Institute of Mental Health, Bethesda, MD, USA;
9. Laboratory of Neurogenomic Biomarkers, Centre for Integrative Biology (CIBIO), University of Trento, Trento, Italy;
10. University of North Carolina at Chapel Hill, NC, USA
11. Karolinska Institutet, Stockholm, Sweden
12. Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA, USA;
13. Systems Biology, Sage Bionetworks, Seattle, WA, USA;
14. Section of Genetic Medicine, Department of Medicine, University of Chicago, Chicago, Illinois, USA
15. The Lundbeck Foundation Initiative for Integrative Psychiatric Research, iPSYCH, Denmark.
16. Center for Integrative Sequencing, Aarhus University, Aarhus, Denmark.

Abstract

Transcriptomic imputation approaches offer an opportunity to test associations between disease and gene expression in otherwise inaccessible tissues, such as brain, by combining eQTL reference panels with large-scale genotype data. These genic associations could elucidate signals in complex GWAS loci and may disentangle the role of different tissues in disease development. Here, we use the largest eQTL reference panel for the dorso-lateral pre-frontal cortex (DLPFC), collected by the CommonMind Consortium, to create a set of gene expression predictors and demonstrate their utility. We applied these predictors to 40,299 schizophrenia cases and 65,264 matched controls, constituting the largest transcriptomic imputation study of schizophrenia to date. We also computed predicted gene expression levels for 12 additional brain regions, using publicly available predictor models from GTEx. We identified 413 genic associations across 13 brain regions. Stepwise conditioning across the genes and tissues identified 71 associated genes (67 outside the MHC), with the majority of associations found in the DLPFC, and of which 14/67 genes did not fall within previously genome-wide significant loci. We identified 36 significantly enriched pathways, including hexosaminidase-A deficiency, and multiple pathways associated with porphyric disorders. We investigated developmental expression patterns for all 67 non-MHC associated genes using BRAINSPAN, and identified groups of genes expressed specifically pre-natally or post-natally.

Introduction

Genome-wide association studies (GWAS) have yielded large lists of disease-associated loci. Despite this, progress in identifying the causal variants driving these associations, particularly for complex psychiatric disorders such as schizophrenia, has lagged much further behind. Interpreting associated variants and loci is therefore vital to understanding how genetic variation contributes to disease pathology. Expression Quantitative Trait Loci (eQTLs), which are responsible for a substantial proportion of gene expression variance, have been posited as a potential link between associated loci and disease susceptibility¹⁻⁵, and indeed have yielded results for a host of complex traits⁶⁻⁹. Consequently, numerous methods to identify and interpret co-localisation of eQTLs and GWAS loci have been developed¹⁰⁻¹³. However, these methods require simplifying assumptions about genetic architecture (i.e., one causal variant per GWAS locus) and/or linkage disequilibrium, may be underpowered or overly conservative, especially in the presence of allelic heterogeneity, and have not yet yielded substantial insights into existing or novel loci.

Biologically relevant information can be extracted by transcriptomic investigations, as recently described by the CommonMind Consortium¹⁴ (CMC), thanks to detailed RNA-sequencing in a large cohort of genotyped individuals with schizophrenia and bipolar disorder¹⁴. These analyses however are underpowered to detect with statistical confidence differential expression of genes mapping at schizophrenia (SCZ) risk loci, due to the small effects predicted by GWAS combined with the difficulty of obtaining adequate sample sizes of neurological tissues¹⁴. Still, such methods do not necessarily identify all risk variation in GWAS loci. Transcriptomic imputation is an alternative approach that leverages large eQTL reference panels to bridge the gap between large-scale genotyping studies and biologically useful transcriptome studies^{15,16}. This approach seeks to identify and codify the relationships between genotype and gene expression in matched panels of individuals, then impute the genetic component of the transcriptome into large-scale genotype-only datasets, such as case-control GWAS cohorts, which enables investigation of disease-associated gene expression changes. This will allow us to study genes with modest effect sizes, likely representing a large proportion of genomic risk for psychiatric disorders^{14,17}.

The access to the large collection of dorso-lateral pre-frontal cortex (DLPFC) gene expression data collected by the CommonMind Consortium¹⁴ affords us a unique opportunity to study and codify relationships between genotype and gene expression. Here, we present a novel set of gene expression predictor models, built using CommonMind Consortium DLPFC data¹⁴. We compare different regression approaches to building these models (including elastic net¹⁵, Bayesian sparse linear mixed models and ridge regression¹⁶, and using max eQTLs), and benchmark performance of these predictors against existing GTEx prediction models. We applied our CMC DLPFC predictors and 12 GTEx-derived neurological prediction models to predict gene expression in schizophrenia GWAS data, obtained through collaboration with the Psychiatric Genomics Consortium (PGC) schizophrenia working group, the “CLOZUK2” cohort, and the iPSYCH-GEMS schizophrenia working group. We identified 413 genome-wide significant genic associations with schizophrenia in our PGC+CLOZUK2 sample, constituting 67 independent associations outside the MHC region. We demonstrate the relevance of these associations to schizophrenia aetiopathology using gene set enrichment analysis, and by examining the effects of manipulation of these genes in mouse models. Finally, we investigated spatio-temporal expression of these genes using a developmental transcriptome dataset, and identified distinct spatio-temporal patterns of expression across our associated genes.

Results

Prediction Models based on CommonMind Consortium DLPFC expression

Using matched genotype and gene expression data from the CommonMind Consortium Project, we developed DLPFC genetically regulated gene expression (GREX) prediction models. We systematically compared four approaches to building predictors^{15,16} within a cross-validation framework. Elastic net regression had a higher distribution of cross-validation R^2 (R_{cv}^2) and higher mean R_{cv}^2 values (Supplementary Figures 1, 2A) than all other methods. We therefore used elastic net regression to build our prediction models. We compared prediction models created using elastic net regression on SVA-corrected and uncorrected data¹⁴. The distribution of R_{cv}^2 values for the SVA-based models was significantly higher than for the un-corrected data^{14,18} (ks-test; $p < 2.2 \times 10^{-16}$; Supplementary figure 1B-C). In total, 10,929 genes were predicted with elastic net cross-validation $R_{cv}^2 > 0.01$ in the SVA-corrected data and were included in the final predictor database (mean $R_{cv}^2 = 0.076$).

To test the predictive accuracy of the CMC-derived DLPFC models, and to benchmark this against existing GTEx-derived prediction models, genetically-regulated gene expression (GREX) was calculated in an independent DLPFC RNA-sequencing dataset (the Religious Orders Study Memory and Ageing Project, ROSMAP¹⁹). We compared predicted GREX to measured ROSMAP gene expression for each gene (Replication R^2 or R_R^2) for the CMC-derived DLPFC models and twelve GTEx-derived brain tissue models^{15,20,21} (Figure 1, Supplementary Figure 2B). CMC-derived DLPFC models had higher average R_R^2 values (Mean $R_R^2 = 0.056$), more genes with $R_R^2 > 0.01$, and significantly higher overall distributions of R_R^2 values than any of the twelve GTEx models (ks-test, $p < 2.2 \times 10^{-16}$ across all analyses; Figure 1). Median R_R^2 values were significantly correlated with sample size of the original tissue set ($\rho = 0.92$, $p = 7.2 \times 10^{-6}$), the number of genes in the prediction model ($\rho = 0.9$, $p = 2.6 \times 10^{-5}$), and the number of significant ‘eGenes’ in each tissue type ($\rho = 0.95$, $p = 5.5 \times 10^{-7}$; Figure 1C). Notably, these correlations persist after removing obvious outliers (Figure 1C).

To estimate trans-ancestral prediction accuracy, genetically regulated gene expression was calculated for 162 African-American individuals and 280 European individuals from the NIMH Human Brain Collection Core (HBCC) dataset (supplementary figure 2B). R_R^2 values were

higher on average in Europeans than African-Americans (average $R_{R_EUR}^2 = 0.048$, $R_{R_AA}^2 = 0.040$), but were significantly correlated between African-Americans and Europeans ($\rho=0.78$, $p<2.2 \times 10^{-16}$, Pearson test; supplementary figure 3).

Application of Transcriptomic Imputation to Schizophrenia

We used CMC DLPFC and the 12 GTEx-derived brain tissue prediction models to impute genetically regulated expression levels (GREX) of 19,661 unique genes in cases and controls from the PGC-SCZ GWAS study²². Predicted expression levels were tested for association with schizophrenia. Additionally, we applied CMC and GTEx-derived prediction models to summary statistics from 11 PGC cohorts (for which raw genotypes were unavailable) and the CLOZUK2 cohort. Meta-analysis was carried out across all PGC-SCZ and CLOZUK2 cohorts using an odds-ratio based approach in METAL. Our final analysis included 40,299 cases and 65,264 controls (Figure 2A).

We identified 413 genome-wide significant associations, representing 256 genes in 13 tissues (Figure 3A). The largest number of associations were detected in the CMC DLPFC GREX data (Figure 3C; 49 genes outside the MHC, 69 genes overall). We sought replication of our CMC DLPFC SCZ-associations in an independent dataset of 4,133 cases and 24,788 controls in collaboration with the iPSYCH-GEMS SCZ working group (Figure 2B). We found significant correlation of effect sizes ($p=1.784 \times 10^{-04}$; $\rho=0.036$) and $-\log_{10}$ p-values ($p=1.073 \times 10^{-05}$; $\rho=0.043$) between our discovery (PGC+CLOZUK2) and replication (iPSYCH-GEMS) samples. Non-MHC Genes reaching genome-wide significance in our discovery sample (49 genes) were significantly more likely to reach nominal significance in the replication sample, and had significantly more consistent directions of effect than might be expected by chance (binomial test, $p=2.42 \times 10^{-05}$, $p=0.044$). (Suppl. info).

To identify the top independent associations within genomic regions, which include multiple associations for a single gene across tissues, or multiple nearby genes, we partitioned genic associations into 58 groups defined based on genomic proximity and applied stepwise forward conditional analysis within each group (Supplementary Table 1). In total, 67 genes remained genome-wide significant after conditioning (Table 1; Figure 3A-B). The largest signal was

identified in the CMC DLPFC predicted expression data (24 genes; Figure 3C), followed by the Putamen (7 genes). 19/67 genes did not lie within 1Mb of a previously genome-wide significant GWAS locus²² (shown in bold, Table 1); of these, 5/19 genes were within 1Mb of a locus which approached genome-wide significance ($p < 5 \times 10^{-7}$). The remaining 14 genes all fall within nominally significant PGC-SCZ GWAS loci ($p < 8 \times 10^{-4}$), but did not reach genome-wide significance.

Implicated genes highlight SCZ-associated molecular pathways and gene set analyses

We tested for overlap between our non-MHC SCZ-associated genes and 8,657 genesets comprised of 1) hypothesis-driven pathways and 2) general molecular database pathways. We corrected for multiple testing using the Benjamin-Hochberg false discovery rate (FDR) correction²³.

We identified three significantly associated pathways in our hypothesis-driven analysis (Table 2). Targets of the fragile-X mental retardation protein formed the most enriched pathway (FMRP; $p = 1.96 \times 10^{-8}$). Loss of FMRP inhibits synaptic function, is comorbid with autism spectrum disorder, and causes intellectual disability, as well as psychiatric symptoms including anxiety, hyperactivity and social deficits²⁴. Enrichment of this large group of genes has been observed frequently, in the original CommonMind analysis¹⁴, by colleagues investigating the same PGC and CLOZUK2 samples²⁶ as well as by investigators studying autism^{24,27}. There was a significant enrichment among our SCZ associated genes and genes that have been shown to be intolerant to loss-of-function mutations²⁸ ($p = 5.86 \times 10^{-5}$) as well as with CNVs associated with bipolar disorder²⁹ ($p = 7.92 \times 10^{-8}$), in line with a recent variant-based study of the same individuals²⁶.

Next, we performed an agnostic search for overlap between our schizophrenia-associated genes and ~ 8,500 molecular pathways collated from large, publicly available databases. 33 pathways were significantly enriched after FDR correction (Table 2, Suppl. Table 2), including a number of pathways with some prior literature in psychiatric disease. We identified an enrichment with porphyrin metabolism ($p = 1.03 \times 10^{-4}$). Deficiencies in porphyrin metabolism lead to “Porphyrria”, an adult-onset metabolic disorder with a host of associated psychiatric symptoms, in particular

episodes of violence and psychosis^{30–35}. Five pathways potentially related to porphyrin metabolism, regarding abnormal iron level in the spleen, liver and kidney are also significantly enriched, including 2/5 of the most highly enriched pathways ($p < 2.0 \times 10^{-4}$). The PANTHER and REACTOME pathways for Heme biosynthesis and the GO pathway for protoporphyrinogen IX metabolic process, which are implicated in the development of porphyric disorders, are also highly enriched ($p = 2.2 \times 10^{-4}$, 2.6×10^{-4} , 4.1×10^{-4}), although do not pass FDR-correction.

Hexosaminidase activity was enriched ($p = 3.47 \times 10^{-5}$) in our results; this enrichment is not driven by a single highly-associated gene; rather, every single gene in the HEX-A pathway is nominally significant in the SCZ association analysis (Supplementary Table 2). Deficiency of hexosaminidase A (HEX-A) results in serious neurological and mental problems, most commonly presenting in infants as “Tay-Sachs” disease³⁶. Adult-onset HEX-A deficiency presents with neurological and psychiatric symptoms, notably including onset of psychosis and schizophrenia³⁷. Five pathways corresponding to Ras- and Rab- signaling, protein regulation and GTPase activity were enriched ($p < 6 \times 10^{-5}$). These pathways have a crucial role in neuron cell differentiation³⁸ and migration³⁹, and have been implicated in the development of schizophrenia and autism^{40–43}. We also find significant enrichment with protein phosphatase type 2A regulator activity ($p = 5.24 \times 10^{-5}$), which was associated with MDD and across MDD, BPD and SCZ in the same large integrative analysis⁴⁴, and has been implicated in antidepressant response and serotonergic neurotransmission⁴⁵.

Predicted gene expression changes are consistent with functional validation studies

To test the functional impact of our SCZ-associated predicted gene expression changes (GREX), we performed two in-silico analyses. First, we compared directions of effect in our meta-analysis to those in the CMC analysis of differentially expressed genes between SCZ cases and controls. This analysis highlighted six loci where expression levels of a single gene putatively affected schizophrenia risk. All six of these genes are nominally significant in our DLPFC analysis, and two (*CLCN3* and *FURIN*) reach genome-wide significance. In the conditional analysis across all brain regions, one additional gene (*SNXI9*) reaches genome-wide significance. The direction of effect for all six genes matches the direction of gene expression changes observed in the original CMC paper, indicating that gene expression estimated in the imputed transcriptome reflects

measured expression levels in brains of individuals with Schizophrenia. Further, this observation is consistent with a model where the differential expression signature observed in CMC is caused by genetics rather than environment.

The original CMC analysis identified 21 eSNP genes using SHERLOCK^{14,46}, of which 17 were present in our CMC DLPFC analysis. 14/17 genes reached nominal significance (significantly more than expected by chance, $p=3.6 \times 10^{-16}$), and 11 reached genome-wide significance (binomial p-value 6.04×10^{-55}). Additionally, 31 regions contained genes ranked highly by Sherlock in the original CMC analysis (supplementary data file 2 in Fromer, M. *et al.* Gene expression elucidates functional impact of polygenic risk for schizophrenia. *Nat. Neurosci.* 19, 1442–1453 (2016)¹⁴). Of these, 14 regions lay near one of our CMC DLPFC associated genes, and 13/14 regions had common genes between SHERLOCK and PrediXcan analyses. Five loci included multiple SHERLOCK genes; in every instance we are able to specifically identify one or two associated genes from the longer SHERLOCK list.

To understand the impact of altered expression of our 67 SCZ-associated genes, we performed an in-silico analysis of mouse mutants, by collating large, publicly available mouse databases^{47–51}. We identified mutant mouse lines lacking expression of 37/67 of our SCZ-associated genes, and obtained 5,333 phenotypic data points relating to these lines, including 1,170 related to behavioral, neurological or craniofacial phenotypes. 25/37 genes were associated with at least one behavioral, neurological or related phenotype (Supplementary table 3). We repeated this analysis for genes identified in 366 GWAS, including any GWAS for which at least ten mutant mouse lines exist (105 GWAS). SCZ-associated genes were more likely to be associated with behavior, brain development and nervous system phenotypes than genes in these GWAS sets ($p=0.057$).

Spatiotemporal expression of SCZ-associated genes indicated distinct patterns of risk throughout development

We assessed expression of our SCZ-associated genes throughout development using BRAINSPAN⁵². Data were partitioned into eight developmental stages (four pre-natal, four post-natal), and four brain regions^{29,52} (Figure 4A). We noted that SCZ-associated genes were

significantly co-expressed, in both pre-natal and post-natal development and in all four brain regions, based on local connectedness⁵³ (Figure 4B), global connectedness⁵³ (i.e., average path length between genes, supplementary Figure 6), and network density (i.e., number of edges, supplementary Figure 7). Examining pairwise gene expression correlation (suppl. Fig 8) and gene co-expression networks (suppl. Fig 9) for each spatiotemporal point indicated that the same genes do not drive this co-expression pattern throughout development; rather, it appears that separate groups of genes drive early pre-natal, late pre-natal and post-natal clustering.

To visualize this, we calculated Z scores of gene expression for each SCZ-associated gene, across all 32 time-points (Figure 5). Genes clustered into four groups (supplementary fig 10), with distinct spatio-temporal expression signatures. The largest cluster (Cluster A, Figure 5A; 29 genes) spanned early to late-mid pre-natal development (4-24 weeks post conception), either across the whole brain (22 genes) or in regions 1-3 only (7 genes). 12 genes were expressed in late pre-natal development (Figure 5D; 25-38 pcw); 10 genes were expressed in regions 1-3, post-natally and in the late pre-natal period (Figure 5C), and 15 genes were expressed throughout development (Figure 5B), either specifically in region four (nine genes) or throughout the brain (six genes). We used a stratified qq-plot approach⁵⁴ to examine whether SNPs in cis-regions of genes in these four clusters are differentially enriched in psychiatric disorders. SNPs in cis-regions of genes in the two pre-natal clusters are more highly enriched than SNPs in cis-regions of genes in post-natal clusters, and compared to all SNPs, in childhood-onset disorders (ASD and ADHD, supplementary figure 13), but not adult-onset disorders (BPD and MDD, data not shown).

We noticed a relationship between patterns of gene expression and the likelihood of behavioral, neurological or related phenotypes in our mutant mouse model database. Mutant mice lacking genes expressed exclusively pre-natally in humans, or genes expressed pre- and post-natally, were more likely to have any behavioral or neurological phenotypes than mutant mice lacking expression of genes expressed primarily in the third trimester or post-natally ($p=1.7 \times 10^{-04}$) (supplementary figure 11).

Discussion

In this study, we present gene expression prediction models for the dorso-lateral pre-frontal cortex (DLPFC), constructed using CommonMind Consortium genotype and gene expression data. These prediction models may be applied to either raw data or summary statistics, in order to yield gene expression information in large data sets, and across a range of tissues. This has the significant advantage of allowing researchers to access transcriptome data for non-peripheral tissues, at scales currently prohibited by the high cost of RNA sequencing, and circumventing distortions in measures of gene expression stemming from errors of measurement or environmental influences. Since disease status may alter gene expression but not the germline profile, analyzing genetically regulated expression ensures that we identify only the causal direction of effect between gene expression and disease¹⁵. Large, imputed transcriptomic datasets represent the first opportunity to study the role of subtle gene expression changes (and therefore modest effect sizes) in disease development.

There are some inherent limitations to this approach. The accuracy of transcriptomic imputation (TI) is reliant on access to large eQTL reference panels, and it is therefore vital that efforts to collect and analyze these samples continue. TI has exciting advantages for gene discovery as well as downstream applications^{15,55,56}; however, the relative merits of existing methodologies are as yet under-explored. Our analysis suggests that, overall, sparser elastic net models better capture gene expression regulation than BSLMM; at the same time, the improved performance of elastic net over max-eQTL models suggests that a single eQTL model is over-simplified^{2,15}. Fundamentally, transcriptomic imputation methods model only the genetically regulated portion of gene expression, and so cannot capture or interpret variance of expression induced by environment or lifestyle factors, which may be of particular importance in psychiatric disorders. Given the right study design, analyzing genetic components of expression together with observed expression could open doors to better study the role of gene expression in disease.

Sample size and tissue matching contribute to accuracy of TI results. Our CMC-derived DLPFC prediction models had higher average validation R^2 values in external DLPFC data than GTEx-derived brain tissue models. Notably, the model with the second highest percent of genes passing the R^2 threshold is the Thyroid, which has the largest sample size among the GTEx brain

prediction models. When looking at mean R^2 values, the second highest value comes from the GTEx Frontal Cortex, despite the associated small sample size, implying at least some degree of tissue specificity of eQTLs architecture.

We were able to compare TI accuracy in European and African-American individuals, and found that our models were applicable to either ethnicity with only a small decrease in accuracy. Common SNPs shared across ethnicities have important effects on gene expression, and as such we expect GREX to have consistency across populations. There is a well-documented dearth of exploration of genetic associations in non-European cohorts^{57,58} We believe that these analyses should be carried out in non-European cohorts.

We applied the CMC DLPFC prediction models, along with 12 GTEx-derived brain expression prediction models, to schizophrenia cases and controls from the PGC2 and CLOZUK2 collections, constituting the largest transcriptomic analysis of schizophrenia to date. Predicted gene expression levels were calculated for 19,661 unique genes across brain regions (Figure 1C) and tested for association with SCZ case-control status. We identified 413 significant associations, constituting 67 independent associations. We found significant replication of our CMC DLPFC associations in a large independent replication cohort, in collaboration with the iPSYCH-GEMS consortium. A recent TWAS study of 30 GWAS summary statistic traits⁵⁵ identified 38 non-MHC genes associated at tissue-level significance with SCZ in CMC- and GTEx-derived brain tissues (ie, matching those used in our study). Of these, 26 also reach genome-wide significance in our study, although in many instances these genes are not identified as the lead independent associated gene following our conditional analysis. Among our 67 SCZ-associated genes, 19 were novel, i.e. did not fall within 1Mb of a previous GWAS locus (including 5/7 of the novel brain genes identified in the recent TWAS analysis).

We used conditional analyses to identify independent associations within loci. These analyses clarify the most strongly associated genes and tissues (Table 1), while we note that nearly collinear gene-tissue pairs could also represent causal associations. The tissues highlighted allowed us to tabulate apparently independent contributions to SCZ risk from different brain

regions, even though their transcriptomes are highly correlated generally. We find DLPFC and Cerebellum effects, as well as from Putamen, Caudate and Nucleus Accumbens Basal Ganglia.

We used these genic associations to search for enrichments with molecular pathways and gene sets, and identified 36 significant enriched pathways. Among novel pathways, we identified a significant association with HEX-A deficiency. Despite the well-studied and documented symptomatic overlap between adult-onset HEX-A deficiency and schizophrenia, we believe that this is the first demonstration of shared genetics between the disorders. Notably, this overlap is not driven by a single highly-associated gene which is shared by both disorders; rather, every single gene in the HEX-A pathway is nominally significant in the SCZ association analysis, and five genes have $p < 1 \times 10^{-3}$, indicating that there may be substantial shared genetic aetiology between the two disorders that warrants further investigation. Additionally, we identified a significant overlap between our SCZ-associated genes and a number of pathways associated with porphyrin metabolism. Porphyrin disorders have been well characterized and are among early descriptions of “schizophrenic” and psychotic presentations of schizophrenia, as described in the likely eponymous mid-19th century poem “Porphyria’s Lover”, by Robert Browning⁵⁹, and have been cited as a likely diagnosis for the various psychiatric and metabolic ailments of Vincent van Gogh^{60–65} and King George III⁶⁶.

Finally, we assessed patterns of expression for the 67 SCZ-associated genes throughout development using spatio-temporal transcriptomic data obtained from BRAINSPAN. We identified four clusters of genes, with expression in four distinct spatiotemporal regions, ranging from early pre-natal to strictly post-natal expression. There are plausible hypotheses and genetic evidence for SCZ disease development in adolescence, given the correlation with age of onset, as well as prenatally, supported by genetic overlap with neurodevelopmental disorders^{67–69} as well as the earlier onset of cognitive impairments^{70–73}. Understanding the temporal expression patterns of SCZ-associated genes can help to elucidate gene development and trajectory, and inform research and analysis design. Identification of SCZ-associated genes primarily expressed prenatally is striking given our adult eQTL reference panels, and may reflect common eQTL architecture across development, which is known to be partial^{74–76}; therefore, our results should spur interest in extending TI data and/or methods to early development⁷⁴. Identification of SCZ-

associated genes primarily expressed in adolescence and adult-hood is of particular interest for direct analysis of the brain transcriptome in adult psychiatric cases.

eQTL data have been recognized for nearly a decade as potentially important for understanding complex genetic variation. Nicolae et al¹ showed that common variant-common disease associations are strongly enriched for genetic regulation of gene expression. Therefore, integrative approaches combining transcriptomic and genetic association data have great potential. Current TI association analyses increase power for genetic discovery, even while many open areas of TI remain to be developed, such as leveraging additional data types such as chromatin modifications⁷⁷ (e.g. methylation, histone modification), imputing different tissues or different exposures (e.g. age, smoking, trauma) and modeling trans/coexpression effects. It remains critical to leverage TI associations to provide insights into specific disease mechanisms. Here, the accelerated identification of disease associated genes allows the detection of novel pathways and distinct spatiotemporal patterns of expression in schizophrenia risk.

Online Methods (Limit 3,000 words, at end of manuscript, currently 2,064)

Creating gene expression predictors for the dorso-lateral pre-frontal cortex

eQTL Data

Genotype and RNAseq data were obtained for 538 European individuals through the CommonMind Project¹⁴. RNA-seq data were generated from post-mortem human dorsolateral prefrontal cortex (DLPFC). The gene expression matrix was normalized to log(counts per million) using voom. Adjustments were made for known covariates (including sample ascertainment, quality, experimental parameters, ancestry) and surrogate variables, using linear modelling with voom-derived regression weights. Details on genotyping, imputation and RNA-seq generation may be found in the CommonMind Consortium flagship paper¹⁴.

A 1% MAF cut-off was applied. Variants were filtered to remove any SNPs in high LD ($r^2 > 0.9$), indels, and all variants with ambiguous ref/alt alleles. All protein coding genes on chromosomes 1-22 with at least one cis-SNP after these QC steps were included in this analysis. SNPs in trans have been shown not to provide a substantial improvement in prediction accuracy¹⁵ and were not included here.

Building gene expression prediction databases

Gene expression prediction models were created following the “PrediXcan” method¹⁵. Matched genotype and gene expression data were used to identify a set of variants that influence gene expression (Supplementary Figure 2A). Weights for these variants are calculated using regression in a ten-fold cross-validation framework. All cross-validation folds were balanced for diagnoses, ethnicity, and other clinical variables.

All SNPs within the cis-region (± 1 mb) of each gene were included in the regression analysis. Accuracy of prediction was estimated by comparing predicted expression to measured expression, across all 10 cross-validation folds; this correlation was termed cross-validation R^2 or R_{cv}^2 . Genes with $R_{cv}^2 > 0.01$ ($\sim p < 0.05$) were included in our final predictor database.

Prediction models were compared across four different regression methods; elastic net (prediXcan), ridge regression (using the TWAS method¹⁶), Bayesian sparse linear mixed modelling (BSLMM; TWAS), and linear regression using the best eQTL for each gene (Supplementary Figure 1A). Mean R_{cv}^2 values were significantly higher for elastic net regression (mean $R_{cv}^2=0.056$) than for eQTL-based prediction (mean $R_{cv}^2=0.025$), BSLMM (mean $R_{cv}^2=0.021$) or Ridge Regression (mean $R_{cv}^2=0.020$). The distribution of R_{cv}^2 values was also significantly higher for elastic net regression than for any other method (ks-test, $p<2.2\times10^{-16}$).

Replication of gene expression prediction models in independent data

Predictive accuracy of CMC DLPFC models were tested in two independent datasets. First, we used data from the Religious Orders Study and Memory and Aging Project (ROSMAP¹⁹). This study included genotype data and DLPFC RNA-seq data⁷⁸ for 451 individuals of European descent (Supplementary Figure 2B).

DLPFC genetically-regulated expression (GREX) was calculated using the CMC DLPFC predictor models. Correlation between RNA-seq expression and CMC DLPFC GREX (“Replication R^2 values” or R_R^2) was used as a measure of predictive accuracy. R_R^2 was calculated including correction for ten ancestry components, as follows:

Equation 1: R_R^2 calculation.

$$R_{R1}^2 = (M \sim GREX + PC_1 + PC_2 + \dots + PC_{10})$$

$$R_{R2}^2 = (M \sim PC_1 + PC_2 + \dots + PC_{10})$$

$$R_R^2 = R_{R1}^2 - R_{R2}^2$$

Where:

M	Measured expression (RNA-seq)
$GREX$	GREX imputed expression
PC_n	n^{th} Principal Component

A small number of genes (158) had very low predictive accuracy and were removed from further analyses. Cross-validation R^2 (R_{cv}^2) values and R_R^2 values were highly correlated ($\rho=0.62$, $p<2.2e-16$; Supplementary Figure 3A). 55.7% of CMC DLPFC genes had R_R^2 values > 0.01 .

Prediction accuracy was also assessed for 11 publicly available GTEx neurological predictor databases, and R_R^2 values used to compare to CMC DLPFC performance. CMC DLPFC models had higher average R_R^2 values, more genes with $R_R^2 > 0.01$, and significantly higher overall distributions of R_R^2 values than any of the twelve GTEx brain tissue models (ks-test, $p < 2.2e-16$; Figure 1A,B).

To estimate trans-ancestral prediction accuracy, genetically regulated gene expression was calculated for 162 African-American individuals and 280 European individuals from the NIMH Human Brain Collection Core (HBCC) dataset⁷⁹ (Supplementary Figure 2C). Predicted gene expression levels were compared to DLPFC expression levels measured using microarray. There was a significant correlation between the European and African-American samples for R_{CV}^2 values and R_R^2 values ($\rho = 0.66, 0.56$; Supplementary figure 3B-C). R_R^2 values were higher on average in Europeans, but were significantly correlated between African-Americans and Europeans ($\rho = 0.78, p < 2.2e-16$, Pearson test; supplementary figure 3D).

Extension to Summary Statistics

Transcriptomic Imputation may be applied to summary statistics instead of raw dosages, in instances where raw data is unavailable. However, this method suffers from slightly reduced accuracy, requires covariance matrices calculated in an ancestrally-matched reference population⁸⁰ (usually only possible for European cohorts), and precludes testing of endophenotypes within the data, and so should not be applied when raw data is available.

We assessed concordance between CMC DLPFC transcriptomic imputation results using summary-statistics (MetaXcan⁸⁰) and raw genotypes (PrediXcan¹⁵) using nine European and three Asian PGC-SCZ cohorts²² for which both data types were available. Cohorts were chosen to encompass a range of case : control ratios, to test previous suggestions that accuracy is reduced in unbalanced cohorts⁸⁰. Covariances for all variants included in the DLPFC predictor models were computed using MetaXcan⁸⁰. For all European cohorts, Pearson correlation of log-10 p-values and effect sizes was above 0.95. The mean correlation was 0.963 (Supplementary Figure 4). There was no correlation between total sample size, case-control ratio, p-value or

effect-size. Seven genes were removed due to discordant p-values. For the three Asian cohorts tested, the mean correlation was 0.91 (Supplementary Figure 5).

Concordance was also tested for the same nine European PGC-SCZ cohorts, across 12 neurological GTEx prediction databases. All correlations were significant ($\rho > 0.95$, $p < 2.2 \times 10^{-16}$). There was a significant correlation between p-value concordance and case-control ratio ($\rho = 0.37$, $p = 7.606 \times 10^{-15}$). 114 genes had discordant p-values between the two methods and were excluded from future analyses.

Application to Schizophrenia

Dataset Collection

We obtained 53 discovery cohorts for this study, including 40,299 SCZ cases and 65,264 controls (Figure 2). 52/53 cohorts (35,079 cases, 46,441 controls) were obtained through collaboration with the Psychiatric Genomics Consortium, and are described in the 2014 PGC Schizophrenia GWAS²². The remaining cohort, referred to as CLOZUK2, constitutes the largest single cohort of individuals with Schizophrenia (5,220 cases and 18,823 controls), collected as part of an effort to investigate treatment-resistant Schizophrenia²⁶.

50/53 datasets included individuals of European ancestry, while three datasets include individuals of Asian ancestry (1,836 cases, 3,383 controls). All individuals were ancestrally matched to controls. Information on genotyping, quality control and other data management issues may be found in the original papers describing these collections^{22,26}. All sample collections complied with ethical regulations. Details regarding ethical compliance and consent procedures may be found in the original manuscripts describing these collections^{22,26}.

Access to dosage data was available for 44/52 PGC-SCZ cohorts. The remaining PGC cohorts, and the CLOZUK2 cohort provided summary statistics. Three European PGC cohorts were trio-based, rather than case-control.

Additionally, we tested for replication of our CMC DLPFC associations in an independent dataset of 4,133 cases and 24,788 controls obtained through collaboration with the iPSYCH-

GEMS schizophrenia working group (effective sample size 14,169.5; Figure 2B, supplementary information).

Transcriptomic Imputation and association testing

Transcriptomic Imputation was carried out individually for each case-control PGC-SCZ cohort with available dosage data (44/52 cohorts). Predicted gene expression levels were computed using the DLPFC predictors described in this manuscript, as well as for 11 other brain tissues prediction databases created using GTEx tissues^{15,20,21,81} (Figure 1C). Associations between predicted gene expression values and case-control status were calculated using a linear regression test in R. Ten ancestry principal components were included as covariates. Association tests were carried out independently for each cohort, across 12 brain tissues.

For the 8 PGC cohorts with no available dosage data, the three PGC trio-based analyses, and the CLOZUK2 cohort, a summary-statistic based transcriptomic imputation approach was used (“MetaXcan”), as described previously.

Meta-analysis

Meta-analysis was carried out across all 53 cohorts using METAL⁸². Cochran’s Q test for heterogeneity was implemented in METAL^{82,83}, and a heterogeneity p-value threshold of $p > 1 \times 10^{-3}$ applied to results. A conservative significance threshold was applied to these data, correcting for the total number of genes tested across all tissues (121,611 gene-region tests in total). This resulted in a genome-wide significance threshold of 4.1×10^{-7} .

Effect sizes and direction of effect quoted in this manuscript refer to changes in predicted expression in cases compared to controls i.e., genes with negative effect sizes have decreased predicted expression in cases compared to controls.

Identifying independent associations

We identified a number of genomic regions which contained multiple gene associations and/or genes associated across multiple tissues. We identified 58 of these regions, excluding the MHC, based on distance between associated genes, and verified them using visual inspection. In order

to identify independent genic associations within these regions, we carried out a stepwise forward conditional analysis following “GCTA-COJO” theory⁸⁴ using “CoCo” (<https://github.com/theboocock/coco/>), an R implementation of GCTA-COJO. CoCo allows the specification of custom correlation matrices by the user (for example, ancestrally specific LD matrices). For each region, we generated a predicted gene expression correlation matrix for all significant genes ($p \leq 1 \times 10^{-6}$), as the root-effective sample size (N_{eff} , eqn 2) weighted average correlation across all cohorts where we had access to dosage data.

Equation 2: Effective Sample Size, N_{eff}

$$N_{\text{eff}} = \frac{4}{\left(\frac{1}{N_{\text{cases}}} + \frac{1}{N_{\text{controls}}}\right)}$$

Forward stepwise conditional analysis of all significant genes was carried out using joint linear regression modeling. First, the top-ranked gene was added to the model, then the next most significant gene in a joint model is added if significant at a given p-value threshold, and so on until either all genes are added to the model, or no joint statistic reaches the significance threshold.

We calculated effect sizes and odds ratios for SCZ-associated genes by adjusting “CoCo” betas to have unit variance (Table 1, eqn. 3).

Equation 3: GREX Beta adjustment

$$\beta = \beta_{\text{CoCo}} \times \sqrt{\text{GVAR}}$$

Where GVAR is the variance of the GREX predictor for each gene.

Gene set Analyses

Pathway analyses were carried out using an extension to MAGMA⁸⁵. P-values were assigned to genes using the most significant p-value achieved by each gene in the meta-analysis. We then carried out a competitive gene-set analysis test using these p-values, using two gene sets:

1. 159 gene sets with prior hypotheses for involvement in SCZ development, including loss-of-function intolerant genes, CNV-intolerant genes, targets of the fragile-X mental

retardation protein, CNS related gene sets, and 104 behavioural and neurological pathways from the Mouse Genome Informatics database^{14,26,67,86}.

2. An agnostic analysis, including ~8,500 gene sets collated from publicly available databases including GO^{87,88}, KEGG⁸⁹, REACTOME⁹⁰, PANTHER^{91,92}, BIOCARTA⁹³ and MGI⁴⁸. Sets were filtered to include only gene sets with at least ten genes.

Significance levels were adjusted across all pathways included in either test using the Benjamini-Hochberg “FDR” correction in R²³.

Coexpression of SCZ genes throughout development

We investigate spatiotemporal expression of our associated genes using publicly available developmental transcriptome data, obtained from the BRAINSPAN consortium⁹⁴. We partitioned these data into biologically relevant spatio-temporal data sets⁹⁵, corresponding to four general brain regions; the frontal cortex, temporal and parietal regions, sensory-motor regions, and subcortical regions (Figure 4A⁹⁶), and eight developmental time-points (four pre-natal, four post-natal)⁹⁵.

First, we tested for correlation of gene expression for all SCZ-associated genes at each spatiotemporal time-point. Genes with pearson correlation coefficients ≥ 0.8 or ≤ -0.8 were considered co-expressed. 100,000 iterations of this analysis were carried out using random gene sets with equivalent expression level distributions to the SCZ-associated genes. For each gene set, a gene co-expression network was created, with edges connecting all co-expressed genes. Networks were assessed using three criteria; first, the number of edges within the network, as a crude measure of connectedness; second, the Watts-Strogatz average path length between nodes, as a global measure of connectedness across all genes in the network⁵³; third, the Watts-Strogatz clustering coefficient, to measure tightness of the clusters within the network⁵³. For each spatio-temporal time point, we plotted gene-pair expression correlation (suppl. Fig 8) and co-expression networks (suppl. Fig 9).

For each of the 67 SCZ-associated genes, we calculated average expression at each spatiotemporal point. We then calculated Z-Score of expression specificity using these values,

and plotted Z-Scores to visually examine patterns of gene expression throughout development and across brain regions. Clusters were formally identified using a dendrogram cut at height 10 (Suppl. Fig 10).

In-silico replication of SCZ-associated genes in mouse models

We downloaded genotype, knock-out allele information and phenotyping data for ~10,000 mouse mutant models from five large mouse phenotyping and genotyping projects; Mouse Genome Informatics (MGI⁴⁸), EuroPhenome^{47,97}, Mouse Genome Project (MGP^{47,49}), International Mouse Phenotyping Consortium (IMPC⁵⁰), and Infection and Immunity Immunophenotyping (3I⁹⁸). Where possible, we also downloaded raw phenotyping data regarding specific assays. In total, we obtained 175,012 phenotypic measurements, across 10,288 mutant mouse models. We searched for any mouse lines with phenotypes related to behavior (natural, observed, stereotypic or assay-induced); cognition or working memory; brain, head or craniofacial dysmorphology; retinal or eye morphology, and/or vision or visual dysfunction or impairment; ear morphology or hearing dysfunction or impairment; neural tube defects; brain and/or nervous system development; abnormal nociception.

We compared the prevalence of psychiatric phenotypes in mutant mice for our SCZ-associated genes to the prevalence among other disease-associated gene sets. We selected 366 GWAS gene sets, and removed any for which fewer than ten mutant mouse models were included in our databases, leaving 105 gene sets. We compared the prevalence of 13 different categories of psychiatric phenotypes, relating to adrenal gland, behavior, brain development, craniofacial dysmorphology, ear/auditory phenotypes, eye dysmorphology, head dysmorphology, nervous system development, abnormal nociception, seizures, thyroid gland, vision phenotypes. For each GWAS gene set, we counted the number of categories with at least one phenotype, and compared to the number in our SCZ-associated gene set to obtain an empirical p-value.

Data Availability

Our CMC-derived DLPFC prediction models will be made publicly available.

Acknowledgements

Data were generated as part of the CommonMind Consortium supported by funding from Takeda Pharmaceuticals Company Limited, F. Hoffman-La Roche Ltd and NIH grants R01MH085542, R01MH093725, P50MH066392, P50MH080405, R01MH097276, RO1-MH-075916, P50M096891, P50MH084053S1, R37MH057881 and R37MH057881S1, HHSN271201300031C, AG02219, AG05138 and MH06692.

Brain tissue for the study was obtained from the following brain bank collections: the Mount Sinai NIH Brain and Tissue Repository, the University of Pennsylvania Alzheimer's Disease Core Center, the University of Pittsburgh NeuroBioBank and Brain and Tissue Repositories and the NIMH Human Brain Collection Core. CMC Leadership: Pamela Sklar, Joseph Buxbaum (Icahn School of Medicine at Mount Sinai), Bernie Devlin, David Lewis (University of Pittsburgh), Raquel Gur, Chang-Gyu Hahn (University of Pennsylvania), Keisuke Hirai, Hiroyoshi Toyoshiba (Takeda Pharmaceuticals Company Limited), Enrico Domenici, Laurent Essioux (F. Hoffman-La Roche Ltd), Lara Mangravite, Mette Peters (Sage Bionetworks), Thomas Lehner, Barbara Lipska (NIMH).

ROSMAP study data were provided by the Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago. Data collection was supported through funding by NIA grants P30AG10161, R01AG15819, R01AG17917, R01AG30146, R01AG36836, U01AG32984, U01AG46152, the Illinois Department of Public Health, and the Translational Genomics Research Institute.

The iPSYCH-GEMS team would like to acknowledge funding from the Lundbeck Foundation (grant no R102-A9118 and R155-2014-1724), the Stanley Medical Research Institute, an Advanced Grant from the European Research Council (project no: 294838), the Danish Strategic Research Council the Novo Nordisk Foundation for supporting the Danish National Biobank resource, and grants from Aarhus and Copenhagen Universities and University Hospitals, including support to the iSEQ Center, the GenomeDK HPC facility, and the CIRRAU Center.

The Genotype-Tissue Expression (GTEx) Project was supported by the [Common Fund](#) of the Office of the Director of the National Institutes of Health, and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. The data used for the analyses described in this manuscript were obtained from the [GTEx Portal](#) on 09/05/16. BrainSpan: Atlas of the Developing Human Brain [Internet]. Funded by ARRA Awards 1RC2MH089921-01, 1RC2MH090047-01, and 1RC2MH089929-01.

CommonMind Consortium Working Group

Jessica S Johnson 1, Hardik R Shah 2,3, Lambertus L Klein 4, Kristen K Dang 5, Benjamin A Logsdon 5, Milind C Mahajan 2,3, Lara M Mangravite 5, Hiroyoshi Toyoshiba 6, Raquel E Gur 7, Chang-Gyu Hahn 8, Eric Schadt 2,3, David A Lewis 4, Vahram Haroutunian 1,5,9,10, Mette A Peters 5, Barbara K Lipska 11, Joseph D Buxbaum 1, 12, 13, Keisuke Hirai 14, Thanneer M Perumal 5, Laurent Essioux 15,

1. Division of Psychiatric Genomics, Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, New York, USA
2. Institute for Genomics and Multiscale Biology, Icahn School of Medicine at Mount Sinai, New York, New York, USA
3. Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, New York, USA
4. Department of Psychiatry, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania, USA
5. Sage Bionetworks, Seattle, Washington, USA
6. Integrated Technology Research Laboratories, Pharmaceutical Research Division, Takeda Pharmaceutical Company Limited, Fujisawa, Kanagawa, Japan
7. Neuropsychiatry Section, Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA
8. Neuropsychiatric Signaling Program, Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA
9. Psychiatry, JJ Peters Virginia Medical Center, Bronx, New York, USA.
10. Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, New York, USA
11. Human Brain Collection Core, National Institutes of Health, NIMH, Bethesda, Maryland, USA
12. Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, New York, USA.
13. Seaver Autism Center for Research and Treatment, Icahn School of Medicine at Mount Sinai, New York, New York, USA
14. CNS Drug Discovery Unit, Pharmaceutical Research Division, Takeda Pharmaceutical Company Limited, Fujisawa, Kanagawa, Japan
15. F. Hoffman-La Roche Ltd

iPSYCH-GEMS SCZ working group

Anders D. Børglum 1,2,3, Ditte Demontis 1,2,3, Veera Manikandan Rajagopal 1,2,3, Thomas D. Als 1,2,3, Manuel Mattheisen 1,2,3, Jakob Grove 1,2,3,4, Thomas Werge 1,7,8, Preben Bo Mortensen 1,2,9,10, Carsten Bøcker Pedersen 1,9,10, Esben Agerbo 1,9,10, Marianne Giørtz Pedersen 1, 9, 10, Ole Mors 1,6, Merete Nordentoft 1, 11, David M. Hougaard 1,5, Jonas Bybjerg-Grauholm 1,5, Marie Bækvad-Hansen 1,5, Christine Søholm Hansen 1,5

1. iPSYCH, The Lundbeck Foundation Initiative for Integrative Psychiatric Research, Denmark
2. iSEQ, Center for Integrative Sequencing, Aarhus University, Aarhus, Denmark
3. Department of Biomedicine - Human Genetics, Aarhus University, Aarhus, Denmark
4. Bioinformatics Research Centre, Aarhus University, Aarhus, Denmark
5. Center for Neonatal Screening, Department for Congenital Disorders, Statens Serum Institut, Copenhagen, Denmark
6. Psychosis Research Unit, Aarhus University Hospital, Risskov, Denmark
7. Institute of Biological Psychiatry, MHC Sct. Hans, Mental Health Services Copenhagen, Roskilde, Denmark
8. Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark
9. National Centre for Register-Based Research, Aarhus University, Aarhus, Denmark
10. Centre for Integrated Register-based Research, Aarhus University, Aarhus, Denmark
11. Mental Health Services in the Capital Region of Denmark, Mental Health Center Copenhagen, University of Copenhagen, Copenhagen, Denmark

Potential Conflicts of Interest: TW has acted as advisor and lecturer to H. Lundbeck A/S

Schizophrenia Working Group of the Psychiatric Genomics Consortium

Stephan Ripke 1,2, Benjamin M. Neale1,2,3,4, Aiden Corvin 5, James T. R. Walters 6, Kai-How Farh1, Peter A. Holmans 6,7, Phil Lee1,2,4, Brendan Bulik-Sullivan1,2, David A. Collier 8,9, Hailiang Huang 1,3, Tune H. Pers 3,10,11, Ingrid Agartz 12,13,14, Esben Agerbo 15,16,17, Margot Albus 18, Madeline Alexander 19, Farooq Amin 20,21, Silviu A. Bacanu 22, Martin Begemann 23, Richard A. Belliveau Jr 2, Judit Bene 24,25, Sarah E. Bergen 2,26, Elizabeth Bevilacqua 2, Tim B. Bigdeli 22, Donald W. Black 27, Richard Bruggeman28, Nancy G. Buccola29, Randy L. Buckner30,31,32, William Byerley33, Wiepke Cahn34, Guiqing Cai35,36, DominiqueCampion37, Rita M. Cantor38, Vaughan J. Carr39,40, Noa Carrera6, Stanley V. Catts39,41, Kimberly D. Chambert2, Raymond C. K. Chan42, Ronald Y. L. Chen43, Eric Y. H. Chen43,44, Wei Cheng45, Eric F. C. Cheung46, Siow Ann Chong47, C. Robert Cloninger48, David Cohen49, Nadine Cohen50, Paul Cormican5, Nick Craddock6,7, James J. Crowley51, David Curtis52,53, Michael Davidson54, Kenneth L. Davis36, Franziska Degenhardt55,56, Jurgen Del Favero57, Ditte Demontis17,58,59, Dimitris Dikeos60, Timothy Dinan61, Srdjan Djurovic14,62, Gary Donohoe5,63, Elodie Drapeau36, Jubao Duan64,65, Frank Dudbridge66, Naser Durmishi67, Peter Eichhammer68, Johan Eriksson69,70,71, Valentina Escott-Price6, Laurent Essioux72, Ayman H. Fanous73,74,75,76, Martilias S. Farrell51, Josef Frank77, Lude Franke78, Robert Freedman79, Nelson B. Freimer80, Marion Friedl81, Joseph I. Friedman36, Menachem Fromer1,2,4,82, Giulio Genovese2, Lyudmila Georgieva6, Ina Giegling81,83, Paola Giusti-Rodri'guez51, Stephanie Godard84, Jacqueline I. Goldstein1,3, Vera Golimbet85, Srihari Gopal86, Jacob Gratten87, Lieuwe de Haan88, Christian Hammer23, Marian L. Hamshere6, Mark Hansen89, Thomas Hansen17,90, Vahram Haroutunian36,91,92, Annette M. Hartmann81, Frans A. Henskens39,93,94, Stefan Herms55,56,95, Joel N. Hirschhorn3,11,96, Per Hoffmann55,56,95, Andrea Hofman55,56, Mads V. Hollegaard97, David M. Hougaard97, Masashi Ikeda98, Inge Joa99, Antonio Julia 100, Rene S. Kahn34, Luba Kalaydjieva101,102, Sena Karachanak-Yankova103, Juha Karjalainen78, David Kavanagh6, Matthew C. Keller104, James L. Kennedy105,106,107, Andrey Khrunin108, Yunjung Kim51, Janis Klovins109, James A. Knowles110, Bettina Konte81, Vaidutis Kucinskas111, Zita Ausrele Kucinskiene111, Hana Kuzelova-Ptackova112, Anna K. Kahler26, Claudine Laurent19,113, Jimmy Lee Chee Keong47,114, S. Hong Lee87, Sophie E. Legge6, Bernard Lerer115, Miaoxin Li43,44,116, Tao Li117, Kung-Yee Liang118, Jeffrey Lieberman119, Svetlana Limborska108, Carmel M. Loughland39,120, Jan Lubinski121, Jouko Lonnqvist122, Milan Macek Jr112, Patrik K. E. Magnusson26, Brion S. Maher123, Wolfgang Maier124, Jacques Mallet125, Sara Marsal100, Manuel Mattheisen17,58,59,126, Morten Mattingsdal14,127, Robert W. McCarley128,129, ColmMcDonald130, Andrew M. McIntosh131,132, Sandra Meier77, Carin J. Meijer88, Bela Meleg24,25, Ingrid Melle14,133, Raquelle I. Mesholam-Gately128,134, Andres Metspalu135, Patricia T. Michie39,136, Lili Milani135, Vihra Milanova137, Younes Mokrab8, Derek W. Morris5,63, Ole Mors17,58,138, Kieran C. Murphy139, Robin M. Murray140, Inez Myin-Germeys141, Bertram Muller-Myhsok142,143,144, Mari Nelis135, Igor Nenadic145, Deborah A. Nertney146, Gerald Nestadt147, Kristin K. Nicodemus148, Liene Nikitina-Zake109, Laura

Nisenbaum¹⁴⁹, Annelie Nordin¹⁵⁰, Eadbhard O’Callaghan¹⁵¹, Colm O’Dushlaine², F. Anthony O’Neill¹⁵², Sang-Yun Oh¹⁵³, Ann Olincy⁷⁹, Line Olsen^{17,90}, Jim Van Os^{141,154}, Psychosis Endophenotypes International Consortium¹⁵⁵, Christos Pantelis^{39,156}, George N. Papadimitriou⁶⁰, Sergi Papiol²³, Elena Parkhomenko³⁶, Michele T. Pato¹¹⁰, Tiina Paunio^{157,158}, Milica Pejovic-Milovancevic¹⁵⁹, Diana O. Perkins¹⁶⁰, Olli Pietiläinen^{158,161}, Jonathan Pimm⁵³, Andrew J. Pocklington⁶, John Powell¹⁴⁰, Alkes Price^{3,162}, Ann E. Pulver¹⁴⁷, Shaun M. Purcell⁸², Digby Quested¹⁶³, Henrik B. Rasmussen^{17,90}, Abraham Reichenberg³⁶, Mark A. Reimers¹⁶⁴, Alexander L. Richards⁶, Joshua L. Roffman^{30,32}, Panos Roussos^{82,165}, Douglas M. Ruderfer^{6,82}, Veikko Salomaa⁷¹, Alan R. Sanders^{64,65}, Ulrich Schall^{39,120}, Christian R. Schubert¹⁶⁶, Thomas G. Schulze^{77,167}, Sibylle G. Schwab¹⁶⁸, Edward M. Scolnick², Rodney J. Scott^{39,169,170}, Larry J. Seidman^{128,134}, Jianxin Shi¹⁷¹, Engilbert Sigurdsson¹⁷², Teimuraz Silagadze¹⁷³, Jeremy M. Silverman^{36,174}, Kang Sim⁴⁷, Petr Slominsky¹⁰⁸, Jordan W. Smoller^{2,4}, Hon-Cheong So⁴³, Chris C. A. Spencer¹⁷⁵, Eli A. Stahl^{3,82}, Hreinn Stefansson¹⁷⁶, Stacy Steinberg¹⁷⁶, Elisabeth Stogmann¹⁷⁷, Richard E. Straub¹⁷⁸, Eric Strengman^{179,34}, Jana Strohmaier⁷⁷, T. Scott Stroup¹¹⁹, Mythily Subramaniam⁴⁷, Jaana Suvisaari¹²², Dragan M. Svrakic⁴⁸, Jin P. Szatkiewicz⁵¹, Erik Soderman¹², Srinivas Thirumalai¹⁸⁰, Draga Toncheva¹⁰³, Sarah Tosato¹⁸¹, Juha Veijola^{182,183}, John Waddington¹⁸⁴, Dermot Walsh¹⁸⁵, Dai Wang⁸⁶, Qiang Wang¹¹⁷, Bradley T. Webb²², Mark Weiser⁵⁴, Dieter B. Wildenauer¹⁸⁶, Nigel M. Williams⁶, Stephanie Williams⁵¹, Stephanie H. Witt⁷⁷, Aaron R. Wolen¹⁶⁴, Emily H. M. Wong⁴³, Brandon K. Wormley²², Hualin Simon Xi¹⁸⁷, Clement C. Zai^{105,106}, Xuebin Zheng¹⁸⁸, Fritz Zimprich¹⁷⁷, Naomi R. Wray⁸⁷, Kari Stefansson¹⁷⁶, Peter M. Visscher⁸⁷, Wellcome Trust Case-Control Consortium²¹⁸⁹, Rolf Adolfsson¹⁵⁰, Ole A. Andreassen^{14,133}, Douglas H. R. Blackwood¹³², Elvira Bramon¹⁹⁰, Joseph D. Buxbaum^{35,36,91,191}, Anders D. Børglum^{17,58,59,138}, Sven Cichon^{55,56,95,192}, Ariel Darvasi¹⁹³, Enrico Domenici¹⁹⁴, Hannelore Ehrenreich²³, Tonu Esko^{3,11,96,135}, Pablo V. Gejman^{64,65}, Michael Gill⁵, Hugh Gurling⁵³, Christina M. Hultman²⁶, Nakao Iwata⁹⁸, Assen V. Jablensky^{39,102,186,195}, Erik G. Jonsson^{12,14}, Kenneth S. Kendler¹⁹⁶, George Kirov⁶, Jo Knight^{105,106,107}, Todd Lencz^{197,198,199}, Douglas F. Levinson¹⁹, Qing Qin S. Li⁸⁶, Jianjun Liu^{188,200}, Anil K. Malhotra^{197,198,199}, Steven A. McCarroll^{2,96}, Andrew McQuillin⁵³, Jennifer L. Moran², Preben B. Mortensen^{15,16,17}, Bryan J. Mowry^{87,201}, Markus M. Nothen^{55,56}, Roel A. Ophoff^{38,80,34}, Michael J. Owen^{6,7}, Aarno Palotie^{2,4,161}, Carlos N. Pato¹¹⁰, Tracey L. Petryshen^{2,128,202}, Danielle Posthuma^{203,204,205}, Marcella Rietschel⁷⁷, Brien P. Riley¹⁹⁶, Dan Rujescu^{81,83}, Pak C. Sham^{43,44,116}, Pamela Sklar^{82,91,165}, David St Clair²⁰⁶, Daniel R. Weinberger^{178,207}, Jens R. Wendland¹⁶⁶, Thomas Werge^{17,90,208}, Mark J. Daly^{1,2,3}, Patrick F. Sullivan^{26,51,160} & Michael C. O’Donovan^{6,7}

1. Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, Massachusetts 02114, USA. 2. Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, Massachusetts 02142, USA. 3. Medical and Population Genetics

841 Program, Broad Institute of MIT and Harvard, Cambridge, Massachusetts 02142, USA. 4.
842 Psychiatric and Neurodevelopmental Genetics Unit, Massachusetts General Hospital, Boston,
843 Massachusetts 02114, USA. 5. Neuropsychiatric Genetics Research Group, Department of
844 Psychiatry, Trinity College Dublin, Dublin 8, Ireland. 6. MRC Centre for Neuropsychiatric
845 Genetics and Genomics, Institute of Psychological Medicine and ClinicalNeurosciences, School
846 of Medicine, Cardiff University, Cardiff CF24 4HQ,UK. 7. NationalCentre for Mental Health,
847 Cardiff University, Cardiff CF244HQ,UK. 8. Eli Lilly and Company Limited, Erl Wood Manor,
848 Sunninghill Road, Windlesham, Surrey GU20 6PH, UK. 9. Social, Genetic and Developmental
849 Psychiatry Centre, Institute of Psychiatry, King’s College London, London SE58AF,UK. 10.
850 Center for BiologicalSequence Analysis, Department of Systems Biology, Technical University
851 of Denmark, DK-2800, Denmark. 11. Division of Endocrinology and Center for Basic and
852 Translational Obesity Research, Boston Children’s Hospital, Boston, Massachusetts 02115,USA.
853 12. Department of Clinical Neuroscience, Psychiatry Section, Karolinska Institutet, SE-17176
854 Stockholm, Sweden. 13. Department of Psychiatry, Diakonhjemmet Hospital, 0319 Oslo,
855 Norway. 14. NORMENT, KG Jebsen Centre for Psychosis Research, Institute of Clinical
856 Medicine, University of Oslo, 0424 Oslo, Norway. 15. Centre for Integrative Register-based
857 Research, CIRRAU, Aarhus University, DK-8210 Aarhus, Denmark. 16. National Centre for
858 Register-based Research, Aarhus University, DK-8210Aarhus,Denmark. 17. The Lundbeck
859 Foundation Initiative for Integrative Psychiatric Research, iPSYCH, Denmark. 18. State Mental
860 Hospital, 85540 Haar, Germany. 19. Department of Psychiatry and Behavioral Sciences,
861 Stanford University, Stanford, California 94305, USA. 20. Department of Psychiatry and
862 Behavioral Sciences, Atlanta Veterans Affairs Medical Center, Atlanta, Georgia 30033, USA.
863 21. Department of Psychiatry and Behavioral Sciences, Emory University, Atlanta, Georgia
864 30322, USA. 22. Virginia Institute for Psychiatric and Behavioral Genetics, Department of
865 Psychiatry, Virginia Commonwealth University, Richmond, Virginia 23298, USA. 23. Clinical
866 Neuroscience, Max Planck Institute of Experimental Medicine, Gottingen 37075, Germany. 24.
867 Department of Medical Genetics, University of Pécs, Pécs H-7624, Hungary. 25. Szentagothai
868 Research Center, University of Pécs, Pécs H-7624, Hungary. 26. Department of Medical
869 Epidemiology and Biostatistics, Karolinska Institutet, Stockholm SE-17177, Sweden. 27.
870 Department of Psychiatry, University of Iowa Carver College of Medicine, Iowa City, Iowa
871 52242, USA. 28. University Medical Center Groningen, Department of Psychiatry, University of
872 Groningen NL-9700 RB, The Netherlands. 29. School of Nursing, Louisiana State University
873 Health Sciences Center, New Orleans, Louisiana 70112, USA. 30. Athinoula A. Martinos
874 Center, Massachusetts General Hospital, Boston, Massachusetts 02129, USA. 31. Center for
875 Brain Science, Harvard University, Cambridge, Massachusetts 02138, USA. 32. Department of
876 Psychiatry, Massachusetts General Hospital, Boston, Massachusetts 02114, USA. 33.
877 Department of Psychiatry, University of California at San Francisco, San Francisco, California
878 94143, USA. 34. University Medical Center Utrecht, Department of Psychiatry, Rudolf Magnus
879 Institute of Neuroscience, 3584 Utrecht, The Netherlands. 35. Department of Human

880 Genetics, Icahn School of Medicine at Mount Sinai, New York, New York 10029, USA. 36.
881 Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, New York
882 10029, USA. 37. Centre Hospitalier du Rouvray and INSERM U1079 Faculty of Medicine,
883 76301 Rouen, France. 38. Department of Human Genetics, David Geffen School of Medicine,
884 University of California, Los Angeles, California 90095, USA. 39. Schizophrenia Research
885 Institute, Sydney NSW2010, Australia. 40. School of Psychiatry, University of New
886 South Wales, Sydney NSW2031, Australia. 41. Royal Brisbane and Women's Hospital,
887 University of Queensland, Brisbane, St Lucia QLD 4072, Australia. 42. Institute of Psychology,
888 Chinese Academy of Science, Beijing 100101, China. 43. Department of Psychiatry, Li Ka Shing
889 Faculty of Medicine, The University of Hong Kong, Hong Kong, China. 44. State Key
890 Laboratory for Brain and Cognitive Sciences, Li Ka Shing Faculty of Medicine, The University
891 of Hong Kong, Hong Kong, China. 45. Department of Computer Science, University of North
892 Carolina, Chapel Hill, North Carolina 27514, USA. 46. Castle Peak Hospital, Hong Kong, China.
893 47. Institute of Mental Health, Singapore 539747, Singapore. 48. Department of Psychiatry,
894 Washington University, St. Louis, Missouri 63110, USA. 49. Department of Child and
895 Adolescent Psychiatry, Assistance Publique Hopitaux de Paris, Pierre and Marie Curie Faculty of
896 Medicine and Institute for Intelligent Systems and Robotics, Paris 75013, France. 50. Blue Note
897 Biosciences, Princeton, New Jersey 08540, USA. 51. Department of Genetics, University of
898 North Carolina, Chapel Hill, North Carolina 27599-7264, USA. 52. Department of Psychological
899 Medicine, Queen Mary University of London, London E1 1BB, UK. 53. Molecular Psychiatry
900 Laboratory, Division of Psychiatry, University College London, London WC1E 6JJ, UK. 54.
901 Sheba Medical Center, Tel Hashomer 52621, Israel. 55. Department of Genomics, Life and Brain
902 Center, D-53127 Bonn, Germany. 56. Institute of Human Genetics, University of Bonn, D-53127
903 Bonn, Germany. 57. Applied Molecular Genomics Unit, VIB Department of Molecular Genetics,
904 University of Antwerp, B-2610 Antwerp, Belgium. 58. Centre for Integrative Sequencing, iSEQ,
905 Aarhus University, DK-8000 Aarhus C, Denmark. 59. Department of Biomedicine, Aarhus
906 University, DK-8000 Aarhus C, Denmark. 60. First Department of Psychiatry, University of
907 Athens Medical School, Athens 11528, Greece. 61. Department of Psychiatry, University
908 College Cork, Co. Cork, Ireland. 62. Department of Medical Genetics, Oslo University Hospital,
909 0424 Oslo, Norway. 63. Cognitive Genetics and Therapy Group, School of Psychology and
910 Discipline of Biochemistry, National University of Ireland Galway, Co. Galway, Ireland. 64.
911 Department of Psychiatry and Behavioral Neuroscience, University of Chicago, Chicago, Illinois
912 60637, USA. 65. Department of Psychiatry and Behavioral Sciences, North Shore University
913 Health System, Evanston, Illinois 60201, USA. 66. Department of Non-Communicable Disease
914 Epidemiology, London School of Hygiene and Tropical Medicine, London WC1E 7HT, UK. 67.
915 Department of Child and Adolescent Psychiatry, University Clinic of Psychiatry, Skopje 1000,
916 Republic of Macedonia. 68. Department of Psychiatry, University of Regensburg, 93053
917 Regensburg, Germany. 69. Department of General Practice, Helsinki University Central Hospital,
918 University of Helsinki P.O. Box 20, Tukholmankatu 8 B, FI-00014, Helsinki, Finland. 70.
919 Folkhälsan Research Center, Helsinki, Finland, Biomedicum Helsinki 1, Haartmaninkatu 8, FI-

00290, Helsinki, Finland. 71. National Institute for Health and Welfare, P.O. Box 30, FI-00271
Helsinki, Finland. 72. Translational Technologies and Bioinformatics, Pharma Research
and Early Development, F. Hoffman-La Roche, CH-4070 Basel, Switzerland. 73. Department of
Psychiatry, Georgetown University School of Medicine, Washington DC 20057, USA. 74.
Department of Psychiatry, Keck School of Medicine of the University of Southern California,
Los Angeles, California 90033, USA. 75. Department of Psychiatry, Virginia Commonwealth
University School of Medicine, Richmond, Virginia 23298, USA. 76. Mental Health Service
Line, Washington VA Medical Center, Washington DC 20422, USA. 77. Department of Genetic
Epidemiology in Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim,
University of Heidelberg, Heidelberg, D-68159 Mannheim, Germany. 78. Department of
Genetics, University of Groningen, University Medical Centre Groningen, 9700 RB Groningen,
The Netherlands. 79. Department of Psychiatry, University of Colorado Denver, Aurora,
Colorado 80045, USA. 80. Center for Neurobehavioral Genetics, Semel Institute for
Neuroscience and Human Behavior, University of California, Los Angeles, California 90095,
USA. 81. Department of Psychiatry, University of Halle, 06112 Halle, Germany. 82. Division of
Psychiatric Genomics, Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New
York, New York 10029, USA. 83. Department of Psychiatry, University of Munich, 80336,
Munich, Germany. 84. Departments of Psychiatry and Human and Molecular Genetics,
INSERM, Institut de Myologie, Hôpital de la Pitié-Salpêtrière, Paris 75013, France. 85. Mental
Health Research Centre, Russian Academy of Medical Sciences, 115522 Moscow, Russia. 86.
Neuroscience Therapeutic Area, Janssen Research and Development, Raritan, New
Jersey 08869, USA. 87. Queensland Brain Institute, The University of Queensland, Brisbane,
Queensland, QLD 4072, Australia. 88. Academic Medical Centre University of Amsterdam,
Department of Psychiatry, 1105 AZ Amsterdam, The Netherlands. 89. Illumina, La Jolla,
California, California 92122, USA. 90. Institute of Biological Psychiatry, Mental Health Centre
Sct. Hans, Mental Health Services Copenhagen, DK-4000, Denmark. 91. Friedman Brain
Institute, Icahn School of Medicine at Mount Sinai, New York, New York 10029, USA. 92. J. J.
Peters VA Medical Center, Bronx, New York, New York 10468, USA. 93. Priority Research
Centre for Health Behaviour, University of Newcastle, Newcastle NSW 2308, Australia. 94.
School of Electrical Engineering and Computer Science, University of Newcastle, Newcastle
NSW 2308, Australia. 95. Division of Medical Genetics, Department of Biomedicine, University
of Basel, Basel CH-4058, Switzerland. 96. Department of Genetics, Harvard Medical School,
Boston, Massachusetts, Massachusetts 02115, USA. 97. Section of Neonatal Screening and
Hormones, Department of Clinical Biochemistry, Immunology and Genetics, Statens Serum
Institut, Copenhagen DK-2300, Denmark. 98. Department of Psychiatry, Fujita Health
University School of Medicine, Toyoake, Aichi, 470-1192, Japan. 99. Regional Centre for
Clinical Research in Psychosis, Department of Psychiatry, Stavanger University Hospital, 4011
Stavanger, Norway. 100. Rheumatology Research Group, Vall d'Hebron Research Institute,
Barcelona 08035, Spain. 101. Centre for Medical Research, The University of Western Australia,
Perth WA6009, Australia. 102. The Perkins Institute for Medical Research, The University of

960 Western Australia, Perth WA6009, Australia. 103. Department of Medical Genetics, Medical
 961 University, Sofia 1431, Bulgaria. 104. Department of Psychology, University of Colorado
 962 Boulder, Boulder, Colorado 80309, USA. 105. Campbell Family Mental Health Research
 963 Institute, Centre for Addiction and Mental Health, Toronto, Ontario M5T 1R8, Canada. 106.
 964 Department of Psychiatry, University of Toronto, Toronto, Ontario M5T 1R8, Canada. 107.
 965 Institute of Medical Science, University of Toronto, Toronto, Ontario M5S1A8, Canada. 108.
 966 Institute of Molecular Genetics, Russian Academy of Sciences, Moscow 123182, Russia. 109.
 967 Latvian Biomedical Research and Study Centre, Riga, LV-1067, Latvia. 110. Department of
 968 Psychiatry and Zilkha Neurogenetics Institute, Keck School of Medicine at University of
 969 Southern California, Los Angeles, California 90089, USA. 111. Faculty of Medicine, Vilnius
 970 University, LT-01513 Vilnius, Lithuania. 112. Department of Biology and Medical Genetics,
 971 2nd Faculty of Medicine and University Hospital Motol, 150 06 Prague, Czech Republic. 113.
 972 Department of Child and Adolescent Psychiatry, Pierre and Marie Curie Faculty of Medicine,
 973 Paris 75013, France. 114. Duke-NUS Graduate Medical School, Singapore 169857. 115.
 974 Department of Psychiatry, Hadassah-Hebrew University Medical Center, Jerusalem 91120,
 975 Israel. 116. Centre for Genomic Sciences, The University of Hong Kong, Hong Kong, China.
 976 117. Mental Health Centre and Psychiatric Laboratory, West China Hospital, Sichuan
 977 University, Chengdu, 610041 Sichuan, China. 118. Department of Biostatistics, Johns Hopkins
 978 University Bloomberg School of Public Health, Baltimore, Maryland 21205, USA. 119.
 979 Department of Psychiatry, Columbia University, New York, New York 10032, USA. 120.
 980 Priority Centre for Translational Neuroscience and Mental Health, University of Newcastle,
 981 Newcastle NSW 2300, Australia. 121. Department of Genetics and Pathology, International
 982 Hereditary Cancer Center, Pomeranian Medical University in Szczecin, 70-453 Szczecin,
 983 Poland. 122. Department of Mental Health and Substance Abuse Services; National Institute for
 984 Health and Welfare, P.O.BOX30, FI-00271 Helsinki, Finland. 123. Department of Mental Health,
 985 Bloomberg School of Public Health, Johns Hopkins University, Baltimore, Maryland 21205,
 986 USA. 124. Department of Psychiatry, University of Bonn, D-53127 Bonn, Germany. 125. Centre
 987 National de la Recherche Scientifique, Laboratoire de Génétique Moléculaire de la
 988 Neurotransmission et des Processus Neurodégénératifs, Hôpital de la Pitié-Salpêtrière 75013
 989 Paris, France. 126. Department of Genomics Mathematics, University of Bonn, D-53127 Bonn,
 990 Germany. 127. Research Unit, Sørlandet Hospital, 4604 Kristiansand, Norway. 128. Department
 991 of Psychiatry, Harvard Medical School, Boston, Massachusetts 02115, USA. 129. VA Boston
 992 Health Care System, Brockton, Massachusetts 02301, USA. 130. Department of Psychiatry,
 993 National University of Ireland Galway, Co. Galway, Ireland. 131. Centre for Cognitive Ageing
 994 and Cognitive Epidemiology, University of Edinburgh, Edinburgh EH16 4SB, UK. 132. Division
 995 of Psychiatry, University of Edinburgh, Edinburgh EH16 4SB, UK. 133. Division of Mental
 996 Health and Addiction, Oslo University Hospital, 0424 Oslo, Norway. 134. Massachusetts Mental
 997 Health Center Public Psychiatry Division of the Beth Israel Deaconess Medical Center, Boston,
 998 Massachusetts 02114, USA. 135. Estonian Genome Center, University of Tartu, Tartu 50090,
 999 Estonia. 136. School of Psychology, University of Newcastle, Newcastle NSW2308, Australia.

1000 137. First Psychiatric Clinic, Medical University, Sofia 1431, Bulgaria. 138. Department P,
1001 Aarhus University Hospital, DK-8240 Risskov, Denmark. 139. Department of Psychiatry, Royal
1002 College of Surgeons in Ireland, Dublin 2, Ireland. 140. King's College London, London
1003 SE58AF, UK. 141. Maastricht University Medical Centre, South Limburg Mental Health
1004 Research and Teaching Network, EURON, 6229HX Maastricht, The Netherlands. 142. Institute of
1005 Translational Medicine, University of Liverpool, Liverpool L69 3BX, UK. 143. Max Planck
1006 Institute of Psychiatry, 80336 Munich, Germany. 144. Munich Cluster for Systems Neurology
1007 (SyNergy), 80336 Munich, Germany. 145. Department of Psychiatry and Psychotherapy, Jena
1008 University Hospital, 07743 Jena, Germany. 146. Department of Psychiatry, Queensland Brain
1009 Institute and Queensland Centre for Mental Health Research, University of Queensland,
1010 Brisbane, Queensland, St Lucia QLD 4072, Australia. 147. Department of Psychiatry and
1011 Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, Maryland
1012 21205, USA. 148. Department of Psychiatry, Trinity College Dublin, Dublin 2, Ireland. 149. Eli
1013 Lilly and Company, Lilly Corporate Center, Indianapolis, 46285 Indiana, USA. 150. Department
1014 of Clinical Sciences, Psychiatry, Umeå University, SE-901 87 Umeå, Sweden. 151. DETECT
1015 Early Intervention Service for Psychosis, Blackrock, Co. Dublin, Ireland. 152. Centre for Public
1016 Health, Institute of Clinical Sciences, Queen's University Belfast, Belfast BT12 6AB, UK. 153.
1017 Lawrence Berkeley National Laboratory, University of California at Berkeley, Berkeley,
1018 California 94720, USA. 154. Institute of Psychiatry, King's College London, London SE5
1019 8AF, UK. 155. A list of authors and affiliations appear in the Supplementary Information. 156.
1020 Melbourne Neuropsychiatry Centre, University of Melbourne & Melbourne Health, Melbourne,
1021 Vic 3053, Australia. 157. Department of Psychiatry, University of Helsinki, P.O. Box 590, FI-
1022 00029 HUS, Helsinki, Finland. 158. Public Health Genomics Unit, National Institute for Health
1023 and Welfare, P.O. BOX 30, FI-00271 Helsinki, Finland. 159. Medical Faculty, University of
1024 Belgrade, 11000 Belgrade, Serbia. 160. Department of Psychiatry, University of North Carolina,
1025 Chapel Hill, North Carolina 27599-7160, USA. 161. Institute for Molecular Medicine Finland,
1026 FIMM, University of Helsinki, P.O. Box 20 FI-00014, Helsinki, Finland. 162. Department of
1027 Epidemiology, Harvard School of Public Health, Boston, Massachusetts 02115, USA. 163.
1028 Department of Psychiatry, University of Oxford, Oxford, OX3 7JX, UK. 164. Virginia Institute
1029 for Psychiatric and Behavioral Genetics, Virginia Commonwealth University, Richmond,
1030 Virginia 23298, USA. 165. Institute for Multiscale Biology, Icahn School of Medicine at Mount
1031 Sinai, New York, New York 10029, USA. 166. Pharma Therapeutics Clinical Research, Pfizer
1032 Worldwide Research and Development, Cambridge, Massachusetts 02139, USA. 167.
1033 Department of Psychiatry and Psychotherapy, University of Göttingen, 37073 Göttingen,
1034 Germany. 168. Psychiatry and Psychotherapy Clinic, University of Erlangen, 91054 Erlangen,
1035 Germany. 169. Hunter New England Health Service, Newcastle NSW2308, Australia. 170. School
1036 of Biomedical Sciences, University of Newcastle, Newcastle NSW2308, Australia. 171. Division
1037 of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, Maryland 20892,
1038 USA. 172. University of Iceland, Landspítali, National University Hospital, 101 Reykjavik,
1039 Iceland. 173. Department of Psychiatry and Drug Addiction, Tbilisi State Medical University

1040 (TSMU), N33,0177 Tbilisi, Georgia. 174. Research and Development, Bronx Veterans Affairs
1041 Medical Center, New York, New York 10468, USA. 175. WellcomeTrust Centre for Human
1042 Genetics, Oxford OX3 7BN, UK. 176. deCODE Genetics, 101 Reykjavik, Iceland. 177/
1043 Department of Clinical Neurology, Medical University of Vienna, 1090 Wien, Austria. 178.
1044 Lieber Institute for Brain Development, Baltimore, Maryland 21205, USA. 179. Department of
1045 Medical Genetics, University Medical Centre Utrecht, Universiteitsweg 100,3584CG, Utrecht,
1046 The Netherlands. 180. Berkshire Healthcare NHS Foundation Trust, Bracknell RG12 1BQ, UK.
1047 181. Section of Psychiatry, University of Verona, 37134 Verona, Italy. 182. Department of
1048 Psychiatry, University of Oulu, P.O. Box 5000, 90014, Finland. 183. University Hospital of
1049 Oulu, P.O. Box 20, 90029 OYS, Finland. 184. Molecular and Cellular Therapeutics, Royal
1050 College of Surgeons in Ireland, Dublin 2, Ireland. 185. Health Research Board, Dublin 2,
1051 Ireland. 186. School of Psychiatry and Clinical Neurosciences, The University of Western
1052 Australia, Perth WA6009, Australia. 187. Computational Sciences CoE, Pfizer Worldwide
1053 Research and Development, Cambridge, Massachusetts 02139, USA. 188. Human Genetics,
1054 Genome Institute of Singapore, A*STAR, Singapore 138672. 189. A list of authors and
1055 affiliations appear in the Supplementary Information. 190. University College London, London
1056 WC1E 6BT, UK. 191. Department of Neuroscience, Icahn School of Medicine at Mount
1057 Sinai, New York, New York 10029, USA. 192. Institute of Neuroscience and Medicine (INM-1),
1058 Research Center Juelich, 52428 Juelich, Germany. 193. Department of Genetics, The Hebrew
1059 University of Jerusalem, 91905 Jerusalem, Israel. 194. NeuroscienceDiscovery and Translational
1060 Area, Pharma Research and Early Development, F. Hoffman-La Roche, CH-4070 Basel,
1061 Switzerland. 195. Centre for Clinical Research in Neuropsychiatry, School of Psychiatry and
1062 Clinical Neurosciences, The University of Western Australia, Medical Research Foundation
1063 Building, Perth WA6000, Australia. 196. Virginia Institute for Psychiatric and Behavioral
1064 Genetics, Departments of Psychiatry and Human and Molecular Genetics, Virginia
1065 Commonwealth University, Richmond, Virginia 23298, USA. 197. The Feinstein Institute for
1066 Medical Research, Manhasset, New York 11030, USA. 198. The Hofstra NS-LIJ School of
1067 Medicine, Hempstead, New York 11549, USA. 199. The Zucker Hillside Hospital, Glen Oaks,
1068 New York 11004, USA. 200. Saw Swee Hock School of Public Health, National University of
1069 Singapore, Singapore 117597, Singapore. 201. Queensland Centre for Mental Health Research,
1070 University of Queensland, Brisbane 4076, Queensland, Australia. 202. Center for Human Genetic
1071 Research and Department of Psychiatry, Massachusetts General Hospital, Boston, Massachusetts
1072 02114, USA. 203. Department of Child and Adolescent Psychiatry, Erasmus University Medical
1073 Centre, Rotterdam 3000, The Netherlands. 204. Department of Complex Trait Genetics,
1074 Neuroscience Campus Amsterdam, VU University Medical Center Amsterdam,
1075 Amsterdam 1081, The Netherlands. 205. Department of Functional Genomics, Center for
1076 Neurogenomics and Cognitive Research, Neuroscience Campus Amsterdam, VU University,
1077 Amsterdam 1081, The Netherlands. 206. University of Aberdeen, Institute of Medical Sciences,
1078 Aberdeen AB25 2ZD, UK. 207. Departments of Psychiatry, Neurology, Neuroscience and
1079 Institute of Genetic Medicine, Johns Hopkins School of Medicine, Baltimore, Maryland 21205,

1080 USA. 208. Department of Clinical Medicine, University of Copenhagen, Copenhagen 2200,
1081 Denmark.

References

1. Nicolae, D. L. *et al.* Trait-Associated SNPs Are More Likely to Be eQTLs: Annotation to Enhance Discovery from GWAS. *PLoS Genet.* **6**, e1000888 (2010).
2. Dobbyn, A. *et al.* Co-localization of Conditional eQTL and GWAS Signatures in Schizophrenia. *Rev.* (2017).
3. Gilad, Y., Rifkin, S. A. & Pritchard, J. K. Revealing the architecture of gene regulation: the promise of eQTL studies. *Trends Genet.* **24**, 408–415 (2008).
4. Cookson, W., Liang, L., Abecasis, G., Moffatt, M. & Lathrop, M. Mapping complex disease traits with global gene expression. *Nat. Rev. Genet.* **10**, 184–194 (2009).
5. Albert, F. W. & Kruglyak, L. The role of regulatory variation in complex traits and disease. *Nat. Rev. Genet.* **16**, 197–212 (2015).
6. Moffatt, M. F. *et al.* Genetic variants regulating ORMDL3 expression contribute to the risk of childhood asthma. *Nature* **448**, 470–473 (2007).
7. Speliotes, E. K. *et al.* Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat. Genet.* **42**, 937–948 (2010).
8. Dubois, P. C. A. *et al.* Multiple common variants for celiac disease influencing immune gene expression. *Nat. Genet.* **42**, 295–302 (2010).
9. Libioulle, C. *et al.* Novel Crohn disease locus identified by genome-wide association maps to a gene desert on 5p13.1 and modulates expression of PTGER4. *PLoS Genet.* **3**, e58 (2007).
10. Giambartolomei, C. *et al.* Bayesian Test for Colocalisation between Pairs of Genetic Association Studies Using Summary Statistics. *PLoS Genet.* **10**, e1004383 (2014).
11. Boocock, J., Giambartolomei, C. & Stahl, E. A. COLOC2. (2016).
12. Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nat. Genet.* **48**, 481–487 (2016).
13. Pickrell, J. K. *et al.* Detection and interpretation of shared genetic influences on 42 human traits. *Nat. Genet.* **48**, 709–717 (2016).
14. Fromer, M. *et al.* Gene expression elucidates functional impact of polygenic risk for schizophrenia. *Nat. Neurosci.* **19**, 1442–1453 (2016).
15. Gamazon, E. R. *et al.* A gene-based association method for mapping traits using reference transcriptome data. *Nat. Genet.* **47**, 1091–8 (2015).

- 1113 16. Gusev, A. *et al.* Integrative approaches for large-scale transcriptome-wide association
1114 studies. *Nat. Genet.* **48**, 245–52 (2016).
- 1115 17. Geschwind, D. H. & Flint, J. Genetics and genomics of psychiatric disease. *Science* (80-
1116). **349**, (2015).
- 1117 18. Leek, J. T., Johnson, W. E., Parker, H. S., Jaffe, A. E. & Storey, J. D. The sva package for
1118 removing batch effects and other unwanted variation in high-throughput experiments.
1119 *Bioinformatics* **28**, 882–3 (2012).
- 1120 19. A. Bennett, D., A. Schneider, J., Arvanitakis, Z. & S. Wilson, R. Overview and Findings
1121 from the Religious Orders Study. *Curr. Alzheimer Res.* **9**, 628–645 (2012).
- 1122 20. Mele, M. *et al.* The human transcriptome across tissues and individuals. *Science* (80-
1123). **348**, 660–665 (2015).
- 1124 21. (GTEx Consortium). GTEx Portal. (2015). at
1125 <<http://gtexportal.org/home/documentationPage>>
- 1126 22. Ripke, S. *et al.* Biological insights from 108 schizophrenia-associated genetic loci. *Nature*
1127 **511**, 421–427 (2014).
- 1128 23. Benjamin, Y. & Hochberg, Y. Controlling the False Discovery Rate: a Practical and
1129 Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society* 289–300
1130 (1995).
- 1131 24. Darnell, J. C. *et al.* FMRP stalls ribosomal translocation on mRNAs linked to synaptic
1132 function and autism. *Cell* **146**, 247–61 (2011).
- 1133 25. Fromer, M. *et al.* *Gene Expression Elucidates Functional Impact of Polygenic Risk for*
1134 *Schizophrenia*. *bioRxiv* (Cold Spring Harbor Labs Journals, 2016). doi:10.1101/052209
- 1135 26. Pardiñas, A. F. *et al.* Common schizophrenia alleles are enriched in mutation-intolerant
1136 genes and maintained by background selection. *bioRxiv* 68593 (2016).
1137 doi:10.1101/068593
- 1138 27. Sanders, S. J. First glimpses of the neurobiology of autism spectrum disorder. *Curr. Opin.*
1139 *Genet. Dev.* **33**, 80–92 (2015).
- 1140 28. Exome Aggregation Consortium *et al.* *Analysis of protein-coding genetic variation in*
1141 *60,706 humans*. *bioRxiv* (Cold Spring Harbor Labs Journals, 2015). doi:10.1101/030338
- 1142 29. Malhotra, D. *et al.* High frequencies of de novo CNVs in bipolar disorder and
1143 schizophrenia. *Neuron* **72**, 951–963 (2011).

- 1144 30. Bautista, O., Vázquez-Caubet, J. C., Zhivago, E. A. & Dolores Sáiz, M. From metabolism
1145 to psychiatric symptoms: psychosis as a manifestation of acute intermittent porphyria. *J.*
1146 *Neuropsychiatry Clin. Neurosci.* **26**, E30 (2014).
- 1147 31. Zimmermann, M., Bonaccorso, C., Valerius, C. & Hamann, G. F. [Acute intermittent
1148 porphyria. A clinical chameleon: case study of a 40-year-old female patient]. *Nervenarzt*
1149 **77**, 1501–5 (2006).
- 1150 32. Ventura, P. *et al.* A challenging diagnosis for potential fatal diseases: recommendations
1151 for diagnosing acute porphyrias. *Eur. J. Intern. Med.* **25**, 497–505 (2014).
- 1152 33. Pischik, E. & Kauppinen, R. An update of clinical management of acute intermittent
1153 porphyria. *Appl. Clin. Genet.* **8**, 201–14 (2015).
- 1154 34. Kumar, B. Acute intermittent porphyria presenting solely with psychosis: a case report
1155 and discussion. *Psychosomatics* **53**, 494–8 (2012).
- 1156 35. Bonnot, O. *et al.* Diagnostic and treatment implications of psychosis secondary to
1157 treatable metabolic disorders in adults: a systematic review. *Orphanet J. Rare Dis.* **9**, 65
1158 (2014).
- 1159 36. Kaback, M. M. & Desnick, R. J. *Hexosaminidase A Deficiency*. *GeneReviews*(®)
1160 (University of Washington, Seattle, 1993).
- 1161 37. Saleh, O. Late Onset Tay-Sachs Disease Pre senting as a Brief Psychotic Disorder with
1162 Catatonia: A Case Report and Review of Literature.
- 1163 38. Skaper, S. D. in *Brain Protection in Schizophrenia, Mood and Cognitive Disorders* (ed.
1164 Ritsner, M. S.) 135–165 (Springer Science & Business Media, 2010, 2010).
- 1165 39. Castellano, E. *et al.* RAS signalling through PI3-Kinase controls cell migration via
1166 modulation of Reelin expression. *Nat. Commun.* **7**, 11245 (2016).
- 1167 40. Gururajan, A. & Buuse, M. van den. Is the mTOR-signalling cascade disrupted in
1168 Schizophrenia? *J. Neurochem.* **129**, 377–387 (2014).
- 1169 41. Ritsner, M. S. *Brain Protection in Schizophrenia, Mood and Cognitive Disorders*.
1170 (Springer Science & Business Media, 2010, 2010). doi:10.1007/978-90-481-8553-5
- 1171 42. Enriquez-Barreto, L. & Morales, M. The PI3K signaling pathway as a pharmacological
1172 target in Autism related disorders and Schizophrenia. *Mol. Cell. Ther.* **4**, 2 (2016).
- 1173 43. Glessner, J. T. *et al.* Strong synaptic transmission impact by copy number variations in
1174 schizophrenia. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 10584–9 (2010).

- 1175 44. Network and Pathway Analysis Subgroup of Psychiatric Genomics Consortium.
1176 Psychiatric genome-wide association study analyses implicate neuronal, immune and
1177 histone pathways. *Nat. Neurosci.* **18**, 199–209 (2015).
- 1178 45. Bauman, A. L. *et al.* Cocaine and antidepressant-sensitive biogenic amine transporters
1179 exist in regulated complexes with protein phosphatase 2A. *J. Neurosci.* **20**, 7571–8
1180 (2000).
- 1181 46. He, X. *et al.* Sherlock: detecting gene-disease associations by matching patterns of
1182 expression QTL and GWAS. *Am J Hum Genet* **92**, 667–680 (2013).
- 1183 47. Ayadi, A. *et al.* Mouse large-scale phenotyping initiatives: overview of the European
1184 Mouse Disease Clinic (EUMODIC) and of the Wellcome Trust Sanger Institute Mouse
1185 Genetics Project. *Mamm. Genome* **23**, 600–610 (2012).
- 1186 48. MGI-About the Mouse Genome Informatics database resource. at
1187 <<http://www.informatics.jax.org/mgihome/projects/aboutmgi.shtml>>
- 1188 49. Keane, T. M. *et al.* Mouse genomic variation and its effect on phenotypes and gene
1189 regulation. *Nature* **477**, 289–294 (2011).
- 1190 50. About KOMP | IMPC.
- 1191 51. Howe, D. G. *et al.* ZFIN, the Zebrafish Model Organism Database: increased support for
1192 mutants and transgenics. *Nucleic Acids Res.* **41**, D854–D860 (2013).
- 1193 52. TRANSCRIPTOME PROFILING BY RNA SEQUENCING AND EXON
1194 MICROARRAY TISSUE ACQUISITION AND QUALIFICATION.
- 1195 53. Watts, D. J. & Strogatz, S. H. Collective dynamics of ‘small-world’ networks. *Nature* **393**,
1196 440–442 (1998).
- 1197 54. Schork, A. J. *et al.* All SNPs Are Not Created Equal: Genome-Wide Association Studies
1198 Reveal a Consistent Pattern of Enrichment among Functionally Annotated SNPs. *PLoS*
1199 *Genet.* **9**, e1003449 (2013).
- 1200 55. Mancuso, N. *et al.* Integrating Gene Expression with Summary Association Statistics to
1201 Identify Genes Associated with 30 Complex Traits. *Am. J. Hum. Genet.* **100**, 473–487
1202 (2017).
- 1203 56. Gottlieb, A., Daneshjou, R., DeGorter, M., Montgomery, S. & Altman, R. Population-
1204 specific imputation of gene expression improves prediction of pharmacogenomic traits for
1205 African Americans. *bioRxiv* 115451 (2017). doi:10.1101/115451

- 1206 57. Need, A. & Goldstein, D. B. Next generation disparities in human genomics: concerns and
1207 remedies. *Trends Genet* **25**, 489–94 (2009).
- 1208 58. Popejoy, A. & Fullerton, S. Genomics is failing on diversity. *Nature* **538**, 161–164 (2016).
- 1209 59. Browning, R. in *The Poems of Robert Browning* (eds. Porter, C. & Clarke, H. A.) 257–271
1210 (Thomas Y. Cromwell and Company, 1896).
- 1211 60. Loftus, L. S. & Arnold, W. N. Vincent van Gogh's illness: acute intermittent porphyria?
1212 *BMJ* **303**, 1589–91
- 1213 61. Strik, W. K. [The psychiatric illness of Vincent van Gogh]. *Nervenarzt* **68**, 401–9 (1997).
- 1214 62. Arnold, W. N. The illness of Vincent van Gogh. *J. Hist. Neurosci.* **13**, 22–43 (2004).
- 1215 63. Hughes, J. R. A reappraisal of the possible seizures of Vincent van Gogh. *Epilepsy Behav.*
1216 **6**, 504–10 (2005).
- 1217 64. Bhattacharyya, K. B. & Rai, S. The neuropsychiatric ailment of Vincent Van Gogh. *Ann.*
1218 *Indian Acad. Neurol.* **18**, 6–9 (2014).
- 1219 65. Correa, R. Vincent van Gogh: A pathographic analysis. *Med. Hypotheses* **82**, 141–144
1220 (2014).
- 1221 66. Peters, T. J. & Beveridge, A. The madness of King George III: a psychiatric re-
1222 assessment. *Hist. Psychiatry* **21**, 20–37 (2010).
- 1223 67. Szatkiewicz, J. P. *et al.* Copy number variation in schizophrenia in Sweden. *Mol.*
1224 *Psychiatry* **19**, 762–773 (2014).
- 1225 68. Fromer, M. *et al.* De novo mutations in schizophrenia implicate synaptic networks. *Nature*
1226 **506**, 179–184 (2014).
- 1227 69. Consortium, C.-D. G. of the P. G. Identification of risk loci with shared effects on five
1228 major psychiatric disorders: a genome-wide analysis. *Lancet* **381**, 1371–1379 (2013).
- 1229 70. Keefe, R. S. E. & Fenton, W. S. How Should DSM-V Criteria for Schizophrenia Include
1230 Cognitive Impairment? *Schizophr. Bull.* **33**, 912–920 (2007).
- 1231 71. Reichenberg, A. *et al.* Static and Dynamic Cognitive Deficits in Childhood Preceding
1232 Adult Schizophrenia: A 30-Year Study. *Am. J. Psychiatry* **167**, 160–169 (2010).
- 1233 72. Gold, J. M. Cognitive deficits as treatment targets in schizophrenia. *Schizophr. Res.* **72**,
1234 21–28 (2004).
- 1235 73. Cannon, M. *et al.* Evidence for Early-Childhood, Pan-Developmental Impairment Specific
1236 to Schizophreniform Disorder. *Arch. Gen. Psychiatry* **59**, 449 (2002).

74. Parikshak, N. N., Gandal, M. J., Geschwind, D. H. & Angeles, L. Systems biology and gene networks in neurodevelopmental and neurodegenerative disorders. *Nat. Rev. Genet.* **16**, 441–458 (2015).
75. Glass, D. *et al.* Gene expression changes with age in skin, adipose tissue, blood and brain. *Genome Biol.* **14**, R75 (2013).
76. Colantuoni, C. *et al.* Temporal dynamics and genetic control of transcription in the human prefrontal cortex. *Nature* **478**, 519–523 (2012).
77. Gusev, A. *et al.* Transcriptome-wide association study of schizophrenia and chromatin activity yields mechanistic disease insights. *bioRxiv* 67355 (2016). doi:10.1101/067355
78. ROSMAP Study - syn3219045. at <<https://www.synapse.org/#!Synapse:syn3219045>>
79. NIMH » Human Brain Collection Core (HBCC). at <<https://www.nimh.nih.gov/labs-at-nimh/research-areas/research-support-services/hbcc/human-brain-collection-core-hbcc.shtml>>
80. Barbeira, A. *et al.* MetaXcan: Summary Statistics Based Gene-Level Association Method Infers Accurate PrediXcan Results. *bioRxiv* (2016).
81. Ardlie, K. G. *et al.* The Genotype-Tissue Expression (GTEx) pilot analysis: Multitissue gene regulation in humans. *Science* (80-.). **348**, 648–660 (2015).
82. Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* **26**, 2190–2191 (2010).
83. COCHRAN, W. G. THE COMPARISON OF PERCENTAGES IN MATCHED SAMPLES. *Biometrika* **37**, 256–266 (1950).
84. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide complex trait analysis. *Am. J. Hum. Genet.* **88**, 76–82 (2011).
85. de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: generalized gene-set analysis of GWAS data. *PLoS Comput. Biol.* **11**, e1004219 (2015).
86. Kirov, G. *et al.* De novo CNV analysis implicates specific abnormalities of postsynaptic signalling complexes in the pathogenesis of schizophrenia. *Mol. Psychiatry* **17**, 142–53 (2012).
87. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat. Genet.* **25**, 25–29 (2000).
88. The Gene Ontology Consortium. Gene Ontology Consortium: going forward. *Nucleic*

1268 *Acids Res.* **43**, D1049-1056 (2014).

1269 89. Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic*

1270 *Acids Res.* **28**, 27–30 (2000).

1271 90. Croft, D. *et al.* The Reactome pathway knowledgebase. *Nucleic Acids Res.* **42**, D472-7

1272 (2014).

1273 91. Thomas, P. D. *et al.* PANTHER: a library of protein families and subfamilies indexed by

1274 function. *Genome Res.* **13**, 2129–2141 (2003).

1275 92. Mi, H., Muruganujan, A. & Thomas, P. D. PANTHER in 2013: modeling the evolution of

1276 gene function, and other gene attributes, in the context of phylogenetic trees. *Nucleic*

1277 *Acids Res.* **41**, D377-86 (2013).

1278 93. BioCarta. MSigDB Collections. (2017).

1279 94. Miller, J. A. *et al.* Transcriptional landscape of the prenatal human brain. *Nature* **508**,

1280 199–206 (2014).

1281 95. Lin, G. N. *et al.* Spatiotemporal 16p11.2 Protein Network Implicates Cortical Late Mid-

1282 Fetal Brain Development and KCTD13- Cul3-RhoA Pathway in Psychiatric Diseases.

1283 *Neuron* **85**, 742–754 (2015).

1284 96. Bahl, E., Koomar, T. & Michaelson, J. J. cerebroViz: An R package for anatomical visu-

1285 alization of spatiotemporal brain data. *Bioinformatics* **33**, btw726 (2016).

1286 97. van der Weyden, L., White, J. K., Adams, D. J. & Logan, D. W. The mouse genetics

1287 toolkit: revealing function and mechanism. *Genome Biol.* **12**, 224 (2011).

1288 98. Home | 3i Consortium - Wellcome Trust Sanger Institute. at

1289 <<http://www.immunophenotype.org/>>

1290 99. Mors, O., Perto, G. P. & Mortensen, P. B. The Danish Psychiatric Central Research

1291 Register. *Scand. J. Public Health* **39**, 54–57 (2011).

1292 100. Børghlum, A. D. *et al.* Genome-wide study of association and interaction with maternal

1293 cytomegalovirus infection suggests new schizophrenia loci. *Mol. Psychiatry* **19**, 325–33

1294 (2014).

1295 101. Hollegaard, M. V *et al.* Robustness of genome-wide scanning using archived dried blood

1296 spot samples as a DNA source. *BMC Genet.* **12**, 58 (2011).

1297 102. Illumina. illumina GenCall Data Analysis Software. Illumina Tech Note. (2005). at

1298 <https://www.illumina.com/Documents/products/technotes/technote_gencall_data_analysi

1299 s_software.pdf>
1300 103. Korn, J. M. *et al.* Integrated genotype calling and association analysis of SNPs, common
1301 copy number polymorphisms and rare CNVs. *Nat. Genet.* **40**, 1253–60 (2008).
1302 104. Goldstein, J. I. *et al.* zCall: a rare variant caller for array-based genotyping: genetics and
1303 population analysis. *Bioinformatics* **28**, 2543–2545 (2012).
1304




Figure 1: Replication of DLPFC prediction models in independent data.

Measured gene expression (ROSMAP RNA-seq) was compared to predicted genetically-regulated gene expression for CMC DLPFC and 12 GTEx predictor databases. Replication R^2 values are significantly higher for the DLPFC than for the 12 GTEx brain expression models.

- Distribution of R_R^2 values of CMC DLPFC predictors in ROSMAP data. Mean $R_R^2 = 0.056$. 47.7% of genes have $R_R^2 \geq 0.01$.
- Distribution of R_R^2 values of 12 GTEx predictors in ROSMAP data.
- Table of sample sizes and p-val thresholds for CMC DLPFC and GTEx data. Number of samples, number of genes in the prediXcan model and number of eGenes are all significantly correlated with predictor performance in ROSMAP data.




Figure 2: Analysis outline.

- A) Discovery Samples. 41 PGC-SCZ cohorts had available raw genotypes (i). Predicted DLPFC gene expression was calculated in each cohort using prediXcan (ii) and tested for association with case-control status (iii). 11 PGC cohorts (3 trio, 8 case-control) and the CLOZUK2 cohort had only summary statistics available (iv). MetaXcan was used to calculate DLPFC associations for each cohort (v). Results were meta-analysed across all 53 cohorts (vi). This procedure was repeated for 12 GTEx prediction models.
- B) Replication Samples. iPSYCH-GEMS samples were collected in 25 waves (i). Predicted DLPFC gene expression was calculated in each wave separately using prediXcan (ii) and merged for association testing (iii). A mega-analysis was run across all 25 waves, using wave membership as a covariate in the regression (iv)




Figure 3: SCZ associations results

- A) 413 genes are associated with SCZ across 12 brain tissues
- B) 67 genes remain significant outside the MHC after stepwise conditional analysis
- C) Number of genome-wide significant loci, outside the MHC region, identified in each brain region. Abbreviations are as follows; CB- Cerebellum; CX- Cortex; FL- Frontal Cortex; DLPFC- Dorso-lateral pre-frontal cortex; CB HEMI- Cerebellar Hemisphere; HIP- Hippocampus; PIT- Pituitary Gland; HTH- Hypothalamus; NAB- Nucleus Accumbens (Basal Ganglia); PUT- Putamen (Basal Ganglia); CAU- Caudate (Basal Ganglia); CNG- Anterior Cingulate Cortex




Figure 4: SCZ-associated genes are co-expressed throughout development and across brain regions

- A) Brain tissues selected for each of four BRAINSPAN regions. Region 1: IPC, V1C, ITC, OFC, STC, A1C; Region 2: S1C, M1C, DFC, VFC, MFC; Region 3: HIP, AMY, STR; Region 4: CB
- B) Average clustering coefficients were calculated for all pairs of SCZ-associated genes, and compared to permuted gene networks to obtain empirical significance levels.




Figure 5: Gene expression patterns for SCZ-associated genes cluster into four groups, relating to distinct spatiotemporal expression.

Brain regions are shown in figure 5a.

- A. 29 genes are expressed in the early-mid pre-natal period (4-24 post-conception weeks)
- B. 15 genes are expressed throughout development; sub-clusters correspond to either specific expression in region 4, or expression across the brain
- C. Ten genes are expressed in the late-prenatal (25-38pcw) and post-natal period
- D. 12 genes are expressed in the late pre-natal period (25-39pcw)

Table 1: SCZ-associated genes

<i>Gene name</i>	Tissue	BETA	P	GVAR	Adjusted BETA	Adjusted OR
<i>GNL3</i>	GTEX Cerebellum	0.037	1.39E-11	0.115	0.012	1.012
<i>THOC7</i>	GTEX Cerebellum	-0.113	5.77E-10	0.010	-0.011	0.989
<i>NAGA</i>	GTEX Cerebellum	0.122	1.12E-09	0.009	0.011	1.011
<i>TAC3</i>	GTEX Cerebellum	-0.868	8.03E-08	0.000	-0.015	0.985
<i>CHRNA2</i>	GTEX Cerebellum	-0.016	1.63E-07	0.395	-0.010	0.990
<i>ACTR5</i>	GTEX Cerebellum	0.208	3.88E-07	0.019	0.029	1.029
<i>INO80E</i>	GTEX Frontal Cortex	0.130	7.25E-12	0.009	0.012	1.013
<i>PLPPR5</i>	GTEX Frontal Cortex	-0.672	2.58E-09	0.006	-0.053	0.948
<i>FAM205A</i>	GTEX Frontal Cortex	0.043	1.21E-08	0.061	0.011	1.011
<i>AC110781.3</i>	GTEX Thyroid	0.342	1.31E-13	0.002	0.014	1.014
<i>IMMP2L</i>	GTEX Thyroid	-0.073	7.09E-12	0.046	-0.016	0.984
<i>IGSF9B</i>	GTEX Thyroid	-0.024	3.05E-07	0.156	-0.010	0.991
<i>NMRAL1</i>	GTEX Thyroid	0.038	4.03E-07	0.060	0.009	1.009
<i>HIF1A</i>	CMC DLPEC	11.130	7.52E-14	0.000	0.148	1.159
<i>TMEM29</i>	CMC DLPEC	11.207	9.27E-14	0.000	0.168	1.183
<i>ST7-014</i>	CMC DLPEC	10.170	5.79E-13	0.001	0.318	1.374
<i>H2AFY2</i>	CMC DLPEC	10.962	3.60E-12	0.000	0.191	1.211
<i>STARD3</i>	CMC DLPEC	10.740	5.90E-12	0.001	0.304	1.355
<i>CTC-471F3.5</i>	CMC DLPEC	8.535	1.11E-11	0.000	0.104	1.110
<i>SF3A1</i>	CMC DLPEC	8.651	1.32E-11	0.000	0.083	1.086
<i>ZNF512</i>	CMC DLPEC	10.312	1.32E-11	0.001	0.261	1.298
<i>FURIN</i>	CMC DLPEC	-0.084	2.22E-11	0.022	-0.012	0.988
<i>INHBA-AS1</i>	CMC DLPEC	8.399	2.24E-11	0.000	0.127	1.135
<i>SF3B1</i>	CMC DLPEC	0.099	6.14E-11	0.014	0.012	1.012
<i>EFTUD1P1</i>	CMC DLPEC	-0.092	1.81E-10	0.017	-0.012	0.988
<i>MLH1</i>	CMC DLPEC	2.840	2.10E-10	0.001	0.069	1.071
<i>GATAD2A</i>	CMC DLPEC	-0.044	2.18E-10	0.071	-0.012	0.988
<i>METTL1</i>	CMC DLPEC	9.357	2.23E-10	0.000	0.166	1.181

<i>DMC1</i>	CMC	DLPFC	7.229	4.48E-10	0.000	0.130	1.139
<i>RAD51D</i>	CMC	DLPFC	7.612	2.11E-09	0.000	0.111	1.117
<i>RERE</i>	CMC	DLPFC	2.847	6.32E-09	0.000	0.036	1.037
<i>PCCB</i>	CMC	DLPFC	-0.044	2.05E-08	0.054	-0.010	0.990
<i>CLCN3</i>	CMC	DLPFC	0.141	2.96E-08	0.005	0.010	1.010
<i>ATG101</i>	CMC	DLPFC	8.086	4.90E-08	0.007	0.695	2.005
<i>JRK</i>	CMC	DLPFC	0.032	1.25E-07	0.091	0.010	1.010
<i>PTPRU</i>	CMC	DLPFC	-0.077	1.60E-07	0.016	-0.010	0.990
<i>MARCKS</i>	CMC	DLPFC	0.398	2.05E-07	0.001	0.015	1.015
<i>TCF4</i>	GTEX	Anterior Cingulate Cortex	-0.059	5.22E-13	0.051	-0.013	0.987
<i>DGKD</i>	GTEX	Anterior Cingulate Cortex	-0.937	2.63E-11	0.001	-0.022	0.979
<i>CIQTNF4</i>	GTEX	Anterior Cingulate Cortex	-0.173	1.37E-09	0.010	-0.017	0.983
<i>PTPNA</i>	GTEX	Anterior Cingulate Cortex	-0.243	1.77E-07	0.002	-0.010	0.990
<i>FXR1</i>	GTEX	Caudate Basal Ganglia	0.439	5.40E-12	0.001	0.017	1.017
<i>ZDHHC1</i>	GTEX	Caudate Basal Ganglia	0.354	5.36E-08	0.001	0.011	1.012
<i>PDE4D</i>	GTEX	Cerebellar Hemisphere	0.365	6.81E-11	0.001	0.013	1.013
<i>DRD2</i>	GTEX	Cerebellar Hemisphere	-0.182	2.47E-10	0.004	-0.012	0.988
<i>PTPNM2</i>	GTEX	Cerebellar Hemisphere	-0.065	2.21E-09	0.028	-0.011	0.989
<i>RINT1</i>	GTEX	Cerebellar Hemisphere	0.086	6.32E-09	0.016	0.011	1.011
<i>SRMS</i>	GTEX	Cerebellar Hemisphere	-0.440	3.08E-08	0.001	-0.011	0.989
<i>SETD6</i>	GTEX	Cerebellar Hemisphere	-0.043	1.05E-07	0.054	-0.010	0.990
<i>APOPT1</i>	GTEX	Cortex	-0.074	1.24E-10	0.026	-0.012	0.988
<i>VSIG2</i>	GTEX	Cortex	-0.092	6.01E-09	0.013	-0.011	0.989
<i>SDCC4G8</i>	GTEX	Cortex	-0.069	3.88E-07	0.002	-0.003	0.997
<i>PIK3C2A</i>	GTEX	Cortex	-0.040	4.04E-07	0.365	-0.024	0.976
<i>AS3MT</i>	GTEX	Frontal Cortex	0.594	5.65E-17	0.001	0.017	1.017
<i>FOXN2</i>	GTEX	Hippocampus	-0.250	2.65E-07	0.021	-0.036	0.964
<i>RASIP1</i>	GTEX	Nucleus Accumbens Basal Ganglia	0.055	3.80E-08	0.034	0.010	1.010
<i>TCF23</i>	GTEX	Nucleus Accumbens Basal Ganglia	-0.076	4.83E-08	0.019	-0.010	0.990
<i>TTCL4</i>	GTEX	Nucleus Accumbens Basal Ganglia	-0.089	4.84E-08	0.013	-0.010	0.990
<i>TYW5</i>	GTEX	Putamen Basal Ganglia	-0.080	2.63E-13	0.035	-0.015	0.985

<i>SNX19</i>	GTEx	Putamen Basal Ganglia	0.031	1.31E-12	0.179	0.013	1.013
<i>CIART</i>	GTEx	Putamen Basal Ganglia	0.090	6.78E-10	0.017	0.012	1.012
<i>SH2D7</i>	GTEx	Putamen Basal Ganglia	0.096	7.89E-09	0.013	0.011	1.011
<i>DGULO</i>	GTEx	Putamen Basal Ganglia	0.255	8.26E-08	0.002	0.011	1.011
<i>CI2orf76</i>	GTEx	Putamen Basal Ganglia	0.031	2.27E-07	0.095	0.010	1.010
<i>LIRC374</i>	GTEx	Putamen Basal Ganglia	-0.035	2.69E-07	0.076	-0.010	0.991
<i>AC005841.1</i>	GTEx	Pituitary	0.162	3.28E-09	0.005	0.011	1.011
<i>RPS17</i>	GTEx	Pituitary	0.035	4.03E-08	0.082	0.010	1.010

MHC Region:

<i>BTNL1A1</i>	GTEx	Caudate Basal Ganglia	-0.2606	1.6666E-22			
<i>VARS2</i>	GTEx	Anterior Cingulate Cortex	0.0747019	7.4821E-15			
<i>HST1H3H</i>	GTEx	Putamen Basal Ganglia	-1.105982	3.2236E-10			
<i>NUDT3</i>	GTEx	Nucleus Accumbens Basal Ganglia	0.10378753	6.546E-09			

Table 2: Significantly enriched pathways and gene sets

Analysis	Gene Set	Comp P	FDR P
Hypothesis driven	FMRP-targets	1.96x10 ⁻⁰⁸	3.097x10 ⁻⁰⁶
	BP denovo CNV	7.92x10 ⁻⁰⁸	6.257x10 ⁻⁰⁶
	HIGH LOF intolerant	5.86x10 ⁻⁰⁵	0.00309
Agnostic	Increased spleen iron level	2.72x10 ⁻⁰⁸	0.000245
	Decreased IgM level	6.80x10 ⁻⁰⁷	0.00307
	Condensed chromosome	1.99x10 ⁻⁰⁶	0.00598
	Chromosome	2.80x10 ⁻⁰⁶	0.00632
	Abnormal spleen iron level	6.79x10 ⁻⁰⁶	0.00765
	Mitotic Anaphase	6.39 x10 ⁻⁰⁶	0.00765
	Mitotic Metaphase and Anaphase	5.13 x10 ⁻⁰⁶	0.00765
	Resolution of Sister Chromatid Cohesion	5.82 x10 ⁻⁰⁶	0.00765
	Increased liver iron level	1.03 x10 ⁻⁰⁵	0.0103
	Separation of Sister Chromatids	1.28 x10 ⁻⁰⁵	0.0115
	Regulation of Rab GTPase activity	1.78 x10 ⁻⁰⁵	0.0123
	Regulation of Rab protein signal transduction	1.78 x10 ⁻⁰⁵	0.0123
	Protein phosphorylated amino acid binding	1.75x10 ⁻⁰⁵	0.0123
	Chromosome	2.57x10 ⁻⁰⁵	0.0165
	Hexosaminidase activity	3.47x10 ⁻⁰⁵	0.0174
	Abnormal learningmemoryconditioning	3.11x10 ⁻⁰⁵	0.0174
	Abnormal liver iron level	3.47x10 ⁻⁰⁵	0.0174
	Mitotic Prometaphase	2.99x10 ⁻⁰⁵	0.0174
	M Phase	3.70x10 ⁻⁰⁵	0.0176
	Positive regulation of Rab GTPase activity	5.93x10 ⁻⁰⁵	0.0232
	Rab GTPase activator activity	5.93x10 ⁻⁰⁵	0.0232
	Protein phosphatase type 2A regulator activity	5.24x10 ⁻⁰⁵	0.0232
	Replicative senescence	5.44x10 ⁻⁰⁵	0.0232
	Condensed nuclear chromosome	7.11x10 ⁻⁰⁵	0.0267
	Ubiquitin-specific protease activity	0.000104	0.0335
	Ras GTPase activator activity	9.61x10 ⁻⁰⁵	0.0335
	Metabolism of porphyrins	0.000103	0.0335
	Kinetochore	0.000103	0.0335
	Decreased physiological sensitivity to xenobiotic	0.000127	0.0381
	Antigen Activates B Cell Receptor Leading to Generation of Second Messengers	0.000124	0.0381
	Phosphoprotein binding	0.000146	0.0424
	Abnormal dorsal-ventral axis patterning	0.000152	0.0429