

***Cyfp1* haploinsufficiency increases compulsive-like behavior and modulates palatable food intake: Implications for Prader-Willi Syndrome**

Richard K. Babbs, Ph.D.^{*}, Qiu T. Ruan, B.S.^{*,†,‡}, Julia C. Kelliher, B.S.^{*}, Jacob A. Beierle, B.S.^{*,‡}, Melanie M. Chen, Ashley X. Feng^{*}, Stacey L. Kirkpatrick, B.A.^{*}, Fabiola A. Benitez^{*}, Fred A. Rodriguez^{*}, Johanne J. Pierre^{*}, Jeya Anandakumar^{*}, Vivek Kumar, Ph.D.[§], Megan K. Mulligan, Ph.D.^{**}, Camron D. Bryant, Ph.D.^{*}

Affiliations

^{*} Laboratory of Addiction Genetics, Department of Pharmacology and Experimental Therapeutics and Psychiatry, Boston University School of Medicine, 72 E. Concord St., L-606C, Boston, MA 02118 USA

[†] NIGMS Training Program in Biomolecular Pharmacology, Boston University School of Medicine, 72 E. Concord St., Boston, MA 02118 USA

[‡] Boston University's Transformative Training Program in Addiction Science (TTPAS), Biomedical Genetics, Boston University School of Medicine, 72 E. Concord St., E-200, Boston, MA 02118 USA

[§] The Jackson Laboratory, 600 Main St., Bar Harbor, ME 04609USA

^{**} Department of Genetics, Genomics, and Informatics, University of Tennessee Health Science Center, 71 S. Manassas St, Memphis, TN 38163 USA

Running title: *Cyfp1* deletion and palatable food intake

Five key words: FMRP, overeating, C57BL/6 substrains, obsessive-compulsive, Fragile X

***Corresponding Author**

Camron D. Bryant, Ph.D.

Director, Laboratory of Addiction Genetics

Department of Pharmacology and Experimental Therapeutics

72 E. Concord St.

L-606C

Boston, MA 02118

E: camron@bu.edu

P: (617) 638-4489

F: (617) 638-4329

ABSTRACT

Binge eating (BE) is a heritable trait associated with eating disorders and involves rapid consumption of large quantities of food. We identified cytoplasmic FMRP-interacting protein 2 (*Cytip2*) as a major genetic factor underlying BE and concomitant compulsive-like behaviors in mice. *CYFIP2* is a gene homolog of *CYFIP1* - one of four paternally-deleted genes in patients with the more severe Type I Prader-Willi Syndrome (PWS). PWS is a neurodevelopmental disorder where 70% of cases involve paternal deletion of 15q11-q13. PWS symptoms include hyperphagia, obesity (if untreated), cognitive deficits, and obsessive-compulsive behaviors. We tested whether *Cytip1* haploinsufficiency (+/-) would enhance premorbid compulsive-like behavior and palatable food (PF) intake in a parent-of-origin-selective manner. We tested *Cytip1*^{+/-} mice on a C57BL/6N (N) background that were homozygous for the BE-associated missense mutation in *Cytip2* (S968F) as well as mice that we backcrossed to homozygosity for the C57BL/6J (J) allele at *Cytip2* (*Cytip2*^{J/J}). *Cytip1*^{+/-} mice showed increased compulsive-like behavior on both backgrounds, increased PF consumption on the *Cytip2*^{N/N} background in a paternally-enhanced manner, and decreased PF consumption in male *Cytip1*^{+/-} mice on the *Cytip2*^{J/J} background in a maternally selective manner. In the hypothalamus, there was a maternally-enhanced reduction of *Cytip1* transcription, but a paternally-enhanced reduction in CYFIP1 protein. In the nucleus accumbens, there was a maternally-enhanced reduction in CYFIP1 protein. Together, increased compulsive-like behavior, parent-of-origin-, and genetic background-dependent effects of *Cytip1* haploinsufficiency on PF consumption implicate *CYFIP1* in behaviors in neurodevelopmental disorders involving reduced expression of *CYFIP1*, including PWS, Fragile X Syndrome, and 15q11.2 Microdeletion Syndrome.

INTRODUCTION

Binge eating (**BE**) refers to the rapid consumption of large quantities of food and is accompanied by feelings of loss of control. Binge eating disorder (**BED**) is a psychiatric disorder with a lifetime prevalence of 3.5% in women and 2% in men¹. Both BED² and BE are heritable³. However, genome-wide association studies have yet to identify genetic risk factors associated with BE⁴. The first genome-wide significant loci were recently identified for anorexia nervosa (comprising restricted eating)⁵ and bipolar disorder with BE behavior PRR5-ARHGAP8⁶. Additional genome-wide significant loci will likely be uncovered for BE-associated disorders with increasing sample sizes and power⁷.

We used quantitative trait locus (**QTL**) mapping and gene knockout in C57BL/6 mouse substrains to identify cytoplasmic FMR1-interacting protein 2 (*Cyfp2*) as a major genetic factor underlying BE and compulsive-like behaviors⁸. The QTL capturing increased palatable food (PF) intake mapped to a missense mutation in *Cyfp2* in the C57BL/6N strain (S968F; “*Cyfp2*^{M1N}”) that is hypothesized to act as a gain-of-function mutation⁹. Accordingly, mice with one copy of a null allele and one copy of the missense allele of *Cyfp2* showed a reduction in BE toward the phenotypic direction of the wild-type C57BL/6J level⁸. This same missense SNP in *Cyfp2* was first associated with reduced behavioral sensitivity to cocaine⁹, which could indicate a common neurobiological mechanism involving synaptic plasticity within the mesocorticolimbic dopamine reward pathway^{10,11} that affects the hedonic component of PF consumption^{12,13}.

Cyfp2 and the gene homolog *Cyfp1* code for proteins that interact with the RNA binding protein Fragile X Mental Retardation Protein (**FMRP**) and are part of the canonical WAVE regulatory complex and transduce activity-dependent Rac signaling in regulating actin dynamics during neuronal development and synaptic plasticity¹⁴. CYFIP1 expression is necessary for the maintenance and stabilization of neuronal dendritic arborization and morphological complexity¹⁵. In humans, *CYFIP1* resides within a non-imprinted region on chromosome 15 (15q11.2) that contains four genes *TUBGCP5*, *NIPA1*, *NIPA2*, and *CYFIP1*¹⁶. The syntenic region in mice is located on chromosome 7C (55.4 Mb - 56 Mb). Haploinsufficiency of 15q11.2 underlies Microdeletion Syndrome (**MDS**) which can comprise developmental delay (speech, motor), reduced cognitive function, dysmorphic features,

intellectual disability, autism, ADHD, obsessive-compulsive disorder, and schizophrenia¹⁷. At least one case study of 15q11.2 MDS reported hypotonia, increased food craving and obesity, and obsessive-compulsive disorder¹⁸. *CYFIP1* haploinsufficiency is implicated in multiple symptoms of 15q11.2 MDS. Preclinical models of *Cyfp1* haploinsufficiency demonstrate perturbations in synaptic activity during neural development, activity-dependent plasticity, dendritic morphology, and fear learning¹⁹⁻²².

The 15q11.2 region is also paternally-deleted in a subset of individuals with a more severe form (Type I) of Prader-Willi Syndrome (**PWS**), a neurodevelopmental disorder defined genetically by paternal deletion of 15q11-q13 in a majority of cases²³. Extreme hyperphagia due to lack of satiety is the most defining and debilitating feature of PWS that is difficult to treat and emerges during childhood, leading to obesity if left untreated. Food-related obsessive-compulsive (**OC**) behaviors are common in PWS; however, OC symptoms unrelated to food are also frequent²⁴, and include repetitive, ritualistic behaviors, perseverative speech, counting, adaptive impairment, need to tell, ask, or know, ordering and arranging, repeating rituals, and self-mutilation²⁵⁻²⁷. Genetic deletion in PWS involves either the shorter paternal deletion (Type II) of 15q11-q13 or a larger, paternal Type I deletion that also includes the 0.5 Mb 15q11.2 MDS region comprising four genes: *TUBGCP5*, *NIPA1*, *NIPA2*, and *CYFIP1*^{16,28}. Type I PWS is associated with reduced transcription of these genes and a more severe neurodevelopmental and neuropsychiatric profile, including reduced cognition, increased risk of autism and schizophrenia, and increased severity and lack of control over OC behaviors (e.g., grooming and bathing, arranging objects, object hoarding, checking) that interfere with social functioning^{16,18,28-30}.

Decreased *CYFIP1* expression is also implicated in the Prader-Willi Phenotype (**PWP**) of a subset of individuals with Fragile-X Syndrome (**FXS**). FXS is the most common genetic cause of intellectual disability and autism and is caused by a CGG trinucleotide repeat expansion within the fragile X mental retardation 1 (*FMR1*) gene that is located on the X chromosome and codes for FMRP, a major interacting protein of *CYFIP* proteins³¹. Interestingly, ten percent of FXS individuals also exhibit a PWP in the absence structural or imprinting differences in 15q11-q13. The PWP includes hallmark hyperphagia, lack of satiation, obesity, and more severe behavioral problems, such as OC behaviors and an increased rate of autism^{32,33}. The cause of the PWP is unknown, although one logical candidate

gene is *CYFIP1*, given its association with PWS and its interaction with FMRP³¹. PWP-presenting individuals with FXS show a two-to four-fold decrease in *CYFIP1* transcription compared to FXS individuals without PWP³³. There was also a two-fold decrease in *Cyfp1* gene transcription in a mouse model of FXS³⁴.

Because of the association of the gene homolog *Cyfp2* with BE⁸ and because both *CYFIP1* deletion and reduced *CYFIP1* expression are associated with PWS (Type I) and hyperphagia in the PWP (FXS) respectively, here, we tested the hypothesis that *Cyfp1* haploinsufficiency would increase premorbid compulsive-like behavior and consumption of palatable food (**PF**) in our BE paradigm^{8,35,36}. We tested the effect of *Cyfp1* deletion on two different *Cyfp2* genetic backgrounds. Additionally, because a recent study demonstrated a parental origin (**PO**) effect of *Cyfp1* haploinsufficiency on hippocampal synaptic transmission, learning, and anxiety-like behavior¹⁹, we tested for a PO effect of *Cyfp1* deletion on compulsive-like behavior and PF intake. To gain insight into the molecular mechanisms underlying the PO effect of *Cyfp1* deletion on PF intake, we examined transcription of *Cyfp1*, *Cyfp2*, and *Magel2* - a nearby imprinted gene within the syntenic, canonical PWS region implicated in hyperphagia and obesity. Additionally, we examined *CYFIP1* protein expression in the hypothalamus and nucleus accumbens as a function of both *Cyfp1* genotype and PO. Finally, because OC behaviors are associated with BE³⁷⁻³⁹ and hyperphagia in PWS⁴⁰, we employed a battery of tests to assess anxiety-like and compulsive-like behaviors and post-BE training behaviors, including compulsive-like eating and concomitant behaviors in the light/dark conflict test⁸ in *Cyfp1* haploinsufficient mice.

MATERIALS AND METHODS

Mice

All experiments were performed in accordance with the National Institutes of Health Guidelines for the Use of Laboratory Animals and were approved by the Institutional Animal Care and Use Committee at Boston University. Mice were 50-100 days old at the first day of testing. A minimum

sample size of N = 20 per Genotype per Treatment was employed for behavioral studies based on power analysis of PF intake from the *Cytip2* study⁸ (see **Supplementary Information** for additional details). Mice heterozygous for a null deletion in exons 4 through 6 of *Cytip1* (*Cytip1*^{+/-}) were propagated on two different C57BL/6 genetic backgrounds (**Fig.1**): (1) the BE-prone C57BL6/N background, and (2) a mixed background whereby mice were homozygous for the BE-resistant C57BL6/J background at the *Cytip2* locus. Details regarding generation of mice and *Cytip1* and *Cytip2* genotyping are provided in the **Supplementary Information**.

Premorbid anxiety- and compulsive-like behavioral battery

Because of the link between anxiety, compulsivity and pathological overeating³⁹ and because OC behavior is associated with eating disorders^{41,42}, we incorporated a behavioral battery to assess differences in premorbid anxiety-like and compulsive-like behaviors in experimentally naïve, *Cytip1*^{+/-} mice. Mice were tested in the behavioral battery and were either sacrificed afterward (mice on *Cytip2*^{N/N} background) or were subsequently trained for BE (mice on the *Cytip2*^{J/J} background). Mice were assayed in the battery with one test per day over five days in the following order: 1) open field; 2) elevated plus maze; 3) marble burying; 4) hole board; 5) mist-induced grooming. Procedural details are provided in the **Supplementary Information**. Testing was conducted between 0800 and 1300 h. The experimenters responsible for running the mice, video tracking, data curation, and analysis were blinded to Genotype for each cohort.

Digging, burrowing, and burying of objects with bedding are highly correlated behaviors associated with survival. Inherent burying of non-aversive stimuli can be distinguished from defensive burying of aversive, noxious stimuli⁴³ and is not related to anxiety, stimulus novelty, or locomotor activity levels. Rather, inherent marble burying is an indirect measure of the natural tendency to dig⁴⁴⁻⁴⁶. Marble burying represents persistent, repetitive behavior that is resistant to habituation and is proposed to model obsessive/compulsive behavior^{45,47}. This behavior is used to screen pharmacotherapeutics for OCD⁴⁸. Genetic variance in marble burying among inbred mouse strains is heritable but is genetically uncorrelated with anxiety-like behaviors and is thus, mediated by distinct genetic factors⁴⁵.

The hole board test is commonly used to assess anxiety⁴⁹, novelty-seeking⁵⁰, and repetitive behavior⁵¹. Moreover, head-dipping activity in the hole board test has been shown to be a valid predictor of reward-associated behaviors such as nicotine self-administration⁵², and cocaine-induced conditioned place preference⁵⁰.

BE procedure and light/dark conflict test of compulsive eating

Mice were trained in an intermittent, limited access procedure to detect genetic differences in BE^{8,35}. For details, see **Supplementary Information**. Briefly, mice were tested for side preference on D1 and D22. In the intervening days, mice were confined to a food-paired and non-food-paired side on alternating days (Tues-Fri). Cages were assigned to either the PF or Chow group in a counterbalanced design in order to ensure equal distribution across Sex, Genotype, Treatment, and PO. On D23, mice were assessed for compulsive eating and associated behaviors, as previously described⁸ (**Supplementary information**). The experimenters responsible for running the mice, video tracking, data curation, and analysis were blinded to Genotype for each cohort.

Hypothalamus dissections for real-time quantitative PCR (qPCR)

We chose a subset of Chow-trained, PF-naïve mice (n = 7-9 per Genotype per PO; both sexes) on the *Cytip2*^{N/N} background or untrained, PF-naïve mice on a *Cytip2*^{J/J} background (n= 8-12 per Genotype per PO; both sexes) to examine baseline (PF-naïve) gene transcription between *Cytip1*^{N/-} versus *Cytip1*^{N/N} mice and PO. We examined *Cytip1*, *Cytip2*, and *Magel2* transcript levels in the hypothalamus, a brain region important for hyperphagia in PWS⁴⁰ and for the effects of *Magel2* deletion^{53,54} on eating behavior and homeostatic function⁵⁵. Haploinsufficiency of *MAGEL2* is associated with PWS-like hyperphagia in humans^{54,56,57}.

On D24, brains from Chow-trained mice (*Cytip2*^{N/N} background) were harvested and the hypothalamus was free form dissected by pinching the entire structure from the ventral surface with forceps while using the anterior commissure and mammillary bodies as landmarks. Tissue was stored

in RNAlater Solution (Invitrogen, Carlsbad, CA USA) at 4°C. After five days, the tissue was dried and transferred to a -80°C freezer.

Real-time quantitative PCR (qPCR)

Total RNA from hypothalamus was extracted and processed for qPCR as described^{8,35,58}. Briefly, oligo-dT primers were used to synthesize cDNA. PCR reactions were conducted on the StepOne Plus 96-Well Real-Time PCR machine (Life Technologies, Foster City, CA, USA) in technical triplicates and averaged (SD < 0.5). Plates were balanced across Genotype, PO, and Sex. We report the difference in expression in *Cyfp1*^{+/-} relative to *Cyfp1*^{+/+} using the $2^{-(\Delta\Delta CT)}$ method⁵⁹. Primer sequences are provided in the **Supplementary Information**.

Western blot of hypothalamus and nucleus accumbens

Hypothalamus was dissected as described above. Nucleus accumbens punches were harvested using 1.2 mm punches of ventral forebrain centered around anterior commissure from the first 4 mm of brain section in a brain matrix. Samples were processed and analyzed for quantity of CYFIP1 protein. Detailed methods can be found in the **supplementary material**. Because we found no effect of Treatment for either brain region, data were collapsed across Treatment for analysis.

Analysis

Statistical analyses were conducted using R (<https://www.r-project.org>). For the compulsive behavioral tests, two-tailed unpaired t-tests were used to detect effects of Genotype for all behaviors except marble burying behaviors which were analyzed by non-parametric Mann-Whitney U tests. Slope analyses were conducted as previously described^{8,60} using GraphPad Prism 7 (GraphPad Software, La Jolla, CA USA). We analyzed food intake using mixed model ANOVAs with Genotype, Treatment, and Sex as independent variables, and Day as a repeated measure using the “aov” function in R. Additionally, we assessed the effect of PO (maternal, paternal) using mixed-model ANOVAs with Genotype, Treatment, and PO as independent variables and Day as a repeated measure. To address

issues of non-normality or unequal variance, we included additional non-parametric analyses to support key findings (**Supplementary Results**).

RESULTS

***Cytip1* haploinsufficiency increases compulsive-like behaviors**

Sample sizes are listed in **Supplementary Table 1**. **Figure 1** illustrates the breeding scheme for *Cytip1*^{+/-} mice on two *Cytip2* genetic backgrounds: *Cytip2*^{N/N} and *Cytip2*^{J/J}. Because symptomatic severity is worse in individuals with Type I PWS (which includes the *CYFIP1* deletion) and because compulsivity was negatively correlated with *CYFIP1* expression in PWS patients with Type I deletions¹⁶, we hypothesized that *Cytip1*^{+/-} mice would exhibit greater compulsive-like behavior.

In the marble burying test, *Cytip1*^{N/-} mice on the *Cytip2*^{N/N} genetic background showed a greater number of marbles that were at least 50% buried and a greater average percentage of marbles buried than *Cytip1*^{N/N} mice (**Fig.2A-C**) which are commonly used measures of marble burying⁶¹. This result was replicated in *Cytip1*^{J/-} on the *Cytip2*^{J/J} background (**Fig.2D,E**). *Cytip1* deletion on the *Cytip2*^{N/N} genetic background did not induce a change in any other behaviors within the battery (**Supplementary Table 2**, all *ps* > 0.05). However, on the *Cytip2*^{J/J} background, *Cytip1*^{J/-} showed a greater number of head dips in the hole board test than *Cytip1*^{J/J}, (**Supplementary Fig.1C**), further supporting increased compulsive-like behaviors induced by *Cytip1* haploinsufficiency.

Because *CYFIP1* is paternally-deleted in Type I PWS and because PO effects of *Cytip1* deletion on synaptic transmission and behavior were reported¹⁹, we next investigated the effect of PO of *Cytip1* deletion on anxiety-like and compulsive-like behaviors. There was no effect of PO or interaction with *Cytip1* Genotype on marble burying or any other behaviors within the battery (data not shown). To summarize, *Cytip1* haploinsufficiency induced a selective increase in compulsive-like marble burying regardless of genetic background, as well as an increase in compulsive-like head-dipping in the hole board test in mice on the *Cytip2*^{J/J} genetic background.

Effect of *Cytip1* haploinsufficiency on PF intake depends on *Cytip2* genetic background

In testing the hypothesis that *Cytip1* haploinsufficiency would increase PF intake in our intermittent, limited access BE and CPP paradigm (**Fig.3A**), we first found that PF-trained mice of both genotypes on the *Cytip2^{N/N}* background consumed significantly more food than Chow-trained mice (**Fig.3B**) – this result was reflected by slopes of escalation that were significantly greater than zero in both PF-trained genotypes but not in the Chow-trained genotypes (**Fig.3C**). As predicted, *Cytip1^{N/-}* mice consumed more PF than *Cytip1^{N/N}* mice, but not more Chow (**Fig.3B**). Finally, PF-trained *Cytip1^{N/-}* mice showed a greater y-intercept than all other groups (**Fig. 3C**), indicating an initial higher level of consumption that persisted throughout the study.

In examining the effect of *Cytip1* haploinsufficiency on food intake on a *Cytip2^{J/J}* genetic background, we observed less overall PF intake, as expected, relative to the *Cytip2^{N/N}* background [$t(155) = 2.4$; $p = 0.02$]⁸. Moreover, PF-trained mice again showed greater intake than Chow-trained mice (**Fig.3D**). However, when examining the effect of Genotype on the *Cytip2^{J/J}* background, the results were unexpected. Interestingly, *Cytip1^{J/-}* mice showed *less* PF intake than their wild-type *Cytip1^{J/J}* counterparts (**Fig.3D**) and did not escalate over time whereas wild-type mice escalated as indicated by a significant, positive slope (**Fig.3E**).

To further dissect the unexpected decrease in PF intake in *Cytip1^{J/-}* mice, we examined female and males separately. Overall, there was greater PF intake in females compared to males (**Fig.3F**). Surprisingly, male *Cytip1^{J/-}* mice completely accounted for the decrease in PF consumption in *Cytip1^{J/-}* mice (**Fig.3F**). Furthermore, neither female nor male *Cytip1^{J/-}* showed an escalation in PF consumption whereas their wild-type *Cytip1^{J/J}* counterparts of both sexes showed positive slopes (**Fig.3G**).

PO- and sex-dependent effects of *Cytip1* haploinsufficiency on PF intake

We next investigated the effect of PO of *Cytip1* deletion on food intake in light of a recent study demonstrating a PO effect of *Cytip1* deletion on emotional learning and synaptic transmission¹⁹. We focused on PF intake rather than Chow intake based on the above results.

For mice on the *Cytip2^{N/N}* background, paternal *Cytip1* deletion induced greater PF consumption in all offspring (**Figs.4A,B**). Paternally-deleted *Cytip1^{N/-}* mice showed a greater y-intercept than both

Cyfp1^{N/N} groups (**Fig.4C**), indicating overall greater intake across time. Wild-type *Cyfp1*^{N/N} offspring derived from families with paternal *Cyfp1* deletion also showed a greater y-intercept (**Fig.4C**), thus confirming an overall effect of parental *Cyfp1* genotype on neurobehavioral expression of all offspring within those families.

In examining PO effects in *Cyfp2*^{J/J} mice, maternal *Cyfp1* deletion accounted for the overall Genotype effect of decreased PF intake (**Fig4.D,E**). Additionally, only wild-type mice showed evidence for escalated intake and *Cyfp1*^{J/-} mice showed no significant slope, regardless of PO. Because we identified a Sex x Genotype x PO x Day interaction in mice with the *Cyfp2*^{J/J} background, we sought to identify the source of these interactions. When separated by Sex, paternally-deleted female *Cyfp1*^{J/-} mice showed *enhanced* PF intake as indicated by an increase in y-intercept (**Fig.4G-I**) whereas maternally-deleted male *Cyfp1*^{J/-} mice showed markedly *decreased* PF intake as indicated both by reduced PF consumption and no escalation over time (**Fig.4J-L**). To summarize, we observed markedly different effects of *Cyfp1* deletion on PF intake that depended on *Cyfp2* genetic background, PO, and Sex. Despite changes in PF intake across Genotype and PO, we did not detect any effects of Genotype or PO on body weight. (**Supplementary Fig.2**). Thus, homeostatic differences are unlikely to explain the above findings.

Conditioned food reward in *Cyfp1*^{+/-} mice

In examining CPP on the *Cyfp2*^{N/N} genetic background, there was no effect of *Cyfp1* Genotype or Treatment in the ANOVA model in mice from either genetic background (**Supplementary Fig.3A,B**). However, when we considered PF treatment alone, there was increased PF-CPP in *Cyfp1*^{N/-} versus *Cyfp1*^{N/N} mice that was in line with increased PF consumption (**Supplementary Fig.3A**). For *Cyfp1*^{J/-} mice on the *Cyfp2*^{J/J} background, there was no genotypic difference (**Supplementary Fig.3B**). In considering PO effects on PF-CPP, there was no effect of Genotype, PO, or interaction on either *Cyfp2* genetic background (**Supplementary Fig.3C,D**).

Compulsive-like eating in the light/dark conflict test in *Cyfp1*^{+/-} mice

Because we observed increased compulsive-like behavior in *Cytip1*^{+/-} mice and because increased PF consumption can become compulsive-like³⁹, we next examined post-training compulsive-like eating using the light/dark conflict test (**Fig.5A**)⁸. PF-trained mice showed greater compulsive-like consumption than Chow-trained mice (**Fig.5B,D**). Furthermore, females showed greater PF consumption than males on both genetic backgrounds (**Fig.5C,E**). *Cytip1* deletion on the *Cytip2*^{N/N} background had no effect on PF consumption (**Fig.5B,C**), regardless of PO (**Supplementary Fig.4A-C**). For the *Cytip2*^{J/J} background, *Cytip*^{J/-} mice showed reduced PF consumption that was driven primarily by the males (**Fig.5D,E**), similar to PF intake during training (**Fig.3F,G**). However, unlike the PO dependency of the reduced PF observed during training (**Fig.4J-L**), the male-specific reduction in PF intake during the compulsive-like eating test did not depend on PO (**Supplementary Fig.4D,E**).

Reduced transcription of *Cytip1* but not *Cytip2* or *Magel2* in the hypothalamus of *Cytip1*^{+/-} mice

We hypothesized that the PO- and genetic background-dependent effects of *Cytip1* deletion on PF intake could involve differences in hypothalamic gene transcription of *Cytip1* and perhaps *Cytip2*⁸. We also examined *Magel2* which is a nearby imprinted gene that is located within the syntenic, canonical (Type II) PWS locus and has been implicated in changes in eating behavior in mouse models of PWS⁵⁵ and PWS-like hyperphagia in humans^{54,56,57}. **Supplementary Table 3** lists the qPCR results as a function of both *Cytip1* haploinsufficiency and PO. As expected, *Cytip1* deletion significantly reduced gene transcription. When assessed on the *Cytip2*^{N/N} background, the reduction in *Cytip1* transcription in *Cytip1*^{N/-} mice was similar following maternal versus paternal *Cytip1* deletion (**Supplementary Table 3A**). However, when assessed on the *Cytip2*^{J/J} background, the decrease in *Cytip1* expression was enhanced following maternal *Cytip1* deletion (**Supplementary Table 3B**) which was consistent with the enhanced reduction in PF consumption in maternally-deleted *Cytip1*^{J/-} mice on the *Cytip2*^{J/J} background (**Fig.4**). Finally, there was no effect of *Cytip1* Genotype or PO on transcription of *Cytip2* or *Magel2* (**Supplementary Table 3**).

Reduced CYFIP1 protein in *Cytip1*^{+/-} mice depends on PO

We next investigated the effect of *Cytip1* haploinsufficiency on protein expression in *Cytip2*^{J/J} mice following completion of BE training. In the hypothalamus, we found a decrease in CYFIP1 protein in paternally-deleted *Cytip1*^{J/-} mice (**Supplementary Fig.5A**). When females and males were considered separately, we found a similar decrease in females resulting from paternal deletion (**Supplementary Fig.5B**), but no difference in males (**Fig.5C**). In contrast, in the nucleus accumbens, CYFIP1 expression was reduced in maternally-deleted *Cytip1*^{J/-} mice (**Fig.5D**). When we considered females and males separately, we did not detect any differences in CYFIP1 protein (**Fig.5E,F**).

DISCUSSION

Cytip1 haploinsufficiency increased OC-like behavior on two different *Cytip2* genetic backgrounds (**Figs.1-2**) and altered PF consumption and *Cytip1* gene expression, depending on genetic background (*Cytip2*), Sex, and PO (**Figs.3-5**). These findings identify a significant contribution of reduced CYFIP1 expression to OC-like behaviors and PF intake that has relevance for neurodevelopmental disorders, including Type II PWS or the PWP (**FXS**). The selective increase in sweetened PF but not Chow intake during training (**Fig.3**) and during the test for compulsive-like eating (**Fig.5**) is consistent with increased preference for sweetened PF in PWS⁶²⁻⁶⁴.

OC-like behavior and PF intake in *Cytip1*^{+/-} mice

The selective increase in OC-like but not anxiety-like behavior following *Cytip1* deletion (**Fig.2**; **Supplementary Table 2**) is consistent with a lack of genetic correlation between marble burying and anxiety and supports marble burying as a repetitive, perseverative-like behavior⁴⁵. Nevertheless, there is likely an anxiety-like component to marble burying⁶⁵ as there is with OC behaviors in humans. For the *Cytip1*^{J/J} genetic background, the increase in head-dipping behavior in the hole board task in *Cytip1*^{J/-} mice further supports an increase in OC-like behaviors associated with *Cytip1* haploinsufficiency⁶⁶.

An increase in marble burying can predict BE^{67,68} and thus, our results suggest that human genetic polymorphisms affecting CYFIP1 expression could modulate OC behavior and risk for eating disorders^{41,42}. OC behaviors are associated with both BE^{37,38} and PWS hyperphagia⁴⁰. However, in our

studies, there was no clear relationship between OC-like behaviors and PF intake. Although *Cytip1* haploinsufficiency increased marble burying on both backgrounds (**Fig.2**) and increased PF consumption on the *Cytip2*^{N/N} background, it *decreased* PF consumption on the *Cytip2*^{J/J} background (**Fig.3**) while on the same background increasing OC-like head-dipping behavior (**Supplementary Fig.1**). Thus, our results show dissociable effects of *Cytip1* deletion on OC-like behavior and the direction in modulation of PF intake. Notably, PWS patients show an increase in non-food OC behaviors, which is exacerbated in Type I PWS patients (who all have the *CYFIP1* deletion)^{16,18,28-30}. *CYFIP1* haploinsufficiency could conceivably enhance OC behaviors and affect eating behaviors through separate, independent neural mechanisms. Additionally, although *CYFIP1* deletion is associated with the more severe, Type I form of PWS, we are not aware of any studies showing increased severity of hyperphagia in Type I PWS. Thus, *CYFIP1* genotype could act more generally as a modifier of PWS hyperphagia rather than distinguishing PWS subtypes.

Effect of *Cytip1* deletion on PF intake depends on genetic background

As predicted, mice on a *Cytip2*^{N/N} versus *Cytip2*^{J/J} background showed greater overall PF consumption (**Fig.3B-E**)⁶⁹. Additionally, *Cytip1*^{N/-} mice showed a greater increase in PF consumption and escalation (**Fig.3D,E**). In contrast, *Cytip1*^{J/-} mice on the lower PF-consuming *Cytip2*^{J/J} background showed a *decrease* in escalation of PF intake (**Fig.3D,E**). This result was driven by the surprising induction of BE in wild-type *Cytip1*^{J/J} males (**Fig.3F,G**), specifically males derived from maternal *Cytip1*^{J/-} deletion (**Fig.4J,L**). The escalated intake in wild-type *Cytip1*^{J/J} males was opposite to the low-level BE on the *Cytip2*^{J/J} background⁸ and is inconsistent with a lack of BE in the parental B6J strain, especially males^{8,35}. While we bred the *Cytip2* locus to be fixed for the J allele, heterozygous N alleles segregating elsewhere in the genome could contribute to BE in *Cytip2*^{J/J} mice. In support, F2 mice possessing the *Cytip2*^{J/J} genotype showed greater PF consumption than the parental B6J strain⁸. However, heterozygosity at other loci cannot fully explain the current results, since only a specific subset of wild-type males (maternal *Cytip1*^{J/-} - derived) showed an anomalous induction of escalation in consumption. Thus, wild-type *Cytip1*^{J/J} males could be especially sensitive to social influences of

maternal-pup and/or pup-pup interactions in the maternally-deleted *Cytip1*^{J/-} environment, ultimately explaining escalation of PF intake.

In stark contrast to male *Cytip1*^{J/-} mice with the *maternal* deletion, female *Cytip1*^{J/-} mice (**Fig.3F,G**) with the *paternal* *Cytip1*^{J/-} deletion showed a markedly enhanced, initial consumption of PF that obfuscated detection of escalation over time (**Fig.4H,I**). In considering sex differences in PWS, male PWS patients with 15q11-q13 deletions showed greater food-related preoccupation with food, impaired satiety, and other food-related negative behaviors in the absence of differences in OC or other non-food-related behaviors⁷⁰. It is unclear whether sex differences in food-related problems differ depending on Type I versus Type II PWS deletions. In addition to behavioral differences, female PWS patients showed higher levels of circulating insulin and HOMA-IR, indicating greater insulin resistance and decreased adiponectin levels⁷¹. Together, our results illustrate the importance of considering Sex when investigating maternal versus paternal gene haploinsufficiency in eating behavior and the potential relevance to PWS in humans.

Interactions of PO and offspring genotype in offspring behavior

The increase in PF consumption in all offspring from paternal *Cytip1*^{N/-} deletion (**Fig. 4B vs. Fig.4A**) and the robust increase in PF consumption in wild-type *Cytip1*^{J/J} males derived from maternal but not paternal *Cytip1*^{J/-} deletion (**Fig.4J-L**) highlight potential genetic interactions with social environment in explaining behavioral variance⁷². Maternal versus paternal *Cytip1* deletion could affect social interactions with the dam and sire or with the pups. Of direct relevance to *Cytip1*, maternal deletion of the *Fmr1* gene (coding for FMRP) can induce neurobehavioral phenotypes in wild-type offspring and enhance phenotypes in mutant offspring, including locomotor hyperactivity, reduced behavioral response to D2 dopamine receptor activation, and enhanced behavioral response to GABA-B receptor activation⁷³. In addition, males can demonstrate paternal pup retrieval⁷⁴ and thus, *Cytip1* deletion could also affect sire-pup contact. Given the association between *CYFIP1* deletion and social deficits in both 15q11.2 MDS and PWS but also FXS, autism, and schizophrenia¹⁴, it is plausible that

Cyfp1 deletion in the dam or sire affects social dynamics in the offspring in a PO-specific and genotype (offspring)-dependent manner, leading to long-term neurobehavioral effects.

Hedonic hypothesis of *Cyfp1* modulation of PF intake

Despite differences in PF consumption as a function of *Cyfp1* Genotype, PO, Sex, and Genetic Background, there were no genotypic differences in body weight (**Supplementary Fig.2**), suggesting that *Cyfp1* haploinsufficiency could modulate PF intake through a non-homeostatic, perhaps hedonic mechanism involving altered sensory or affective processing of sweetened PF^{12,13}. The selective changes in PF consumption and conditioned reward (at least in *Cyfp1*^{N/-} mice; **Supplementary Fig.3A**) support the hedonic hypothesis and are consistent with enhanced preference for sweetened food in individuals with Type I PWS⁶²⁻⁶⁴. The *Cyfp2*^{N/N} mutation is associated with cocaine neurobehavioral sensitivity and plasticity⁹ and compulsive-like BE⁸. Furthermore, differences in *Cyfp2* mRNA expression were genetically correlated with differences in cocaine self-administration⁷⁵. Here, we observed PO-dependent decreases in CYFIP1 protein in both nucleus accumbens and hypothalamus (**Supplementary Fig.5**). Previous transcriptome analysis of the striatum from *Cyfp2*^{N/-} versus *Cyfp2*^{N/N} genotypes identified “morphine addiction” and “cocaine addiction” as two of the top five KEGG enrichment terms⁸. FMRP, a major interacting protein with CYFIP1/2, is involved in reward processing⁷⁶ and cocaine neurobehavioral plasticity⁷⁷. We hypothesize that *Cyfp1* deletion/polymorphisms could alter the hedonic effects of PF consumption via the dopaminergic mesolimbic reward circuitry and interact with other haploinsufficient genes underlying hypothalamic, homeostatic mechanisms of hyperphagia in PWS to modulate food intake^{78,79}.

Sex-dependent PO effects on *Cyfp1* expression and behavior

There is no published evidence that *Cyfp1* is imprinted and while our analysis of *Cyfp1* transcript and protein levels indicate PO-dependent effects, the direction was not always consistent with maternal imprinting and was dependent on Sex and the particular brain region (**Supplementary Table 3; Supplementary Fig.5**). A recent study of nearly 100 phenotypes showed that most complex

traits exhibit PO effects and that non-imprinted KO alleles (e.g., *Cytip1*) can induce extensive PO effects by interacting in *trans* with imprinted loci throughout the genome to affect gene networks⁸⁰. If a *trans*-acting genomic mechanism underlies the effects of *Cytip1* haploinsufficiency, it would have to be co-inherited faithfully with the maternal or paternal deletion to explain the PO effects on behavior. An obvious candidate mechanism could involve inheritance of sex-dependent gene expression originating from sex chromosomes (and potential sex chromosome variants between substrains that would explain genetic background-dependent effects) that interacts with *Cytip1* deletion. This is particularly relevant to CYFIP proteins given that *Fmr1*, which codes for FMRP, the interacting protein of CYFIP that regulates protein translation and is located on the X chromosome. The human *FMR1* gene undergoes X-inactivation⁸¹; thus, females and males should have equivalent levels of gene dosage, transcription, and translation of FMRP. Nevertheless, 10-25% of human genes and 3-7% of mouse genes show variable degrees of X inactivation^{82,83}. *Fmr1* could undergo variable x-inactivation depending on the tissue and time point. Furthermore, genetic polymorphisms between B6J and B6NJ on the X chromosome could affect the expression of genes that act as modifiers of *Cytip1* transcription or modifiers of X-inactivation and account for the background-dependent PO effects of *Cytip1* deletion on behavior. The use of the four core genotypes model (XX, XY, XX-male, XY-female) could be used to test the involvement of sex chromosomes in PO- and Sex-dependent effects of *Cytip1* deletion on PF intake - a recent study using this genetic model identified a contribution of sex chromosomes to operant reinforcement for PF⁸⁴.

Conclusion

Our preclinical findings provide evidence that reduced CYFIP1 expression could contribute to OC and eating behaviors in PWS and other neurodevelopmental disorders. Future genomic studies of multiple brain regions, cell types, and developmental time points could inform molecular mechanisms of eating behaviors on different genetic backgrounds and potential interaction of *Cytip1* deletion with gene expression on sex chromosomes. Furthermore, PO effects of *Cytip1* deletion could be tested for behavioral and genomic interactions with imprinted genes in PWS models of hyperphagia (e.g., deletion

of *Magel2* or *Snord116*). Such efforts could improve upon existing PWS models that, like our findings, have historically lacked obesity⁸⁵ and could inform pharmacotherapeutic treatment of eating behavior tailored to a particular subtype of neurodevelopmental syndrome.

ACKNOWLEDGEMENTS

Funding was provided by NIH/NIDA R21DA038738 (C.D.B.), NIH/NIDA R01DA039168, NIDA Diversity Scholars Network (F.R.), Burroughs-Wellcome Fund Transformative Training Program in Addiction Science (TTPAS; 1011479), NIH/NIGMS T32GM008541, and NIH/NIDA U01DA041668 (V.K.). We thank Dr. Rachel Wevrick for providing us with the primer sequences used for gene expression analysis of *Magel2*. We would also like to acknowledge Dr. Lynn Deng and Matthew Au of the Boston University Analytical Instrumentation Core Facility (S10OD023663) for their support in conducting the qPCR studies.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

References

1. Hudson JI, Hiripi E, Pope HG,Jr, Kessler RC. The prevalence and correlates of eating disorders in the national comorbidity survey replication. *Biol Psychiatry* 2007; **61**: 348-58.
2. Mitchell KS, Neale MC, Bulik CM, Aggen SH, Kendler KS, Mazzeo SE. Binge eating disorder: A symptom-level investigation of genetic and environmental influences on liability. *Psychol Med* 2010; **40**: 1899-906.
3. Bulik CM, Sullivan PF, Kendler KS. Genetic and environmental contributions to obesity and binge eating. *Int J Eat Disord* 2003; **33**: 293-8.
4. Yilmaz Z, Hardaway JA, Bulik CM. Genetics and epigenetics of eating disorders. *Adv Genomics Genet* 2015; **5**: 131-50.
5. Hinney A, Kesselmeier M, Jall S, Volckmar AL, Focker M, Antel J, et al. Evidence for three genetic loci involved in both anorexia nervosa risk and variation of body mass index. *Mol Psychiatry* 2017; **22**: 192-201.
6. McElroy SL, Winham SJ, Cuellar-Barboza AB, Colby CL, Ho AM, Sicotte H, et al. Bipolar disorder with binge eating behavior: A genome-wide association study implicates PRR5-ARHGAP8. *Transl Psychiatry* 2018; **8**: 40,017-0085-3.
7. Huckins LM, Hatzikotoulas K, Southam L, Thornton LM, Steinberg J, Aguilera-McKay F, et al. Investigation of common, low-frequency and rare genome-wide variation in anorexia nervosa. *Mol Psychiatry* 2017; .
8. Kirkpatrick SL, Goldberg LR, Yazdani N, Babbs RK, Wu J, Reed ER, et al. Cytoplasmic FMR1-interacting protein 2 is a major genetic factor underlying binge eating. *Biol Psychiatry* 2017; **81**: 757-69.

9. Kumar V, Kim K, Joseph C, Kourrich S, Yoo SH, Huang HC, et al. C57BL/6N mutation in cytoplasmic FMRP interacting protein 2 regulates cocaine response. *Science* 2013; **342**: 1508-12.
10. Bello NT, Hajnal A. Dopamine and binge eating behaviors. *Pharmacol Biochem Behav* 2010; **97**: 25-33.
11. Berridge KC. Wanting and liking: Observations from the neuroscience and psychology laboratory. *Inquiry (Oslo)* 2009; **52**: 378.
12. DiLeone RJ, Taylor JR, Picciotto MR. The drive to eat: Comparisons and distinctions between mechanisms of food reward and drug addiction. *Nat Neurosci* 2012; **15**: 1330-5.
13. Lutter M, Nestler EJ. Homeostatic and hedonic signals interact in the regulation of food intake. *J Nutr* 2009; **139**: 629-32.
14. Abekhoukh S, Bardoni B. CYFIP family proteins between autism and intellectual disability: Links with fragile X syndrome. *Front Cell Neurosci* 2014; **8**: 81.
15. Pathania M, Davenport EC, Muir J, Sheehan DF, Lopez-Domenech G, Kittler JT. The autism and schizophrenia associated gene CYFIP1 is critical for the maintenance of dendritic complexity and the stabilization of mature spines. *Transl Psychiatry* 2014; **4**: e374.
16. Bittel DC, Kibiryeva N, Butler MG. Expression of 4 genes between chromosome 15 breakpoints 1 and 2 and behavioral outcomes in prader-willi syndrome. *Pediatrics* 2006; **118**: e1276-83.
17. Cox DM, Butler MG. The 15q11.2 BP1-BP2 microdeletion syndrome: A review. *Int J Mol Sci* 2015; **16**: 4068-82.
18. Doornbos M, Sikkema-Raddatz B, Ruijvenkamp CA, Dijkhuizen T, Bijlsma EK, Gijsbers AC, et al. Nine patients with a microdeletion 15q11.2 between breakpoints 1 and 2 of the prader-willi critical region, possibly associated with behavioural disturbances. *Eur J Med Genet* 2009; **52**: 108-15.

19. Chung L, Wang X, Zhu L, Towers AJ, Cao X, Kim IH, et al. Parental origin impairment of synaptic functions and behaviors in cytoplasmic FMRP interacting protein 1 (Cyfip1) deficient mice. *Brain Res* 2015; **1629**: 340-50.
20. Hsiao K, Harony-Nicolas H, Buxbaum JD, Bozdagi-Gunal O, Benson DL. Cyfip1 regulates presynaptic activity during development. *J Neurosci* 2016; **36**: 1564-76.
21. Oguro-Ando A, Rosensweig C, Herman E, Nishimura Y, Werling D, Bill BR, et al. Increased CYFIP1 dosage alters cellular and dendritic morphology and dysregulates mTOR. *Mol Psychiatry* 2015; **20**: 1069-78.
22. Bozdagi O, Sakurai T, Dorr N, Pilorge M, Takahashi N, Buxbaum JD. Haploinsufficiency of Cyfip1 produces fragile X-like phenotypes in mice. *PLoS One* 2012; **7**: e42422.
23. Angulo MA, Butler MG, Cataletto ME. Prader-willi syndrome: A review of clinical, genetic, and endocrine findings. *J Endocrinol Invest* 2015; .
24. State MW, Dykens EM, Rosner B, Martin A, King BH. Obsessive-compulsive symptoms in prader-willi and "prader-willi-like" patients. *J Am Acad Child Adolesc Psychiatry* 1999; **38**: 329-34.
25. Dykens EM, Leckman JF, Cassidy SB. Obsessions and compulsions in prader-willi syndrome. *J Child Psychol Psychiatry* 1996; **37**: 995-1002.
26. Stein DJ, Keating J, Zar HJ, Hollander E. A survey of the phenomenology and pharmacotherapy of compulsive and impulsive-aggressive symptoms in prader-willi syndrome. *J Neuropsychiatry Clin Neurosci* 1994; **6**: 23-9.
27. Feurer ID, Dimitropoulos A, Stone WL, Roof E, Butler MG, Thompson T. The latent variable structure of the compulsive behaviour checklist in people with prader-willi syndrome. *J Intellect Disabil Res* 1998; **42 (Pt 6)**: 472-80.

28. Butler MG, Bittel DC, Kibiryeva N, Talebizadeh Z, Thompson T. Behavioral differences among subjects with prader-willi syndrome and type I or type II deletion and maternal disomy. *Pediatrics* 2004; **113**: 565-73.
29. Milner KM, Craig EE, Thompson RJ, Veltman MW, Thomas NS, Roberts S, et al. Prader-willi syndrome: Intellectual abilities and behavioural features by genetic subtype. *J Child Psychol Psychiatry* 2005; **46**: 1089-96.
30. Zarcone J, Napolitano D, Peterson C, Breidbord J, Ferraioli S, Caruso-Anderson M, et al. The relationship between compulsive behaviour and academic achievement across the three genetic subtypes of prader-willi syndrome. *J Intellect Disabil Res* 2007; **51**: 478-87.
31. Schenck A, Bardoni B, Moro A, Bagni C, Mandel JL. A highly conserved protein family interacting with the fragile X mental retardation protein (FMRP) and displaying selective interactions with FMRP-related proteins FXR1P and FXR2P. *Proc Natl Acad Sci U S A* 2001; **98**: 8844-9.
32. Muzar Z, Lozano R, Kolevzon A, Hagerman RJ. The neurobiology of the prader-willi phenotype of fragile X syndrome. *Intractable Rare Dis Res* 2016; **5**: 255-61.
33. Nowicki ST, Tassone F, Ono MY, Ferranti J, Croquette MF, Goodlin-Jones B, et al. The prader-willi phenotype of fragile X syndrome. *J Dev Behav Pediatr* 2007; **28**: 133-8.
34. Stefan M, Portis T, Longnecker R, Nicholls RD. A nonimprinted prader-willi syndrome (PWS)-region gene regulates a different chromosomal domain in trans but the imprinted pws loci do not alter genome-wide mRNA levels. *Genomics* 2005; **85**: 630-40.
35. Goldberg LR, Kirkpatrick SL, Yazdani N, Luttik KP, Lacki OA, Babbs RK, et al. Casein kinase 1-epsilon deletion increases mu opioid receptor-dependent behaviors and binge eating. *Genes Brain Behav* 2017; .

36. Babbs RK, Kelliher JC, Scotellaro JL, Yazdani N, Ruan QT, Mulligan MK, et al. Genetic differences in the behavioral organization of binge eating, conditioned food reward, and compulsive-like eating in C57BL/6J and DBA/2J strains. *BioRxiv* 2017; .
37. Wilfley DE, Citrome L, Herman BK. Characteristics of binge eating disorder in relation to diagnostic criteria. *Neuropsychiatr Dis Treat* 2016; **12**: 2213-23.
38. Kessler RM, Hutson PH, Herman BK, Potenza MN. The neurobiological basis of binge-eating disorder. *Neurosci Biobehav Rev* 2016; **63**: 223-38.
39. Moore CF, Sabino V, Koob GF, Cottone P. Pathological overeating: Emerging evidence for a compulsivity construct. *Neuropsychopharmacology* 2017; **42**: 1375-89.
40. Griggs JL, Sinnayah P, Mathai ML. Prader-willi syndrome: From genetics to behaviour, with special focus on appetite treatments. *Neurosci Biobehav Rev* 2015; **59**: 155-72.
41. Micali N, Hilton K, Nakatani E, Heyman I, Turner C, Mataix-Cols D. Is childhood OCD a risk factor for eating disorders later in life? A longitudinal study. *Psychol Med* 2011; **41**: 2507-13.
42. Cavallini MC, Bertelli S, Chiapparino D, Riboldi S, Bellodi L. Complex segregation analysis of obsessive-compulsive disorder in 141 families of eating disorder probands, with and without obsessive-compulsive disorder. *Am J Med Genet* 2000; **96**: 384-91.
43. De Boer SF, Koolhaas JM. Defensive burying in rodents: Ethology, neurobiology and psychopharmacology. *Eur J Pharmacol* 2003; **463**: 145-61.
44. Njung'e K, Handley SL. Evaluation of marble-burying behavior as a model of anxiety. *Pharmacol Biochem Behav* 1991; **38**: 63-7.

45. Thomas A, Burant A, Bui N, Graham D, Yuva-Paylor LA, Paylor R. Marble burying reflects a repetitive and perseverative behavior more than novelty-induced anxiety. *Psychopharmacology (Berl)* 2009; **204**: 361-73.
46. Gyertyan I. Analysis of the marble burying response: Marbles serve to measure digging rather than evoke burying. *Behav Pharmacol* 1995; **6**: 24-31.
47. Deacon RM. Digging and marble burying in mice: Simple methods for in vivo identification of biological impacts. *Nat Protoc* 2006; **1**: 122-4.
48. Hoffman KL, Cano-Ramirez H. Lost in translation? A critical look at the role that animal models of obsessive compulsive disorder play in current drug discovery strategies. *Expert Opin Drug Discov* 2017; 1-10.
49. do-Rego JC, Viana AF, Le Maitre E, Deniel A, Rates SM, Leroux-Nicollet I, et al. Comparisons between anxiety tests for selection of anxious and non anxious mice. *Behav Brain Res* 2006; **169**: 282-8.
50. Arenas MC, Daza-Losada M, Vidal-Infer A, Aguilar MA, Minarro J, Rodriguez-Arias M. Capacity of novelty-induced locomotor activity and the hole-board test to predict sensitivity to the conditioned rewarding effects of cocaine. *Physiol Behav* 2014; **133**: 152-60.
51. Moy SS, Nadler JJ, Poe MD, Nonneman RJ, Young NB, Koller BH, et al. Development of a mouse test for repetitive, restricted behaviors: Relevance to autism. *Behav Brain Res* 2008; **188**: 178-94.
52. Abreu-Villaca Y, Queiroz-Gomes Fdo E, Dal Monte AP, Filgueiras CC, Manhaes AC. Individual differences in novelty-seeking behavior but not in anxiety response to a new environment can predict nicotine consumption in adolescent C57BL/6 mice. *Behav Brain Res* 2006; **167**: 175-82.

53. Mercer RE, Michaelson SD, Chee MJ, Atallah TA, Wevrick R, Colmers WF. Magel2 is required for leptin-mediated depolarization of POMC neurons in the hypothalamic arcuate nucleus in mice. *PLoS Genet* 2013; **9**: e1003207.
54. Tacer KF, Potts PR. Cellular and disease functions of the prader-willi syndrome gene MAGEL2. *Biochem J* 2017; **474**: 2177-90.
55. Bervini S, Herzog H. Mouse models of prader-willi syndrome: A systematic review. *Front Neuroendocrinol* 2013; **34**: 107-19.
56. Schaaf CP, Gonzalez-Garay ML, Xia F, Potocki L, Gripp KW, Zhang B, et al. Truncating mutations of MAGEL2 cause prader-willi phenotypes and autism. *Nat Genet* 2013; **45**: 1405-8.
57. Fountain MD, Schaaf CP. Prader-willi syndrome and schaaaf-yang syndrome: Neurodevelopmental diseases intersecting at the MAGEL2 gene. *Diseases* 2016; **4**: 10.3390/diseases4010002.
58. Yazdani N, Parker CC, Shen Y, Reed ER, Guido MA, Kole LA, et al. Hnrnp1 is A quantitative trait gene for methamphetamine sensitivity. *PLoS Genet* 2015; **11**: e1005713.
59. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nature protocols* 2008; **3**: 1101-8.
60. Babbs RK, Wojnicki FH, Corwin RL. Assessing binge eating. an analysis of data previously collected in bingeing rats. *Appetite* 2012; **59**: 478-82.
61. Angoa-Perez M, Kane MJ, Briggs DI, Francescutti DM, Kuhn DM. Marble burying and nestlet shredding as tests of repetitive, compulsive-like behaviors in mice. *J Vis Exp* 2013; **(82)**:50978. doi: 50978.

62. Martinez Michel L, Haqq AM, Wismer WV. A review of chemosensory perceptions, food preferences and food-related behaviours in subjects with prader-willi syndrome. *Appetite* 2016; **99**: 17-24.
63. Hinton EC, Holland AJ, Gellatly MS, Soni S, Owen AM. An investigation into food preferences and the neural basis of food-related incentive motivation in prader-willi syndrome. *J Intellect Disabil Res* 2006; **50**: 633-42.
64. Taylor RL, Caldwell ML. Type and strength of food preferences of individuals with prader-willi syndrome. *J Ment Defic Res* 1985; **29 (Pt 1)**: 109-12.
65. Albelda N, Joel D. Animal models of obsessive-compulsive disorder: Exploring pharmacology and neural substrates. *Neurosci Biobehav Rev* 2012; **36**: 47-63.
66. Takeda H, Tsuji M, Matsumiya T. Changes in head-dipping behavior in the hole-board test reflect the anxiogenic and/or anxiolytic state in mice. *Eur J Pharmacol* 1998; **350**: 21-9.
67. Satta V, Scherma M, Giunti E, Collu R, Fattore L, Fratta W, et al. Emotional profile of female rats showing binge eating behavior. *Physiol Behav* 2016; **163**: 136-43.
68. Freund N, Thompson BS, Norman KJ, Einhorn P, Andersen SL. Developmental emergence of an obsessive-compulsive phenotype and binge behavior in rats. *Psychopharmacology (Berl)* 2015; **232**: 3173-81.
69. Kirkpatrick SL, Bryant CD. Behavioral architecture of opioid reward and aversion in C57BL/6 substrains. *Front Behav Neurosci* 2015; **8**: 450.
70. Gito M, Ihara H, Ogata H, Sayama M, Murakami N, Nagai T, et al. Gender differences in the behavioral symptom severity of prader-willi syndrome. *Behav Neurol* 2015; **2015**: 294127.

71. Irizarry KA, Bain J, Butler MG, Ilkayeva O, Muehlbauer M, Haqq AM, et al. Metabolic profiling in prader-willi syndrome and nonsyndromic obesity: Sex differences and the role of growth hormone. *Clin Endocrinol (Oxf)* 2015; **83**: 797-805.
72. Baud A, Mulligan MK, Casale FP, Ingels JF, Bohl CJ, Callebert J, et al. Genetic variation in the social environment contributes to health and disease. *PLoS Genet* 2017; **13**: e1006498.
73. Gleason G, Zupan B, Toth M. Maternal genetic mutations as gestational and early life influences in producing psychiatric disease-like phenotypes in mice. *Front Psychiatry* 2011; **2**: 25.
74. Liu HX, Lopatina O, Higashida C, Fujimoto H, Akther S, Inzhutova A, et al. Displays of paternal mouse pup retrieval following communicative interaction with maternal mates. *Nat Commun* 2013; **4**: 1346.
75. Dickson PE, Miller MM, Calton MA, Bubier JA, Cook MN, Goldowitz D, et al. Systems genetics of intravenous cocaine self-administration in the BXD recombinant inbred mouse panel. *Psychopharmacology (Berl)* 2015; .
76. Fish EW, Krouse MC, Stringfield SJ, Diberto JF, Robinson JE, Malanga CJ. Changes in sensitivity of reward and motor behavior to dopaminergic, glutamatergic, and cholinergic drugs in a mouse model of fragile X syndrome. *PLoS One* 2013; **8**: e77896.
77. Smith LN, Jedynak JP, Fontenot MR, Hale CF, Dietz KC, Taniguchi M, et al. Fragile X mental retardation protein regulates synaptic and behavioral plasticity to repeated cocaine administration. *Neuron* 2014; **82**: 645-58.
78. Heksch R, Kamboj M, Anglin K, Obrynba K. Review of prader-willi syndrome: The endocrine approach. *Transl Pediatr* 2017; **6**: 274-85.

79. Holsen LM, Savage CR, Martin LE, Bruce AS, Lepping RJ, Ko E, et al. Importance of reward and prefrontal circuitry in hunger and satiety: Prader-willi syndrome vs simple obesity. *Int J Obes (Lond)* 2012; **36**: 638-47.
80. Mott R, Yuan W, Kaisaki P, Gan X, Cleak J, Edwards A, et al. The architecture of parent-of-origin effects in mice. *Cell* 2014; **156**: 332-42.
81. Kirchgessner CU, Warren ST, Willard HF. X inactivation of the FMR1 fragile X mental retardation gene. *J Med Genet* 1995; **32**: 925-9.
82. Carrel L, Brown CJ. When the lyon(ized chromosome) roars: Ongoing expression from an inactive X chromosome. *Philos Trans R Soc Lond B Biol Sci* 2017; **372**: 10.1098/rstb.2016.0355.
83. Balaton BP, Brown CJ. Escape artists of the X chromosome. *Trends Genet* 2016; **32**: 348-59.
84. Seu E, Groman SM, Arnold AP, Jentsch JD. Sex chromosome complement influences operant responding for a palatable food in mice. *Genes Brain Behav* 2014; **13**: 527-34.
85. Rodriguez JA, Zigman JM. Hypothalamic loss of Snord116 and prader-willi syndrome hyperphagia: The buck stops here? *J Clin Invest* 2018; .
86. Skarnes WC, Rosen B, West AP, Koutsourakis M, Bushell W, Iyer V, et al. A conditional knockout resource for the genome-wide study of mouse gene function. *Nature* 2011; **474**: 337-42.

FIGURE LEGENDS

Figure 1: Generation of the *Cytip1* knockout allele and breeding scheme for *Cytip1* haploinsufficient mice on *Cytip2*^{N/N} and *Cytip2*^{J/J} genetic backgrounds. (A): A schematic of the knockout first allele for KOMP generation of *Cytip1*^{N/-} mice was obtained from the International Mouse Phenotyping Consortium (IMPC) website ([http://www.mousephenotype.org/data/alleles/MGI:1338801/tm2a\(EUCOMM\)Wtsi](http://www.mousephenotype.org/data/alleles/MGI:1338801/tm2a(EUCOMM)Wtsi)). Mice containing floxed alleles flanking exons 4 through 6 were generated from embryonic stem cells on a C57BL/6N background by the International Knockout Mouse Consortium and were crossed to global Cre-expressing mice, yielding offspring heterozygous for constitutive deletions in exons 4 through 6. We derived mice heterozygous for the null deletion on a C57BL/6N background using sperm obtained from The Jackson Laboratory. **(B): Left panel:** In the first study, we re-derived *Cytip1*^{N/-} mice on an isogenic C57BL/6N background. **Right panel:** All mice were homozygous for the N allele (N/N) at *Cytip2* which contains a missense mutation that we previously showed was associated with a marked enhancement of binge eating (BE), accounting for approximately 50% of the genetic variance in parental strain BE⁸. We maintained this colony on an isogenic C57BL/6N background by breeding *Cytip1*^{N/-} mice with C57BL/6NJ mice (black bars; N/N) ordered from The Jackson laboratory. **(C):** In the second study, we generated another colony on a mixed background. The primary goal was to monitor and replace the BE-associated N/N *Cytip2* alleles with C57BL/6J (J/J) alleles via backcrossing *Cytip1*^{N/-} mice to C57BL/6J (white bars; J/J) for three and four generations and assess the effect of *Cytip1* deletion on BE on a mixed N3 and N4 background containing a fixed, BE-resistant, homozygous J/J genotype at *Cytip2*⁸. Mixed-color bars illustrate hypothetical recombination events that accumulate through backcrossing to C57BL/6J (white).

Figure 2. Increase in premorbid, OC-like marble burying in *Cytip1*^{N/-} and *Cytip1*^{J/-} mice. (A): Schematic of the marble burying apparatus. **(B,C):** *Cytip1*^{N/-} mice buried more marbles with greater than 50% coverage than wild-type *Cytip1*^{N/N} mice [B: U(102) = 1050; p = 0.031; two-tailed], and had a greater average percentage of burial across the marbles [C; F(1,96) = 7.1; p = 0.009]. **(D,E):** Similarly,

Cyfp1^{J/-} mice buried more marbles with greater than 50% coverage than J/J mice [**D**: U(137) = 1884; $p = 0.019$, two-tailed], and also had a greater average percentage of burial [**E**: $F(1,134) = 4.2$; $p = 0.042$]. Data are presented as mean $\pm$ SEM.

Figure 3. PF consumption in *Cyfp1*^{N/-} and *Cyfp1*^{J/-} mice. (A): The conditioned place preference chamber used for food consumption training had a smooth-textured non-food-paired side (left) and a rough-textured food-paired side. **(B):** Both wild-type *Cyfp1*^{N/N} and *Cyfp1*^{N/-} mice trained with PF in the CPP chamber ate more food over time than Chow-trained mice [# main effect of Treatment: $F(1,932) = 274.7$; $p < 2 \times 10^{-16}$]. There was also a main effect of Sex [$F(1,932) = 30.4$; $p = 4.5 \times 10^{-8}$], a Genotype x Treatment interaction [$F(1,932) = 4.7$; $p = 0.03$], and a Treatment x Sex interaction [$F(1,932) = 22.3$; $p = 2.7 \times 10^{-6}$]. **(C):** Both PF-trained genotypes exhibited slopes that were significantly greater than zero (*Cyfp1*^{N/N}: $m = 0.009 \pm 0.003$, $p = 0.024$; *Cyfp1*^{N/-}: $m = 0.005 \pm 0.002$, $p = 0.045$, respectively) indicating escalation in PF intake over time. Neither Chow-trained group showed escalation. Moreover, PF-trained *Cyfp1*^{N/-} mice showed a significantly greater y-intercept than all other groups (\$: $p < 0.008$), indicating consistently greater overall food consumption throughout the study. **(D):** When we examined the same behaviors in *Cyfp1*^{J/-} versus *Cyfp1*^{J/J} mice, we observed a main effect of Treatment [$F(1,824) = 200.4$; $p = 2 \times 10^{-16}$], indicating that PF-trained mice consumed more food over time. We also observed a main effect of Genotype [$F(1,824) = 4.2$; $p = 0.04$], but in contrast to *Cyfp1*^{N/-} mice, *Cyfp1*^{J/-} consumed less food than their wild-type *Cyfp1*^{J/J} counterparts. Additionally, we observed a main effect of Sex [$F(1,542) = 27.0$, $p = 3.0 \times 10^{-7}$], Day [$F(1,824) = 17.9$; $p = 2.6 \times 10^{-5}$], a Sex x Treatment interaction [$F(1,824) = 6.3$; $p = 0.01$], a Treatment x Day interaction [$F(1,824) = 16.9$; $p = 4.3 \times 10^{-5}$], a Genotype x Day interaction [$F(1,824) = 6.7$; $p = 0.01$], and most importantly, a Sex x Treatment x Genotype interaction [$F(1,824) = 11.9$; $p = 0.0006$]. **(E):** In examining escalation in food intake over time, only PF-trained *Cyfp1*^{J/J} mice exhibited a positive non-zero slope [#; $F(1,244) = 21.0$; $p < 0.0001$], indicating escalation in consumption over time and further supporting reduced food intake in *Cyfp1*^{J/-} mice. **(F,G):** In order to better understand the interactions with Sex in PF-trained mice, we examined PF consumption in *Cyfp1*^{J/-} versus *Cyfp1*^{J/J} mice in females and males. We found main

effects of Sex [$F(1,446) = 11.2$; $p = 0.0009$], Genotype [$F(1,446) = 4.2$; $p = 0.04$], and Day [$F(1,446) = 21.3$; $p = 5.2 \times 10^{-6}$]. We also observed a Genotype x Day interaction [$F(1,446) = 4.8$; $p = 0.03$], a Sex x Genotype interaction [$F(1,446) = 7.8$; $p = 0.005$], a Gene x PO interaction [$F(1,446) = 10.6$; $p = 0.001$], and a Sex x Genotype x PO interaction [$F(1,446) = 5.7$; $p = 0.02$]. **(G)** In examining escalation in PF consumption, we found that both female and male wild-type *Cytip1*^{J/J} mice had positive slopes (#; both p s < 0.009 vs. zero) whereas both female and male *Cytip1*^{J/-} mice did not have any significant slope (both p s > 0.08). However, the *Cytip1*^{J/-} females had the greatest y-intercept ($p < 0.0001$) indicating the greatest initial consumption. Data are presented as mean $\pm$ SEM.

Figure 4. Effect of parent-of-origin (PO) on PF consumption in *Cytip1*^{N/-} and *Cytip1*^{J/-} mice. (A,B):

In considering mice on a *Cytip2*^{N/N} background, there was a main effect of Genotype [$F(1,464) = 12.3$; $p = 0.0005$], PO [$F(1,464) = 9.0$; $p = 0.003$], Sex [$F(1,464) = 39.4$; $p = 8.0 \times 10^{-10}$], and Day [$F(1,464) = 20.8$; $p = 6.5 \times 10^{-6}$]. The effect of Genotype was explained by *Cytip1*^{N/-} mice consuming more PF than *Cytip1*^{N/N} mice [*; $t(29) = 2.1$; uncorrected $p = 0.046$ vs. *Cytip1*^{N/N} on D4]. The effect of Sex was explained by females consuming more PF than males (not shown). The effect of PO was explained by offspring derived from parents with a paternal *Cytip1* deletion **(B)** consuming a greater amount of PF than offspring derived from parents with a maternal *Cytip1* deletion **(A)**. **(C)**: No differences were observed among the groups in the slopes of escalation in PF consumption [$F(3,16) = 0.7$ $p = 0.56$]; however, paternally-deleted *Cytip1*^{N/-} mice (open squares) showed a greater overall consumption than either of the *Cytip1*^{N/N} wild-type groups as indicated by a greater y-intercept (\$: both p 's < 0.02) and *Cytip1*^{N/N} mice derived from families with a paternal deletion showed a greater y-intercept than *Cytip1*^{N/N} mice with a maternal deletion ($p = 0.046$). **(D,E)**: In considering the effects of *Cytip1* deletion and PO on a *Cytip2*^{J/J} background, there was a main effect of Genotype [$F(1,446) = 4.1$; $p = 0.04$], Sex [$F(1,446) = 11.8$; $p = 0.0006$], Day [$F(1,446) = 21.3$; $p = 5.2 \times 10^{-6}$] and importantly, there was a Genotype x PO interaction [$F(1,454) = 10.4$; $p = 0.001$] that reflected less PF consumption in *Cytip1*^{J/-} mice with the maternal *Cytip1*^{J/-} deletion **(D)**; * $t(41) = 2.4$; uncorrected $p = 0.02$ vs. *Cytip1*^{J/J} on D11) but not paternal *Cytip1*^{J/-} deletion **(E)**. **(F)**: We observed significant escalation in consumption in both *Cytip1*^{J/J} wild-type

groups [maternal: $F(1,136) = 12.6$; $p = 0.0005$; paternal: $F(1,106) = 9.6$; $p = 0.003$]. Neither *Cytip1*^{J/-} group had a significant non-zero slope (both p s > 0.09). Moreover, *Cytip1*^{J/-} mice with a paternal deletion had a greater y-intercept than all three of the other groups ($\$$; all p s < 0.0004). **(G-L)**: We also found a Sex x Genotype interaction [$F(1,446) = 9.3$, $p = 0.002$], a Gene x PO interaction [$F(1,446) = 9.1$; $p = 0.003$], and a Genotype x Sex x PO x Day interaction [$F(1,446) = 5.7$; $p = 0.02$]. To understand the source of these interactions, we next separated PO effects of *Cytip1*^{J/-} by Sex. **(G-I)**: In the females, we observed a main effect of Day [$F(1,220) = 10.6$; $p = 0.001$] and a Genotype x PO interaction [$F(1,220) = 4.9$; $p = 0.03$], **(G,H)**. Moreover, both *Cytip1*^{J/J} wild-type groups showed a significant escalation, regardless of PO (**I**; both p s < 0.02). Interestingly, although the *Cytip1*^{J/-} mice with a paternal deletion had the greatest y-intercept ($\$$; $p < 0.0001$), they had a negative slope of escalation (**I**; $-3.4 \times 10^{-5} \pm 7.5 \times 10^{-5}$) indicating initially higher PF consumption but a progressive decrease in consumption over time. **(J-L)**: For males there was an effect of Genotype [$F(1,226) = 13.6$; $p = 0.0003$], Day [$F(1,226) = 10.7$; $p = 0.001$], a Genotype x PO interaction [$F(1,226) = 5.7$; $p = 0.02$], and a Genotype x PO x Day interaction [$F(1,226) = 3.9$; $p = 0.049$]. that was explained by maternally-deleted *Cytip1*^{J/-} mice eating markedly less PF than their *Cytip1*^{J/J} wild-type counterparts or, i.e., the induction of robust escalation in consumption in the wild-type *Cytip1*^{J/J} males (**J**, * $p < 0.03$ vs *Cytip1*^{J/J} on D16). In contrast, paternally-deleted *Cytip1*^{J/-} mice showed no difference relative to their wild-type *Cytip1*^{J/J} counterparts (**K**). In examining the slopes of escalation in all four groups, only the wild-type *Cytip1*^{J/J} males coming from maternal deletion showed a significant escalation over time (**L**; $p = 0.01$). Data are presented as mean $\pm$ SEM.

Figure 5. Compulsive-like PF intake in the light/dark conflict test in *Cytip1*^{N/-} and *Cytip1*^{J/-} mice.

(A): A cartoon of the apparatus for the light/dark conflict test of compulsive-like PF consumption is shown. **(B)**: For the *Cytip2*^{N/N} background, there was a main effect of Training Treatment [$F(1,148) = 31.1$; $p = 1 \times 10^{-7}$], indicating that PF-trained mice showed increased PF intake. However, there was no effect of Genotype [$F(1,148) = 0.1$; $p = 0.7$] or interaction of Genotype with Training Treatment on consumption [$F(1,148) = 2.3$; $p = 0.13$]. **(C)**: In examining PF-trained mice alone, females showed

increased intake [$*F(1,72) = 13.6$; $p = 0.0004$]; however, there was no effect of Genotype [$F(1,72) = 1.2$; $p = 0.3$] or Genotype x Sex interaction [$F(1,72) = 0.7$; $p = 0.4$]. **(D)**: For mice on the *Cytip2*^{J/J} genetic background, there was a main effect of Training Treatment [$*F(1,132) = 10.4$; $p = 0.002$], indicating greater PF intake in PF-trained mice. There was also a main effect of Genotype [$F(1,132) = 8.9$; $p = 0.003$]. PF-trained *Cytip1*^{J/-} mice consumed less than PF-trained *Cytip1*^{J/J} mice [$†$; $t(75) = 2.5$; $p = 0.02$]. **(E)**: In considering only PF-trained mice, females showed greater overall intake [$*F(1,69) = 4.1$; $p = 0.047$] and *Cytip1*^{J/-} mice showed overall less intake [$F(1,69) = 4.9$; $p = 0.03$]. However, only male *Cytip1*^{J/-} mice consumed less than male *Cytip1*^{J/J} mice [$‡$; $t(37) = 2.7$; $p = 0.01$]. Data are presented as mean $\pm$ SEM.

FIG.2




FIG.3




FIG.4




FIG.5

