

**Unique mapping of structural and functional connectivity on cognition**

**Abbreviated title: Unique mapping of SC and FC on cognition**

Zimmermann, J.\* , Griffiths, J.G., McIntosh, A.R.

*Baycrest Health Sciences, Rotman Research Institute, 3560 Bathurst St, Toronto, Ontario,*

*M6A 2E1, Canada*

## Abstract

The unique mapping of structural and functional brain connectivity (SC, FC) on cognition is currently not well understood. It is not clear whether cognition is mapped via a global connectome pattern or instead is underpinned by several sets of distributed connectivity patterns. Moreover, we also do not know whether the pattern of SC and of FC that underlie cognition are overlapping or distinct. Here, we study the relationship between SC and FC and an array of psychological tasks in 609 subjects from the Human Connectome Project (HCP). We identified several sets of connections that each uniquely map onto different aspects of cognitive function. We found a small number of distributed SC and a larger set of cortico-cortical and cortico-subcortical FC that express this association. Importantly, SC and FC each show unique and distinct patterns of variance across subjects and differential relationships to cognition. The results suggest that a complete understanding of connectome underpinnings of cognition calls for a combination of the two modalities.

## Significance Statement

Structural connectivity (SC), the physical white-matter inter-regional pathways in the brain, and functional connectivity (FC), the temporal co-activations between activity of brain regions, have each been studied extensively. Little is known, however, about the distribution of variance in connections as they relate to cognition. Here, in a large sample of subjects (N = 609), we showed that two sets of brain-behavioural patterns capture the correlations between SC, and FC with a wide range of cognitive tasks, respectively. These brain-behavioural patterns reveal distinct sets of connections within the SC and the FC network and provide new evidence that SC and FC each provide unique information for cognition.

## Introduction

In neuroscience, big data initiatives such as the Human Connectome Project (HCP) acquire connectomic and phenotypic data from a large number of individuals in an effort to understand how brain networks relate to individual behaviour (Van Essen et al., 2013; Fornito, 2016). Connectomes can represent either structural connectivity (SC), the white-matter inter-regional pathways estimated from diffusion-weighted MR imaging (Baldassarre et al., 2012), or functional connectivity (FC), the patterns of temporal dependencies between regional activity measurements such as blood oxygen level dependent functional magnetic resonance imaging (fMRI) timeseries. The most commonly studied form of FC, resting-state FC (rsFC), is measured in the absence of an explicit task. It represents meaningful coordinated fluctuations that have been related to task FC and performance (Mennes et al., 2010; Baldassarre et al., 2012; Stevens and Spreng, 2014).

Both SC (Matejko et al., 2013; Willmes et al., 2014; Moeller et al., 2015; Klein et al., 2016) and rsFC (Song et al., 2008; Song et al., 2009; Pamplona et al., 2015; Santarnechchi et al., 2015; Hearne et al., 2016; Smith, 2016; Ferguson MA, 2017; Pezoulas et al., 2017) have been linked to cognitive functioning, including higher order cognitive processes such as fluid and crystallized intelligence, visuospatial processing (Ponsoda et al., 2017), or numerical cognition (Matejko et al., 2013; Willmes et al., 2014; Moeller et al., 2015; Klein et al., 2016).

What remains unclear, however, is the specificity of the behaviourally-relevant aspect of the connectome. That is, can individual variability across connectomes be explained by a single set of connections that is highly predictive of cognitive function *more generally*? Or are there multiple sets of connections that *each predict different*

aspects of cognitive functioning? The former suggests that connectomes are relevant for understanding general differences in global cognitive functioning whilst the latter suggests that individual differences in particular networks can inform specific types of cognitive functioning, such as attention, memory, or executive function. Evidence of both views exists. For instance, Rosenberg et al. (2013) showed that a particular set of connectivity patterns predicted individuals' attention ability, supporting the notion of specific connectome-behaviour associations. The connections identified were specific to attention ability and did not predict cognition more generally. In contrast, other research has shown that a single mode of covariation can capture relationships between a distributed set of functional connections and a wide set of behavioural and demographic variables (Smith et al., 2015). The same question can be asked for SC.

Currently, little is known about the overlap between the two modalities in their mapping to individual differences in cognition. Given the limitations of mapping individual SC to FC (Koch et al., 2002; Skudlarski et al., 2008; Honey et al., 2009; Zimmermann et al., 2016; Zimmermann et al., 2018), we might expect that the two modalities provide unique and distinct sources of cognitive-related variability (Duda et al., 2010; Hirsiger et al., 2016). On the other hand, cognition arises from an interplay of structure and function, and so a degree of overlap in the spatial pattern of networks that give rise to cognitive function is expected.

In the present study, we examine how cortical and subcortical SC and rsFC from 609 subjects from the Human Connectome Project relates to a wide range of cognitive functions, including working memory, executive function/cognitive flexibility, processing speed, fluid intelligence, episodic memory, and attention/inhibitory control. We examined

1) whether there exists a single set of connections that generally map onto cognition, or rather several sets of connections that map onto different aspects of cognition, and 2) whether the patterns of connectivity that map onto cognition are independent and unique for SC versus rsFC or whether they provide common information. We used Partial Least Squares to map orthogonal patterns of brain-behavioural relationships (McIntosh and Lobaugh, 2004; Krishnan et al., 2011). This method is comparable to canonical correlation, however, is more suitable for neuroimaging data as it is robust to collinearity within the datasets.

## Methods

### *Subjects and behavioural measures*

The sample included 609 genetically unrelated subjects from the Q7 HCP release (M/F = 269/340 female; age range = 22-36) (Van Essen et al., 2013). The research was performed in compliance with the Code of Ethics of the World Medical Association (Declaration of Helsinki). All subjects provided written informed consent, approved by the ethics committee in accordance with guidelines of HCP WU-Minn. In the current study, 11 behavioural measures of cognitive function were correlated with SC and FC (described below). Cognitive measures covered a range of processes. Note that the working memory 0-back and 2-back tasks each included 2 measures: accuracy and reaction time. The measures are described in Table 1 below and in detail in the following: <https://wiki.humanconnectome.org/display/PublicData/HCP+Data+Dictionary+Public+500+Subject+Release>. Note that cognitive scores were age adjusted.

Table 1. Cognitive tests and corresponding function

Test	Function	M (SD), range
ListSort_AgeAdj	Working memory (List sorting)	103 (13), 63-141
PicSeq_AgeAdj	Episodic Memory (Picture sequence memory)	106 (17), 56-135
CardSort_AgeAdj	Executive Function / Cognitive flexibility (Dimensional Change card sort)	102 (10), 58-123
Flanker_AgeAdj	Executive Function/Inhibition (Flanker Task)	102 (10), 74-124
ProcSpeed_AgeAdj	Processing Speed (Pattern Completion Processing Speed)	103 (19), 45-149
WM_Task_2bk_Acc	Working Memory accuracy (2-back task)	83 (11), 37-100
WM_Task_2bk_Median_RT	Working Memory reaction time (2-back task)	970 (143), 602-1440
WM_Task_0bk_Acc	Working Memory accuracy (0-back task)	89 (11), 48-100
WM_Task_0bk_Median_RT	Working memory reaction time	775 (139), 506-1571

---

(0-back task)		
Fluid Intelligence correct		
PMAT24_A_CR	responses	17 (5), 5-24
(Penn Progressive Matrices)		
PMAT24_A_RTCR	Fluid Intelligence reaction time, correct responses	15454 (2976), 1989-61641
(Penn Progressive Matrices)		

---

### ***MRI acquisition and preprocessing***

Data acquisition details for the WU-Minn HCP corpus (Van Essen et al., 2013) were described in detail elsewhere (Smith et al., 2013; Sotiropoulos et al., 2013; Ugurbil et al., 2013). The present study made use of resting-state BOLD fMRI, multi-shell (multiple non zero b-values) high-resolution dwMRI, and structural (T1-weighted MRI) data from the HCP minimally preprocessed pipeline (Glasser et al., 2013; Smith et al., 2013).

Diffusion data with a spin-echo multiband EPI sequence, with 111 slices of 1.25 mm isotropic voxels (Feinberg et al., 2010; Sotiropoulos et al., 2013; Ugurbil et al., 2013), distortion corrected (Andersson et al., 2003; Andersson and Sotiropoulos, 2015) were used. Segmentation and parcellation was performed on the basis of the high-resolution T1-weighted image (voxel size: 0.7 mm isotropic) of each subject using FreeSurfer (Fischl, 2012), automatically parcellating the brain into 83 cortical and subcortical ROIs (41 per hemisphere, plus brainstem) according to the Lausanne 2008 atlas (Daducci et al., 2012). An 83 x 83 connectivity matrix was formed, representing for each pair of regions their reconstructed pathways. Deterministic tractography using Dipy software (Garyfallidis et al.,

2014) was performed on minimally preprocessed data using a method similar to that of Hagmann et al (2008) and Cammoun et al (2012). Streamlines were computed from 60 equally spaced points within each voxel on the grey-white matter interface via the EuDX algorithm (Garyfallidis et al (2012)). Streamlines that were shorter than 10mm or longer than 250mm were discarded, as well as those that did not terminate at the grey-white matter interface. The strength of the reconstructed connections was measured as streamline density, computed as the number of tractography streamlines that touched both cortical regions (Hagmann et al., 2008).

Whole brain echo-planar images (EPI) (TR = 720 ms, 2 mm<sup>3</sup> voxels) (Moeller et al., 2010; Ugurbil et al., 2013) with denoising procedures from the resting-state FIX denoised dataset were used (Glasser et al., 2013; Smith et al., 2013). Data included both phase encoding acquisition directions (L-R, R-L) (acquisition time = 14 mins, 33 sec). We computed average regional timeseries from these voxels, based on the 83 cortical and subcortical regions. Pearson's correlations were calculated for each pair of timeseries to compute the FC between all regions; these were then Fisher Z-transformed, and the two (L-R and R-L) pairs of matrices were averaged for each subject. The resulting FC matrices were 83x83 for the 609 subjects. Global signal regression was not performed for comparability with previous resting state studies. Note that age was regressed from SC and Fisher's z-transformed FC, and residuals were used for analysis.

### ***Experimental Design and Statistical Analysis***

#### **Cognitive Measures**

The 11 cognitive measures were correlated (Pearson's) with one another to quantify the relationship between them. The resulting p-value cross-correlation matrix was of size 11 x 11. The p-value of each correlation value was corrected for multiple comparisons using FDR (Matlab function `fdr_bky`) (Benjamini, 2006). We also wanted to understand whether the cognitive measures describe a single global cognitive component of intelligence, or whether numerous components are required to capture the different subtypes of cognitive functioning. To this end, we conducted a principal components analysis (PCA) of the cognitive measures (Matlab function `princomp`). To assess the significance of the resulting eigenvalues, we permuted the cognitive measures 100 times (scrambled across cognitive measures and subjects) and decomposed (PCA) the permuted cognitive matrices to obtain a null distribution of eigenvalues for each PC. We then calculated the p-value per PC eigenvalue as the proportion of times the permuted eigenvalue exceeded the obtained eigenvalue.

#### *Correlation between cognition and connectivity*

We used Partial Least Squares (PLS) for neuroimaging (McIntosh and Lobaugh, 2004; Krishnan et al., 2011) in order to assess the multidimensional connectome-cognition relationships. We used the Behavioral PLS correlation function described there, ran in Matlab using custom code. As the connectivity matrices are symmetric, we used the vectorized upper triangle of the connectivity matrices from the 609 subjects. Connections that were 0 for 66% of subjects or more were not included in the analysis, resulting in 870 remaining connections for SC and 3403 connections for FC. Vectorized connectomes were then stacked resulting in a subjects\*connections brain matrix, for SC = 609\*870, and for

FC = 609\*3403. The behavioural matrix was subjects\*behavioural measures, of size 609\*11.

The brain and behavioural matrices were then cross-correlated to start the PLS analysis. PLS is multivariate analysis method comparable to canonical correlation, and captures maximally covarying brain-behaviour relationships in mutually orthogonal latent variables (LVs). The significance of LVs is assessed via permutation tests (1000 iterations) of the singular values from singular value decomposition (SVD) of the brain and behavioural correlation matrices, and reliability of each connectivity estimate to the LV is assessed via bootstrap resampling (3000 iterations). The reliability of each connection's loading onto the brain-behaviour relationship in each LV is represented as a bootstrap ratio, the ratio of a connection's weight over its estimated standard error. The ratio can be considered equivalent to a z-score, but is used to impart reliability rather than significance. A connection with a positive high bootstrap ratio contributes positively and reliably to the brain-behaviour correlation obtained for that LV, whereas a connection with a negative high bootstrap ratio contributes negatively and reliably to the brain-behaviour relationship. Bootstrapping is also used to construct confidence intervals on the brain-behaviour correlations.

### SC-Cognition vs FC-Cognition

In order to compare the connections that contributed to the SC-cognition relationship and those that contributed to the FC-cognition relationship, we calculated the scalar dot product (Matlab function dot) between the brain scores ("U") from the PLS SVD expressed in the FC-cognition and those expressed in the SC-cognition, across all

significant LVs. We did the same thing for cognitive measures between analyses, using Pearson's correlations instead of dot products. Note that these correlations were corrected for multiple comparisons using FDR (Matlab function `fdr_bky`) (Benjamini, 2006).

## Results

### Cognitive measures

The cognitive scores correlated positively amongst one another, with the exception of the WM reaction time tasks, which correlated negatively with all other cognitive tests (See Figure 1). All correlations were significant (all  $p < 0.001$ , MC corrected, See Methods).




Figure 1. Correlation (Pearson's) amongst the cognitive tests.

A PCA of the 11 cognitive test scores yielded three significant PCs, where significance was assessed by permutation testing (See Methods) ( $\lambda = 3.5072, 1.8679$ , and  $1.2026$ , % variance explained = 61, 17, 7 respectively) (Figure 2). All cognitive tests

loaded onto the first PC, with an emphasis on the speed-accuracy trade-off for the two WM tests. Thus, participants who were more accurate were also slower on these WM tests. The second PC emphasized the PMAT tests; note that reaction time for correct responses (PMAT RT) and the number of correct responses (PMAT CR) were positively correlated. The third PC emphasized similarities between the two WM RT tests and the remaining tests, in opposition to the two WM accuracy measures.




Figure 2. PCA of the 11 cognitive tests. Shown here are the principal component coefficients (loadings) on each PC. A) PC1 B) PC2 C) PC3.

### **Correlation between cognition and connectivity**

PLS analyses identified two significant LVs that describe the relationship between SC and cognition, and FC and cognition each, respectively.

#### ***SC-Cognition***

The two LVs that captured the SC-cognition association revealed two distinct patterns of cognitive functions that mapped onto two sets of unique structural connections (Figure 3). (LV1 SC: 37.22% of total covariance, singular value = 2.53,  $p = 0.04$ , LV2 SC:

14.81% of total covariance, singular value = 1.6,  $p = 0.03$ ). For both LVs, a small number of connections across cortical and subcortical regions were stable by bootstrap. Connections loaded positively and negatively onto each of the SC-cognition LVs (See bootstrap ratios in Figure 3B, 3D). For the cognitive measures, the first LV strongly expressed the full array of cognitive tests, with a speed-accuracy trade-off for the WM tests. The second LV expressed primarily the two PMAT tests.




Figure 3. SC-Cognition A) LV1 correlations between SC and cognition, with CIs from bootstrap resampling, B) LV1 bootstrap ratios, these are connection loadings on the SC-cognition relationship, bootstrap threshold = -2.5, 2.5. Connections with positive bootstrap ratios contribute positively to the SC-cognition correlation, negative bootstrap ratios

contribute negatively to the SC-cognition correlation. Right and left hemisphere regions, and subcortical and cortical regions, are separated by a space in the circular plot. C) LV2 correlations between SC and cognition D) LV2 bootstrap ratios

### ***FC-Cognition***

The two LVs that captured the FC-cognition association revealed two distinct patterns of cognitive functions that map onto two sets of functional connections (Figure 4). (LV1 FC: 41.5% of total covariance, singular value = 6.45,  $p = 0.01$  LV2 FC: 33% of total covariance, singular value = 5.75,  $p = 0.046$ ).

The first LV revealed FC-cognition correlations expressed across an array of cognitive tests, and a large number of cortical, mainly fronto-parietal connections (as well as insula, transverse temporal) that load negatively onto this relationship (See bootstrap ratios in Figure 4B, 4D). Thus, better performance on card-sorting, processing speed, WM accuracy, as well as better performance on the three RT measures (better performance = lower RT) was associated with lower FC. Only 4 connections loaded negatively onto this LV, 3 of which were connections of the R entorhinal. The second LV revealed FC-cognition correlations that were expressed primarily with the PMAT cognitive tests. A large number of inter-hemispheric cortico-cortical and cortico-subcortical connections loaded positively onto this LV. Thus, better performance on the PMAT tasks correlated with higher FC.




Figure 4. FC-Cognition A) LV1 correlations between FC and cognition, with CIs from bootstrap resampling, B) LV1 bootstrap ratios, these are connection loadings on the FC-Cognition relationship, bootstrap threshold = -3, 3. Connections with high bootstrap ratios contribute positively to the FC-cognition correlation, low bootstrap ratios contribute negatively to the FC-cognition correlation. Right and left hemisphere regions, and

subcortical and cortical regions, are separated by a space in the circular plot. B) LV2 correlations between FC and cognition D) LV2 bootstrap ratios.

### ***SC-Cognition vs FC-Cognition***

As evident from Fig 3 and 4, there were more connections in the FC that correlate with cognition compared to in the SC. This can be shown as the number of connections that exceeded the bootstrap ratio threshold for each connection-cognition LV, expressed as a proportion of the number of connections that entered the PLS analysis (3403 vs 870, FC, SC). The proportion of connections that exceeded the bootstrap threshold were as follows, FC: LV1 = 15%, LV2 = 9%; SC: LV1 = 2%, LV2 = 1%. We hypothesized that this may be due to the difference in total between-subjects variance in the SC and FC. To this end, we conducted a SVD of the subject-wise SC (609 subjects \* 870 connections) as well as the subject-wise FC (609 subjects \* 3403 connections) and compared the sum of the squared singular values corrected by the number of connections that entered the analysis, in order to account for the sparsity of the SC ( $\sum S^2_{\text{corrected}}$ ). Indeed, we found that the total corrected variance in subject-wise SC was smaller than in the subject-wise FC ( $\sum S^2_{\text{SC\_corrected}} = 0.0198$ ,  $\sum S^2_{\text{FC\_corrected}} = 0.0294$ ).

We observed that a very different pattern of connections in the SC related to cognition than in the FC. This can be expressed quantitatively as the dot product of the PLS brain scores (“U”) from the FC-cognition SVD and from the SC-cognition SVD, across both LVs. These dot products were close to zero (Table 2). This comparison was made for the behaviour contributions to each LV as well. To this end, we calculated the dot product of the PLS cognitive scores (“V”) from the FC-cognition SVD and the cognitive

scores from the SC-cognition SVD (Table 3). In order to map which cognitive PCs were expressed in which LVs, we correlated the cognitive PC loadings with the SC-cognition and FC-cognition correlations on each LV (Table 4). Note that these correlations were all significant,  $p < 0.001$ , MC corrected (See Methods). We found that some cognitive PCs were particularly strongly captured in specific connectivity-cognition LVs. For example, the first cognitive PC was expressed very strongly in the first SC-cognition LV. Yet, even the second SC-cognition LV correlated well with this PC.

Table 2. Dot product between brain scores for FC-cognition and SC-cognition across both LVs.

FC-Cognition		
SC-Cognition	LV1	LV2
LV1	0.0043	0.0792
LV2	0.0159	-0.0419

Table 3. Dot product between behavioural scores for FC-cognition and SC-cognition across both LVs.

FC-Cognition		
SC-Cognition	LV1	LV2
LV1	0.51	0.83
LV2	0.38	-0.07

Table 4. Correlations of cognitive PCs loadings and SC-Cognition and FC-Cognition LVcorrs.

Cognition	SC-Cog		FC-Cog	
	LV1	LV2	LV1	LV2
PC1	0.9825	0.7415	0.6798	0.6473
PC2	-0.2102	-0.4327	-0.8290	0.3725
PC3	-0.5635	-0.2377	-0.0664	-0.5344

## Discussion

In the present study, we compared whole-brain cortical and subcortical SC and rsFC from 609 subjects from the Human Connectome Project with individual cognitive function across an array of 11 measures, including measures of working memory (WM n-back accuracy and RT, and ListSort), executive function/cognitive flexibility (CardSort), processing speed (ProcSpeed), fluid intelligence (PMAT), episodic memory (PicSeq), and attention/inhibitory control (Flanker). We first showed that the cognitive measures mapped onto three principal components (PCs) reflecting the heterogenous nature of functions measured via the tests. PC1 was a global cognition component that emphasized a speed-accuracy trade-off for the two WM tests. PC2 expressed fluid intelligence (PMAT). PC3 emphasized a more deliberate, slower, executive function/cognitive flexibility and attention/inhibitory control.

The conclusions of the study are two-fold: 1) that unique sets of connections map onto specific components of cognitive function, and 2) that SC and FC each capture

independent and unique connections that relate to cognition. Importantly, previous approaches to studying connectivity-cognition relations have constrained the dimensionality via PCA before conducting Canonical Correlation Analysis (CCA) (Smith, 2016), which constrains how the behaviour can be projected into the brain space. In contrast, we compared the conjunction of brain and behaviour without first constraining dimensionality by PCA. Moreover, unlike CCA, PLS is robust to the collinearity that may be present in connectivity data. Thus, our method may be more sensitive in being able to extract the dimensions that relate brain and behaviour.

### ***Unique sets of connections map onto specific components of cognitive functions***

First, we showed that SC and rsFC each capture multiple sets of connection-cognition associations. Two LVs for SC, and two for FC, each expressed a distinct and unique set of connections and components of cognitive function. This is in line with the view that distinct sets of connections are required to support heterogeneous components of cognitive function, implying more specialized connectivity (Rosenberg et al., 2013) as opposed to the view that a global cognitive factor can be captured via a single set of connections (Malpas et al., 2016; Smith, 2016).

The two SC-cognition LVs each captured a limited, unique set of primarily intra-hemispheric cortical and subcortical connections. The first SC-cognition LV mapped almost perfectly onto the first global cognition PC, which can be attributed to the low variability across SCs. The second SC-cognition LV strongly expressed fluid intelligence (PMAT test) via a set of distributed cortical and subcortical connections.

The two FC-cognition LVs each captured a large set of unique inter and intra-hemispheric connections. The connectivity-cognition correlations we found were comparable to those identified previously with the HCP dataset (i.e.,  $r = 0.045$ ) (Hearne et al., 2016). The first FC-cognition LV expressed fronto-parietal, visual, and cingulate cortical connections, as well as transverse temporal, insula and L hippocampus connections to the rest of the cortex. rsFC in these regions was negatively associated with processing speed, executive function/cognitive flexibility, and working memory performance. The second FC-cognition LV was expressed via widely distributed cortico-cortical and cortico-subcortical connections, with strong contributions from the bilateral caudate and putamen connections to cortical regions. A large number of inter-hemispheric connections were expressed in this LV. The PMAT fluid intelligence measures were particularly dominant, with higher rsFC correlating with higher PMAT correct response as well as reaction time scores, so that subjects with higher rsFC were more accurate, but slower. We note that for both SC and FC, the PMAT measures were strongly expressed in the second LVs. The PMAT is an abbreviated version of the Raven's Progressive Matrices (Prabhakaran et al., 1997; Gray et al., 2003; Wendelken et al., 2008), and is a measure of fluid intelligence that has been previously linked to individual differences in FC (Finn et al., 2015; Hearne et al., 2016; Smith, 2016; Ferguson MA, 2017).

The patterns of connectivity that were correlated with cognitive variability in our study were consistent with those observed previously. On the side of SC, the limited, widespread connections that associated with cognition were consistent with what had previously been reported by Ponsoda, Martinez et al (2017), where only 36 connections distributed across the entire brain predicted higher order cognitive functions. A number of

these connections mapped closely onto the SC connections that we identified across the two LVs. On the side of the rsFC, connectivity distributed across the cortex likely supports cognitive function (Ferguson MA, 2017). Positive rsFC-cognition associations have typically been identified in fronto-parietal regions (Hearne et al., 2016), and negative associations in the default mode and dorsal attention network including visual and cingulate-parietal connectivity (Song et al., 2008; Song et al., 2009; Pamplona et al., 2015; Santarnecchi et al., 2015). Our first and second FC-cognition LV captured these negative and positive associations, respectively. Our second FC-cognition LV expressed several cortico-subcortical connections, primarily between the putamen, caudate, thalamus and the rest of the cortex. This is not surprising, as the cortical control of behaviour is mediated via several cortico-striatal-thalamic-cortical circuits (Alexander et al., 1986; Peters et al., 2016) that can be identified via structural and functional imaging (Seeley et al., 2007; Metzger et al., 2010). Connectivity from subcortical areas including the striatum have previously been tied to individual differences in phenotype and behaviour (Vaidya and Gordon, 2013). The cortico-striatal-thalamic cognitive control loop also includes the brainstem (Peters et al., 2016), which exhibited cognition-related connectivity with the caudate in our LV2. Moreover, the striatum (putamen and caudate) is specifically involved in learning, storing and processing memories (Packard and Knowlton, 2002), and higher WM performance has been tied to higher connectivity between the cingulo-opercular network and putamen rsFC (Tu et al., 2012). In our first LV, the only subcortical region that had negative rsFC associations with cognition was the L hippocampus, consistent with prior work (Salami et al., 2014). Insular connectivity, primarily with anterior cingulate

regions, was also particularly prominent in the first LV, a region that may be important for reactive attentional control (Jiang et al., 2015).

Interestingly, in a sample of 317 HCP subjects, Hearne et al. (2016) identified only positive network-level associations between rsFC and two cognitive measures (one of these was the same PMAT measure used in the present study). As the authors themselves note, this may be because their network-level approach overlooks any existing edge level connectivity-cognition relationships that may exist (Song et al., 2008; Song et al., 2009; Pamplona et al., 2015; Santarnecci et al., 2015). We replicated the positive association between rsFC and PMAT scores that was found previously (Hearne et al., 2016) within our second FC-cognition LV, where fluid intelligence was emphasized as the strongest cognitive correlate.

### ***SC and FC each capture independent and unique connections that relate to cognition***

Second, we showed that that SC and FC each captured independent and complementary features of the connectome that were linked to cognitive function. The connections that expressed the SC-cognition association did not overlap with behaviourally-relevant connections in the FC, evidenced by the comparison of the spatial pattern of brain scores between the two analyses. While a similar suggestion has previously been made for SC versus task FC (Duda et al., 2010) and rsFC in select pathways (Hirsiger et al., 2016), it has not as of yet been examined in whole-brain SC-rsFC in as large a sample as ours. We found that far fewer connections within the SC compared to the FC associated with cognition, even when correcting for the sparsity of the SC. This was likely due to the comparably smaller amount of total variance in the SC

across subjects. The limited, yet distributed nature of the SC network that varies with cognition has previously been reported (Ponsoda et al., 2017). Yet, we note that the amount of covariance accounted for by SC and by FC towards cognition was comparable. This suggests that both modalities are equally important for understanding individual differences in cognition.

It is important to consider that the imperfect association between SC and FC (Koch et al., 2002; Skudlarski et al., 2008; Honey et al., 2009) may impose limitations on the amount of overlapping information that can be provided by the two modalities. Moreover, the large repertoire of cognitive brain function is made possible by virtue of the dynamic nature of FC, which arises from a static SC backbone (Park and Friston, 2013). In this vein, an investigation into functional connectivity dynamics may help describe how the spatial contributions of SC and rsFC to cognition fluctuate over time. This investigation is, however, beyond the scope of this study.

## References

Alexander GE, DeLong MR, Strick PL (1986) Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Ann Rev Neurosci* 9:357-381.

Andersson JL, Sotiroopoulos SN (2015) Non-parametric representation and prediction of single- and multi-shell diffusion-weighted MRI data using Gaussian processes. *NeuroImage* 122:166-176.

Andersson JL, Skare S, Ashburner J (2003) How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging. *NeuroImage* 20:870-888.

Baldassarre A, Lewis CM, Committeri G, Snyder AZ, Romani GL, Corbetta M (2012) Individual variability in functional connectivity predicts performance of a perceptual task. *Proceedings of the National Academy of Sciences of the United States of America* 109:3516-3521.

Benjamini Y, Krieger, A.M. & Yekutieli, D. (2006) Adaptive linear step-up procedures that control the false discovery rate. *Biometrika* 93:491-507.

Daducci A, Gerhard S, Griffa A, Lemkadem A, Cammoun L, Gigandet X, Meuli R, Hagmann P, Thiran JP (2012) The connectome mapper: an open-source processing pipeline to map connectomes with MRI. *PloS one* 7:e48121.

Duda JT, McMillan C, Grossman M, Gee JC (2010) Relating structural and functional connectivity to performance in a communication task. *Med Image Comput Comput Assist Interv* 13:282-289.

Feinberg DA, Moeller S, Smith SM, Auerbach E, Ramanna S, Gunther M, Glasser MF, Miller KL, Ugurbil K, Yacoub E (2010) Multiplexed echo planar imaging for sub-second whole brain fMRI and fast diffusion imaging. *PloS one* 5:e15710.

Ferguson MA AJ, Spreng RN (2017) Fluid and flexible minds: Intelligence reflects synchrony in the brain's intrinsic network architecture. *Network Neuroscience* 1:192-207.

Finn ES, Shen X, Scheinost D, Rosenberg MD, Huang J, Chun MM, Papademetris X, Constable RT (2015) Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. *Nature neuroscience* 18:1664-1671.

Fischl B (2012) FreeSurfer. *NeuroImage* 62:774-781.

Fornito A (2016) Fundamentals of brain network analysis / Alex Fornito, Andrew Zalesky, Edward T. Bullmore. Amsterdam ; Boston: Elsevier/Academic Press.

Glasser MF, Sotiroopoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, Xu J, Jbabdi S, Webster M, Polimeni JR, Van Essen DC, Jenkinson M, Consortium WU-MH (2013) The minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage* 80:105-124.

Gray JR, Chabris CF, Braver TS (2003) Neural mechanisms of general fluid intelligence. *Nature neuroscience* 6:316-322.

Hagmann P, Cammoun L, Gigandet X, Meuli R, Honey CJ, Wedeen V, Sporns O (2008) Mapping the structural core of human cerebral cortex. *Plos Biology* 6:1479-1493.

Hearne LJ, Mattingley JB, Cocchi L (2016) Functional brain networks related to individual differences in human intelligence at rest. *Sci Rep* 6:32328.

Hirsiger S, Koppelmans V, Merillat S, Liem F, Erdeniz B, Seidler RD, Jancke L (2016) Structural and functional connectivity in healthy aging: Associations for cognition and motor behavior. *Human brain mapping* 37:855-867.

Honey CJ, Sporns O, Cammoun L, Gigandet X, Thiran JP, Meuli R, Hagmann P (2009) Predicting human resting-state functional connectivity from structural connectivity. *Proceedings of the National Academy of Sciences of the United States of America* 106:2035-2040.

Jiang J, Beck J, Heller K, Egner T (2015) An insula-frontostriatal network mediates flexible cognitive control by adaptively predicting changing control demands. *Nat Commun* 6:8165.

Klein E, Suchan J, Moeller K, Karnath HO, Knops A, Wood G, Nuerk HC, Willmes K (2016) Considering structural connectivity in the triple code model of numerical cognition: differential connectivity for magnitude processing and arithmetic facts. *Brain Struct Funct* 221:979-995.

Koch MA, Norris DG, Hund-Georgiadis M (2002) An investigation of functional and anatomical connectivity using magnetic resonance imaging. *NeuroImage* 16:241-250.

Krishnan A, Williams LJ, McIntosh AR, Abdi H (2011) Partial Least Squares (PLS) methods for neuroimaging: a tutorial and review. *NeuroImage* 56:455-475.

Malpas CB, Genc S, Saling MM, Velakoulis D, Desmond PM, O'Brien TJ (2016) MRI correlates of general intelligence in neurotypical adults. *J Clin Neurosci* 24:128-134.

Matejko AA, Price GR, Mazzocco MM, Ansari D (2013) Individual differences in left parietal white matter predict math scores on the Preliminary Scholastic Aptitude Test. *NeuroImage* 66:604-610.

McIntosh AR, Lobaugh NJ (2004) Partial least squares analysis of neuroimaging data: applications and advances. *NeuroImage* 23 Suppl 1:S250-263.

Mennes M, Kelly C, Zuo XN, Di Martino A, Biswal BB, Castellanos FX, Milham MP (2010) Inter-individual differences in resting-state functional connectivity predict task-induced BOLD activity. *NeuroImage* 50:1690-1701.

Metzger CD, Eckert U, Steiner J, Sartorius A, Buchmann JE, Stadler J, Tempelmann C, Speck O, Bogerts B, Abler B, Walter M (2010) High field fMRI reveals thalamocortical integration of segregated cognitive and emotional processing in mediodorsal and intralaminar thalamic nuclei. *Front Neuroanat* 4:138.

Moeller K, Willmes K, Klein E (2015) A review on functional and structural brain connectivity in numerical cognition. *Frontiers in human neuroscience* 9:227.

Moeller S, Yacoub E, Olman CA, Auerbach E, Strupp J, Harel N, Ugurbil K (2010) Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial parallel imaging with application to high spatial and temporal whole-brain fMRI. *Magnetic resonance in medicine : official journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine* 63:1144-1153.

Packard MG, Knowlton BJ (2002) Learning and memory functions of the Basal Ganglia. *Annual review of neuroscience* 25:563-593.

Pamplona GS, Santos Neto GS, Rosset SR, Rogers BP, Salmon CE (2015) Analyzing the association between functional connectivity of the brain and intellectual performance. *Frontiers in human neuroscience* 9:61.

Park HJ, Friston K (2013) Structural and functional brain networks: from connections to cognition. *Science* 342:1238411.

Peters SK, Dunlop K, Downar J (2016) Cortico-Striatal-Thalamic Loop Circuits of the Salience Network: A Central Pathway in Psychiatric Disease and Treatment. *Front Syst Neurosci* 10:104.

Pezoulas VC, Zervakis M, Michelogiannis S, Klados MA (2017) Resting-State Functional Connectivity and Network Analysis of Cerebellum with Respect to Crystallized IQ and Gender. *Frontiers in human neuroscience* 11:189.

Ponsoda V, Martinez K, Pineda-Pardo JA, Abad FJ, Olea J, Roman FJ, Barbey AK, Colom R (2017) Structural brain connectivity and cognitive ability differences: A multivariate distance matrix regression analysis. *Human brain mapping* 38:803-816.

Prabhakaran V, Smith JA, Desmond JE, Glover GH, Gabrieli JD (1997) Neural substrates of fluid reasoning: an fMRI study of neocortical activation during performance of the Raven's Progressive Matrices Test. *Cogn Psychol* 33:43-63.

Rosenberg M, Noonan S, DeGutis J, Esterman M (2013) Sustaining visual attention in the face of distraction: a novel gradual-onset continuous performance task. *Atten Percept Psychophys* 75:426-439.

Salami A, Pudas S, Nyberg L (2014) Elevated hippocampal resting-state connectivity underlies deficient neurocognitive function in aging. *Proceedings of the National Academy of Sciences of the United States of America* 111:17654-17659.

Santarnecchi E, Tatti E, Rossi S, Serino V, Rossi A (2015) Intelligence-related differences in the asymmetry of spontaneous cerebral activity. *Human brain mapping* 36:3586-3602.

Seeley WW, Menon V, Schatzberg AF, Keller J, Glover GH, Kenna H, Reiss AL, Greicius MD (2007) Dissociable intrinsic connectivity networks for salience processing and executive control. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 27:2349-2356.

Skudlarski P, Jagannathan K, Calhoun VD, Hampson M, Skudlarska BA, Pearlson G (2008) Measuring brain connectivity: diffusion tensor imaging validates resting state temporal correlations. *NeuroImage* 43:554-561.

Smith S (2016) Linking cognition to brain connectivity. *Nature neuroscience* 19:7-9.

Smith SM, Nichols TE, Vidaurre D, Winkler AM, Behrens TE, Glasser MF, Ugurbil K, Barch DM, Van Essen DC, Miller KL (2015) A positive-negative mode of population covariation links brain connectivity, demographics and behavior. *Nature neuroscience* 18:1565-1567.

Smith SM et al. (2013) Resting-state fMRI in the Human Connectome Project. *NeuroImage* 80:144-168.

Song M, Liu Y, Zhou Y, Wang K, Yu C, Jiang T (2009) Default network and intelligence difference. *Conf Proc IEEE Eng Med Biol Soc* 2009:2212-2215.

Song M, Zhou Y, Li J, Liu Y, Tian L, Yu C, Jiang T (2008) Brain spontaneous functional connectivity and intelligence. *NeuroImage* 41:1168-1176.

Sotiropoulos SN, Jbabdi S, Xu J, Andersson JL, Moeller S, Auerbach EJ, Glasser MF, Hernandez M, Sapiro G, Jenkinson M, Feinberg DA, Yacoub E, Lenglet C, Van Essen DC, Ugurbil K, Behrens TE, Consortium WU-MH (2013) Advances in diffusion MRI acquisition and processing in the Human Connectome Project. *NeuroImage* 80:125-143.

Stevens WD, Spreng RN (2014) Resting-state functional connectivity MRI reveals active processes central to cognition. *Wiley Interdiscip Rev Cogn Sci* 5:233-245.

Tu PC, Hsieh JC, Li CT, Bai YM, Su TP (2012) Cortico-striatal disconnection within the cingulo-opercular network in schizophrenia revealed by intrinsic functional connectivity analysis: a resting fMRI study. *NeuroImage* 59:238-247.

Ugurbil K et al. (2013) Pushing spatial and temporal resolution for functional and diffusion MRI in the Human Connectome Project. *NeuroImage* 80:80-104.

Vaidya CJ, Gordon EM (2013) Phenotypic variability in resting-state functional connectivity: current status. *Brain connectivity* 3:99-120.

Van Essen DC, Smith SM, Barch DM, Behrens TE, Yacoub E, Ugurbil K, Consortium WU-MH (2013) The WU-Minn Human Connectome Project: an overview. *NeuroImage* 80:62-79.

Wendelken C, Nakhabenko D, Donohue SE, Carter CS, Bunge SA (2008) "Brain is to thought as stomach is to ??": investigating the role of rostral-lateral prefrontal cortex in relational reasoning. *Journal of cognitive neuroscience* 20:682-693.

Willmes K, Moeller K, Klein E (2014) Where numbers meet words: a common ventral network for semantic classification. *Scand J Psychol* 55:202-211.

Zimmermann J, Griffiths J, Schirner M, Ritter P, McIntosh AR (2018) Subject-specificity of the correlation between large-scale structural and functional connectivity. *bioRxiv*.

Zimmermann J, Ritter P, Shen K, Rothmeier S, Schirner M, McIntosh AR (2016) Structural architecture supports functional organization in the human aging brain at a regionwise and network level. *Human brain mapping* 37:2645-2661.

## Acknowledgments

Data were provided by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University. The authors acknowledge the support of the NSERC grant (RGPIN-2017-06793). The authors declare no competing financial interests.