

The SERM/SERD Bazedoxifene Disrupts ESR1 Helix 12 to Overcome Acquired Hormone Resistance in Breast Cancer Cells

Sean W. Fanning¹, Rinath Jeselsohn^{2,3}, Venkatasubramanian Dharmarajan⁴, Christopher G. Mayne^{5,6}, Mostafa Karimi⁷, Gilles Buchwalter², Rene Houtman⁸, Weiyi Toy⁹, Colin E. Fowler¹, Muriel Lainé¹, Kathryn E Carlson⁵, Teresa A. Martin⁵, Jason Nowak⁴, Jerome Nwachukwu⁴, David J. Hosfield¹, Sarat Chandralapathy⁹, Emad Tajkhorshid⁶, Kendall W. Nettles⁴, Patrick R. Griffin⁴, Yang Shen⁷, John A. Katzenellenbogen⁵, Myles Brown^{2,3}, Geoffrey L. Greene¹

1: Ben May Department for Cancer Research, University of Chicago, Chicago, IL 60637 USA
2: Center for Functional Cancer Epigenetics, Dana-Farber Cancer Institute, Boston, MA 02215 USA
3: Department of Medical Oncology, Dana Farber Cancer Institute, Boston, MA 02215 USA
4: Department of Molecular Medicine, The Scripps Research Institute, Jupiter, FL 33458, USA
5: Department of Chemistry, University of Illinois at Urbana-Champaign, Urbana, IL 61801 USA
6: Department of Biochemistry, College of Medicine, Center for Biophysics and Computational Biology, and Beckman Institute for Advanced Science and Technology, University of Illinois at Urbana-Champaign, Urbana, IL 61801 USA
7: Department of Electrical and Computer Engineering, TEES-AgriLife Center for Bioinformatics and Genomic Systems Engineering, Texas A&M University, College Station, TX 77843 USA
8: PamGene International B.V, 's-Hertogenbosch, Netherlands 5211
9: Human Oncology and Pathogenesis Program, Memorial Sloan-Kettering Cancer Center (MSKCC), New York, NY 10065, USA

Running Title: Bazedoxifene inhibits ER α with activating mutations.

Precis: Bazedoxifene's SERD activities enable it to resist the impact of activating ESR1 mutations.

Keywords (5): ESR1 Somatic Mutations, Breast Cancer

Significance: Bazedoxifene (BZA) is a potent orally available antiestrogen that is clinically approved for use in hormone replacement therapy (DUAVEE). We explore the efficacy of BZA to inhibit activating somatic mutants of ER α that can arise in metastatic breast cancers after prolonged exposure to aromatase inhibitors or tamoxifen therapy. Breast cancer cell line, biophysical, and structural data show that BZA disrupts helix 12 of the ER α ligand binding domain to achieve improved potency against Y537S and D538G somatic mutants compared to 4-hydroxytamoxifen.

Financial Support:

SWF: Susan G. Komen Foundation PDF14301382
SWF, BDG, CEF, ML, GLG, WT, SC: Department of Defense Breakthrough Award W81XWH-14-1-0360
JAK: NIH DK015556, Breast Cancer Research Foundation BCRF-17-083
SC: NCI Cancer Center Support Grant (CCSG, P30 CA08748) and NIH R01CA204999
YS: NIH R35GM124952 and NSF CCF-1546278

Conflicts of Interest: Giles Buchwalter is an employee and shareholder at Celgene. Rene Houtman is an employee at PamGene International.

Abstract

Acquired resistance to endocrine therapy remains a significant clinical burden for breast cancer patients. Somatic mutations in the *ESR1* (estrogen receptor alpha (ER α) gene ligand-binding domain (LBD) represent a recognized mechanism of acquired resistance. Antiestrogens with improved efficacy versus tamoxifen might overcome the resistant phenotype in ER+ breast cancers. Bazedoxifene (BZA) is a potent antiestrogen that is clinically approved for use in hormone replacement therapies. We find BZA possesses improved inhibitory potency against the Y537S and D538G ER α mutants compared to tamoxifen and has additional inhibitory activity in combination with the CDK4/6 inhibitor palbociclib. In addition, comprehensive biophysical and structural biology studies show that BZA's selective estrogen receptor degrading (SERD) properties that override the stabilizing effects of the Y537S and D538G ER α mutations.

Introduction

Estrogen receptor alpha (ER α) plays critical roles in the etiology, treatment and prevention of the majority of breast cancers [1]. Due to the high degree of efficacy and wide therapeutic indices of endocrine therapies, patients may receive such treatments for progressive disease over the course of several years [2]. Unfortunately, the majority of ER+ metastatic breast cancers that initially respond to endocrine treatment will become refractory despite continued ER α expression [2]. Selective estrogen receptor modulators (SERMs) like tamoxifen are antagonistic in the breast and agonistic in the bone and endometrium. SERM agonist activity stems from tissue-specific co-regulator binding in the presence of tamoxifen [3]. In addition, somatic mutations to *ESR1* (gene for ER α) ligand binding domain (LBD) were identified in 25-30% of patients who previously received endocrine treatment [2, 4-6]. Y537S and D538G are the two most prevalent mutations, and pre-clinical studies show that these mutations confer hormone-free transcriptional activity and relative resistance to tamoxifen and fulvestrant treatment [2, 4-6]. Both mutants enable constitutive ER α activity by favoring the agonist-like conformation of the receptor activating function-2 (AF-2) surface and significantly reduce hormone and 4-hydroxytamoxifen (the active metabolite of tamoxifen) binding affinities [7, 8].

Endocrine treatments with improved efficacy could potentially overcome resistance engendered by the activating somatic mutants and other mechanisms. In pre-clinical studies, fulvestrant (FULV, a selective estrogen receptor degrader (SERD) and complete antiestrogen) at high concentrations was the only molecule that reduced the Y537S and D538G ER α mutant transcriptional activity to basal levels [2, 4-6]. However, its clinical efficacy is limited by poor





solubility and oral bioavailability [9-11]. Consequently, new complete antiestrogens with improved oral bioavailability and pharmacokinetics, including AZD9496, GDC-0927, and RAD1901, are currently in development, although their suitability as long-term endocrine therapies remains to be determined [10, 12-14].

Here, we explore whether bazedoxifene (BZA), a potent antiestrogen that retains some SERM properties, shows activity against breast cancer cells that express ESR1 somatic mutants. We chose BZA because it has been extensively studied in clinical trials and is approved for the use in combination with conjugated equine estrogens for hormone replacement therapy in postmenopausal women (DUAVEE, Pfizer) in the US and for the prevention of osteoporosis as a single agent in Europe [9, 15]. Importantly, it displayed strong antagonist and SERD profiles in the breast while retaining beneficial agonist properties in the bone and did not stimulate endometrial tissue in pre-clinical studies [9, 16, 17]. Further, BZA showed good oral bioavailability and improved pharmacokinetics compared with fulvestrant (FULV) [9, 18].

In this study, breast cancer reporter gene assays reveal the inhibitory capacity of BZA against the ER α mutants compared to the SERM 4-hydroxytamoxifen (4-OHT) and SERD FULV in several ER $^{+}$ breast cancer cell lines (MCF-7, ZR75, T47D). We further assessed the ability of BZA to induce the degradation of WT, Y537S, and D538G somatic mutant ER α in MCF7 cells. Additionally, because inhibitors of CDK4/6 combined with antiestrogens are approved for first line therapy and beyond in metastatic ER $^{+}$ breast cancers [19-21], we examined whether the CDK4/6 inhibitor, palbociclib, can be used in combination with BZA to enhance the inhibition of breast cancer cell proliferation. Importantly, comprehensive structural and biophysical studies provide additional molecular insights into the chemical differences between BZA, 4-OHT, and raloxifene (RAL, another SERM) that appear to underlie the SERD properties of BZA and its

106 improved inhibitory efficacy against the Y537S and D538G mutants in breast cancer cells. **Table**
 107 **1** shows the chemical structures of the molecules examined in this study and summarizes their
 108 clinical indications.

109 **Table 1:** Competitive inhibitors of estrogen receptor alpha.

Antiestrogen	Class	Approved Clinical Indications
 4-Hydroxytamoxifen (4-OHT)	SERM	• Adjuvant treatment for ER+ breast cancers [22]. • Metastatic Breast Cancer [23]. • Ductal Carcinoma in Situ [24]. • Reduction in Breast Cancer Incidence in High Risk Women [25].
 Raloxifene (RAL)	SERM	• Osteoporosis in postmenopausal women [26]. • Reduction in Breast Cancer Incidence in High Risk Women[27].
 Fulvestrant (FULV)	SERD	• First-line therapy for metastatic breast cancer [28]. • Postmenopausal women with progressive breast cancer following other antiestrogen therapy [29, 30].
 Bazedoxifene (BZA)	SERM/SERD	• In combination with conjugated equine estrogens (DUAVEE) to prevent postmenopausal osteoporosis [15].

Results

Bazedoxifene Displays SERD Activity in MCF-7 Cells that Express WT ER α

To assess the ability of BZA to inhibit WT ER α in breast cancer cells, we examined its impact on ER α transcriptional activity, degradation and cell growth in MCF-7 cells. 4-hydroxytamoxifen (4-OHT) was used as a representative SERM and FULV was used as a representative SERD (**Figure 1**). In MCF7 cells that expressed an ERE-luciferase reporter gene, BZA was a more potent inhibitor of WT ER α transcription than either 4-OHT or FULV (inhibition of luciferase IC₅₀ for BZA = 0.12 nM, 4-OHT = 0.39 nM and FULV = 0.76 nM) (**Figure 1A**). To test the effect of BZA on endogenous WT ER α transcriptional activity, qPCR was used to quantify the relative mRNA levels of known ER target genes, including cyclin D1, c-myc, CA12, and GREB1, in MCF-7 cells treated with estradiol (E2) or with E2 in combination with BZA, 4-OHT or FULV at 10⁻⁸ and 10⁻⁶ M (antagonistic mode). For cyclin D1, 4-OHT increased the mRNA level at 10⁻⁸ M and showed little effect at 10⁻⁶ M, while both FULV and BZA decreased mRNA levels at 10⁻⁶ M (**Figure 1B**). The agonist activity of tamoxifen at low concentrations has been described previously [31]. BZA increased c-myc mRNA levels at 10⁻⁸ M while it significantly decreased c-myc mRNA at 10⁻⁶ M (**Figure 1C**). Presumably this effect is similar to the behavior of low-level tamoxifen stimulation and merits further examination. Interestingly, 10⁻⁶ M BZA showed the greatest reduction in mRNA levels of both CA12 and GREB compared to 4-OHT and FULV (**Figure 1, D and E**).

As BZA was shown to behave as a SERM/SERD in previous studies [9, 17], we next tested the activity of BZA as an inducer of ER α degradation and observed dose-dependent ER

degradation in MCF7 cells. Overall, BZA mediated similar levels of ER α degradation compared to FULV (**Figure 1F**). In terms of cell growth inhibition, BZA showed an improved IC₅₀ compared to 4-OHT and in the same range as fulvestrant (BZA IC₅₀ = 2.4x10⁻¹⁰ M, FULV IC₅₀ = 3.1x10⁻¹⁰ M and 4-OHT IC₅₀ = 1.19x10⁻⁹ M (**Figure 1G**). Together, these data indicate that BZA degrades WT ER α in breast cancer cells and is more effective at inhibiting ER transcription and cell growth than 4-OHT and FULV.




Figure 1: The inhibitory potency of BZA in MCF-7 cells. **A)** ER α transcriptional reporter gene assay in cells treated with BZA, FULV, and 4-OHT. **B)** Relative cyclin D1 mRNA levels of in MCF-7 cells treated with E2 plus 4-OHT, FULV, or BZA vehicle and normalized to E2. **C)** Relative c-myc mRNA levelsMCF-7 cells treated with E2 plus 4-OHT, FULV, or BZA vehicle and normalized to E2. **D)**

Relative GREB1 mRNA levels in MCF-7 cells treated with E2 plus 4-OHT, FULV, or BZA vehicle and normalized to E2. **E)** Relative CA12 mRNA levels of MCF-7 cells treated with E2 plus 4-OHT, FULV, or BZA vehicle and normalized to E2. **F)** ER α degradation in MCF-7 cells with increasing doses of BZA, or FULV normalized to β -actin. **G)** Inhibition of cell growth with increasing concentrations of BZA, FULV, or 4-OHT.

BZA is a Potent Inhibitor of Activating Somatic Mutants of ER α in Breast Cancer Cells

We next tested the activity of BZA in MCF7 cells that ectopically expressed the Y537S mutant ER α to determine the inhibition of Y537S mutant cell growth. BZA demonstrated an increased potency compared to FULV and 4-OHT, with an IC₅₀ of 1×10^{-10} M vs 2×10^{-9} M and 7×10^{-9} M, respectively (**Figure 2A**). In addition, qPCR data showed that BZA inhibited the transcription of ER α target genes cyclin D1, c-Myc, and PR, in cells expressing the Y537S mutant, in a dose-dependent manner, confirming the on-target effects of BZA in the presence of the mutation (**Figure 2B**).

To evaluate the ability of BZA to induce WT and mutant ER α degradation in breast cancer cells, we treated MCF-7 cells that ectopically expressed HA-tagged WT, Y537S and D538G ER α with BZA and other ligands for comparison. Levels of WT and mutant ER α were quantified using immunoblots with an anti-HA antibody. Cells were treated with 10 nM E2, or 100 nM 4-OHT, 100 nM BZA, 100 nM RAL, 1 μ M FULV or vehicle for 24 hours before immunostaining; 1 μ M FULV was chosen because it was the minimal concentration necessary to achieve maximal ER α degradation. All data were normalized to vehicle-treated cells. In cells expressing HA-WT ER α , BZA and FULV induced degradation of the receptor to similar levels while the amount of the receptor increased upon 4-OHT treatment and was slightly reduced with E2 and RAL (**Figure 2C and D**). Interestingly, for the Y537S mutant, ER α expression remained unchanged for E2 and FULV, while it increased for 4-OHT and RAL. Y537S ER α also increased with BZA but less so than RAL or 4-OHT. Surprisingly, BZA degraded the D538G

ER α mutant to a greater extent than FULV while 4-OHT and RAL both increased its expression after 24 hours. It should be noted that BZA and FULV elicited consistent WT and mutant ER α degradation across all replicates (**Supplemental Figure 1**). However, 4-OHT and RAL elicited slight variations in the actin-normalized quantity of ER α after 24-hours. Overall, these data suggest that BZA degrades WT and D538G ER α in MCF7 cells, but that the Y537S mutant is resistant to degradation. However, the levels of Y537S ER α in the BZA treated cells were still reduced compared to 4-OHT and RAL treatment, consistent with the reduced activities demonstrated by these compounds in MCF-7 reporter gene assays [2].




Figure 2: The ability of BZA to disrupt Y537S and D538G ER α activity. **A)** Cell growth in MCF-7 cells with DOX-induced Y537S ER α expression. **B)** Inhibition of ER α target genes in DOX-induced Y537S ER α expressed MCF-7 cells with increasing doses of BZA. **C)** Representative immunoblot of HA-ER α WT, 537S, or D538G treated with E2, 4-OHT, BZA, RAL, or FULV for 24 hours. **D)** Representative counts of HA-ER α from the immunoblot normalized to actin.

Dual Treatment with BZA and Palbociclib

CDK4/6 inhibitors have emerged as potent agents in the treatment of metastatic ER+ breast cancer in combination with endocrine treatment. Combined endocrine treatment with a CDK4/6 inhibitor is now the standard of care in either first or second line treatment of metastatic ER+ breast cancer [19-21]. Because BZA showed increased activity over FULV and 4-OHT, we explored whether the activity of BZA combined with the CDK4/6 inhibitor, palbociclib (PB), in multiple ER positive cell lines (MCF-7, ZR75, T47D) and long-term, estrogen-deprived (LTED) ER+ MCF7 cells that mimic resistance to aromatase inhibitors. For the first three cell lines, the combination of BZA and PB demonstrated the greatest arrest in cellular proliferation, whereas for the LTED cells it was comparable to PB+FULV (**Figure 3A-D**). Additionally, reduced proliferation of MCF-7 cells expressing the Y537S mutant was observed for the BZA+PB treatment compared to all other treatments (**Figure 3E and F**). Transcriptional reporter gene assays in MCF7 cells showed that: 1. BZA had superior activity in the inhibition of ER transcriptional activity compared to fulvestrant. In addition, palbociclib does not affect ER transcriptional activity either as a single agent or in combination with BZA (**Figure 3G and 3H**). Similarly, immunoblotting for ER showed that treatment with PB does not affect BZA or FULV induced degradation of ER α (**Figure 3I**). In sum, these data show that dual inhibition of CDK4/6 with PB and ER α with BZA is an effective combination with significant activity against breast cancer cells expressing WT or constitutively active mutant ER α .




Figure 3: Combination treatment with CDK4/6 inhibitor and BZA. **A)** Cell growth inhibit with MCF7 breast cancer cells. **B)** ZR75 breast cancer cells. **C)** T47D breast cancer cells. **D)** LTED breast cancer cells. **E)** Non-induced MCF7 breast cancer cells with a dox-inducible ERα Y537S mutant. **F)** MCF7 cells expressing ERα Y537S. **G)** Dose-response curves for inhibition of ERα transcriptional activity in the presence of BZA, PALBO, and FULV in MCF7 cells. **H)** ERα transcriptional reporter gene assays for combination treatments. **I)** ERα stability resulting from combination treatments.

Coregulator Binding Specificity and Affinities of WT and Mutant ERα with BZA, 4-OHT, and FULV

Because hormone regulated coactivator recruitment is crucial for ERα genomic action and inhibition of coactivator recruitment is a key aspect of SERM-mediated ERα antagonism [32], we tested the effects of 4-OHT, BZA and FULV on co-regulator binding. We applied the Microarray Assay for Real-time Coregulator-Nuclear receptor Interaction (MARCoNI), which

allows the quantification of binding affinity of a nuclear receptor with co-regulator peptides. To determine the effect of 4-OHT, BZA and FULV on co-regulator binding to WT, Y537S and D538G ER α , MCF7 cells that ectopically express HA-tagged WT, Y537S, or D538G ER α were used in conjunction with an HA antibody to detect ER binding to the co-regulator array. Experiments were performed under E2 stimulated conditions for WT ER and under apo conditions for mutant ER α . Overall, dose-dependent inhibition of co-regulator binding was observed for the majority of co-regulator peptides with the three drugs (**Supplemental Figure 2A**). A comparison of EC₅₀ levels for inhibition of co-regulator binding of the three ER antagonists showed that EC₅₀ levels for FULV in both the WT and mutant cells was higher, as expected given the mechanism of action of FULV, compared to SERMs. The 4-OHT and BZA EC₅₀s were higher in the presence of the Y537S and D538G mutations. Collectively, these results show that differences in antagonistic activity manifested by the three endocrine treatments are reflected by changes in co-regulator binding. There are significant differences among these drugs in their antagonistic activity on WT-ER and mutant ER.

SERMs and SERDs Abolish Hormone-Independent ER α -Coactivator Binding *In Vitro* and Reverse Hormone Recruitment of Coactivators

To further dissect the molecular basis for the reduced BZA, 4-OHT, and RAL potency/efficacy observed with mutant ERs, biochemical coactivator recruitment and competitive ligand-binding experiments were performed. As described previously [8], Förster resonance energy transfer (FRET) assays were used to evaluate the interaction of wild type and Y537S and D538G mutant ER α with steroid receptor coactivator 3 (SRC3), a key coregulator in breast cancer cells. The nuclear receptor recognition domain (NRD) of SRC3 and LBD of the ERs, were used in these experiments.

Previously, we showed that in the absence of hormone, SRC3 did not bind to WT ER α LBD, whereas Y537S ER α bound SRC3 markedly in the absence of E2 with a 10-fold reduced affinity, and D538G ER α bound SRC3 with a 100-fold reduced affinity compared to hormone bound WT receptor [8]. To better ascertain the potency of ligands to inhibit coactivator recruitment, we titrated the ligands into fixed concentrations of LBD and SRC3 and monitored LBD-SRC3 interaction by a FRET assay; the three samples were primed with E2 to get a measurable signal from WT and D538G ER. 4-OHT, BZA and RAL reversed the binding of SRC3 NRD to the two mutant ERs and WT-ER α with similar potencies (**Supplemental Figure 2B**). Together, these data show that BZA, 4-OHT, and RAL inhibit both the basal and E2-stimulated recruitment of SRC3 coregulator by the WT and mutant ER α *in vitro*.

BZA, RAL and FULV Elicit Similar Reduced Binding Affinities for Y537S and D538G Compared to WT ER α LBD

To examine what role alterations in binding affinity may play in this reduced potency, competitive [3 H]-E2 ligand binding assays ligand-binding experiments were used to examine the effect of Y537S and D538G mutations on BZA, RAL, and FULV ER binding affinities *in vitro*. We previously showed that the E2 and 4-OHT binding was significantly reduced for both the Y537S and D538G mutants [8]. Affinities were reported using K_i values calculated from IC_{50} values using the Cheng-Prusoff equation (**Supplemental Table 2**) [33]. The affinities of RAL, BZA and FULV for the ER α mutants were reduced 9 to 27-fold relative to WT-ER α . It should be noted that the binding affinity of 4-OHT remained the highest compared to RAL, BZA and FULV against the mutant LBDs. The binding affinities of all tested antiestrogens were somewhat more reduced in D538G than in Y537S. However, FULV and BZA demonstrated the highest potencies in the transcriptional reporter gene assays, even though they exhibited reduced

affinities compared to 4-OHT. Therefore, our data indicate that other factors beyond the reduced binding affinity of mutant LBDs for SERM or SERD must play a role in their decreased potency.

The SERD Properties of BZA Arise from its Disruption of Helix 12

X-ray crystallography was used to reveal the structural details of BZA's antiestrogen properties. An x-ray crystal structure for the WT ER α LBD, in complex with BZA was solved to 2.4 Å with two dimers in the asymmetric unit (ASU) (PDB: 4XI3). The BZA ligand and H12 are well resolved in each monomer in the ASU; poorly resolved residues were not included in the model. **Supplemental Figure 3** shows the observed difference map density for the BZA ligand for chain A. We were unable to obtain diffraction quality crystals with BZA or RAL in complex with either the Y537S or D538G mutant LBDs.

Clear structural differences are apparent compared to the previously published molecular modeling and docking simulations of the ER α LBD-BZA complex [34]. Specifically, the C3 methyl on the indole ring of BZA appears to shift the core of the molecule away from M386, L391 and L428 and towards H12. BZA is most structurally similar to RAL; however, BZA displays more SERD-like behavior [9]. **Figure 4** shows a superposition of the ER α LBD-BZA or RAL x-ray crystal structures. Interestingly, the distal phenol on BZA appears to form a hydrogen bond with improved binding geometry to H524 compared to the distal phenol of RAL. This suggests that the ketone on the RAL linker constrains the phenyl group, sterically precluding the adoption of an ideal hydrogen bonding geometry with H524. The core differences for BZA broadcast down the linker arm to alter its vector towards H12 where the azepan ring now pushes against V534 and P535 (**Figure 4B**). The alterations to V534 and P535 propagate to H12 which appears displaced out of the AF-2 cleft into a less stable orientation.




Figure 4: Structural Basis for the SERD properties exhibited by BZA. A) Overlay of BZA (cyan) with RAL (grey) x-ray crystal structures showing differences in H11-12 loop and H12 orientation. B) Hydrogen bonds (dashed lines) formed by BZA and RAL in the binding pocket and highlighting differences in H11-12 loop and H12 conformation. PDBs: 1ERR and 4XI3.

BZA Binding Conformation is Energetically Favored Compared to RAL

Quantum mechanical calculations were employed to determine whether inherent differences in the BZA and RAL ligands accounted for differences in antagonist potency. A significant energetic shift was observed over the time-course of the scans, revealing that the BZA arm can adopt a set of torsion angles with significantly reduced energetic penalties compared to RAL (**Figure 5A**). Importantly, the energetic minima well is significantly broader for BZA compared to RAL, indicating that the angles adopted by BZA in the ER α ligand-binding site are

more favorable than RAL. Furthermore, an energetic penalty of approximately 1 kcal/mol would be incurred by RAL to adopt the same conformation observed for BZA in the x-ray crystal structure. Together these data show that the BZA ligand itself possesses physical properties that are more favorable to impact ER α H12 compared to RAL.

BZA and FULV Reduce the Impact of Y537S and D538G Mutations on Helix 12 Dynamics

SERDs competitively bind to the ER α LBD and destabilize helix 12 (H12), leading to proteosomal degradation, while SERMs push the helix into the AF2 cleft to block coregulator binding [8]. Furthermore, AZD9496, a newer orally available SERD pushes H12 into the AF2 cleft but destabilizes the helix [13]. Previous work showed that the Y537S and D538G mutants, in complex with 4-OHT, adopts an altered antagonist conformation with respect to the WT-4-OHT complex [8]. Here, we explored how Y537S and D538G ER α LBD mutations impact H12 mobility in the BZA complex using differential hydrogen-deuterium exchange mass spectrometry (HDX-MS). For comparisons we chose E2 as a representative hormone, FULV as a representative SERD, and 4-OHT as a representative SERM. Average time-dependent amide-deuterium uptake kinetics is indicative of conformational flexibility in proteins with highly dynamic regions being more susceptible to solvent deuterium exchange compared to conformationally rigid regions. As expected, addition of E2 resulted in an increased protection against exchange in H12 (inferred as increased stability or less dynamic), and this protection was enhanced for the Y537S and D538G mutants (**Supplemental Figure 4**). Also, FULV treatment led to increased D₂O uptake in H12 (interpreted as destabilization) of regions near H12 in both the WT and D538G mutant (**Supplemental Figure 5**), consistent with its SERD-like properties. Unfortunately, we were unable to collect HDX data for the Y537S mutant with FULV because it precipitated out of solution. Similar to previously published data [8], binding of 4-OHT resulted

in decreased stability of H12 with the Y537S compared to D538G and WT receptor (Supplemental Figure 6), suggesting that these mutants resist the ability of the SERM to alter their structure. Interestingly, addition of BZA did not increase the stability (lesser protection) of the region near H12 to as great of an extent as 4-OHT in WT and mutant ER α LBDs, suggesting that BZA-bound ER α adopts a less stable antagonist conformation than 4-OHT-bound ER α (Supplemental Figure 7), consistent with the crystal structure. These data suggest that, while their antagonist conformations are altered by Y537S or D538G mutations, the ER α -BZA and FULV complexes maintain potency because they resist the stabilizing impact of the mutations better than 4-OHT.

How Y537S and D538G ER α LBD Mutants Alter the BZA Antagonist Structure

To understand the structural basis for the reduced BZA degradation of Y537S compared to WT ER α , atom-level explicit-solvent molecular dynamics (MD) simulations for the LBD of ER α Y537S-BZA and D538G-BZA were performed using the WT-BZA crystal structure as template. When compared to the WT-4-OHT structure (PDB: 3ERT), the D538G-4-OHT structure displayed a significantly altered H11-12 loop, which leads to a perturbed H12 antagonist state (Figure 5B) [8].

MD simulations show that H12 of both BZA-bound mutants remain close to an antagonist conformation (Figure 5B, D). Furthermore, H12 conformational fluctuations observed in MD simulations are less pronounced in Y537S-BZA than they are in D538G-BZA (Figure 5C), which echoes the aforementioned reduced H12 dynamics observed in HDX-MS data and agrees with our result that Y537S reduces BZA activity more than D538G does. Over the course of the simulations, the Y537S-BZA structure shows a hydrogen bond formation

342 between E380 and S537 (**Figure 5B**), which could contribute to fewer H12 conformational
343 fluctuations in Y537S ER α . It should be noted that this hydrogen bond is not sterically allowed
344 in the WT-BZA crystal structure. This predicted hydrogen bond may stabilize H12 in the AF-2
345 cleft relative to WT. The simulated H11-12 loop conformations for both Y537S-BZA and
346 D538G-BZA resemble WT-BZA or D538G-4-OHT, being closer to the ligand binding site, than
347 they do to WT-4-OHT. The D538G mutation further increases the conformational variance of
348 the H11-12 loop compared to Y537S (**Figure 5C**), which can be explained by the lack of a
349 hydrogen bond in Y537S and the greater backbone conformational freedom allowed by the
350 mutant glycine residue. Additionally, the H11-12 loop of both Y537S and D538G appears
351 further away from the protein core and BZA compared to that of WT (**Figure 5B, D**). We
352 hypothesize that the varied conformations and increased dynamics of the H11-12 loop in both
353 ER α mutants makes it more difficult for BZA to maintain sufficient interactions with the loop to
354 disrupt the ER antagonist conformation. Together, these data show that both mutations produce a
355 stable antagonist conformation, especially at H12, and reduce the SERD-like properties of BZA
356 by lessening its ability to disrupt the H11-12 loop and stabilize H12 in the AF-2 cleft.




Figure 5: A) QM scan of RAL and BZA arm torsion angle energetics. **B)** Overlaid representative calculated structure of Y537S-BZA (cyan) closest to the last 50-ns average from molecular dynamics (MD) simulation and the WT-BZA structure, vertical lines represent the ligand binding conformation in the x-ray crystal structure. Nitrogen atoms of the azepane part of BZA for the last 50-ns MD ensemble are shown in blue spheres near H12. Loop H11-12 is also shown for crystal structures of WT-4-OHT (wheat; PDB: 3ERT) and D538G-4-OHT (yellow; PDB: 4Q50). **C)** Root mean squared fluctuations of ERα LBD residues, including the C-end of H11 (resi. 519-527), the loop H11-12 (resi. 528-535), and H12 (resi. 536-544), in the last 50-ns MD simulations for Y537S-BZA (cyan) and D538G-BZA (magenta). **D)** Overlay of a representative structure of D538G-BZA (magenta) from MD simulations and the same WT-BZA structure (gray) as in B).

Discussion

Somatic mutation to *ESR1* following prolonged estrogen-deprivation therapy represents a newly appreciated mechanism of acquired hormone resistance in metastatic breast cancer. The two most prevalent mutations, Y537S and D538G, give rise to a dysfunctional receptor that escapes hormone regulation and has decreased sensitivity to inhibition by 4-OHT and fulvestrant [8]. Newly characterized orally available pure antiestrogens (*e.g.*, GDC-0927, AZ9496, and RAD1901) are emerging as potentially potent inhibitors of these mutants, but their long-term clinical utility is unknown and their side effect profiles have not been studied in large patient populations [12, 14]. Because BZA is already clinically approved for use in hormone replacement therapy and is a potent ER α antagonist in the breast, an agonist in bone, and neutral in the endometrium with long-term safety data in thousands of patients, we explored its ability to inhibit the Y537S and D538G ER α somatic mutants in breast cancer [9].

We first quantified the ability of BZA to disrupt WT and mutant ER α transcription, induce degradation, and inhibit cell growth in MCF-7 cells. These data show that BZA has increased activity, compared to the SERD FULV and SERM 4-OHT, toward inhibition of cell growth and ER α transcriptional activity. Further, BZA elicited degradation of WT receptor, although not as much as FULV. Importantly, BZA reduced the levels of Y537S ER α relative to RAL and 4-OHT. Together these data suggest that the ability of BZA to degrade ER α confers an inhibitory advantage over 4-OHT in the context of the somatic mutants. Additionally, combined treatment with BZA and the CDK4/6 inhibitor palbociclib resulted in additional inhibitory potency of cellular proliferation in breast cancer cells expressing both WT and Y537S ER α without impacting BZA's action. As such, combination BZA-CDK4/6 treatment shows significant potential in cell line experiments and merits further evaluation.

Comprehensive structural-biochemical investigations were undertaken to understand the basis for the SERD-like properties of BZA and its improved potency against the Y537S and D538G ER α LBD mutants compared to 4-OHT. Biochemical coactivator studies using recombinant ER α LBD demonstrated that BZA could inhibit the both basal and E2-induced recruitment of coregulators *in vitro*, while ligand-binding assays showed that the Y537S and D538G mutants had significantly reduced affinities for BZA, 4-OHT, RAL, and FULV. Interestingly, an HDX MS comparison of BZA, 4-OHT, and FULV suggested that FULV and BZA can both resist the impact of somatic mutation on their antagonist binding conformations compared to 4-OHT. An x-ray crystal structure of the WT-BZA complex revealed that differences in the core of the molecule translated to an altered vector of the linker arm, resulting in contacts with the H11-12 loop and a perturbed H12 antagonist binding mode. This less stable H12 antagonist conformation likely explains the SERM/SERD properties of BZA, wherein it allows H12 to adopt an antagonist conformation (like a SERM) that is somewhat destabilized, although not as destabilized as when FULV is bound.

Atomistic MD simulations were used to examine the molecular basis for the observed decrease in degradation of ER α within breast cancer cells expressing the Y537S and D538G mutants. Both mutants stabilized the antagonist conformation of H12 in the AF-2 cleft, while the Y537S appeared to be the more stable of the two by forming a hydrogen bond between S537 and E380. Interestingly, E380Q has also been found to be a recurrent ESR1 mutation able to confer endocrine resistance. MD simulations also suggest that these mutants have varied conformations and increased dynamics of loop H11-12, which potentially reduces BZA's ability to disrupt ER antagonist conformations. Together these results suggest that BZA retains its SERM antagonist properties within breast cancer cells expressing Y537S and D538G mutant ER α to a greater

extent than 4-OHT but its SERD-like properties are diminished. The reduced potency on the mutants likely derives from the stabilization of the apo receptor in the agonist conformation, which reduces the on rate of ligand binding [8], reflecting that the agonist conformer blocks ligand exchange [35].

Overall, our findings show the molecular basis for the SERD-like activity of BZA and its potential advantage with and without CDK4/6 inhibitor, versus 4-OHT, to inhibit the Y537S and D538G ER α mutants. Importantly, interrogating the structural details of BZA-ER α LBD binding suggests that molecules with improved pharmacological profiles that specifically disrupt the H11-12 loop at H12 will have clear advantages against breast cancer cells expressing WT, Y537S, and D538G ER α . **Figure 6** summarizes how BZA achieves SERM/SERD activity compared to SERM or agonist compounds. In fact, the newer SERM/SERDs and SERDs with improved pharmacologic profiles (*e.g.*, AZ9496) appear to do so using a similar mechanism [13]. Therefore, further preclinical evaluation of BZA with and without a CDK4/6 inhibitor should be conducted in context of breast cancer cells expressing the Y537S and D538G somatic mutant ER α . In addition, the ability of other new SERDs to withstand the impact of Y537S and D538G mutations on their antagonist binding modes should be investigated.




Figure 6: Summary of BZA's SERM/SERD activity compared to SERMs and agonist compounds.

Materials Methods

Breast Cancer Cellular Reporter Gene Assays

Luciferase reporter assay system (Promega) was used to monitor luciferase activity in MCF7 cells with stable expression of ERE luciferase per the manufacturer's recommendations, using a single tube luminometer (BD Monolight 2010). MCF7 cells were plated in 6-well plates and treated with increasing doses of BZA, FULV or 4-OHT (0 / 0.01 / 0.1 / 1 / 10 / 100 / 1000nM) in complete medium for 24 hours. All studies were performed in triplicates, and luciferase results are reported as relative light units (RLU) and normalized with β -Galactosidase activity using Mammalian β -gal assay kit (Thermo Scientific). Mycoplasma was tested every 6 weeks in these cells and no mycoplasma was detected in the MCF-7 cell lines using the MycoAlert Mycoplasma Detection Kit (Lonza). MCF-7 cells were purchased directly from ATCC and the studies were completed within 1 year of purchase. Cells reached a maximum of 30 passages during these studies.

ER α Stability Immunoblots

Tet-ON MCF7 cells lines (HA-ER α WT, HA-ER α Y537S, HA-ER α D538G) were cultured in DMEM supplemented with 10% fetal bovine serum (FBS), 2 mM L-glutamate, 1% penicillin-streptomycin, 100 μ g/mL genitacin, and 100 μ g/mL hygromycin. Before western blotting experiments, 300,000 cells were plated in each well of a 6-well culture dish and cultured for 48 hours in DMEM supplemented with 10% charcoal-stripped FBS (SFBS), 2 mM L-glutamate, 1% penicillin-streptomycin, and 0.2 μ g/mL doxycycline. Compounds were diluted in ethanol or DMSO. Cells were treated with either 10 nM estradiol (Sigma), 1 μ M FULV (Tocris), 100 nM Ral (Sigma), 100 nM BZA (Pfizer), 100 nM 4-OHT (Tocris), or vehicle (ethanol) for 24 hours.

Subsequently, cells were washed twice with 1 mL of ice-cold PBS, harvested via scraping, and pelleted at 4 °C. Cells were resuspended in 50 µL lysis buffer [0.1% CHAPS, 50 mM HEPES (pH 8.0), 120 mM NaCl, 1 mM EDTA, 10 mM Na pyrophosphate, and 10 mM glycerophosphate; supplemented with a protease inhibitor cocktail solution III (CalBiochem)]. Cells were lysed via 3 freeze-thaw cycles. Lysates were then pelleted at 4 °C, and 100 µg of protein was mixed with standard 2x Laemmli buffer. Samples were loaded onto a 10% SDS-polyacrylamide electrophoresis gel, transferred after electrophoresis onto nitrocellulose membrane, and immunoblotted using anti-HA-tag (Cell Signaling C29F4) and anti-actin (Santa Cruz Biotechnology AC-15) antibodies. Membranes were probed using anti-rabbit 800 nm (Rockland) and anti-mouse 680 nm (Rockland) and imaged on the Licor Odyssey. Membrane intensities were normalized to actin control and quantified using ImageStudio. Mycoplasma was tested every 6 weeks in these cells and no mycoplasma was detected in any of the Tet-ON MCF-7 cell lines using the MycoAlert Mycoplasma Detection Kit (Lonza).

Co-Regulator Interaction Profiling

This method has been described previously [36]. Cell lysates of MCF7 cells expressing HA-tagged WT-ER, Y537S mutant ER and D538G mutant ER were quantified by ELISA (Active Motif, USA) to enable equimolar input. An array with a set of immobilized peptides representing coregulator-derived NR-binding motifs was incubated with a reaction mixture of crude lysate, vehicle (2% DMSO) with or without 1 µM 17β-estradiol (E2), increasing concentrations of BZA, 4-OHT or FULV (0.1, 1, 10, 100, 1000 nM) and anti-HIS-Alexa488 (Qiagen, USA). Incubation was performed for 40 minutes at 20 °C, followed by removal of unbound receptor by washing and generation of a TIFF image of each array using a PamStation96 (PamGene International).

Image processing and quantification of ER α binding to each peptide on the array was performed by Bionavigator software (PamGene International).

Coactivator Binding Assays

Protein preparation for TR-FRET. Expression, purification, and site-specific labeling of the ER α -LBD was performed as described previously [37]. Generation of the nuclear receptor interaction domain (NRD) of human SRC3 coactivator has also been described previously [38]. ER α LBD and the SRC3 fragment were labeled with Mal-dPEG4-biotin (Quanta BioDesign, Powell, OH), and 5-iodoacetamido fluorescein (Molecular Probes, Invitrogen, Eugene, OR), respectively.

Coactivator Recruitment with Ligand Titration. To raise the background level of SRC3 NRD binding, the ER α LBD preparation (1 nM ER-LBD, 0.25 nM SaTb, 100 nM SRC3-fluorescein) was primed with 20 nM E2, and then increasing ligand concentrations (from 3×10^{-12} to 3×10^{-7} M) were added. Diffusion-enhanced FRET was determined by a parallel incubation without biotinylated ER-LBD and subtracted as a background signal. The time-resolved Förster resonance energy transfer measurements were performed with a Victor X5 plate reader (Perkin Elmer, Shelton, CT) with an excitation filter at 340/10 nm and emission filters for terbium and fluorescein at 495/20 and 520/25 nm, respectively, with a 100- μ s delay [39-41]. The data, representing 2–3 replicate experiments, each with duplicate points, was analyzed using GraphPad Prism 4, and are expressed as the IC₅₀ in nM.

Ligand Binding Assays

The dissociation constant, (K_d) of estradiol binding to each ER was measured by saturation binding with [³H]17 β -E₂ and Scatchard plot analysis [42], as described previously [8, 43].

Relative binding affinities (RBA) were determined by a competitive radiometric ligand binding assay with 2 nM [³H]E2 as tracer [8]. Incubations were at 0 °C for 18–24 hr. Hydroxyapatite was used to adsorb the receptor-ligand complex, and unbound ligand was washed away. The determination of the RBA values is reproducible in independent experiments with a CV of 0.3, and the values shown represent the average ± range or SD of two or more separate determinations. K_i was determined by the Cheng-Prusoff equation [33] $K_i = IC_{50}/(1 + [tracer\ total/K_d\ of\ tracer])$.

Hydrogen/Deuterium Exchange-Mass Spectrometry (HDX-MS)

Solution-phase amide HDX experiments were carried out with a fully automated system (CTC HTS PAL, LEAP Technologies, Carrboro, NC; housed inside a 4°C cabinet) as described previously [8] with slight modifications.

Peptide Identification. Peptides were identified using tandem MS (MS^2 or MS/MS) experiments performed with a LTQ linear ion trap mass spectrometer (LTQ Orbitrap XL with ETD, ThermoFisher, San Jose, CA) over a 70 min gradient. Product ion spectra were acquired in a data-dependent mode and the five most abundant ions were selected for the product ion analysis per scan event. The MS/MS *.raw data files were converted to *.mgf files and then submitted to MASCOT ver2.3 (Matrix Science, London, UK) for peptide identification. The maximum number of missed cleavage was set at 4 with the mass tolerance for precursor ions +/- 0.6 Da and for fragment ions +/- 8ppm. Oxidation to Methionine was selected for variable modification. Pepsin was used for digestion and no specific enzyme was selected in the MASCOT during the search. Peptides included in the peptide set used for HDX detection had a MASCOT score of 20 or greater. The MS/MS MASCOT search was also performed against a decoy (reverse) sequence

and false positives were ruled out. The MS/MS spectra of all the peptide ions from the MASCOT search were further manually inspected and only the unique charged ions with the highest MASCOT score were used in estimating the sequence coverage and included in HDX peptide set.

HDX-MS analysis. For differential HDX experiments, 5 μ L of a 10 μ M ER α LBD (Apo or in complex with 10-fold excess compound) was diluted to 25 μ L with D₂O-containing HDX buffer (20 mM Tris-HCl pH 8.0, 150 mM NaCl, 5% glycerol, 1 mM DTT) and incubated at 4 °C for 10 s, 30 s, 60 s, 900 s, and 3600 s. Following on-exchange, unwanted forward or backward exchange is minimized, and the protein is denatured by dilution to 50 μ L with 0.1% TFA in 5 M urea with 50mM TCEP (held at 4 °C, pH 2.5). Samples are then passed across an immobilized pepsin column (prepared in house) at 50 μ L min⁻¹ (0.1% TFA, 15 °C), and the resulting peptides are trapped onto a C₈ trap cartridge (Thermo Fisher, Hypersil Gold). Peptides were then gradient eluted (5% CH₃CN to 50% CH₃CN, 0.3% formic acid over 6 min, 4 °C) across a 1mm $\times$ 50 mm C₁₈ analytical column (Hypersil Gold, Thermo Fisher) and electrosprayed directly into a high resolution orbitrap mass spectrometer (LTQ Orbitrap XL with ETD, Thermo Fisher). Percent deuterium exchange values for peptide isotopic envelopes at each time point were calculated and processed using HDX Workbench [44]. Each HDX experiment was carried out in triplicate with a single preparation of each protein-ligand complex. The intensity weighted mean m/z centroid value of each peptide envelope was calculated and subsequently converted into a percentage of deuterium incorporation. This is accomplished by determining the observed averages of the undeuterated and using the conventional formula described elsewhere [45]. Statistical significance for the differential HDX data is determined by an unpaired t-test for each time point, a procedure that is integrated into the HDX Workbench software

[44]. Corrections for back-exchange were made on the basis of an estimated 70% deuterium recovery and accounting for 80% final deuterium concentration in the sample (1:5 dilution in D₂O HDX buffer).

X-ray Crystallographic Analysis of the WT ER α LBD-BZA Complex

The 6 $\times$ His-TEV-tagged ER α -L372S, L536S double mutant LBD was expressed in *E.coli* BL21(DE3) and purified as described [46]. LBD (10 mg/mL) and incubated with 1 mM BZA overnight at 4 °C. LBD-BZA was crystallized using vapor diffusion by hanging drop in 33% PEG 3,350, 100 mM Tris pH 6.6, and 250 mM MgCl₂. Diffraction data were collected at the Canadian Light Source at beamline 08ID-1 at a wavelength of 0.97 Å. Indexing, scaling, and structure refinement were performed as described [8]. **Supplemental Table 1** shows data collection and refinement statistics. Final coordinates were deposited in the Protein Databank with the accession code 4XI3.

Quantum Mechanical Calculations

Torsion scans were performed on the bond connecting the internal substituents to the central core for each ligand. The ligand coordinates were extracted from x-ray crystal structures of BZA (PDB code 4XI3) and RAL (PDB code 2QXS) and all hydrogens were added. Relaxed potential energy surface scans in which the remainder of the structure is geometry optimized at each torsion step were prepared and analyzed using the torsion scan module of the Force Field Toolkit [47] (ffTK) plugin of VMD [48]. Quantum mechanical calculations were performed using Gaussian G09 [49] at the MP2 level of theory with a 6-31G* basis set. Both ligands were scanned using a bidirection technique originating from the crystal structure conformation and scanning outward in the (+) and (-) directions independently. The BZA ligand was scanned in 4-

degree increments while the RAL ligand required a smaller 2-degree step size to avoid discontinuities due to broader conformational changes when taking larger steps.

Molecular Dynamics Simulations

Ligand parameterization. A 3D structure of BZA (without hydrogen atoms) was first built using the computer program GaussView (version 4.1.2; part of the computer program Gaussian 03 [50]). The remaining ligand parameterization was carried out as described [8].

Structure preparation, molecular dynamics, data visualization and analysis. WT-BZA (PDB: 4XI3) was used as a template to construct starting structures of Y537S-BZA and D538G-BZA. Specifically, chains A and C were chosen among the three dimers in 4XI3 for having the least missing residues in the loop H11-12 region, with ions removed and water molecules retained. Side chain atoms of mutation sites (residue 537 and 538, respectively) were also replaced with the mutant residues. Otherwise, structures were prepared, molecular dynamics were calculated, and data were analyzed/visualized as described [8].

Acknowledgements

Special thanks to Dr. Richard Walter and James Gorin for assistance with structure data collection. The Canadian Light Source is supported by the Canada Foundation for Innovation, Natural Sciences and Engineering Research Council of Canada, the University of Saskatchewan, the Government of Saskatchewan, Western Economic Diversification Canada, the National Research Council Canada, and the Canadian Institutes of Health Research. Research reported in this publication was supported by the Susan G. Komen Foundation (S.W.F), Department of Defense Breakthrough Award (G.L.G and S.C.), The Virginia and D.K. Ludwig Fund for Cancer Research (GLG), National Institute of General Medical Sciences of the National Institutes of Health under Award Number R35GM124952 (Y.S.) and by the National Science Foundation under Award Number CCF-1546278 (Y.S.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the National Science Foundation. Portions of MD simulations were conducted with high performance research computing resources provided by Texas A&M University (<http://hprc.tamu.edu/>).

References Cited

1. Frasor, J., et al., *Selective Estrogen Receptor Modulators*. Cancer Research, 2004. **64**(4): p. 1522.
2. Toy, W., et al., *ESR1 ligand-binding domain mutations in hormone-resistant breast cancer*. Nat Genet, 2013. **45**(12): p. 1439-45.
3. Shang, Y. and M. Brown, *Molecular Determinants for the Tissue Specificity of SERMs*. Science, 2002. **295**(5564): p. 2465.
4. Jeselsohn, R., et al., *ESR1 mutations-a mechanism for acquired endocrine resistance in breast cancer*. Nat Rev Clin Oncol, 2015. **12**(10): p. 573-83.
5. Jeselsohn, R., et al., *Emergence of constitutively active estrogen receptor-alpha mutations in pretreated advanced estrogen receptor-positive breast cancer*. Clin Cancer Res, 2014. **20**(7): p. 1757-67.
6. Niu, J., et al., *Incidence and clinical significance of ESR1 mutations in heavily pretreated metastatic breast cancer patients*. Onco Targets Ther, 2015. **8**: p. 3323-8.
7. Nettles, K.W., et al., *NFkappaB selectivity of estrogen receptor ligands revealed by comparative crystallographic analyses*. Nat Chem Biol, 2008. **4**(4): p. 241-7.
8. Fanning, S.W., Mayne, C.G., Dharmarajan, V., Carlson, K.E., Martin, T.A., Novick, S.J., Toy, W., Green, B., Panchamukhi, S., Katzenellenbogen, B.S., Tajkhorshid, E., Griffin, P.R., Shen, Y., Chandarlapaty, S., Katzenellenbogen, J.A., Greene, G.L., *Estrogen Receptor Alpha Somatic Mutations Y537S and D538G Confer Breast Cancer Endocrine Resistance by Stabilizing the Activating Function-2 Binding Conformation*. eLife, 2016.
9. Wardell, S.E., et al., *Bazedoxifene exhibits antiestrogenic activity in animal models of tamoxifen-resistant breast cancer: implications for treatment of advanced disease*. Clin Cancer Res, 2013. **19**(9): p. 2420-31.
10. Garner, F., et al., *RAD1901: a novel, orally bioavailable selective estrogen receptor degrader that demonstrates antitumor activity in breast cancer xenograft models*. Anticancer Drugs, 2015. **26**(9): p. 948-56.
11. van Kruchten, M., et al., *Measuring Residual Estrogen Receptor Availability during Fulvestrant Therapy in Patients with Metastatic Breast Cancer*. Cancer Discovery, 2015. **5**(1): p. 72.
12. Lai, A., et al., *Identification of GDC-0810 (ARN-810), an Orally Bioavailable Selective Estrogen Receptor Degradar (SERD) that Demonstrates Robust Activity in Tamoxifen-Resistant Breast Cancer Xenografts*. J Med Chem, 2015. **58**(12): p. 4888-904.
13. De Savi, C., et al., *Optimization of a Novel Binding Motif to (E)-3-(3,5-Difluoro-4-((1R,3R)-2-(2-fluoro-2-methylpropyl)-3-methyl-2,3,4,9-tetra hydro-1H-pyrido[3,4-b]indol-1-yl)phenyl)acrylic Acid (AZD9496), a Potent and Orally Bioavailable Selective Estrogen Receptor Downregulator and Antagonist*. J Med Chem, 2015. **58**(20): p. 8128-40.
14. Toy, W., et al., *Activating ESR1 Mutations Differentially Affect the Efficacy of ER Antagonists*. Cancer Discov, 2016.
15. Tikoo, D. and M. Gupta, *Duavee: a tissue-selective estrogen complex for menopausal symptoms and prevention of osteoporosis*. International Journal of Basic & Clinical Pharmacology, 2015. **4**(2): p. 391.
16. Komm, B.S., et al., *Bazedoxifene acetate: a selective estrogen receptor modulator with improved selectivity*. Endocrinology, 2005. **146**(9): p. 3999-4008.
17. Lewis-Wambi, J.S., et al., *The Selective Estrogen Receptor Modulator Bazedoxifene Inhibits Hormone-Independent Breast Cancer Cell Growth and Down-Regulates Estrogen Receptor α and Cyclin D1*. Molecular Pharmacology, 2011. **80**(4): p. 610-620.

- 661 18. Biskobing, D.M., *Update on bazedoxifene: a novel selective estrogen receptor modulator*. Clin
662 Interv Aging, 2007. **2**(3): p. 299-303.
- 663 19. Dean, J.L., et al., *Therapeutic CDK4/6 inhibition in breast cancer: key mechanisms of response*
664 *and failure*. Oncogene, 2010. **29**(28): p. 4018-4032.
- 665 20. Wardell, S.E., et al., *Efficacy of SERD/SERM Hybrid-CDK4/6 Inhibitor Combinations in Models of*
666 *Endocrine Therapy-Resistant Breast Cancer*. Clinical Cancer Research, 2015. **21**(22): p. 5121.
- 667 21. Yang, C., et al., *Acquired CDK6 amplification promotes breast cancer resistance to CDK4/6*
668 *inhibitors and loss of ER signaling and dependence*. Oncogene, 2017. **36**(16): p. 2255-2264.
- 669 22. *Tamoxifen for early breast cancer: an overview of the randomised trials*. Early Breast Cancer
670 *Trialists' Collaborative Group*. (0140-6736 (Print)).
- 671 23. Lipton, A., et al., *Randomized Trial of Aminoglutethimide & versus & Tamoxifen in Metastatic Breast Cancer*. Cancer Research, 1982. **42**(8 Supplement): p. 3434s.
- 672 24. Allred, D.C., et al., *Adjuvant tamoxifen reduces subsequent breast cancer in women with*
673 *estrogen receptor-positive ductal carcinoma in situ: a study based on NSABP protocol B-24*.
674 (1527-7755 (Electronic)).
- 675 25. Visvanathan, K., et al., *American society of clinical oncology clinical practice guideline update on*
676 *the use of pharmacologic interventions including tamoxifen, raloxifene, and aromatase inhibition*
677 *for breast cancer risk reduction*. (1527-7755 (Electronic)).
- 678 26. Messalli, E.M. and C. Scaffa, *Long-term safety and efficacy of raloxifene in the prevention and*
679 *treatment of postmenopausal osteoporosis: an update*. International Journal of Women's
680 Health, 2009. **1**: p. 11-20.
- 681 27. Cauley, J.A., et al., *Continued breast cancer risk reduction in postmenopausal women treated*
682 *with raloxifene: 4-year results from the MORE trial*. Multiple outcomes of raloxifene evaluation.
683 (0167-6806 (Print)).
- 684 28. Howell, A., et al., *Comparison of Fulvestrant Versus Tamoxifen for the Treatment of Advanced*
685 *Breast Cancer in Postmenopausal Women Previously Untreated With Endocrine Therapy: A*
686 *Multinational, Double-Blind, Randomized Trial*. Journal of Clinical Oncology, 2004. **22**(9): p.
687 1605-1613.
- 688 29. Osborne, C.K., et al., *Double-blind, randomized trial comparing the efficacy and tolerability of*
689 *fulvestrant versus anastrozole in postmenopausal women with advanced breast cancer*
690 *progressing on prior endocrine therapy: results of a North American trial*. 2002(0732-183X
691 (Print)).
- 692 30. Howell, A., et al., *Fulvestrant, formerly ICI 182,780, is as effective as anastrozole in*
693 *postmenopausal women with advanced breast cancer progressing after prior endocrine*
694 *treatment*. 2002(0732-183X (Print)).
- 695 31. Horwitz, K.B., Y. Koseki, and W.L. McGuire, *Estrogen Control of Progesterone Receptor in Human*
696 *Breast Cancer: Role of Estradiol and Antiestrogen**. Endocrinology, 1978. **103**(5): p. 1742-1751.
- 697 32. Liao, L., et al., *Molecular structure and biological function of the cancer-amplified nuclear*
698 *receptor coactivator SRC-3/AIB1*. The Journal of Steroid Biochemistry and Molecular Biology,
699 2002. **83**(1-5): p. 3-14.
- 700 33. Yung-Chi Cheng, W.H.P., *Relationship between the inhibition constant (K_i) and the concentration*
701 *of inhibitor which causes 5- per cent inhibition (I_{50}) of an enzymatic reaction*. Biochemical
702 Pharmacology, 1973. **22**: p. 3099-3108.
- 703 34. Lewis-Wambi, J.S., et al., *The selective estrogen receptor modulator bazedoxifene inhibits*
704 *hormone-independent breast cancer cell growth and down-regulates estrogen receptor alpha*
705 *and cyclin D1*. Mol Pharmacol, 2011. **80**(4): p. 610-20.
- 706

35. Sonoda, M.T., et al., *Ligand dissociation from estrogen receptor is mediated by receptor dimerization: evidence from molecular dynamics simulations*. Molecular endocrinology (Baltimore, Md.), 2008. **22**(7): p. 1565-1578.
36. Koppen, A., et al., *Nuclear receptor-coregulator interaction profiling identifies TRIP3 as a novel peroxisome proliferator-activated receptor gamma cofactor*. Mol Cell Proteomics, 2009. **8**(10): p. 2212-26.
37. Tamrazi, A., et al., *Estrogen Receptor Dimerization: Ligand Binding Regulates Dimer Affinity and Dimer Dissociation Rate*. Molecular Endocrinology, 2002. **16**(12): p. 2706-2719.
38. Kim, S.H., et al., *A Proteomic Microarray Approach for Exploring Ligand-initiated Nuclear Hormone Receptor Pharmacology, Receptor Selectivity, and Heterodimer Functionality*. Molecular & Cellular Proteomics, 2005. **4**(3): p. 267-277.
39. Tamrazi, A., et al., *Coactivator Proteins as Determinants of Estrogen Receptor Structure and Function: Spectroscopic Evidence for a Novel Coactivator-Stabilized Receptor Conformation*. Molecular Endocrinology, 2005. **19**(6): p. 1516-1528.
40. Moore, T.W., J.R. Gunther, and J.A. Katzenellenbogen, *Estrogen receptor alpha/co-activator interaction assay - TR-FRET*. Methods in molecular biology (Clifton, N.J.), 2015. **1278**: p. 545-553.
41. Jeyakumar, M., et al., *Exploration of Dimensions of Estrogen Potency: PARSING LIGAND BINDING AND COACTIVATOR BINDING AFFINITIES*. The Journal of Biological Chemistry, 2011. **286**(15): p. 12971-12982.
42. Scatchard, G., *THE ATTRACTIONS OF PROTEINS FOR SMALL MOLECULES AND IONS*. Annals of the New York Academy of Sciences, 1949. **51**(4): p. 660-672.
43. Hurth, K.M., et al., *Ligand-Induced Changes in Estrogen Receptor Conformation As Measured by Site-Directed Spin Labeling*. Biochemistry, 2004. **43**(7): p. 1891-1907.
44. Pascal, B.D., et al., *HDX Workbench: Software for the Analysis of H/D Exchange MS Data*. Journal of the American Society for Mass Spectrometry, 2012. **23**(9): p. 10.1007/s13361-012-0419-6.
45. Zhang, Z. and D.L. Smith, *Determination of amide hydrogen exchange by mass spectrometry: a new tool for protein structure elucidation*. Protein Science : A Publication of the Protein Society, 1993. **2**(4): p. 522-531.
46. Sharma, N., et al., *Exploring the Structural Compliancy versus Specificity of the Estrogen Receptor Using Isomeric Three-Dimensional Ligands*. ACS Chemical Biology, 2017. **12**(2): p. 494-503.
47. Mayne, C.G., et al., *Rapid parameterization of small molecules using the Force Field Toolkit*. Journal of computational chemistry, 2013. **34**(32): p. 10.1002/jcc.23422.
48. Humphrey, W., A. Dalke, and K. Schulten, *VMD: Visual molecular dynamics*. Journal of Molecular Graphics, 1996. **14**(1): p. 33-38.
49. Frisch, M.J., et al., *Gaussian 09, Revision E.01*. 2009: Wallingford CT.
50. Frisch, M.J., et al., *Gaussian 03, Revision C.02*.
51. Zhao, Y., et al., *Structurally Novel Antiestrogens Elicit Differential Responses from Constitutively Active Mutant Estrogen Receptors in Breast Cancer Cells and Tumors*. Cancer Research, 2017. **77**(20): p. 5602.

749

751

752

753

754

755

Fanning, S.W.

eLife

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770 **Supplementary Figures**

	ER α LBD-BZA
Data Collection	
Space Group	P1
a, b, c (Å)	53.57, 59.17, 94.14
α , β , γ (°)	86.76, 75.36, 63.03
Resolution Range	50-2.49
Number of	

Fanning, S.W.

eLife

Reflections	
(all/unique)	63,978/29,080
I/ σ (highest resolution)	1.35
R _{merge}	8.0
Completeness (%)	97.6
Redundancy	2.2
Refinement	
Rwork/Rfree	21.1/29.3
No. Residues/Chain	
ER α LBD D538G	241
GRIP Peptide	0
Water	5
Ligand	1
RMSD	
Bond lengths (Å)	0.010
Bond angles (°)	1.575
Chiral volume	0.1016
Ramachandran plot statistics	
Preferred number (%)	837 (97.44%)
Additional allowed (%)	18 (2.10%)
Outliers (%)	4 (0.47%)

771 **Supplemental Table 1:** Crystallographic data collection and refinement statistics.


772

773

774

775

776



Supplemental Figure 1: Replicate experiments of HA-ERα levels in MCF-7 cells upon treatment with E2, 4-OHT, RAL, fulvestrant (FULV), or BZA for 24 hours. A) Replicate immunoblot to that shown in **Figure 1** probing for HA-ERα and actin in MCF-7 cells upon treatment. B) Counts from immunoblot A normalized to actin. C) Third replicate immunoblot to that shown in **Figure 1** probing for HA-ERα and actin in MCF-7 cells upon treatment. D) Counts from immunoblot C normalized to actin.


Supplemental Table 2: Ligand binding affinities for WT, Y537S, and D538G mutant ERα LBD..

Ligand/Mutant	K_d (nM)^a		
	WT	Y537S	D538G
E₂*	0.22 ± 0.11	1.40 ± 0.54	1.77 ± 0.66
	K_i (nM)^b		
4-OHT*	0.12 ± 0.003	2.64 ± 0.4	3.28 ± 0.7
RAL	0.30 ± 0.05	3.59 ± 1.0	3.77 ± 1.0
BZA	0.37 ± 0.01	3.50 ± 0.6	5.53 ± 0.7
Fulvestrant*	0.13 ± 0.03	3.68 ± 0.8	5.06 ± 1.2

^a Measured directly by Scatchard Analysis using [³H]E₂.

^b Calculated using the Cheng-Prusoff equation from the IC₅₀ values determined in a competitive radiometric binding analysis using [³H]E₂ as a tracer.

*Indicates previously published data [8, 51].




Supplemental Figure 2: Coactivator recruitment and inhibition of WT, Y537S, and D538G ERα LBD.

A) EC₅₀ quartiles for cells treated with 4-OHT (red), BZA (blue), or FULV (green). **B)** *In vitro* quantification of the effect of ligands on promoting (E2) or inhibiting (4-OHT, RAL, BZA) the binding of SRC3-NRD to recombinant expressed WT, Y537S, or D538G ERα LBD. To be able to measure a signal from all three receptors, they were first primed with 20 nME2 before adding ligand. IC₅₀ values (nM) are shown next to the legend for each protein.

Fanning, S.W.

eLife



817

818 **Supplemental Figure 3: 2 | Fo-Fc | difference map of BZA in the ER α ligand binding pocket contoured**

819 **to 1.5 σ .**


820




832

Fanning, S.W.

eLife




Supplemental Figure 5: HDX MS for WT and D538G ER α LBD in complex with FULV. Deuterium uptake for each peptide is calculated as the average of % D for the 6 time points (10s, 30s, 60s, 300s, 900s and 3600s) and the difference in average % D values between the Apo-ER α and ER α -FULV bound samples is shown as a heat map with a color code given at the bottom of the figure (warm colors for deprotection and cool colors for protection). Peptides are colored only if they show a >5% difference (less or more protection) in average deuterium uptake between the two states and the software employs a paired two-tailed student's t-test-based coloring scheme (p-value < 0.05 for two consecutive time points or a p-value < 0.01 for a single time point) to distinguish real protection differences from inherent variation in the data. Grey color represents no significant change (0-5%) between the two states.



Supplemental Figure 6: HDX MS for WT, Y537S, and D538G ERα LBD in complex with 4-OHT.

Deuterium uptake for each peptide is calculated as the average of % D for the 6 time points (10s, 30s, 60s, 300s, 900s and 3600s) and the difference in average % D values between the Apo-ERα and ERα-FULV bound samples is shown as a heat map with a color code given at the bottom of the figure (warm colors for deprotection and cool colors for protection). Peptides are colored only if they show a >5% difference (less or more protection) in average deuterium uptake between the two states and the software employs a paired two-tailed student's t-test-based coloring scheme (p-value < 0.05 for two consecutive time points or a p-value < 0.01 for a single time point) to distinguish real protection differences from inherent variation in the data. Grey color represents no significant change (0-5%) between the two states.

Sequence	Charge	Start	End	WT+BZA	Y537S+BZA	D538G+BZA
LDAPPL	1	320	327	-37 (5)	-32 (3)	-26 (5)
DAEPPL	1	321	327	-34 (6)	-31 (4)	-24 (5)
YSEYDPTTRPSE	2	328	339	-2 (3)*	0 (2)*	-2 (5)*
YSEYDPTTRPSEASM	2	328	342	-1 (3)*	0 (2)*	-2 (5)*
MGLLTNL	1	343	349	-40 (5)	-35 (2)	-21 (5)
ADRELVHMINWAKR/PGF	4	350	367	-13 (3)	-14 (2)	-9 (4)
ADRELVHMINWAKR/PGFVD	3	350	369	-17 (6)	-13 (2)	-8 (3)
VDLTLDQVHL	2	368	378	-1 (2)*	-5 (2)*	-6 (3)
VDLTLDQVHLE	2	368	380	-2 (2)*	-4 (1)*	-5 (3)
IGLVWRSMHPGKL	3	389	402	-12 (3)	-8 (1)	-7 (4)
LVWRSMHPGKL	3	391	402	-15 (4)	-10 (2)	-9 (4)
LVWRSMHPGKL	3	391	404	-18 (5)	-15 (7)	-10 (5)
LVWRSMHPGKL	3	391	408	-20 (6)	-14 (3)	-11 (7)
LVWRSMHPGKL	3	391	409	-21 (7)	-14 (3)	-11 (5)
VWRSMHPGKL	3	392	402	-14 (7)	-11 (2)	-9 (4)
VWRSMHPGKL	3	392	410	-21 (7)	-14 (2)	-12 (6)
LFAPNLL	1	403	410	-36 (7)	-24 (2)	-15 (6)
LLDRNQKCVGEM	2	409	421	-12 (3)	-10 (3)	-9 (7)
DRNQKCVGEM	2	411	421	-14 (3)	-12 (3)	-10 (5)
VEIFDL	1	422	428	-14 (3)	-6 (2)	-4 (3)*
LATSSRFMMNLQGEFF	3	429	445	-3 (2)*	-5 (4)	-6 (7)
RMMNLQGEFF	2	436	445	-1 (10)*	-4 (2)*	-9 (7)
SSTLSLEKDHHRVLDKDTOTL	3	463	486	-2 (2)*	-1 (1)*	-3 (4)*
KSLLEKDHHRVLDKDTOTL	3	467	486	-4 (4)*	-3 (4)*	-2 (5)*
IHLMAKAGLTQQQHQRLAQL	3	487	507	-3 (4)*	-2 (1)*	-1 (4)*
IHLMAKAGLTQQQHQRLAQLL	3	487	508	-4 (2)*	-2 (2)*	-3 (4)*
MAKAGLTQQQHQRLAQL	3	490	507	-4 (3)*	-2 (2)*	-4 (4)*
MAKAGLTQQQHQRLAQLL	3	490	508	-4 (3)*	-2 (2)*	-4 (4)*
MAKAGLTQQQHQRLAQLLL	3	490	509	5 (4)*	-2 (1)*	-4 (4)*
LILSHIRHMSNKGMEHL	2	508	525	-12 (7)	-7 (2)	-5 (6)
LILSHIRHMSNKGMEHL	3	509	525	-14 (5)	-7 (2)	-5 (6)
LILSHIRHMSNKGMEHL	3	510	525	-15 (5)	-7 (2)	-7 (6)
YSMKCKNVVPLSDLL	2	526	540	-6 (3)	-1 (3)*	-1 (5)*
KCKNVVPLVGLL	2	529	540	-8 (3)	1 (3)*	-1 (5)*
LEMIDAHLHAPTS	2	541	554	-3 (3)*	0 (2)*	2 (5)*
LEMIDAHLHAPTS	3	541	554	-4 (3)*	1 (3)*	2 (5)*
EMIDAHLHAPTS	2	542	554	0 (3)*	2 (3)*	1 (5)*
LDHLHAPTS	2	544	554	0 (3)*	1 (3)*	1 (5)*
DAHLHAPTS	2	545	554	1 (3)*	1 (3)*	0 (5)*



Supplemental Figure 7: HDX MS for WT, Y537S, and D538G ER α LBD in complex with BZA. Deuterium uptake for each peptide is calculated as the average of % D for the 6 time points (10s, 30s, 60s, 300s, 900s and 3600s) and the difference in average % D values between the Apo-ER α and ER α -FULV bound samples is shown as a heat map with a color code given at the bottom of the figure (warm colors for deprotection and cool colors for protection). Peptides are colored only if they show a >5% difference (less or more protection) in average deuterium uptake between the two states and the software employs a paired two-tailed student's t-test-based coloring scheme (p-value < 0.05 for two consecutive time points or a p-value < 0.01 for a single time point) to distinguish real protection differences from inherent variation in the data. Grey color represents no significant change (0-5%) between the two states.