

Assessing distinct patterns of cognitive aging using tissue-specific brain age prediction based on diffusion tensor imaging and brain morphometry

Geneviève Richard^{a,b,c,*}, Knut Kolskår^{a,b,c}, Anne-Marthe Sanders^{a,b,c}, Tobias Kaufmann^a, Anders Petersen^d, Nhat Trung Doan^a, Jennifer Monereo Sánchez^a, Dag Alnæs^a, Kristine M. Ulrichsen^{a,b,c}, Erlend S. Dørum^{a,b,c}, Ole A. Andreassen^a, Jan Egil Nordvik^c, Lars T. Westlye^{a,b,*}

^a NORMENT, KG Jebsen Centre for Psychosis Research, Division of Mental Health and Addiction, Oslo University Hospital & Institute of Clinical Medicine, University of Oslo, Norway

^b Department of Psychology, University of Oslo, Norway

^c Sunnaas Rehabilitation Hospital HT, Nesodden, Norway

^d Center for Visual Cognition, Department of Psychology, University of Copenhagen, Copenhagen, Denmark

***Corresponding authors:**

Geneviève Richard & Lars T. Westlye

Email: genevieve.richard@medisin.uio.no, l.t.westlye@psykologi.uio.no

Postal address: Oslo University Hospital, P.O.Box 4956 Nydalen, 0424 OSLO, Norway

Telephone: +47 23 02 73 50, Fax: +47 23 02 73 33

Short title:

Tissue-specific brain age prediction

Key words:

Brain age, machine learning, white matter, gray matter, DTI, T1

Disclosures:

The authors declare no competing financial interests.

Abstract

Multimodal imaging enables sensitive measures of the architecture and integrity of the human brain, but the high-dimensional nature of advanced brain imaging features poses inherent challenges for the analyses and interpretations. Multivariate age prediction reduces the dimensionality to one biologically informative summary measure with potential for assessing deviations from normal lifespan trajectories. A number of studies documented remarkably accurate age prediction, but the differential age trajectories and the cognitive sensitivity of distinct brain tissue classes have to a lesser extent been characterized.

Exploring differential brain age models driven by tissue-specific classifiers provides a hitherto unexplored opportunity to disentangle independent sources of heterogeneity in brain biology. We trained machine-learning models to estimate brain age using various combinations of FreeSurfer based morphometry and diffusion tensor imaging based indices of white matter microstructure in 612 healthy controls aged 18-87 years. To compare the tissue-specific brain ages and their cognitive sensitivity we applied each of the 11 models in an independent and cognitively well-characterized sample (n=265, 20-88 years). Correlations between true and estimated age in our test sample were highest for the most comprehensive brain morphometry ($r=0.83$, CI:0.78-0.86) and white matter microstructure ($r=0.79$, CI:0.74-0.83) models, confirming sensitivity and generalizability. The deviance from the chronological age were sensitive to performance on several cognitive tests for various models, including spatial Stroop and symbol coding, indicating poorer performance in individuals with an over-estimated age. Tissue-specific brain age models provide sensitive measures of brain integrity, with implications for the study of a range of brain disorders.

Introduction

Increasing age is a major risk factor for cognitive decline and neurodegeneration, and deviating lifespan trajectories in brain structure and function is a sensitive marker in several common neurological and mental disorders (Cole & Franke 2017). The maturing and aging brain is highly heterogeneous in term of individual trajectories and in term of brain regions and mechanisms involved (Fjell et al. 2013; Westlye et al. 2010b). Understanding the individual determinants and heterogeneity of the developing and aging brain is imperative for identifying persons at risk for various brain disorders, and for developing and applying effective and targeted treatments.

Exploring different modalities acquired by magnetic resonance imaging (MRI) provide a powerful tool to investigate age-related differences in both gray- and white- matter tissue classes across brain regions. However, the richness and complexity of the information provided by advanced imaging pipelines challenges its interpretation. Together, the multifactorial age-related variability and the richness of imaging measures have motivated the development of biologically informative summary measures based on brain imaging data. Using machine-learning to estimate the biological age of the brain based on neuroimaging data is one such approach (Cole & Franke 2017; Cole et al. 2018; Kaufmann et al. 2018). Deviation from the normative trajectory is a highly relevant biomarker for the integrity of the brain in healthy and clinical populations (Marquand et al. 2016; Wolfers et al. in press). Brain age gap is a heritable trait showing regionally specific genetic overlaps with major brain disorders, including schizophrenia and multiple sclerosis (Kaufmann et al. 2018), and accumulating evidence supports increased brain age in several clinical groups, including patients with schizophrenia (Kaufmann et al. 2018; Schnack et al. 2016), Alzheimer's disease (Amoroso et al. 2017; Kaufmann et al. 2018), HIV (Cole et al. 2017b; Kuhn et al. 2018) multiple sclerosis (Kaufmann et al. 2018) and cardiovascular risk factors (Franke et al. 2013; Habes et al. 2016). Indeed, while individuals with brains estimated as younger than their chronological age have been shown to be more physically active (Steffener et al. 2016), augmented brain age has been associated with poor health (Ronan et al. 2016), poor cognitive performance (Liem et al. 2017), early neurodegenerative diseases (Gaser et al. 2013), and increased mortality (Cole et al. 2017a). Less is known about the regional heterogeneity, i.e. to which degree different brain regions, systems or compartments show differential aging patterns and sensitivity to cognitive performance. Brain gray and white matter compartments, which can be assessed and quantified using T1-weighted imaging and diffusion tensor

imaging (DTI), respectively, comprise distinct tissue classes with largely differential biological and environmental modifiers and age trajectories (Bennett et al. 2010; Cao et al. 2017; Fjell et al. 2013; Salat et al. 2005; Storsve et al. 2014; Westlye et al. 2010a; Westlye et al. 2010b). Therefore, allowing for differential brain age models for these distinct classes provides an opportunity to disentangle independent sources of heterogeneity in brain aging.

Thus, to identify common and unique aging patterns with potentially differential sensitivity to cognitive function, we aimed to test the complementary value of tissue-specific prediction by comparing brain age estimated using different combinations of FreeSurfer based morphometric measures (regional cortical thickness, surface area and volume) and white matter microstructure features (DTI based fractional anisotropy and mean, radial and axial diffusivity) across the brain. Based on previous studies on brain aging, we expected high accuracy and generalizability of the age prediction models (Cole & Franke 2017). Since tissue specific brain age models capture biologically distinct information, we anticipated that the different FreeSurfer based brain morphometry and white matter microstructure models would only partly reflect common variance, and therefore provide complementary information with differential sensitivity to cognitive performance. Given that brain age predictions might be sensitive to the overall integrity of the brain (Liem et al. 2017), we anticipated that individuals with an over-estimated brain age would show lower cognitive performance, in particular among the elderly, and that the tissue-specific brain age models would show partly differential cognitive sensitivity.

To ensure generalizability, we trained the models in a large publicly available training set (n=612, 18-87 years) and validated their performance using 10-fold cross-validation before applying to an independent and well characterized test set (n=265, 20-88 years). We assessed the cognitive sensitivity using linear and non-linear models with performance on a range of paper-and-pencil and computerized tests comprising different large-scale cognitive domains (processing speed, executive functioning, working memory, attention, and general intellectual abilities) and cognitive scores based on computational models as dependent variables and age, sex and brain age gap (BAG, estimated brain age minus chronological age) as independent variables. For transparency, we report results both at an uncorrected level and corrected using false discovery rate (FDR) and Bonferroni methods to control the error rate.

Materials and methods

Table 1 summarizes key demographics. We included data from healthy volunteers from two independent cohorts: (1) the Cambridge Centre for Ageing and Neuroscience (Cam-CAN) sample (<http://www.mrc-cbu.cam.ac.uk/datasets/camcan/>; (Shafte et al. 2014; Taylor et al. 2017)) and (2) StrokeMRI, which is an ongoing study on the determinants of stroke recovery, brain health and successful aging (Dorum et al. 2016; Dorum et al. 2017). Figure 1 shows the age distribution for each sample.

Volunteers were recruited to Cam-CAN through a large-scale collaborative research project funded by the Biotechnology and Biological Sciences Research Council (BBSRC, grant number BB/H008217/1), the UK Medical Research Council and University of Cambridge. For more information, see www.cam-can.org. Among the 650 datasets made available, 17 were excluded based on missing or poor quality DTI data and 21 due to poor T1-weighted data quality. Data from the remaining 612 individuals (age 18-87, mean = 54.41, SD = 18.26, 314 females) were included.

Healthy individuals were recruited to StrokeMRI through advertisement in newspapers, social media and word-of-mouth. All participants completed a comprehensive cognitive assessment, multimodal MRI and blood sampling for clinical biochemical analysis, various biomarkers and genotyping. MRI and cognitive assessments were performed on two subsequent days. Exclusion criteria included history of stroke, dementia, or other neurologic and psychiatric diseases, alcohol- and substance abuse, medications significantly affecting the nervous system and counter indications for MRI. In addition, individuals scoring lower than 25 on the Montreal Cognitive Assessment (MoCA; Nasreddine et al. 2005) were assessed for inclusion based on their age, level of education and performance on other cognitive tests. No participants were excluded based on a single low score. A neuroradiologist reviewed all scans and 14 participants with clinically significant abnormalities were excluded. Additional exclusion criteria included missing or incomplete MRI or cognitive data (n=2), or poor quality images (n=20). The remaining 265 participants (age 20-88, mean = 56.95, SD = 14.84, 168 females) were included in further analyses. The study was approved by the Regional Committee for Medical and Health Research Ethics (South-East Norway), and conducted in accordance with the Helsinki declaration. All subjects signed an informed consent prior to participating and received a compensation for their participation.




Fig. 1 Histogram of the age distribution for each sample.

Cognitive assessment in StrokeMRI

Cognitive performance was assessed with a set of neuropsychological and computerized tests assumed to be sensitive to cognitive aging, including the MoCA, the vocabulary and matrix subtests of the Wechsler Abbreviated Scale of Intelligence (WASI; Wechsler 1999), the California Verbal Learning Test (CVLT-II; Delis et al. 2000), and the Delis-Kaplan Executive Function System (D-KEFS) color word interference test (Stroop; Delis et al. 2001). We included several computerized tests from the Cognitive Assessment at Bedside for iPad (CABPad; Willer et al. 2016), including motor speed, verbal fluency (phonological and semantic), working memory, spatial Stroop (executive control of attention), spatial attention span, and symbol digit coding tests. In addition, in order to assess the specificity of cognitive associations using computation modeling, we included three mathematically independent parameters based on the Theory of Visual Attention (TVA; Bundesen 1990; Bundesen & Habekost 2008), including short-term memory storage (K), processing speed (C), perceptual threshold (t_0). These parameters based on computational modeling of response patterns have been shown to be sensitive to age, brain structure and function in healthy individuals (Espeseth et al. 2014; Wiegand et al. 2018) and a range of brain disorders (Habekost 2015; Habekost & Starrfelt 2009). Here, we used a TVA-based modeling of a whole report (Sperling 1960), in which six letters were briefly presented for different exposure durations and the participant's task was to accurately report as many letters as possible. Task error rate was also assessed (i.e. number of incorrect letters out of reported letters).

MRI acquisition

Cam-CAN participants were scanned on a 3T Siemens TIM Trio scanner with a 32-channel head-coil at Medical Research Council (UK) Cognition and Brain Sciences Unit (MRC-

CBSU) in Cambridge, UK. DTI data was acquired using a twice-refocused spin echo sequence with the following parameters a repetition time (TR) of 9100 ms, echo time (TE) of 104 ms, field of view (FOV) of 192 x 192 mm, voxel size: 2 mm³, 66 axial slices using 30 directions with b= 1000 s/mm², 30 directions with b= 2000 s/mm², and 3 b=0 images (Shafto et al. 2014). Only the b=[0,1000] were used in the current analysis. High-resolution 3D T1-weighted data was acquired using a magnetization prepared rapid gradient echo (MPRAGE) sequence with the following parameters: TR: 2250 ms, TE: 2.99 ms, inversion time (TI): 900 ms, flip angle (FA): 9°, FOV of 256 x 240 x 192mm; voxel size =1 mm³ isotropic, GRAPPA acceleration factor of 2, scan time 4:32 minutes (Shafto et al. 2014).

StrokeMRI participants were scanned on a 3T GE 750 Discovery MRI scanner with a 32-channel head coil at Oslo University Hospital. Paddings were used to reduce head motion. DTI data were acquired using an echo planar imaging (EPI) sequence with the following parameters: TR/TE/flip angle: 8150 ms/83.1 ms/90°, FOV: 256 x 256 mm, slice thickness: 2 mm, in-plane resolution: 2 mm, 60 directions (b=1000 s/mm²) and 5 b=0 volumes, scan time: 8:58 min. In addition, 7 b=0 volumes with reversed phase-encoding direction were acquired. High-resolution T1-weighted data was acquired using a 3D IR-prepared FSPGR (BRAVO) with the following parameters: repetition time: 8.16 ms, echo time: 3.18 ms, flip angle: 12°, voxel size: 1 × 1 × 1 mm, field of view: 256 x 256 mm, 188 sagittal slices, scan time: 4:43 minutes.

DTI processing and analysis

Diffusion MRI data from both samples were processed locally using the Oxford Center for Functional Magnetic Resonance Imaging of the Brain (FMRIB) Software Library (FSL) (<http://www.fmrib.ox.ac.uk/fsl>). To correct for geometrical distortions, motion and eddy currents, data were preprocessed using topup (<https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/topup>) and eddy (<https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/eddy>) respectively (Andersson et al. 2003; Smith et al. 2004). Topup uses information from the reversed phase-encoded image, resulting in pairs of images (blip-up, blip-down) with distortions going in opposite directions. From these image pairs the susceptibility-induced off-resonance field was estimated and the two images were combined into a single corrected one (Andersson et al. 2003; Smith et al. 2004). This step was performed on StrokeMRI data only. Eddy detects and replaces slices affected by signal loss due to bulk motion during diffusion encoding, which is performed within an integrated framework along with correction for susceptibility induced distortions, eddy

currents and motion (Andersson & Sotiropoulos 2016). Although these processing steps have been shown to strongly increase the temporal signal-to-noise ratio (tSNR) (Doan et al. 2017), we performed additional visual inspection to identify and remove poor quality data.

Fractional anisotropy (FA), eigenvector, and eigenvalue maps were calculated using dtifit in FSL. Mean diffusivity (MD) was defined as the mean of all three eigenvalues, radial diffusivity (RD) as the mean of the second and third eigenvalue, and axial diffusivity (AD) as the principal eigenvalue.

Voxelwise analysis of FA, MD, AD and RD were carried out using Tract-Based Spatial Statistics (TBSS; Smith et al. 2006), part of FSL (Smith et al. 2004). First, all subjects' FA data were aligned to a common space using the nonlinear registration tool FNIRT (Andersson et al. 2007a; Andersson et al. 2007b). Next, the mean FA image was created and thinned to create a mean FA skeleton, which represents the centers of all tracts common to all participants. Each subject's aligned FA data was then projected onto this skeleton and the resulting data fed into voxelwise cross-subject statistics. The same warping and skeletonization was repeated for MD, AD and RD. We thresholded and binarized the mean FA skeleton at $FA > 0.2$. For each individual, we calculated the mean skeleton FA, MD, AD and RD, as well as mean values within 23 regions of interest (ROIs) based on two probabilistic white matter atlases provided with FSL, i.e. the CBM-DTI-81 white-matter labels atlas and the JHU white-matter tractography atlas (Hua et al. 2008; Mori et al. 2005; Wakana et al. 2007), yielding a total of 96 DTI features per individual.

T1 processing

T1-weighted images were processed using FreeSurfer 5.3 (<http://surfer.nmr.mgh.harvard.edu>; (Dale et al. 1999)) including brain extraction, intensity normalization, automated tissue segmentation, generation of white and pial surfaces (Dale et al. 1999). All reconstructions were visually assessed and corrected as appropriate. Cortical parcellation was performed using the Desikan–Killiany atlas (Desikan et al. 2006; Fischl et al. 2004) and subcortical segmentation was performed based on a probabilistic atlas (Fischl et al. 2002). In addition to global features (intracranial volume, total surface area, whole-cortex mean thickness), mean thickness, total surface area, and volume for each cortical ROI, as well as the volume of subcortical structures were computed yielding a set of 251 FreeSurfer based features.

Age prediction

Eleven different models were trained to estimate age based on the feature sets described above (one based on FreeSurfer T1 features, one based on WM DTI features, one including all T1 and DTI features, in addition to eight models based on a smaller subset of features, including models based on FA, MD, AD, RD, sub-cortical volume, volume, area and thickness to further explore the modality specificity of the estimations).

Due to the systematic differences in the brain features across scanners as well as non-linear effects of age, we fit a generalized additive model (GAM; Hastie 2017) for each feature and regressed out the effects of scanning site and age², accounting all models for age and sex. In addition, we regressed out the estimated total intracranial volume from the area and volume features. Next, for each model, we created a training data matrix by concatenating all the features for all participants in the training sample (Cam-CAN), which were used as input to estimate age. We used the *xgboost* framework in R (<http://xgboost.readthedocs.io/en/latest/R-package/xgboostPresentation.html>), an efficient and scalable implementation of gradient boosting machine learning techniques, to build the prediction models. The following parameters were used: learning rate (η) = 0.1, nround = 5000, gamma = 1, max_depth = 6, subsample=0.5. To estimate the performance of our models, we used a 10-fold cross-validation procedure within the training sample and repeated the cross-validation step 1000 times to provide a robust estimate of model predictive accuracy. Next, we tested the performance of our trained models by predicting age in unseen healthy subjects in the test sample (StrokeMRI).

For each feature set, we calculated the correlation between the predicted and the chronological age as a measure of the model performance, in addition to the mean absolute error (MAE, in years). For each individual, we calculated the discrepancy between the estimated and the chronological age, i.e. the BAG, for each model. The MAE was calculated from the BAG for each model. Since we were interested in the effect of BAG independently of age, the effect of age was regressed out for each BAG using linear models.

Statistical analysis

Statistical analysis was performed using R (<http://www.r-project.org>). For cognitive data, we used *outlierTest* from the *car* package (Fox & Weisberg 2011) to identify the most extreme observations based on a linear model, including age and sex. Twenty-five observations were identified as outliers and treated as missing values based on a Bonferroni corrected $p < 0.05$.

To visualize the associations between the cognitive tests and to form cognitive domain scores based on the correlation patterns, we performed hierarchical clustering using the default setting of the heatmap.2 package in gplots (Warnes et al. 2016), which uses hclust (Müllner 2013) to form clusters based on the complete linkage method. Briefly, this is a step-wise clustering process that merges the two nearest clusters until only one single cluster remains, maximizing distance between individuals components between two clusters.

For each cognitive measure and summary score based on the clusters formed from the clustering step above, we used linear models to test for the effect of age and sex. Since cognitive performance may show non-linear associations with age, we performed an additional analysis including both age and age² in the models. Then, for each test showing a significant association with age, we tested whether adding BAG to the models lead to an improved model fit. More specifically, we tested for differential associations with cognitive function by comparing the parameter estimates for the different BAG models using Fisher z-transformation. To test the assumption that increased BAG is more relevant for cognitive function among the elderly, we tested for age by BAG interactions on cognitive performance. For transparency, we report both uncorrected p-values and p-values adjusted using FDR (Benjamini & Hochberg 1995; Wright 1992) and Bonferroni correction using a factor of 495 (11 brain gaps and 45 cognitive features).

Results

Brain age prediction

10-fold cross-validation on the training sample revealed high correlations between chronological and predicted age for the DTI based white matter microstructure ($r=0.87$) and FreeSurfer based morphometric ($r=0.88$) models. Likewise, the correlations for FA ($r=.76$), MD ($r=.80$), AD ($r=.83$), RD ($r=.78$), sub-volume ($r=.84$), volume ($r=.80$), area ($r=.70$) and thickness ($r=.79$) based models also confirmed reasonable model performance.

Most models accurately predicted age in the independent test set. Figure 2A shows a correlation matrix for the 11 BAGs. Figure 2B shows the correlations between the chronological age and the predicted age in the test sample for each model with their confidence intervals, ranging from ($r=.86$, CI: .82-.89, MAE= 6.14) for the combined model to $r=.58$ (CI: .49-.65, MAE=10.24) for the model based on area. Figure 2 (D to F) show the estimated age from the three models that performed best among the 11 feature sets, i.e. the

combined DTI and T1 feature models ($r=.86$, CI:.82-.89, MAE= 6.14), the 251 FS T1 features ($r=.83$, CI:.78-.86, MAE= 6.76), and the 96 WM DTI features ($r=.79$, CI:.74-.83, MAE=7.28).




Fig. 2 Comparison between the 11 BAG models. (a) Heatmap of the correlation between different BAGs. (b) Correlations between the chronological age and the predicted age in the test sample for each model with their confidence intervals. (c) Mean and standard error of the 45 p-values ($-\log_{10}(p)$) for the cognitive scores and composite scores for each row (i.e. BAGs), with a higher mean representing a stronger global association across tests. (d) Correlation between the chronological age of each subjects and the combined age, (e) the brain morphometry age, and (f) the white matter microstructure age.

Cognitive assessments and associations with BAGs

Table 1 summarizes descriptive statistics and associations with age and sex for each of the 49 cognitive scores, derived features and domain scores. Linear models revealed 45 significant associations with age after correcting for multiple comparisons, with the strongest effect sizes for the symbol coding test, motor speed, spatial Stroop and spatial attention span. Since non-linear models revealed significant associations with age² only with the color word Stroop 3 (inhibition) and its derived scores (See supplementary Table S1), the main models presented here are linear in order to keep the model to its simplest form. Figure 3 shows a correlation matrix across all normalized cognitive scores with the variables sorted according to the

hierarchical clustering. Several variables were highly correlated, and the clustering solution generally suggested seven broad cognitive domains including (Cluster 1) memory and learning (CVLT, attention span, MoCA), (Cluster 2) visual processing speed (TVA processing speed and perceptual threshold), (Cluster 3) verbal skills (phonological and semantic flow), (Cluster 4) attentional control and speed (spatial Stroop), (Cluster 5) executive control and speed (color-word Stroop), (Cluster 6) reasoning and psychomotor speed (matrix, symbol coding and motor speed, short-term memory storage (TVA-parameter K)), and (Cluster 7) working memory.




Fig. 3 Hierarchical clustering of the cognitive features. Each cognitive score was normalized and when required the scores were multiplied by -1 to ensure that positive scores represent good performance. The higher panel shows the dendrogram resulting from the hierarchical clustering of the scores in 7 cognitive domains.

Table 2 shows summary statistics for the associations between cognitive performance and BAG using linear models, including age and sex as covariates. Figure 4 shows a heatmap of the association between cognitive scores and brain age gaps for which the significant associations have been marked with an asterisk. Supplementary Table S1 and Fig. S1 shows




Fig. 5 Scatter plots of the 2 strongest associations between cognitive measures and BAG. The color gradient represents the age where lighter color is assigned to older individuals, and darker color to younger individuals. All associations indicate worse performance with higher brain age gap.

Discussion

Brain aging is highly heterogeneous, and expanding our understanding of the biological determinants of human aging is imperative for reducing the burden of age-related cognitive decline and neurodegenerative disorders. An estimate of an individual's deviation from the expected lifespan trajectory in brain structure and function may provide a sensitive measure of individual brain integrity and health, both in presumably healthy individuals and in patients suffering from various brain disorders.

The biological heterogeneity of the brain strongly suggests that the concept of a single brain age is too simple, and that tissue-specific brain age models may provide increased sensitivity and specificity in relation to cognitive and mental functions. In line with this view, our main findings demonstrate that different combinations of FreeSurfer based brain morphometry and DTI based white matter microstructural indices can be used to accurately predict the age of individuals, but that the shared variance from the different models suggest that they reflect partly non-overlapping processes of brain aging. Further, the results revealed partly differential sensitivity to cognitive performance; with the strongest cumulative associations across cognitive tests for brain age gaps estimated using RD. Even though our data provide no strong evidence of independent associations with cognitive performance in the current sample of healthy individuals, tissue specific age prediction models might better

inform us about the individual determinants and heterogeneity of the aging brain compared to models collapsing several brain compartments by potentially capturing distinct measures of brain aging.

Brain age prediction

For the age prediction models, our results demonstrated that the 11 different combinations of FreeSurfer based morphometric measures (regional cortical thickness, surface area and volume) and white matter microstructure features (diffusion tensor imaging (DTI) based fractional anisotropy and mean, radial and axial diffusivity) across the brain age models accurately predicted the age of an individual with a mean absolute error between 6.14 and 10.23 years. Brain morphometry and white matter microstructure models had a MAE of 6.76 and 7.28 respectively, which correspond with previous publications (Cole et al. 2016; Han et al. 2014; Valizadeh et al. 2017). In general, combining features and modalities increased the performance, and the highest performing model included a combination of both brain morphometry and white matter microstructure (mean absolute error of 6.14 years). Moreover, the correlations between the different brain age gaps suggested a relatively low level of shared variance (mean correlation = 0.51, SD=0.13). Together these findings support the notion that tissue specific brain age models capture biologically distinct information. This is in line with the characteristic lifespan patterns of global linear decreases in gray matter volume and the nonlinear trajectories of total white matter volume and DTI based metrics of white matter microstructure (Cox et al. 2016; Fjell et al. 2013; Ge et al. 2002; Liu et al. 2017; Raz et al. 2010; Westlye et al. 2010b), highlighting that the different compartments carry unique biological information and that combining different modalities lead to a better estimation the age of individuals (Cherubini et al. 2016; Liem et al. 2017; Madan & Kensinger 2018).

Cognitive associations

We performed a comprehensive cognitive assessment of the test sample, confirming previous evidence of substantial age-related differences in cognitive performance across a range of tests and domains. Hierarchical clustering of the cognitive features indicated a characteristic pattern of covariance, largely reflecting broad cognitive domains, including memory and learning, visual processing speed, verbal skills, attentional and executive control, reasoning and psychomotor speed, and working memory. Ninety percent of the included cognitive features showed age-differences, with the largest effect sizes observed for speed-based

measures, such as symbol coding test, which measures mental and visuo-motor speed (Willer et al. 2016). This is in line with the well-established literature on age-related decline in information processing speed in healthy aging (Bennett et al. 2010; Craik & Salthouse 2008; Harada et al. 2013). Importantly, not only tasks measuring reaction time, but also various TVA measures based on computational modeling, such as short-term memory storage (K), processing speed (C), and perceptual threshold (t_0) showed strong associations with age, in line with previous studies (Espeseth et al. 2014; Habekost 2015; Habekost et al. 2013; McAvinue et al. 2012; Wiegand et al. 2018).

Based on the assumption that brain age captures variance related to the integrity of the brain, we anticipated that individuals with an over-estimated age would show lower cognitive performance, and that the tissue-specific brain age models would show partly differential sensitivity. To test these hypotheses, we used linear models to explore the associations between cognitive performance and BAG, with age and sex as covariates, and directly compared the parameter estimates from the different brain age models. We found a significant association between performance on several tests and BAG beyond the age associations, indicating lower performance in individuals with higher BAG. Briefly, one significant association was found for WM DTI, five for combined BAG, two for the sub-volume, one for the RD and one for the MD BAG. The strongest associations were found with the spatial Stroop congruent trials, and number of responses. These findings support that the deviance between the estimated age and the chronological age captures relevant biological information regarding the cognitive performance of an individual. Whereas we found no significantly different associations between brain age models, the association with symbol digit coding test was only seen for WM DTI BAG, while associations with Stroop 3 and 4 were observed only for sub-volume BAG, suggesting some specificity that should be investigated in future studies including larger samples and a broader spectrum of mental health, cognitive and brain phenotypes, both across healthy and clinical samples. We speculate that the contributions of the different modalities in predicting age and the associations with both cognitive performance, but also age-related illnesses vary across the age-span, as it does during maturational age (Brown et al. 2012). Thus, future studies might benefit from investigating modality specific brain-age estimation using specific age range.

Limitations

The present findings do not come without limitations. First, although reducing the dimensionality of complex brain imaging data to a biologically informative brain age is a powerful method to assess deviations from normal lifespan trajectories in brain health, findings from this data reduction method are limited in specificity. Here, we attempted to both reduce the complexity of the information while keeping some modality specificity measured by different MRI parameters. Finding a balance between specificity and precision represents an interesting challenge for future studies. Moreover, causality and individual level trajectories cannot be established based on cross-sectional data. Therefore, future longitudinal studies are needed to inform us about the relevance of the differential trajectories of the tissue-specific brain age prediction with implications for the study of a range of brain disorders. Next, although the age distribution of the test sample is irrelevant to the individual prediction accuracy, the relative overrepresentation of older individuals in the test sample is a limitation when investigating interactions between BAG and age. Thus, although the lack of brain by BAG interactions on cognitive function did not support our hypothesis that increased BAG is more relevant for cognitive function among the elderly, future studies including individuals across a broader range of function are needed to characterize the lifespan dynamics in the associations between brain and behavior. Although we covered a relatively broad spectrum of structural brain features, the link between imaging based indices of brain structure and brain function is elusive, and brain age models including other brain imaging features, including functional measures, might provide a sensitive supplement to the current models. Lastly, whereas the results showed some numerical differences in the cognitive sensitivity of the different combinations of FreeSurfer based morphometry and white matter microstructure models, these differences were not statistically significant, and the hypothesis that tissue specific models provide increased specificity in terms of cognitive associations remains to be further explored in future studies.

In conclusion, we have demonstrated that models based on different combinations of brain morphometry and white matter microstructural indices provide partly differential information about the aging brain, emphasizing the relevance of tissue-specific brain age models in the study of brain and mental function in health and disease.

Acknowledgements

This study was supported by the Norwegian ExtraFoundation for Health and Rehabilitation (2015/FO5146), the Research Council of Norway (249795, 248238), the South-Eastern Norway Regional Health Authority (2014097, 2015044, 2015073), Sunnaas Rehabilitation Hospital, and the Department of Psychology, University of Oslo, and approved by the Regional Committee for Medical and Health Research Ethics (South-East). Volunteers were recruited to StrokeMRI through advertisement in newspapers, social media and word-of-mouth. In addition, volunteers were recruited to Cam-CAN through a large-scale collaborative research project funded by the Biotechnology and Biological Sciences Research Council (BBSRC, grant number BB/H008217/1), the UK Medical Research Council and University of Cambridge.

References

- Amoroso N, Diacono D, Fanizzi A, La Rocca M, Monaco A, Lombardi A, Guaragnella C, Bellotti R, Tangaro S, and Alzheimer's Disease Neuroimaging I. 2017. Deep learning reveals Alzheimer's disease onset in MCI subjects: results from an international challenge. *J Neurosci Methods*. 10.1016/j.jneumeth.2017.12.011
- Andersson JLR, Jenkinson M, and Smith S. 2007a. TR07BP1: Non-linear optimisation. FMRIB Analysis Group Technical Reports: FMRIB Analysis Group.
- Andersson JLR, Jenkinson M, and Smith S. 2007b. TR07JA2: Non-linear registration, aka spatial normalization. FMRIB Analysis Group Technical Reports: FMRIB Analysis Group.
- Andersson JLR, Skare S, and Ashburner J. 2003. How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging. *Neuroimage* 20:870-888. 10.1016/S1053-8119(03)00336-7
- Andersson JLR, and Sotiropoulos SN. 2016. An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging. *Neuroimage* 125:1063-1078. 10.1016/j.neuroimage.2015.10.019
- Benjamini Y, and Hochberg Y. 1995. Controlling the False Discovery Rate - a Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B-Methodological* 57:289-300.
- Bennett IJ, Madden DJ, Vaidya CJ, Howard DV, and Howard JH, Jr. 2010. Age-related differences in multiple measures of white matter integrity: A diffusion tensor imaging study of healthy aging. *Hum Brain Mapp* 31:378-390. 10.1002/hbm.20872
- Brown TT, Kuperman JM, Chung Y, Erhart M, McCabe C, Hagler DJ, Jr., Venkatraman VK, Akshoomoff N, Amaral DG, Bloss CS, Casey BJ, Chang L, Ernst TM, Frazier JA, Gruen JR, Kaufmann WE, Kenet T, Kennedy DN, Murray SS, Sowell ER, Jernigan TL, and Dale AM. 2012. Neuroanatomical assessment of biological maturity. *Curr Biol* 22:1693-1698. 10.1016/j.cub.2012.07.002
- Bundesden C. 1990. A theory of visual attention. *Psychol Rev* 97:523-547.
- Bundesden C, and Habekost T. 2008. Principles of Visual Attention: Linking Mind and Brain. *Oxford University Press*. 10.1093/acprof:oso/9780198570707.001.0001
- Cao B, Mwangi B, Passos IC, Wu MJ, Keser Z, Zunta-Soares GB, Xu DP, Hasan KM, and Soares JC. 2017. Lifespan Gyrification Trajectories of Human Brain in Healthy Individuals and Patients with Major Psychiatric Disorders. *Scientific Reports* 7. ARTN 511

10.1038/s41598-017-00582-1

Cherubini A, Caligiuri ME, Peran P, Sabatini U, Cosentino C, and Amato F. 2016. Importance of Multimodal MRI in Characterizing Brain Tissue and Its Potential Application for Individual Age Prediction. *Ieee Journal of Biomedical and Health Informatics* 20. 10.1109/Jbhi.2016.2559938

Cole JH, and Franke K. 2017. Predicting Age Using Neuroimaging: Innovative Brain Ageing Biomarkers. *Trends Neurosci.* 10.1016/j.tins.2017.10.001

Cole JH, Marioni RE, Harris SE, and Deary IJ. 2018. Brain age and other bodily 'ages': implications for neuropsychiatry. *Mol Psychiatry*. 10.1038/s41380-018-0098-1

Cole JH, Poudel RPK, Tsagkrasoulis D, Caan MWA, Steves C, Spector TD, and Montana G. 2016. Predicting brain age with deep learning from raw imaging data results in a reliable and heritable biomarker. *ARXIV*. 2016arXiv161202572C

Cole JH, Ritchie SJ, Bastin ME, Valdes Hernandez MC, Munoz Maniega S, Royle N, Corley J, Pattie A, Harris SE, Zhang Q, Wray NR, Redmond P, Marioni RE, Starr JM, Cox SR, Wardlaw JM, Sharp DJ, and Deary IJ. 2017a. Brain age predicts mortality. *Mol Psychiatry*. 10.1038/mp.2017.62

Cole JH, Underwood J, Caan MWA, De Francesco D, van Zoest RA, Leech R, Wit FWNM, Portegies P, Geurtsen GJ, Schmand BA, van der Loeff MFS, Franceschi C, Sabin CA, Majoie CBLM, Winston A, Reiss P, Sharp DJ, and Collaboration C. 2017b. Increased brain-predicted aging in treated HIV disease. *Neurology* 88:1349-1357. 10.1212/Wnl.00000000000003790

Cox SR, Ritchie SJ, Tucker-Drob EM, Liewald DC, Hagenaars SP, Davies G, Wardlaw JM, Gale CR, Bastin ME, and Deary IJ. 2016. Ageing and brain white matter structure in 3,513 UK Biobank participants. *Nat Commun* 7:13629. 10.1038/ncomms13629

Craik FIM, and Salthouse TA. 2008. *The handbook of aging and cognition*. New York: Psychology Press.

Dale AM, Fischl B, and Sereno MI. 1999. Cortical surface-based analysis - I. Segmentation and surface reconstruction. *Neuroimage* 9:179-194. DOI 10.1006/nimg.1998.0395

Delis DC, Kaplan E, and Kramer JH. 2001. *Delis-Kaplan Executive Function System: Technical Manual*. San Antonio, TX: Harcourt Assessment Company.

Delis DC, Kramer JH, Kaplan E, and Ober BA. 2000. *California Verbal Learning Test- Second Edition (CVLT-II)*. San Antonio, TX: Psychological Corporation.

Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM, Maguire RP, Hyman BT, Albert MS, and Killiany RJ. 2006. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* 31:968-980. 10.1016/j.neuroimage.2006.01.021

Doan NT, Engvig A, Persson K, Alnaes D, Kaufmann T, Rokicki J, Cordova-Palomera A, Moberget T, Braekhus A, Barca ML, Engedal K, Andreassen OA, Selbaek G, and Westlye LT. 2017. Dissociable diffusion MRI patterns of white matter microstructure and connectivity in Alzheimer's disease spectrum. *Scientific Reports* 7. ARTN 45131 10.1038/srep45131

Dorum ES, Alnaes D, Kaufmann T, Richard G, Lund MJ, Tonnesen S, Sneve MH, Mathiesen NC, Rustan OG, Gjertsen O, Vatn S, Fure B, Andreassen OA, Nordvik JE, and Westlye LT. 2016. Age-related differences in brain network activation and co-activation during multiple object tracking. *Brain and Behavior* 6. 10.1002/brb3.533

Dorum ES, Kaufmann T, Alnaes D, Andreassen OA, Richard G, Kolskar KK, Nordvik JE, and Westlye LT. 2017. Increased sensitivity to age-related differences in brain functional connectivity during continuous multiple object tracking compared to resting-state. *Neuroimage* 148:364-372. 10.1016/j.neuroimage.2017.01.048

Espeseth T, Vangkilde SA, Petersen A, Dyrholm M, and Westlye LT. 2014. TVA-based assessment of attentional capacities-associations with age and indices of brain white matter microstructure. *Front Psychol* 5:1177. 10.3389/fpsyg.2014.01177

Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, van der Kouwe A, Killiany R, Kennedy D, Klaveness S, Montillo A, Makris N, Rosen B, and Dale AM. 2002. Whole brain segmentation: Automated labeling of neuroanatomical structures in the human brain. *Neuron* 33:341-355. Doi 10.1016/S0896-6273(02)00569-X

Fischl B, van der Kouwe A, Destrieux C, Halgren E, Segonne F, Salat DH, Busa E, Seidman LJ, Goldstein J, Kennedy D, Caviness V, Makris N, Rosen B, and Dale AM. 2004. Automatically parcellating the human cerebral cortex. *Cerebral Cortex* 14:11-22. 10.1093/cercor/bhg087

Fjell AM, Westlye LT, Grydeland H, Amlie I, Espeseth T, Reinvang I, Raz N, Holland D, Dale AM, Walhovd KB, and Alzheimer Disease Neuroimaging I. 2013. Critical ages in the life course of the adult brain: nonlinear subcortical aging. *Neurobiol Aging* 34:2239-2247. 10.1016/j.neurobiolaging.2013.04.006

- 609 Fox J, and Weisberg S. 2011. An {R} Companion to Applied Regression. In: Second,
610 editor. Thousand Oaks, CA: Sage.
- 611 Franke K, Gaser C, Manor B, and Novak V. 2013. Advanced BrainAGE in older adults
612 with type 2 diabetes mellitus. *Frontiers in Aging Neuroscience* 5. ARTN 90
613 10.3389/fnagi.2013.00090
- 614 Gaser C, Franke K, Kloppel S, Koutsouleris N, Sauer H, and Initia AsDN. 2013.
615 BrainAGE in Mild Cognitive Impaired Patients: Predicting the Conversion to Alzheimer's
616 Disease. *PLoS One* 8. ARTN e67346
617 10.1371/journal.pone.0067346
- 618 Ge Y, Grossman RI, Babb JS, Rabin ML, Mannon LJ, and Kolson DL. 2002. Age-
619 related total gray matter and white matter changes in normal adult brain. Part I: volumetric
620 MR imaging analysis. *AJNR Am J Neuroradiol* 23:1327-1333.
- 621 Habekost T. 2015. Clinical TVA-based studies: a general review. *Front Psychol* 6:290.
622 10.3389/fpsyg.2015.00290
- 623 Habekost T, and Starrfelt R. 2009. Visual attention capacity: a review of TVA-based
624 patient studies. *Scand J Psychol* 50:23-32. 10.1111/j.1467-9450.2008.00681.x
- 625 Habekost T, Vogel A, Rostrup E, Bundesen C, Kyllingsbaek S, Garde E, Ryberg C, and
626 Waldemar G. 2013. Visual processing speed in old age. *Scand J Psychol* 54:89-94.
627 10.1111/sjop.12008
- 628 Habes M, Guray E, Toledo JB, Zhang T, Bryan RN, Janowitz D, Doshi J, von
629 Sarnowski B, Hegenscheid K, Voelzke H, Schminke U, Hoffmann W, Grabe HJ, and
630 Davatzikos C. 2016. Non-Resilient Brain Aging in Association with Cardiovascular Risk and
631 White Matter Hyperintensities: The Ship Study. *Alzheimer's & Dementia* 12:P226-P227.
632 10.1016/j.jalz.2016.06.407
- 633 Han CE, Peraza LR, Taylor JP, and Kaiser M. 2014. Predicting Age across Human
634 Lifespan Based on Structural Connectivity from Diffusion Tensor Imaging. *2014 Ieee*
635 *Biomedical Circuits and Systems Conference (Biocas)*:137-140.
- 636 Harada CN, Natelson Love MC, and Triebel KL. 2013. Normal cognitive aging. *Clin*
637 *Geriatr Med* 29:737-752. 10.1016/j.cger.2013.07.002
- 638 Hastie T. 2017. gam: Generalized Additive Models.
- 639 Hua K, Zhang JY, Wakana S, Jiang HY, Li X, Reich DS, Calabresi PA, Pekar JJ, van
640 Zijl PCM, and Mori S. 2008. Tract probability maps in stereotaxic spaces: Analyses of white

matter anatomy and tract-specific quantification. *Neuroimage* 39:336-347.

10.1016/j.neuroimage.2007.07.053

Kaufmann T, van der Meer D, Doan NT, Schwarz E, Lund MJ, Agartz I, Alnæs D, Barch DM, Baur-Streubel R, Bertolino A, Bettella F, Beyer MK, Bøen E, Borgwardt S, Brandt CL, Buitelaar J, Celius EG, Cervenka S, Conzelmann A, Córdova-Palomera A, Dale AM, de Quervain DJF, Di Carlo P, Djurovic S, Dørum ES, Eisenacher S, Elvsashagen T, Espeseth T, Fatouros-Bergman H, Flyckt L, Franke B, Frei O, Haatveit B, Haberg AK, Harbo HF, Hartman CA, Heslenfeld D, Hoekstra PJ, Høgestøl EA, Jernigan T, Jonassen R, Jönsson EG, Kirsch P, Kloszewska I, Kolskar K-K, Landrø NI, Le Hellard S, Lesch K-P, Lovestone S, Lundervold A, Lundervold AJ, Maglanoc LA, Malt UF, Mecocci P, Melle I, Meyer-Lindenberg A, Moberget T, Norbom LB, Nordvik JE, Nyberg L, Oosterlaan J, Papalino M, Papassotiropoulos A, Pauli P, Pergola G, Persson K, Richard G, Rokicki J, Sanders A-M, Selbæk G, Shadrin AA, Smeland OB, Soininen H, Sowa P, Steen VM, Tsolaki M, Ulrichsen KM, Vellas B, Wang L, Westman E, Ziegler GC, Zink M, Andreassen OA, and Westlye LT. 2018. Genetics of brain age suggest an overlap with common brain disorders. *bioRxiv*.

Kuhn T, Kaufmann T, Doan NT, Westlye LT, Jones J, Nunez RA, Bookheimer SY, Singer EJ, Hinkin CH, and Thames AD. 2018. An augmented aging process in brain white matter in HIV. *Hum Brain Mapp*. 10.1002/hbm.24019

Liem F, Varoquaux G, Kynast J, Beyer F, Masouleh SK, Huntenburg JM, Lampe L, Rahim M, Abraham A, Craddock RC, Riedel-Heller S, Luck T, Loeffler M, Schroeter ML, Witte AV, Villringer A, and Margulies DS. 2017. Predicting brain-age from multimodal imaging data captures cognitive impairment. *Neuroimage* 148:179-188.

10.1016/j.neuroimage.2016.11.005

Liu K, Yao S, Chen K, Zhang J, Yao L, Li K, Jin Z, and Guo X. 2017. Structural Brain Network Changes across the Adult Lifespan. *Front Aging Neurosci* 9:275.

10.3389/fnagi.2017.00275

Madan CR, and Kensinger EA. 2018. Predicting age from cortical structure across the lifespan. *Eur J Neurosci* 47:399-416. 10.1111/ejn.13835

Marquand AF, Rezek I, Buitelaar J, and Beckmann CF. 2016. Understanding Heterogeneity in Clinical Cohorts Using Normative Models: Beyond Case-Control Studies. *Biol Psychiatry* 80:552-561. 10.1016/j.biopsych.2015.12.023

- 672 McAvinue LP, Habekost T, Johnson KA, Kyllingsbaek S, Vangkilde S, Bundesen C,
673 and Robertson IH. 2012. Sustained attention, attentional selectivity, and attentional capacity
674 across the lifespan. *Atten Percept Psychophys* 74:1570-1582. 10.3758/s13414-012-0352-6
- 675 Mori S, Wakana S, van Zijl PCM, and Nague-Poetscher LM. 2005. MRI Atlas of
676 Human White Matter. In: Science E, editor. 1st Edition ed: Elsevier Science. p 276.
- 677 Müllner D. 2013. {fastcluster}: Fast Hierarchical, Agglomerative Clustering Routines
678 for {R} and {Python}. *Journal of Statistical Software* 53:1-18.
- 679 Nasreddine ZS, Phillips NA, Bedirian V, Charbonneau S, Whitehead V, Collin I,
680 Cummings JL, and Chertkow H. 2005. The montreal cognitive assessment, MoCA: A brief
681 screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*
682 53:695-699. DOI 10.1111/j.1532-5415.2005.53221.x
- 683 Raz N, Ghisletta P, Rodrigue KM, Kennedy KM, and Lindenberger U. 2010.
684 Trajectories of brain aging in middle-aged and older adults: regional and individual
685 differences. *Neuroimage* 51:501-511. 10.1016/j.neuroimage.2010.03.020
- 686 Ronan L, Alexander-Bloch AF, Wagstyl K, Farooqi S, Brayne C, Tyler LK, Fletcher
687 PC, and Cam-CAN. 2016. Obesity associated with increased brain age from midlife.
688 *Neurobiology of Aging* 47:63-70. 10.1016/j.neurobiolaging.2016.07.010
- 689 Salat DH, Tuch DS, Greve DN, van der Kouwe AJW, Hevelone ND, Zaleta AK, Rosen
690 BR, Fischl B, Corkin S, Rosas HD, and Dale AM. 2005. Age-related alterations in white
691 matter microstructure measured by diffusion tensor imaging. *Neurobiology of Aging* 26:1215-
692 1227. 10.1016/j.neurobiolaging.2004.09.017
- 693 Schnack HG, van Haren NE, Nieuwenhuis M, Hulshoff Pol HE, Cahn W, and Kahn RS.
694 2016. Accelerated Brain Aging in Schizophrenia: A Longitudinal Pattern Recognition Study.
695 *Am J Psychiatry* 173:607-616. 10.1176/appi.ajp.2015.15070922
- 696 Shafto MA, Tyler LK, Dixon M, Taylor JR, Rowe JB, Cusack R, Calder AJ, Marslen-
697 Wilson WD, Duncan J, Dalgleish T, Henson RN, Brayne C, Matthews FE, and Cam-CAN.
698 2014. The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) study protocol: a
699 cross-sectional, lifespan, multidisciplinary examination of healthy cognitive ageing. *Bmc*
700 *Neurology* 14. ARTN 204
701 10.1186/s12883-014-0204-1
- 702 Smith SM, Jenkinson M, Johansen-Berg H, Rueckert D, Nichols TE, Mackay CE,
703 Watkins KE, Ciccarelli O, Cader MZ, Matthews PM, and Behrens TEJ. 2006. Tract-based

- spatial statistics: Voxelwise analysis of multi-subject diffusion data. *Neuroimage* 31:1487-1505. 10.1016/j.neuroimage.2006.02.024
- Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TEJ, Johansen-Berg H, Bannister PR, De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J, Vickers J, Zhang YY, De Stefano N, Brady JM, and Matthews PM. 2004. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* 23:S208-S219. 10.1016/j.neuroimage.2004.07.051
- Sperling G. 1960. The information available in brief visual presentations. *Psychological Monographs: General and Applied* 74:1-29. 10.1037/h0093759
- Steffener J, Habeck C, O'Shea D, Razlighi Q, Bherer L, and Stern Y. 2016. Differences between chronological and brain age are related to education and self-reported physical activity. *Neurobiology of Aging* 40:138-144. 10.1016/j.neurobiolaging.2016.01.014
- Storsve AB, Fjell AM, Tamnes CK, Westlye LT, Overbye K, Aasland HW, and Walhovd KB. 2014. Differential Longitudinal Changes in Cortical Thickness, Surface Area and Volume across the Adult Life Span: Regions of Accelerating and Decelerating Change. *Journal of Neuroscience* 34:8488-8498. 10.1523/Jneurosci.0391-14.2014
- Taylor JR, Williams N, Cusack R, Auer T, Shafto MA, Dixon M, Tyler LK, Cam C, and Henson RN. 2017. The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) data repository: Structural and functional MRI, MEG, and cognitive data from a cross-sectional adult lifespan sample. *Neuroimage* 144:262-269. 10.1016/j.neuroimage.2015.09.018
- Valizadeh SA, Hanggi J, Merillat S, and Jancke L. 2017. Age prediction on the basis of brain anatomical measures. *Hum Brain Mapp* 38:997-1008. 10.1002/hbm.23434
- Wakana S, Caprihan A, Panzenboeck MM, Fallon JH, Perry M, Gollub RL, Hua KG, Zhang JY, Jiang HY, Dubey P, Blitz A, van Zijl P, and Mori S. 2007. Reproducibility of quantitative tractography methods applied to cerebral white matter. *Neuroimage* 36:630-644. 10.1016/j.neuroimage.2007.02.049
- Warnes GR, Bolker B, Bonebakker L, Gentleman R, Liaw WHA, Lumley T, Maechler M, Magnusson A, Moeller S, Schwartz M, and Venables B. 2016. *gplots: Various R Programming Tools for Plotting Data*.
- Wechsler D. 1999. *Wechsler Abbreviated Scale of Intelligence (WASI)*: Psychological Corporation.
- Westlye LT, Walhovd KB, Dale AM, Bjornerud A, Due-Tønnessen P, Engvig A, Grydeland H, Tamnes CK, Ostby Y, and Fjell AM. 2010a. Differentiating maturational and

aging-related changes of the cerebral cortex by use of thickness and signal intensity. *Neuroimage* 52:172-185. 10.1016/j.neuroimage.2010.03.056

Westlye LT, Walhovd KB, Dale AM, Bjornerud A, Due-Tønnessen P, Engvig A, Grydeland H, Tamnes CK, Ostby Y, and Fjell AM. 2010b. Life-Span Changes of the Human Brain White Matter: Diffusion Tensor Imaging (DTI) and Volumetry. *Cerebral Cortex* 20:2055-2068. 10.1093/cercor/bhp280

Wiegand I, Lauritzen MJ, Osler M, Mortensen EL, Rostrup E, Rask L, Richard N, Horwitz A, Benedek K, Vangkilde S, and Petersen A. 2018. EEG correlates of visual short-term memory in older age vary with adult lifespan cognitive development. *Neurobiol Aging* 62:210-220. 10.1016/j.neurobiolaging.2017.10.018

Willer L, Pedersen PM, Forchhammer HB, and Christensen HK. 2016. Cognitive assessment at bedside for iPad: A preliminary validation of a novel cognitive test for stroke patients. *European Stroke Journal* 1:294-301. 10.1177/2396987316665233

Wolfers T, Doan NT, Kaufmann T, Alnæs D, Moberget T, Agartz I, Buitelaar J, Ueland T, Melle I, Beckmann CF, Franke B, Andreassen OA, Westlye LT, and Marquand A. in press. Extensive interindividual differences in schizophrenia and bipolar disorder: mapping biological heterogeneity in reference to normative brain ageing *JAMA Psychiatry*.

Wright SP. 1992. Adjusted P-Values for Simultaneous Inference. *Biometrics* 48:1005-1013. Doi 10.2307/2532694

Figure legends

Fig. 1 Histogram of the age distribution for each sample.

Fig. 2 Comparison between the 11 BAG models. (a) Heatmap of the correlation between different BAGs. (b) Correlations between the chronological age and the predicted age in the test sample for each model with their confidence intervals. (c) Mean and standard error of the 45 p-values ($-\log_{10}(p)$) for the cognitive scores and composite scores for each row (i.e. BAGs), with a higher mean representing a stronger global association across tests. (d) Correlation between the chronological age of each subjects and the combined age, (e) the brain morphometry age, and (f) the white matter microstructure age.

Fig. 3 Hierarchical clustering of the cognitive features. Each cognitive score was normalized and when required the scores were multiplied by -1 to ensure that positive scores represent good performance. The higher panel shows the dendrogram resulting from the hierarchical clustering of the scores in 7 cognitive domains.

Fig. 4 Heatmap of the association between cognitive scores and brain age gaps. The color scale depicts the minus log of the p-values ($-\log_{10}(p)$) for each association. The association marked with a small star represents significant associations after FDR correction, and the one marked with a big star shows significant associations after Bonferroni correction.

Fig. 5 Scatter plots of the 2 strongest associations between cognitive measures and BAG. The color gradient represents the age where lighter color is assigned to older individuals, and darker color to younger individuals. All associations indicate worse performance with higher brain age gap.

Table legends

Table 1. Demographics and cognitive information. * significant associations between cognitive measures with age after FDR correction, ** significant associations between cognitive measures with age after Bonferroni correction

Table 2. Cognitive associations with Brain Age Gap (BAG) – statistics. * FDR significant ** Bonferroni significant

Supplementary Materials

Fig. S1. Heatmap of the association between cognitive scores and brain age gaps using non-linear models, including age, age² and sex as covariates. The color scale depicts the minus log of the p-values ($-\log_{10}(p)$) for each association. The association marked with a small star represents significant associations after FDR correction, and the one marked with a big star shows significant associations after Bonferroni correction.

Table S1. Cognitive associations with Brain Age Gap (BAG) using non-linear models, including age, age² and sex as covariates – statistics. * FDR significant ** Bonferroni significant.

TISSUE-SPECIFIC BRAIN AGE PREDICTION 29

835 Tables

836 Demographics and cognitive information

	Cam-CAN	StrokeMRI Mean (SD)	Range (IQR)	Main effect Age <i>t</i> (<i>p</i>)	Main effect Sex <i>t</i> (<i>p</i>)
Total N (% females)	612 (51.3%)	265 (63.4%)			
Mean age (SD)	54.41 (18.26)	56.95 (14.84)			
Age range	18-87	20-88			
MoCA	-	27.60 (1.72)	21 – 30 (2)	-4.57 (<0.001)**	-2.32 (0.021)
WASI words	-	65.27 (6.60)	44 – 79 (10)	4.72 (<0.001)**	0.10 (0.920)
WASI matrix	-	25.39 (5.64)	7 – 35 (6)	-7.60 (<0.001)**	-0.28 (0.776)
CVLT learning 1-5	-	48.92 (11.37)	17 – 73 (15.5)	-5.05 (<0.001)**	-5.26 (<0.001)
CVLT interference	-	5.53 (2.15)	0 – 13 (3)	-4.33 (<0.001)**	-0.41 (0.681)
CVLT recall	-	10.83 (3.42)	0 – 16 (5)	-6.50 (<0.001)**	5.94 (<0.001)
CVLT delayed recall	-	11.39 (3.44)	0 – 16 (5)	-4.97 (<0.001)**	-5.51 (<0.001)
CVLT recognition hit	-	14.70 (1.50)	8 – 16 (2)	-2.62 (0.0093)*	-2.68 (0.008)
CVLT recognition errors	-	3.79 (3.92)	0 – 18 (4)	5.22 (<0.001)**	4.18 (<0.001)
CVLT recog misses	-	1.30 (1.49)	0 – 8 (2)	2.62 (0.0093)*	2.68 (0.008)
CVLT recog false alarm	-	2.46 (3.48)	0 – 18 (3)	4.45 (<0.001)**	3.59 (0.0004)
CVLT recog correct rejection	-	44.20 (3.92)	30 – 48 (4)	-5.22 (<0.001)**	-4.18 (<0.001)
CVLT d'	-	2.97 (0.72)	0.97 – 3.90 (1.11)	-5.01 (<0.001)**	-4.50 (<0.001)
STROOP 1	-	31.14 (5.66)	21 – 50 (7)	5.05 (<0.001)**	2.44 (0.015)
STROOP 2	-	22.12 (3.49)	14 – 35 (4)	2.89 (0.004)*	2.27 (0.024)
STROOP 3	-	55.86 (14.13)	10 – 108 (15)	7.55 (<0.001)**	2.97 (0.003)
STROOP 4	-	61.74 (14.85)	33 – 117 (19)	7.51 (<0.001)**	1.77 (0.078)
STROOP mean 1 and 2	-	26.54 (4.16)	18.5 – 42 (5)	4.47 (<0.001)**	2.47 (0.014)
STROOP 3 minus mean 1 and 2	-	81.94 (16.51)	34.5 – 145 (18.5)	7.31 (<0.001)**	3.02 (0.003)
STROOP 4 minus mean 1 and 2	-	87.64 (16.73)	53.5 – 142 (24)	7.52 (<0.001)**	1.85 (0.066)
CP – Right motor speed	-	79.56 (23.34)	34 – 153 (32)	-12.25 (<0.001)**	-0.36 (0.716)
CP – Left motor speed	-	81.36 (17.80)	39 – 131 (26)	-12.07 (<0.001)**	0.20 (0.842)
CP – FAS Phonological flow	-	54.70 (14.53)	14 – 95 (19.75)	-0.61 (0.541)	-2.58 (0.011)
CP – FAS Semantic flow	-	51.00 (10.14)	27 – 81 (13)	-2.93 (0.004)*	-3.93 (<0.001)

TISSUE-SPECIFIC BRAIN AGE PREDICTION 30

CP – Visual WM forward ls	-	4.23 (1.01)	2 – 7 (2)	-5.31 (<0.001)**	0.29 (0.774)
CP – Visual WM forward ss	-	5.45 (1.87)	1 – 10 (3)	-6.59 (<0.001)**	-0.25 (0.803)
CP – Visual WM backward ls	-	3.80 (1.28)	0 – 8 (1)	-4.60 (<0.001)**	-1.85 (0.065)
CP – Visual WM backward ss	-	4.56 (2.08)	0 – 12 (3)	-5.48 (<0.001)**	-1.02 (0.309)
CP – Visual WM ss	-	9.96 (3.57)	1 – 21 (4)	-7.04 (<0.001)**	-0.95 (0.342)
CP – Spatial stroop congruent (ms)	-	674.42 (132.77)	410 – 1159 (181)	8.52 (<0.001)**	-1.03 (0.304)
CP – Spatial stroop incongruent (ms)	-	929.52 (198.01)	462 – 1827 (269)	9.41 (<0.001)**	-0.75 (0.451)
CP – Spatial stroop Errors	-	2.17 (2.41)	0 – 11 (3)	0.73 (0.463)	1.59 (0.113)
CP – Spatial stroop numb of reps	-	119.63 (16.64)	55 – 166 (22)	-9.67 (<0.001)**	1.23 (0.219)
CP – Spatial stroop incong – cong (ms)	-	252 (110)	20 – 678 (134.5)	5.73 (<0.001)**	-0.68 (0.498)
CP – Spatspan ls	-	5.37 (1.78)	1 – 10 (2)	-9.12 (<0.001)**	-4.88 (<0.001)
CP – Spatspan tot	-	29.87 (12.43)	3 – 55 (18)	-9.28 (<0.001)**	-4.66 (<0.001)
CP – Coding corr	-	54.50 (12.11)	24 – 88 (16)	-16.69 (<0.001)**	-2.46 (0.015)
CP – Coding error	-	0.67 (0.99)	0 – 5 (1)	-1.10 (0.271)	1.56 (0.121)
TVA – Short-term memory storage (<i>K</i>)	-	3.38 (0.77)	1.46 – 5.53 (1.09)	-7.75 (<0.001)**	-1.52 (0.129)
TVA – Processing speed (<i>C</i>)	-	31.55 (14.07)	5.99 – 89.67 (14.75)	-4.69 (<0.001)**	0.41 (0.6847)
TVA – Perceptual threshold (<i>t₀</i>)	-	23.01 (14.05)	0 – 79.75 (17.59)	5.72 (<0.001)**	-1.94 (0.053)
TVA – Error rate	-	0.10 (0.06)	0.0035 – 0.3316 (0.0983)	-1.35 (0.177)	0.67 (0.502)
Cluster 1	-	-	-	-7.19 (<0.001)**	-5.16 (<0.001)
Cluster 2	-	-	-	-7.28 (<0.001)**	1.61 (0.110)
Cluster 3	-	-	-	-2.01 (0.045)*	-3.99 (<0.001)
Cluster 4	-	-	-	-9.98 (<0.001)**	1.25 (0.212)
Cluster 5	-	-	-	-6.86 (<0.001)**	-2.56 (0.011)
Cluster 6	-	-	-	-15.79 (<0.001)**	-1.08 (0.282)
Cluster 7	-	-	-	-6.50 (<0.001)**	-0.77 (0.440)

Table 1. Demographics and cognitive information. * significant associations between cognitive measures with age after FDR correction, ** significant associations between cognitive measures with age after Bonferroni correction

TISSUE-SPECIFIC BRAIN AGE PREDICTION 31

843 Cognitive associations with Brain Age Gap (BAG) – statistics

Test	Adj R ² no-BAG	BAG	Main effect Age <i>t</i> (<i>p</i>)	Main effect Sex <i>t</i> (<i>p</i>)	Main effect BAG <i>t</i> (<i>p</i>)	Adj R ²
MoCA	0.0907	T1	-4.5596 (<0.001)	-2.3145 (0.021)	-0.124 (0.901)	0.0878
		DTI	-4.5599 (<0.001)	-2.3155 (0.021)	1.5914 (0.113)	0.0966
		Combined	-4.5653 (<0.001)	-2.3176 (0.021)	-0.4626 (0.644)	0.0885
WASI words	0.0731	T1	4.7118 (<0.001)	0.1020 (0.919)	-0.2169 (0.828)	0.0704
		DTI	4.7056 (<0.001)	0.1121 (0.911)	-0.8126 (0.417)	0.0727
		Combined	4.7091 (<0.001)	0.1041 (0.917)	-0.4827 (0.630)	0.0711
WASI matrix	0.1791	T1	-7.6061 (<0.001)	-0.2785 (0.781)	-0.9158 (0.361)	0.1793
		DTI	-7.6610 (<0.001)	-0.2624 (0.793)	-1.6546 (0.099)	0.1854
		Combined	-7.6128 (<0.001)	-0.2726 (0.785)	-1.1102 (0.268)	0.1806
CVLT learning 1-5	0.1810	T1	-5.0373 (<0.001)	-5.2514 (<0.001)	-0.2505 (0.802)	0.1750
		DTI	-5.0418 (<0.001)	-5.2533 (<0.001)	-0.3608 (0.719)	0.1753
		Combined	-5.0387 (<0.001)	-5.2522 (<0.001)	-0.2492 (0.803)	0.1750
CVLT interference	0.0664	T1	-4.3256 (<0.001)	-0.4062 (0.685)	-0.9588 (0.339)	0.0626
		DTI	-4.3218 (<0.001)	-0.4104 (0.682)	-0.2391 (0.811)	0.0594
		Combined	-4.3202 (<0.001)	-0.4101 (0.682)	-0.1875 (0.851)	0.0594
CVLT recall	0.2438	T1	-6.4897 (<0.001)	-5.9257 (<0.001)	-0.4868 (0.627)	0.2397
		DTI	-6.4885 (<0.001)	-5.9257 (<0.001)	-0.1245 (0.901)	0.2391
		Combined	-6.5080 (<0.001)	-5.9373 (<0.001)	-1.1114 (0.268)	0.2427
CVLT delayed recall	0.1850	T1	-4.9636 (<0.001)	-5.4973 (<0.001)	0.1421 (0.887)	0.1808
		DTI	-4.9611 (<0.001)	-5.4969 (<0.001)	0.224 (0.823)	0.1809
		Combined	-4.9655 (<0.001)	-5.4954 (<0.001)	-0.3038 (0.762)	0.1810
CVLT recognition hits	0.0494	T1	-2.6125 (0.010)	-2.6822 (0.008)	-0.8586 (0.391)	0.0486
		DTI	-2.6144 (0.010)	-2.6786 (0.008)	0.0946 (0.925)	0.0459
		Combined	-2.6212 (0.009)	-2.6854 (0.008)	-1.0724 (0.285)	0.0501
CVLT recognition errors	0.1526	T1	5.2227 (<0.001)	4.1850 (<0.001)	-0.8471 (0.398)	0.1528
		DTI	5.2115 (<0.001)	4.1755 (<0.001)	-0.5651 (0.573)	0.1514
		Combined	5.2139 (<0.001)	4.1740 (<0.001)	-0.2537 (0.800)	0.1506
CVLT recog misses	0.0494	T1	2.6125 (0.010)	2.6822 (0.008)	0.8586 (0.391)	0.0486
		DTI	2.6144 (0.010)	2.6786 (0.008)	-0.0946 (0.925)	0.0459
		Combined	2.6212 (0.009)	2.6854 (0.008)	1.0724 (0.285)	0.0501
CVLT recog false alarm	0.1150	T1	4.4519 (<0.001)	3.5827 (<0.001)	-0.776 (0.439)	0.1146
		DTI	4.4378 (<0.001)	3.5803 (<0.001)	-0.5207 (0.603)	0.1134
		Combined	4.4418 (<0.001)	3.5788 (<0.001)	-0.3488 (0.728)	0.1129
CVLT recog correct rejection	0.1526	T1	-5.2227 (<0.001)	-4.1850 (<0.001)	0.8471 (0.398)	0.1528
		DTI	-5.2115 (<0.001)	-4.1755 (<0.001)	0.5651 (0.573)	0.1514
		Combined	-5.2139 (<0.001)	-4.1740 (<0.001)	0.2537 (0.800)	0.1506
CVLT d'	0.1566	T1	-5.0074 (<0.001)	-4.4914 (<0.001)	0.3628 (0.717)	0.1536
		DTI	-5.0021 (<0.001)	-4.4969 (<0.001)	0.8538 (0.394)	0.1556
		Combined	-5.0038 (<0.001)	-4.4902 (<0.001)	0.1699 (0.865)	0.1533
STROOP 1	0.1118	T1	5.1466 (<0.001)	2.4999 (0.013)	2.6939 (0.008)	0.1299

TISSUE-SPECIFIC BRAIN AGE PREDICTION 32

		DTI	5.0968 (<0.001)	2.4769 (0.014)	1.6664 (0.097)	0.1147
		Combined	5.2111 (<0.001)	2.5317 (0.012)	3.3767 (<0.001)*	0.1434
STROOP 2	0.0477	T1	2.8868 (0.004)	2.2619 (0.025)	0.1557 (0.876)	0.0433
		DTI	2.8768 (0.004)	2.2489 (0.025)	-0.4639 (0.643)	0.0440
		Combined	2.8949 (0.004)	2.2713 (0.024)	0.4976 (0.619)	0.0442
STROOP 3	0.2104	T1	7.5930 (<0.001)	2.9898 (0.003)	1.5092 (0.133)	0.2109
		DTI	7.6511 (<0.001)	3.0224 (0.003)	2.231 (0.027)	0.2190
		Combined	7.6793 (<0.001)	3.0233 (0.003)	2.5768 (0.011)	0.2240
STROOP 4	0.1887	T1	7.5403 (<0.001)	1.7884 (0.075)	1.2397 (0.216)	0.1906
		DTI	7.5847 (<0.001)	1.8121 (0.071)	1.7368 (0.084)	0.1953
		Combined	7.6387 (<0.001)	1.8247 (0.069)	2.3662 (0.019)	0.2033
STROOP mean 1 and 2	0.0949	T1	4.5089 (<0.001)	2.5033 (0.013)	1.5875 (0.114)	0.0978
		DTI	4.4750 (<0.001)	2.4760 (0.014)	0.3927 (0.695)	0.0894
		Combined	4.5432 (<0.001)	2.5399 (0.012)	2.0254 (0.044)	0.1034
STROOP 3 minus mean 1 and 2	0.2051	T1	7.3383 (<0.001)	3.0427 (0.003)	1.1397 (0.256)	0.2021
		DTI	7.3613 (<0.001)	3.0703 (0.002)	1.3546 (0.177)	0.2038
		Combined	7.4197 (<0.001)	3.1063 (0.002)	2.1881 (0.030)	0.2130
STROOP 4 minus mean 1 and 2	0.1936	T1	7.5360 (<0.001)	1.8671 (0.063)	0.8763 (0.382)	0.1919
		DTI	7.5297 (<0.001)	1.8697 (0.063)	0.6331 (0.527)	0.1907
		Combined	7.6081 (<0.001)	1.9215 (0.056)	1.7531 (0.081)	0.1993
CP – Right motor speed	0.3695	T1	-12.2893 (<0.001)	-0.3592 (0.720)	-1.5504 (0.122)	0.3676
		DTI	-12.2318 (<0.001)	-0.3612 (0.718)	-0.3435 (0.732)	0.3620
		Combined	-12.3125 (<0.001)	-0.3587 (0.720)	-1.8139 (0.071)	0.3697
CP – Left motor speed	0.3630	T1	-12.1437 (<0.001)	0.2100 (0.834)	-1.9945 (0.047)	0.3634
		DTI	-12.0669 (<0.001)	0.2081 (0.835)	-0.8704 (0.385)	0.3555
		Combined	-12.2516 (<0.001)	0.2149 (0.830)	-2.9047 (0.004)	0.3740
CP – FAS Semantic flow	0.0840	T1	-2.9562 (0.003)	-3.9454 (<0.001)	-2.0826 (0.038)	0.0960
		DTI	-2.9607 (0.003)	-3.9388 (<0.001)	-2.0997 (0.037)	0.0963
		Combined	-2.9513 (0.004)	-3.9389 (<0.001)	-1.8308 (0.068)	0.0926
CP – Visual WM forward ls	0.0936	T1	-5.3071 (<0.001)	0.2850 (0.776)	-0.5838 (0.560)	0.0906
		DTI	-5.3392 (<0.001)	0.2963 (0.767)	-1.7204 (0.087)	0.0999
		Combined	-5.3059 (<0.001)	0.2853 (0.776)	-0.3127 (0.755)	0.0897
CP – Visual WM forward ss	0.1416	T1	-6.5795 (<0.001)	-0.2502 (0.803)	-0.2158 (0.829)	0.1375
		DTI	-6.6000 (<0.001)	-0.2448 (0.807)	-1.1695 (0.243)	0.1420
		Combined	-6.5786 (<0.001)	-0.2496 (0.803)	-0.02 (0.984)	0.1373
CP – Visual WM backward ls	0.0852	T1	-4.5941 (<0.001)	-1.8511 (0.065)	-0.1047 (0.917)	0.0820
		DTI	-4.6170 (<0.001)	-1.8545 (0.065)	-1.3334 (0.184)	0.0884
		Combined	-4.6051 (<0.001)	-1.8550 (0.065)	-0.8013 (0.424)	0.0843
CP – Visual WM backward ss	0.1022	T1	-5.4741 (<0.001)	-1.0181 (0.310)	-0.2721 (0.786)	0.1015
		DTI	-5.4971 (<0.001)	-1.0179 (0.310)	-1.3043 (0.193)	0.1072
		Combined	-5.4898 (<0.001)	-1.0215 (0.308)	-1.0074 (0.315)	0.1048
CP – Visual WM ss	0.1607	T1	-7.0322 (<0.001)	-0.9515 (0.342)	-0.3013 (0.763)	0.1591
		DTI	-7.0622 (<0.001)	-0.9511 (0.342)	-1.3634 (0.174)	0.1649

TISSUE-SPECIFIC BRAIN AGE PREDICTION 33

		Combined	-7.0399 (<0.001)	-0.9528 (0.342)	-0.6665 (0.506)	0.1603
CP – Spatial stroop congruent	0.2288	T1	8.6156 (<0.001)	-1.0080 (0.314)	2.1921 (0.029)	0.2288
		DTI	8.6687 (<0.001)	-1.0021 (0.317)	2.6995 (0.007)	0.2362
		Combined	8.8278 (<0.001)	-0.9828 (0.327)	3.9007 (<0.001)**	0.2588
CP – Spatial stroop incongruent	0.2548	T1	9.5489 (<0.001)	-0.7429 (0.458)	2.6569 (0.008)	0.2700
		DTI	9.5931 (<0.001)	-0.7587 (0.449)	2.8817 (0.004)	0.2735
		Combined	9.7197 (<0.001)	-0.7378 (0.461)	3.8071 (<0.001)**	0.2903
CP – Spatial stroop numb of reps	0.2731	T1	-9.7755 (<0.001)	1.2211 (0.223)	-2.2212 (0.027)	0.2753
		DTI	-9.8507 (<0.001)	1.2328 (0.219)	-2.9614 (0.003)	0.2859
		Combined	-9.9891 (<0.001)	1.2198 (0.224)	-3.8816 (<0.001)**	0.3027
CP – Spatial stroop incong – cong	0.1012	T1	5.7663 (<0.001)	-0.6595 (0.510)	1.5611 (0.120)	0.1134
		DTI	5.7466 (<0.001)	-0.6678 (0.505)	0.9705 (0.333)	0.1081
		Combined	5.7568 (<0.001)	-0.6584 (0.511)	1.2056 (0.229)	0.1099
CP – Spatspan ls	0.3055	T1	-9.1038 (<0.001)	-4.8656 (<0.001)	-0.032 (0.975)	0.3009
		DTI	-9.1746 (<0.001)	-4.9104 (<0.001)	-1.5749 (0.117)	0.3077
		Combined	-9.1043 (<0.001)	-4.8663 (<0.001)	-0.075 (0.940)	0.3009
CP – Spatspan total	0.3057	T1	-9.2664 (<0.001)	-4.6439 (<0.001)	0.1074 (0.915)	0.3024
		DTI	-9.3260 (<0.001)	-4.6815 (<0.001)	-1.3773 (0.170)	0.3076
		Combined	-9.2686 (<0.001)	-4.6461 (<0.001)	-0.0612 (0.951)	0.3024
CP – Coding corr	0.5387	T1	-16.7647 (<0.001)	-2.5004 (0.013)	-1.6149 (0.108)	0.5352
		DTI	-17.0893 (<0.001)	-2.5467 (0.012)	-3.3998 (<0.001)*	0.5510
		Combined	-17.0071 (<0.001)	-2.5604 (0.011)	-3.0056 (0.003)	0.5467
TVA - Short-term memory storage (K)	0.2013	T1	-7.7691 (<0.001)	-1.5196 (0.130)	-1.1179 (0.265)	0.1981
		DTI	-7.8117 (<0.001)	-1.5383 (0.125)	-2.0302 (0.043)	0.2070
		Combined	-7.7525 (<0.001)	-1.5195 (0.130)	-0.9537 (0.341)	0.1970
TVA - Perceptual threshold (t_0)	0.0764	T1	5.7303 (<0.001)	-1.9470 (0.053)	0.9617 (0.337)	0.1141
		DTI	5.7333 (<0.001)	-1.9444 (0.053)	1.1066 (0.270)	0.1152
		Combined	5.7523 (<0.001)	-1.9587 (0.051)	1.8346 (0.068)	0.1226
TVA - Processing speed (C)	0.1304	T1	-4.6692 (<0.001)	0.3969 (0.692)	0.8093 (0.419)	0.0723
		DTI	-4.6800 (<0.001)	0.4053 (0.686)	0.1402 (0.889)	0.0699
		Combined	-4.6827 (<0.001)	0.3944 (0.694)	0.8916 (0.374)	0.0728
Cluster 1	0.2470	T1	-7.1741 (<0.001)	-5.1567 (<0.001)	-0.1927 (0.847)	0.2440
		DTI	-7.1623 (<0.001)	-5.1410 (<0.001)	0.3683 (0.713)	0.2443
		Combined	-7.1805 (<0.001)	-5.1641 (<0.001)	-0.3879 (0.699)	0.2443
Cluster 2	0.1720	T1	-7.2680 (<0.001)	1.6030 (0.110)	-0.1013 (0.919)	0.1687
		DTI	-7.2785 (<0.001)	1.6062 (0.110)	-0.6549 (0.513)	0.1701
		Combined	-7.2740 (<0.001)	1.6104 (0.109)	-0.6382 (0.524)	0.1700
Cluster 3	0.0698	T1	-2.0177 (0.045)	-3.9824 (<0.001)	-0.8103 (0.419)	0.0686
		DTI	-2.0337 (0.043)	-3.9969 (<0.001)	-1.84 (0.067)	0.0783
		Combined	-2.0185 (0.045)	-3.9877 (<0.001)	-0.9765 (0.330)	0.0697
Cluster 4	0.2783	T1	-10.1319 (<0.001)	1.2314 (0.219)	-2.5436 (0.012)	0.2937
		DTI	-10.1479 (<0.001)	1.2377 (0.217)	-2.5207 (0.012)	0.2933
		Combined	-10.3013 (<0.001)	1.2196 (0.224)	-3.6163 (<0.001)*	0.3113

TISSUE-SPECIFIC BRAIN AGE PREDICTION 34

Cluster 5	0.1772	T1	-6.8872 (<0.001)	-2.5902 (0.010)	-1.1084 (0.269)	0.1779
		DTI	-6.8667 (<0.001)	-2.5805 (0.010)	-0.5825 (0.561)	0.1750
		Combined	-6.9577 (<0.001)	-2.6481 (0.009)	-1.9103 (0.057)	0.1858
Cluster 6	0.5092	T1	-15.9345 (<0.001)	-1.1148 (0.266)	-1.8971 (0.059)	0.5145
		DTI	-15.9719 (<0.001)	-1.1080 (0.269)	-2.0875 (0.038)	0.5160
		Combined	-16.0156 (<0.001)	-1.1196 (0.264)	-2.459 (0.015)	0.5193
Cluster 7	0.1399	T1	-6.4852 (<0.001)	-0.7736 (0.440)	-0.3433 (0.732)	0.1369
		DTI	-6.5210 (<0.001)	-0.7689 (0.443)	-1.6007 (0.111)	0.1452
		Combined	-6.4926 (<0.001)	-0.7759 (0.439)	-0.63 (0.529)	0.1379

Table 2. Cognitive associations with BAG – statistics. * FDR significant ** Bonferroni significant.

TISSUE-SPECIFIC BRAIN AGE PREDICTION 36

889 **Supplementary Tables**

890 Cognitive associations with Brain Age Gap (BAG) using non-linear models, including age,
891 age² and sex as covariates – statistics.

Test	Adj R ² no-BAG	BAG	Main effect Age <i>t(p)</i>	Main effect Age ² <i>t(p)</i>	Main effect Sex <i>t(p)</i>	Main effect BAG <i>t(p)</i>	Adj R ²
MoCA	0.088	T1	-4.5563 (<0.001)	-0.2752 (0.7834)	-2.1792 (0.0302)	-0.1146 (0.9088)	0.0845
		DTI	-4.5575 (<0.001)	-0.3078 (0.7585)	-2.1727 (0.0307)	1.5937 (0.1122)	0.0934
		Combined	-4.561 (<0.001)	-0.2413 (0.8095)	-2.1894 (0.0295)	-0.4399 (0.6604)	0.0851
WASI words	0.0792	T1	4.7079 (<0.001)	-1.5614 (0.1197)	0.4698 (0.6389)	-0.1719 (0.8637)	0.0756
		DTI	4.7017 (<0.001)	-1.5532 (0.1216)	0.4775 (0.6334)	-0.7843 (0.4336)	0.0778
		Combined	4.7058 (<0.001)	-1.5325 (0.1266)	0.4655 (0.642)	-0.3554 (0.7226)	0.076
WASI matrix	0.1825	T1	-7.6085 (<0.001)	-1.3223 (0.1873)	0.048 (0.9618)	-0.8787 (0.3804)	0.1817
		DTI	-7.6627 (<0.001)	-1.311 (0.1911)	0.0606 (0.9517)	-1.6227 (0.1059)	0.1876
		Combined	-7.6132 (<0.001)	-1.2678 (0.206)	0.0407 (0.9676)	-1.0117 (0.3126)	0.1825
CVLT learning 1-5	0.1757	T1	-5.0181 (<0.001)	0.5198 (0.6036)	-5.2162 (<0.001)	-0.2662 (0.7903)	0.1727
		DTI	-5.0228 (<0.001)	0.5204 (0.6032)	-5.2181 (<0.001)	-0.3723 (0.71)	0.1729
		Combined	-5.0196 (<0.001)	0.5363 (0.5922)	-5.2205 (<0.001)	-0.2959 (0.7675)	0.1727
CVLT interference	0.0593	T1	-4.3175 (<0.001)	-0.062 (0.9506)	-0.3788 (0.7051)	-0.9545 (0.3407)	0.0589
		DTI	-4.3143 (<0.001)	-0.086 (0.9316)	-0.3772 (0.7064)	-0.2366 (0.8132)	0.0558
		Combined	-4.3124 (<0.001)	-0.075 (0.9403)	-0.3794 (0.7047)	-0.1796 (0.8576)	0.0557
CVLT recall	0.2401	T1	-6.4676 (<0.001)	0.6091 (0.543)	-5.8925 (<0.001)	-0.5048 (0.6141)	0.2379
		DTI	-6.4667 (<0.001)	0.5965 (0.5514)	-5.8892 (<0.001)	-0.1381 (0.8902)	0.2371
		Combined	-6.4859 (<0.001)	0.6973 (0.4863)	-5.9255 (<0.001)	-1.1682 (0.2438)	0.2412
CVLT delayed recall	0.1822	T1	-4.9433 (<0.001)	0.6624 (0.5083)	-5.4919 (<0.001)	0.1193 (0.9052)	0.179
		DTI	-4.9411 (<0.001)	0.6618 (0.5087)	-5.4912 (<0.001)	0.2079 (0.8354)	0.1791
		Combined	-4.9454 (<0.001)	0.6981 (0.4857)	-5.4986 (<0.001)	-0.3668 (0.7141)	0.1794
CVLT recognition hits	0.0464	T1	-2.615 (0.0095)	-0.3694 (0.7121)	-2.5144 (0.0125)	-0.8444 (0.3992)	0.0453
		DTI	-2.6175 (0.0094)	-0.3994 (0.6899)	-2.504 (0.0129)	0.1039 (0.9173)	0.0427
		Combined	-2.622 (0.0093)	-0.3004 (0.7641)	-2.5328 (0.0119)	-1.0382 (0.3002)	0.0467
CVLT recognition errors	0.1567	T1	5.2609 (<0.001)	1.4118 (0.1592)	3.7395 (<0.001)	-0.8964 (0.3709)	0.1561
		DTI	5.2485 (<0.001)	1.3947 (0.1643)	3.7336 (<0.001)	-0.5993 (0.5495)	0.1546
		Combined	5.2514 (<0.001)	1.4099 (0.1598)	3.7277 (<0.001)	-0.3837 (0.7015)	0.1539
CVLT recog misses	0.0464	T1	2.615 (0.0095)	0.3694 (0.7121)	2.5144 (0.0125)	0.8444 (0.3992)	0.0453
		DTI	2.6175 (0.0094)	0.3994 (0.6899)	2.504 (0.0129)	-0.1039 (0.9173)	0.0427
		Combined	2.622 (0.0093)	0.3004 (0.7641)	2.5328 (0.0119)	1.0382 (0.3002)	0.0467
CVLT recog false alarm	0.1191	T1	4.4883 (<0.001)	1.4053 (0.1612)	3.1557 (0.0018)	-0.8236 (0.411)	0.1179
		DTI	4.4728 (<0.001)	1.3906 (0.1656)	3.1567 (0.0018)	-0.5544 (0.5798)	0.1167
		Combined	4.4784 (<0.001)	1.4159 (0.158)	3.1475 (0.0018)	-0.4786 (0.6326)	0.1164
CVLT recog correct rejection	0.1567	T1	-5.2609 (<0.001)	-1.4118 (0.1592)	-3.7395 (<0.001)	0.8964 (0.3709)	0.1561
		DTI	-5.2485 (<0.001)	-1.3947 (0.1643)	-3.7336 (<0.001)	0.5993 (0.5495)	0.1546
		Combined	-5.2514 (<0.001)	-1.4099 (0.1598)	-3.7277 (<0.001)	0.3837 (0.7015)	0.1539
CVLT d'	0.1548	T1	-5.0157 (<0.001)	-0.7018 (0.4835)	-4.1938 (<0.001)	0.3855 (0.7002)	0.1519

TISSUE-SPECIFIC BRAIN AGE PREDICTION 37

		DTI	-5.0104 (<0.001)	-0.7106 (0.478)	-4.1973 (<0.001)	0.8695 (0.3854)	0.1539
		Combined	-5.0122 (<0.001)	-0.7079 (0.4796)	-4.1897 (<0.001)	0.2343 (0.8149)	0.1516
STROOP 1	0.1181	T1	5.2178 (<0.001)	1.8817 (0.061)	1.9786 (0.0489)	2.6477 (0.0086)	0.1385
		DTI	5.1698 (<0.001)	1.915 (0.0566)	1.9488 (0.0524)	1.6361 (0.103)	0.1239
		Combined	5.2707 (<0.001)	1.7146 (0.0876)	2.0444 (0.0419)	3.243 (0.0013)	0.1499
STROOP 2	0.0565	T1	2.955 (0.0034)	1.8769 (0.0617)	1.7273 (0.0853)	0.0898 (0.9286)	0.0528
		DTI	2.9455 (0.0035)	1.897 (0.059)	1.7093 (0.0886)	-0.5228 (0.6016)	0.0538
		Combined	2.9595 (0.0034)	1.8439 (0.0664)	1.7389 (0.0833)	0.3307 (0.7411)	0.0531
STROOP 3	0.2519	T1	7.9387 (<0.001)	4.0215 (<0.001)**	2.0391 (0.0425)	1.4584 (0.146)	0.2552
		DTI	8.0033 (<0.001)	4.0506 (<0.001)**	2.0666 (0.0398)	2.245 (0.0256)	0.2635
		Combined	8.0064 (<0.001)	3.9001 (<0.001)**	2.0915 (0.0375)	2.3542 (0.0193)	0.265
STROOP 4	0.2092	T1	7.727 (<0.001)	2.7144 (0.0071)	1.1078 (0.269)	1.1959 (0.2329)	0.2105
		DTI	7.7745 (<0.001)	2.7322 (0.0067)	1.1277 (0.2605)	1.7316 (0.0846)	0.2154
		Combined	7.8094 (<0.001)	2.5945 (0.01)	1.1659 (0.2448)	2.2006 (0.0287)	0.221
STROOP mean 1 and 2	0.1065	T1	4.6014 (<0.001)	2.1809 (0.0301)	1.8851 (0.0606)	1.5137 (0.1314)	0.1111
		DTI	4.5693 (<0.001)	2.2194 (0.0274)	1.8487 (0.0657)	0.3205 (0.7489)	0.1033
		Combined	4.6255 (<0.001)	2.0629 (0.0402)	1.9411 (0.0534)	1.8349 (0.0677)	0.1148
STROOP 3 minus mean 1 and 2	0.2486	T1	7.6928 (<0.001)	4.0734 (<0.001)**	2.0223 (0.0442)	1.0347 (0.3018)	0.2488
		DTI	7.7161 (<0.001)	4.0759 (<0.001)**	2.0485 (0.0416)	1.2741 (0.2038)	0.2504
		Combined	7.7517 (<0.001)	3.9419 (<0.001)**	2.0994 (0.0368)	1.8804 (0.0612)	0.2561
STROOP 4 minus mean 1 and 2	0.2225	T1	7.7912 (<0.001)	3.224 (0.0014)*	1.0274 (0.3052)	0.7875 (0.4317)	0.2213
		DTI	7.7853 (<0.001)	3.233 (0.0014)*	1.0273 (0.3053)	0.5573 (0.5778)	0.2203
		Combined	7.8437 (<0.001)	3.1135 (0.0021)*	1.0945 (0.2748)	1.4951 (0.1362)	0.2263
CP - Right motor speed	0.3621	T1	-12.2583 (<0.001)	0.4241 (0.6718)	-0.4499 (0.6532)	-1.56 (0.12)	0.3656
		DTI	-12.201 (<0.001)	0.3813 (0.7033)	-0.4415 (0.6592)	-0.3491 (0.7273)	0.3599
		Combined	-12.2818 (<0.001)	0.5259 (0.5994)	-0.4741 (0.6358)	-1.8479 (0.0658)	0.368
CP - Left motor speed	0.3586	T1	-12.1901 (<0.001)	-1.3758 (0.1701)	0.5329 (0.5946)	-1.9551 (0.0517)	0.3656
		DTI	-12.1163 (<0.001)	-1.4124 (0.159)	0.5396 (0.5899)	-0.8474 (0.3976)	0.3579
		Combined	-12.2849 (<0.001)	-1.2158 (0.2252)	0.4997 (0.6177)	-2.7995 (0.0055)	0.3752
CP - FAS semantic flow	0.1047	T1	-3.0413 (0.0026)	-2.5748 (0.0106)	-3.2544 (0.0013)	-2.0265 (0.0437)	0.1153
		DTI	-3.0473 (0.0025)	-2.6035 (0.0098)	-3.2424 (0.0013)	-2.0796 (0.0385)	0.116
		Combined	-3.0316 (0.0027)	-2.4889 (0.0134)	-3.2639 (0.0012)	-1.642 (0.1018)	0.1105
CP - Visual WM forward ls	0.0897	T1	-5.3048 (<0.001)	-0.2803 (0.7795)	0.3424 (0.7323)	-0.5645 (0.5729)	0.0873
		DTI	-5.3364 (<0.001)	-0.2747 (0.7838)	0.352 (0.7252)	-1.71 (0.0885)	0.0966
		Combined	-5.3039 (<0.001)	-0.2873 (0.7742)	0.3445 (0.7307)	-0.2836 (0.7769)	0.0864
CP - Visual WM forward ss	0.1388	T1	-6.5947 (<0.001)	-0.6439 (0.5202)	-0.0916 (0.9271)	-0.1756 (0.8608)	0.1355
		DTI	-6.6142 (<0.001)	-0.6289 (0.53)	-0.0903 (0.9281)	-1.1522 (0.2503)	0.1399
		Combined	-6.5945 (<0.001)	-0.657 (0.5118)	-0.0873 (0.9305)	0.0421 (0.9664)	0.1354
CP - Visual WM backward ls	0.0852	T1	-4.6076 (<0.001)	-0.9343 (0.351)	-1.567 (0.1184)	-0.068 (0.9458)	0.0816
		DTI	-4.6299 (<0.001)	-0.9132 (0.362)	-1.5757 (0.1163)	-1.3135 (0.1902)	0.0878
		Combined	-4.6163 (<0.001)	-0.8736 (0.3831)	-1.5842 (0.1144)	-0.7237 (0.4699)	0.0835
CP - Visual WM backward ss	0.1026	T1	-5.4768 (<0.001)	-0.619 (0.5365)	-0.8352 (0.4044)	-0.2473 (0.8049)	0.0993

TISSUE-SPECIFIC BRAIN AGE PREDICTION 38

		DTI	-5.4994 (<0.001)	-0.604 (0.5464)	-0.8389 (0.4023)	-1.2897 (0.1983)	0.105
		Combined	-5.4903 (<0.001)	-0.5466 (0.5852)	-0.8555 (0.3931)	-0.9558 (0.3401)	0.1023
CP - Visual WM ss	0.1618	T1	-7.0453 (<0.001)	-0.9494 (0.3433)	-0.6911 (0.4902)	-0.2639 (0.7921)	0.1588
		DTI	-7.0749 (<0.001)	-0.9354 (0.3504)	-0.6944 (0.4881)	-1.3432 (0.1804)	0.1645
		Combined	-7.051 (<0.001)	-0.9076 (0.365)	-0.7016 (0.4836)	-0.5866 (0.558)	0.1597
CP - Spatial stroop congruent	0.2152	T1	8.6103 (<0.001)	0.5061 (0.6132)	-1.099 (0.2728)	2.1677 (0.0311)	0.2265
		DTI	8.6646 (<0.001)	0.539 (0.5904)	-1.1012 (0.2719)	2.6844 (0.0078)	0.234
		Combined	8.8148 (<0.001)	0.27 (0.7874)	-1.0172 (0.3101)	3.8562 (<0.001)**	0.2561
CP - Spatial stroop incongruent	0.2498	T1	9.5215 (<0.001)	-0.3129 (0.7546)	-0.6399 (0.5228)	2.6625 (0.0083)	0.2674
		DTI	9.5663 (<0.001)	-0.2497 (0.803)	-0.6711 (0.5028)	2.8797 (0.0043)	0.2708
		Combined	9.6908 (<0.001)	-0.5285 (0.5976)	-0.5807 (0.5619)	3.8325 (<0.001)*	0.2883
CP - Spatial stroop numb of reps	0.2613	T1	-9.7578 (<0.001)	-0.1542 (0.8776)	1.2196 (0.2237)	-2.2091 (0.0281)	0.2725
		DTI	-9.8334 (<0.001)	-0.1782 (0.8587)	1.2369 (0.2173)	-2.951 (0.0035)	0.2831
		Combined	-9.967 (<0.001)	0.0905 (0.928)	1.1582 (0.2479)	-3.8672 (<0.001)**	0.2999
CP - Spatial stroop incong - cong	0.1076	T1	5.7387 (<0.001)	-0.9619 (0.337)	-0.4011 (0.6887)	1.5968 (0.1116)	0.1131
		DTI	5.7189 (<0.001)	-0.9105 (0.3635)	-0.4223 (0.6732)	0.9815 (0.3273)	0.1075
		Combined	5.7304 (<0.001)	-1.0014 (0.3176)	-0.3888 (0.6978)	1.2835 (0.2005)	0.1099
CP - Spatspan ls	0.3027	T1	-9.0803 (<0.001)	0.8225 (0.4116)	-4.9183 (<0.001)	-0.0628 (0.95)	0.3
		DTI	-9.1511 (<0.001)	0.8555 (0.3931)	-4.9703 (<0.001)	-1.5904 (0.113)	0.307
		Combined	-9.0818 (<0.001)	0.8301 (0.4073)	-4.9202 (<0.001)	-0.1448 (0.885)	0.3
CP - Spatspan total	0.3033	T1	-9.2413 (<0.001)	0.5919 (0.5545)	-4.6451 (<0.001)	0.085 (0.9324)	0.3006
		DTI	-9.3007 (<0.001)	0.6245 (0.5329)	-4.6899 (<0.001)	-1.3875 (0.1665)	0.3059
		Combined	-9.2443 (<0.001)	0.6028 (0.5472)	-4.6491 (<0.001)	-0.1118 (0.9111)	0.3006
CP - Coding corr	0.5307	T1	-16.7381 (<0.001)	-0.3279 (0.7433)	-2.3396 (0.0201)	-1.596 (0.1117)	0.5335
		DTI	-17.0629 (<0.001)	-0.3481 (0.7281)	-2.3804 (0.018)	-3.3875 (<0.001)*	0.5495
		Combined	-16.9736 (<0.001)	-0.1442 (0.8855)	-2.4411 (0.0153)	-2.976 (0.0032)	0.5449
TVA - Short-term memory storage (K)	0.1941	T1	-7.7533 (<0.001)	-0.0891 (0.9291)	-1.4529 (0.1475)	-1.1129 (0.2668)	0.1949
		DTI	-7.7958 (<0.001)	-0.0894 (0.9288)	-1.471 (0.1426)	-2.0247 (0.044)	0.2039
		Combined	-7.7352 (<0.001)	-0.0343 (0.9726)	-1.4655 (0.144)	-0.9449 (0.3456)	0.1938
TVA - Perceptual threshold (t ₀)	0.1146	T1	5.7716 (<0.001)	0.9866 (0.3248)	-2.1312 (0.034)	0.9264 (0.3551)	0.1141
		DTI	5.7764 (<0.001)	1.0081 (0.3144)	-2.1342 (0.0338)	1.0951 (0.2745)	0.1153
		Combined	5.784 (<0.001)	0.8478 (0.3973)	-2.1074 (0.0361)	1.7412 (0.0829)	0.1217
TVA - Processing speed (C)	0.0699	T1	-4.66 (<0.001)	0.1122 (0.9107)	0.3556 (0.7225)	0.804 (0.4222)	0.0686
		DTI	-4.6708 (<0.001)	0.135 (0.8927)	0.3579 (0.7207)	0.1387 (0.8898)	0.0662
		Combined	-4.6734 (<0.001)	0.0619 (0.9507)	0.3657 (0.7149)	0.8815 (0.3789)	0.0691
Cluster 1	0.2446	T1	-7.1703 (<0.001)	-0.4829 (0.6296)	-4.8899 (<0.001)	-0.1702 (0.865)	0.2416
		DTI	-7.1589 (<0.001)	-0.5024 (0.6158)	-4.8709 (<0.001)	0.3825 (0.7024)	0.242
		Combined	-7.1752 (<0.001)	-0.4542 (0.6501)	-4.8989 (<0.001)	-0.3391 (0.7349)	0.2419
Cluster 2	0.1695	T1	-7.2665 (<0.001)	-0.5147 (0.6072)	1.6797 (0.0943)	-0.0814 (0.9352)	0.1662
		DTI	-7.2772 (<0.001)	-0.5154 (0.6068)	1.6829 (0.0937)	-0.6514 (0.5154)	0.1676
		Combined	-7.2707 (<0.001)	-0.4611 (0.6451)	1.6725 (0.0957)	-0.5914 (0.5548)	0.1674
Cluster 3	0.0767	T1	-2.065 (0.0399)	-1.6934 (0.0916)	-3.4589 (<0.001)	-0.7735 (0.4399)	0.0753

TISSUE-SPECIFIC BRAIN AGE PREDICTION 39

Cluster 4	0.2757	DTI	-2.0812 (0.0384)	-1.693 (0.0917)	-3.4728 (<0.001)	-1.8207 (0.0698)	0.0849
		Combined	-2.0636 (0.0401)	-1.6405 (0.1021)	-3.4746 (<0.001)	-0.848 (0.3972)	0.0757
		T1	-10.1014 (<0.001)	0.3766 (0.7068)	1.098 (0.2733)	-2.5518 (0.0113)	0.2912
Cluster 5	0.2124	DTI	-10.118 (<0.001)	0.31 (0.7568)	1.1208 (0.2635)	-2.5199 (0.0124)	0.2908
		Combined	-10.2703 (<0.001)	0.5794 (0.5629)	1.0357 (0.3014)	-3.6468 (<0.001)*	0.3095
		T1	-7.1604 (<0.001)	-3.4563 (<0.001)*	-1.6891 (0.0924)	-1.0197 (0.3089)	0.2125
Cluster 6	0.51	DTI	-7.1405 (<0.001)	-3.4709 (<0.001)*	-1.6759 (0.095)	-0.5011 (0.6167)	0.21
		Combined	-7.2104 (<0.001)	-3.3405 (0.001)*	-1.7586 (0.0799)	-1.6393 (0.1024)	0.2177
		T1	-15.9382 (<0.001)	-1.0984 (0.2732)	-0.8171 (0.4147)	-1.8518 (0.0653)	0.5149
Cluster 7	0.1381	DTI	-15.9778 (<0.001)	-1.1226 (0.2627)	-0.8054 (0.4214)	-2.0589 (0.0406)	0.5165
		Combined	-16.0092 (<0.001)	-0.9591 (0.3385)	-0.8532 (0.3944)	-2.3608 (0.019)	0.5191
		T1	-6.5016 (<0.001)	-0.6594 (0.5102)	-0.5961 (0.5516)	-0.3018 (0.7631)	0.135
		DTI	-6.5363 (<0.001)	-0.643 (0.5208)	-0.5958 (0.5518)	-1.5824 (0.1148)	0.1432
		Combined	-6.5065 (<0.001)	-0.6229 (0.5339)	-0.6062 (0.5449)	-0.5676 (0.5708)	0.1358

892 **Table S1.** Cognitive associations with Brain Age Gap (BAG) using non-linear models,
893 including age, age² and sex as covariates – statistics. * FDR significant ** Bonferroni
894 significant.
895